Regulation of Ferroptosis in Obesity: Muscle Type-Specific Effects of Dietary Restriction and Exercise

Fujue Ji[1, 2], Yu Jin Park[1], Haesung Lee[1, 2], and Jong-Hee Kim[1, 2, #]

[1] Major in Sport Science, Division of Sport Industry and Science, College of Performing Arts and Sport,

Hanyang University, 222 Wangsimni-ro, Seongdong-gu, Seoul, Republic of Korea

[2] BK21 FOUR Human-Tech Convergence Program, Hanyang University, 222 Wangsimni-ro, Seongdong-gu,

Seoul 04763, Republic of Korea

[#] Corresponding Author

carachel07@hanyang.ac.kr

Background

Obesity is a significant global health issue and a risk factor for numerous diseases. Ferroptosis, an iron-dependent regulated cell death, is triggered by iron overload and the excessive accumulation of lipid peroxidation mediated by reactive oxygen species. Studies has identified a strong association between ferroptosis and obesity. Additionally, dietary restriction (DR) and DR combined with exercise (DR+Ex) are effective strategies for managing obesity and ferroptosis. However, the regulation of ferroptosis and its signaling pathways in skeletal muscle under conditions of obesity, DR, and DR+Ex remains poorly understood.

Methods

Mice were divided into four groups: normal diet, high-fat diet, high-fat DR, and high-fat DR+Ex. All mice were fed ad libitum with either a normal or high-fat diet for the first 14 weeks, followed by the respective interventions for the subsequent 8 weeks. Mice muscle ferroptosis were examined by immunohistochemistry, Hematoxylin & Eosin, Masson's trichrome, Prussian blue staining, and Western-Immunoblot.

Results

The high-fat diet resulted in increased inflammatory cell infiltration, fibrosis, and iron accumulation. Red and white muscles showed increased expression of 4-HNE, regulated by GPX4 and NCAO4, respectively. DR and DR+Ex reduced downstream 4-HNE expression by regulating GPX4 in red muscle.

Conclusion

Red and white muscles respond to obesity-induced ferroptosis through different signaling pathways. The regulation of ferroptosis by DR and DR+Ex is muscle type-specific. Specifically, red muscle is more sensitive to the regulation of ferroptosis signaling by DR and DR+Ex compared to white muscle.

Keywords: Obesity, Ferroptosis, Muscle type-specific, Dietary restriction, Exercise

1. Introduction

Obesity has emerged as a significant global health issue, affecting millions of individuals worldwide. Its detrimental effects on human health are well-documented, with numerous studies linking it to chronic diseases such as cardiovascular disorders, type 2 diabetes, and certain types of cancer (Blüher, 2019). Recent research has found that various chronic diseases caused by obesity are related to its induction of multiple cell death pathways, including ferroptosis (Stockwell et al., 2017).

Ferroptosis is an iron-dependent form of regulated cell death caused by iron overload and the excessive accumulation of lipid peroxidation products mediated by reactive oxygen species (ROS) (Dixon et al., 2012). This unique cell death mechanism has been implicated in the pathogenesis of various diseases across different organs and tissues, including neurodegenerative disorders, liver diseases, and kidney injury (Stockwell & Jiang, 2019). Studies have found that obesity can lead to tissue or organ damage and functional decline by regulating ferroptosis. Specifically, obesity induces ferroptosis signaling, leading to varying degrees of damage and functional impairment in the heart (Pei et al., 2021), liver (Luo et al., 2020), kidneys (Luo et al., 2020),and pancreas (Wei et al., 2020).

Unexpectedly, recent studies have highlighted the close association between ferroptosis and skeletal muscle-related disorders (Wang et al., 2020; Wang et al., 2022). Ferroptosis not only induces skeletal muscle cell death but also inhibits their proliferation and differentiation (Ding et al., 2021; Guerrero-Hue et al., 2019; Huang et al., 2021). Consequently, it is strongly associated with conditions such as sarcopenia, rhabdomyolysis, rhabdomyosarcoma, and exhaustive exercise-induced fatigue (Dächert et al., 2020; Guerrero-Hue et al., 2019; Huang et al., 2021; Xiao et al., 2022). These skeletal muscle disorders significantly impair patients' quality of life and physical function, and can lead to acute and chronic diseases such as diabetes, bone fractures, hypercalcemia, and acute kidney injury (Liu et al., 2014; Oliveira & Vaz, 2015; Tsekoura et al., 2017; Welch et al., 2020). However, the regulatory effect of obesity on ferroptosis in skeletal muscle is currently unclear.

Dietary restriction (DR) has been extensively studied for its potential health benefits, including increased lifespan, improved metabolic health, and reduced risk of age-related diseases (Mattson et al., 2017). DR not only ameliorates obesity-induced skeletal muscle dysfunction (Suga et al., 2014) but also reduces the excessive accumulation of ROS (Bevilacqua et al., 2005; Walsh et al., 2014). ROS causes lipid peroxidation, ultimately leading to ferroptosis by reacting with polyunsaturated fatty acids in phospholipid membranes (Han et al., 2020; Li et al., 2020). The inhibitory effect of DR on obesity and lipid peroxidation may be a potential mechanism to regulate the ferroptosis signaling pathway in skeletal muscle. Exercise (Ex) is widely recognized for its positive impact on health (Ruegsegger & Booth, 2018). Ex not only improves obesity- and ROS-induced skeletal muscle dysfunction (Ostler et al., 2014) but also modulates the expression of ferroptosis signaling in the brain, kidney, and nerves (Chen et al., 2023; Liu et al., 2022; Xiang et al., 2023). Dietary restriction combined with exercise (DR+Ex) is acknowledged as a method that more effectively counteracts the negative effects of obesity and lipid peroxidation than DR or Ex alone (Khachatoorian & Samara, 2018; Kim et al., 1996; Kvedaras et al., 2020; Li et al., 2017). However, the regulatory effects of DR and DR+Ex on obesity-induced ferroptosis in skeletal muscle remain unclear.

Given the critical role of skeletal muscle in overall health and the emerging evidence linking

79 ferroptosis to muscle disorders, it is imperative to investigate the regulation of ferroptosis in the
80 context of obesity, DR, and DR+Ex. Understanding the molecular mechanisms underlying these
81 processes could provide valuable insights into the development of targeted interventions for
82 obesity-related muscle disorders and promote overall health and well-being.

2. Methods

2.1 Animals and Experimental Design

Male C57BL/6N mice (14 months old) were randomly divided into four experimental groups:
Normal diet (ND, n=7), High-fat diet (HF, n=7), 20% high-fat dietary restriction (HFR, n=7), and
20% high-fat dietary restriction with voluntary wheel running exercise (HFRV, n=7). For the first
4 months, all mice were fed either a standard chow diet (ND; Teklad Global 2018, Envigo Inc.) or
a high-fat diet (HF; 45% kcal from fat, D12451, Research Diets Inc.). In the subsequent 4 months,
dietary restriction (DR) and dietary restriction combined with exercise (DR+Ex) interventions
were applied to the respective groups (Figure 1). The mice were housed in a controlled
environment, maintained at a temperature of 22 ± 2 °C with 50–60% humidity, and subjected to a
12-hour light/dark cycle. All animals had ad libitum access to water and were allowed physical
activity. The experimental procedures were approved by the Institutional Animal Care and Use
Committee (IACUC) of Hanyang University (HYU 2021-0239A).

Figure 1. Schematic overview experimental study design..

2.2 Body Composition

98 Mice were anesthetized with 40 mg/kg ketamine and 0.8 mg/kg medetomidine and analyzed for
99 body composition using an InAlyzer Dual-energy X-ray Absorptiometry (DEXA) system (Micro
100 Photonics Inc., PA, USA). The parameters measured included total mass (g), lean mass ratio (%),
101 fat mass (g), lean mass (g), fat content in tissue (%), and fat mass ratio (%).

2.3 Skeletal Muscle Function Tests

All mice underwent skeletal muscle function tests before euthanasia. Skeletal muscle function was
assessed in four groups of mice using tests for walking speed, endurance, physical activity, and
grip strength (Figure 1).
Rota-Rod test: Walking speed was assessed using the Rota-Rod test (Leem et al., 2022). An
adaptation period was conducted during the first three days, involving a pre-test exercise at 5 rpm
for 1 minute, once per day. The official test utilized an acceleration mode, where the speed

increased from 5 to 50 rpm over a 5-minute period (Model 76-0770, Harvard Apparatus Inc.,
Holliston, MA, USA). The latency to fall from the device was recorded. Each mouse was tested
three times with a ten-minute interval between tests, and the best performance out of the three
trials was used as the outcome measure.

Treadmill test: Endurance capacity was evaluated using a treadmill test (Alghannam et al., 2016).
Adaptation was performed three days before the official test, consisting of running at 5 cm/s for 5
minutes, once per day, on a 0-degree slope. During the official test, the treadmill speed increased
by 1 cm/s every 20 seconds starting from 5 cm/s, with a 0-degree slope. The trial ended when the
mouse touched the shock pad (set at 0.6 mA) three times.

Voluntary wheel test: Physical activity was measured using the voluntary wheel test (Manzanares
et al., 2018). Running distance was recorded using a voluntary wheel (MAN86130, Lafayette
Instrument Company, Lafayette, IN, USA), with each wheel rotation corresponding to a distance
of 0.4 m. The average running distance over a 5-day period was recorded for each mouse.

Inverted-cling grip test: Grip strength was assessed using the inverted-cling grip test (Graber et al.,
2013). Adaptation was conducted prior to the official test, with the adaptive test performed once a
day for three days. During the official test, mice were placed in the center of a wire mesh screen,
and a stopwatch was started. The screen was then rotated to an inverted position over 2 seconds,
with the mouse's head descending first, and held 40-50 cm above a padded surface. The time until
the mouse fell was recorded. This measurement was repeated three times with ten-minute intervals
between tests, and the maximum reading was recorded.

2.4 Preservation and Preparation of Skeletal Muscle
The anesthetized mice were euthanized by cervical dislocation. Following euthanasia, the
gastrocnemius (GAS) muscles were carefully dissected from the mice. The left leg muscles were
fixed in 10% formalin for 24 hours, then dehydrated and embedded in paraffin. Sections were
prepared for Hematoxylin and Eosin (H&E) staining, Prussian blue staining, Masson's trichrome
staining, and immunohistochemistry (IHC). The right leg muscles were snap-frozen in liquid
nitrogen and stored at –80°C for subsequent Western-immunoblot (WB) analyses.

2.5 Hematoxylin and Eosin (H&E) staining
Rehydrated tissue sections (5 μm) were stained using ClearView™ hematoxylin (MA0101010,
StatLab, McKinney, TX, USA) for 2-3 minutes, rinsed in tap water for 30-60 seconds,
differentiated in 1% hydrochloric acid alcohol for 5-10 seconds, washed in tap water for 3-5
minutes, and stained with ClearView™ eosin (MA0101015, StatLab, McKinney, TX, USA) for
1-2 minute. Subsequently, the sections were dehydrated, cleared, and mounted with neutral resin.
Images were acquired using a slide scanner (Axio Scan. Z1, Zeiss) at 200x magnification. For
each mouse, a single section of each skeletal muscle was randomly photographed to obtain 5
images, each covering an area of 0.271 square millimeters. The average of these five images was
used to represent the cross-sectional area (CSA) of that skeletal muscle for the mouse. The CSA of
each muscle fiber was automatically analyzed using "Cellpose," an open-source deep
learning-based segmentation tool (Stringer et al., 2021).

2.6 Prussian Blue Staining
After deparaffinization and rehydration, the paraffin tissue sections (5 μm) were stained using

Prussian Blue staining kit (G1422, Beijing Solarbio Science & Technology Co., Ltd.). Images were acquired using a slide scanner (Axio Scan. Z1, Zeiss) at 200x magnification. For each mouse, a single section of each skeletal muscle was randomly photographed to obtain 5 images, each covering an area of 0.271 square millimeters. The average of these five images was used to represent the value of that iron accumulation for the mouse. ImageJ software was used for the analysis.

2.7 Masson's Trichrome Staining

After deparaffinization and rehydration, the paraffin tissue sections (5 μm) were stained using Masson's trichrome staining kit (G1340, Beijing Solarbio Science & Technology Co., Ltd.). Images were acquired using a slide scanner (Axio Scan. Z1, Zeiss) at 200x magnification. For each mouse, a single section of each skeletal muscle was randomly photographed to obtain 5 images, each covering an area of 0.271 square millimeters. The average of these five images was used to represent the ECM value of that skeletal muscle for the mouse. ImageJ software was used for analysis. A threshold was established to remove the blank background, the ECM signal was isolated using color deconvolution and thresholding, and the area of the ECM signal was subsequently compared with the net area of the muscle section using ImageJ software.

2.8 Western-Immunoblot (WB)

The red and white GAS were dissected separately (Chen et al., 2015). The muscles were lysed using an EzRIPA Lysis Kit (WSE-7420, ATTO). Protein concentration was determined using the Pierce™ Bicinchoninic Acid Protein Assay Kit (Thermo Scientific, Waltham, MA, USA). Equal amounts of protein samples (40 μg) were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis and electro-transferred to nitrocellulose membranes (Bio-Rad Laboratories, Hercules, CA, USA). The membranes were blocked with 5% non-fat milk dissolved in Tris-buffered saline with Tween-20 (TBST; 10 mM Tris, 150 mM NaCl, and 0.1% Tween-20; pH 7.6) for 1.5 hours at room temperature and then incubated with primary antibodies (Table 1) overnight at 4°C. Subsequently, the membranes were incubated with horseradish peroxidase-conjugated secondary antibodies (Table 1) for 1.5 hours at room temperature. The targeted bands were quantified by densitometry using ImageJ software.

2.9 Immunohistochemistry (IHC)

Following the immunohistochemistry protocol with modifications based on Magaki et al., paraffin-embedded tissue sections (5 μm) were deparaffinized and rehydrated (Magaki et al., 2019). Heat-induced antigen retrieval was performed using 0.01 mol/L sodium citrate buffer (pH 6.0, C1010, Beijing Solarbio Science & Technology Co., Ltd.) in a microwave oven for 20 minutes at 100°C. The sections were then washed twice with Phosphate-buffered saline (PBS; 0.137M NaCl, 0.0027M KCL, 0.010mM Na_2HPO_4, and 0.002mM KH_2PO_4; pH 7.2) for 5 minutes each and blocked with 3% hydrogen peroxide (4104-4400, DAE JUNG) for 15 minutes twice to remove endogenous peroxidases. Following three additional 5-minute washes with PBS, the sections were blocked with 100-200 μl of blocking buffer (5% Bovine serum albumin in PBS) for 30 minutes at room temperature. After blocking, the sections were washed three times with PBS for 5 minutes each and incubated with the primary antibody (Table 1) overnight at 4°C. The sections were then washed three times with PBS for 5 minutes each and incubated with the

secondary antibody (Table 1) in a humidified chamber for 60 minutes at room temperature.
Following three more 5-minute washes with PBS, the sections were incubated with the Pierce™
DAB Substrate Kit (Thermo Scientific, #34002) for 6 minutes and washed three times with PBS
for 5 minutes each. Finally, the sections were rehydrated and mounted. Images were acquired
using a slide scanner (Axio Scan. Z1, Zeiss) at 200x magnification. For each mouse, a single
section of each skeletal muscle was randomly photographed to obtain 5 images, each covering an
area of 0.271 square millimeters. The average of these five images was used to represent the
antibody expression value of that skeletal muscle for the mouse. ImageJ software was used for
analysis.

Table 1. Primary and secondary antibodies.

Antibodies	Source	Vender	Catalog	Dilution (WB)	Dilution (IHC)
P53	Rabbit	Cell signaling	#30313	1:500	
SLC7A11	Rabbit	Abcam	ab175186	1:1000	
GAPDH	Rabbit	Cell signaling	#5174	1:3000	
GPX4	Rabbit	Abcam	ab125066	1:1000	
AMPK	Rabbit	Cell signaling	#5831	1:500	
p-AMPK	Rabbit	Cell signaling	#2535	1:500	
TFR1	Rabbit	Abcam	ab84036	1:500	
NCOA4	Rabbit	Invitrogen	#PA5-115626	1:500	
4-HNE	Rabbit	Abcam	ab46545	1:1000	1:200
MDA	Mouse	Abcam	ab243066	1:250	1:100
Adiponectin	Rabbit	Cell signaling	#2789	1:1000	
Leptin	Rabbit	Invitrogen	PA1-052	1:500	
Goat anti Rabbit	Goat	Invitrogen	#G-21234	1:5000	1:500
m-IgG Fc	-	SANTA	sc-525409	1:5000	1:25

2.10 Statistical Analysis
Statistical analyses were performed using GraphPad Prism software (version 9). Comparisons
between two groups were conducted using an unpaired Student's t-test. For comparisons among
three groups, a one-way analysis of variance (ANOVA) followed by a least significant difference
(LSD) post-hoc test was employed. To assess the effects of muscle type (white vs. red) and dietary
conditions (HF vs. DR vs. DR+Ex), a two-way ANOVA with Bonferroni's post-hoc test was
applied. Outliers were identified and removed using the Z-score approach prior to conducting
statistical analyses to ensure the robustness and reliability of the results. All results are expressed
as the mean ± standard deviation (SD). Statistical significance was set at $p < 0.05$, with asterisks
indicating the following levels of significance: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p <$
0.0001.

3. Results
3.1 High-Fat Diet-Induced Obesity Evokes Dysfunction and Ferroptosis in Skeletal Muscle
To investigate whether obesity regulates ferroptosis in skeletal muscle, we induced obesity in mice
by feeding them a high-fat diet for 14 weeks. All the data are shown in Figure 2.
DEXA analysis revealed that high-fat diet intake led to increases in body weight ($P < 0.0001$), lean

mass (P = 0.0021), fat mass (P < 0.0001), fat mass ratio (P < 0.0001), and fat mass in tissues (P < 0.0001), while the lean mass ratio decreased (P < 0.0001) (Figure 2A-H). These indicators are widely used to evaluate and identify mouse models of obesity (Chen et al., 2012; Koza et al., 2006; Y. Yang et al., 2014; Yeu et al., 2019). Consistent with impaired skeletal muscle function in obese mice (Graham et al., 2019; Vellers et al., 2017), we found that a high-fat diet significantly reduced walking speed (P = 0.0424), grip strength (P = 0.0053), and endurance in mice (P = 0.0217) (Figure 2I-K). These findings further validate the successful establishment of the obese mouse model.

Histological features of ferroptosis include the accumulation of inflammatory cells in organs and tissues (Zhang et al., 2023) and fibrosis (Yu et al., 2020). H&E staining revealed that although the CSA of HF skeletal muscle significantly increased (P = 0.00210), it was accompanied by substantial inflammatory cell infiltration (Figure 2M and O). Additionally, Masson's trichrome staining indicated a significant increase in skeletal muscle fibrosis in HF compared to NDC (P = 0.0110) (Figure 2N and P). Ferroptosis is highly associated with lipid peroxidation and abnormal iron accumulation (Dixon et al., 2012). As products of lipid peroxidation, malondialdehyde (MDA) and 4-hydroxynonenal (4-HNE) can reflect the extent of ferroptosis in cells (Chen et al., 2021; Tang et al., 2021). IHC analysis showed elevated levels of lipid peroxidation products 4-HNE (P = 0.0320) and MDA (P = 0.0150) in HF skeletal muscle compared to NDC (Figure 2Q-S). Furthermore, Prussian blue staining revealed abnormal iron accumulation in HF skeletal muscle (P = 0.0210) (Figure 2T-V). These data indicate that high-fat diet induces obesity, leading to skeletal muscle dysfunction and ferroptosis in mice.

I

J

K

L

NDC

NDC

HF

HF

H&E Staining

Masson Staining

M

N

O

P

IHC: Negative control

NDC

NDC

HF

HF

IHC: 4-HNE

IHC: MDA

Q

R

NDC

S T U

Figure 2. High-fat diet-induced obesity evokes dysfunction and ferroptosis in skeletal muscle. (A) Representative DEXA-scanned images of NDC and HF mice; skeletal muscle, fat tissue, and bone are shown in blue, red, and white, respectively. (B) Weekly body weight changes (group × weeks): NDC vs. HF (, 54-72 weeks). (C-H) Quantitative data of final total mass, lean mass, lean mass ratio, fat mass, fat mass ratio, and fat mass in tissue, respectively, were acquired by DEXA scanning. (I-L) Skeletal muscle function tests: (I) Walking speed test. (J) Strength test. (K) Endurance test. (L) Physical activity test. (M-N) Representative images showing H&E staining and Masson's trichrome staining of gastrocnemius. The collagen is indicated in blue and the muscle in red. (O) Skeletal muscle cross-sectional area in NDC and HF. (P) Skeletal muscle collagen area in NDC and HF. (Q) Representative images of IHC. (R) 4-HNE expression levels in skeletal muscle. (S) MDA expression levels in skeletal muscle. (T) Representative images of Prussian blue staining. The iron is indicated in blue and the muscle in red. (U) Iron percentage in skeletal muscle. Significant differences are denoted by asterisks: p < 0.05 (*), p < 0.01 (**), p < 0.001 (***), and p < 0.0001 (****). All values are presented as mean ± SD. Normal diet (ND, n=7), high-fat diet (HF, n=7). Scale bar = 50 μm.*

3.2 Muscle Type-Specific Regulation of Ferroptosis by GPX4 and NCOA4 in High-Fat Diet-Induced Obesity.

Given the observed ferroptosis in skeletal muscle as a result of obesity, we employed WB analysis to investigate the impact of obesity on the expression of representative proteins in five ferroptosis-related signaling pathways: amino acid metabolism, lipid metabolism, iron metabolism, lipophagy, and ferritinophagy in different muscle types. All the data are shown in Figure 3.

Under normal diet conditions, the expression of Glutathione peroxidase 4 (GPX4) (P = 0.0073) and Nuclear receptor coactivator 4 (NCOA4) (P = 0.0007) was significantly higher, and 4-HNE (P = 0.002) was lower in white muscle compared to red muscle (Figure 3A and H). Under high-fat diet conditions, the expression of NCOA4 was significantly lower in white muscle compared to red muscle (P < 0.0001) (Figure 3G). Additionally, within the same muscle type, a high-fat diet significantly decreased the expression of GPX4 (P = 0.0007) and NCOA4 (P = 0.0015) and increased 4-HNE (P < 0.0001) in white muscle compared to a normal diet (P < 0.05) (Figure 3A, G, and H). Although a high-fat diet did not alter the expression of GPX4 in red muscle, it significantly increased the expression of NCOA4 (P < 0.0001) and 4-HNE (P = 0.0375) (Figure 3A, G, and H). These data show muscle type-specific regulation of ferroptosis by GPX4 and NCOA4 in high-fat diet-induced obese mice. Specifically, obesity regulates ferroptosis by decreasing the expression of GPX4 in white muscle and increasing the expression of NCOA4 in red muscle.

Figure 4. Muscle type-specific regulation of ferroptosis by GPX4 and NOCA4 in high-fat diet-induced obese mice. WB was performed for protein expression levels of ferroptosis signaling pathways, (A): Glutathione peroxidase 4 (GPX4), (B): P53, (C): Solute Carrier Family 7 Member 11 (SLC7A11), (D): AMP-activated protein kinase (AMPK), (E): Ras-related protein Rab-7a (Rab7), (F): Telomeric repeat-binding factor 1 (TRF1), (G) Nuclear Receptor Coactivator 4 (NCOA4), (H): 4-Hydroxynonenal (4-HNE), and (I): Malonaldehyde (MDA). Significant differences are denoted by asterisks: $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***) and $p < 0.0001$ (****). All values are presented as mean ± SD. Normal diet (ND, n=7), High-fat diet (HF, n=7).

3.3 DR and DR+Ex Show Some Improvements and Regulation in Body Composition and Ferroptosis in Obese Mice.

Previous studies have extensively reported the positive effects of DR and DR+Ex on obesity, lipid peroxidation, ROS, and even ferroptosis (Bevilacqua et al., 2005; Chen et al., 2023; Li et al., 2017; Liu et al., 2022; Walsh et al., 2014). Therefore, we investigated the effects of DR and DR+Ex intervention on ferroptosis in the skeletal muscle of high-fat diet-induced obese mice. All the data are shown in Figure 4.

Results showed significant differences in body weight between the HF and HFRV groups at 70 weeks (P = 0.0435) (Figure 4B). Compared to the HF, HFRV significantly reduced total mass (P = 0.0061), lean mass (P = 0.0171), and fat mass in mice (P = 0.0159), but there were no changes in lean mass ratio, fat mass ratio, or fat mass in tissues (Figure 4C-H). DR and DR+Ex did not appear to effectively improve skeletal muscle function in high-fat diet-induced obese mice (Figure 4I-L).

Additionally, H&E staining showed that DR and DR+Ex reduced inflammatory cell infiltration in muscle fibers; however, DR+Ex also decreased the CSA of skeletal muscle (P = 0.0296) (Figure

4M and O). Masson's trichrome staining and Prussian blue staining indicated fibrosis and iron in skeletal muscle were not significant changes (Figure 4N, P, T, and U). Lastly, IHC analysis showed no significant differences in the expression of 4-HNE and MDA among the HF, HFR, and HFRV groups (Figure 4Q-S). These data suggest that intervention of DR and DR+Ex has limited effects on improving and regulating body composition and ferroptosis in high-fat diet-induced obese mice.

3.4 Muscle Type-Specific Regulation of Ferroptosis by GPX4 Through DR and DR+Ex in High-Fat Diet-Induced Obese Mice

CR and CR+Ex have regulatory effects on ferroptosis in the skeletal muscle of high-fat diet-induced obese mice. However, the detailed regulatory effects on muscle type-specific ferroptosis are not well understood. Therefore, it is worthwhile to further explore the muscle type-specific regulation of ferroptosis by these interventions through different pathways. The results of the WB analysis are shown in Figure 5.

Under the same conditions (HFRV), the expression of 4-HNE in white muscle was significantly higher compared to red muscle (P < 0.0001) (Figure 5H). In red muscle, the expression of GPX4 was significantly higher in the HFRV group compared to the HF group (P = 0.0471) (Figure 5A); the expression of 4-HNE decreased sequentially in the HF, HFR, and HFRV groups, with significant differences (P < 0.0001 and P = 0.0136) (Figure 5H). In white muscle, the expression of AMP-activated protein kinase (AMPK) was significantly lower in the HFRV group compared to the HF (P = 0.0086) and HFR (P = 0.0126) groups (Figure 5D), and there were significant differences in the expression of 4-HNE between the HFR and HFRV groups (P = 0.0427) (Figure 5H). These data suggest that the regulation of ferroptosis by DR and DR+Ex intervention in high-fat diet-induced obese mice is muscle type-specific. Specifically, DR+Ex inhibits obesity-induced skeletal muscle ferroptosis by increasing the expression of GPX4 in red muscle.

G   H   I

Figure 6. Through CR and CR+Ex muscle type-specific regulation of ferroptosis by GPX4 in high-fat diet-induced

obese mice. WB was performed for protein expression levels of ferroptosis signaling pathways, (A): Glutathione

peroxidase 4 (GPX4), (B): P53, (C): Solute Carrier Family 7 Member 11 (SLC7A11), (D): AMP-activated protein

kinase (AMPK), (E): Ras-related protein Rab-7a (Rab7), (F): Telomeric repeat-binding factor 1 (TRF1), (G)

Nuclear Receptor Coactivator 4 (NCOA4), (H): 4-Hydroxynonenal (4-HNE), and (I): Malonaldehyde (MDA).

Significant differences are denoted by asterisks: p < 0.05 (*), p < 0.01 (**), p < 0.001 (***) and p < 0.0001

(****). All values are presented as mean ± SD. High-fat diet (HF, n=7), 20% high-fat diet dietary restriction (HFR,

n=7), and 20% high-fat diet dietary restriction + voluntary wheel running exercise (HFRV, n=7).

4. Discussion

In recent years, the issue of dysfunction in cells, tissues, and organs caused by obesity has

garnered widespread attention. Obesity-induced ferroptosis is considered one of the primary

causes of these dysfunctions (Ganz, 2012; Ma et al., 2017; Orr et al., 2014; Zhao et al., 2022).

Understanding the regulatory mechanisms of ferroptosis in the context of obesity is of significant

clinical importance for related diseases. This study investigated the impact of obesity on

ferroptosis in specific skeletal muscle types and the response mechanisms of its signaling

pathways to DR and DR+Ex. Our research found that obesity can induce ferroptosis in skeletal

muscle, while DR and DR+Ex can regulate this process. More importantly, the regulation of

ferroptosis by obesity and its response to intervention are muscle type-specific. These findings

enhance our understanding of the mechanisms of obesity and ferroptosis, providing a scientific

basis for further in-depth studies of these diseases and cell death mechanisms.

4.1 Appropriateness of an Obesity Mouse Model.

Mice at 14 months of age are approximately equivalent to humans aged 35-45 years (Justice et al.,

2016). During human aging, the prevalence of obesity increases with age. Research has shown

that after the age of 40, the prevalence of obesity significantly increases (Chooi et al., 2019). In

our study, mice began a high-fat diet at 14 months to simulate continuous obesity starting around

age 40 in humans. We induced an obese mouse model through 4 months of a high-fat diet

(D12451, Research Diets Inc.). This diet has been widely used in the construction of obese mouse

models (Armani et al., 2014; Han et al., 2006; Yongbin Yang et al., 2014). Although there is no

precise definition for high-fat diet-induced obese mouse models, the vast majority of studies use at

least 8 weeks of ad libitum high-fat diet to create obese mouse models (Buettner et al., 2007; Han

et al., 2006; Hariri & Thibault, 2010).

Additionally, after analyzing the body composition of the mice using DEXA, we found significant

increases in body weight, lean mass, fat mass, fat mass in tissues, and fat mass percentage

following the high-fat diet. These changes in body composition are consistent with the

characteristics of obese mice reported in previous studies (Lin et al., 1977; Mandarim-de-Lacerda

et al., 2021; Rogers & Webb, 1980). Moreover, obesity leads to a decline in certain skeletal muscle

functions, which is also in line with prior research, indicating that dysfunctional skeletal muscle is

a hallmark of obesity (Shortreed et al., 2009; Zhang et al., 2015). Therefore, we conclude that a

4-month high-fat diet can successfully induce obesity, providing an experimental animal model for

subsequent study.

4.2 High-Fat Diet-Induced Obesity Evokes Ferroptosis in Mouse Skeletal Muscle

Through H&E staining, we found that high-fat diet-induced obesity leads to inflammatory cell

infiltration in skeletal muscle. In the obese state, the high expression levels of pro-inflammatory

cytokines, such as TNF-α, IL-6, and IL-1β, are primarily secreted by increased infiltrating

inflammatory cells (Hotamisligil, 2006; Weisberg et al., 2003). These findings are consistent with

what we observed in the skeletal muscles of high-fat diet-induced obese mice.

Recent studies suggest that not only obesity but also inflammatory cell infiltration is a hallmark of

ferroptosis. Ferroptosis causes lipid peroxidation of cell membranes and cell rupture, and the

release of cellular contents (such as high-mobility group protein B1) can activate a local

inflammatory response, leading to inflammatory cell infiltration. For instance, research by

Friedmann Angeli et al. (Friedmann Angeli et al., 2014) demonstrated that ferroptosis leads to

acute renal failure, accompanied by significant inflammatory cell infiltration. In the liver, where

ferroptosis occurs, inflammatory cell infiltration is also present, as shown in the hemochromatosis

mouse model study by Wang et al. (Wang et al., 2017). Inflammation not only hallmark of

ferroptosis but also promotes it.

Oxidative stress and iron metabolism disorders in the inflammatory response can exacerbate

ferroptosis, while oxidized lipid products released during ferroptosis can further promote

inflammation. The study by Dixon et al. (Dixon et al., 2012) pointed out the potential

inflammatory role of ferroptosis in various diseases. These research findings strongly support that

ferroptosis is accompanied by inflammatory cell infiltration. In summary, ferroptosis can induce a

local inflammatory response by releasing cellular contents, while inflammatory cell infiltration

can exacerbate ferroptosis, forming a vicious cycle. This interaction has important pathological

significance in various diseases, and further understanding of this mechanism will help develop

new therapeutic strategies.

Masson's trichrome staining revealed that high-fat diet-induced obesity exacerbates skeletal

muscle fibrosis. In the obese state, increased levels of oxidative stress in skeletal muscle induce

oxidative damage in muscle cells, activate fibroblasts, and promote fibrosis (Furukawa et al.,

2004). Inflammatory factors and oxidative stress induced by obesity not only cause damage to

muscle cells but also activate fibroblasts in muscle tissue. These activated fibroblasts secrete large

amounts of extracellular matrix components (ECM), such as collagen, leading to increased muscle

fibrosis (Lieber & Ward, 2013). These findings are consistent with what we observed in the

skeletal muscles of high-fat diet-induced obese mice.

Recently, the relationship between ferroptosis and tissue organ fibrosis has been explored in

several studies. Ferroptosis, an iron-dependent form of cell death, releases oxidized lipids and

cellular contents that can activate fibroblast proliferation, thereby promoting fibrosis. Specifically,

cellular contents released during ferroptosis, such as high-mobility group protein B1, can activate

fibroblasts and bone marrow-derived fibroblast-like cells. These cells, once activated, secrete large

amounts of ECM components like collagen, leading to tissue fibrosis(Kang et al., 2016; Wenzel et
al., 2017). For example, ferroptosis is considered a key pathological mechanism in liver and heart
fibrosis (Fang et al., 2020; Friedmann Angeli et al., 2014).
Moreover, ROS and lipid peroxides produced during ferroptosis not only directly cause cell
damage but also induce inflammatory responses that further exacerbate fibrosis. For instance,
ROS generated by ferroptosis can activate nuclear factor kappa B (NF-κB) and other
pro-inflammatory signaling pathways, promoting the release of pro-fibrotic factors (Dixon et al.,
2012; Stockwell et al., 2017). Synthesizing these studies, it is evident that ferroptosis exacerbates
tissue fibrosis through inducing inflammatory responses, activating fibroblasts, and increasing
oxidative stress. These mechanisms provide important theoretical foundations for understanding
the role of ferroptosis in fibrotic diseases and developing new therapeutic strategies.

Prussian blue staining revealed a significant increase in iron accumulation in the skeletal muscles
of high-fat diet-induced obese mice. Obesity is often accompanied by systemic metabolic
disorders, including altered iron metabolism. Studies have shown that obese individuals often
exhibit lower serum iron levels and elevated hepcidin levels (Ganz, 2012). This increase in
hepcidin inhibits iron absorption and release, leading to iron accumulation in the body (Ganz,
2012). Additionally, pro-inflammatory factors secreted in obesity, such as TNF-α and IL-6, not
only stimulate hepcidin synthesis but also directly affect iron metabolism, increasing iron
accumulation in cells and tissues (Hotamisligil, 2006; Weisberg et al., 2003). These findings are
consistent with what we observed in the skeletal muscles of high-fat diet-induced obese mice.
Ferroptosis is an iron-dependent form of cell death, and its extent can be reflected by the amount
of iron accumulation in cells, tissues, and organs. Iron accumulation is a critical trigger for
ferroptosis. Excess intracellular iron generates large amounts of ROS through the Fenton Reaction,
which induce lipid peroxidation and thereby trigger ferroptosis (Dixon et al., 2012; Stockwell et
al., 2017). These research findings strongly support that ferroptosis is accompanied by iron
accumulation.

IHC revealed a significant increase in the expression of 4-HNE and MDA in the skeletal muscles
of high-fat diet-induced obese mice. 4-HNE is a highly reactive aldehyde compound produced by
lipid peroxidation, primarily derived from the peroxidation of ω-6 polyunsaturated fatty acids
(Ayala et al., 2014). It can form adducts with proteins, DNA, and phospholipids, altering their
structure and function, thereby causing cellular damage (Murdolo et al., 2023). Studies have
shown that the levels of 4-HNE are significantly elevated in obese individuals, indicating the role
of lipid peroxidation in obesity (Guo et al., 2017; Murdolo et al., 2023). Similarly, MDA, another
product of lipid peroxidation, is commonly used to measure oxidative stress levels. High levels of
MDA indicate lipid damage to cell membranes and altered cellular function (Ahmed et al., 2016;
Savira et al., 2020). As with 4-HNE, elevated levels of MDA have been observed in obese
individuals (Ahmed et al., 2016; Savira et al., 2020).
Ferroptosis is accompanied by the accumulation of lipid peroxides and oxidative stress (Dixon et
al., 2012). This process is closely related to the expression of 4-HNE and MDA, as both
compounds are markers of lipid peroxidation (He et al., 2022; Park et al., 2021). During
iron-catalyzed lipid peroxidation, large amounts of ROS are produced, with 4-HNE and MDA
being key lipid peroxides. These compounds can react with phospholipids, proteins, and DNA in

cell membranes, leading to membrane damage, loss of protein function, and DNA damage,
ultimately resulting in cell death (He et al., 2022; Weber & Lanci, 2021). Iron generates highly
reactive hydroxyl radicals (•OH) through the Fenton Reaction, which further induce lipid
peroxidation, increasing the production of 4-HNE and MDA, enhancing oxidative stress within
cells, and promoting ferroptosis (Endale et al., 2023). Furthermore, 4-HNE is not only a product of
lipid peroxidation but also an important signaling molecule that can regulate various cellular
processes, including inflammatory responses, cell proliferation, and apoptosis (Park et al., 2021).
In summary, 4-HNE and MDA play critical roles as markers of lipid peroxidation in the process of
ferroptosis.
In summary, high-fat diet-induced obesity leads to inflammation cell infiltration, fibrosis, iron
metabolism dysregulation, and increased lipid peroxides in skeletal muscle. These negative factors
collectively contribute to the occurrence of ferroptosis in skeletal muscle.

4.3 Regulation of GPX4 in White Muscle by Obesity.

To better explore the regulatory mechanisms of ferroptosis in skeletal muscle induced by obesity,
we used WB to examine the expression of ferroptosis signaling pathways in different types of
muscle. GPX4 plays a key regulatory role in ferroptosis through amino acid metabolism by
inhibiting the cell death process via the reduction of lipid peroxides accumulation (Stockwell et al.,
2017). The deficiency or inactivation of GPX4 leads to an increase in lipid peroxides, thereby
triggering ferroptosis (W. S. Yang et al., 2014).
Under normal dietary conditions, the expression of GPX4 is significantly higher in white muscle
compared to red muscle. This may be due to the higher metabolic activity and oxidative stress
levels in white muscle, which require more antioxidant protection to maintain cellular homeostasis
(Brennan, 2008; Kitase et al., 2018). Furthermore, the fiber type of white muscle has higher
oxidative metabolic demands and ROS generation, further promoting the increased expression of
GPX4 to prevent damage caused by lipid peroxidation (Wesolowski et al., 2022).
However, under the high-fat diet-induced obesity, the expression of GPX4 in white muscle
significantly decreases, while its expression in red muscle remains unaffected. This may be due to
the higher sensitivity of white muscle to lipid peroxidation. A high-fat diet increases oxidative
stress in white muscle, leading to GPX4 depletion, whereas red muscle, with its stronger
antioxidant capacity, can maintain GPX4 levels (Katunga et al., 2015; Zhuang et al., 2021).
Additionally, higher metabolic activity and lipid accumulation in white muscle may also
contribute to the significant decrease in GPX4 expression (Shi et al., 2024).

4.4 Regulation of NCOA4 in Red Muscle by Obesity.

NCOA4 mediates ferritinophagy, the process of degrading ferritin to release stored iron, thereby
increasing intracellular free iron content. This promotes iron-catalyzed Fenton Reactions, leading
to lipid peroxidation and ferroptosis (Dowdle et al., 2014; Mancias et al., 2014). Studies have
shown that the absence of NCOA4 significantly reduces the occurrence of ferroptosis, further
demonstrating its crucial role in the ferroptosis process (Dowdle et al., 2014; Mancias et al.,
2014).
Under normal conditions, the expression of NCOA4 is significantly higher in white muscle
compared to red muscle. This may be due to the higher metabolic activity and iron demand in
white muscle, necessitating higher levels of NCOA4 to regulate iron metabolism and storage, thus

meeting the energy needs during high-intensity, short-duration contractions (Muckenthaler et al.,
2017; Palstra et al., 2013). Additionally, NCOA4 plays a vital role in maintaining iron homeostasis
and promoting muscle function in metabolically active white muscle by releasing iron from
ferritin through ferritinophagy (Gehrer et al., 2023).

However, a high-fat diet induced obesity impacts the expression of NCOA4 differently in red and
white muscles, with an increase in red muscle and a significant decrease in white muscle. White
muscle primarily relies on anaerobic metabolism and is more sensitive to lipid peroxidation and
metabolic stress. The increased metabolic stress caused by a high-fat diet enhances the white
muscle cells' need for oxidative stress defense, possibly downregulating NCOA4 expression to
reduce ferritinophagy activities, thereby protecting the cells damage from further ferroptosis. This
hypothesis requires further in-depth research.

4.5 Red and White Muscle Regulate 4-HNE Through NCOA4 and GPX4
Under normal conditions, the expression of 4-HNE in white muscle is lower than in red muscle.
This may be due to the higher oxidative metabolism rate and mitochondrial content in red muscle,
leading to increased generation of ROS, which trigger more lipid peroxidation reactions and
produce more 4-HNE (Morris et al., 2008; Paglialunga et al., 2015). Additionally, the antioxidant
defense mechanisms in red muscle are relatively weaker, resulting in lower clearance efficiency of
lipid peroxidation products like 4-HNE, further increasing its accumulation in red muscle (Hart et
al., 2018).

A high-fat diet-induced obesity leads to increased lipid accumulation and oxidative stress in both
red and white muscles, significantly enhancing the production of 4-HNE (Hart et al., 2018; Shi et
al., 2024). This diet also weakens antioxidant defense mechanisms, such as reducing the activity
of glutathione and superoxide dismutase, further exacerbating lipid peroxidation (Hart et al., 2018).
It is of paramount importance to note that the skeletal muscle type specific expression of GPX4
and NCOA4 as upstream and regulatory signals of 4-HNE was significantly modulated by
high-fat-induced obesity. Consequently, we propose that the upregulation of 4-HNE in red and
white muscle due to high-fat induced obesity is caused by significant alterations in the expression
of NCOA4 and GPX4, separately.

4.6 Obesity and Muscle Type do not Alter MDA Expression
After a high-fat diet, the expression of 4-HNE significantly increased in both red and white
muscles, whereas the expression of MDA did not show significant changes. This discrepancy may
be due to the differences in the generation and metabolism processes of 4-HNE and MDA. The
production of 4-HNE is highly influenced by ROS. A high-fat diet increases lipid peroxidation
reactions and ROS production, thereby significantly elevating 4-HNE levels (Morris et al., 2008;
Ramana et al., 2014). Additionally, 4-HNE is highly reactive and can form stable adducts with
various biomolecules such as proteins, DNA, and phospholipids, leading to its accumulation in
tissues (Ayala et al., 2014; Morris et al., 2008). However, the production of MDA is relatively
stable, and its metabolic pathways are more diverse and efficient. For example, MDA can be
metabolized through various pathways, such as being converted into water-soluble products by
glutathione-S-transferase for excretion or being transformed into less reactive compounds
(Esterbauer et al., 1991). These metabolic pathways make it less likely for MDA levels to show
significant changes under high-fat diet conditions (Del Rio et al., 2005).

Furthermore, some studies suggest that MDA is more likely to be rapidly produced and cleared
under acute oxidative stress conditions rather than accumulating under chronic conditions (Del
Rio et al., 2005). Previous studies have found that, after a high-fat diet, the expression of 4-HNE
significantly increased in some cells, tissues, and organs, while the expression of MDA did not
show significant changes (Bastin et al., 1990; Chung et al., 2018; Maciejczyk et al., 2018; Tanaka
et al., 2008). These studies indicate that due to the more diverse and efficient metabolic and
clearance mechanisms of MDA, its levels may not significantly change in certain tissues and
organs.
Additionally, experimental techniques can affect the expression of MDA. When using IHC and
WB to detect MDA, IHC showed a significant increase in MDA after a high-fat diet, whereas WB
did not show the same results. This discrepancy may arise from differences in the sensitivity and
specificity of the detection methods (Del Rio et al., 2005). IHC can detect local accumulation of
MDA in specific cell types or tissue regions, while WB measures overall protein levels, which
may dilute signals of local high expression. Furthermore, differences in sample processing and
extraction methods may lead to dilution or degradation of locally high concentrations of MDA in
WB, affecting the detection results. These factors collectively explain why IHC can show a
significant increase in MDA after a high-fat diet, while WB does not detect the same changes.

4.8 Regulation of Ferroptosis by DR and DR+Ex
We found that intervention with a DR did not improve the obesity-induced changes in body
composition and skeletal muscle dysfunction. The main reasons may include the following points.
Firstly, obesity is often accompanied by chronic low-grade inflammation, which can lead to
skeletal muscle dysfunction and adverse changes in body composition (Pedersen & Febbraio,
2008). A 20% DR may not be sufficient to significantly reduce inflammation levels. Secondly, a
high-fat diet leads to insulin resistance (Samuel & Shulman, 2012). Insulin resistance not only
affects glucose metabolism but also reduces muscle protein synthesis, thereby impacting skeletal
muscle function (Wilkinson et al., 2013). Thus, a 20% DR may not be enough to significantly
improve insulin sensitivity. Additionally, the metabolic disorders caused by obesity usually require
a longer duration and a greater degree of dietary intervention to reverse. A 20% high-fat dietary
restriction in terms of time and extent may not be sufficient to achieve the desired effects (Astrup
et al., 2008). Intervention with a 20% DR+Ex partially improved body composition but not
skeletal muscle function. This is because exercise can increase energy expenditure, promote fat
oxidation, and improve metabolic function (Hill & Wyatt, 2005). However, improving skeletal
muscle function requires a longer duration and specific types of stimuli, such as resistance training,
rather than just aerobic exercise (Kelley, 1998).

H&E staining showed that an DR and DR+Ex reduced inflammatory cell infiltration in skeletal
muscle. This is due to the anti-inflammatory effects of DR and Ex. However, Masson's trichrome
staining showed that these interventions did not reduce fibrosis in the skeletal muscles of obese
mice, likely because the pathological process of fibrosis is complex and difficult to significantly
affect with short-term interventions (Gillies & Lieber, 2011). Additionally, dietary adjustments
may not have been comprehensive enough, and the type and intensity of exercise may have been
insufficient to effectively inhibit fibrosis formation (Bonnard et al., 2008). Similarly, these
interventions did not reduce iron accumulation in the skeletal muscles of obese mice. This could

be due to insufficient dietary adjustments and exercise intensity and type, which may not have
significantly impacted iron metabolism (McClung & Karl, 2009). Moreover, iron accumulation is
a long-term process, and short-term interventions are unlikely to change existing iron deposits
(Domellöf et al., 2018). Finally, the diet may have lacked key nutrients such as vitamin C, which
can affect iron absorption and regulation (Zimmermann & Hurrell, 2007).

After DR and DR+Ex, the expression of 4-HNE and MDA in the skeletal muscles of obese mice
did not decrease. This may be attributed to the complexity of metabolic adaptation, oxidative
stress, and antioxidant mechanisms, as well as the diverse effects of different interventions on
skeletal muscle. Firstly, diet and exercise may lead to complex metabolic adaptations in skeletal
muscle. Studies have shown that although a high-fat diet induces obesity and insulin resistance, it
does not significantly alter the transcriptome expression in skeletal muscle, indicating that gene
expression changes under such interventions are minimal and difficult to detect (de Wilde et al.,
2009). Secondly, in the state of obesity, the levels of oxidative stress may already be very high,
and DR or DR+Ex alone may not be sufficient to significantly reduce the production of 4-HNE
and MDA. Research has shown that while DR and Ex improve metabolic indicators and skeletal
muscle function, they do not significantly alter the levels of specific oxidative stress markers (J. H.
Lee et al., 2021). Lastly, the effects of exercise and diet on skeletal muscle are diverse. Some
studies suggest that DR and Ex can improve obesity and insulin resistance, but the specific effects
on mitochondrial function and oxidative stress in skeletal muscle are complex, requiring further
research to clarify the comprehensive effects of these interventions (Pattanakuhar et al., 2019).
These factors may collectively contribute to the stability of 4-HNE and MDA levels in skeletal
muscle.

4.9 DR+Ex Regulated GPX4 Expression in Red Muscle

Although no significant regulatory effects of DR and DR+Ex on ferroptosis signals (4-HNE and
MDA) in the wholly skeletal muscles of obese mice were found, some regulatory trends did
emerge. Therefore, we used WB analysis to examine the regulatory mechanisms of DR and
DR+Ex on ferroptosis signals in different muscle types. The results showed that after DR+Ex, the
expression of GPX4 in red muscle significantly increased in obese mice, whereas it did not in
white muscle. This may be related to the oxidative stress, metabolic demands, and specific gene
regulatory mechanisms of different skeletal muscle types.
Firstly, red muscle primarily relies on aerobic metabolism and is rich in mitochondria, adapting to
prolonged low-intensity exercise. Therefore, its antioxidant mechanisms are more active to cope
with higher levels of oxidative stress. Studies have shown that antioxidant mechanisms in red
muscle are more prominent, potentially leading to the easier expression of GPX4 in response to
oxidative stress in this environment (Rinnankoski-Tuikka et al., 2012). Secondly, DR and Ex lead
to increased metabolic activity in skeletal muscle, requiring more antioxidant protection to prevent
lipid peroxidation damage (Civitarese et al., 2007; Galassetti et al., 2006). As a key antioxidant
enzyme, GPX4 is upregulated in this context, while white muscle, which primarily relies on
glycolysis for energy, has lower oxidative stress, and its GPX4 expression maybe does not
significantly increase. Lastly, DR and Ex can regulate gene expression through specific signaling
pathways such as AMPK and PGC-1α. These pathways are more active in red muscle (Gouspillou
et al., 2014; Jørgensen et al., 2007; Pengam et al., 2020) and are related to the expression of

antioxidant enzymes.

In summary, the increased expression of GPX4 in red muscle may be due to its high sensitivity to
oxidative stress, higher metabolic demands, and specific gene regulation induced by exercise. In
contrast, white muscle, with its different metabolic characteristics and lower oxidative stress levels,
did not show the same changes in GPX4 expression.

4.10 DR and DR+Ex Regulated AMPK Expression in White Muscle

We found that after DR+Ex in obese mice, the expression of AMPK decreased in white muscle but
not in red muscle. This may be related to the metabolic differences between muscle fiber types,
exercise-induced specific regulation, and the different impacts of diet and exercise on metabolic
regulation.

Red muscle primarily relies on aerobic metabolism, is rich in mitochondria, and is adapted to
prolonged low-intensity exercise, maintaining high AMPK activity to meet its metabolic demands
and antioxidant capacity. Under conditions of exercise and dietary restriction, AMPK expression
remains stable in red muscle. In contrast, white muscle primarily relies on glycolysis for energy,
has lower oxidative stress, and therefore requires less AMPK, leading to a possible decrease in its
expression (Jørgensen et al., 2007).

Additionally, exercise regulates gene expression through specific signaling pathways such as
AMPK and PGC-1α, which are more active and abundant in red muscle, thereby maintaining its
expression levels. However, in white muscle, lower AMPK activity leads to a reduction in its
expression (Sriwijitkamol et al., 2006). Lastly, the effects of a high-fat diet and exercise on
metabolic regulation differ between muscle fiber types. In red muscle, dietary restriction and
exercise can maintain high metabolic activity and antioxidant capacity, whereas in white muscle,
these regulatory effects are weaker, leading to a decrease in AMPK expression (J.-H. Lee et al.,
2021).

4.11 DR and DR+Ex Led to a Decrease in 4-HNE Expression in Red Muscle

After DR and DR+Ex in obese mice, the expression of 4-HNE decreased in red muscle but not in
white muscle. This difference is related to the metabolic differences between skeletal muscle types,
specific gene regulation, and the specific effects of diet and exercise on red muscle.

Red muscle primarily relies on aerobic metabolism, is rich in mitochondria, and is adapted to
prolonged low-intensity exercise. Consequently, its metabolic activity and oxidative stress are
higher, necessitating stronger antioxidant protection mechanisms. Studies have shown that red
muscle significantly increases antioxidant enzyme activity under DR and Ex, reducing the
production of oxidative stress markers such as 4-HNE (Jørgensen et al., 2007). Moreover, exercise
regulates gene expression through specific signaling pathways which are more active in red
muscle. This promotes the expression and activity of antioxidant enzymes, further reducing levels
of oxidative stress markers (Sriwijitkamol et al., 2006).

We found that as an upstream signal and regulatory molecule of 4-HEN, the expression of GPX4
in red muscle significantly increased after intervention. Therefore, we propose that DR and
DR+Ex can regulate ferroptosis by altering the expression of GPX4 in red muscle, which in turn
modulates the downstream expression of 4-HEN. However, AMPK, also an upstream regulatory
molecule of 4-HEN, showed inconsistent expression changes with 4-HNE in white muscle. After a
high-fat diet, the regulatory effect of AMPK on 4-HNE in white muscle was impaired, likely due

to metabolic disturbances caused by obesity. This warrants further investigation in future studies.

5. Conclusion

5. Conclusion

Obesity induced by a high-fat diet leads to ferroptosis in the skeletal muscle of mice. The
mechanism is muscle type-specific, with NCOA4 expression upregulated in red muscle and GPX4
expression upregulated in white muscle, ultimately triggering ferroptosis. Additionally, DR and
DR+Ex can partially mitigate ferroptosis in the skeletal muscle of obese mice induced by a
high-fat diet. This mechanism is achieved by regulating the expression of GPX4 in red muscle.
Overall, we have identified muscle type-specific regulatory mechanisms of ferroptosis, providing
molecular biology-related scientific data for the prevention and treatment of related diseases in the
future.

6. limitation

The study focuses on the gastrocnemius muscle, which may not fully represent the effects of
dietary restriction and exercise on other muscle groups, as different muscles may respond
differently to these interventions. Additionally, the duration of dietary restriction and exercise
intervention was limited to eight weeks. Longer-term studies are necessary to understand the
chronic effects and sustainability of these interventions on ferroptosis and overall muscle health.
Furthermore, incorporating advanced experimental tools, such as single-cell sequencing, could
provide a more detailed and microscopic analysis of the results.

7. WB representative picture

8. Ethical approval and consent to participate

All the procedures followed in this experiment were approved by the Institutional Animal Care
and Use Committee (IACUC) of Hanyang University (HYU 2021-0066A).

9. Consent for publication

Not applicable.

10. Availability of data and materials

The data used to support the findings of this study are presented here. Any further data requirements are available from the corresponding author upon request.

11. Competing interests

No conflicts of interest, financial or otherwise, have been declared by the author(s).

12. Funding

This research was supported by the National Research Foundation of Korea (NRF-2020R1F1A1061726).

13. Author's contributions

F.J.J. and J.H.K. conceived and designed the study. F.J.J., Y.J.P. and H.s.L. performed the experiments. F.J.J. and H.s.L. analyzed the data and prepared the figures. F.J.J. and J.H.K. interpreted the results, drafted, edited, and revised the manuscript. J.H.K. acquired funding. All authors approved the final version of the manuscript.

14. Acknowledgments

We thank Jian Guo (Hanyang University, Korea) for his generous help in dissecting the mice.

We are grateful to Prof. Dr. Gwang-woong Go (Hanyang University, Korea) for generously providing the DEXA machine for body composition analysis of the mice.

We thank the support of Skill Learning from Kaixin Doctor and MASCU (Medical Association with Science, Creativity, and Unity), Inc, Shenzhen, China (mascu_forever@163.com).

15. Reference

Ahmed, S., Maher, F., & Naji, N. (2016). Effect of leptin and oxidative stress in the blood of obese individuals. *Biochem Anal Biochem*, *5*(288), 2161-1009.

Alghannam, A. F., Jedrzejewski, D., Tweddle, M., Gribble, H., Bilzon, J. L., & Betts, J. A. (2016). Reliability of Time to Exhaustion Treadmill Running as a Measure of Human Endurance Capacity. *Int J Sports Med*, *37*(3), 219-223. https://doi.org/10.1055/s-0035-1555928

Armani, A., Cinti, F., Marzolla, V., Morgan, J., Cranston, G. A., Antelmi, A., Carpinelli, G., Canese, R., Pagotto, U., & Quarta, C. (2014). Mineralocorticoid receptor antagonism induces browning of white adipose tissue through impairment of autophagy and prevents adipocyte dysfunction in high-fat-diet-fed mice. *The FASEB Journal*, *28*(8), 3745-3757.

Astrup, A., Dyerberg, J., Selleck, M., & Stender, S. (2008). Nutrition transition and its relationship to the development of obesity and related chronic diseases. *Obes Rev*, *9 Suppl 1*, 48-52. https://doi.org/10.1111/j.1467-789X.2007.00438.x

Ayala, A., Muñoz, M. F., & Argüelles, S. (2014). Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxidative Medicine and Cellular Longevity*, *2014*.

Bastin, J., Djouadi, F., Geloso, J. P., & Merlet-Benichou, C. (1990). Postnatal development of oxidative enzymes in various rat nephron segments: effect of weaning on different diets. *Am J Physiol*, *259*(6 Pt 2), F895-901. https://doi.org/10.1152/ajprenal.1990.259.6.F895

Bevilacqua, L., Ramsey, J. J., Hagopian, K., Weindruch, R., & Harper, M.-E. (2005). Long-term caloric restriction increases UCP3 content but decreases proton leak and reactive oxygen species production in rat skeletal muscle mitochondria. *American Journal of Physiology-Endocrinology and Metabolism*, *289*(3), E429-E438.

Blüher, M. (2019). Obesity: global epidemiology and pathogenesis. *Nature Reviews Endocrinology*, *15*(5), 288-298.

Bonnard, C., Durand, A., Peyrol, S., Chanseaume, E., Chauvin, M. A., Morio, B., Vidal, H., & Rieusset, J. (2008). Mitochondrial dysfunction results from oxidative stress in the skeletal muscle of diet-induced insulin-resistant mice. *J Clin Invest*, *118*(2), 789-800. https://doi.org/10.1172/jci32601

Brennan, K. M. (2008). *Antioxidant gene expression and mitochondrial function during [beta]-oxidation in beef cattle* Washington State University].

Buettner, R., Schölmerich, J., & Bollheimer, L. C. (2007). High-fat diets: modeling the metabolic disorders of human obesity in rodents. *Obesity*, *15*(4), 798-808.

Chen, D., Wang, Y., & Chin, E. R. (2015). Activation of the endoplasmic reticulum stress response in skeletal muscle of G93A*SOD1 amyotrophic lateral sclerosis mice. *Front Cell Neurosci*, *9*, 170. https://doi.org/10.3389/fncel.2015.00170

Chen, J., Zhu, T., Yu, D., Yan, B., Zhang, Y., Jin, J., Yang, Z., Zhang, B., Hao, X., & Chen, Z. (2023). Moderate intensity of treadmill exercise rescues TBI-induced ferroptosis, neurodegeneration, and cognitive impairments via suppressing STING pathway. *Molecular Neurobiology*, *60*(9), 4872-4896.

Chen, W., Wilson, J. L., Khaksari, M., Cowley, M. A., & Enriori, P. J. (2012). Abdominal fat analyzed by DEXA scan reflects visceral body fat and improves the phenotype description and the assessment of metabolic risk in mice. *Am J Physiol Endocrinol Metab*, *303*(5), E635-643. https://doi.org/10.1152/ajpendo.00078.2012

Chen, X., Comish, P. B., Tang, D., & Kang, R. (2021). Characteristics and Biomarkers of Ferroptosis. *Front Cell Dev Biol*, *9*, 637162. https://doi.org/10.3389/fcell.2021.637162

Chooi, Y. C., Ding, C., & Magkos, F. (2019). The epidemiology of obesity. *Metabolism*, *92*, 6-10. https://doi.org/10.1016/j.metabol.2018.09.005

Chung, A., Gurtu, S., Chakravarthi, S., Moorthy, M., & Palanisamy, U. D. (2018). Geraniin Protects High-Fat Diet-Induced Oxidative Stress in Sprague Dawley Rats. *Front Nutr*, *5*, 17. https://doi.org/10.3389/fnut.2018.00017

Civitarese, A. E., Carling, S., Heilbronn, L. K., Hulver, M. H., Ukropcova, B., Deutsch, W. A., Smith, S. R., & Ravussin, E. (2007). Calorie restriction increases muscle mitochondrial biogenesis in healthy humans. *PLoS medicine*, *4*(3), e76.

Dächert, J., Ehrenfeld, V., Habermann, K., Dolgikh, N., & Fulda, S. (2020). Targeting ferroptosis in rhabdomyosarcoma cells. *Int J Cancer*, *146*(2), 510-520. https://doi.org/10.1002/ijc.32496

de Wilde, J., Smit, E., Mohren, R., Boekschoten, M. V., de Groot, P., van den Berg, S. A., Bijland, S., Voshol, P. J., van Dijk, K. W., de Wit, N. W., Bunschoten, A., Schaart, G., Hulshof, M. F., & Mariman, E. C. (2009). An 8-week high-fat diet induces obesity and insulin resistance with small changes in the muscle transcriptome of C57BL/6J mice. *J Nutrigenet Nutrigenomics*, *2*(6), 280-291. https://doi.org/10.1159/000308466

Del Rio, D., Stewart, A. J., & Pellegrini, N. (2005). A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *Nutr Metab Cardiovasc Dis*, *15*(4), 316-328. https://doi.org/10.1016/j.numecd.2005.05.003

Ding, H., Chen, S., Pan, X., Dai, X., Pan, G., Li, Z., Mai, X., Tian, Y., Zhang, S., Liu, B., Cao, G., Yao, Z., Yao, X., Gao, L., Yang, L., Chen, X., Sun, J., Chen, H., Han, M., . . . Xie, L. (2021). Transferrin receptor 1 ablation in satellite cells impedes skeletal muscle regeneration through activation of ferroptosis. *J Cachexia Sarcopenia Muscle*, *12*(3), 746-768. https://doi.org/10.1002/jcsm.12700

Dixon, S. J., Lemberg, K. M., Lamprecht, M. R., Skouta, R., Zaitsev, E. M., Gleason, C. E., Patel, D. N., Bauer, A. J., Cantley, A. M., & Yang, W. S. (2012). Ferroptosis: an iron-dependent form of nonapoptotic cell death. *cell*, *149*(5), 1060-1072.

Domellöf, M., Szitanyi, P., Simchowitz, V., Franz, A., & Mimouni, F. (2018). ESPGHAN/ESPEN/ESPR/CSPEN guidelines on pediatric parenteral nutrition: Iron and trace minerals. *Clin Nutr*, *37*(6 Pt B), 2354-2359. https://doi.org/10.1016/j.clnu.2018.06.949

Dowdle, W. E., Nyfeler, B., Nagel, J., Elling, R. A., Liu, S., Triantafellow, E., Menon, S., Wang, Z., Honda, A., Pardee, G., Cantwell, J., Luu, C., Cornella-Taracido, I., Harrington, E., Fekkes, P., Lei, H., Fang, Q., Digan, M. E., Burdick, D., . . . Murphy, L. O. (2014). Selective VPS34 inhibitor blocks autophagy and uncovers a role for NCOA4 in ferritin degradation and iron homeostasis in vivo. *Nature cell biology*, *16*(11), 1069-1079. https://doi.org/10.1038/ncb3053

Endale, H. T., Tesfaye, W., & Mengstie, T. A. (2023). ROS induced lipid peroxidation and their role in ferroptosis. *Frontiers in cell and Developmental Biology*, *11*.

Esterbauer, H., Schaur, R. J., & Zollner, H. (1991). Chemistry and biochemistry of 4-hydroxynonenal, malonaldehyde and related aldehydes. *Free Radic Biol Med*, *11*(1), 81-128. https://doi.org/10.1016/0891-5849(91)90192-6

Fang, X., Cai, Z., Wang, H., Han, D., Cheng, Q., Zhang, P., Gao, F., Yu, Y., Song, Z., & Wu, Q. (2020). Loss of cardiac ferritin H facilitates cardiomyopathy via Slc7a11-mediated ferroptosis. *Circulation research*, *127*(4), 486-501.

Friedmann Angeli, J. P., Schneider, M., Proneth, B., Tyurina, Y. Y., Tyurin, V. A., Hammond, V. J., Herbach, N., Aichler, M., Walch, A., & Eggenhofer, E. (2014). Inactivation of the ferroptosis regulator Gpx4 triggers acute renal failure in mice. *Nature cell biology*, *16*(12), 1180-1191.

Galassetti, P. R., Nemet, D., Pescatello, A., Rose-Gottron, C., Larson, J., & Cooper, D. M. (2006). Exercise, caloric restriction, and systemic oxidative stress. *Journal of investigative medicine*, *54*(2), 67-75.

Ganz, T. (2012). Macrophages and systemic iron homeostasis. *Journal of innate immunity*, *4*(5-6), 446-453.

Gehrer, C. M., Mitterstiller, A. M., Grubwieser, P., Meyron-Holtz, E. G., Weiss, G., & Nairz, M. (2023). Advances in Ferritin Physiology and Possible Implications in Bacterial Infection. *Int J Mol Sci*, *24*(5). https://doi.org/10.3390/ijms24054659

Gillies, A. R., & Lieber, R. L. (2011). Structure and function of the skeletal muscle extracellular matrix. *Muscle Nerve*, *44*(3), 318-331. https://doi.org/10.1002/mus.22094

Gouspillou, G., Sgarioto, N., Norris, B., Barbat-Artigas, S., Aubertin-Leheudre, M., Morais, J. A., Burelle, Y., Taivassalo, T., & Hepple, R. T. (2014). The relationship between muscle fiber type-specific PGC-1α content and mitochondrial content varies between rodent models and humans. *PloS one*, *9*(8), e103044. https://doi.org/10.1371/journal.pone.0103044

Graber, T. G., Ferguson-Stegall, L., Kim, J. H., & Thompson, L. V. (2013). C57BL/6 neuromuscular
healthspan scoring system. *J Gerontol A Biol Sci Med Sci*, *68*(11), 1326-1336.
https://doi.org/10.1093/gerona/glt032

Graham, L. C., Grabowska, W. A., Chun, Y., Risacher, S. L., Philip, V. M., Saykin, A. J., Rizzo, S. J. S.,
Howell, G. R., & Initiative, A. s. D. N. (2019). Exercise prevents obesity-induced cognitive
decline and white matter damage in mice. *Neurobiology of aging*, *80*, 154-172.

Guerrero-Hue, M., García-Caballero, C., Palomino-Antolín, A., Rubio-Navarro, A., Vázquez-Carballo, C.,
Herencia, C., Martín-Sanchez, D., Farré-Alins, V., Egea, J., Cannata, P., Praga, M., Ortiz, A.,
Egido, J., Sanz, A. B., & Moreno, J. A. (2019). Curcumin reduces renal damage associated with
rhabdomyolysis by decreasing ferroptosis-mediated cell death. *Faseb j*, *33*(8), 8961-8975.
https://doi.org/10.1096/fj.201900077R

Guo, L., Zhang, X.-M., Zhang, Y.-B., Huang, X., & Chi, M.-H. (2017). Association of 4-hydroxynonenal
with classical adipokines and insulin resistance in a Chinese non-diabetic obese population.
Nutricion hospitalaria, *34*(2), 363-368.

Han, C., Lee, S., Lim, S., Kong, H., Kim, S., Lee, S., & Chang, J. (2006). Studies on production of high fat
diet induced obesity C57BL/6NCrjBgi mice. *Laboratory Animal Research*, *22*(3), 221-226.

Han, C., Liu, Y., Dai, R., Ismail, N., Su, W., & Li, B. (2020). Ferroptosis and its potential role in human
diseases. *Frontiers in pharmacology*, *11*, 239.

Hariri, N., & Thibault, L. (2010). High-fat diet-induced obesity in animal models. *Nutrition research*
reviews, *23*(2), 270-299.

Hart, C. R., Layec, G., Trinity, J. D., Kwon, O. S., Zhao, J., Reese, V. R., Gifford, J. R., & Richardson, R. S.
(2018). Increased skeletal muscle mitochondrial free radical production in peripheral arterial
disease despite preserved mitochondrial respiratory capacity. *Exp Physiol*, *103*(6), 838-850.
https://doi.org/10.1113/ep086905

He, F., Huang, X., Wei, G., Lin, X., Zhang, W., Zhuang, W., He, W., Zhan, T., Hu, H., & Yang, H. (2022).
Regulation of ACSL4-catalyzed lipid peroxidation process resists cisplatin ototoxicity.
Oxidative Medicine and Cellular Longevity, *2022*.

Hill, J. O., & Wyatt, H. R. (2005). Role of physical activity in preventing and treating obesity. *J Appl*
Physiol (1985), *99*(2), 765-770. https://doi.org/10.1152/japplphysiol.00137.2005

Hotamisligil, G. S. (2006). Inflammation and metabolic disorders. *Nature*, *444*(7121), 860-867.

Huang, Y., Wu, B., Shen, D., Chen, J., Yu, Z., & Chen, C. (2021). Ferroptosis in a sarcopenia model of
senescence accelerated mouse prone 8 (SAMP8). *Int J Biol Sci*, *17*(1), 151-162.
https://doi.org/10.7150/ijbs.53126

Jørgensen, S. B., Treebak, J. T., Viollet, B., Schjerling, P., Vaulont, S., Wojtaszewski, J. F., & Richter, E. A.
(2007). Role of AMPKα2 in basal, training-, and AICAR-induced GLUT4, hexokinase II, and
mitochondrial protein expression in mouse muscle. *American Journal of*
Physiology-Endocrinology and Metabolism, *292*(1), E331-E339.

Justice, J. N., Cesari, M., Seals, D. R., Shively, C. A., & Carter, C. S. (2016). Comparative Approaches to
Understanding the Relation Between Aging and Physical Function. *J Gerontol A Biol Sci Med*
Sci, *71*(10), 1243-1253. https://doi.org/10.1093/gerona/glv035

Kang, R., Zeng, L., Xie, Y., Yan, Z., Zhou, B., Cao, L., Klionsky, D. J., Tracey, K. J., Li, J., & Wang, H. (2016).
A novel PINK1-and PARK2-dependent protective neuroimmune pathway in lethal sepsis.
Autophagy, *12*(12), 2374-2385.

Katunga, L. A., Gudimella, P., Efird, J. T., Abernathy, S., Mattox, T. A., Beatty, C., Darden, T. M., Thayne, K.
A., Alwair, H., Kypson, A. P., Virag, J. A., & Anderson, E. J. (2015). Obesity in a model of gpx4
haploinsufficiency uncovers a causal role for lipid-derived aldehydes in human metabolic
disease   and   cardiomyopathy.   *Mol   Metab*,   *4*(6),   493-506.
https://doi.org/10.1016/j.molmet.2015.04.001

Kelley, G. A. (1998). Aerobic exercise and bone density at the hip in postmenopausal women: a
meta-analysis. *Prev Med*, *27*(6), 798-807. https://doi.org/10.1006/pmed.1998.0360

Khachatoorian, Y., & Samara, A. (2018). Differential effects of dietary restriction combined with
exercise vs dietary restriction alone on visceral and subcutaneous adipose tissues: A
systematic review. *Obesity Medicine*, *9*, 7-17.

Kim, J. D., McCarter, R. J., & Yu, B. P. (1996). Influence of age, exercise, and dietary restriction on
oxidative stress in rats. *Aging (Milano)*, *8*(2), 123-129. https://doi.org/10.1007/bf03339566

Kitase, Y., Vallejo, J. A., Gutheil, W., Vemula, H., Jähn, K., Yi, J., Zhou, J., Brotto, M., & Bonewald, L. F.
(2018). β-aminoisobutyric Acid, l-BAIBA, Is a Muscle-Derived Osteocyte Survival Factor. *Cell
Rep*, *22*(6), 1531-1544. https://doi.org/10.1016/j.celrep.2018.01.041

Koza, R. A., Nikonova, L., Hogan, J., Rim, J. S., Mendoza, T., Faulk, C., Skaf, J., & Kozak, L. P. (2006).
Changes in gene expression foreshadow diet-induced obesity in genetically identical mice.
PLoS Genet, *2*(5), e81. https://doi.org/10.1371/journal.pgen.0020081

Kvedaras, M., Minderis, P., Krusnauskas, R., & Ratkevicius, A. (2020). Effects of ten-week 30% caloric
restriction on metabolic health and skeletal muscles of adult and old C57BL/6J mice.
Mechanisms of Ageing and Development, *190*, 111320.

Lee, J.-H., Zhang, D., Kwak, S.-E., Shin, H.-E., & Song, W. (2021). Effects of exercise and a high-fat,
high-sucrose restriction diet on metabolic indicators, Nr4a3, and mitochondria-associated
protein expression in the gastrocnemius muscles of mice with diet-induced obesity. *Journal of
Obesity & Metabolic Syndrome*, *30*(1), 44.

Lee, J. H., Zhang, D., Kwak, S. E., Shin, H. E., & Song, W. (2021). Effects of Exercise and a High-Fat,
High-Sucrose Restriction Diet on Metabolic Indicators, Nr4a3, and Mitochondria-Associated
Protein Expression in the Gastrocnemius Muscles of Mice with Diet-Induced Obesity. *J Obes
Metab Syndr*, *30*(1), 44-54. https://doi.org/10.7570/jomes20043

Leem, Y. H., Park, J. S., Park, J. E., Kim, D. Y., & Kim, H. S. (2022). Neurogenic effects of rotarod walking
exercise in subventricular zone, subgranular zone, and substantia nigra in MPTP-induced
Parkinson's disease mice. *Sci Rep*, *12*(1), 10544. https://doi.org/10.1038/s41598-022-14823-5

Li, C., Feng, F., Xiong, X., Li, R., & Chen, N. (2017). Exercise coupled with dietary restriction reduces
oxidative stress in male adolescents with obesity. *Journal of sports sciences*, *35*(7), 663-668.

Li, J., Cao, F., Yin, H.-l., Huang, Z.-j., Lin, Z.-t., Mao, N., Sun, B., & Wang, G. (2020). Ferroptosis: past,
present and future. *Cell Death & Disease*, *11*(2), 88.

Lin, P.-Y., Romsos, D. R., & Leveille, G. A. (1977). Food intake, body weight gain, and body composition
of the young obese (ob/ob) mouse. *The Journal of Nutrition*, *107*(9), 1715-1723.

Liu, C. K., Leng, X., Hsu, F. C., Kritchevsky, S. B., Ding, J., Earnest, C. P., Ferrucci, L., Goodpaster, B. H.,
Guralnik, J. M., Lenchik, L., Pahor, M., & Fielding, R. A. (2014). The impact of sarcopenia on a
physical activity intervention: the Lifestyle Interventions and Independence for Elders Pilot
Study (LIFE-P). *J Nutr Health Aging*, *18*(1), 59-64. https://doi.org/10.1007/s12603-013-0369-0

Liu, T., Cui, Y., Dong, S., Kong, X., Xu, X., Wang, Y., Wan, Q., & Wang, Q. (2022). Treadmill training reduces cerebral ischemia-reperfusion injury by inhibiting ferroptosis through activation of SLC7A11/GPX4. *Oxidative Medicine and Cellular Longevity*, *2022*.

Luo, Y., Chen, H., Liu, H., Jia, W., Yan, J., Ding, W., Zhang, Y., Xiao, Z., & Zhu, Z. (2020). Protective effects of ferroptosis inhibition on high fat diet-induced liver and renal injury in mice. *International Journal of Clinical and Experimental Pathology*, *13*(8), 2041.

Ma, X., Pham, V. T., Mori, H., MacDougald, O. A., Shah, Y. M., & Bodary, P. F. (2017). Iron elevation and adipose tissue remodeling in the epididymal depot of a mouse model of polygenic obesity. *PloS one*, *12*(6), e0179889.

Maciejczyk, M., Żebrowska, E., Zalewska, A., & Chabowski, A. (2018). Redox Balance, Antioxidant Defense, and Oxidative Damage in the Hypothalamus and Cerebral Cortex of Rats with High Fat Diet-Induced Insulin Resistance. *Oxid Med Cell Longev*, *2018*, 6940515. https://doi.org/10.1155/2018/6940515

Magaki, S., Hojat, S. A., Wei, B., So, A., & Yong, W. H. (2019). An Introduction to the Performance of Immunohistochemistry. *Methods Mol Biol*, *1897*, 289-298. https://doi.org/10.1007/978-1-4939-8935-5_25

Mancias, J. D., Wang, X., Gygi, S. P., Harper, J. W., & Kimmelman, A. C. (2014). Quantitative proteomics identifies NCOA4 as the cargo receptor mediating ferritinophagy. *Nature*, *509*(7498), 105-109.

Mandarim-de-Lacerda, C. A., Del Sol, M., Vazquez, B., & Aguila, M. B. (2021). Mice as an animal model for the study of adipose tissue and obesity. *Int J Morphol*, *39*(6), 1521-1528.

Manzanares, G., Brito-da-Silva, G., & Gandra, P. G. (2018). Voluntary wheel running: patterns and physiological effects in mice. *Braz J Med Biol Res*, *52*(1), e7830. https://doi.org/10.1590/1414-431x20187830

Mattson, M. P., Longo, V. D., & Harvie, M. (2017). Impact of intermittent fasting on health and disease processes. *Ageing research reviews*, *39*, 46-58.

McClung, J. P., & Karl, J. P. (2009). Iron deficiency and obesity: the contribution of inflammation and diminished iron absorption. *Nutr Rev*, *67*(2), 100-104. https://doi.org/10.1111/j.1753-4887.2008.00145.x

Morris, R. T., Laye, M. J., Lees, S. J., Rector, R. S., Thyfault, J. P., & Booth, F. W. (2008). Exercise-induced attenuation of obesity, hyperinsulinemia, and skeletal muscle lipid peroxidation in the OLETF rat. *J Appl Physiol (1985)*, *104*(3), 708-715. https://doi.org/10.1152/japplphysiol.01034.2007

Muckenthaler, M. U., Rivella, S., Hentze, M. W., & Galy, B. (2017). A Red Carpet for Iron Metabolism. *cell*, *168*(3), 344-361. https://doi.org/10.1016/j.cell.2016.12.034

Murdolo, G., Bartolini, D., Tortoioli, C., Vermigli, C., Piroddi, M., & Galli, F. (2023). Accumulation of 4-Hydroxynonenal Characterizes Diabetic Fat and Modulates Adipogenic Differentiation of Adipose Precursor Cells. *International Journal of Molecular Sciences*, *24*(23), 16645.

Oliveira, A., & Vaz, C. (2015). The role of sarcopenia in the risk of osteoporotic hip fracture. *Clin Rheumatol*, *34*(10), 1673-1680. https://doi.org/10.1007/s10067-015-2943-9

Orr, J. S., Kennedy, A., Anderson-Baucum, E. K., Webb, C. D., Fordahl, S. C., Erikson, K. M., Zhang, Y., Etzerodt, A., Moestrup, S. K., & Hasty, A. H. (2014). Obesity alters adipose tissue macrophage iron content and tissue iron distribution. *Diabetes*, *63*(2), 421-432.

Ostler, J. E., Maurya, S. K., Dials, J., Roof, S. R., Devor, S. T., Ziolo, M. T., & Periasamy, M. (2014). Effects
of insulin resistance on skeletal muscle growth and exercise capacity in type 2 diabetic mouse
models. *American Journal of Physiology-Endocrinology and Metabolism*, *306*(6), E592-E605.

Paglialunga, S., Ludzki, A., Root-McCaig, J., & Holloway, G. P. (2015). In adipose tissue, increased
mitochondrial emission of reactive oxygen species is important for short-term high-fat
diet-induced insulin resistance in mice. *Diabetologia*, *58*(5), 1071-1080.
https://doi.org/10.1007/s00125-015-3531-x

Palstra, A. P., Beltran, S., Burgerhout, E., Brittijn, S. A., Magnoni, L. J., Henkel, C. V., Jansen, H. J., Van
Den Thillart, G. E., Spaink, H. P., & Planas, J. V. (2013). Deep RNA sequencing of the skeletal
muscle transcriptome in swimming fish. *PloS one*, *8*(1), e53171.

Park, M. W., Cha, H. W., Kim, J., Kim, J. H., Yang, H., Yoon, S., Boonpraman, N., Yi, S. S., Yoo, I. D., &
Moon, J.-S. (2021). NOX4 promotes ferroptosis of astrocytes by oxidative stress-induced lipid
peroxidation via the impairment of mitochondrial metabolism in Alzheimer's diseases. *Redox*
biology, *41*, 101947.

Pattanakuhar, S., Sutham, W., Sripetchwandee, J., Minta, W., Mantor, D., Palee, S., Pratchayasakul, W.,
Chattipakorn, N., & Chattipakorn, S. C. (2019). Combined exercise and calorie restriction
therapies restore contractile and mitochondrial functions in skeletal muscle of obese-insulin
resistant rats. *Nutrition*, *62*, 74-84. https://doi.org/10.1016/j.nut.2018.11.031

Pedersen, B. K., & Febbraio, M. A. (2008). Muscle as an endocrine organ: focus on muscle-derived
interleukin-6. *Physiol Rev*, *88*(4), 1379-1406. https://doi.org/10.1152/physrev.90100.2007

Pei, Z., Liu, Y., Liu, S., Jin, W., Luo, Y., Sun, M., Duan, Y., Ajoolabady, A., Sowers, J. R., & Fang, Y. (2021).
FUNDC1 insufficiency sensitizes high fat diet intake-induced cardiac remodeling and
contractile anomaly through ACSL4-mediated ferroptosis. *Metabolism*, *122*, 154840.

Pengam, M., Moisan, C., Simon, B., Guernec, A., Inizan, M., & Amérand, A. (2020). Training protocols
differently affect AMPK-PGC-1α signaling pathway and redox state in trout muscle. *Comp*
Biochem Physiol A Mol Integr Physiol, *243*, 110673.
https://doi.org/10.1016/j.cbpa.2020.110673

Ramana, K. V., Srivastava, S., & Singhal, S. S. (2014). Lipid peroxidation products in human health and
disease 2014. *Oxid Med Cell Longev*, *2014*, 162414. https://doi.org/10.1155/2014/162414

Rinnankoski-Tuikka, R., Silvennoinen, M., Torvinen, S., Hulmi, J. J., Lehti, M., Kivelä, R., Reunanen, H., &
Kainulainen, H. (2012). Effects of high-fat diet and physical activity on pyruvate
dehydrogenase kinase-4 in mouse skeletal muscle. *Nutrition & metabolism*, *9*, 1-13.

Rogers, P., & Webb, G. P. (1980). Estimation of body fat in normal and obese mice. *British Journal of*
Nutrition, *43*(1), 83-86.

Ruegsegger, G. N., & Booth, F. W. (2018). Health benefits of exercise. *Cold Spring Harbor perspectives*
in medicine, *8*(7), a029694.

Samuel, V. T., & Shulman, G. I. (2012). Mechanisms for insulin resistance: common threads and missing
links. *cell*, *148*(5), 852-871. https://doi.org/10.1016/j.cell.2012.02.017

Savira, M., Rusdiana, S. S. W., & Syahputra, M. (2020). Comparison Malondialdehyde (MDA) Level
between Obesity Non Metabolic Syndrome and Obesity with Metabolic Syndrome Patients.
Age, *53*(11.3), 44.55-10.48.

Shi, Y., Zhong, L., Liu, Y., Xu, S., Dai, J., Zhang, Y., & Hu, Y. (2024). Dietary sanguinarine supplementation
recovers the decrease in muscle quality and nutrient composition induced by high-fat diets of

grass carp (Ctenopharyngodon idella). *Animal Nutrition*, *17*, 208-219. https://doi.org/https://doi.org/10.1016/j.aninu.2024.04.001

Shortreed, K. E., Krause, M. P., Huang, J. H., Dhanani, D., Moradi, J., Ceddia, R. B., & Hawke, T. J. (2009). Muscle-specific adaptations, impaired oxidative capacity and maintenance of contractile function characterize diet-induced obese mouse skeletal muscle. *PloS one*, *4*(10), e7293.

Sriwijitkamol, A., Ivy, J. L., Christ-Roberts, C., DeFronzo, R. A., Mandarino, L. J., & Musi, N. (2006). LKB1-AMPK signaling in muscle from obese insulin-resistant Zucker rats and effects of training. *American Journal of Physiology-Endocrinology and Metabolism*, *290*(5), E925-E932.

Stockwell, B. R., Angeli, J. P. F., Bayir, H., Bush, A. I., Conrad, M., Dixon, S. J., Fulda, S., Gascón, S., Hatzios, S. K., & Kagan, V. E. (2017). Ferroptosis: a regulated cell death nexus linking metabolism, redox biology, and disease. *cell*, *171*(2), 273-285.

Stockwell, B. R., & Jiang, X. (2019). A physiological function for ferroptosis in tumor suppression by the immune system. *Cell Metabolism*, *30*(1), 14-15.

Stringer, C., Wang, T., Michaelos, M., & Pachitariu, M. (2021). Cellpose: a generalist algorithm for cellular segmentation. *Nature Methods*, *18*(1), 100-106. https://doi.org/10.1038/s41592-020-01018-x

Suga, T., Kinugawa, S., Takada, S., Kadoguchi, T., Fukushima, A., Homma, T., Masaki, Y., Furihata, T., Takahashi, M., & Sobirin, M. A. (2014). Combination of exercise training and diet restriction normalizes limited exercise capacity and impaired skeletal muscle function in diet-induced diabetic mice. *Endocrinology*, *155*(1), 68-80.

Tanaka, Y., Aleksunes, L. M., Yeager, R. L., Gyamfi, M. A., Esterly, N., Guo, G. L., & Klaassen, C. D. (2008). NF-E2-related factor 2 inhibits lipid accumulation and oxidative stress in mice fed a high-fat diet. *J Pharmacol Exp Ther*, *325*(2), 655-664. https://doi.org/10.1124/jpet.107.135822

Tang, D., Chen, X., Kang, R., & Kroemer, G. (2021). Ferroptosis: molecular mechanisms and health implications. *Cell Research*, *31*(2), 107-125. https://doi.org/10.1038/s41422-020-00441-1

Tsekoura, M., Kastrinis, A., Katsoulaki, M., Billis, E., & Gliatis, J. (2017). Sarcopenia and Its Impact on Quality of Life. *Adv Exp Med Biol*, *987*, 213-218. https://doi.org/10.1007/978-3-319-57379-3_19

Vellers, H. L., Letsinger, A. C., Walker, N. R., Granados, J. Z., & Lightfoot, J. T. (2017). High Fat High Sugar Diet Reduces Voluntary Wheel Running in Mice Independent of Sex Hormone Involvement. *Front Physiol*, *8*, 628. https://doi.org/10.3389/fphys.2017.00628

Walsh, M. E., Shi, Y., & Van Remmen, H. (2014). The effects of dietary restriction on oxidative stress in rodents. *Free Radical Biology and Medicine*, *66*, 88-99.

Wang, H., An, P., Xie, E., Wu, Q., Fang, X., Gao, H., Zhang, Z., Li, Y., Wang, X., & Zhang, J. (2017). Characterization of ferroptosis in murine models of hemochromatosis. *Hepatology*, *66*(2), 449-465.

Wang, H., Liu, C., Zhao, Y., & Gao, G. (2020). Mitochondria regulation in ferroptosis. *European journal of cell biology*, *99*(1), 151058.

Wang, Y., Zhang, Z., Jiao, W., Wang, Y., Wang, X., Zhao, Y., Fan, X., Tian, L., Li, X., & Mi, J. (2022). Ferroptosis and its role in skeletal muscle diseases. *Frontiers in molecular biosciences*, *9*, 1051866.

Weber, D., & Lanci, R. (2021). Lipid peroxidation and cancer. *Cancer Adv*, *4*(6), 24.

Wei, S., Qiu, T., Yao, X., Wang, N., Jiang, L., Jia, X., Tao, Y., Wang, Z., Pei, P., & Zhang, J. (2020). Arsenic
induces pancreatic dysfunction and ferroptosis via mitochondrial ROS-autophagy-lysosomal
pathway. *Journal of hazardous materials*, *384*, 121390.

Weisberg, S. P., McCann, D., Desai, M., Rosenbaum, M., Leibel, R. L., & Ferrante, A. W. (2003). Obesity
is associated with macrophage accumulation in adipose tissue. *The Journal of clinical*
investigation, *112*(12), 1796-1808.

Welch, A. A., Hayhoe, R. P. G., & Cameron, D. (2020). The relationships between sarcopenic skeletal
muscle loss during ageing and macronutrient metabolism, obesity and onset of diabetes.
Proc Nutr Soc, *79*(1), 158-169. https://doi.org/10.1017/s0029665119001150

Wenzel, S. E., Tyurina, Y. Y., Zhao, J., Croix, C. M. S., Dar, H. H., Mao, G., Tyurin, V. A., Anthonymuthu, T.
S., Kapralov, A. A., & Amoscato, A. A. (2017). PEBP1 wardens ferroptosis by enabling
lipoxygenase generation of lipid death signals. *cell*, *171*(3), 628-641. e626.

Wesolowski, L. T., Semanchik, P. L., & White-Springer, S. H. (2022). Beyond antioxidants: Selenium and
skeletal   muscle   mitochondria.   *Front   Vet   Sci*,   *9*,   1011159.
https://doi.org/10.3389/fvets.2022.1011159

Wilkinson, D. J., Hossain, T., Hill, D. S., Phillips, B. E., Crossland, H., Williams, J., Loughna, P.,
Churchward-Venne, T. A., Breen, L., Phillips, S. M., Etheridge, T., Rathmacher, J. A., Smith, K.,
Szewczyk, N. J., & Atherton, P. J. (2013). Effects of leucine and its metabolite
β-hydroxy-β-methylbutyrate on human skeletal muscle protein metabolism. *J Physiol*, *591*(11),
2911-2923. https://doi.org/10.1113/jphysiol.2013.253203

Xiang, Y.-Y., Baek, K.-W., Won, J.-H., Park, Y., Kim, J.-S., Xiang, Y.-Y., Baek, K.-W., Won, J.-H., Park, Y., &
Kim, J.-S. (2023). Effects of Lifelong Aerobic Exercise on Ferroptosis-Related Gene Expressions
in Kidney of Aged Mice. *Exercise Science*, *32*(4), 410-418.

Xiao, R., Wei, Y., Zhang, Y., Xu, F., Ma, C., Gong, Q., Gao, J., & Xu, Y. (2022). Trilobatin, a Naturally
Occurring Food Additive, Ameliorates Exhaustive Exercise-Induced Fatigue in Mice:
Involvement of Nrf2/ARE/Ferroptosis Signaling Pathway. *Front Pharmacol*, *13*, 913367.
https://doi.org/10.3389/fphar.2022.913367

Yang, W. S., SriRamaratnam, R., Welsch, M. E., Shimada, K., Skouta, R., Viswanathan, V. S., Cheah, J. H.,
Clemons, P. A., Shamji, A. F., & Clish, C. B. (2014). Regulation of ferroptotic cancer cell death
by GPX4. *cell*, *156*(1), 317-331.

Yang, Y., Smith, D. L., Jr., Keating, K. D., Allison, D. B., & Nagy, T. R. (2014). Variations in body weight,
food intake and body composition after long-term high-fat diet feeding in C57BL/6J mice.
Obesity (Silver Spring), *22*(10), 2147-2155. https://doi.org/10.1002/oby.20811

Yang, Y., Smith Jr, D. L., Keating, K. D., Allison, D. B., & Nagy, T. R. (2014). Variations in body weight,
food intake and body composition after long-term high-fat diet feeding in C57BL/6J mice.
Obesity, *22*(10), 2147-2155.

Yeu, J., Ko, H. J., Kim, D., Ahn, Y., Kim, J., Lee, W., Jung, I., Suh, J., & Lee, S. J. (2019). Evaluation of
iNSiGHT VET DXA (Dual-Energy X-ray Absorptiometry) for assessing body composition in
obese rats fed with high fat diet: a follow-up study of diet induced obesity model for 8 weeks.
Lab Anim Res, *35*, 2. https://doi.org/10.1186/s42826-019-0004-2

Yu, Y., Jiang, L., Wang, H., Shen, Z., Cheng, Q., Zhang, P., Wang, J., Wu, Q., Fang, X., & Duan, L. (2020).
Hepatic transferrin plays a role in systemic iron homeostasis and liver ferroptosis. *Blood, The*
Journal of the American Society of Hematology, *136*(6), 726-739.

Zhang, X., Ma, Y., Lv, G., & Wang, H. (2023). Ferroptosis as a therapeutic target for
inflammation-related intestinal diseases. *Front Pharmacol*, *14*, 1095366.
https://doi.org/10.3389/fphar.2023.1095366

Zhang, Y., Fischer, K. E., Soto, V., Liu, Y., Sosnowska, D., Richardson, A., & Salmon, A. B. (2015).
Obesity-induced oxidative stress, accelerated functional decline with age and increased
mortality in mice. *Archives of biochemistry and biophysics*, *576*, 39-48.

Zhao, X., Si, L., Bian, J., Pan, C., Guo, W., Qin, P., Zhu, W., Xia, Y., Zhang, Q., & Wei, K. (2022). Adipose
tissue macrophage-derived exosomes induce ferroptosis via glutathione synthesis inhibition
by targeting SLC7A11 in obesity-induced cardiac injury. *Free Radical Biology and Medicine*,
182, 232-245.

Zhuang, A., Yang, C., Liu, Y., Tan, Y., Bond, S. T., Walker, S., Sikora, T., Laskowski, A., Sharma, A., de Haan,
J. B., Meikle, P. J., Shimizu, T., Coughlan, M. T., Calkin, A. C., & Drew, B. G. (2021). SOD2 in
skeletal muscle: New insights from an inducible deletion model. *Redox Biol*, *47*, 102135.
https://doi.org/10.1016/j.redox.2021.102135

Zimmermann, M. B., & Hurrell, R. F. (2007). Nutritional iron deficiency. *Lancet*, *370*(9586), 511-520.
https://doi.org/10.1016/s0140-6736(07)61235-5