**Metabolic Segregation and Characteristic Gene Clusters of the Key Functional Populations in Anaerobic Digestion Consortia**

Yubo Wang[1,2#], Ruoqun Zhang[1], Chunxiao Wang[2], Weifu Yan[2], Tong Zhang[2], Feng Ju[1#]

Author Affiliation:

1. Institute of Advanced Technology, Westlake Institute for Advanced Study, Hangzhou 310024, Zhejiang Province, China

2.Environmental Microbiome Engineering and Biotechnology Laboratory, The University of Hong Kong, Pokfulam, China

[#]Corresponding authors:

Yubo Wang, Email: Y.Wang-43@tudelft.nl, phone: +31(06)17877127

Feng Ju, Email: jufeng@westlake.edu.cn, Ph: +86-571-87380995

**Abstract**

Enrichment experiment integrated with genome-centric meta-omics analysis revealed function

partition among populations in the three functional niches of the methanogenic bioprocess:

primary fermentation, secondary fermentation (syntrophic acetogenesis) hydrogenotrophic and

acetoclastic methanogenesis. Accumulation of $H_2$ as a metabolic intermediate originates

mainly from primary fermentation. Formate, rather than $H_2$, plays a key role in secondary

fermentation, mediating interspecies electron transfer from the syntrophic oxidizing bacteria to

the formate-utilizing methanogens. Such segregation of the electron carrier pools by $H_2$ and

formate in primary fermentation and in secondary fermentation, respectively, can benefit the

syntrophic oxidizers with a higher tolerance to the accumulation of $H_2$ from the co-occurring

primary fermentation. It is interesting to note that, under $H_2$ concentration as high as 50,975

ppm, degradation of either butyrate or propionate was not completely inhibited. Metabolic

specificity and characteristic genes/gene clusters are noted for anaerobes of different function

niches enriched in this study, i.e., primary fermenting bacteria Clostridium FI21,22, syntrophic

butyrate oxidizer in *Syntrophomoas* FI3,4,7, syntrophic propionate oxidizer *Smithellaceae*

DES18-25, and formate- and $H_2$-utilizing methanogen *Methanobacteriaceae* EA7.

**Keywords:** Anaerobic digestion, Enriched microbiota, Genome-centric meta-omics, Function

partition, Characteristic gene clusters.

## 1. Introduction

Anaerobic digestion (AD) is of central importance in the field of environmental biotechnology since 1980s. It is one of the most cost-effective approaches to stabilize organic components in high-strength waste streams, aiding in bioenergy production and biosolids reduction[1, 2, 3]. However, challenges persist in addressing process upsets such as over-acidification and foam formation[4]. Fine-tuned control and optimization of the AD process depend on a deeper understanding of the sequential microbial reactions mediated by anaerobe populations across the four major functional niches: hydrolysis (conversion of polymers to monomers), primary fermentation/acidogenesis (conversion of monomers to $H_2$/HCOOH, acetate and other reduced fatty acids), secondary fermentation (conversion of reduced fatty acids to acetate and $H_2$/HCOOH) and methanogenesis ($CH_4$ generation mainly from acetate and $H_2$/HCOOH)[5].

Due to the limited amount of energy generated, secondary fermentation is often regarded as the bottleneck step in the AD bioprocess[6]. One of the objectives of this study is to chart the physiology of anaerobes in this specific function niche, specifically, syntrophic butyrate oxidizing bacteria (SBOBs) and syntrophic propionate oxidizing bacteria (SPOBs) in the AD consortia. As key intermediates in the anaerobic mineralization of organic compounds, propionate and butyrate can account for 20-43% of the total methane formation[2].

One of the well-accepted metabolic traits of the SBOBs and the SPOBs is their thermodynamic sensitivity to $H_2$, with an estimated $H_2$ concentration of ~300 ppm being high enough to inhibit the anaerobic oxidation of propionate or butyrate[7]. In the established scheme of methanogenic carbon cycling, the metabolism of almost all substrates, such as carbohydrates and amino acids, generates $H_2$[8]. To our knowledge, there has been little systematic characterization on the dynamics (generation and turnover) of $H_2$ during the AD bioprocess[9] and little evaluation has been conducted on its impact on the conversion of volatile fatty acids (VFAs) beyond theoretical calculations [10]. Inhibition concentration of $H_2$ characterized at 2.03

kPa for anaerobic butyrate oxidation and at 11.1 kPa for propionate oxidation are much higher

than 300 ppm calculated theoretically [7, 11]. Additionally, evidence is accumulating that suggests

that formate plays a more important role than $H_2$ as an interspecies electron carrier in syntrophic

oxidation [12]. This study aims to address this knowledge gap by combining the characterization

of $H_2$ dynamics with the degradation dynamics of VFAs in lab-scale AD reactors.

Full-scale anaerobic digesters rarely harbor high numbers of syntrophic bacteria[3],

although they do contain highly abundant primary fermenting populations. Given that

syntrophic oxidation of reduced fatty acids proceeds at the thermodynamic limit of life[11],

another scientific question to be addressed in this study is: are the SBOBs and SPOBs obligate

syntroph, or they more commonly are opportunists capable of fermentation flexibility [11, 13]?

Many uncharacterized and uncultivated bacteria in the AD consortia have yet-to-be-identified

relationships with the two trophic groups, i.e., primary fermentation *vs.* secondary fermentation,

and an in-depth understanding on the metabolic versatility/specificity of the syntrophic

populations could help elucidate the intricate ecology that supports the methanogenic

conversion of organic compounds.

In this study, to obtain consortia dominated by primary fermenting bacteria, SBOBs and

SPOBs, enrichment experiment was conducted in three sets of reactors, using glucose, butyrate

and propionate as the substrates, respectively. Metabolic versatility of the enriched anaerobes

in the three function guilds was evaluated by their metabolic capacity towards and

transcriptome response to the presence of either the fermentative substrates (i.e., glucose,

fumarate, crotonate) or the reduced short-chain fatty acids (SCFAs, i.e., propionate, butyrate).

Dynamics of the generation and turnover of $H_2$ as an intermediate metabolite, and also the

impact of $H_2$ accumulation on the syntrophic oxidation of propionate and butyrate, were

monitored throughout the batch experiments.

**2. Material and methods**

 *2.1 Enrichment of primary fermenting bacteria, SBOBs and SPOBs*

Three sets of AD consortia were enriched for the degradation of glucose, propionate and

butyrate, respectively. Duplicate enrichments utilizing each substrate were processed in 1000-

ml serum bottles with working volume of 500 ml. Hereafter, the consortia enriched with

glucose as substrate will be referred to as 'Consortia-G', the consortia enriched with butyrate

as substrate will be referred to as 'Consortia-B', and the consortia enriched with propionate as

substrate will be referred to as 'Consortia-P'.

The inoculum used in this enrichment experiment was surplus sludge sampled from Chengxi

wastewater treatment plant (Hangzhou, China). A detailed description on the seed sludge used,

the feeding strategy, and methods for the physiochemical characterization of the biogas

contents and the volatile fatty acids can be found in the supporting information.

*2.2 Characterization on the metabolic specificity of the primary fermenting bacteria, SBOBs*

*and SPOBs*

To characterize the metabolic specificity of the primary fermenting bacteria, SBOBs and

SPOBs, experiment was designed as summarized in Figure S2, to test the metabolic response

of the enriched primary fermenting bacteria, SBOBs and SPOBs to different substrates applied.

Take the test conducted on SBOBs enriched in Consortia-B as an example: after 219 days

enrichment, the 1000 ml (500 ml *2) sludge from two parallel reactors of Consortia-B, in which

SBOBs have been enriched, are combined and equally distributed into eight 200 ml serum

bottles (each with a 125 ml working volume). In the subsequent batch tests, the primary

substrate was sequentially switched from n-butyrate, to propionate, to glucose, to crotonate,

and   to   the   combined   co-substrate   mix   of   propionate:n-butyrate:glucose

(COD:COD:COD=1:1:1). Degradation profile (Figure S3, S4, S5) of the varying substrates by

the enriched SBOBs, and transcriptome response of the enriched SBOBs to the varying

substrates applied was charted.

 *2.3 Biomass sampling, Nucleic Acid Extraction, and meta-omics analysis*

     To profile the succession of the microbial community during the enrichment process,

 sludge samples were archived weekly to biweekly for sequencing-based analysis. Genomic

 DNA from the seed sludge (day-0), and from sludge of the enriched consortia collected on day-

 85, day-134 and day-219 were subjected to metagenome sequencing. When metatranscriptome

 analysis is relevant, sludge samples (15 ml) collected at each time point were flash frozen in

 liquid nitrogen and stored in -80 ℃ freezer prior to total RNA extraction. Metatranscriptome

 sequencing was conducted on 20 of the sludge samples collected in section 2.2 (Table S1,

 Figure S3, S4, S5). A more detailed documentation on the genome-centric meta-omics analysis,

 the methodologies for 16S rRNA gene (v4-v5 region) based microbiome profiling and the

 corresponding principal coordinate analysis (PCoA) are provided in the supporting information.

 *3. Results and discussion*

 *3.1 Effects of glucose, butyrate and propionate as the primary substrate on the succession of*

      *the anaerobic consortia enriched*

 Slow growth rates of the SBOBs and the SPOBs, and the selecting pressure imposed by SCFAs

 as the primary substrate was reflected by the significant decrease in the volatile suspended

 solids (VSS, a surrogate measure of the microbial biomass) in the Consortia-B and Consortia-

 P enriched. As summarized in Figure S6, at the same organic loading rate in terms of COD/L/d,

 in contrast to ~43% increment in VSS from 8.0 g/L to 11.7 g/l in Consortia-G, both Consortia-

 B and Consortia-P were observed with a significant decrease in VSS by ~50% from 8.0 g/l to

 4.1~4.4 g/l. Despite of the varying VSS contents, equally promising methanogenic conversion

 of glucose, butyrate and propionate, respectively, was observed in the three consortia, i.e.,

 Consortium-G, Consortia-B and Consortia-P (Figure S1).

     A distinct shift in the composition of the microbiome is noted during the enrichment

 process. A detailed summary on the taxonomy affiliation of the top 15 genera and their relative

abundance over the course of the enrichment can be found in Figure 1. Additionally

summarized in Figure 2 is the PCoA analysis on the composition of the three consortia enriched.

Over the course of the 219-days enrichment, genera that get characteristically enriched in

Consortia-G are: *Clostridium sensu stricto3*, *Propionimicrobium* and an unclassified genus in

the family of Propionibacteriaceae (g_un_f_Propionibacteriaceae). The one genus that gets

characteristically enriched in Consortia-P is *Smithella*, and the one genus that gets

characteristically enriched in Consortia-B is *Syntrophomonas*. Herein, we speculate that

*Clostridium sensu stricto3*, *Propionimicrobium* and g_un_f_Propionibacteriaceae are primary

fermenting bacteria utilizing glucose; *Smithella* is the putative SPOB oxidizing propionate

anaerobically, and *Syntrophomonas* the putative SBOB oxidizing butyrate anaerobically.

As is summarized in Figure S7-a, after 219 days enrichment, the top 15 genera accounting

for 75~83% of the populations in the three enriched consortia are of only ~4% relative

abundance in the seed inoculum. Among these top 15 genera are two archaeal methanogen

genera: *Methanosaeta* and *Methanobacterium*, and a much higher relative abundance of these

two methanogen genera, especially *Methanobacterium* (Figure S7-b), were annotated in

Consortia-B (20%) and Consortia-P (22%) than in Consortia-G (6%).

***3.2 Genome-centric identification of the primary fermenting bacteria, SBOBs and SPOBs***

***enriched***

317 bacterial metagenome assembled genomes (MAGs) are recovered from the metagenome

datasets. A summary of their taxonomy affiliation, their relative abundance and their

transcription activity can be found in Figure 3 and Appendix file 1. In the following genome-

centric meta-omics analysis, we specifically focus on those 17 bacterial MAGs that have

transcription response (>=0.5% of total mRNA clean reads mapped) to at least one of the

substrates provided, i.e., glucose, propionate, butyrate, fumarate and crotonate. These 17

MAGs include FI21 and FI22 (hereafter, these two MAGs will be referred to as FI21, 22)

affiliated with the genus of *Clostridium_T*; DES18, DES19, DES20, DES21, DES22, DES23,

DES24, and DES25 (hereafter, these eight MAGs in the same genus of g_Fen_1166 will be

referred to as DES18-25) affiliated with the family of *Smithellaceae*; FI3, FI4 and FI7 (hereafter,

these three MAGs will be referred to as FI3,4,7) affiliated with the genus of *Syntrophomoas_B*;

ER1, ER2 and ER4 (hereafter, these three MAGs will be referred to as ER1,2,4) that are

annotatable only at the phylum level as *Eremiobacterota*; and TM1 affiliated with the species

*Mesotoga*.

Considering that FI21,22 are among the populations that get specifically enriched with

glucose as substrate in Consortia-G (datapoint G_DNA in Figure 3), DES18-25 with propionate

as substrate in Consortia-P (datapoint P_DNA in Figure 3) and FI3,4 with butyrate as substrate

in Consortia-B (datapoint B_DNA in Figure 3), we tentatively assume that FI21, 22 are the

primary fermenting bacteria utilizing glucose, FI3,4,7 are the SBOBs degrading butyrate, and

DES18-25 are the SPOBs degrading propionate.

Function niches of FI21, 22 as the primary fermenting bacteria, FI3,4,7 as SBOBs, and

DES18-25 as SPOBs were confirmed by the transcriptome analysis. Specifically, when glucose

is provided as the primary substrate to Consortia-G (datapoints gg2 in Figure 3), or as co-

substrate to Consortia-B (datapoints bc4 in Figure 3) and Consortia-P (datapoints pc4 in Figure

3), FI21, 22 demonstrate the highest transcription activity in all the three consortia enriched,

irrespective of their very low abundance in Consortia-B and Consortia-P. Consortia-G cannot

use either butyrate or propionate efficiently, probably due to the absence of the SBOB and

SPOB populations in the community. When butyrate was provided as the substrate to

Consortia-B (datapoint bb4 in Figure 3) or to Consortia-P (datapoint pb5 in Figure 3), FI3,4,7

are noted with the highest transcription activity in both consortia. When propionate was

provided as the primary substrate to either Consortia-B (datapoint bp5 in Figure 3) or

Consortia-P (datapoint pp4 in Figure 3), consistent and relatively high transcription activity of

DES18-25 was observed in both consortia.

The other four MAGs, ER1,2,4 and TM1, did not show consistent transcriptome response

to the presence of any specific substrate tested, and they might be fermentative opportunists in

the consortia, utilizing cell debris or other extracellular substances. A more detailed

interpretation on the function niches of these four populations can be found in supporting

information.

***3.3 Genes and gene clusters mediating either Flavin based electron bifurcation (FBEB) or***

***reverse electron transfer (RET) in primary fermentation, in butyrate oxidation and in***

***propionate oxidation***

Transcription patterns of the key functional genes in bacterial MAGs of the three function

niches are summarized in Figure 4. And summarized in Table 1 are the genes/gene clusters

mediating either the flavin-based electron bifurcation (FBEB) or reverse electron transfer (RET)

that are the featuring biochemistry in primary fermentation, syntrophic butyrate oxidation and

syntrophic propionate oxidation.

The high transcription activity of the FBEB *HydABC* gene cluster in FI21,22 suggests the

involvement of electron confurcation of $Fd_{red}$ and NADH for $H_2$ generation during acetate

generation (Figure 5a) in primary fermentation of the AD bioprocess where hydrogenotrophic

methanogens keep the $H_2$ concentration low.

Butyryl-CoA dehydrogenase (*Bcd_EtfAB*) are of high transcription activity in both

FI21,22 and in FI3,4,7. In FI21,22, *Bcd_EtfA1B1* neighbors with the GroupB [FeFe]

hydrogenase, mediating the NADH-dependent FBEB biochemistry of crotonyl-CoA reduction

to butyryl-CoA, and the electrons in the $Fd_{red}$ generated will be further diverted to $H_2$ during

butyrate generation in primary fermentation (Figure 5a). In FI3,4,7, *Bcd_EtfA2B2* clusters with

the membrane-bound oxidoreductase (*EMO*), mediating the proton motive force (PMF)-

dependent RET from butyryl-CoA to membranous methylmenaquinone (MMKH2) [14]. This is

one of the featuring biochemistry in anaerobic butyrate oxidation (Figure 5b).

The *hydABC* gene cluster is also annotated and with transcription activity in both SBOBs

FI3,4,7 and in SPOBs DES18-25 (Figure 4). While comparing with the *hydABC* gene cluster

in the primary fermenting bacteria *Clostridium_T* FI21,22, mutations are noted in the *hydB*

subunit in the *hydABC* gene cluster of both SBOBs and SPOBs. As visualized in Figure 6, same

as that in *S. aciditrophicus* and in *S. wolfei*, the *hydB* subunits in *Syntrophomoas* FI3,4,7 and

in *Smithelacea* DES18-25 are all in lack of the C-terminal end 2*[4Fe-4S] function domains.

Activity of the FBEB-type *hydABC* requires the presence of both $Fd_{red}$ and NADH [15], while

biochemical characterization revealed that enzyme products of the *hydABC* gene cluster in both

*S. aciditrophicus* and *S. wolfei* cannot interact with Fdred and can use NADH as direct electron

donor for the reduction of $H^+$ [16] [17]. We therefore interpret the *hydABC* gene cluster in

*Syntrophomoas* FI3,4,7 and in *Smithelacea* DES18-25 enriched in this study are also of the

non-FBEB type and can use NADH as the direct electron donor for $H_2$ generation.

The SPOB populations, DES18-25, that get specifically enriched in this study are of a

same undefined genus in the family of *Smithelacea*.  Propionate conversion via the dismutation

pathway, i.e., two propionate molecules being first converted to acetate and butyrate, has been

reported only in *Smithella propionica* [18]. And methylmalonyl-CoA pathway is found in all other

SPOBs characterized so far [3]. As is summarized in Figure 4, both the presence and transcription

activity of genes involved in the β-oxidation of butyrate and in the methylmalonyl-CoA

pathway are observed in *Smithelacea* DES18-25.

One of the interesting transcription patterns noted in DES18-25 is of the

*FdhB_MvhD_HdrABC* gene cluster. Transcription activity and proteome signal of the

*FdhB_MvhD_HdrABC* gene cluster in SPOBs have been reported in previous studies [19, 20].

There are two hypotheses regarding the function roles of this *FdhB_MvhD_HdrABC* gene

cluster and the corresponding enzyme complex: it may either be involved in sulfate reduction

[20] or may possess an uncharacterized FBEB activity [19]. We tentatively favor the second

hypothesis for two reasons: 1) it was reported that the gram-negative bacteria *Smithella* cannot

reduce sulfate [3], further, in this study, to exclude the interference of sulfate as electron acceptor,

we replaced $NaSO_4$ with $NaHCO_3$ in the solution medium; and 2) the *FdhB_MvhD_HdrABC*

gene cluster showed high transcriptional activity almost exclusively under the presence of

propionate as substrate. We cannot exclude the possibility that this *FdhB_MvhD_HdrABC*

enzyme complex is relevant in mediating the anaerobic oxidation of succinate to fumarate,

which is thermodynamically the most challenging step in anaerobic propionate oxidation *via*

the methylmalonyl-CoA pathway [6].

Transcription patterns of the key functional genes as summarized in Figure 4 also

suggested metabolic specificity of the primary fermenting bacteria, the SBOBs and the SPOBs

enriched in this study. A lack of fermentation capacity of the SBOBs and SPOBs was indicated

by their quite low transcription activity when fermentative substances (substrate denoted as

'S3' in Figure 4) were provided as substrate. In all the three consortia enriched, when glucose

was supplied as the primary substrate or as co-substrate, regardless of their either high or low

relative abundance in the community, the two fermenting bacteria FI21, 22 were always

identified as transcriptionally most active. Specifically, VSS equivalents of FI21,22 were

estimated to be of only 13 ug-VSS and 25 ug-VSS in Consortia-B and Consortia-P, respectively

(Table S2), being two orders of magnitude lower than that of the putative SBOB populations

FI3,4,7 (9817 ug-VSS) in Consortia_B and that of the putative SPOB populations DES18-25

(18204 ug-VSS) in Consortia_P; but still, when glucose was provided as co-substrate, this

minor amount of FI21,22 dominated in both Consortia_B and Consortia_P the transcription

activity of genes involved in glycolysis and pyruvate fermentation (grid M03 in Figure 4), of

genes involved in the fatty acid synthesis (grid M04 in Figure 4), and of gene encoding the

ferredoxin-dependent [FeFe] hydrogenase during butyrate fermentation (Figure 5a) .

**3.4 Segregation of $H_2$ and formate metabolism in primary and secondary fermentation**

As is discussed in the above section, in primary fermenting bacteria FI21,22, genes encoding hydrogenase ($H_2$ase), i.e., *hydABC* and ferredoxin-dependent [FeFe] $H_2$ase, are annotated and transcribed, mediating electron transfer to $H_2$ during either acetate or butyrate fermentation, and no formate dehydrogenase (*FDH*) was annotated or transcribed. Formate in primary fermentation originates from pyruvate formate lyase (*PFL*), although the transcriptional activity of *PFL* is much lower than that of pyruvate ferredoxin oxidoreductase (*PFOR*), indicating $H_2$ is the dominant electron carrier in primary fermentation.

In contrast, uniquely transcribed in the secondary fermenting populations, i.e., SBOBs FI3,4,7 and SPOBs DES18-22, are *FDH*. The cytosolic formate dehydrogenase (cFDH) catalyze $CO_2/H^+$ reduction to HCOOH with electrons from NADH, while the membrane-bound formate dehydrogenase (mFDH) catalyze proton motive force (PMF)-dependent reverse electron transfer (RET) for $CO_2/H^+$ reduction to HCOOH with electrons from the membranous quinone (M)MKH$_2$. Both cFDH and mFDH are of high transcription activity in SBOBs and SPOBs.

Formate hydrogenlyase (*FHL*) is not annotated or transcribed in either the primary fermenting populations or the secondary fermenting populations. Bacterial *FHL* mediate the conversion of formate to $H_2$, and its absence suggests restricted biochemical interconversion of formate and $H_2$. In other words, metabolism of $H_2$ and formate as electron carriers in anaerobic digestion bioprocess are segregated.

One of the ecological relevance of such segregation of the $H_2$ and formate metabolism in AD bioprocess could be that the 'H$_2$-tolerant' primary fermentation and the 'H$_2$-sensitive'secondary fermentation can proceed simultaneously, with $H_2$ being the electron carrier in the former, and formate as the predominant electron carrier in the latter. Under high concentrations of $H_2$ emitted from primary fermentation, the 'H$_2$-sensitive' secondary fermentation can proceed with formate as dominant electron carrier.

Our wet-lab observation aligns well with the genome-centric transcriptome analysis.

Accumulation of $H_2$ in high concentrations (5,000-70,000 ppm, Figure S8) was observed only

when fermentative substances (i.e., glucose and fumarate) are present as either primary

substrate or as co-substrate. $H_2$ accumulation was never detected when only either propionate

or butyrate was supplied as the primary substrate.

If we assume that $H_2$ is the only electron carrier in the oxidation of propionate and butyrate,

at pH=7.0±0.5, (35±1) $^o$C, acetate, propionate and butyrate concentration at 10 mM, $CH_4$ and

$CO_2$ content in the gaseous phase at 70% and 30% as averaged in this study,  the acetogenic

oxidation of propionate and butyrate would become thermodynamically unfavorable when the

gaseous $H_2$ content is above 132 ppm and 134 ppm, respectively (Figure S9). However, when

substrate mix as glucose:propionate:butyrate =1:1:1 (COD:COD:COD) was provided,

intermittent $H_2$ accumulation to >30000 ppm was detected in both Consortia-B and Consortia-

P, which was far above the theoretical inhibition concentration of $H_2$ as calculated. And at the

time points when such high $H_2$ concentrations were detected, the degradation of propionate and

butyrate were not completely inhibited, as indicated by the decline of their concentrations

summarized in Figure 7.

Degradation of propionate and butyrate under high $H_2$ concentrations suggests that $H_2$ is

not the sole electron carrier and formate can play a dominant role in mediating interspecies

electron transfer during the syntrophic metabolism, which is suggested also by the high

transcription activity of *FDH* in SBOBs and SPOBs. However, we did not measure significant

accumulation of formate during either butyrate or propionate degradation. Formate

concentration below the detection limit (1 mM) is probably due to rapid interspecies formate

transfer and efficient utilization of formate by the coupling methanogens in the consortia

enriched.

**3.5 Formate- and H₂-utilizing methanogen and acetoclastic methanogen identified in the enriched consortia**

Among the 39 archaeal MAGs recovered (Figure 7), 4 archaeal MAGs are of the highest transcriptional activity, accounting for (79±9) % of the mRNA gene transcripts mapped to the 39 archaeal MAGs. These 4 archaeal MAGs are: EA7 affiliated with the family of *Methanobacteriaceae*, and HA1, HA2, HA3 (hereafter, termed as HA1,2,3) affiliated with the species of *Methanothrix soehngenii*. In the following genome-centric interrogation, we focused specifically on these 4 archaeal MAGs, to elaborate on the methanogenic activity of the enriched consortia.

According to the transcription profile as summarized in Figure 8 and Figure S10, EA7 is a formate- and H2- utilizing methanogen and HA1,2,3 are acetoclastic methanogens. *Methanobacteriaceae* EA7 dominated the transcription activity of genes encoding Formate dehydrogenase (coenzyme F420-coupled, denotated as *FdhAB*), formylmethanofuran dehydrogenase (*fwd*), and coenzyme F420 hydrogenase (*frh*). In comparison, *Methanothrix soehngenii* HA1,2,3 dominated the mRNA transcripts of genes encoding the acetyl-CoA decarbonylase (*cdhDE*) and the carbon-monoxide dehydrogenase (*cdhAB*). The acetoclastic archaeal MAGs HA1,2,3 are of high identity to strain *Methanothrix soehngenii*, while the hydrogenotrophic archaeal MAG EA7 are annotatable only at the family level of *Methanobacteriacea,* indicating novelty of this hydrogenotrophic methanogen population enriched.

Formate as an important electron carrier in the syntrophic cooperation between the SBOB/SPOB and the methanogen archaea is hinted also by the high transcription activity, in EA7, of the *fdhC* gene encoding formate transporter, and also the presence and the high transcription activity of the *FdhAB_mvhD_HdrABC* gene cluster. The formate-centered biochemistry mediated by this *FdhAB_mvhD_hdrABC* enzyme complex is reaction (1). Also

annotated in EA7 is the presence and transcription activity of the *MvhAG_MvhD_HdrABC*

gene cluster, mediating the $H_2$-centered reaction (2).

$$2HCOOH + Fd_{ox} + CoM-S-S-CoB = 2H^+ + 2CO_2 + Fd_{red} + CoM-SH + CoB-SH \quad (1)$$

$$2H_2 + Fd_{ox} + CoM-S-S-CoB = 2H^+ + Fd_{red} + CoM-SH + CoB-SH \quad (2)$$

***3.6 Metabolic specificity of the key functional populations in the AD consortia***

According to characterization of the three AD consortia enriched in this study, there is a clear

function partition among primary fermentation, syntrophic butyrate oxidation, syntrophic

propionate oxidation, hydrogenotrophic methanogenesis, and acetoclastic methanogenesis.

Judging by the featuring gene/gene clusters as illustrated in the above sections, function

partition among different populations could be clearly seen in Figure 10. In whichever of the

three consortia enriched, when butyrate was provided as the primary substrate or as a co-

substrate, (100±0) % of the *Bcd_etfAB_EMO* gene transcripts were expressed by the SBOBs.

When propionate was provided as the primary substrate or as a co-substrate, (90±5) % of the

Succinate dehydrogenase (*Sdh)* gene transcripts were expressed by the SPOBs. When glucose

was provided as the primary substrate or as a co-substrate, the primary fermenting populations

contributed (97±2)% of the transcripts of genes involved in the glycolysis &fermentation

metabolism, e.g., the *PFOR* gene.

*cFDH* and *mFDH* were exclusively annotated and transcribed in syntrophic populations

(SBOBs FI3,4,7 and SPOBs DES18-22). Coenzyme F420-dependent formate dehydrogenase

(*FdhAB*) exclusively in the hydrogenotrophic methanogen EA7. This again highlighted formate

as the electron carrier between syntrophic oxidizers and hydrogenotrophic methanogens.

The acetoclastic methanogens *Methanothrix* HA1,2,3 and the hydrogenotrophic

*Methanobacteriacea* EA7 equally dominant the overall $CH_4$ generation, as these two groups of

methanogens each contributed ~50% of the *mcrA* gene transcripts. The *Methanobacteriaceae*

EA7 dominated methanogenic conversion of formate and $H_2$ by contributing (95±2) % of the

transcripts of the functional genes, e.g., *frhA*, *fdhC* genes. The *Methanothrix soehngenii*

HA1,2,3 dominated methanogenic conversion of acetate by contributing (92±4) % of the

transcripts of functional genes, e.g., the *cdhA* gene.

**Conclusions**

Revealed in this study is the segregation of $H_2$- and formate-metabolism in anaerobic digestion

bioprocess, with $H_2$ being dominant as electron carrier in primary fermentation, and formate

being dominant in secondary fermentation. Such segregation benefit the anaerobic oxidation

of butyrate and propionate with higher tolerance to $H_2$ accumulated as metabolic intermediate

from primary fermentation.

Metabolic specificity was noted for the key functional populations in the AD consortia.

Genome-centric transcriptional characterization highlighted the characteristic genes/gene

clusters of the different functional groups. specifically, *Bcd_etfAB_H₂ase* in primary

fermenting bacteria *Clostridium_T*, *Bcd_etfAB_EMO* in the syntrophic butyrate-oxidizing

bacteria (SBOBs) in the genus of *Syntrophomonas_B*, *Sdh* and *FdhB_mvhD_hdrABC* in the

syntrophic   propionate-oxidizing   bacteria   (SPOBs)   affiliated   to   *Smithellaceae*,

*FdhC_FdhAB_mvhD_hdrABC* and *mvhABG_mvhD_hdrABC* in the one hydrogenotrophic

methanogen affiliated to *Methanobacteriaceae*, and *cdhA* in the acetoclastic methanogen

species *Methanothrix soehngenii*.

Genome-centric interpretation at the level of gene clusters can deliver higher resolution

in genotype-phenotype correlation. At the gene cluster level, genetic machineries of primary

fermenting bacteria generating butyrate and SBOB degrading butyrate can be distinguished. A

membrane-bound EMO subunit neighboring with  etfAB subunits is essential for reverse

electron transfer in SBOB.

Mutations  in  the  *hydB*  subunit  are  noted  in  the  *hydABC*  gene  cluster  of  both

*Syntrophomonas_B* SBOBs and *Smithellaceae* SPOBs enriched in this study. Such mutations

probably render H$^+$ reduction by NADH as direct electron donor during syntrophic oxidation

of butyrate and metabolism. And only the *hydABC* enzyme complex in primary fermenting

bacteria are of true FBEB activity and can interact with both NADH and Fd$_{red}$ for H$^+$ reduction.

**Data Availability Statement**

Metagenome DNA (10 datasets in total) and metatranscriptome RNA (20 data sets in total)

sequencing data are available in the NCBI Sequence Read Achieve (SRA) database under the

Bioproject accession number of PRJNA886957. The 17 bacterial MAGs and 4 archaeal MAGs

analyzed in this study are accessible in NCBI under the Bioproject accession number of

PRJNA886046, function genes annotated in these 21 MAGs and their transcription activity

could be found in Appendix file 1. 16S rRNA gene amplicon sequences (v4-v5 region) have

also been deposited in NCBI under the Bioproject accession number of PRJNA885972.

**Declaration of Competing Interest**

396 The authors declare that there is no competing interest.

**Abbreviations**

SBOBs - syntrophic butyrate oxidizing bacteria; SPOBs - syntrophic propionate oxidizing

bacteria; AD - anaerobic digestion; VFAs - volatile fatty acids; SCFAs - short-chain fatty acids;

COD - chemical oxygen demand; PCoA - principle coordinate analysis; VSS - volatile

suspended solids; MAGs - metagenome assembled genomes; FBEB – flavin based electron

bifurcation; RET – reverse electron transfer; PMF – proton motive force; PFL – pyruvate

formate lyase; PFOR – pyruvate ferredoxin oxidoreductase; FHL – formate hydrogenlyase;

EMO – membrane-bound oxidoreductase.

**Acknowledgments**

This work was supported by the Zhejiang Provincial Natural Science Foundation of China

under Grant No. LR22D010001 and Key R&D Program of Zhejiang (2022C03075) and

Research Center for Industries of the Future (RCIF). Technical assistance from Ms. Yisong Xu

in laboratory affairs and Mr. Guoqing Zhang in maintaining the in-house computing facility

are greatly appreciated.

**Authors' contributions**

YW conceptualized the research and discussed with FJ. YW conducted the wet-lab experiment,

the meta-omics data analysis, and wrote the manuscript. RZ assisted with the wet-lab

experiment. CW, WY, TZ and FJ contributed in the interpretation of the results and revision of

the manuscript.

**Table1.** Presence (+) and absence(-) of the gene clusters mediating either flavin based electron bifurcation (FBEB) or reverse electron transfer
(RET) in primary fermentation and in syntrophic oxidation of butyrate and propionate, their cellular component and the corresponding
biochemistry**.**

| Gene/ Gene clusters | Cellular component | Biochemistry | Primary fermenting bacteria | SBOB | SPOB |
|---|---|---|---|---|---|
| | | | FI21,22 | FI3,4,7 | DES18-25 |
| hydA_hydB_hydC | Cytoplasm | $Fd_{red} + NADH + 4H^+ = 2H_2 + NAD^+ + Fd_{ox}$ (in primary fermentation); OR $NADH + 2H^+ = H_2 + NAD^+$ (in syntrophic oxidation) | + | + | + |
| Bcd_EtfA_EtfB_H$_2$ase | Cytoplasm | $2NADH + Crotonyl\text{-}CoA + Fd_{ox} = 2\ NAD^+ + Butyryl\text{-}CoA + Fd_{red}$ | + | - | - |
| cFDH | Cytoplasm | $NADH + CO_2 + H^+ = NAD^+ + HCOOH$ | - | + | + |
| mFDH | Membrane-bound | $(M)MKH_2 + CO_2 + PMF = HCOOH + (M)MK$ | - | + | + |
| FdhB_mvhD_HdrABC | Cytoplasm | UNKNOWN | - | - | + |
| Bcd_EtfB_EMO_EtfA | Membrane-bound | $Butyryl\text{-}CoA + MMK + PMF = Crotonyl\text{-}CoA + MMKH_2$ | - | + | - |
| SdhA_SdhB_SdhC | Membrane-bound | $Succinate + (M)MK + PMF = Fumarate + (M)MKH_2$ | - | - | + |

M06 are hydrogenase genes, in M07 are genes encoding the the electron transfer flavoprotein,
in M08 is the ATP synthase gene, in M09 are genes encoding the type IV pili, flagellar and
fimbria assembly protein, in M10 are genes encoding the protease, murein-degrading enzymes
and amylase etc., in M11 are genes encoding the synthesis of protein, amino acids, and
glycogen. Full names of genes denoted with abbreviations could be found in the Appendix file
1.

**Figure 5.** a) Acetate and butyrate metabolism in primary fermenting bacteria *Clostridium*
FI21,22 and the corresponding gene clusters encoding enzymes mediating the flavin-based
electron bifurcation; b) proton motive force (PMF) dependent butyryl-CoA oxidation to
crotonyl-CoA, the biochemistry and the corresponding gene clusters.

**Figure 6.** Summary of the function domains in the *hydB* subunit in the flavin based electron
bifurcation (FBEB) type of *hydABC* in the primary fermenting populations (upper box) and in
the non-FBEB type *hydABC* in the secondary fermenting populations. Name of the reference
microbe strains in which the catalytic activity of the enzyme complex has been characterized
are highlighted in blue. Names of the anaerobe populations enriched in study are shown in
grey. Provided in the brackets are the gene accession number of the corresponding *hydB* gene.

**Figure 7.** Degradation profile of propionate (HPr) and butyrate (HBut), and dynamics of $H_2$
concentration when glucose:propionate:butyrate = 1:1:1 (COD:COD:COD) was provided as substrate
in a) Consortia-P, b) Consortia-B, and c) Consortia-G. d) Gibbs free energy change calculated for the
degradation of propionate, n-butyrate and iso-butyrate in Consortia-P and Consortia-B , respectively,
assuming that $H_2$ is the only electron carrier generated from the syntrophic oxidation of butyrate or
propionate.

**Figure 8.** Maximum likelihood tree of the 39 archaeal MAGs recovered in this study. The inner
heatmap summarized the meta-RNA based transcription activity of each MAG. A detailed
information on each meta-RNA datapoint could be found in Table S1. The tree was constructed
using FastTree (maximum-likelihood phylogeny) [21] based on a set of 122 concatenated
universal single-copy proteins for *Archaea* [22]. Branches of the phylogenetic tree are coloured
according to their taxonomy affiliation, and those light purple dots indicate bootstrap values
>=0.9.

**Figure 9.** Transcription activity of the key functional genes annotated in the
Methanobacteriaceae MAG EA7, and in the three *Methanothrix* MAGs HA1,2,3. The heatmap
represents the transcription activity of the corresponding genes, and the data given alongside
the heatmap bar are square root of the RNA-RPKM values. Full names of genes denoted with
abbreviations could be found in the Appendix file 1. Genes grouped in grid R1 are those
specifically involved in hydrogenotrophic methanogenesis, genes grouped in grid R2.1-R2.3
are those specifically involved in acetoclastic methanogenesis, and genes grouped in grid R3
are genes shared by both the hydrogenotrophic methanogenesis and acetoclastic
methanogenesis

**Figure 10.** Contribution of the putative fermenter bacteria (*Clostridium_T* FI21,22), the
putative SBOBs (*Syntrophomonadaceae* FI3,4,7), the putative SPOBs (*Smithellaceae*, DES18-
25) to the mRNA transcripts of the core genes (left, coloured by genes) involved in the butyrate
metabolism (*EtfAB_EMO*), the propionate metabolism (*Sdh*, Succinate dehydrogenase) and the
pyruvate fermentation (*PFOR,* Pyruvate-ferredoxin oxidoreductase), under the stimuli of
butyrate, propionate and glucose, respectively. And also, when there is no $H_2$ accumulation,
contribution of *Methanobacteriaceae* EA7 and *Methanothrix_soehngenii* HA1,2,3 to the
mRNA transcripts of the core genes involved in the methanogenesis metabolism, i.e., *fdhA*:
formate dehydrogenase (coenzyme F420) alpha subunit, *fdhC*: formate transporter, *cdhA*:
acetyl-CoA decarbonylase, and *mcrA*: methyl coenzyme M reductase. Colours represent
different genes, and the width of each ribbon represents the contribution ratios of each microbe
group to the mRNA transcripts of each gene, the ratios were averaged across the stimuli results

in the three enriched consortia: Consortia-G, Consortia-B and Consortia-P. cFDH and mFDH

refer to the cytosolic and membrane-bound formate dehydrogenase, respectively.

## References

(1) Ariesyady, H. D.; Ito, T.; Okabe, S. Functional bacterial and archaeal community structures of major trophic groups in a full-scale anaerobic sludge digester. *Water Res* **2007**, *41* (7), 1554-1568. DOI: 10.1016/j.watres.2006.12.036. Jain, M. Anaerobic Membrane Bioreactor as Highly Efficient and Reliable Technology for Wastewater Treatment—A Review. *Advances in Chemical Engineering and Science* **2018**, *08* (02), 82-100. DOI: 10.4236/aces.2018.82006. Speece, R. E. Anaerobic biotechnology for industrial wastewater treatment. *ES&T Features* **1983**, *17* (9), 416-417. DOI: 10.1021/es00115a001,. Macarie, H. Overview of the application of anaerobic treatment to chemical and petrochemical wastewaters. *Water Science and Technology* **2000**, *42* (5-6), 201-214. DOI: 10.2166/wst.2000.0515.

(2) Bryant, R. I. M. a. M. P. Metabolic activity of fatty acid-oxidizing bacteria and the contribution of acetate, propionate, butyrate and CO2 to methanogenesis in cattle waste at 40 and 60 ºC. *Applied and Environmental Microbiology* **1981**, *41* (6), 1363-1373.

(3) Stams, A. J.; Sousa, D. Z.; Kleerebezem, R.; Plugge, C. M. Role of syntrophic microbial communities in high-rate methanogenic bioreactors. *Water Sci Technol* **2012**, *66* (2), 352-362. DOI: 10.2166/wst.2012.192 From NLM Medline.

(4) Moeller, L.; Zehnsdorf, A. Process upsets in a full-scale anaerobic digestion bioreactor: over-acidification and foam formation during biogas production. *Energy, Sustainability and Society* **2016**, *6* (1). DOI: 10.1186/s13705-016-0095-7.

(5) C.Adinarayana Reddy, M. P. B., M.J.Wolin. Characteristics of S Organism Isolated from Methanobacillus omelianskii. *Journal of Bacteriology* **1972**, *109* (2), 539-545. McInerney, M. J.; Struchtemeyer, C. G.; Sieber, J.; Mouttaki, H.; Stams, A. J.; Schink, B.; Rohlin, L.; Gunsalus, R. P. Physiology, ecology, phylogeny, and genomics of microorganisms capable of syntrophic metabolism. *Ann N Y Acad Sci* **2008**, *1125*, 58-72. DOI: 10.1196/annals.1419.005 From NLM Medline. Michael J. McInerney, M. P. B., and Norbert Pfenning. Anaerobic Bacterium that Degrades Fatty Acids in Syntrophic Association with Methanogens. *Archives of Microbiology* **1979**, *122*, 129-135. Narihiro, T.; Nobu, M. K.; Kim, N. K.; Kamagata, Y.; Liu, W. T. The nexus of syntrophy-associated microbiota in anaerobic digestion revealed by long-term enrichment and community survey. *Environ Microbiol* **2015**, *17* (5), 1707-1720. DOI: 10.1111/1462-2920.12616. Nobu, M. K.; Narihiro, T.; Rinke, C.; Kamagata, Y.; Tringe, S. G.; Woyke, T.; Liu, W. T. Microbial dark matter ecogenomics reveals complex synergistic networks in a methanogenic bioreactor. *ISME J* **2015**, *9* (8), 1710-1722. DOI: 10.1038/ismej.2014.256. Rudolf K. Thauer, K. J., Karl Decker. Energy Conservation in Chemotrophic Anaerobic Bacteria. *Bacteriological Reviews* **1977**, *41* (1), 100-180. Schink, B. Energetics of Syntrophic Cooperation in Methanogenic Degradation.

*Microbiology and Molecular Biology Reviews* **1997**, *61* (2), 262-280.

(6) Westerholm, M.; Calusinska, M.; Dolfing, J. Syntrophic propionate-oxidizing bacteria in methanogenic systems. *FEMS Microbiol Rev* **2022**, *46* (2). DOI: 10.1093/femsre/fuab057  From NLM Medline.

(7) Nobu, M. K.; Narihiro, T.; Mei, R.; Kamagata, Y.; Lee, P. K. H.; Lee, P. H.; McInerney, M. J.; Liu, W. T. Catabolism and interactions of uncultured organisms shaped by eco-thermodynamics in methanogenic bioprocesses. *Microbiome* **2020**, *8* (1), 111. DOI: 10.1186/s40168-020-00885-y.

(8) de Kok, S.; Meijer, J.; van Loosdrecht, M. C.; Kleerebezem, R. Impact of dissolved hydrogen partial pressure on mixed culture fermentations. *Appl Microbiol Biotechnol* **2013**, *97* (6), 2617-2625. DOI: 10.1007/s00253-012-4400-x  From NLM Medline.

(9) Birgitte K. Ahring, P. W. Kinetics of Butyrate, Acetate, and Hydrogen Metabolism in a Thermophilic, Anaerobic, Butyrate-Degrading Triculture. *Appl Environ Microbiol* **1987**, *53*, 434-439.

(10) R. Conrad, B. S., and T.J. Phelps. Thermodynamics of $H_2$-consuming and H2-producing metabolic reactions in diverse methanogenic environments under in situ conditions. *FEMS Microbiol Eco* **1986**, *38*, 353-360.

(11) Sieber, J. R.; McInerney, M. J.; Gunsalus, R. P. Genomic insights into syntrophy: the paradigm for anaerobic metabolic cooperation. *Annu Rev Microbiol* **2012**, *66*, 429-452. DOI: 10.1146/annurev-micro-090110-102844  From NLM Medline.

(12) de Bok, F. A.; Luijten, M. L.; Stams, A. J. Biochemical evidence for formate transfer in syntrophic propionate-oxidizing cocultures of Syntrophobacter fumaroxidans and Methanospirillum hungatei. *Appl Environ Microbiol* **2002**, *68* (9), 4247-4252. DOI: 10.1128/AEM.68.9.4247-4252.2002  From NLM Medline. Leslie A. Day, E. L. K., Dallas R. Fonseca, Kyle C. Costa. Interspecies Formate Exchange Drives Syntrophic Growth of Syntrophotalea carbinolica and Methanococcus maripaludis. *Appl Environ Microbiol* **2022**, *88*.

(13) Worm, P.; Koehorst, J. J.; Visser, M.; Sedano-Nunez, V. T.; Schaap, P. J.; Plugge, C. M.; Sousa, D. Z.; Stams, A. J. M. A genomic view on syntrophic versus non-syntrophic lifestyle in anaerobic fatty acid degrading communities. *Biochim Biophys Acta* **2014**, *1837* (12), 2004-2016. DOI: 10.1016/j.bbabio.2014.06.005  From NLM Medline.

(14) Agne, M.; Estelmann, S.; Seelmann, C. S.; Kung, J.; Wilkens, D.; Koch, H. G.; van der Does, C.; Albers, S. V.; von Ballmoos, C.; Simon, J.; et al. The missing enzymatic link in syntrophic methane formation from fatty acids. *Proc Natl Acad Sci U S A* **2021**, *118* (40). DOI: 10.1073/pnas.2111682118  From NLM Medline.

(15) Buckel, W.; Thauer, R. K. Flavin-Based Electron Bifurcation, A New Mechanism of Biological Energy Coupling. *Chem Rev* **2018**, *118* (7), 3862-3886. DOI:

10.1021/acs.chemrev.7b00707  From NLM Medline.

(16) Losey, N. A.; Mus, F.; Peters, J. W.; Le, H. M.; McInerney, M. J. Syntrophomonas

wolfei Uses an NADH-Dependent, Ferredoxin-Independent [FeFe]-Hydrogenase To

Reoxidize NADH. *Appl Environ Microbiol* **2017**, *83* (20). DOI: 10.1128/AEM.01335-

17  From NLM Medline.

(17) Losey, N. A.; Poudel, S.; Boyd, E. S.; McInerney, M. J. The Beta Subunit of Non-

bifurcating NADH-Dependent [FeFe]-Hydrogenases Differs From Those of

Multimeric Electron-Bifurcating [FeFe]-Hydrogenases. *Front Microbiol* **2020**, *11*, 1109.

DOI: 10.3389/fmicb.2020.01109  From NLM PubMed-not-MEDLINE.

(18) Yitai Liu, D. L. B., Henry C. Aldrich, Gwendolyn R. Drake, David R. Boone.

Characterization of the anaerobic propionate-degrading syntrophs Smithella propionica

gen. nov., sp. nov. and Syntrophobacter wolinii. *International Journal of Systematic*

*Bacteriology* **1999**, *49*, 545-556. de Bok, F. A.; Stams, A. J.; Dijkema, C.; Boone, D. R.

Pathway of propionate oxidation by a syntrophic culture of Smithella propionica and

Methanospirillum hungatei. *Appl Environ Microbiol* **2001**, *67* (4), 1800-1804. DOI:

10.1128/AEM.67.4.1800-1804.2001  From NLM Medline.

(19) Sedano-Nunez, V. T.; Boeren, S.; Stams, A. J. M.; Plugge, C. M. Comparative

proteome analysis of propionate degradation by Syntrophobacter fumaroxidans in pure

culture and in coculture with methanogens. *Environ Microbiol* **2018**, *20* (5), 1842-1856.

DOI: 10.1111/1462-2920.14119  From NLM Medline.

(20) Hao, L.; Michaelsen, T. Y.; Singleton, C. M.; Dottorini, G.; Kirkegaard, R. H.;

Albertsen, M.; Nielsen, P. H.; Dueholm, M. S. Novel syntrophic bacteria in full-scale

anaerobic digesters revealed by genome-centric metatranscriptomics. *ISME J* **2020**, *14*

(4), 906-918. DOI: 10.1038/s41396-019-0571-0  From NLM Medline.

(21) Price, M. N.; Dehal, P. S.; Arkin, A. P. FastTree: computing large minimum

evolution trees with profiles instead of a distance matrix. *Mol Biol Evol* **2009**, *26* (7),

1641-1650. DOI: 10.1093/molbev/msp077.

(22) Chaumeil, P. A.; Mussig, A. J.; Hugenholtz, P.; Parks, D. H. GTDB-Tk: a toolkit

to classify genomes with the Genome Taxonomy Database. *Bioinformatics* **2019**. DOI:

10.1093/bioinformatics/btz848.

**a**

Glucose

$2NAD^+$ $\rightarrow$ $2ATP$
$2NADH$

*Clostridium*
**FI 21,22**

*Group A3 Electron confurcating [FeFe]H₂ase hydABC*

$4H^+ + Fd_{red} + NADH = 2H_2 + NAD^+ + Fd_{ox}$

2 Pyruvate

*hydA* → *hydB* → *hydC*

*PFL*     *PFOR*     *PFOR*

$2H^+$   $Fd_{ox}$   $Fd_{ox}$   $NAD^+$

$H_2$   $Fd_{red}$   $Fd_{red}$   *Rnf*   $NADH$

$2NADH$
$2NAD^+$

$2Fd_{ox}$

$2Fd_{red}$   *hydABC*   $4H_2$

$8H^+$

2Acetyl-CoA + 2CO₂    PMF=0.3ATP    2Acetyl-CoA + 2CO₂

$\rightarrow 2ATP$

2Acetate

Acetoacetyl-CoA   $NADH$ / $NAD^+$

Hydroxybutyryl-CoA

2Acetyl-CoA + **2HCOOH**

$\rightarrow 2ATP$

2Acetate

Crotonyl-CoA

$H_2O$

$2NADH$   $Fd_{ox}$   $2H^+$

*EtfAB_Bcd*   *H₂ase_2*   $H_2$

$2NAD^+$   $Fd_{red}$

Butyrate-CoA    Crotonyl-CoA + NADH + Fd_{ox} = Butyrate-CoA + Fd_{red} + 2 NAD^+

$\rightarrow ATP$     Fd_{red} + 2 H^+ = Fd_{ox} + H_2

Butyrate

*Electron bifurcating Bcd_EtfAB gene cluster*

Butyryl-CoA dehydrogenase → EtfB1 → EtfA1 → 3-hydroxybutyryl-CoA dehydrogenase ← ← → Ferredoxin-NADP reductase subunit alpha → 4Fe-4S ferredoxin-type oxidoreductase → *GroupB [FeFe]H₂ase* Ferredoxin-dependent hydrogenases →

**b**

*Syntrophomoas*
**FI 3,4,7**

Butyryl-CoA + MMK + PMF = Crotonyl-CoA + MMKH₂

MMKH₂ + CO₂ + H⁺ + PMF = HCOO⁻ + MMK

$HCOO^-$   $CO_2 + H^+$

$1 Na^+/H^+$    *mFDH*    $1 Na^+/H^+$

*BcdEfAB_EMO*   $MMKH_2$

$MMK$

Butyryl-CoA

Crotonyl-CoA

Butyryl-CoA dehydrogenase →   Butyryl-CoA: acetate-CoA transferase → 3-hydroxybutyryl-CoA dehydrogenase →

*Bcd gene cluster*    Acetyl-CoA acetyltransferase →    Crotonyl-CoA hydratase

*fdnG selenocysteine-containing* → → *fdnH (Fe-S ferredoxin type)* → *fdnI* → *fdhE* →

Molybdopterin molybdenumtransferase    *Fdh-N type formate dehydrogenase*

EtfB2 → EtfA2 → EMO →

*EtfAB_EMO*

**In FBEB-type *hydABC***     $4H^+ + Fd_{red} + NADH = 2H_2 + NAD^+ + H^+ + Fd_{ox}$

*Thioredoxin-like [2Fe-2S] ferredoxin*    *FMN-binding domain*    *SLBB*    *2Fe-2S binding region*    *2 * [4Fe-4S] binding domain*

*T. maritima* [WP_004082447.1]

*FI21,22* [AFKLIHEH_01411]

**In *hyd4BC* gene cluster of SBOB and SPOB**     $2H^+ + NADH = H_2 + NAD^+ + H^+$

*S. aciditrophicus* [WP_011417005.1]

*S. wolfei* [WP_011640437.1]

*FI3,4,7 [KHOEJDGE_01517]*

*DES18-25[BLLFMGGA_00367]*

*Thioredoxin-like [2Fe-2S] ferredoxin*    *FMN-binding domain*    *SLBB*    *2Fe-2S binding region*    *FAD binding domain*

| Consortia-P fed with G:B:P=1:1:1 | | | | | | Gibbs free energy change[1] for the acetogenic degradation of | | |
|---|---|---|---|---|---|---|---|---|
| Time (h) | $H_2$ (ppm) | Hpr (mg/l) | n-Hbut (mg/l) | HAc (mg/l) | iso-Hbut (mg/l) | Hpr | n-Hbut | iso-Hbut |
| 2 | 20 | 312±2 | 269±2 | 17±1 | 9±0.1 | -23 | -38 | -29 |
| 4 | 5114 | 316±2 | 273±2 | 55±0.2 | 18±0.1 | 23 | 50 | 57 |
| 6 | 19903 | 289±12 | 281±7 | 89±2 | 20±5 | 35 | 73 | 80 |
| 10 | 34937 | 264±5 | 257±5 | 133±5 | 24±1 | 40 | 81 | 87 |
| 24 | 174 | 217±5 | 153±2 | 255±8 | 40±1 | 2 | 4 | 7 |
| 32 | 116 | 164±1 | 91±1 | 248±1 | 38±0.2 | -1 | -2 | 1 |
| 49 | 51 | 120±1 | 39±0.3 | 199±2 | 34±0.4 | -7 | -13 | -13 |

| Consortia-B fed with G:B:P=1:1:1 | | | | | | Gibbs free energy change for the acetogenic degradation of | | |
|---|---|---|---|---|---|---|---|---|
| Time (h) | $H_2$ (ppm) | Hpr (mg/l) | n-Hbut (mg/l) | HAc (mg/l) | iso-Hbut (mg/l) | Hpr | n-Hbut | iso-Hbut |
| 2 | 353 | 264±6 | 399±10 | 29±0.6 | 1±0.03 | 7 | 16 | 16 |
| 4 | 18747 | 275±4 | 371±4 | 70±1 | 34±0.1 | 71 | 79 | 79 |
| 6 | 50975 | 264±6 | 416±9 | 95±2 | 43±0.3 | 87 | 96 | 96 |
| 10 | 30000 | 258±1 | 386±4 | 101±1 | 21±0.1 | 42 | 50 | 50 |
| 24 | 103 | 232±5 | 203±1 | 174±5 | 0 | -7 | -3 | -3 |
| 32 | 107 | 216±0.4 | 109±1 | 182±2 | 0 | -5 | -3 | -3 |
| 49 | 51 | 172±2 | 43±4 | 91±1 | 0 | -16 | -16 | -16 |