Formation of brain-wide neural geometry during visual item recognition in monkeys

He Chen[†1,2], Jun Kunimatsu[†3,4,5], Tomomichi Oya[6,7], Yuri Imaizumi[8], Yukiko Hori[9], Masayuki Matsumoto[3,4], Yasuhiro Tsubo[10], Okihide Hikosaka[5], Takafumi Minamimoto[9], Yuji Naya[1,11,12], Hiroshi Yamada*[34]

Abbreviated title: Brain-wide geometric change in neural space

[1]School of Psychological and Cognitive Sciences, Peking University, No. 52, Haidian Road, Haidian District, Beijing 100805, China

[2]Department of Biological Structure and Washington National Primate Research Center, University of Washington, Seattle, WA 98195, USA

[3]Division of Biomedical Science, Institute of Medicine, University of Tsukuba, 1-1-1 Tenno-dai, Tsukuba, Ibaraki 305-8577, Japan

[4]Transborder Medical Research Center, University of Tsukuba, 1-1-1 Tenno-dai, Tsukuba, Ibaraki 305-8577, Japan

[5]Laboratory of Sensorimotor Research, National Eye Institute, National Institutes of Health, Bethesda, MD 20892, USA

[6]The Brain and Mind Institute, University of Western Ontario, London, N6A 3K7, Canada

[7]Department of Physiology and Pharmacology, University of Western Ontario, N6A 3K7, London, Canada

[8]Medical Sciences, University of Tsukuba, 1-1-1 Tenno-dai, Tsukuba, Ibaraki 305-8577, Japan

[9]Advanced Neuroimaging Center, National Institutes for Quantum Science and Technology, 4-9-1 Anagawa, Inage-ku, Chiba 263-8555 Japan

[10]College of Information Science and Engineering, Ritsumeikan University, 1-1-1 Noji-

Higashi, Kusatsu, Shiga, 525-8577, Japan

[11]IDG/McGovern Institute for Brain Research at Peking University, No. 52, Haidian Road,

Haidian District, Beijing 100805, China

[12]Beijing Key Laboratory of Behavior and Mental Health, Peking University, No. 52,

Haidian Road, Haidian District, Beijing 100805, China

†These authors contributed equally to this work.

*Correspondence to: Hiroshi Yamada, Ph.D.

Division of Biomedical Science, Institute of Medicine, University of Tsukuba

1-1-1 Tennodai, Tsukuba, Ibaraki, 305-8577 Japan.

Tel/Fax: 81-29-853-6013; e-mail: h-yamada@md.tsukuba.ac.jp

Acknowledgments

The authors thank Takashi Kawai, Ryo Tajiri, Yoshiko Yabana, Yuki Suwa, and Shiho

Nishino for their technical assistance. Monkeys FU was provided by NBRP "Japanese

Monkeys" through the National Bio Resource Project of the MEXT, Japan. Funding: This

research was supported by JSPS KAKENHI (Grant Numbers JP:15H05374, 22H04832),

JST Moonshot R&D JPMJMS2294 (H.Y.), and the National Natural Science Foundation

of China (Grant 31871139) (Y.N.).

Conflict of interest: The authors declare no competing of interests.

Keywords: brain, neural population dynamics, monkey, vision

Author Contributions H.Y. conceptualized the study. H.Y., Y.N., T.M., O.H., and J.K.

designed the experiments. H.Y., H.C., J.K., Y.I., and T.M. performed the experiments. H.Y.

and Y.T. developed the analytical tools. H.Y., H.C., J.K., and Y.H. analyzed the data. H.Y.,

H.C., and J.K. wrote the manuscript. All authors edited and approved the final version of

the manuscript.

Data availability: All data and analysis codes used in this study are available in the

supporting files.

Exp.1: Analysis code: Analy_mainchen.r and AnalyCue_PCAchen.r

Data: ChenMatData_All1506_inter_ranked.csv for 50 ms time bin

Chen_CellListAll1506.csv for cell type list

Exp. 2: Analysis code, Analy_mainkuni.r and AnalyCue_PCAkuni.r

Data: KuniMatData_All116_nointer_ranked.csv for 50 ms time bin

Exp. 3: Analysis code, Analy_mainhori.r and AnalyCue_PCAhori.r

Data: HoriMatData_All150.csv for 50ms time bin

Exp. 4: Analysis code, Analy_mainyamadai.r and AnalyCue_PCAyamada.r

Data: MatDataPCAProMag_50ms2021Sep09.txt for 50 ms data

CellList_DSVSOFCmOFC2019_0116.csv for cell type list

Common code for all experiments (1 to 4):

ParaData_BS1k50ms3ave.csv: Bootstrap replicate Data for geometric features.

ParaData_BS1k50ms3aveLissague2final.csv: bootstrap replicate Data for

Lissajous curve.

Analysis code for PCA: Analy_pca1.0final.r.

Abstract (150/150)

Neural dynamics reflect canonical computations that relay and transform information in the brain. Previous studies have identified the neural population dynamics in many individual brain regions as a trajectory geometry, a manifest of some computational motifs. However, whether these populations share particular geometric patterns across brain-wide neural populations remains unclear. Here, by mapping neural dynamics widely across temporal/frontal/limbic regions in the cortical and subcortical structures of monkeys, we show that 10 neural populations, including 2,500 neurons, propagate visual item information in a stochastic manner. We found that the visual inputs predominantly evoked rotational dynamics in the higher-order visual area, the TE and its downstream striatum tail, while curvy/straight dynamics appeared more frequently downstream in the orbitofrontal/hippocampal network. These geometric changes were not deterministic but rather stochastic according to their respective emergence rates. These results indicated that visual information propagates as a heterogeneous mixture of stochastic neural population signals in the brain.

Introduction

Visual inputs activate a large number of neurons in the brain that construct numerous neural networks to process information in an environment (*1-3*). This brain-wide activity change reflects the information processing embedded in each individual neural circuit; however, limitations of spatial and temporal resolution in the measurements of circuitry activity disrupt our understanding of brain-wide visual information processing (*4-9*). Under this limitation, considerable attempts have been made toward understanding how the brain processes information using a variety of developing theoretical frameworks (*10-15*).

One of the analytic frameworks developed within the last decade is state-space analysis (*16*) that provides a mechanistic structure of information processed in the lower-dimensional space of a neural population (*17-19*). This analytical tool identified dynamic neural population structures that reflect information processing for general biological features (*20, 21*) and allowed us to describe those features as a neural geometry in a fine time resolution (*13-15*) in the sub-second order. A large number of unidentifiable neural circuits may process information moment-by-moment (*6*), and they may form a population geometry, such as rotational (*18*), curvy (*19, 22*), or straight geometries (*23*), as the typical and basic features of dynamics. A recent finding suggests that the combination of neural population geometries may be the key to processing information to transform sensory inputs into memory (*24*). Recent studies have extended these analytical frameworks (*25-29*); however, because of poor comparisons among brain-wide neural populations, how the brain processes information in the form of geometry remains elusive.

To examine how brain-wide neural dynamics are formed to process visual information, we accumulated the neural population data of monkeys from four laboratories that contained 10 neural populations, including 2,500 neurons across temporal/frontal/limbic networks. We applied targeted dimensional reduction, with a bootstrap resampling

technique that detects and replicates neural modulation dynamics in a low-dimensional

neural space. Following a parametric bootstrap analysis using the Lissajous curve function,

our cross-study comparison revealed that a gradual shift in neural population geometries

occurred throughout the temporal-to-frontal regions in a stochastic manner.

Results

We compared the trajectory geometries across many neural populations widely distributed

in the brain from the output brain regions in the ventral visual pathway (*30-33*) and its

downstream brain regions that may access memories associated with a visual stimulus.

These included ten brain regions that were accumulated from nine monkeys examined in

the four laboratories, from the higher-order visual area TE and their downstream brain

regions in cortical, subcortical, and limbic structures, such as the temporal/orbitofrontal

cortices, striatum, and hippocampus (HPC) (Fig. 1A, No. 1 to 10). A total of 2,500 neurons

accumulated across the four behavioral tasks (Fig. S1), in which visual items provided

monkeys with position and/or reward information during the active (Exps. 1 and 3), and

passive (Exps. 2 and 4) behavioral responses. Using state-space analysis, we

characterized the types of neural population geometry that appeared in the lower

dimensional neural space, which describes how neural modulation by the task parameters

of interest processes information at the population level (*23, 29*).

We found all three types of geometric patterns, rotational, curvy, and straight

geometries, in the top two dimensions (Fig. 1B-D, see Fig. S2 for performance as the

percentage of variance explained), including unclear structures based on visual inspection

(Fig. 1E). All 10 neural populations showed a significant structure at the principal

component (PC)1-2 plane based on shuffle controls (Fig. S3, $P < 0.05$ for all PC1 and

PC2). PC3 expression was not significantly different in some neuronal populations (Fig.

S3, Exps. 2 and 4; compare the black and gray dots for each PC). These identified

geometric structures appeared to be distributed from complex to simple, reflecting the

circuit distance to the visual input (Fig. 1). For example, the TE (Fig. 1B, No. 1, A11 plane

in A) and its downstream region striatum tail (STRt) (Fig. 1B, No. 2, A11 plane in A),

showed rotational geometries during visual item recognition. In more detail, at the

beginning of information processing after the visual item presentation (Fig. 1B left, see

green **s** at time = 0), the neural state was positioned at around the center of the PC1-2

plane, and then rotated at approximately 0.2 s and started to move the second quadrant

with the counterclockwise rotation going back close to the initial point (see green **e** 0.6 s

after visual item onset). An opposite rotation was observed for the worst item, with a

smaller change (orange). These rotational structures were also observed at the STRt on

a similar timescale (Fig. 1B, right).

In the downstream brain regions, such as the perirhinal cortex (PRC) and caudate

body (CDb), rotational or curvy dynamics were observed (Fig. 1C, No. 3, PRC, A11 plane

in A and No. 4, CDb, A23 plane in A), which were characterized by a half rotation ending

at the opposite neural space and end points deviating from the initial point, although it was

unclear whether they showed rotational or curvy dynamics. In contrast, straight dynamics

were observed in brain regions far away from the visual inputs in the HPC (Fig. 1A, No. 5

in the A11 plane, Fig. 1D) and central part of the orbitofrontal cortex (cOFC) (Fig. 1A, No.

7 in the A32 plane, Fig. 1D). In addition, the ventral striatum (VS) showed straight

dynamics (Fig. 1A, No. 6 in the A23 plane, Fig. 1D), although some structures could not

be clearly determined (Fig. 1E, parahippocampal cortex, PHC, medial orbitofrontal cortex,

mOFC). The straight dynamics also showed a geometric change back and forth along the

straight trajectory (Fig. 1D). These qualitative observations based on visual inspection

suggest that neural population structures may change dynamically through visual

processing, and the shift of neural population geometries might occur throughout the

cortical and sub-cortical structures across the temporal/frontal/limbic network.

Evaluation of geometric patterns based on their selected features

To quantify these geometric patterns occurring at approximately half a second, we

estimated indices for the characteristics of dynamic neural changes in the low-dimensional

neural state (Fig. 2A). They were as follows: accumulated angle difference weighted by

deviance, $\Sigma d\,\theta$, which reflects a measure similar to the centrifugal force (see Materials

and Methods for details, Fig. 2A, top); mean distance of vectors (d, Fig. 2A, top); rotational

speed ($\Theta/0.1$ s, Fig. 2A, bottom); and distance between start and end of trajectory (d_{S-E},

Fig. 2A, bottom). Following the replication of the neural population geometries based on

the bootstrap resampling technique (see Materials and Methods), we calculated these

parameter values for each replicated neural population.

We found that across the 10 neural populations, these indices captured geometric

features to some extent, similar to the rotational geometries observed in the TE and STRt

(Fig. 2B-E). For instance, identified clusters based on the dendrogram and principal

component analysis (PCA) (Fig. 2B-C) showed that they possess a rotational

characteristic with high rotational speed (Fig. 2D, right, red), large $\Sigma d\,\theta$ (Fig. 2D, middle,

red), large d (Fig. 2D, middle, red), and small d_{S-E} (Fig. 2D, left, red), which occupy more

than 90% of the STRt population in the best item condition (Fig. 2E, see also Fig. 1B, right

green trajectory). A smaller rotational structure characterized by smaller values of $\Sigma d\,\theta$

was also captured by another cluster (Fig. 2D and E, shallow red), which occupied

approximately 90% of the STRt population in the worst item condition (Fig. 2E, see also

Fig. 1B right, orange trajectory). These rotational features were observed in other temporal

brain regions (Fig. 2E, see reddish, more than 50% in TE, 20-40% in PRC and PHC), but

merely observed at the frontal/limbic brain regions, such as HPC (less than 10% in all

remaining brain regions). In contrast, curvy/straight dynamics were observed in other

clusters in the downstream brain regions (Fig. 2E, green and blue).

Collectively, in each of the 10 brain regions, a cluster with high rotational speed

occupied the STRt and half of the TE populations (Fig. 2D, reddish), while the

curvy/straight dynamics (Fig. 2D, blue and green) occupied most of the replicates in the

remaining cortical and subcortical regions (Fig. 2E, blue and green).

Parameterization of geometric patterns using Lissajous curve function

To parameterize these geometric features in more detail, we fit the Lissajous curve

function (*34*) to the replicated data, which can mathematically capture all rotational, curvy,

and straight dynamics simultaneously. In the Lissajous function, any two-dimensional

geometric features represented by *F(x, y)* are captured using the following equations:

$x = Ax \cos (\omega x \, t(i) + \Phi x) + bx$ (1)

$y = Ay \cos (\omega y \, t(i) + \Phi y) + by$ (2)

where ω and Φ represent cycle of the rotation and their deviance as a function of time, t(i)

(Fig. 3A). For the horizontal and vertical axes, *Ax* and *Ay* define the amplitude of the

trajectory, respectively, whereas *bx* and *by* define its location. t(i) takes the values from 0

to 0.6 s in all the four experiments, and thus, one cycle of the trajectory is represented as

0 to 3.33 π. In this function, rotational dynamics is represented by the same ω among x

and y formulas, and half cycle differences in Φ values between *x* and *y* formulas (Fig. 3A,

left, see also Fig. S4A for more details). We note that this rotational example represent

less than one cycle due to $\omega x = 3.0$. In contrast, straight dynamics are represented with

the same ω and also same Φ values between the two formulas (Fig. 3A, right, see also

Fig. S4C). Curvy dynamics are represented with some difference of ω and same Φ values

(Fig. 3A, middle, see also Fig. S4B). We fitted this Lissajous curve function to each of the 20,000 bootstrap replicates derived from 10 neural populations (see Materials and Methods, 1000 replicates times 10 populations times two conditions). For instance, three replicated examples obtained from HPC population were well captured by the Lissagous curve function, as rotational-to-straight trajectories (Fig. 3B). We obtained all the estimates for these parameters (Fig. 3C). Thereafter, we applied clustering to these data (Fig. 3D-F) to specify the type of dynamics that appeared in the 10 brain regions as a function of the Lissajous curve parameters.

We found that the rotational dynamics (Fig. 3G-H, clusters C1-C3, reddish) appeared at the TE and STRt, which occupied high percentages of these neural populations (Fig. 3H, approximately 70%), and they were also observed in more than 50% of temporal brain regions. Cluster 3 seemed to have the intermediate characteristics between rotational and curvy structures; if we define this cluster as the curvy one, the rotational percentage becomes low in the PRC, PHC, and HPC (30-40%), but not in the STRt or TE (50-70%). Curvy structures were predominantly observed in the CDb population in more than 40% (Fig. 3G and H, Clusters 3 and 4, greenish). These clusters were observed not deterministically but stochastically, as seen in the predominant percentages of intermediate features between curvy and straight dynamics (Fig. 3G and H, Cluster 6) and almost straight dynamics (Fig. 3G and H, Clusters 7-10, bluish). Even with the STRt, the neural population contained curvy or straight dynamics in more than 20%. These heterogeneous mixtures of replicated signals in each population suggested that neural dynamics emerged in a stochastic manner with a functional gradient in the temporal/frontal/limbic networks of cortical and subcortical structures. A brain-wide neural population may propagate item information as a heterogeneous mixture within approximately half a second.

Discussion

Collectively, our results revealed that the parts of the orbitofrontal cortex (cOFC and mOFC) and their target subcortical brain region VS predominantly showed curvy, straight, and intermediate dynamics (Fig. 3G-H). These dynamics exhibited maximum modulation at approximately 0.3 sec after visual onset, except in the slowest VS dynamics (Fig. 1D). Rotational dynamics were observed at the temporal cortices and their connected striatal regions, at the relatively shorter latency around 0.20 s when rotation started (Fig. 1B). Remarkably, the rotational dynamics were observed at different neural proportions of replicated population in a stochastic manner (Fig. 3H), whereas different monkeys performed active (Exps. 1 and 3), and passive (Exps. 2 and 4) behavioral tasks. In contrast, straight dynamics started its geometric change approximately after 0.2 s of the visual onset (Fig. 1D, see the distance between initial point S and 0.2 s location), indicating that they follow the rotational dynamics. Taken together, these three dynamics were distinctive in terms of geometric patterns and their dynamic changes over time (Fig. 4B for summary), in which a rotational/curvy change was followed by a change in straight dynamics.

A previous study showed that rotational dynamics have been uncovered broadly in the primary sensory (*24, 35*) and motor (*36*) cortices, which are closer to the inputs and outputs of the brain, such as motor unit activity (*37*). Other studies have shown that the prefrontal cortex (*38, 39*) and parietal cortex (*22, 35*) exhibit curvy dynamics. We found curvy dynamics in the CDb, where action information was transferred from the cortices (*40*). These characteristic differences in the individual previous studies were not deterministic, similar to the stochastic differences among the populations observed in the present study, which may reflect the neural computations processed in the brain.

In the present study, we specifically focused on neural dynamics in two core areas: i)

low-dimensional geometries and ii) neural modulation dynamics. First, although these data

were obtained from four different laboratories using distinctive behavioral tasks during the

passive and active responses of monkeys, the low-dimensional features of neural

dynamics are thought to be preserved across mammals in the brain-wide network (*41, 42*).

A recent study provides clear evidence that different animal species share and preserve

their neural geometries during behavioral tasks (*41*). It is suggested that a low-dimensional

manifold in the neural state space might be one of the representational states of

biologically relevant information, similar to many combinations of physical properties in the

world (*43*). Second, the dynamics in neural modulations examined here are comparable

with those using the standard analytic frameworks in the rate coding model, which has

provided a huge amount of knowledge corresponding to low-dimensional neural activity

modulation in the literature, such as the Gabor function in the visual cortices (*44, 45*),

movement direction and muscle force in the motor cortex (*46, 47*), reward value in the

parietal cortex (*48*), and comprehension of the location of animals during navigation in the

HPC (*49*). Thus, in our analysis, the dynamics of these well-known brain region features

were compared as the geometric patterns across neural populations during visual

recognition (*29*).

One concern with our approach is that there may be limitations in data interpretation

in terms of data sharing and comparisons across different behavioral tasks and different

individual animals. Is it possible to compare the neural population trajectory using

accumulated data across animals and tasks with a certain analytical tool? In previous

literature, when we analyzed neural modulation using a linear regression model,

comparisons of firing rate modulations were accepted by most neurophysiologists for

different animals as well as different behavioral tasks. In one study, a challenge was made

to compare the neural modulation dynamics as the trajectory geometry between different

laboratories' data (*50*). Thus, the neural trajectory would be comparable among our shared

data with greater deliberation.

Our findings would add to the emerging literature describing how visual inputs alter

the brain-wide neural dynamics associated with visual memories by connecting neural

geometry types and their alignments across many brain regions. Previous studies have

shown the existence of different types of neural population dynamics in each individual

study (*22, 24, 35, 36, 38, 39*). Although some of these dynamics may reflect task demand,

as observed in the dorsolateral prefrontal cortex (*38, 51*), they are difficult to disentangle

from changes in behavior and neural activity levels and may involve some transformation

of information for behavioral responses (*52*). It is possible that the dorsal and motor-related

brain regions have this type of flexibility in their dynamics, as partly observed in this study

in the CDb, where curvy dynamics were predominantly observed (Fig. 4B).

Our results raise the possibility that geometric features determine the important neural

mechanisms widely observed in the brain. For instance, the stochastic gradient relative

distance to the visual input may reflect the dynamics of the neural circuitry in half a second

(Fig. 4B). The unidimensional straight dynamics in the hippocampal-frontal circuitry may

reflect memory access during visual recognition, such as location and reward. The

rotational dynamics might reflect the visual recognition process, during which recurrent

feedback signals change the circuit dynamics. Future studies should test the underlying

mechanisms and define whether engagement is best considered a change in behavior

and/or task context for whole-brain neural population activity. Regardless of the

mechanism, the shift of modulation structures in the lower-dimensional neural space could

play a fundamental role in brain-wide information processing, such as transforming visual

feature recognition to memory access.

Materials and Methods

Subjects and experimental procedures

Nine rhesus monkeys were used in the present study (Exp. 1: *Macaca mulatta,* A, 9.3 kg,

male; *Macaca mulatta,* D, 9.5 kg, male; Exp 2: Macaca mulatta, WK, 12.0 kg, male;

Macaca mulatta, SP, 7.0 kg, male; Exp 3: *Macaca mulatta,* BI, 8.2 kg, male; *Macaca*

mulatta, FG, 11.0 kg, male; *Macaca mulatta,* ST, 5.2 kg, male; Exp 4: *Macaca mulatta,*

SUN, 7.1 kg, male; *Macaca fuscata*, FU, 6.7 kg, female). All experimental procedures were

approved by the Institutional Animal Care and Use Committee of Laboratory Animals

approved by Peking University (Exp. 1, project number Psych-YujiNaya-1), and Animal

Care and Use Committee of the National Eye Institute, and complied with the Public Health

Service Policy on the Humane Care and Use of Laboratory Animals (Exp. 2, protocol

number NEI-622), the Animal Ethics Committee of the National Institutes for Quantum

Science and Technology (Exp. 3, protocol no. 11-1038-11), and the Animal Care and Use

Committee of the University of Tsukuba (Exp. 4, protocol no H30.336). All procedures

were performed in compliance with the US Public Health Service Guide for the Care and

Use of Laboratory Animals.

Behavioral task

Exp. 1. Item-location-retention (ILR) task. The animals performed the task under dim light

conditions in an electromagnetically shielded room. The task started with an encoding

phase, which was initiated by the animal pulling a lever and fixating on a white square

(0.6°) presented within one of four quadrants at 12.5° (monkey A) or 10° (monkey D) from

the center of the touchscreen (3M™ MicroTouch™ Display M1700SS, 17 in), situated

approximately 28 cm from the subjects. The eye position was monitored using an infrared

digital camera with a sampling frequency of 120 Hz (ETL-200, ISCAN). After fixation for

0.6 s, one of six items (3.0° for monkey A and 2.5° for monkey D, radius) was presented

in the same quadrant as a sample stimulus for 0.3 s, followed by another 0.7 s fixation on

the white square. If the fixation was successfully maintained (typically < 2.5°), the encoding

phase ended with the presentation of a single drop of water.

The encoding phase was followed by a blank interphase delay interval of 0.7–1.4 s

during which no fixation was required. The response phase was initiated using a fixation

dot presented at the center of the screen. One of the six items was then presented at the

center for 0.3 s, as a cue stimulus. After another 0.5 s delay period, five disks were

presented as choices, including a blue disk in each quadrant and a green disk in the center.

When the cue stimulus was the same as the sample stimulus, the animal was required to

make a choice by touching the blue disk in the same quadrant as the sample (i.e., the

match condition). Otherwise, the subject was required to choose the green disk (i.e., non-

match condition). If the animal made the correct choice, four–eight drops of water were

provided as a reward; otherwise, an additional 4 s was added to the standard inter-trial

interval (1.5–3 s). The number of reward drops was increased to encourage the animal to

maintain good performance in the latter phase of a daily recording session, which was

typically conducted in blocks (e.g., a minimal set of 60 trials with equal numbers of visual

items presented in a match/nonmatch condition). During the trial, a large gray square (48°

on each side) was presented at the center of the display as a background. At the end of

the trial, all stimuli disappeared, and the entire screen displayed a light red color during

the inter-trial interval. The start of a new trial was indicated by the reappearance of a large

gray square on the display, at which point the monkey could pull the lever, triggering the

appearance of a white fixation dot.

In the match condition, sample stimuli were chosen pseudo-randomly from six well-

learned visual items, and each item was presented pseudo-randomly within four quadrants,

resulting in 24 (6 × 4) configuration patterns. In the non-match condition, the location of

the sample stimulus was randomly chosen from the four quadrants, and the cue stimulus

was randomly chosen from the remaining five items that differed from the sample. The

match and non-match conditions were randomly presented in a ratio of 4:1, resulting in 30

(24 + 6) configuration patterns. The same six stimuli were used during all the recording

sessions.

Exp. 2. Scene-based object-value task. Animals learned the scene-object associations.

After the monkeys fixated on the red-square fixation point on the scene image for 0.6–1 s,

the fixation cue disappeared, and two visual items (objects of different values) appeared

simultaneously in a different hemifield (for training and neuronal testing) or the same

hemifield (for pharmacological experiments). A reward was given after the monkeys made

a saccade to the stimulus and maintained fixation for 0.2 s. Half of the fractal visual items

were associated with a large reward (0.3 mL), and the other half were associated with a

small reward (0.1 mL). This reward association changed depending on the scene (Fig. S1

D). *Passive Viewing Task.* One of the two scene images was presented for 0.8 s randomly.

If the monkey fixated on a central red square, two to four fractals were presented

sequentially on the scene image within the neuron's receptive field (presentation time, 0.4

s; interstimulus interval, 0.4 s; Fig. S1C). A liquid reward (0.2 mL) was delivered 0.3 s after

the last object was presented. Thus, reward occurrence was not associated with any of

the visual items. Each item was presented at least seven times per session.

Exp. 3. Delayed reward tasks. The monkeys were seated on a primate chair inside a dark,

sound-attenuated, electrically shielded room. A touch-sensitive bar was mounted on the

chair. The visual stimuli were displayed on a computer video monitor placed in front of the

animals. Each of the six cues was associated with a combination of reward size (1 drop;

3 or 4 drops) and reward delay (0, 3.3, and 6.9 s). The trials began when the monkey

touched the bar. A visual cue appeared, and the monkey released a bar when a red spot

(waiting signal) turned green (go signal) after a variable interval. If the monkey released

the bar 0.2–1 s after this go signal, the trial was considered correct and the spot turned

blue (correct signal). A liquid reward of a small (1 drop, approximately 0.1 mL) or large

amount (3 drops, except for monkey BI, 4 drops) was delivered immediately (0.3 ± 0.1 s)

or with an additional delay of either 3.3 ± 0.6 s or 6.9 ± 1.2 s after correct release of the

bar. The cues were chosen with equal probability and were independent of the preceding

reward condition. Anticipatory bar releases (before or no later than 0.2 s after the

appearance of the go signal) and failure to release the bar within 1 s of the appearance of

the go signal were counted as errors. In the error trials, the trial was terminated

immediately, all visual stimuli disappeared, and following inter trial interval (1 s), the trial

was repeated; that is, the reward size/delay combination remained the same as that in the

error trial. Behavioral control and data acquisition were performed using a real-time

experimentation system (REX) (*53*). The Neurobehavioral Systems Presentation software

was used to display the visual stimuli (Neurobehavioral Systems).

Exp. 4. Cued lottery tasks. The animals performed one of two visually cued lottery tasks:

a single-cue or a choice task. Neuronal activity was only recorded during the single-cue

task.

Animals performed the task under dim lighting conditions in an electromagnetically

shielded room. Eye movements were measured using a video camera system at 120 Hz

(EyeLink, SR Research). Visual stimuli were generated using a liquid-crystal display at 60

Hz, placed 38 cm from the monkey's face when seated. At the beginning of the single-cue

task trials, the monkeys had 2 s to align their gaze within 3° of a 1°-diameter gray central

fixation target. After a fixation for 1 s, a pie chart was presented for 2.5 s, to provide

information regarding the probability and magnitude of rewards in the same location as

the central fixation target. The probability and magnitude of the rewards were associated

with the number of blue and green 8° pie chart segments, ranging from 0.1 to 1.0 mL in

0.1 mL increments for magnitude, and 0.1 to 1.0 in 0.1 increments for probability. Following

a 0.2 s interval from the removal of the pie chart, a 1 kHz or 0.1 kHz tone of 0.15 s duration

was provided to indicate reward or no-reward outcomes, respectively. After a 0.2 s interval

following the high tone, a fluid reward was delivered, whereas no rewards were delivered

following the low tone. An inter-trial interval of 4–6 s was used. During the choice task,

animals were instructed to choose one of two peripheral pie charts, each of which

indicated either the probability or magnitude of an upcoming reward. The two target

options were presented for 2.5 s at 8° to the left or right of the central fixation location. The

animals received a fluid reward as indicated by the green pie chart of the chosen target,

with the probability indicated by the blue pie chart. Otherwise, no reward was delivered.

A total of 100 pie charts composed of 10 levels of probability and magnitude of

rewards were used in the experiments. In the single-cue task, 100 pie charts were

presented once in random order. In the choice task, two pie charts were randomly

assigned to the two options. During one electrophysiological recording session,

approximately 30–60 trial blocks of the choice task were interleaved with 100–120 trial

blocks of the single-cue task.

Electrophysiological recordings and data preprocessing

Exp. 1. To record the single-unit activity, we used a 16-channel vector array microprobe

(V1 X 16-Edge, NeuroNexus), 16-channel U-Probe (Plexon), tungsten tetrode probe

(Thomas RECORDING), or single-wire tungsten microelectrode (Alpha Omega).

Electrophysiological signals were amplified, bandpass-filtered (200–6000 Hz), and

monitored. Single-neuron activity was isolated based on spike waveforms, either online or

offline. For both clustering and offline sorting, the activities of all single neurons were sampled when the activity of an isolated neuron demonstrated a good signal-to-noise ratio (>2.5). The signal-to-noise ratio was visually checked by calculating the range of background noise against the spike amplitude, which was monitored online using the OmniPlex Neural Data Acquisition System, or offline using the sorter software Plexon. The recorded neurons were not blinded. The sample sizes required to detect the effect sizes (numbers of recorded neurons, recorded trials in a single neuron, and monkeys) were estimated based on previous studies (*31, 54*). Neural activity was recorded during 60–240 trials of the ILR task. We recorded 590 hippocampal neurons, among which the recording sites appeared to cover all subdivisions (i.e., the dentate gyrus, CA3, CA1, and subicular complex).

Exp. 2. We used conventional techniques to record the single-neuron activity in the STRt, including the caudate and putamen tails. A tungsten microelectrode (1–3 MΩ Frederic hair; 0.5-1.5 MΩ Alpha Omega Engineering) was used to record single-neuron activity. The recording site was determined using a grid system that allowed electrode penetration at 1 mm intervals. We amplified and filtered (0.3 to 10 kHz; Model 1800, A-M Systems; Model MDA-4I, BAK) signals obtained from the electrodes and collected at 1 kHz. Single neurons were isolated online using custom voltage–time window discriminator software (Blip; available at http://www.robilis.com/blip/). The presumed medium spiny neurons were identified based on their low baseline activity (<3 spikes/s) and broad action potentials (*55*). The recorded neurons were not blinded. The sample sizes required to detect the effect sizes (numbers of recorded neurons, recorded trials in a single neuron, and monkeys) were estimated based on previous studies (*56, 57*). Neural activity was recorded during 10–30 trials of the *passive viewing task*. We recorded 115 medium spiny neurons in the STRt. In Exp. 2, only a single-neuron recording was performed online. We note that

we termed the scene and object for two visual stimuli in our previous study (*58*), but here

we termed them scene and item.

Exp. 3. Conventional techniques were used to record single-neuron activity in the dorsal

part of the head of the caudate nucleus (CD). A tungsten microelectrode (1.1–1.5 MΩ,

Microprobes for Life Science; 1.0 MΩ, Alpha Omega Engineering Ltd.) was used to record

single-neuron activity. The electrophysiological signals were amplified and monitored

using a TDT recording system (RZ2, Tucker-Davis Technologies, TDT). Single-neuron

activity was manually isolated based on the online spike waveforms. The activity of all

single neurons was sampled from the activity of presumed projection neurons, which are

characterized as having a low spontaneous discharge rate (<2 spikes/s) outside the task

context and exhibiting phasic discharges in relation to one or more behavioral task events

(Yamada et al, 2016). Neural activity was recorded during 100–120 trials per block in the

delayed-reward task. We recorded the CD of the left or right hemisphere in each of the

three monkeys in the experiment, with 150 CD neurons (51, 31, and 68 from the BI, FG,

and ST, respectively).

Exp. 4. Conventional techniques were used to record single-neuron activity in the DS, VS,

cOFC (area 13M), and mOFC (area 14o). A tungsten microelectrode (1–3 MΩ, FHC) was

used to record single-neuron activity. Electrophysiological signals were amplified, band-

pass filtered (50–3,000 Hz), and monitored using a TDT recording system (RZ5D, Tucker-

Davis Technologies, TDT). Single-neuron activity was manually isolated based on the

online spike waveforms. The activity of all single neurons was sampled when the activity

of an isolated neuron demonstrated a good signal-to-noise ratio (>2.5). The signal-to-noise

ratio was calculated online as the ratio of the spike amplitude to the baseline voltage range

on the oscilloscope. The recorded neurons were not blinded. The sample sizes required

to detect the effect sizes (numbers of recorded neurons, recorded trials in a single neuron,

and monkeys) were estimated based on previous studies (*59-61*). Neural activity was

recorded during 100–120 trials of the single-cue task. Neural activity was not recorded

during selection trials. We recorded the neurons of a single right hemisphere in each of

the two monkeys: 194 DS neurons (98 and 96 from monkeys SUN and FU, respectively),

144 VS neurons (89 SUN and 55 FU), 190 cOFC neurons (98 SUN and 92 FU), and 158

mOFC neurons (64 SUN and 94 FU). In Exp. 1, only a single-neuron recording was

performed online.

Statistical analysis

For statistical analysis, we used the statistical software package MATLAB (MathWorks,

Exps. 1 and 2), and R (Exps. 3 and 4) for conventional analyses such as linear regression

and ANOVA. To analyze the regression matrix using PCA, we used R software. All

statistical tests for the neural analyses were two-tailed.

Behavioral analysis

No new behavioral results were included; however, the procedure for the behavioral

analysis was as follows:

Exp. 1. We previously reported that two monkeys learned to retain the item and location

information of a sample stimulus (*62*). Here, we describe the analysis steps used to check

whether the monkey used both item and location information to perform the task.

To examine this, we compared the animals' actual correct rates during the recording

to random correct rates (chi-square test). The ILR response phase had five options,

resulting in a 20% random correct rate. If the animal used an incorrect strategy, such as

only retaining the location information of the sample stimulus and ignoring the item

information, the correct rate for the match condition would be 100% and that for the

nonmatch condition would be 0. Based on the above considerations, we examined the

correct rates of the two animals in the match and nonmatch conditions, respectively. In

general, the average correct rates for both animals in the match and nonmatch conditions

were well above chance levels after training.

Exp. 2. We previously reported that two monkeys switched their behavior depending on

the value of the item based on the scene (*58*). Here, we describe how to check whether

the monkey learned both the scene and item information. We calculated the correct rate

for the scene-based object-value task. Because the two scenes appear in random

sequences, the monkey must switch object choice if the scene has changed. After

performing more than 160 trials, the correct rate reached a plateau above chance. The

monkey was able to switch object choices immediately after the scene changed. Once the

monkeys learned this extensively, their choice behavior became automatic, as the choice

tended to occur even when the reward was not delivered after saccades to high-valued

items according to the scene.

Exp. 3. We previously reported that the three monkeys behaved based on temporally

discounted values that integrated both delay and reward size information provided by

visual stimuli (*63*). Here, we describe an analysis to check how monkeys discount reward

values for delay and reward information. Error rates in task performance were calculated

by dividing the total number of errors by the total number of trials for each reward condition

and then averaged across all sessions. The average error rates were fitted to the inverse

function of reward size with hyperbolic temporal discounting: $E = 1+kD/aR$ (*E*: average

error rates, *D*: delay, *R*: reward size, *k*: discounting factor, *a*: incentive impact), and

exponential temporal discounting: $E = e^{-kd}/aR$. We used the 'optim' function in R, evaluated

the goodness of fit of the two models by least-squares minimization, and compared the

models by leave-one-out cross-validation as described previously (Minamimoto et al.,

2009).

Exp. 4. We previously reported that monkey behavior depends on expected values,

defined as the probability time magnitude (*23*). We described the analysis steps to check

whether the monkey's behavior reflected task parameters, such as reward probability and

magnitude. Importantly, we showed that the monkey's choice behavior reflected the

expected values of the rewards, that is, the probability multiplied by the magnitude. For

this purpose, the percentage choosing the right-side option was analyzed in the pooled

data using a general linear model with a binomial distribution:

$Pchooses_R = 1/(1 + e^{-z})$ (3)

where the relationship between $Pchooses_R$ and Z is given by the logistic function in each

of the following three models: number of pie segments (M1), probability and magnitude

(M2), and expected values (M3).

M1: $Z = b_0 + b_1 Npie_L + b_2 Npie_R$ (4)

where b_0 is the intercept, and $Npie_L$ and $Npie_R$ are the number of pie segments contained

in the left and right pie chart stimuli, respectively. The values of b_0 to b_2 are free parameters

and estimated by maximizing the log likelihood.

M2: $Z = b_0 + b_1 P_L + b_2 P_R + b_3 M_L + b_4 M_R$ (5)

where b_0 is the intercept; P_L and P_R are the probabilities of rewards for the left and right

pie chart stimuli, respectively; and M_L and M_R are the magnitudes of rewards for the left

and right pie chart stimuli, respectively. The values of b_0 to b_4 are free parameters and

estimated by maximizing the log likelihood.

M3: $Z = b_0 + b_1 EV_L + b_2 EV_R$ (6)

where b_0 is the intercept and EV_L and EV_R are the expected values of rewards as

probability multiplied by magnitude for the left and right pie chart stimuli, respectively. The

values of b_0 to b_2 are free parameters and estimated by maximizing the log likelihood. We

identified the best model to describe monkey behavior by comparing goodness-of-fit

based on Akaike's information criterion and Bayesian information criterion (*64*).

Neural analysis

Peri-stimulus time histograms were constructed for each single-neuron activity aligned at

the onset of the visual stimulus. Average activity curves were smoothed for visual

inspection using a Gaussian kernel ($\sigma = 20, 15, 10$, and 50 ms in Exps. 1–4, respectively),

whereas the Gaussian kernel was not used for statistical tests.

To ensure that the four different datasets were as fair as possible, we used the same

criteria to analyze the neural activity. For the neural analyses, we used the following four

criteria: 1) the same analysis window size, 2) visual response within a short time (0.6 s),

3) neural modulations detected at the same significance level ($P < 0.05$), and 4) a general

linear model (ANOVA in Exps. 1 and 2 and the linear regression in Exp. 3 and 4). The

details of these analytical procedures for the rate coding and dynamic models are shown

below.

Rate-coding model: Conventional analyses to detect neural modulations in each

neuron

Exp. 1. For neural responses during the encoding phase after the sample presentation,

we evaluated the effects of "item" and "location" for each neuron using two-way ANOVA

($P < 0.05$ for each). We analyzed neurons that were tested in at least 60 trials (10 trials

for each stimulus and 15 trials for each location). On average, we tested 100 trials for each

neuron. These results have been previously reported (*62*).

Exp. 2. For neural responses during the appearance of the visual item, we evaluated the

effects of "item" and "scene" for each neuron using paired t-test ($P < 0.05$ with Bonferroni

correction). These results have been previously reported (*58*).

Exp. 3. The neural discharge rates (F) were fitted using a linear combination of the

following variables:

$F = b_0 + b_d \, \text{Delay} + b_m \, \text{Magnitude}$ (7)

where Delay and Magnitude are the delay and magnitude of the reward, respectively, as

indicated by the visual stimulus. b_0 is the intercept. If b_d and b_m were not zero at $P < 0.05$,

the discharge rates were regarded as being significantly modulated by that variable. These

results have been previously reported (*63*).

Exp. 4. The neural discharge rates (F) were fitted using a linear combination of the

following variables:

$F = b_0 + b_p \, \text{Probability} + b_m \, \text{Magnitude}$ (8)

where Probability and Magnitude are the probability and magnitude of the rewards,

respectively, as indicated by the pie chart. b_0 is the intercept. If b_p and b_m were not zero at

$P < 0.05$, the discharge rates were regarded as being significantly modulated by that

variable. These results have been previously reported (*23*).

Population dynamics using principal component analysis

We analyzed neural activity during an identical 0.6 s duration from the sample onset (Exp.

1), item onset (Exp. 2), CUE onset (Exp. 3), and cue onset (Exp. 4). To obtain a time series

of neural firing rates within this time period, we estimated the firing rates of each neuron

for every 0.05 s time bin (without overlap) during the analysis periods. A Gaussian kernel

was not used.

Regression subspace. We used a general linear model to determine how items and

locations (Exp. 1), items and scenes (Exp. 2), delay and magnitude of rewards (Exp. 3),

and the probability and magnitude of the rewards (Exp. 4) affect the activity of each neuron

in the neural populations. Each neural population was composed of all the recorded

neurons in each brain region.

Exp. 1. First, we set six visual items and four locations as categorical variables. We then

described the average firing rate of neuron i at time t as a linear combination of the item

and the location in each neural population:

$$F_{(i,t,k)} = b_{0(i,t)} + b_{1(i,t)}\text{Item}_{(k)} + b_{2(i,t)}\text{Location}_{(k)} \qquad (9)$$

where $F_{(i,t,k)}$ is the average firing rate of neuron i at time t in trial k, $\text{Item}_{(k)}$ is the types of

items cued to the monkey in trial k, and $\text{Location}_{(k)}$ *is* the types of locations cued to the

monkey in trial k. The regression coefficients $b_{0(i,t)}$, $b_{1(i,t)}$, and $b_{2(i,t)}$ describe the degree to

which the firing rates of neuron i depend on the mean firing rates (hence, firing rates

independent of task variables, item, and location), the degree of firing rate in each item

relative to the mean firing rates, and the degree of firing in each location relative to the

mean firing rates, respectively, at a given time t during the trials. The interaction term is

not included in the model.

In the analysis, we performed preference ordering for item and location in each neuron.

$\text{Item}_{(k)}$ and $\text{Location}_{(k)}$ were rank-ordered items and locations, respectively, cued to the

monkey in trial k. Items 1–6 and locations 1–4 were rank-ordered from the most preferred

to least preferred, respectively, defined as the mean firing rate during the entire analysis

time window from 0.08 to 0.6 s. This preference ordering did not change over time t for

each neuron n.

Exp. 2. We first set eight items and two scenes as the categorical variables. We then

described the average firing rate of neuron i at time t as a linear combination of the item

and scene in each neural population:

$F_{(i,t,k)} = b_{0(i,t)} + b_{1(i,t)}\text{Item}_{(k)} + b_{2(i,t)}\text{Scene}_{(k)}$ (10)

where $F_{(i,t,k)}$ is the average firing rate of neuron i at time t in trial k, $\text{Item}_{(k)}$ is the types of

items cued to the monkey in trial k, and $\text{Scene}_{(k)}$ is the types of scene stimuli cued to the

monkey in trial k. The regression coefficients $b_{0(i,t)}$, $b_{1(i,t)}$ and $b_{2(i,t)}$ describe the degree to

which the firing rates of neuron i depend on the mean firing rates (hence, firing rates

independent of task variables, item and scene), the degree of firing rate in each item

relative to the mean firing rates, and the degree of firing in each scene relative to the mean

firing rates, respectively, at a given time t during the trials. The interaction term was not

included in the model.

In the analysis, $\text{Item}_{(k)}$ and $\text{Scene}_{(k)}$ were the rank-ordered item and scene,

respectively, cued to the monkey in trial k. Items 1 to 8 and Scenes 1 and 2 were rank-

ordered from the most preferred to least preferred, respectively, defined as the mean firing

rate during the whole analysis 0.6 s window after the item onset. This preference ordering

did not change over time t for each neuron n.

Exp. 3. We first set the delay and magnitude as 0, 3.3, and 6.9 s and one and three drops

of rewards, respectively, during the behavioral task. In the analysis, we normalized these

values from 0 to 1 divided by the maximum values in each: 0, 0.48, and 1 for delay, and

0.33, 0.66, and 1 for magnitude. This is because these values affect the extent of the

regression subspace between two continuous variables. We then described the average

firing rate of neuron i at time t as a linear combination of the delay and magnitude in each

neural population:

$F_{(i,t,k)} = b_{0(i,t)} + b_{1(i,t)}\text{Delay}_{(k)} + b_{2(i,t)}\text{Magnitude}_{(k)}$ (11)

where $F_{(i,t,k)}$ is the average firing rate of neuron i at time t in trial k, Delay$_{(k)}$ is the normalized

delay to obtain a reward cued to the monkey in trial k, and Magnitude $_{(k)}$ *is* the normalized

number of reward drops cued to the monkey in trial k. The regression coefficients $b_{0(i,t)}$ to

$b_{2(i,t)}$ describe the degree to which the firing rates of neuron i depend on the mean firing

rates (hence, firing rates independent of task variables), delay in rewards, and magnitude

of rewards, respectively, at a given time t during the trials.

Exp. 4. We first set the probability and magnitude as 0.1 to 1.0 and 0.1 to 1.0 mL,

respectively. We did not normalize these values because they were originally prepared

from 0 to 1 originally. We then describe the average firing rate of neuron i at time t as a

linear combination of probability and magnitude in each neural population:

$$F_{(i,t,k)} = b_{0(i,t)} + b_{1(i,t)}\text{Probability}_{(k)} + b_{2(i,t)}\text{Magnitude}_{(k)} \qquad (12)$$

where $F_{(i,t,k)}$ is the average firing rate of neuron i at time t in trial k, Probability$_{(k)}$ is the

probability of the reward cued to the monkey in trial k, and Magnitude$_{(k)}$ *is* the magnitude

of the reward cued to the monkey in trial k. The regression coefficients $b_{0(i,t)}$ to $b_{2(i,t)}$ describe

the degree to which the firing rates of neuron i depend on the mean firing rates (i.e., firing

rates independent of task variables), probability of rewards, and magnitude of rewards,

respectively, at a given time t during the trials.

We used the regression coefficients (i.e., the regression table in the case of ANOVA)

described in Eq. 9–12 to identify how the dimensions of the neural population signals were

composed of information related to the item and location (Exp. 1), item and scene (Exp.

2), delay and magnitude (Exp. 3), and probability and magnitude (Exp. 4) as aggregated

properties of individual neural activity. In this step, an encoding model is constructed in

which the regression coefficients are explained by a temporal structure in the neural

modulation of two categorical variables (Exps. 1 and 2), or two continuous variables (Exps.

3 and 4) at the population level. Our procedures involve targeted dimensionality reduction

using the regression subspace and are aimed at describing neural modulation dynamics

(*29*).

Principal component analysis. We used PCA to identify the dimensions of the neural

population signal in orthogonal spaces composed of two variables in each neural

population of the four experiments. For each neural population, we first prepared a two-

dimensional data matrix X of size $N_{(n)} \times M_{(C \times T)}$. The regression coefficient vectors $b_{1(i,t)}$ and

$b_{2(i,t)}$ in Eq. 9–12, whose rows correspond to the total number of neurons (n) in each neural

population and columns correspond to C, the total number of conditions (that is, 10: six

items and four locations in Exp. 1, 10: eight items and two scenes in Exp. 2, 2: delay and

magnitude in Exp. 3, and 2: probability and magnitude in Exp. 4), and T is the total number

of analysis windows (i.e., 0.6 s divided by the window size bin, 0.05 s, 12 bin). A series of

eigenvectors was obtained by applying PCA once to the data matrix X in each neural

population. The PCs of this data matrix are vectors $v_{(a)}$ of length $N_{(n)}$ and the total number

of recorded neurons if $M_{(C \times T)} > N_{(n)}$; otherwise, the length is $M_{(C \times T)}$. PCs were indexed from

the principal components and explained the most to least variance. The eigenvectors were

obtained using the prcomp () function in R software. We did not include the intercept term

$b_{0(i,t)}$ to focus on the neural modulation by the variables of interest.

Eigenvectors. When we applied PCA to data matrix X, we decomposed the matrix into

eigenvectors and eigenvalues. Each eigenvector had a corresponding eigenvalue. In our

analysis, the eigenvectors at time t represented a vector, for example, in the space of

delay and magnitude in Exp. 3. The eigenvalues at time t for the delay and magnitude

were scalars, indicating the extent of variance in the data in that vector. Thus, the first PC

was the eigenvector with the highest eigenvalue. We analyzed the eigenvectors for the

top two PCs (PC1 and PC2) in the following analyses to describe the geometry in the most

predominant dimension. PCA was applied once to each neural population; thus, the total

variance contained in the data differed among the neural populations.

Shuffle control for PCA. To examine the significance of the population structures described

by PCA, we performed three shuffle controls. The two-dimensional data matrix X was

randomized by shuffling in three ways. In shuffled control 1, matrix X was shuffled by

permutating the allocation of neuron n at time i. This shuffle provided a data matrix X of

size $N_{(n)} \times M_{(C \times T)}$, eliminating the temporal structure of neural modulation by condition C in

each neuron but retaining the neural modulations at time t at the population level. In

shuffled control 2, matrix X was shuffled by permutating the allocation of time i in each

neuron n. This shuffle provided a data matrix X of size $N_{(neuron)} \times M_{(C \times T)}$, eliminating the

neural modulation structure under condition C maintained in each neuron but retaining the

neural modulation in each neuron at the population level. In shuffled control 3, matrix X

was shuffled by permutating the allocation of both time i and neuron n. In these three

shuffle controls, matrix X was estimated to be 1,000 times. PCA performance was

evaluated by constructing the distributions of the explained variances for PC1 to PC12.

The statistical significance of the variances explained by PC1 and PC2 was estimated

based on the 95th percentile of the reconstructed distributions of the explained variance

or bootstrap standard errors (i.e., standard deviation of the reconstructed distribution). We

note that because the significant dimensions of neural populations dynamics differed the

10 neural populations, we analyzed the neural dynamics at the top two dimension, PC1

and 2.

Analysis of eigenvectors. We evaluated the characteristics of the eigenvectors for PC1

and PC2 in each neural population in terms of vector angle, size, and deviation. The

eigenvectors were evaluated for each of the task parameters described above: item and

location in Exp. 1, item and scene in Exp. 2, delay and magnitude in Exp. 3, and probability

and magnitude in Exp. 4. The angle is the vector angle from the horizontal axis from 0º to

360º against the main PCs. The size is the length of the eigenvector. The deviation is the

difference between the vectors. The deviation from the mean vector for each neural

population was estimated. These three eigenvector characteristics were compared among

the populations at $P < 0.05$, using the Kruskal–Wallis test and Wilcoxon rank-sum test with

Bonferroni correction for multiple comparisons. The vector during the first 0.1 s was

extracted from these basic analyses.

To evaluate the neural population geometry using their selected feature, we estimated

the accumulated angle difference weighted by the deviance:

744
$$\sum_{t=S}^{t=E} d\,\theta \qquad (13)$$

where the d is deviation between the vectors at times t and t+1, θ is the angle difference

between vectors at times t and t+1, S is zero, and E is the time period to stop the estimation,

i.e., 0.6 s. This index is analogous to the rotational force accumulated over time. If the

value of the accumulated angle difference was close to zero, the population geometry was

stable, such as a straight or non-dynamic structure, that is, it remained at some point in

the PC1-2 plane.

To quantitatively evaluate the trajectory geometry, we used the Lissajous curve

function, which describes any geometric pattern in a plane using $F(x,y)$:

$x = Ax \cos(\omega x\, t(i) + \Phi x) + bx$ (14)

$y = Ay \cos(\omega y\, t(i) + \Phi y) + by$ (15)

where ω and Φ represent cycle of the rotation and their deviance as a function of time, t(i).

Ax and *Ay* represent the amplitudes of the trajectory, whereas *bx* and *by* represent the

location of the trajectory. For ω, 3.33 π indicates that one cycle since the analysis window

is 0.6 s. Φ is 0 to 2 π for one cycle. We estimated *ωx, Φx, bx, ωy,* and *Φy, by* parameters

by estimating maximum loglikelihood of the model. Nonlinear least squares in the nls()

function in the R program was used. A time series of eigenvectors for PC1 and PC2 in a

0.05 s analysis windows (12 data points) were used with a sliding average between three

time points (hence, 0.15 s time resolution).

Bootstrap resampling and clustering using feature-based parameters. We estimated $\Sigma d\ \theta$,

mean *d*, rotational speed $\Sigma\theta/0.1$s, and $d_{\text{s-e}}$, such as start to end distance using a

parametric bootstrap resampling method (*65*). In each neural population, the neurons

were randomly resampled in duplicate, and a data matrix *X* of size $N_{(neuron)} \times M_{(C \times T)}$ was

obtained. PCA was applied to the data matrix *X.* The time series of eigenvectors was

obtained, and these four features were estimated from the neural trajectory. This

resampling was conducted 1,000 times in each neural population, and the distributions of

these four parameters were obtained.

Following bootstrap resampling, we applied clustering of these parameters based on

PCA and a dendrogram across the replicates in the 10 brain regions, such as 20,000

replicates (10 brain regions times two conditions times 1,000 replicates). Based on this

clustering, proportion of the identified clusters in each brain region was estimated.

Bootstrap resampling and clustering based on Lissajous curve parameters. The Lissajous

curve parameters for the replicated trajectory were estimated using a bootstrap

resampling method (*65*). In each neural population, the neurons were randomly resampled

in duplicate, and a data matrix *X* of size $N_{(neuron)} \times M_{(C \times T)}$ was obtained. PCA was applied to

the data matrix *X.* The time series of eigenvectors was obtained for PC1 and PC2, which

describe the trajectory, and the fitted parameters using the Lissajous curve function were

estimated using the nls() function in R program. This resampling was conducted 1,000

times in each neural population, and the distributions of the Lissajous parameters were

obtained.

Following bootstrap resampling, we applied clustering of these parameters based on

PCA and a dendrogram across the replicates in the 10 brain regions, such as 20,000

replicates (10 brain regions times two conditions times 1,000 replicates). In this process,

the $\omega x/\omega y$ and Φx-Φy were also estimated (Fig. S4), in addition to the ωx, ωy, Φx, and

Φy. Based on this clustering, proportion of the identified clusters in each brain region was

estimated. We used the median of the estimated parameter in a cluster to describe the

trajectory geometries.

References

1.   D. J. Felleman, D. C. Van Essen, Distributed hierarchical processing in the
primate cerebral cortex. *Cereb Cortex* **1**, 1-47 (1991).

2.   H. Chen, Y. Naya, Reunification of Object and View-Center Background
Information in the Primate Medial Temporal Lobe. *Front Behav Neurosci*
15, 756801 (2021).

3.   G. Buzsaki, S. McKenzie, L. Davachi, Neurophysiology of Remembering.
Annu Rev Psychol **73**, 187-215 (2022).

4.   J. B. Goense, N. K. Logothetis, Neurophysiology of the BOLD fMRI signal
in awake monkeys. *Curr Biol* **18**, 631-640 (2008).

5.   S. Hu *et al.*, Real-Time Readout of Large-Scale Unsorted Neural
Ensemble Place Codes. *Cell Rep* **25**, 2635-2642 e2635 (2018).

6.   N. A. Steinmetz, P. Zatka-Haas, M. Carandini, K. D. Harris, Distributed
coding of choice, action and engagement across the mouse brain. *Nature*
576, 266-273 (2019).

7.   J. R. Climer, D. A. Dombeck, Information Theoretic Approaches to
Deciphering the Neural Code with Functional Fluorescence Imaging.
eNeuro **8**, (2021).

8.   A. C. Paulk *et al.*, Large-scale neural recordings with single neuron
resolution using Neuropixels probes in human cortex. *Nat Neurosci* **25**,
252-263 (2022).

9.   J. Manley *et al.*, Simultaneous, cortex-wide dynamics of up to 1 million
neurons reveal unbounded scaling of dimensionality with neuron number.
Neuron **112**, 1694-1709 e1695 (2024).

10.   P. Dayan, L. Abbott, *Theoretical neuroscience: computational and*
mathematical modeling of neural systems. (2001).

11.   P. Sanz-Leon, S. A. Knock, A. Spiegler, V. K. Jirsa, Mathematical
framework for large-scale brain network modeling in The Virtual Brain.
Neuroimage **111**, 385-430 (2015).

12.   R. Yuste, From the neuron doctrine to neural networks. *Nat Rev Neurosci*
16, 487-497 (2015).

13.   S. Vyas, M. D. Golub, D. Sussillo, K. V. Shenoy, Computation Through
Neural Population Dynamics. *Annu Rev Neurosci* **43**, 249-275 (2020).

14.   M. D. Humphries, Strong and weak principles of neural dimension
reduction. *Neurons, Behavior, Data analysis, and Theory* **5**, (2021).

15.   K. V. Shenoy, J. C. Kao, Measurement, manipulation and modeling of
brain-wide neural population dynamics. *Nat Commun* **12**, 633 (2021).

16.   L. M. K. Timothy, B. E. Bona, *State Space Analysis: An Introduction*.
(McGraw-Hill, 1968).

17.   W. Brendel, R. Romo, C. K. Machens, Demixed Principal Component
Analysis. *Advances in Neural Information Processing Systems* **24**, 2654-
2622 (2011).

18.   M. M. Churchland *et al.*, Neural population dynamics during reaching.
Nature **487**, 51-56 (2012).

19.   V. Mante, D. Sussillo, K. V. Shenoy, W. T. Newsome, Context-dependent
computation by recurrent dynamics in prefrontal cortex. *Nature* **503**, 78-84
(2013).

20.   P. Gao, S. Ganguli, On simplicity and complexity in the brave new world of
large-scale neuroscience. *Curr Opin Neurobiol* **32**, 148-155 (2015).

21.   R. Rossi-Pool, R. Romo, Low Dimensionality, High Robustness in Neural
Population Dynamics. *Neuron* **103**, 177-179 (2019).

22.   G. Okazawa, C. E. Hatch, A. Mancoo, C. K. Machens, R. Kiani,
Representational geometry of perceptual decisions in the monkey parietal
cortex. *Cell* **184**, 3748-3761 e3718 (2021).

23.   H. Yamada, Y. Imaizumi, M. Matsumoto, Neural Population Dynamics
Underlying Expected Value Computation. *J Neurosci* **41**, 1684-1698
(2021).

24.   A. Libby, T. J. Buschman, Rotational dynamics reduce interference
between sensory and memory representations. *Nat Neurosci* **24**, 715-726
(2021).

25.   A. Pellegrino, H. Stein, N. A. Cayco-Gajic, Dimensionality reduction
beyond neural subspaces with slice tensor component analysis. *Nat*
Neurosci, (2024).

26.   S. Mukherjee, B. Babadi, Adaptive modeling and inference of higher-order
coordination in neuronal assemblies: A dynamic greedy estimation
approach. *PLoS Comput Biol* **20**, e1011605 (2024).

27.   Y. J. Chang, Y. I. Chen, H. C. Yeh, S. R. Santacruz, Neurobiologically
realistic neural network enables cross-scale modeling of neural dynamics.
Sci Rep **14**, 5145 (2024).

28.   P. Vahidi, O. G. Sani, M. M. Shanechi, Modeling and dissociation of
intrinsic and input-driven neural population dynamics underlying behavior.
Proc Natl Acad Sci U S A **121**, e2212887121 (2024).

29. H. Chen *et al.*, Stable Neural Population Dynamics in the Regression
Subspace for Continuous and Categorical Task Parameters in Monkeys.
eNeuro **10**, (2023).

30. K. S. Saleem, K. Tanaka, Divergent projections from the anterior
inferotemporal area TE to the perirhinal and entorhinal cortices in the
macaque monkey. *J Neurosci* **16**, 4757-4775 (1996).

31. Y. Naya, M. Yoshida, Y. Miyashita, Forward processing of long-term
associative memory in monkey inferotemporal cortex. *J Neurosci* **23**,
2861-2871 (2003).

32. W. A. Suzuki, Y. Naya, The perirhinal cortex. *Annu Rev Neurosci* **37**, 39-
53 (2014).

33. D. Sasikumar, E. Emeric, V. Stuphorn, C. E. Connor, First-Pass
Processing of Value Cues in the Ventral Visual Pathway. *Curr Biol* **28**,
538-548 e533 (2018).

34. K. Palmer, T. Ridgway, O. Al-Rawi, I. Johnson, M. Poullis, Lissajous
figures: an engineering tool for root cause analysis of individual cases--a
preliminary concept. *J Extra Corpor Technol* **43**, 153-156 (2011).

35. Y. Osako *et al.*, Contribution of non-sensory neurons in visual cortical
areas to visually guided decisions in the rat. *Curr Biol* **31**, 2757-2769
e2756 (2021).

36. K. C. Ames, S. I. Ryu, K. V. Shenoy, Neural dynamics of reaching
following incorrect or absent motor preparation. *Neuron* **81**, 438-451
(2014).

37. N. J. Marshall *et al.*, Flexible neural control of motor units. *Nat Neurosci*
25, 1492-1504 (2022).

38. M. C. Aoi, V. Mante, J. W. Pillow, Prefrontal cortex exhibits
multidimensional dynamic encoding during decision-making. *Nat Neurosci*
23, 1410-1420 (2020).

39. M. C. Aoi, J. W. Pillow, Model-based targeted dimensionality reduction for
neuronal population data. *Adv Neural Inf Process Syst* **31**, 6690-6699
(2018).
40. Y. Fan, J. I. Gold, L. Ding, Frontal eye field and caudate neurons make
different contributions to reward-biased perceptual decisions. *Elife* **9**,
(2020).
41. M. Safaie *et al.*, Preserved neural dynamics across animals performing
similar behaviour. *Nature* **623**, 765-771 (2023).
42. S. Melbaum *et al.*, Conserved structures of neural activity in sensorimotor
cortex of freely moving rats allow cross-subject decoding. *Nat Commun*
13, 7420 (2022).
43. C. E. Allen, P. Beldade, B. J. Zwaan, P. M. Brakefield, Differences in the
selection response of serially repeated color pattern characters: standing
variation, development, and evolution. *BMC Evol Biol* **8**, 94 (2008).
44. D. J. Tolhurst, J. A. Movshon, Spatial and temporal contrast sensitivity of
striate cortical neurones. *Nature* **257**, 674-675 (1975).
45. J. P. Jones, L. A. Palmer, An evaluation of the two-dimensional Gabor
filter model of simple receptive fields in cat striate cortex. *J Neurophysiol*
58, 1233-1258 (1987).
46. A. P. Georgopoulos, J. F. Kalaska, R. Caminiti, J. T. Massey, On the
relations between the direction of two-dimensional arm movements and
cell discharge in primate motor cortex. *J Neurosci* **2**, 1527-1537 (1982).
47. E. E. Fetz, P. D. Cheney, Postspike facilitation of forelimb muscle activity
by primate corticomotoneuronal cells. *J Neurophysiol* **44**, 751-772 (1980).
48. M. L. Platt, P. W. Glimcher, Neural correlates of decision variables in
parietal cortex. *Nature* **400**, 233-238 (1999).
49. J. O'Keefe, J. Dostrovsky, The hippocampus as a spatial map. Preliminary
evidence from unit activity in the freely-moving rat. *Brain Res* **34**, 171-175
(1971).

50.   D. Kobak *et al.*, Demixed principal component analysis of neural
925   population data. *Elife* **5**, (2016).

926   51.   J. D. Murray *et al.*, Stable population coding for working memory coexists
927   with heterogeneous neural dynamics in prefrontal cortex. *Proc Natl Acad*
928   *Sci U S A* **114**, 394-399 (2017).

929   52.   R. Rossi-Pool, A. Zainos, M. Alvarez, G. Diaz-deLeon, R. Romo, A
930   continuum of invariant sensory and behavioral-context perceptual coding
931   in secondary somatosensory cortex. *Nat Commun* **12**, 2000 (2021).

932   53.   A. Hays, B. Richmond, L. Optican, Unix-based multiple process system for
933   real-time data aquisition and control. *WESCON* **2**, 1-10 (1982).

934   54.   Y. Naya, W. A. Suzuki, Integrating what and when across the primate
935   medial temporal lobe. *Science* **333**, 773-776 (2011).

936   55.   H. Yamada *et al.*, Characteristics of fast-spiking neurons in the striatum of
937   behaving monkeys. *Neurosci Res* **105**, 2-18 (2016).

938   56.   S. Yamamoto, I. E. Monosov, M. Yasuda, O. Hikosaka, What and where
939   information in the caudate tail guides saccades to visual objects. *J*
940   *Neurosci* **32**, 11005-11016 (2012).

941   57.   J. Kunimatsu, K. Maeda, O. Hikosaka, The Caudal Part of Putamen
942   Represents the Historical Object Value Information. *J Neurosci* **39**, 1709-
943   1719 (2019).

944   58.   J. Kunimatsu, S. Yamamoto, K. Maeda, O. Hikosaka, Environment-based
945   object values learned by local network in the striatum tail. *Proc Natl Acad*
946   *Sci U S A* **118**, (2021).

947   59.   H. Yamada *et al.*, Coding of the long-term value of multiple future rewards
948   in the primate striatum. *J Neurophysiol* **109**, 1140-1151 (2013).

949   60.   H. Yamada, K. Louie, A. Tymula, P. W. Glimcher, Free choice shapes
950   normalized value signals in medial orbitofrontal cortex. *Nat Commun* **9**,
951   162 (2018).

952   61.   X. Chen, V. Stuphorn, Sequential selection of economic good and action

953   in medial frontal cortex of macaques during value-based decisions. *Elife* **4**,

954   (2015).

955   62.   H. Chen, Y. Naya, Forward Processing of Object-Location Association

956   from the Ventral Stream to Medial Temporal Lobe in Nonhuman Primates.

957   *Cereb Cortex* **30**, 1260-1271 (2020).

958   63.   Y. Hori *et al.*, Single caudate neurons encode temporally discounted value

959   for formulating motivation for action. *Elife* **10**, (2021).

960   64.   K. Burnham, D. Anderson, Multimodel inference: understanding AIC and

961   BIC in model selection. *Sociol. Method Res.* **33**, 261–304 (2004).

962   65.   B. Efron, R. J. Tibshirani, *An Introduction to the Bootstrap*. (Chapman &

963   Hall/CRC, 1993).

Fig. 1. Neural population geometries in the visual memory pathway

A, Summary of the neural population obtained from 10 brain regions in eight macaques during the four different behavioral tasks in Exp. 1 to 4. **B-E,** Rotational (**B**), curvy (**C**), straight (**D**), and unclear dynamics (**E**) were detected by visual inspection. In **A-E**, the 10 brain regions are numbered as follows: 1. TE, 2. STRt, 3. PRC, 4. CDb, 5. HPC, 6. VS, 7. cOFC, 8. CDh&b, 9. PHC, and 10. mOFC. The 0.05 s time bin was used for the dynamics analysis.

Fig. 2. Quantitative evaluation of geometric structures according to the rotational features.

A, Schematic depictions of the estimation of accumulated angle difference weighted by

the deviance, $\Sigma d\ \theta$. The accumulated angle difference indicates the degree of geometric

change in terms of the rotational force across time. Vector distance (d), rotational speed

($\theta/0.1s$), and start to endpoint distance (d_{S-E}) were also estimated. **B,** Dendrogram

estimated from these four parameter values based on bootstrap resampling across 10

neural populations. **C,** Percentage of variance explained by PCA of bootstrap resampling

data across 10 neural populations. **D,** Clusters detected among the four parameters based

on PCA. **E,** Percentage of clusters in each of the 10 brain regions. Each neural population

contained two components of neural information: the best and worst conditions (BW) in

Exps. 1 and 3, magnitude and delay of the rewards (MD) in Exp. 2, and magnitude and

probability of rewards (MP) in Exp. 4.

Fig. 3. Quantitative evaluation of geometric structures using the Lissajous curve function.

A, Schematic depictions of trajectory geometries using Lissajous function parameters. For all figures, *ωx* and *ωy* are 3 π. **B,** Three examples of bootstrap replicates for the HPC population fitted by the Lissajous function. *L* indicates the maximum loglikelihood. Estimated parameters were as follows: left, *ωx*, 2.78 π, *Φx*, -0.11 π, *L*, 34.6, *ωy*, 2.96 π, *Φy*, 0.10 π, *L*, 31.6; middle, *ωx*, 2.51 π, *Φx*, -0.03 π, *L*, 24.2, *ωy*, 3.82 π, *Φy*, -0.08 π, *L*,

32.7; right, ωx, 2.21 π, Φx, -0.16 π, L, 26.9, ωy, 2.78 π, Φy, 0.58 π, L, 29.1. **C,** Probability

density estimated for Lissajous parameters obtained from bootstrap replicates in each of

the 10 brain regions times two conditions. Ω and Φ are represented as 0 to 2 π. **D,**

Dendrogram estimated from Lissajous parameter values based on bootstrap resampling

across 10 neural populations times two conditions. **E,** Percentage of variance explained

by PCA of bootstrap resampling data across 10 neural populations times two conditions.

F, Clusters determined using PCA. Data are shown for PC1 to 3. **G,** Reconstructed

trajectory in each cluster based on bootstrap resampling. The trajectories in clusters 1–10

were drawn using the median values of the Lissajous parameters in each cluster. **H,**

Percentage of clusters in each of the 10 brain regions times two conditions. BW: best and

worst conditions. MD: magnitude and delay conditions. MP: magnitude and probability

conditions.

1007

1008

Fig. 4. Summary of the observed dynamics and anatomical connections in the visual

memory pathway.

A, Geometries depicted in the same arbitrary scales on the PC1-2 plane for the eight

neural populations shown in Fig. 1. The start of the trajectory (S) is aligned to describe the

trajectory. E indicates the end of the trajectory at 0.6 s. **B,** Proportion of the clusters defined

in each of the 10 brain regions using the Lissajous curve function. Reddish: rotational,

greenish: curvy, bluish: straight dynamics. Data from CDh&b and CDb were merged (CD).

1016

1017

Fig. S1. Behavioral tasks

A, Sequence of events during the ILR task in Exp. 1. The cue stimulus during the response

phase was the same as the sample stimulus during the encoding phase of the match trial,

whereas the two stimuli differed in the nonmatch trial. Neural activity was analyzed during

0.6 s after sample onset (gray bar). During the task, the monkeys actively responded or

did not respond by making a choice and performing the following action. **B,** Six visual

stimuli and spatial composition of the sample stimulus. **C,** Sequence of events during the

passive viewing task in Exp. 2. During fixation, the visual items were presented

sequentially within two to four times. These visual items were associated with reward or

no-reward outcomes in other behavioral contexts during the learning trials. During the task,

the monkeys were not required to respond, except for fixation to the center (passive).

Neural activity was analyzed during 0.4 s after object onset (gray bars). **D,** Eight visual

items were divided into two groups (items 1–4 and 5–8). In each scene (e.g., scene X),

one group was associated with a large reward and the other with a small reward. This

reward association was reversed in the other scene (e.g., scene Y). In each trial, two items

appeared as shown in **C**, one in items 1–4 and the other in items 5–8 as a random

combination. **E,** Sequence of events during the delayed reward task in Exp. 3. During the

task, the monkeys actively responded to the GO signal by releasing the lever (active).

Neural activity was analyzed during 0.6 s after cue onset (gray bar). **F,** During the task,

six visual items indicated the forthcoming reward size and delay duration to the reward

after the bar release. **G,** Sequence of events during the single-cue task in Exp. 4. A single

visual pie chart with green and blue segments was presented to the monkeys. During the

task, the monkeys were not required to respond, except for fixation to the center (passive).

Neural activity was analyzed during 0.6 s after cue onset. **H,** Payoff matrix: Each of the

magnitudes was fully crossed with each of the probabilities, resulting in a pool of 100

lotteries.

1044

1045

1046 **Fig. S2. Variances explained by PCA in each neural population.**

1047 Cumulative variance explained by PCA in each neural population in Exp. 1 to 4. The 0.05

1048 s time bin was used for the analysis. In Exp. 1 and Exp. 2, categorical task parameters

1049 were used. In Exp. 3 and Exp. 4, continuous task parameters were used.

1050

1051

Fig. S3. Variances explained by PCA under shuffle control.

Boxplot of variances explained by PCA under the three shuffled conditions (see Methods section for details). The plot was not cumulative. A boxplot was made with 1,000 repeats of the shuffle for each condition. Gray plots indicate the percentage of variance explained by PCA. Results using 0.05 s bin data are shown.

1057

Fig. S4. Examples of Lissajous curve represented with the parameter combination.

Rotational, curvy, and straight dynamics are shown against ω ratio and Φ difference. In

this study one cycle is defined with 3.33 π for ω during 0.6 s analysis period (3.33 π × 0.6

= 2 π in the Lissajous function, $A \cos(\omega\, t(i) + \Phi) + b$). Φ from 0 to 2 π defines the phase.

A and b denote the size and location of the curve, respectively. Combination of ω ratio and

Φ difference between x and y determines the shape of trajectory. For example, straight

geometry is defined as the same ω and the same Φ. Rotational dynamics is defined as

the same ω and not identical Φ. Curvy dynamics is defined as the different ω and the

same Φ.

1067

Supplementary Table S4. Summary of the data set and basic analysis.

NO	Experimental number	Brain region	Number of recorded neurons	Recording methods	Number of monkeys used	Behavioral task and standard analysis
1	Exp. 1.	TE	295	Single neuron recording /Multi neuron recording	2	Item-location-retention task Active behavior: match-nonmatch task ANOVA (6 visual stimuli, 4 locations)
2		Perirhinal cortex (PRC)	407			
3		parahippocampal cortex (PHC)	214			
4		Hippocampus (HPC)	590			
5	Exp. 2.	Striatum tail STRt	116	Single neuron recording	2	Scene-based object-value task Passive behavior ANOVA (8 visual stimuli, 2 rewards)
6	Exp. 3.	Caudate body (CDb)	150	Single neuron recording	3	Delayed reward tasks Active behavior: lever release response task Linear regression (delay, magnitude)
7	Exp. 4.	Medial orbitofrontal cortex (mOFC)	158	Single neuron recording	2	Cued lottery tasks Passive behavior Linear regression (probability, magnitude)
8		Central orbitofrontal cortex (cOFC)	190			
9		Ventral striatum (VS)	144			
10		Caudate head and body (CDh&d)	194			