Active control of mitochondrial network morphology by metabolism driven redox state

Gaurav Singh[a#], Vineeth Vengayil[a,e#], Aayushee Khanna[a], Swagata Adhikary[a] and Sunil Laxman[a*]

[a]Institute for Stem Cell Science and Regenerative Medicine, Bangalore, India.

[e]Present address: Perelman School of Medicine, University of Pennsylvania 3400 Civic Center Boulevard Building 421, Philadelphia, PA 19104.

#equal contribution

*Corresponding Author: Sunil Laxman
Address: Institute for Stem Cell Science and Regenerative Medicine (DBT-inStem) GKVK Post Bellary Road Bangalore 560065.
Email for correspondence: sunil@instem.res.in

Competing Interest Statement: No competing interests between the authors.

Classification: Biological Sciences/Cell Biology, Biochemistry/Systems biology

Keywords: mitochondria network, redox state, ROS.

Abstract

Mitochondria are dynamic organelles that change in morphology in different environments. What controls the mitochondrial morphology remains unresolved. Using yeast cells grown in distinct carbon environments requiring mitochondrial respiration, we find that mitochondrial morphology and activity are unrelated. Cells show fragmented as well as networked mitochondrial networks in nutrient environments with high or low mitochondrial activity. Resolving this paradox, we find that mitochondrial morphology is controlled by the intracellular redox state, which depends on the nature of electron entry into the Electron Transport Chain. In conditions with high direct electron entry, reactive oxygen species increase and the cytosolic environment becomes oxidized, leading to rapid mitochondrial fragmentation. Using genetic approaches to reduce electron injection into the ETC, we restore networked morphologies. Using pharmacologic agents to disrupt electron flow, or alter redox state, we demonstrate minute-scale, activity-independent control between networked and fragmented forms. Mitochondria thus actively control their morphology by regulating redox state.

Introduction

Mitochondria are double-membrane bound organelles at the center of energetics and cell state. They play a central role in ATP production through the functioning of the electron transport chain (ETC) and oxidative phosphorylation, coupled with the tricarboxylic acid (TCA) cycle - that also produces precursors for the synthesis of lipids, amino acids, and nucleotides (1–3). The TCA cycle supplies electron donors (NADH and FADH2) to ETC. Electrons flow through the enzyme complexes of the ETC, and generate the proton-motive force across the inner mitochondrial membrane, to drive the ATP synthase for ATP generation. Even as the mitochondria carries out electron transfer in the context of ATP generation, the process of electron transport through the respiratory chain is not completely leak-proof and results in generating reactive oxygen species (ROS), a byproduct of respiration (4, 5). Far from being static biochemical reactors for energetics, mitochondria are extremely dynamic organelles (6) that exhibit a remarkable diversity in form, ranging from a connected network of tubules to round, isolated fragments under different environments, or during cell cycle progression (7). Mitochondria undergo remodeling by fission and fusion to maintain their form. These processes are thought to allow mitochondria to adapt to various environments, or ensure optimal energy production (8, 9). Indeed, how mitochondrial networks respond to diverse environments have attracted intense scientific inquiry, since mitochondrial dysfunction is implicated in a wide range of diseases (10, 11).

However, we still lack an empirical understanding of a relationship between mitochondrial form and function. Assumptions have been made regarding the association of mitochondrial morphology with mitochondrial activity (12–14)). For example, associations have been made with interconnected mitochondrial networks and high respiratory activity, or fragmented mitochondria that have been observed in quiescent or inactive cells (10). Nonetheless, there are contrasting examples where mitochondrial form does not go hand in hand with activity. For example, Romani et al (15) find drastic changes in mitochondrial morphology occur without changes in respiration, or Yu et (16) find conditions where there is mitochondrial fragmentation without changes in activity. In examples from stem cells, different mitochondrial morphologies have been observed, with unclear relationships with bioenergetics or activity (17–19). It therefore remains unresolved as to the underlying basis for what triggers mitochondrial network remodeling.

In contrast, we understand the molecular machinery that regulates mitochondrial form. Mitochondrial morphology is regulated by a balance of fission and fusion processes, through specific proteins and mechanisms that are well conserved across eukaryotic cells, ranging from

yeast to mammals (20). Mitochondrial fission is spearheaded by large dynamin family GTPase, Drp1 (Dnm1 in yeast) that is recruited on mitochondrial membrane by accessory proteins (Mff, Mid49/51 and Fis1), and act in unison with ER and actin-myosin cytoskeleton to divide mitochondria (21). The fusion of outer and inner-mitochondrial membranes is facilitated by mitofusins, Mfn1 and Mfn2, (Fzo1 in yeast) and Opa1 (Mdm1 in yeast). The fission and fusion machinery are heavily regulated (22), but have predominantly been studied in the context of extreme starvation or mitochondrial damage, where mitochondrial quality control systems ensure hyper-fused mitochondria (23), or fragmented mitochondria (24). In summary, the regulators of the fission and fusion process in mitochondria (25) are well studied. However, the initiating signals (26) that determine mitochondrial form remain obscure.

In this study, we address what controls mitochondrial form by examining budding yeast (*S. cerevisiae*) grown in multiple defined carbon-sources, where the respective requirements for mitochondrial function and respiration are established (27). We quantitatively estimate the extent and nature of the mitochondrial networks in these cells, while assessing multiple indicators of mitochondrial outputs. Through this we find that mitochondrial form depends on the collective redox state of the cell, and is independent of activity. We demonstrate that in specific metabolic conditions there is increased direct electron entry into the ETC, which increases ROS. This ROS directly results in fragmented mitochondria. Genetic approaches to reducing direct electron entry results in fused mitochondria, as does decreasing ROS through chemical approaches. Orthogonally increasing ROS due to oxidants rapidly fragments mitochondria. In contrast, a reduced cytosolic environment - which is a consequence of the metabolic state of the cell - leads to networked mitochondria. This control of mitochondrial form is rapid (within minutes), dynamic, fully reversible, and independent of mitochondrial activity. Overall, we propose a systems-level chemical framework for how mitochondria control their morphology by regulating the redox state of the cell, and decouple mitochondrial form from activity. Our study opens direct lines of inquiry on the nature of redox signals that control mitochondrial form.

Results

Mitochondrial network morphology does not correlate with activity

Mitochondria have various morphologies that range from small, fragmented puncta to interconnected tubules found in different cells and conditions (28, 6, 29, 8, 30). These mitochondrial forms dynamically change in response to environmental factors (8, 12, 13, 15, 31, 32). It is currently unclear if mitochondrial form is related to mitochondrial function (Fig. 1A). In part, any correlations made are limited because observations coming from different cell/tissue types or even organisms, and cannot be directly compared. A practical lacuna has been the absence of studies using precisely manipulable systems where mitochondrial morphology can be predictably varied in controlled experimental settings. The 'model' laboratory eukaryotic cell, yeast (*Saccharomyces cerevisiae*), has been an ideal system to identify principles of mitochondrial function (33–38). Yeast cells are metabolically versatile, and grow on different carbon sources with varying requirements of mitochondrial respiration. In glucose, yeast repress their mitochondrial outputs via the Crabtree effect (39). In other carbon sources, yeast cells have higher requirements for mitochondrial respiration to sustain cellular energetics. Therefore, using yeast growing in different carbon sources, it would be possible to directly compare mitochondrial activity and morphology, in precisely defined settings.

We first established such an experimental and analytical pipeline where cells were grown in a defined carbon source, and where we quantitatively assess mitochondrial morphology, as well as activity (Fig 1B). We measured multiple parameters of mitochondrial activity - mitochondrial membrane potential, oxygen consumption, and amounts of a key respiratory complex IV protein Cox2 (Fig 1B), to obtain a composite understanding of activity. The carbon sources used were

glucose (repressed mitochondria), ethanol, glycerol, glycerol-ethanol, and lactic acid (all of which require mitochondrial respiration). We separately obtained high-resolution images of mitochondria, and estimated various network parameters. For accurate imaging and quantification of mitochondrial network, 3D confocal z-stacks were acquired that were then deconvolved. The 3D deconvolved stack is further processed for 3D multidimensional analysis of mitochondrial morphology in ImageJ. Supplementary methods and S1A-C provide detailed information on the imaging pipeline used, and addresses critical aspects of image deconvolution and 3D image processing to obtain clear separation of individual mitochondria, and accurate reconstructions of the mitochondrial networks.

Mitochondrial activity in yeast cells is dictated by how different carbons sources are metabolized. In glucose, cells repress mitochondrial activity (39). In sole carbon sources such as ethanol, lactate or glycerol, the repression due to glucose is released, and cells carry out mitochondrial respiration. We first estimated mitochondrial membrane potential in cells under all these conditions using a standard rosamine based potentiometric dye - Mitotracker red (39, 40). Expectedly, mitochondrial potential was low in glucose (Fig 1C). However, mitochondrial potential showed different extents of increase in glycerol, ethanol, glycerol-ethanol, or lactate (Fig 1C), indicating distinct membrane potentials. We next estimated relative amounts of Cox2 protein, as an indicator of the ETC (39, 41). Glucose had low Cox2 amounts, while a clear hierarchy of Cox2 protein was observed in the other carbon sources (Fig 1D). Finally, we measured basal oxygen consumption (using a Seahorse based assay (39, 42)) in cells in these conditions (Fig 1E). Consistent with the other observations, cells in glucose had very low oxygen consumption, while in each of the other conditions, oxygen consumption was high, but with a clear hierarchy (Fig 1E). Collectively, cells grown in lactate and glycerol-ethanol show the highest levels of mitochondrial activity, cells in ethanol have high activity, and cells growing in glycerol have relatively low activity, as summarized in Fig 1F.

We next asked what was the nature of mitochondrial networks in these conditions. Fig 1G shows the observed mitochondrial morphologies in each of these different carbon sources. We observe that cells in glucose have few but well-connected mitochondrial tubules. In contrast, cells grown in different respiratory carbon sources showed distinct mitochondrial networks (Fig 1G). The morphologies group into - 1) Completely fragmented mitochondria, characterized by short and round tubules as observed in glycerol and lactate, 2) Well networked mitochondria with long and interconnected mitochondrial tubules as observed in ethanol, and 3) Partially networked mitochondria with long connected tubules intermixed with shorter tubules as seen in glycerol-ethanol (Fig 1G). We carried out an unbiased, automated 3D quantification of mitochondrial network (Fig 1H-J) in all the carbon sources, to quantify the visual morphological features based on average branch length, number of branches and branch junctions per mitochondria, average mitochondrial sphericity and form factor (Fig 1H-J, Supplementary S1D and S1E, and see Material and Methods for details).

Summarizing these data, in these diverse media conditions for the same cell type, there is no correlation of mitochondrial morphology with mitochondrial activity. For example, cells grown in glycerol or lactate both show similar, highly fragmented mitochondria but have very low or very high mitochondrial activities respectively. Contrastingly, cells grown in ethanol and glucose both have long, highly networked mitochondrial tubules but high or low mitochondrial activities. Cells growing in glucose or glycerol have low activities, but networked or fragmented mitochondria respectively.

Therefore, in cells growing in a range of commonly occurring growth environments, mitochondrial morphology does not correlate with activity (Fig 1K and S1F).

Fig 1. **Mitochondrial morphology does not correlate with activity in different nutrient environments.** (A) Schematic asking if mitochondrial network morphologies and activities are interconnected. (B) Experiment design to investigate this relationship that involves growing cells in five different carbon sources as indicated, measuring mitochondrial activity using

standardized assays (C-E), and quantifying mitochondrial morphology (G-J). Measurement of multiple measures of mitochondrial activity in different carbon sources:(C) Average Mitotracker fluorescence that reports on mitochondrial membrane potential, measured by single cell imaging. Data represents average intensity per pixel for >100 cells from three independent experiments (mean ±SEM). (D) ETC complex IV subunit Cox2 protein levels measured by western blot using an anti-Cox2 antibody. A representative blot (out of 3 biological replicates, n=3) and their quantifications are shown, and (E) Basal oxygen consumption rate (OCR) corresponding to ~3 x 10^5 cells, from three independent experiments (n=3), normalized to the OD_{600}. (F) Schematic summary of mitochondrial activity of cells grown in different carbon sources. All respiratory carbon sources (glycerol, glycerol ethanol, ethanol and lactate) show higher baseline mitochondrial activity as compared to cells grown in glucose. However, there is considerable heterogeneity in morphology and mitochondrial activities even amongst various respiratory carbon sources. (G) Representative microscopy images of cells with mitochondria targeted with mNeonGreen. Top panel shows the maximum intensity projection of raw 3D z-stack and bottom panel shows 3D renderings of the z-stack after deconvolution. (H-J) Quantitative 3D analysis and comparison of mitochondrial network morphology across different carbon sources. Data represents analysis of more than 200 cells from three independent experiments (mean ±SEM). Scale bar - 2μm. (K) Schematic summary of mitochondrial activity versus respective mitochondrial morphology for cells grown in different carbon sources. Data represents mean ± SD unless otherwise stated. *P < 0.05, **P < 0.01, and ***P< 0.001; n.s., non-significant difference, calculated using unpaired Student's *t* tests.

Cellular redox state correlates with mitochondrial morphology

If mitochondrial activity does not correlate with its shape and network, what might explain the distinct morphological states observed in different metabolic conditions? To ask this question, we wondered what other mitochondrial outputs might be relevant. It is well established that mitochondrial outputs contribute to changing the overall cellular redox state, via the production of reactive oxygen species (ROS) that diffuse out of mitochondria (43–45). Interestingly, it is not as clear under what contexts does mitochondrial activity result in increased ROS. We therefore wondered whether cells growing in distinct carbon sources (where mitochondrial morphologies are very different) had different intracellular redox environments (Fig. 2A). In order to monitor differences in intracellular oxidation levels of cells grown in distinct carbon sources, we first used a well-established dye-based fluorescence assay (DCFDA fluorescence) as a quantitative readout. DCFDA is a cell-permeable non-fluorescent dye that becomes fluorescent upon oxidation inside cells, and is widely used to measure differences in intracellular redox states (15, 16, 46). We observed that cells grown in glycerol and lactate show high DCFDA fluorescence - consistent with a more oxidized intracellular environment - as compared to cells grown in ethanol, ethanol-glycerol or glucose (Fig 2B). This data suggested that cells in glycerol and lactate have a more oxidized cellular state. Contrastingly, cells grown in ethanol, glycerol-ethanol or glucose all exhibited lower DCFDA fluorescence - which would be consistent with a more reduced intracellular environment (Fig 2B). As a control, we supplemented ethanol-grown cells with hydrogen-peroxide (an oxidizing agent), which expectedly led to an immediate increase in DCFDA fluorescence (Fig 2B).

Since oxidation states cannot be interpreted based solely on DCFDA fluorescence, in order to have a quantitative readout of cellular oxidation levels, we next measured relative amounts of total glutathione (GSH) in cells in each of these conditions using mass-spectrometry. GSH is a primary redox buffer in cells, functioning as an antioxidant against ROS. Therefore, higher glutathione amounts would be expected in more oxidizing conditions, as cells counter this environment (15, 47, 48). Notably, we observed significantly higher GSH amounts in lactate and glycerol (Fig 2C) as compared to cells growing in ethanol, ethanol-glycerol, or glucose - suggesting that cells experience a relatively more oxidizing environment in lactate and glycerol. However, DCFDA fluorescence and total GSH amounts are reflections of gross intracellular redox environments. We therefore devised a biochemical assay to directly assess the relative oxidative state of the cytosolic or mitochondrial environment in the different carbon sources. For this, we utilized a maleimide-PEG (49) (polyethylene glycol) based assay (50) to biochemically measure the relative oxidation status of cysteine residues present on select proteins which are located either inside the mitochondrial matrix (isocitrate dehydrogenase - Idh2), or on the mitochondrial surface facing/exposed to the

cytosol (Tom70), or cytosolic GAPDH. The maleimide-PEG will covalently conjugate only to reduced (and not oxidized) cysteine residues, and result in an added mass on the protein that can be observed via mobility changes on a denaturing SDS-PAGE gel. This assay and its expected outputs are illustrated in the schematic diagram (Fig 2D). Through these experiments, we observed that the cytosol facing Tom70 had more reduced cysteines (i.e. more cysteines that can react with the added maleimide-PEG) in cells grown in ethanol or ethanol-glycerol, compared to cells grown in lactate or glycerol (Fig 2E). In contrast, Idh2 (which is present inside the mitochondrial matrix) showed no significant changes in cysteine oxidation across ethanol, ethanol-glycerol, or lactate (Fig 2F). Finally, cytosolic GAPDH showed more reduced cysteines in cells grown in ethanol or ethanol-glycerol, compared to lactate or glycerol (Supplementary S2). Through these data, we collectively conclude that cells growing in ethanol or glycerol-ethanol have a more reduced intracellular environment, while cells in glycerol or lactate have a more oxidized environment.

With these observations, we can now plot relative oxidative states with the mitochondrial morphology (Fig 2G). We observe that mitochondrial morphology in different metabolic conditions correlates with intracellular redox environment. In metabolic conditions where intracellular environment is more oxidized, mitochondria show fragmented morphology whereas metabolic conditions that have a relatively reduced intracellular environment exhibit more networked mitochondria.

Fig 2. Relationship of cellular oxidative state and mitochondrial morphology. (A) Schematic showing whether respiration dependent changes in cellular redox state can explain changes in mitochondrial morphology. (B) The total cellular redox state (measured by DCFDA fluorescence) in cells grown in different carbon sources. Cells were grown in different media as described before and the relative DCFDA fluorescence was measured (see materials and methods). (C) The relative total cellular GSH levels in cells grown in different carbon sources. Cells were grown in different media as described and GSH levels were measured by LC-MS/MS. (D) Schematic showing maleimide-PEG (mPEG) assay for protein redox state estimation. (E) Cytosolic oxidative state in different carbon sources. Cells were grown in different media as described and oxidation state of the cytosol was measured by electrophoretic mobility shift of mPEG bound to cytosolic facing Tom70, in a western blot (using anti-FLAG antibody). (F) Mitochondrial oxidative state in different carbon sources. Cells were grown in different media as described and oxidation state of mitochondria was measured by electrophoretic mobility shift of mPEG bound to mitochondrial Idh2, in a western blot (using anti-FLAG antibody). Data represents mean ± SD from three biological replicates (n=3) for all the experiments. (G) Schematic summary of how mitochondrial morphology correlates with cellular redox state in cells grown in different carbon sources with varied mitochondrial activities. *P < 0.05, **P < 0.01, and ***P< 0.001; n.s., non-significant difference, calculated using unpaired Student's *t* tests for data in (A) and (B) and paired Student's *t* tests for data in (E) and (F).

Mitochondrial morphology is reversibly and rapidly controlled by redox state

From our previous observations, the mitochondrial morphology correlated well with the intracellular redox state, where a more oxidized intracellular environment had fragmented mitochondria. Therefore, we wondered if it is possible to alter mitochondrial morphology in a predictable fashion (Fig 3A), solely by manipulating the redox environment?

To address this question, first we used cells growing in lactate, which is characterized by an oxidized state and fragmented mitochondria. Upon introducing a strong reductant - glutathione or ascorbic acid - to these cells growing in lactate, within ~30 minutes we observed a rapid transition to a fully networked mitochondrial morphology (Fig 3B). In these cells we quantified morphology, intracellular redox state and mitochondrial potential (as a proxy for activity). Notably, we observed that upon the addition of the reductant, the mitochondria rapidly (within 30 min) became networked, the intracellular environment became more reduced, although the mitochondrial activity decreased (Fig 3C). In a complementary experiment, we cultured cells in ethanol (reduced intracellular environment, mitochondria are highly networked). Adding a strong oxidant (H_2O_2) to these cells caused their mitochondria to fragment within 30 minutes (Fig 3D). When we subsequently added a strong reductant (glutathione or ascorbate) to these oxidant-treated cells, the mitochondrial network morphology was fully restored within 30 minutes (Fig 3D, Fig S3). We measured the same parameters: morphology, redox state, and activity. The oxidant induced mitochondrial fragmentation and an oxidized intracellular environment, which were both reversed by the reductant (Fig 3E). Notably, even while the mitochondrial morphology responded to changes in redox state, the mitochondrial activity consistently decreased and did not correlate with changes in morphology (Fig 3E).

Concluding, we find that the mitochondrial network morphology in any given environment can be reversibly controlled by changing the intracellular redox environment. Mitochondrial networks fragment in a more oxidized environment and vice versa. These changes in morphology do not correlate with mitochondrial activity. Instead, the intracellular redox state serves as a reliable correlate for mitochondrial morphology.

Fig 3. Mitochondrial network morphology changes in response to redox perturbations. (A) Schematic showing whether mitochondrial morphology can be reversibly modulated by changing redox state upon addition of reductants or oxidants. (B) Cells grown in lactate upon incubation with Glutathione (GSH) for 30 mins show a reversal in mitochondrial morphology from fragmented to networked (quantified in (C)). (D) Cells grown in ethanol are first incubated with H_2O_2 for 30 mins and then further incubated with GSH for 30 min. Addition of H_2O_2 fragments mitochondria that is then restored upon further addition of GSH. For data shown in panels (B) and (D), changes in mitochondrial morphology, redox state and activity are quantified in (C) and (E) respectively. Mitochondrial Morphology is assessed by quantifying average mitochondrial length per mitochondria, redox state is measured by evaluating changes in DCFDA or DHE fluorescence and Mitotracker fluorescence is used as a proxy for measuring changes in activity. Data represents mean ±SEM (see material and methods for details). Scale bar - 2µm. *P < 0.05, **P < 0.01, and ***P< 0.001; n.s., non-significant difference, calculated using unpaired Student's *t* tests.

Metabolic contexts resulting in direct electron entry have fragmented mitochondria

Our previous observations now raise the question - in different nutrient environments (all of which require mitochondrial activity), what might be a reason for this altered oxidative state. Why might there be a more oxidized cytoplasm in some conditions, and this need not correlate with 'activity'? We therefore reasoned that there might be intrinsic differences in how each carbon source is metabolized by the mitochondria, and lead to changes in ROS. To interrogate this hypothesis, we asked how each of these carbon sources is metabolized by yeast - and how the electron transport chain functions in these different contexts. In the schematic shown in Fig 4A, we summarize how each of these carbon sources are metabolized, with the ETC function placed in context. We observe contrasting processes of how each of these carbon sources drive the ETC (Fig 4A, Fig 4B). Ethanol is a 'classical' carbon source, and when metabolized enters and drives the TCA cycle, from where the reducing equivalents (NADH/FADH2) are produced to drive the ETC (Fig 4A). In contrast, lactate is metabolized by a lactate dehydrogenase (LDH) enzyme first. In *S. cerevisiae*, the primary yeast lactate dehydrogenase - Cyb2 (27, 51, 52) (L-lactate cytochrome-c oxidoreductase) is unusual in that the electrons are not donated to NAD, but to ferricytochrome c, and thus this electron directly enters the respiratory chain via cytochrome c (Fig 4A). Similarly, glycerol is metabolized by the glycerol-3-phosphate dehydrogenase (GPDH enzyme)) (53, 54). The primary yeast GPDH enzyme is Gut2, and this enzyme directly injects an electron into the ETC via a ubiquinone (Fig 4A). Therefore, in yeast, both lactate and glycerol metabolism drive mitochondrial function via direct electron injection into the ETC, via a cytochrome or a ubiquinone (by bypassing complex 1 and/or 2, might be worth mentioning to completely state the point). Both these enzymes are present on the outer leaflet of the inner-mitochondrial membrane. Furthermore, the binding site/pocket of these enzymes for redox acceptors (quinone or cytochrome c) might not be as well protected and shielded as compared to canonical ETC complex I and complex II where the quinone binding pocket is located much deeper. Therefore, we hypothesized that this direct injection of electrons into the ETC by either of these enzymes would result in more electrons interacting with oxygen, leading to higher ROS generation (Fig 4A and Fig 4B). Such mechanisms have been postulated previously (53, 55, 56). We can now observe that in ethanol or glucose (tubular/networked mitochondria) the reducing equivalents produced via the TCA cycle drive the ETC by conventional electron transfer to complex I and complex II (Fig 4B). In contrast, in lactate or glycerol (fragmented mitochondria), there is direct electron injection into the ETC during the metabolism of these carbon sources (Fig 4B).

If differences in direct electron entry determined the mitochondrial morphology, preventing/reducing this direct electron injection (using mutant cells lacking the major forms of the LDH or GPDH enzymes) should restore networked mitochondria. To test this, we first compared the mitochondrial morphologies in WT and Δ*cyb2* cells (note: see methods for growth and experiment details,) in a lactate medium. WT cells showed fragmented mitochondria, as described earlier. In contrast, Δ*cyb2* cells showed a well-networked mitochondrial morphology (Fig 4C). Next, we tested the effect of the loss of Gut2 in yeast cells in glycerol. Note: this is a challenging experiment, so please see methods for growth details. WT cells showed the expected fragmented mitochondria. In contrast, Δ*gut2* cells showed a well-networked mitochondrial morphology (Fig 4D). If this restored network were due to reduced ROS production (from reduced direct electron transfer), the intracellular redox state in these mutants would also be expected to become more reduced. We therefore assessed intracellular redox states in these cells (as compared to WT cells). Consistently, both Δ*cyb2* and Δ*gut2* cells showed a substantially more reduced intracellular state (Fig 4E). Collectively, these data suggest that preventing direct electron entry into ETC will result in more networked mitochondria (Fig 4F).

LDH and mGPDH mutants shows reversal in morphology

Fig 4: Networked mitochondria and reduced environments are restored in LDH and G3PDH mutants. (A) Schematic showing differences in electron injection into ETC when cells are grown on different carbon sources. (B) shows classification of the mode of electron entry into ETC for different carbon sources. Quantification of mitochondrial network morphology in (C) L-lactate dehydrogenase (Cyb2) deletion mutants grown in L-Lactate medium, and (D) glycerol-3- phosphate dehydrogenase (Gut2) deletion mutants grown in glycerol medium. Data represents mean ± SEM. See supplementary

methods for details. (F) The redox state in Cyb2 and Gut2 deletion mutants was estimated by changes in DCFDA fluorescence intensity. Data represent mean ± SD from three biological replicates (n=3). *P < 0.05, **P < 0.01, and ***P< 0.001; n.s., non-significant difference, calculated using unpaired Student's t tests.

Altering electron flow to change ROS controls mitochondrial morphology

Our data suggested that if electron flow, and thereby electron accumulation is modulated, the form of the mitochondrial network could be reversibly controlled. Experimentally, this could be addressed by using a (carefully titrated) inhibition of different ETC complexes. When done carefully and without completely ablating mitochondrial activity, this would result in blocked electron flow, alongside an accumulation/increased release of electrons. The accumulation of electrons should increase ROS. An increase in ROS (regardless of activity) should lead to fragmented mitochondria.

To test this, we selected three distinct inhibitors of electron entry into different complexes of the ETC, which function in yeast. These are: sodium azide (which inhibits complex IV), myxothiazol (which inhibits complex III), and carboxin (which inhibits complex II) (Fig 5A) (57–61). Sodium azide binds heme and copper sites in complex IV to block electron transfer from cytochrome c to complex IV. Myxothiozol blocks electron transfer from ubiquinol to the Riske-iron sulfur complex in complex III. Carboxin binds to the quinone–reduction site of complex II, thereby also preventing electron transfer (Fig 5A). Note that a careful titration of each of these inhibitors had to be first established; in order to optimize the concentrations for this experiment in prototrophic yeast cells (Extensive details are provided in the methods section and Fig. S5). In cells growing in ethanol with well-networked mitochondria, we first added sodium azide. Addition of sodium azide, within 30 min led to fragmented mitochondria, along with a concurrent drop-in mitochondrial activity (Fig 4B). To these cells, we next added a combination of strong reductants (ascorbic acid and mitoTEMPO). Within 30 min, we now observed a complete recovery and restoration of the mitochondrial network (Fig 5B). Notably, the addition of the inhibitor resulted in an increase in ROS, and this increase in intracellular oxidized state was rescued upon adding the reductants. However, the mitochondrial activity that reduced upon addition of sodium azide did not recover upon addition of the reductants (Fig 5B).

We next performed similar experiments with myxothiazol (Fig 5C) and carboxin (Fig 5D), which respectively inhibit complex III and II. In each of these cases, the addition of the inhibitor resulted in a highly fragmented mitochondrial morphology (Fig 5C and 5D). Upon the addition of reductants (ascorbate and mitoTEMPO) this fragmented morphology was restored to a networked morphology. Correspondingly, upon the addition of the respective inhibitors, we also observed an increase in intracellular ROS (Fig 5C and Fig 5D), which is rescued upon the addition of the strong reductants. Note also that there was no correlation of mitochondrial morphology with activity in these contexts.

Summarizing, by controlling the nature of electron flow it is possible to reversibly regulate the mitochondrial network morphology (Fig 5E). Blocking electron flow using inhibitors (of different ETC complexes) results in fragmented mitochondria and increased ROS. If this ROS is scavenged by strong reductants, the same cells will restore their mitochondrial network morphology. Notably, the changes in mitochondrial morphologies observed in the experimental conditions did not correlate with the mitochondrial activity (Fig 5B,5C&5D.)

Fig. 5. Effect of altering the nature of electron flow on mitochondrial morphology. (A, B) Schematic showing how different ETC inhibitors (classified in B) might generate ROS. (C, D, E) Changes in mitochondrial morphology for cells grown in ethanol upon addition of sodium azide (complex IV inhibitor), myxothiazol (complex III inhibitor), and carboxin (complex II inhibitor) respectively. Mitochondrial morphology is restored upon incubation of cells with an antioxidant cocktail of ascorbate (Vit C) and mitoTEMPO (mitochondria targeted superoxide scavenger). Mitochondrial morphology is assessed

by quantifying average mitochondrial length per mitochondria, redox state is measured by evaluating changes in DCFDA or DHE fluorescence (Fig S4) and Mitotracker fluorescence is used as a proxy for measuring changes in activity. Data represents mean ±SEM. Scale bar - 2µm. (E) An illustrative model proposing redox state as an integrated systems level biochemical signal to regulate mitochondrial network morphology. *P < 0.05, **P < 0.01, and ***P< 0.001; n.s., non-significant difference, calculated using unpaired Student's *t* tests.

Discussion

Mitochondria are dynamic organelles that transition between networked and fragmented forms. We used yeast growing in distinct metabolic conditions to unambiguously assess what controls mitochondrial form (networked/fragmented). Our data reveal that mitochondrial form is controlled by the intracellular redox state, which depends on the metabolic state of the cell. A synthesis of these results is summarized in **Fig 5D**, suggesting a systems-level chemical basis of how mitochondrial form is regulated based on intracellular redox state. The intracellular redox state is determined by ROS sources, primarily produced by the mitochondria itself - but which can also come from other sources. The amount of ROS produced depends on the metabolic state, which is determined by the nature of metabolism of the available carbon sources. Based on how certain carbon sources are metabolized, the nature of electron entry and therefore electron flow through the ETC changes. In some metabolic environments, there is increased direct electron entry/injection into the ETC, as seen with the yeast LDH or GPDH during growth in lactate or glycerol. When this happens, there is high ROS generation. As a consequence, there is an oxidized intracellular environment, which results in fragmented mitochondria. If direct electron entry into the ETC is limited (through genetic/pharmacologic means), there is reduced ROS, and a more networked-mitochondria. This explains why cells growing in different carbon sources all of which require mitochondrial respiration, can show highly networked or fragmented mitochondria. In contexts requiring high direct electron transfer that leads to ROS generation (e.g. lactate, glycerol), cells have fragmented mitochondria. Importantly, this ROS-dependent control of mitochondrial form is rapidly reversed (in minutes) by altering only the redox state. For example, reducing the intracellular environment in cells growing in lactate will network the mitochondria, or adding oxidants will fragment mitochondria in cells growing in ethanol.

Our study reiterates how electron flow through the ETC is not unidimensional, illustrating the underappreciated versatility of the ETC machinery. In a 'text-book' scenario where substrates fuel the TCA cycle, the TCA cycle generates reducing equivalents that supply the electrons to drive the ETC. In these cases, electron flow is tightly controlled and with minimal leakage of electrons (and therefore minimal reactive oxygen species generated). In contrast, when cells grow on other carbon sources that require complex metabolism that is coupled with ubiquinone or cytochrome c, electrons enter ETC through direct electron injection (62), leading to increased ROS. In such contexts, the mitochondria will be fragmented, independent of mitochondrial 'activity'. Mechanisms of direct electron injection/entry remain poorly understood. However, there is evidence of its common occurrence in different growth conditions. In addition to examples from this study (the yeast lactate dehydrogenase, or GPDH), such mechanisms have been observed ubiquitously across species. There are enzymes that transfer electrons directly to ubiquinone pool including Proline dehydrogenase (PRODH) involved in proline catabolism, mitochondrial GPDH or GPD2, Dihydro-orotate dehydrogenase (DHOODH) associated with pyrimidine synthesis, Electron transfer flavoprotein:ubiquinone oxidoreductase (ETF: QO) that transfers electrons from beta-oxidation of fats to ubiquinone (53, 63), and these contexts have high ROS generation (64, 62).

Why might such 'direct' electron entry mechanisms exist in the mitochondria? A speculative argument would be that this increases the versatility of mitochondrial functions, to operate in a range of carbon sources that would otherwise not easily be usable. In contexts where the activity TCA cycle is reduced, electron acceptors such as ubiquinone or cytochrome c can receive

electrons from other substrates that can be oxidized easily. This would allow the ETC to continue to function, and sustain the energetic and redox needs of the cell. This would result in metabolically versatile cells that are capable of growing in diverse carbon/nutrient environments, as opposed to specialized cells that grow in a preferred carbon source. In the context of mitochondrial morphology, any mechanism that increases the oxidative environment of the cytosol will lead to the fragmentation of mitochondria. This means exogenous generation of ROS through oxidizing agents, or atypical oxidation of fatty acids (65) or unchecked peroxisomal activity (66) could all result in a fragmented mitochondrial morphology. Such an interpretation can now explain observations from mutants or disease conditions where fragmented mitochondria are observed – without a change in mitochondrial activity (15, 16, 29). Here, it is likely that the intracellular state is more oxidized due to perturbed electron flow from direct electron injection, and/or exogenous ROS generating species. Finally, mutations or inhibition of ETC complexes that block electron flow would also result in an accumulation of electrons, leading to ROS generation through one-electron reduction of molecular oxygen to superoxide, or indirectly by facilitating reverse electron transfer, leading to the fragmentation of the mitochondria (as we exploit in this study). In these contexts, the activity of mitochondria might be compromised (e.g. complex IV inhibition using sodium azide) or remain unchanged (e.g. complex II inhibition using carboxin). Regardless, all these conditions will result in fragmented mitochondria due to increased intracellular oxidation. In any of these conditions, if the intracellular environment can be reduced (for example using reductants such as ascorbic acid, or mitochondria targeted mitoTEMPO) the mitochondrial morphology will restore to a networked form, despite the reduced mitochondrial activity due to the inhibitors. The metabolic versatility and ability of yeast to metabolize different carbon sources was key to be able to dissect relationships of metabolic (redox) state to mitochondrial form. In a metabolically specialized cell, it would be impossible to dissect out how mitochondrial form might be regulated by direct electron entry vs other forms of electron entry that are supported by reducing equivalents derived from the TCA cycle, due to distinct metabolic outputs.

A prediction from this study is the likely existence of signaling mechanisms that sense redox state changes to regulate mitochondrial quality control mechanisms. Such a sensing system will communicate with effectors that determine final mitochondrial form must exist, including regulators of the dynamics of fission vs fusion, rates of mitochondrial turnover, recruitment of cytoskeletal determinants of mitochondrial motility, or all the above. Our study reconciles multiple studies that identify such mechanisms as regulators of mitochondrial form (67, 68, 16, 69), while delineating a path for future inquiry to identify upstream mechanisms that can regulate mitochondrial form based on cellular redox state.

Materials and Methods

The specifics of the media and growth conditions for different cells, mitochondrial imaging and analysis, mitochondrial activity and redox assays, and metabolic analysis are detailed in the Supplementary Materials and Methods section of the SI Appendix.

Cell culture and biochemical assays

A prototrophic CEN.PK strain of *Saccharomyces cerevisiae* was used in all the experiments. Gene deletions, chromosomal C-terminal tagged strains were generated by PCR mediated gene deletion/tagging(70). Mitochondria targeted mNeon strain (Mito-mNeon green) is described in (71). Media compositions and specific growth conditions are described in the Supplementary Materials and Methods section of the SI Appendix. The fluorescence intensity for plate reader based biochemical assays was measured using Thermo Varioskan LUX multimode plate reader.

Detailed protocol is described in SI Appendix. The fluorescence intensities for each sample were normalized using OD_{600} of that sample and relative fluorescence intensity was calculated and plotted.

Protein extraction and Western blotting

Total protein was extracted using trichloroacetic acid (TCA) precipitation and blots were quantified using FIJI software. Detailed protocol is described in SI Appendix.

Basal OCR measurement

Agilent Seahorse XFe24 analyzer was used for basal OCR measurements. The detailed methods are described in SI Appendix.

Estimation of GSH levels by LC-MS/MS

The steady state levels of GSH were estimated by quantitative LC-MS/MS methods as described previously (72). Detailed methodology is described in SI Appendix and peak area measurements are listed in the supplement file.The steady state levels of GSH were estimated by quantitative LC-MS/MS methods as described in (72). Detailed methodology is described in SI Appendix and peak area measurements are listed in the supplement file.

Fluorescence Imaging and Analysis

Fluorescence imaging experiments were performed on live cells on an inverted laser scanning confocal microscope. For all quantifications of mitochondrial morphology, cells containing mneonGreen tagged mitochondria were used. Mitochondrial membrane potential was measured by staining cells with Mitotracker CMXRos. Redox state was measured in live cells by evaluating changes in DCFDA or DHE fluorescence. Custom- written macros and plugins in ImageJ were used for image analysis. Please refer to SI Appendix for detailed imaging pipeline.

Acknowledgments

We acknowledge extensive use of the Central Imaging and Flow Facilities in the NCBS and inStem campus. GS acknowledges a Research Associate Postdoctoral fellowship (DBT-RA) from the Dept. of Biotechnology, Govt. of India. SL acknowledges funding support from a DBT-Wellcome Trust India Alliance Senior Fellowship (IA/S/21/2/505922).

References

1. J. W. Pelley, "7 - Citric Acid Cycle, Electron Transport Chain, and Oxidative Phosphorylation" in *Elsevier's Integrated Review Biochemistry (Second Edition)*, J. W. Pelley, Ed. (W.B. Saunders, 2012), pp. 57–65.

2. J. B. Spinelli, M. C. Haigis, The multifaceted contributions of mitochondria to cellular metabolism. *Nat. Cell Biol.* **20**, 745–754 (2018).

3. I. Martínez-Reyes, N. S. Chandel, Mitochondrial TCA cycle metabolites control physiology and disease. *Nat. Commun.* **11**, 1–11 (2020).

4. D. B. Zorov, M. Juhaszova, S. J. Sollott, Mitochondrial reactive oxygen species (ROS) and ROS-induced ROS release. *Physiol. Rev.* **94**, 909–950 (2014).

5. M. P. Murphy, How mitochondria produce reactive oxygen species. *Biochem. J.* **417**, 1–13 (2009).

6. J. R. Friedman, J. Nunnari, Mitochondrial form and function. *Nature* **505**, 335–343 (2014).

7. A. S. Monzel, J. A. Enríquez, M. Picard, Multifaceted mitochondria : moving mitochondrial science beyond function and dysfunction. **5** (2023).

8. V. Eisner, M. Picard, G. Hajnóczky, Mitochondrial dynamics in adaptive and maladaptive cellular stress responses. *Nat. Cell Biol.* **20**, 755–765 (2018).

9. M. Picard, O. S. Shirihai, B. J. Gentil, Y. Burelle, Mitochondrial morphology transitions and functions: Implications for retrograde signaling? *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **304** (2013).

10. J. Rahman, S. Rahman, Mitochondrial medicine in the omics era. *The Lancet* **391**, 2560–2574 (2018).

11. S. B. Vafai, V. K. Mootha, Mitochondrial disorders as windows into an ancient organelle. *Nature* **491**, 374–383 (2012).

12. B. Westermann, Bioenergetic role of mitochondrial fusion and fission. *Biochim. Biophys. Acta* **1817**, 1833–1838 (2012).

13. T. Wai, T. Langer, Mitochondrial Dynamics and Metabolic Regulation. *Trends Endocrinol. Metab.* **27**, 105–117 (2016).

14. M. Liesa, O. S. Shirihai, Mitochondrial Dynamics in the Regulation of Nutrient Utilization and Energy Expenditure. *Cell Metab.* **17**, 491–506 (2013).

15. P. Romani, *et al.*, Mitochondrial fission links ECM mechanotransduction to metabolic redox homeostasis and metastatic chemotherapy resistance. *Nat. Cell Biol.* **24**, 168–180 (2022).

16. T. Yu, J. L. Robotham, Y. Yoon, Increased production of reactive oxygen species in hyperglycemic conditions requires dynamic change of mitochondrial morphology. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 2653–2658 (2006).

17. W. Zhou, *et al.*, HIF1α induced switch from bivalent to exclusively glycolytic metabolism during ESC-to-EpiSC/hESC transition. *EMBO J.* **31**, 2103–2116 (2012).

18. E. Tsogtbaatar, C. Landin, K. Minter-Dykhouse, C. D. L. Folmes, Energy Metabolism Regulates Stem Cell Pluripotency. *Front. Cell Dev. Biol.* **8**, 1–16 (2020).

19. M. Khacho, *et al.*, Mitochondrial Dynamics Impacts Stem Cell Identity and Fate Decisions by Regulating a Nuclear Transcriptional Program. *Cell Stem Cell* **19**, 232–247 (2016).

20. A. R. Fenton, T. A. Jongens, E. L. F. Holzbaur, Mitochondrial dynamics: Shaping and remodeling an organelle network. *Curr. Opin. Cell Biol.* **68**, 28–36 (2021).

21. A. M. van der Bliek, Q. Shen, S. Kawajiri, Mechanisms of mitochondrial fission and fusion. *Cold Spring Harb. Perspect. Biol.* **5** (2013).

22. A. Pagliuso, P. Cossart, F. Stavru, The ever-growing complexity of the mitochondrial fission machinery. *Cell. Mol. Life Sci.* **75**, 355–374 (2018).

23. A. S. Rambold, B. Kostelecky, N. Elia, J. Lippincott-Schwartz, Tubular network formation protects mitochondria from autophagosomal degradation during nutrient starvation. *Proc. Natl. Acad. Sci.* **108**, 10190–10195 (2011).

24. G. Ashrafi, T. L. Schwarz, The pathways of mitophagy for quality control and clearance of mitochondria. *Cell Death Differ.* **20**, 31–42 (2013).

25. K. L. Cerveny, Y. Tamura, Z. Zhang, R. E. Jensen, H. Sesaki, Regulation of mitochondrial fusion and division. *Trends Cell Biol.* **17**, 563–569 (2007).

26. F. Scialò, D. J. Fernández-Ayala, A. Sanz, Role of Mitochondrial Reverse Electron Transport in ROS Signaling: Potential Roles in Health and Disease. *Front. Physiol.* **8** (2017).

27. S. de Vries, C. A. M. Marres, The mitochondrial respiratory chain of yeast. Structure and biosynthesis and the role in cellular metabolism. *BBA Rev. Bioenerg.* **895**, 205–239 (1987).

28. J. Bereiter-Hahn, M. Voth, Dynamics of Mitochondria in Living Cells: Shape Changes, Dislocation, Fusion, and Fission of Mitochondria. *Microsc. Res. Tech.* **27**, 198–219 (1994).

29. H. Chen, D. C. Chan, Mitochondrial Dynamics in Regulating the Unique Phenotypes of Cancer and Stem Cells. *Cell Metab.* **26**, 39–48 (2017).

30. I. Y. Benador, M. Veliova, M. Liesa, O. S. Shirihai, Mitochondria Bound to Lipid Droplets: Where Mitochondrial Dynamics Regulate Lipid Storage and Utilization. *Cell Metab.* **29**, 827–835 (2019).

31. P. Mishra, D. C. Chan, Mitochondrial dynamics and inheritance during cell division, development and disease. *Nat. Rev. Mol. Cell Biol.* **15**, 634–646 (2014).

32. M. Liesa, O. S. Shirihai, Mitochondrial dynamics in the regulation of nutrient utilization and energy expenditure. *Cell Metab.* **17**, 491–506 (2013).

33. J. M. Shaw, J. Nunnari, Mitochondrial dynamics and division in budding yeast. *Trends Cell Biol.* **12**, 178–184 (2002).

34. K. Okamoto, J. M. Shaw, Mitochondrial morphology and dynamics in yeast and multicellular eukaryotes. *Annu. Rev. Genet.* **39**, 503–536 (2005).

35. S. Meeusen, *et al.*, Mitochondrial Inner-Membrane Fusion and Crista Maintenance Requires the Dynamin-Related GTPase Mgm1. *Cell* **127**, 383–395 (2006).

36. A. Chacinska, C. M. Koehler, D. Milenkovic, T. Lithgow, N. Pfanner, Importing Mitochondrial Proteins: Machineries and Mechanisms. *Cell* **138**, 628–644 (2009).

37. J. R. Friedman, *et al.*, ER tubules mark sites of mitochondrial division. *Science* **334**, 358–362 (2011).

38. C. Malina, C. Larsson, J. Nielsen, Yeast mitochondria: an overview of mitochondrial biology and the potential of mitochondrial systems biology. *FEMS Yeast Res.* **18** (2018).

39. V. Vengayil, S. Niphadkar, S. Adhikary, S. Varahan, S. Laxman, The deubiquitinase Ubp3/Usp10 constrains glucose-mediated mitochondrial repression via phosphate budgeting. *eLife* **12** (2024).

40. B. Chazotte, Labeling mitochondria with mitotracker dyes. *Cold Spring Harb. Protoc.* **6**, 990–992 (2011).

41. F. Fontanesi, I. C. Soto, D. Horn, A. Barrientos, Assembly of mitochondrial cytochrome c-oxidase, a complicated and highly regulated cellular process. *Am. J. Physiol. Cell Physiol.* **291**, C1129-1147 (2006).

42. Gaurav Singh, Geen George, Sufi Raja, Akash Gulyani, A fluorescence lifetime-sensor reveals link between mitochondrial membrane microenvironment and activity. *Manuscr. Prep.* (2020).

43. C. C. Winterbourn, Reconciling the chemistry and biology of reactive oxygen species. *Nat. Chem. Biol.* **4**, 278–286 (2008).

44. C. Lennicke, H. M. Cochemé, Redox metabolism: ROS as specific molecular regulators of cell signaling and function. *Mol. Cell* **81**, 3691–3707 (2021).

45. D. Nolfi-Donegan, A. Braganza, S. Shiva, Mitochondrial electron transport chain: Oxidative phosphorylation, oxidant production, and methods of measurement. *Redox Biol.* **37**, 101674 (2020).

46. A. Wojtala, *et al.*, *Methods to monitor ROS production by fluorescence microscopy and fluorometry*, 1st Ed. (Elsevier Inc., 2014).

47. S. Chakravarthi, C. E. Jessop, N. J. Bulleid, The role of glutathione in disulphide bond formation and endoplasmic-reticulum-generated oxidative stress. *EMBO Rep.* **7**, 271–275 (2006).

48. S. C. Lu, Regulation of glutathione synthesis. *Mol. Aspects Med.* **30**, 42–59 (2009).

49. Y.-J. Lee, G.-D. Chang, Quantitative display of the redox status of proteins with maleimide-polyethylene glycol tagging. *Electrophoresis* **40**, 491–498 (2019).

50. L. Wrobel, A. M. Sokol, M. Chojnacka, A. Chacinska, The presence of disulfide bonds reveals an evolutionarily conserved mechanism involved in mitochondrial protein translocase assembly. *Sci. Rep.* **6**, 27484 (2016).

51. B. Turcotte, X. B. Liang, F. Robert, N. Soontorngun, Transcriptional regulation of nonfermentable carbon utilization in budding yeast. *FEMS Yeast Res.* **10**, 2–13 (2010).

52. S. Passarella, *et al.*, Mitochondria and l-lactate metabolism. *FEBS Lett.* **582**, 3569–3576 (2008).

53. T. Mráček, Z. Drahota, J. Houštěk, The function and the role of the mitochondrial glycerol-3-phosphate dehydrogenase in mammalian tissues. *Biochim. Biophys. Acta - Bioenerg.* **1827**, 401–410 (2013).

54. C. Larsson, *et al.*, The importance of the glycerol 3-phosphate shuttle during aerobic growth ofSaccharomyces cerevisiae. *Yeast* **14**, 347–357 (1998).

55. A. K. O. Boubacar, S. Pethe, J. P. Mahy, F. Lederer, Flavocytochrome b 2: Reactivity of its flavin with molecular oxygen. *Biochemistry* **46**, 13080–13088 (2007).

56. M. D. Brand, The sites and topology of mitochondrial superoxide production. *Exp. Gerontol.* **45**, 466–472 (2010).

57. C. M. Palmeira, A. J. Moreno, *Mitochondrial Bioenergetics: Methods and Protocols* (2012).

58. A. P. Wojtovich, C. O. Smith, C. M. Haynes, K. W. Nehrke, P. S. Brookes, Physiological consequences of complex II inhibition for aging, disease, and the mKATP channel. *Biochim. Biophys. Acta - Bioenerg.* **1827**, 598–611 (2013).

59. F. Bouillaud, Inhibition of Succinate Dehydrogenase by Pesticides (SDHIs) and Energy Metabolism. *Int. J. Mol. Sci.* **24** (2023).

60. A. Y. Andreyev, Y. E. Kushnareva, A. A. Starkov, Mitochondrial metabolism of reactive oxygen species. *Biokhimiya* **70**, 246–264 (2005).

61. C. L. Quinlan, I. V. Perevoschikova, R. L. S. Goncalves, M. Hey-Mogensen, M. D. Brand, *The determination and analysis of site-specific rates of mitochondrial reactive oxygen species production*, 1st Ed. (Copyright © 2013 Elsevier, Inc. All rights reserved., 2013).

62. C. L. Quinlan, I. V. Perevoshchikova, M. Hey-Mogensen, A. L. Orr, M. D. Brand, Sites of reactive oxygen species generation by mitochondria oxidizing different substrates. *Redox Biol.* **1**, 304–312 (2013).

63. M. Alcázar-Fabra, P. Navas, G. Brea-Calvo, Coenzyme Q biosynthesis and its role in the respiratory chain structure. *Biochim. Biophys. Acta BBA - Bioenerg.* **1857**, 1073–1078 (2016).

64. V. Raimondi, F. Ciccarese, V. Ciminale, Oncogenic pathways and the electron transport chain: a dangeROS liaison. *Br. J. Cancer* **122**, 168–181 (2020).

65. M. G. Rosca, *et al.*, Oxidation of Fatty Acids Is the Source of Increased Mitochondrial Reactive Oxygen Species Production in Kidney Cortical Tubules in Early Diabetes. *Diabetes* **61**, 2074–2083 (2012).

66. M. Fransen, M. Nordgren, B. Wang, O. Apanasets, Role of peroxisomes in ROS/RNS-metabolism: Implications for human disease. *Biochim. Biophys. Acta BBA - Mol. Basis Dis.* **1822**, 1363–1373 (2012).

67. B. Morant-Ferrando, *et al.*, Fatty acid oxidation organizes mitochondrial supercomplexes to sustain astrocytic ROS and cognition. *Nat. Metab.* **5**, 1290–1302 (2023).

68. Y. Mi, *et al.*, Loss of fatty acid degradation by astrocytic mitochondria triggers neuroinflammation and neurodegeneration. *Nat. Metab.* **5**, 445–465 (2023).

69. N. Baker, *et al.*, The mitochondrial protein OPA1 regulates the quiescent state of adult muscle stem cells. *Cell Stem Cell* **29**, 1315-1332.e9 (2022).

70. M. S. Longtine, *et al.*, Additional modules for versatile and economical PCR-based gene deletion and modification in Saccharomyces cerevisiae. *Yeast* **14**, 953–961 (1998).

71. N. Dua, A. Seshadri, A. Badrinarayanan, DarT-mediated mtDNA damage induces dynamic reorganization and selective segregation of mitochondria. *J. Cell Biol.* **221** (2022).

72. A. Walvekar, Z. Rashida, H. Maddali, S. Laxman, A versatile LC-MS/MS approach for comprehensive, quantitative analysis of central metabolic pathways. *Wellcome Open Res.* **3**, 122 (2018).