Molecular Tracking and Cultivation Reveal Ammonia Oxidizing Archaea as Integral Members of the Human Skin Microbiome

Alexander Mahnert*[,1], Maximilian Dreer*[,2], Ülkü Perier[2], Michael Melcher[2], Stefanie Duller[1], Adina Lehnen[1], Theodora Goessler[1], Daniela Brunner[1], Thomas Graier[3], Peter Wolf[3], Rafael I. Ponce-Toledo[2,#], Logan Hodgskiss[2], Melina Kerou[+,2], Christine Moissl-Eichinger[+,1,4], and Christa Schleper[+,2]

* Shared authorship
+ Shared corresponding authorship

[1] Medical University of Graz, D&R Institute for Hygiene, Microbiology, and Environmental Medicine, Graz, Austria
[2] University of Vienna, Archaea Biology and Ecogenomics Unit, Vienna, Austria
[3] Medical University of Graz, Department of Dermatology and Venereology, Graz, Austria
[4] BioTechMed Graz, Graz, Austria

Present address: Institut de Systématique, Evolution et Biodiversité (ISYEB), Sorbonne Université, CNRS, Muséum National d'Histoire Naturelle, Paris, France

Abstract

Ammonia oxidizing archaea (AOA) have been repeatedly detected in the human skin microbiome through molecular and biochemical analyses, yet their persistence, physiology, and adaptations remain poorly understood. Here, we describe two cultivated strains, *Candidatus* Nitrosocosmicus epidermidis and *Ca.* Nitrosocosmicus unguis, enriched from human skin samples. These autotrophic strains grow on ammonia and urea as sole energy sources. Genomic islands and expanded gene families in their genomes testify to the capacity of colonizing skin, including specific interactions with host proteins and signaling cascades that distinguish these AOA from their close relatives from soil.

Molecular signatures of *Nitrosocosmicus* ssp. were consistently identified on individuals in cross-sectional and longitudinal cohorts (n=47) with a higher prevalence in sebaceous areas. Co-occurrence network patterns with specific bacteria reinforce the observation that AOA of the genus *Nitrosocosmicus* form a stable component of the healthy skin and represent emerging commensals in the human microbiome.

Introduction

The human body is an ecosystem populated by spatially organized microbial communities composed of bacteria, archaea, viruses/ phages and small eukaryotes, with diverse metabolic activities and interactions: the microbiome [1]. In recent years, numerous studies have led to a better understanding of the mechanisms involved in the complex interplay between the host and its commensals and have revealed the fundamental role of the human microbiome in health and disease [2–4]. However, most studies on human microbial communities have focused exclusively on the bacterial microbiome, while other integral components, such as eukaryotes/fungi, archaea and viruses have remained largely overlooked [5–9]. A systematic metagenomic study has recently revealed the extent of diversity and abundance of archaea in the human gut microbiome [10], where the increased presence of methanogenic archaea affects short-chain fatty acid production and the overall host physiology [11].

The skin microbiome is equally complex and harbors various bacterial species, including *Staphylococcus, Streptococcus, Corynebacterium, and Propionibacterium* (now *Cutibacterium*), and fungi such as Malassezia [12,13]. The composition and diversity of the microbiome varies across different body sites (e.g. oily versus dry skin) and between individuals [14]. As in the gut, microorganisms play essential roles in maintaining skin health as they provide protection against pathogens by competing for nutrients and space, producing antimicrobial compounds and modulating the skin's immune response [15].

Human skin is also colonized by archaea [16,17], in particular by representatives of the ammonia oxidizing archaea (AOA), formerly Thaumarchaeota, now class Nitrososphaeria and phylum Thermoproteota [18–20]. Although rather rare in abundance, archaeal signals were found in human subjects analyzed by 16S rRNA gene sequencing and detection of archaeal *amoA*, the gene for subunit A of the key enzyme ammonia monooxygenase used for ammonia oxidation[16]. AOA-associated Nitrososphaerales signatures were detected in all human subjects and were closely related to those found earlier in cleanrooms and intensive care units [21]. Their presence was confirmed by fluorescence *in situ* hybridization [16], next generation sequencing, and infrared hyperspectral imaging [17]. Although the physiological relevance of AOA to the skin is still subject to speculation, its potential role is intriguing considering that urea and ammonia are components of human sweat and are the main energy sources of ammonia oxidizers. Also, nitric oxide (NO), an important signaling molecule involved in diverse processes in humans, is a byproduct of the ammonia oxidation process [22].

Detection of archaea in human microbiomes is generally hampered by numerous methodological obstacles, such as challenging cultivation [23,24], and molecular detection issues (summarized in [25]), because of the bacteria-centric orientation in sampling, cell lysis, DNA extraction protocols, PCR primers, and classification databases. Consequently, archaeal

signals often remain close to or below the detection limit, making it difficult to understand the

role of archaea on human skin [24]. Such obstacles and the lack of cultivated strains of skin AOA

have left many questions unanswered, with respect to i) abundance, ii) longitudinal stability,

iii) association with human health, iv) physiology and metabolism, v) interaction with the

bacterial microbiome, and vi) specific adaptations to the human skin.

In this manuscript, we present the results of extensive cultivation efforts and molecular-based

analyses targeting ammonia oxidizing archaea of healthy and diseased individuals. This

endeavor resulted in the stable and high enrichment of strains affiliated with the genus

Candidatus Nitrosocosmicus from human skin. Their genomes reveal genomic features that

could facilitate colonization of the skin environment, but also highlight the genomic repertoire

shared among all members of the genus *Ca*. Nitrosocosmicus, a widespread environmental

group of AOA. The signatures of Nitrosocosmicus could be tracked in different cohorts of

healthy volunteers and diseased patients, and correlations with physical parameters and with

bacterial occurrence patterns were identified.

Results

Cultivation of AOA from human skin

Different types of inocula were used to initiate enrichment cultures under aerobic conditions in the presence of ammonia as sole energy source. After several months, amplification and sequencing of archaeal 16S rRNA and *amoA* gene fragments identified the presence of one AOA strain per enrichment. The usage of pooled samples of the face, neck, upper body, back, arms, and legs of individuals resulted in three enrichment cultures: Nitrosocosmicus "T1S", "X2B", and "Z3A". A fourth enrichment culture was obtained from cut fingernail edges as initial inoculum ("R2S").

After repeated transfers over the course of more than a year in medium supplemented with 0.6 mM ammonium, 2 mM of either sodium carbonate (R2S) or sodium bicarbonate (T1S) and an antibiotic mix, two stable cultures displaying near stoichiometric conversion of ammonia to nitrite were obtained within approximately 3-4 weeks (Fig. 1, A-B), with a relative enrichment level of 88-98% based on molecular and microscopic assessments (Extended Data Fig. 1). No bacterial contaminants were detected via amplification of the bacterial 16S rRNA gene (Extended Data Fig. 2), or by light and scanning electron microscopy (Extended Data Fig. 1). Based on the sequenced internal transcribed spacer (ITS) region the remaining contaminant in both enrichments was a single fungal species with 99% sequence identity to *Exophiala* sp. DAOM 216391, a genus of fungi frequently found in manmade environments and known to be an opportunistic pathogen of humans [26,27].

Growth of the two strains, R2S and T1S, correlated with nitrite production in both enrichments, as has been shown previously for other AOA (Fig. 1, A-B) [28–30]. The shortest generation times based on nitrite production were approximately five days at 28°C. After the oxidation of ~500 µM ammonium enrichments R2S and T1S reached cell densities of $2.16*10^5$ ($\pm 1.68*10^4$) and $1.18*10^6$ ($\pm 1.05*10^5$) cells ml^{-1}, respectively. This is similar to the cell densities of *Nitrosocosmicus franklandianus* C13 ($3.8*10^6$ cells ml^{-1}) [31] and *Nitrosocosmicus oleophilus* MY3 ($1.6*10^6$ cells ml^{-1}) [32] after oxidation of ~500 µM ammonium.

In light microscopy, putative AOA were observed as cocci with diameters between 1-2 µm which frequently formed aggregates (Extended Data Fig. 1) and were partly enclosed in putative extracellular polymeric substances (EPS) (Extended Data Fig. 1). These morphological features are comparable with those of previously published members of the genus Nitrosocosmicus [30–33] and were confirmed by scanning electron microscopy (Fig. 1, C-F).

Fig. 1. | Growth and morphology of skin enrichments R2S (A) and T1S (B) displaying near stoichiometric conversion of ammonia (triangles, dashed line) to nitrite (circles, solid line) paralleled by cell growth, estimated by qPCR of archaeal *amoA* gene copies (orange bars). Ammonia and nitrite measurements are averages of biological triplicates. Gene copies of *amo*A are averages of technical triplicates of each biological triplicate. Error bars depict the standard deviation of the mean. Scanning electron micrographs of skin enrichments R2S (C, D) and T1S (E, F) with white arrows indicating AOA cells based on the morphologies seen in Supplementary Figure AOA2 and already published members of the Nitrosocosmicus [30–33]. Dashed white arrows indicate putative extra polymeric substances (D, F). Black arrows indicate the remaining fungal contaminant.

Taxonomic placement and genomic features of the AOA representatives from human skin

Sequencing and assembly of the genomes of the two skin enrichments resulted in one circular contiguous sequence affiliated with AOA for each strain. The genome sizes, G+C content and predicted protein-coding genes of the strains were well within range of other representatives of the genus *Nitrosocosmicus* (Table 1).

Like other representatives of this genus, both strains encoded three identical copies of the 16S and 23S rRNA gene operons, typically indicative of a capacity to respond rapidly to favorable environmental changes [34,35]. The average nucleotide identity (ANI) and average amino acid identity (AAI) values between strains R2S and T1S (87.4% ANI, 86.2% AAI), and between them and other *Nitrosocosmicus* genomes and proteomes were well below the threshold of 95% ANI/AAI proposed for species delineation [36] (Supplementary Tables S1, S2). We therefore propose that they represent two distinct novel species, preliminarily named *Ca.* Nitrosocosmicus unguis R2S (unguis = nail) and *Ca.* Nitrosocosmicus epidermidis T1S ("of the epidermis"), within the candidate genus Nitrosocosmicus*.*

168 **Table 1**: Genomic and growth features of *Ca.* Nitrosocosmicus epidermidis, *Ca.*
169 Nitrosocosmicus unguis compared to other cultivated representatives of the genus
170 *Nitrosocosmicus*

Organism#	Status	Growth temp (°C)	pH	Genome size (Mb)	No. of contigs	DNA GC %	protein coding genes	16S/23S/5S rRNA gene copies	reference
Nitrosocosmicus un-guis R2S	Enriched	28	7	3.21	1	33.77	3597	3/3/1	this study
Nitrosocosmicus epi-dermidis T1S	Enriched	28	7	3.07	1	33.55	3401	3/3/1	this study
N. oleophilus MY3	Pure	30	6.5-7	3.43	1	34.14	3722	3/3/1	Jung et al. 2016
N. franklandianus C13	Pure	40	7	2.84	1	34.07	3025	2/2/1	Lehtovirta-Morley et al. 2016
N. arcticus Kfb	Enriched	28	6	2.65*	23	34	2970	3/3/1	Alves et al. 2019
N. hydrocola G61	Enriched	33	8	2.99	1	33.94	3162	2/2/1	Sauder et al. 2017
N. agrestis SS	Enriched	37	6.5-7	3.22**	43	33.42	3524	2/2/1	Liu et al. 2021

all organisms have Candidatus status

* completeness and contamination values: 98.95%, 1.94%

** completeness and contamination values: 96.1%, 2.91%

A maximum likelihood analysis based on 36 concatenated ribosomal protein marker genes
present in one copy in at least 100 of the 143 genomes of a dataset comprising representative
AOA and non-AOA of Nitrososphaeria confirmed the affiliation of the two novel strains with
the NS-zeta lineage as defined by Alves et al. 2018 [37]. Specifically, the two strains formed a
highly supported cluster with N. arcticus (obtained from arctic soil), N. oleophilus (from coal-
tar contaminated sediment), and other uncharacterized MAGs (Fig 2).

Fig. 2. | Maximum-likelihood phylogenomic tree of AOA and non-AOA reconstructed based
on the concatenated alignment of 36 ribosomal proteins comprising a total of 4,707 amino acid
sites. Numbers in parentheses represent the number of complete genomes and MAGs in each
lineage. NS = Nitrososphaerales (A). Heatmap displaying the detection frequency of specific
functional domains discussed in the text in all analyzed *Ca.* Nitrosocosmicus genomes.
Numbers refer to the number of occurrences of the respective domain in the genome, but do
not always correspond do distinct CDS (i.e. multiple domains on one CDS). Domain IDs refer
to Interpro, Pfam. TIGRfam, Superfamily Database entries (B).

To investigate the specific genomic repertoire of *Ca.* N. unguis and *Ca.* N. epidermidis, we

reconstructed 3,301 orthologous protein families from a dataset including all available

genomes from cultivated strains of the genus *Nitrosocosmicus*: *Ca.* N. franklandianus C13,

Ca. N. oleophilus MY3, N. hydrocola G61, N. agrestis SS, and *Ca.* N. arcticus Kfb (we refrain

from the "Ca." prefix for better readability in the following) (Extended Data: Orthologs). From

these orthologous protein families, 1,592 represented the core genome of the genus,

comprising about 50% of the predicted CDS of the genomes. In line with their phylogenomic

placement, the N. epidermidis shared less protein families with N. oleophilus and N. arcticus

(60), while 125 families were shared between N. unguis and N. oleophilus. The content of

arCOG functional categories did not differ among the Nitrosocosmicus genomes, indicating

that presence and survival on human skin did not require a particular expansion in a functional

category (Extended Data Fig. 3). N. unguis and N. epidermidis shared only 20 protein families

to the exclusion of all others, while both of them shared more protein families with N. oleophilus

(125 and 22 respectively) than any other Nitrosocosmicus genome. The shell (=strain-specific)

proteome of each skin isolate consisted of 17 and 16 families for N. unguis and N. epidermidis

respectively. Below we discuss specific features of the two skin strains and of the genus

Nitrosocosmicus in general, that indicate adaptation to host surfaces and interactions.

A genomic island and expanded gene families implicate factors involved in host
interactions

The genome of N. epidermidis contains genomic islands flanked by integrases, with a range

of proteins potentially involved in host interaction (Extended Data: Annotations). One of these

islands (NET5MPC655_v1_2202- NET5MPC655_v1_2261), encodes a putative

autotransporter-1 (AT-1) family (1.B.12) protein (NET5MPC655_v1_2213) with predicted

extracellular proteins of unknown function. The C-terminus of autotransporter proteins has the

capacity to form β-barrel structures into the outer membrane in order to allow for the export of

diverse N-terminal virulence factors [38]. The N-terminal part of the N. epidermidis

autotransporter protein consists of leucine-rich repeats involved in protein–protein recognition

processes [39]. The island also encodes a putative α-arrestin -like protein

(NET5MPC655_v1_2245), matching both arrestin N-terminal (IPR011021) and arrestin C-

terminal (IPR014752) -like domains [40]. Arrestins are involved in binding and regulating

transmembrane receptors of the 7TM and kinase classes in eukaryotes (including GPCRs in

higher eukaryotes), with crucial physiological functions [40]. Although prokaryotic arrestin-like

proteins have not been characterized, there are examples of pathogens hijacking β-arrestin

signaling pathways of the host to cross epithelial/endothelial barriers [41]. The genomic island

also encodes eight copies of membrane attached serine/threonine protein kinases and four

putative serine-rich adhesins, the importance of which in facilitating binding to various tissue
types and extracellular matrix proteins in streptococci is well documented [42].

The second strain, N. unguis, encodes expanded gene families whose products are likely to
be involved in host interaction (Fig. 2B, Extended Data: Annotations). Among these are seven
cysteine-rich secretory protein (CAP) family proteins, which are only found among few
members of AOA of Nitrosocosmicus, with only one copy in N. epidermidis. Although present
in several bacterial lineages [43], this protein family has only been studied in eukaryotes so far,
with essential functions ranging from adhesion to signaling and regulation of ion channels, and
can be involved either in pathogenicity or in antimicrobial defense [44].

The genome of N. unguis is also more enriched in beta propeller fold- containing proteins
compared to N. epidermidis (Fig. 2B, Extended Data: Annotations) implying different
strategies to interact with the surrounding environment and communities. These proteins can
act as ligand-binding and signaling proteins or catalyze various hydrolytic or oxidoreductase
reactions in the periplasmic space, thereby mediating interactions with the environment, such
as the influenza virus neuraminidase [45]. They are generally found in higher numbers in
Nitrosocosmicus genomes compared to other AOA lineages (Fig. 2B).

A comprehensive description of the genomic repertoire of both strains, including more putative
adaptations to the skin environment can be found in the Supplementary Information and in
Extended Data: Annotations.

Members of the genus *Nitrosocosmicus* are equipped for the colonization of host
surfaces

Different cultivated Nitrosocosmicus strains have been shown to produce biofilms and exhibit
superior adhesion capacities compared to other AOA lineages [32,33], a trait also observed for
many bacterial species residing on the human skin [46]. This is mirrored in the proteomes of
Nitrosocosmicus ssp. which are enriched in acetamidosugar biosynthesis and adhesion-
related proteins, as well as being the only AOA clade encoding LPSE family
lipopolysaccharide exporters (Fig. 2B, Extended Data: Annotations) [32,33]. For example,
Nitrosocosmicus genomes specifically encode one family of extracellular serine-rich repeat
proteins with roles including adhesion and biofilm formation, with 35% sequence identity to
dentin sialoprotein-like proteins from eukaryotes [47].

Furthermore, Nitrosocosmicus genomes are characterized by the presence of a suite of gene
families whose products can directly interact with and modulate other organisms (Fig. 3).
Among these are (i) a putative Zn-dependent endopeptidases related to the RTX toxin family
(1.C.11, e.g. 2 clusters in N. epidermidis) which in Gram-negative bacteria are implicated in
virulence [48,49],

(ii) enrichment of further families of secreted peptidases (e.g. peptidase family M23, Subtilisin-
like serine protease), that can have diverse roles, both in tissue invasion and establishment
but also in modulation of other virulence factors [50], (iii) extracellular or membrane-attached
putative triacylglycerol esterases/lipases (arCOG06923, PF02089) of which the skin strains
encode 2-3 orthologs, that in bacterial colonization/infection of host epithelia enable
catabolism of host fatty acids in order to alter host immune response or serve anabolic
purposes [51,52]. (iv) patatin-like phospholipase domain proteins (arCOG09020) in N. unguis and
N. oleophilus as found in known epithelial colonizing strains [51,52].

Archaeal ammonia oxidizers form a rare, yet prevalent population on human skin

To understand the abundance, prevalence, distribution and stability of archaeal, and
particularly Nitrosocosmicus representatives on human skin, 47 healthy individuals were
investigated on eight defined body locations (cohorts A1 (age 20-40 y), A2 (60-80+)).
According to their microbial profiles (high AOA, low AOA, see methods), 12 of these individuals
were then selected and longitudinally sampled on four body sites for an entire year (cohort B;
study overview: Extended Data Fig. 4 STORM chart). Samples were quantitatively (qPCR)
analyzed using taxonomic (16S rRNA) and functional (*amoA*) marker genes, targeted towards
the detection of archaea and archaeal ammonia oxidizers (AOA). As metagenomic
sequencing failed due to the low biomass and high background in human DNA, diversity
assessments of the skin archaeome were performed via amplicon sequencing to profile the
bacterial, archaeal and AOA-specific signatures. A total of 1,770 amplicons (including
controls) were processed (details see Materials and Methods, Extended Data Fig. 4 STORM
chart).

Throughout the study (cohorts A, B), archaeal signatures were found at least once in all study
participants, but were overall low, according to qPCR measurements (approx. 0.2% relative
abundance, median across all body sites). AOA-specific qPCR signals accounted for 1.54×10^3
gene copies per 300 cm^2 (Fig. 3A). The highest amount of *amoA* gene copies were found in
samples from the forehead (Fig. 3B); there, AOA signatures were found in samples from every
tested person at each time point (prevalence: 100%).

In amplicon sequencing, 9.45% of all samples contained archaea when the universal PCR
primer combination was applied, whereas 69.98% were positive with the archaea-targeted
combination of 16S rRNA gene primers, emphasizing once more the necessity to use archaea-
targeted methodology for optimized detection. Also with this approach, all individuals were
found to be positive for AOA signatures at least once throughout the study period.

Retrieved 16S rRNA gene sequences of the most abundant ASVs grouped well with the
cultivated archaea from human skin (Fig. 3C), because the most prevalent archaeal family

within 326 samples (cohort A and B) was Nitrososphaeraceae (74% positive samples) with
Nitrosocosmicus being the most prominent genus in this family (24%) (Supplementary Table
S3). The joint classification of 16S rRNA to the *amoA* marker gene (see Methods for details)
resulted in a prevalence of 38% for the NS-Zeta clade, across all samples.
302

Fig. 3. | Quantitative analysis (qPCR) of AOA, archaeal and bacterial signatures on human
skin. The median of 16S rRNA gene copies across all sampling sites is shown; data are
corrected for different 16S rRNA gene copy numbers in archaea (median: 1) and bacteria (6);
https://rrndb.umms.med.umich.edu/). B: Gene copies per 300 cm^2 of AOA, Archaea and
Bacteria, logarithmic scale. C: *AmoA* gene copies per 300 cm^2, across sampling sites (A and
B). Maximum-likelihood phylogenetic tree of 16S rRNA gene sequences of AOA and non-
AOA, including most prevalent and abundant ASVs of cohort A and B. Blue sequences
represent ASVs retrieved from amplicon sequencing of skin samples (cohorts A, B). Values in
brackets show the percentage of prevalence of the ASV in our samples (C).

Nitrosocosmicus signatures show longitudinal coherence and unique biogeographic patterns across body sites

Across all eight body sites sampled (cohort A), Nitrososphaeraceae and Nitrosocosmicus signatures were mostly detected on forehead, décolleté, back and arms, indicating a skin physiology-dependent pattern of ammonia-oxidizing archaea in general (Fig. 4, 16S rRNA gene amplicons). High proportions of AOA from the head region could be confirmed quantitatively (qPCR), as highest copy numbers of the *amoA* gene (up to $4.11*10^4$ per 300 cm^2) were retrieved from the head samples (part and forehead; Fig. 3B).

Nitrosocosmicus signatures were stable longitudinally (cohort B, 1 year sampling period), and even individuals that were not Archaea positive at the initial sampling time point became positive over the course of the study period. In addition, none of the participants carrying archaea at the initial time point lost their archaeome during the year of monitoring. Seasons did not have a significant impact on the abundance of AOA (MaAsLin2, q-value = 0.95), indicating that contact with the outer environment, e.g. through contact to soil by gardening work (MaAsLin2, q-value = 0.49), did not influence the skin archaeome. Signatures assigned to Nitrosocosmicus (global mean = 0.17; global variance = 0.09; importance = 0.17) and other genera of the family *Nitrososphaeraceae* (global mean = 0.33; global variance = 0.15; importance = 0.12) achieved high importance in linear regression analysis over time supporting our assumption that AOA represent a coherent longitudinal trait on human skin (Supplementary Table S4). Further information on bacterial ammonia oxidizers can be found in the Supplementary Information.

Fig. 4. | AOA prevalence across all samples from different body sites based on 16S rRNA gene sequences, shown by summed read counts (n=326 samples). The reads obtained were assigned to Nitrosocosmicus and other non-higher resolved genera belonging to the family Nitrososphaeraceae (A). Boxplot showing the abundance of AOA (normalized read counts) per body site (B). Boxplot showing AOA dynamics (normalized read counts) across the three covered seasons. Both B) and C) refer to the data of the longitudinal cohort (1 year, 6 time points, 3 seasons, 4 body sites, 283 samples) (C).

AOA are an integral component of the healthy human skin microbiome

Network analyses were performed to understand the integration of AOA into the skin microbiome. These showed a significant positive correlation of Nitrosocosmicus with integral bacterial members of the human skin microbiome (in particular *Lawsonella* and *Finegoldia* SCNIC sparCC p-adjust 0.26; see also initial in silico metabolic modeling in Extended Data Fig. 5). In contrast to other microbiome components, this interaction was relatively stable across all sampling time points (Fig. 5A, as visualized by the dark color of the node, see legend). A specific integration of Nitrosocosmicus into the skin microbiome network was underscored by the lack of positive correlations of other archaeal signatures that are not common on human skin (e.g., *Methanobrevibacter*).

Fig. 5 | Microbial co-occurrence network between the 40 most abundant archaeal and bacterial taxa (SRS normalized Cmin = 100, TSS and log transformed). Significant <u>positive</u> associations from Spearman correlations are highlighted by connecting edges. Taxa (nodes) were colored according to each sampling event (t1 yellow, t2 red, t3 pink, t4 green, t5 beige, t6 blue), resulting in dark gray colors for longitudinally consistent taxa in the dataset. The location of Nitrosocosmicus within the network is highlighted by a red ellipse (A). Spearman correlation plot of AOA signatures and sebum content of the skin (B).

Ordinary least squares (OLS) regression models were used to find associations between
archaeal taxa and acquired metadata (skin physiology, demographics of the cohort and their
lifestyle as well as health parameters). Considering the compositional nature of our amplicon
datasets we used CLR (center log ratio) transformed values and regressed them with all
numerical and categorical metadata categories while taking care of several potential
confounders (which partly had significant impact on the microbial communities) like a subject's
sex, age, BMI, different sampled body sites and the overall microbial diversity (Shannon
diversity). According to this analysis, signatures of Nitrososphaeraceae (n = 156) were
significantly positively associated (q = 0.02) with qPCR-retrieved quantity for AOA
(Supplementary Table S5). This finding supports the congruence of amplicon-based
qualitative and qPCR-based quantitative molecular analyses.

The abundance of *amoA* genes as detected by qPCR (tp1, n=48) was not significantly
influenced by sex (female vs. male) and age group (group A1 (20-30y) and A2 (>60y)).
However, the abundance of amoA signatures correlated with lower pH (rho= -0.081, p=0.174),
and significantly with age (rho=-0.16, $p=7.7*10^{-3}$), trans-epidermal water-loss ("Tewameter",
rho=0.22, $p=4.8*10^{-4}$), and particularly sebum concentration (rho=0.47, $p=1.0*10^{-16}$) according
to Spearman rank correlations.
382
To assess the possibility whether AOA are involved in psoriatic disease, as hypothesized
earlier [16], 20 subjects were recruited and n=72 samples from lesioned and healthy skin
samples were taken. However, according to differential abundance analysis based on
ANCOM-BC (p adjust = 0.07) and MaAsLin2 (q-value = 0.54) samples from psoriatic skin did
not reveal significantly higher or lower AOA counts.

Based on our observations, we can conclude, that Nitrosocosmicus is an integral and stable
component of the microbial epidermal skin microbiome that is associated with relevant
bacteria via the exchange of ammonia and is characteristic of healthy skin with higher sebum
content, less moisture and lower pH.

Discussion

In this study we demonstrate the consistent occurrence of ammonia oxidizing archaea of the genus Nitrosocosmicus by molecular and cultivation techniques on human skin indicating that organisms of this taxon form an integral component of the skin microbiome. Our evidences include:

(i) high prevalence and specific biogeographic patterns in molecular analyses

(ii) correlation with specific bacterial taxa in correlation networks

(iii) independent enrichments of Nitrosocosmicus strains from human samples, and

(iv) adaptations reflected in their genomes.

Based on their central energy metabolism, the ammonia oxidizing archaea on skin should be involved in removing nitrogenous compounds such as urea and NH_3 that are excreted with human sweat [53], thereby lowering pH [17]. Considering that one of their metabolic byproducts is NO and that the final product of ammonia oxidation is nitrite (NO_2^-), they could also be key players in maintaining NOx homeostasis in the human skin. Since NO is one of the most important signaling molecules in mammalian physiology, and is involved in a range of physiological as well as pathological processes on the human skin, reviewed in [54] and [55], AOA could play a role in skin health or physiology.

Although the overall abundance was relatively low (0.2% of the total microbial signatures in our studies) and our conditions for initial enrichment of AOA were broad enough to include other AOA taxa (such as *Nitrososphaera, Nitrosotalea, Nitrosopumilus*), representatives of the genus Nitrosocosmicus were consistently detected by molecular methods and were the only ones obtained in laboratory cultures. We hypothesize that the actual relative abundance of Nitrosocosmicus could be even higher, as isolation of DNA from Nitrosocosmicus requires harsh cell lysis. However, AOA might also be notoriously low in abundance on the human body due to hygiene practices or clothing preferences since we found sensitivity of AOA to various plastic compounds.

The overall physiology of the genus shows a high congruence with life on the human skin, as Nitrosocosmicus strains exhibit the highest ammonia tolerance among AOA, tolerating between 20 to 200mM of NH_4^+ [33]. This coincides with the elevated concentrations of ammonia (0.6 - 5.8 mM) and urea (22 ± 8 mM) present in human sweat ([56,57]; Table 2.6). Further, among characterized AOA clades, members of Nitrosocosmicus encode the largest repertoire of putative biofilm-forming genes [32,58], and exhibit strong potential for surface attachment [30–33,59], a general characteristic of the skin microbiota, and a desired trait for skin probiotics [60].

It is also intriguing that the genus Nitrosocosmicus is the most abundant AOA taxon recovered
from various plant tissues and was shown to trigger systemic resistance against pathogens
[61,62]. Based on these observations, we can hypothesize that the Nitrosocosmicus genus
among the AOA is particularly suited to colonize a wide range of habitats, including host
surfaces and thus could represent a group of emerging commensals. Parallels with
representatives of the archaeal order Methanomassiliicoccales are obvious, whose
representatives can be found in diverse ecosystems, with specific adaptations that allow them
to thrive in host microbiomes of the gastrointestinal tract [63,64].

It should be noted, that oscillatory behavior of microorganisms, i.e. thriving in natural
environments but also host systems is not rare, however, has been mostly studied in
(opportunistic) pathogens, such as *Vibrio sp.* ([65]; aquatic environments), *Stenotrophomonas*
maltophilia ([66], soil, plant environments), or *Bukholderia pseudomallei* [67].

Further studies of skin archaea will be needed to reveal their evolutionary paths towards
commensalism and to elucidate their roles in the skin microbiome and human health. The
cultures obtained in this study are crucial prerequisites for advancing our knowledge in this
area.
445
446

Methods

Ethics and recruitment of cohorts

Subjects for microbiome analyses were recruited as part of the project "Detection of Archaea on Healthy and Diseased Skin" with the ethical vote EK: 27-289 ex 14/15 (Medical University of Graz) and all ethical requirements were followed and fulfilled. The following exclusion criteria were taken into account: Antibiotics therapy within the last three months, pregnancy, ongoing breast-feeding and non-compliance. The participants were asked to not shower or apply skin cream the day before sampling.

Initial cultivation

Prior to enrichment setups, subjects not working in an experimental laboratory were pre-screened for the presence of *Nitrososphaerales* by targeted 16S rRNA gene amplification (see study cohorts below). Inoccula for enrichment cultures were taken from 3 individuals that showed highest presence of AOA. Sterile polyurethane single sponge swabs (BD BBL™ CultureSwab™ EZ) premoistened with sterile freshwater medium (FWM 8,9) were used to sequentially rub the face, neck, upper body, back, arms, and legs. The polyurethane swab was then removed from the plastic shaft and directly added to FWM (Enrichments T1S, X2B and Z3A). For enrichment R2S, free edges of a subjects' fingernails (two to three clippings each) were directly added to FWM (Enrichment R). Earlier studies had shown the suitability to use fingernail microbiomes to assess the skin microbiome, due to scratching [68].

All initial enrichments were set up in sterile 30 ml polystyrene screw cap tubes (Greiner) containing 20 ml FWM supplemented with 0.5 mM NH4Cl, 20 µl non-chelated trace element solution, 7.5 µM FeNaEDTA, 20 µl vitamin solution as well as 25 µg/ml carbenicillin and 25 µg/ml kanamycin (for detailed medium composition see table 1 in [28,69].

Variations between initial enrichments included the carbon source (2 mM Na_2CO_3 or $NaHCO_3$), pH (7 or 8.5), temperature (28°C or 32°C) supplementation with organic substrates (0.5 mM acetate and 0.5 mM pyruvate or none) and filter sterilized spent growth medium of *Nitrososphaera viennensis* EN76 [28] and Ca. Nitrosocosmicus franklandianus C13 [31] (2.5% (v/v) (Supplementary Table S6). Cultures were incubated aerobically without shaking in the dark and routinely transferred (transfer volume 20% v/v) upon reaching nitrite concentrations of approximately 450 µM. After transferring enrichments R2S and T1S into a medium containing 10 µg/ml chloramphenicol, 10 µg/ml novobiocin and 10 µg/ml streptomycin twice, no bacterial contaminants were detected.

Routine culture maintenance

Enrichments were routinely cultivated in sterile 30 ml polystyrene screw cap tubes containing 18 ml FWM supplemented with 0.6 mM NH_4Cl, 2 mM Na_2CO_3 (R) or $NaHCO_3$ (T1S, X2B, or Z3A), 20 µl non-chelated trace element solution without molybdenum ($Na_2MoO_4 \cdot 2H_2O$), 7.5 µM FeNaEDTA, 20 µl vitamin solution as well as 25 µg/ml carbenicillin and kanamycin and are transferred every two to four weeks upon reaching nitrite concentrations of approximately 450 µM. The inoculation volumes were decreased to 10% (v/v).

The medium of enrichment R2S additionally contained 10 mM MOPS ((3-(N-morpholino)
propanesulfonic acid), titrated to pH 7 with HCl) and had a final pH of 7. Ammonium and nitrite
concentrations were measured colorimetrically by the indophenol method and Griess reagent
respectively, as described previously [28,69]. Aliquots used for ammonium measurements were
frozen at -20°C and measured jointly at a later time point. Cultures were incubated aerobically
at 28°C without shaking in the dark. Solutions were autoclaved (FWM, NH_4Cl) or filter sterilized
with a 0.2 µm filter (trace element solution, vitamin solution, FeNaEDTA, antibiotics) prior to
use (Extended Data Fig. 6: Enrichment Scheme).

Microscopy

Cultures were 10x concentrated before being imaged with phase contrast light microscopy
using 60x and 100x oil immersion objectives. F_{420} autofluorescence was imaged using an F420
filter cube. For scanning electron microscopy six ml of each enrichment at late exponential
phase were concentrated by centrifugation (21000 g, 4°C, 30 min). Pellets were fixed in 2.5%
glutaraldehyde for 2 h, washed three times with 1x PBS and resuspended in 100 µl 1x PBS.
50 µl of the fixed cell suspension were spotted onto 0.01% poly-L-Lysine coated glass slides
(5 mm) and sedimented for 2 h, before being dehydrated in an ethanol series (30-100%, 5 min
each). The dehydrated cells were critical point dried in liquid carbon dioxide and subsequently
coated with gold in a sputter coater. Samples were imaged in a JEOL IT 300 scanning electron
microscope at 20 kV (CIUS imaging facility, Univ. of Vienna).

DNA extraction of cultures
High molecular weight (HMW) DNA of enrichment cultures was extracted by a combination of
chemical lysis based on a modified lysis buffer for *Sulfolobus acidocaldarius* [70] and freeze-
thawing, followed by a phenol chloroform extraction.
Between 0.3 to 0.5 L of each skin enrichment culture were concentrated by filtration (0.22 µm
MF-Millipore™ mixed cellulose ester (MCE) membrane filter), washed off the filter with FWM
and pelleted by centrifugation (21000 g, 4°C, 30 min). Pellets were resuspended in a modified
TENST buffer (20 mM Tris pH 8, 1 mM EDTA, 100 mM NaCl, 5% N-Lauroylsarcosine, 0.12%
Triton X-100) and incubated at 50°C for 1-2 hours. Samples were subsequently subjected to
three rounds of freeze-thawing (5 min at -70°C followed by 5 min at 70°C). Proteinase K was
added to a final concentration of 100 µg/ml and samples were incubated at 65°C for 1-2 hours.
After lysis HMW DNA was extracted by a standard phenol chloroform extraction (2x
Phenol:Chloroform:Isoamylalcohol 25:24:1, 2x Chloroform:Isoamylalcohol 24:1) with low-
speed centrifugation (4500 g, 4°C, 3 min) between steps. DNA was precipitated overnight at
4°C by adding 1µl glycogen as a carrier and 2x volume of a PEG6000 solution (30% PEG6000
(v/v), 1.6 M NaCl). The precipitated DNA was washed twice with 4°C cold 70% EtOH, air-dried
at room temperature for 15-30 min, resuspended in 50-60°C nuclease free water followed by
an incubation at 55°C for up to two hours. The extracted HMW DNA was quantified
fluorometrically with a Qubit™ 2.0 Fluorometer (Invitrogen) and stored at -20°C.

Quantitative PCR
Archaeal *amo*A genes were quantified in triplicate using primers amoA19F and amoA629R as
previously [29,30] with some modifications: 20 µl reactions contained 10 µl Luna® Universal
qPCR Master Mix, 3 µl nuclease free water, 0.5 µm of each primer and 5 µl DNA template.
DNA was amplified in a CFX Real Time PCR System (Bio-Rad) with following cycling

parameters: 95°C for 1 min, 40x cycles of 95°C 15 s denaturation, 60°C 30 s joint annealing-extension and 60°C 30 s fluorescence measurement. Standard dilutions prepared from DNA of each enrichment respectively using primers amoA19F and amoA629R ranged between 10^2 and 10^8 copies µl $^{-1}$. The efficiency of qPCR assays was between 86.06 and 93.04% with an R^2 of $\geq$ 0.996.

Sanger and NGS sequencing of marker genes and genomes

In order to confirm purity and identity of cultures, 16S rRNA genes were amplified by PCR with the archaea specific primer A109f [71], the universal primer Univ1492R [72], the crenarchaea specific primer pair 771F/957R [73] and the fungal ITS region-specific primer pair ITS1F and TW13 [74]. Amplification of the *amoA* genes was performed by PCR using the primers amoA19F[75] and amoA629R[76]. PCR products were purified using the Monarch® PCR & DNA Cleanup Kit from New England Biolabs before being sequenced.

Genome sequencing, assembly and comparative genomics

DNA of our enrichments was shotgun sequenced using the NovaSeq 6000 (paired-end, 150 bp) platform at Novogene. Reads were trimmed and Illumina adapters were removed using Trimmomatic (SLIDINGWINDOW:5:20 LEADING:5 TRAILING:5 MINLEN:50 HEADCROP:6) [77]. To obtain the full genomic sequences of the enriched archaeal communities from the selected cultures, Nanopore sequencing on MinION Mk1C (Oxford Nanopore Technologies plc., UK) was performed. For library preparation, extracted genomic DNA was repaired with the NEBNext Companion Module (New England Biolabs GmbH, GER), and then prepared for sequencing on a chemistry version 14 flow cell (R10.4.1, FLO-MIN114) following the Ligation sequencing gDNA – Native Barcoding Kit 24 V14 (SQK-NBD114.24) and flow cell (FLO-MIN114) protocols.

Nanopore sequencing was performed according to the following configurations: run-time of the MinION Mk1C device was set for 72 hours, with a pore scan frequency of 1.5 hours, a minimum read length of 200 bp, high-accuracy base calling, and active channel selection, reserved pores, read splitting, trimming barcodes and mid-read barcode filtering enabled. The following software versions were used: MinKNOW v22.10.7, Bream v7.3.5, Configuration v5.3.8, Guppy v6.3.9 and MinKNOW Core v5.3.1. Resulting sequencing data was analyzed with the following specifications: First, simplex base calling was carried out on a GPU node at the Life Science Compute Cluster (LiSC) University of Vienna (Austria), using guppy-gpu v6.4.2 and the dna_r10.4.1_e8.2_260bps_sup.cfg model for superior base calling. Secondly, duplex base calling was achieved at a GPU Node at the LiSC University of Vienna (Austria), following the recommended tutorial on: https://github.com/nanoporetech/duplex-tools using dorado v0.1.1 (https://github.com/nanoporetech/dorado) with the dna_r10.4.1_e8.2_260bps_fast@v4.0.0 fast model , followed by finding duplex pairs with duplextools v0.2.20-3.10 (https://github.com/nanoporetech/duplex-tools) and dorado stereo/duplex base calling with the dna_r10.4.1_e8.2_260bps_sup@v4.0.0 superior model. Merged simplex and duplex reads were then investigated by performing quality control using NanoPlot nanocomp v1.20.0 [78] followed by filtering with filtlong v0.2.1 (https://github.com/rrwick/Filtlong) (--min-length 200; --keep_percent 90, and --target_bases 20 million). Filtered reads were then assembled using flye v2.9.1 in --nano_hq and --meta mode with a target genome size of 3 Mbpm [79]. Reads were mapped with minimap2 v2.24 [80], and then polished with racon v1.5.0 (https://github.com/isovic/racon) using the following

settings (-m -8 -x -6 -g -8 -w -500). Finally, consensus contigs were obtained with medaka
v1.7.2 according to the r1041_e82_260bps_sup_g632 model
(https://github.com/nanoporetech/medaka). Basic genome statistics were calculated with
genometools by calling gt seqstat [81], and classified the genomes using gtdbtk v2.1.1 with the
--full_tree option enabled [82]. Genome completeness and contamination were estimated with
checkm2 v1.0.1 including the --allmodels option [83]. After genome comparison using drep
v3.4.2 [84], genomes were finally annotated with eggnog-mapper v2.1.9 [85–90] including prodigal
for gene identification. Marker genes were identified with Metaxa2 2.2 [91] and the remaining
contaminants of enrichments were estimated with kraken2 2.1.2 and bracken 2.8 according to
the PlusPFP database [92,93] (https://benlangmead.github.io/aws-indexes/k2).

Gene prediction and annotation of the MAGs assembled in this study was performed
automatically using the Microscope annotation platform from Genoscope[94], followed by
extensive manual curation. Annotations were supplemented with arCOG assignments from
the archaeal Clusters of Orthologous Genes database (2018 release) [95] , Carbohydrate-active
enzymes assignments (CAZymes) from dbCAN2 (v7.0) [96] and TCDB family assignments from
the Transporter Classification Database [97], using scripts from [98].

Orthologous protein families from a dataset including *Ca.* N. franklandianus C13, *Ca.* N.
oleophilus MY3, *Ca.* N. hydrocola G61, *Ca.* N. agrestis SS, and *Ca.* N. arcticus Kfb and the
two newly sequenced Ca. Nitrosocosmicus isolates R2S and T1S were constructed using
Orthofinder (v.2.5.4) with standard settings [99].
594

Study-cohorts for molecular-based analyses

Forty-seven healthy individuals (group A1: female=11, male=10, age=20-40y; group A2: f=15,
m=11, age=60-85) were recruited for an in-depth analysis of their skin microbiome, and a
subgroup of high vs. low skin archaeal carriers was selected based on quantitative (amoA
gene and 16S rRNA gene qPCR universal and archaea specific primers) and qualitative (16S
rRNA gene amplicons universal and archaea specific primers) observations for longitudinal
investigations (group B1 (from A1): f=2, m=4; group B2 (from A2): f=5, m=1) over the period
of almost two years (6 time points). The sampling sites for A1/A2 were hand, outer arm, crook,
armpit, forehead, part, décolleté and back (n=7), and for B1/B2: forehead, décolleté, arms and
back (n=4). Supplementary Table S7 gives an overview on the demographic characteristics
of all recruited subjects.
Each subject completed a questionnaire with different categories, e.g., height and weight, skin
type, and detailed questions about their respective lifestyle (e.g., eating habits, smoking,
alcohol consumption, skin care, medications, animal contact, and hobbies).
609

Skin measurements and sampling

The forehead and the part, as well as both hands, forearms, crooks and armpits were swabbed
with pre-moistened (0.9 % (v/v) NaCl) swabs (BD BBL ™ Culture Swab ™ EZ; Becton
Dickinson, USA). NaCl was baked at 260°C for 24 h before solving in PCR-grade water to
remove DNA traces. The remaining body sites, including décolleté and back (approximately
300 cm^2), were sampled with a pre-moistened surface-sampling cellulose-sponge kit (VWR,
300-0229P). For the longitudinal sampling (cohort B1/B2, tp2-6), only the forehead was
sampled with the swab. After skin sampling, samples were immediately frozen at -80°C until
further processing. All skin sampling events were accomplished in a lab that never handled AOA.

Measurements of the skin physiology was performed using the Cutometer(R) MPA580
(Courage + Khazaka, Germany), using the Tewameter (to determine the barrier function by
measuring the water loss; 30s, 3 times), the pH-meter, the Sebumeter (to determine the
sebum-content, 30s), and the Corneometer (to determine skin moisture). Each skin site was
measured three times and the mean of the measurements is reported in Supplementary
Tables S8 and S9. Further, the surface temperature of each region was determined using an
infrared thermometer (Future Founder, China).

The interpretation of skin parameters was done based on the recommended specifications by
the manufacturer, which are provided in Supplementary Table S10.

DNA extraction, controls, amplicon sequencing, and qPCR of skin samples

The DNA was extracted from swabs and sponges using the Purelink ™ Microbiome DNA
Purification Kit (Rectal or Environmental Swab Samples; Thermo Fisher Scientific, USA). For
the extraction from sponges (liquid sample extract from décolleté and back) one ml of the
sample was concentrated by centrifugation (20817 x g, 4°C; 10 min). The largest proportion
of the supernatant was removed, and the remaining 70 µl was used for bead beating with 800
µl of the S1 lysis buffer. Material and process controls were included in each step to control
for the DNA contamination of the used reagents. The DNA was stored at -20°C until further
downstream analyses.
Various targets were assessed using amplicon-based analyses: i) the archaeal 16S rRNA
gene region (nested PCR approach as described in [100] and [16]), the ii) amoA targeted approach
(described in [28,101,102]), and the iii) "universal" approach, targeting the microbial 16S rRNA gene
region [103]. All primer sequences and PCR protocols are provided in Supplementary Table S11.
Library construction and next-generation sequencing (Illumina MiSeq®) was performed at the
Core Facility Molecular Biology at the Center for Medical Research (Medical University of
Graz, Austria). In a first step, the DNA concentrations of the generated amplicons were
normalized using a SequalPrepTM normalization plate (Invitrogen) and subsequently each
sample was indexed with a unique barcode sequence by 8 cycles of indexing PCR. These
indexed Samples were pooled and purified by gel cuts before the library was run on an Illumina
MiSeq® instrument and the MiSeq® Reagent Kit v3 with 602 cycles (2 x 301 cycles).

Bioinformatics and data analysis

Resulting demultiplexed fastq reads were processed with QIIME2 [104] versions 2018.11
through 2023.2. After quality control, denoising of reads was achieved with DADA2 [105] and
truncation of forward reads at a length of 230bp and 120bp for reverse reads. Potential
contaminants were identified with decontam v.1.12 [106] using the prevalence mode with a
threshold of 0.5 and removed from downstream analysis. Representative sequences were
classified by a pre-trained Naïve-Bayes classifier based on the curated and trimmed SILVA138
database [107] using RESCRIPt [108] and the feature-classifier classify-sklearn plugin [109]. For
phylogenetic metrics like UniFrac [110], a rooted phylogenetic tree was generated with FastTree2
[111] based on a masked MAFFT alignment. Data normalization was achieved by scaling with
ranked subsampling and the q2-srs QIIME2 plugin [112] at Cmin 50, 100 and 1000 followed by
calculating core metrics for alpha and beta diversity. Feature differentials were determined
with the QIIME2 plugins aldex2 [113], ANCOM [114], ANCOM-BC [114], and the R packages
ANCOM2 (https://github.com/FrederickHuangLin/ANCOM-Code-Archive) and MaAsLin2 [115]
including time as a fixed effect and different subjects as a random effect where applicable,

while feature loadings were determined with the QIIME2 plugin DEICODE [116]. Longitudinal
analysis covering feature volatility over time as well as other numerical metadata categories
like age, BMI or skin physiology measurements including i.e. pH, skin fat or skin wetness were
realized with the q2-longitudinal plugin in QIIME2 [109]. The same plugin was used to calculate
linear-mixed-effect models of Shannon diversity and weighted UniFrac distances along
principal coordinate axis 1. Amplicons of the amoA marker gene were processed in a similar
way without read truncations for DADA2 and read classifications by a pre-trained Naïve-Bayes
classifier based on the amoA reference database [37,117]. Connection of 16S rRNA to amoA
gene information was achieved by following the tutorial at https://github.com/alex-
bagnoud/Linking-16S-to-amoA-taxonomy/. Prediction of metadata categories were realized
with the q2-sample-classifier QIIME2 plugin [109].
674
Custom python code for regressing all metadata categories:
To identify potential associations between taxa and recorded numerical and categorical
metadata an ordinary least squares (OLS) regression fit method was used with python's
statsmodels smf formula (statsmodels.regression.linear_model.OLS - statsmodels_0.15.0
(+8)). Microbial counts of the different investigated cohorts were first clr transformed and
filtered for less than 25% missing values. These values were then used as the response
variable in dependence of each individual, their sex, age, BMI, body site, the overall microbial
diversity indicated by the Shannon index (H'), and where applicable the time of sampling. The
model was then evaluated for each genus and recorded metadata according to their q-values
(FDR-corrected p-values) and estimated regression coefficients (python code available at
GitHub: https://github.com/CME-lab-research/SkinArchaeome).
686
Network analysis:
Network analyses were conducted to identify mutually exclusive and co-occurring clusters of
archaea and bacteria in our merged (archaea-specific and universal) 16S rRNA gene amplicon
dataset. After SRS normalization to Cmin = 100 and TSS and log transformations the 40 most
abundant taxa were represented as nodes within the network where node size reflects taxa
abundance, and edges indicate only positive associations of Spearman correlations between
pairs of taxa. Then, we used PCoA to ordinate the nodes in a two-dimensional plot, such that
correlating nodes were close together and anti-correlating nodes were far apart. In addition,
correlations of the network were further supported by the Sparse Co-Occurrence Network
Investigation for Compositional data (SCNIC) package
(https://github.com/lozuponelab/SCNIC) using the sparCC metric with default settings for
filterings and the R value, and bootstrap adjusted p-values.
699
Phylogenomic tree reconstruction:
A genome database was created comprising the two AOA genomes presented in this study
together with 141 MAGs and completely sequenced genomes (133 AOA and 8 non-AOA
genomes) from IMG, DDBJ, or NCBI databases followed by protein prediction using Prodigal
v2.6.3 [90]. The workflow proposed by Graham et al. (2018) [101] based on the archaeal single-
copy gene collection [118] was followed to identify phylogenetic markers (e-value 10^{-10}) to
reconstruct the phylogenomic tree. A total of 36 ribosomal protein families encoded in at least
100 out of the 143 genomes present in our genome database were selected. Each protein
family was aligned independently using MAFFT 7.520 [119] (L-INS-i algorithm) followed by the
trimming of the alignments using BMGE [120] with default parameters. Trimmed protein families
were concatenated using a custom python script and the concatenated alignment was used

to reconstruct a maximum likelihood (ML) phylogenomic tree in IQ-TREE v2.2.2.7 [121] under
the LG+C20+F+G model with 1000 ultrafast bootstrap replicates.
713
Phylogenetic analysis of 16S rRNA sequences:
The identification of 16S rRNA sequences in our AOA genome database (see above) was
performed using Barrnap v0.9 (https://github.com/tseemann/barrnap). The 16S rRNA
sequence encoded in the MAG GCA_025930575.1 (NS-Epsilon) was used as a query in a
BLASTN search against the NCBI database to improve the sequence representation of this
sub lineage with environmental sequences. Sequences were aligned with the SINA v1.7.2
aligner [122] using the SILVA database SSU Ref NR 99 release 138.1 [107] with default parameters
followed by the alignment of the skin ASVs using MAFFT 7.520 [119] (--addfragments option, L-
INS-i algorithm). The alignment was trimmed manually using AliView [123]. The maximum
likelihood phylogenetic tree was reconstructed using IQ-TREE v2.2.2.7 [121] with the best-fit
model TIM2e+I+R3 chosen by ModelFinder ("-m MFP") [124] according to BIC (Bayesian
Information criterion), SH-like approximate likelihood ratio test ("-alrt 1000") [125] and 1,000
ultrafast bootstrap replicates ("-bb 1000").
727

qPCR specific methods

A SYBR green-based quantitative real-time PCR (qPCR) approach targeting the archaeal
amoA gene (19F-629R) and the 16S rRNA gene of archaea (A806F-A958R) and bacteria
(Bac_331F-Bac_797R) was used to determine the abundance of AOA, all archaea and all
bacteria on selected body sites. Primers, reaction mixtures and thermocycler programs are
listed in Supplementary Table S11. All qPCRs followed the MIQE Guidelines.

qPCRs were performed with the Bio-Rad SsoAdvanced™ Universal SYBR® Green Supermix.
Each reaction contained 1µl of DNA template and 9µl of mastermix. The mastermix for the
archaeal amoA approach consisted of 5µl SsoAdvanced™ Universal SYBR® Green Supermix
(Bio-Rad), 1.25µl of each primer (1.25µM) and 1.5µl of PCR grade water. The mastermix for
the archaeal and bacterial 16S rRNA gene approach consisted of 5µl SsoAdvanced™
Universal SYBR® Green Supermix (Bio-Rad), 0.3µl of each primer (10µM) and 3.4µl of PCR
grade water. Amplifications were carried out in a Bio-Rad CFX96 Touch™ Real-Time PCR
Detection System. The qPCR cycling conditions were as follows: initial denaturation at 95°C
for 1.5min, followed by 45 cycles of denaturation for 15sec at 95°C, annealing for 30sec at
60°C, elongation for 40sec at 72°C, a final denaturation for 15sec at 95°C, 5sec at 60°C and
a storage temperature of 8°C. A plasmid containing the amoA gene of *Nitrososphaera*
viennensis with a starting concentration of 104.1ng/µl (pscA_Ampkan_AmoA, clone 3) was
used as a quantification standard in a 1:10 dilution series from 10^{-2} to 10^{-10}. Product specificity
was confirmed by melting curve analysis. All qPCR experiments were performed in triplicates
and the results were averaged for further calculations. Data from a previous experiment on
the total archaeal community (based on qPCR of the 16S rRNA gene) was also included in
the analyses. To account for the different area sizes of the sampled body sites, copy numbers
were adjusted to 300 cm^2 (amoA gene copies per 300 cm^2; see Supplementary Table S12).
The qPCR files were analyzed using the CFX Manager™ Software, version 3.1 and processed
with the R software, version 3.6.1 and Microsoft Excel.
754

Data availability

All raw sequencing data were submitted to NCBI's Sequence Read Archive (SRA) https://www.ncbi.nlm.nih.gov/sra and are accessible under the following projects: PRJNA1105887 (archaea specific 16S rRNA gene amplicons of cohort A), PRJNA1107153 (universal 16S rRNA gene amplicons of cohort A), PRJNA1107462 (AOA specific amoA gene amplicons of cohort A), PRJNA1107502 (archaea specific 16S rRNA gene amplicons of cohort B), PRJNA1107512 (universal 16S rRNA gene amplicons of cohort B), PRJNA1105887 (archaea specific 16S rRNA gene amplicons of cohort C), PRJNA1108073 (universal 16S rRNA gene amplicons of cohort C), PRJNA1108908 (pooled shotgun metagenomics data of cohort A), PRJNA1108960 (Illumina sequencing data of enrichments), PRJNA1109334 (Nanopore sequencing data of enrichments). Assemblies are publicly available under NCBI BioProjects PRJNA1103140 (Ca. Nitrosocosmicus epidermidis T1S) and PRJNA1103141 (Ca. Nitrosocosmicus unguis). Annotated assemblies are also available at the MicroScope Microbial Genome Annotation and Analysis Platform, developed by the Laboratory of Bioinformatics Analyses for Genomics and Metabolism, Genoscope, Evry, France.

Code availability

The present study did not generate new code, and mentioned tools used for the data analysis were applied with default parameters unless specified otherwise in Methods and our Github repo.

Contributions

AM supervised and performed microbiome sampling and analysis. MD supervised and performed cultivation and microscopy. MM and LH supervised and supported cultivation work. UY performed cultivation. SD, AL, TG, DB performed sampling, sequencing, and data analysis of qPCR and amplicon data. RP performed phylogenetic analyses and genome assembly. TG and PW recruited patients and performed sampling. MK performed genome annotations and contributed to study concept. CME and CS developed the study concept, wrote initial proposal, and wrote, together with MK and AM the manuscript. All authors contributed to the finalization of the manuscript.

Acknowledgments

We acknowledge the technical support by Ignaz Herzeg, Lisa Wink, Polona Mertelj, Sarah Harrer and all collaborators at the Department of Dermatology and Venereology, Medical University of Graz (Andrea Alejandra Catalan Griffiths, Patra Vijaykumar, Isabella Perchthaler, Alexandra Gruber-Wackernagel, Wolfgang Weger, and univ. Prof. Dr. Wolfgang Salmhofer). The work was funded by the Austrian Science Fund FWF Austrian Science Fund (FWF) grant no. P 30796 (given to CME and CS), and supported by FWF grants P 32697, P 30796, SFB F-83, COE 7, given to CME and FWF Z437 to CS. We thank Daniela Gruber from the Core Facility Cell Imaging & Ultrastructure Research (LeWi). We greatly value the computational resources of the MedBioNode provided by the Medical University of Graz and the Life Science Compute Cluster (LiSC) operated by the Computational Systems Biology group at the University of Vienna as well as the Medical University of Graz ZMF Galaxy Team: Core Facility Computational Bioanalytics, Medical

University of Graz, funded by the Austrian Federal Ministry of Education, Science and
Research, Hochschulraum-Strukturmittel 2016 grant as part of BioTechMed Graz.
SD was supported by the local PhD program MolMed.

Conflict of interest

The authors declare no conflict of interest.

References
804

805   1.   Berg, G., Rybakova, D., Fischer, D., Cernava, T., Vergès, M.-C. C., Charles, T.,
806   Chen, X., Cocolin, L., Eversole, K. & Corral, G. H. Microbiome definition re-visited: old
807   concepts and new challenges. *Microbiome* **8,** 1–22 (2020).
808   2.   Fan, Y. & Pedersen, O. Gut microbiota in human metabolic health and disease.
809   *Nature Reviews Microbiology 2020 19:1* **19,** 55–71 (2020). doi: 10.1038/s41579-020-
810   0433-9
811   3.   Flowers, L. & Grice, E. A. The Skin Microbiota: Balancing Risk and Reward. (2020).
812   doi:10.1016/j.chom.2020.06.017 doi: 10.1016/j.chom.2020.06.017
813   4.   Gomaa, E. Z. Human gut microbiota/microbiome in health and diseases: a review.
814   *Antonie Van Leeuwenhoek* **113,** 2019–2040 (2020). doi: 10.1007/s10482-020-01474-
815   7
816   5.   Borrel, G., Brugère, J. F., Gribaldo, S., Schmitz, R. A. & Moissl-Eichinger, C. The
817   host-associated archaeome. *Nat Rev Microbiol* **18,** 622–636 (2020). doi:
818   10.1038/s41579-020-0407-y
819   6.   Bang, C. & Schmitz, R. A. Archaea associated with human surfaces: not to be
820   underestimated. *FEMS Microbiol Rev* **39,** 631–648 (2015). doi:
821   10.1093/femsre/fuv010
822   7.   Gaci, N., Borrel, G., Tottey, W., O'Toole, P. W. & Brugère, J.-F. Archaea and the
823   human gut: New beginning of an old story. *World J Gastroenterol* **20,** 16062 (2014).
824   doi: 10.3748/wjg.v20.i43.16062
825   8.   Nayfach, S., Páez-Espino, D., Call, L., Low, S. J., Sberro, H., Ivanova, N. N., Proal, A.
826   D., Fischbach, M. A., Bhatt, A. S., Hugenholtz, P. & Kyrpides, N. C. Metagenomic
827   compendium of 189,680 DNA viruses from the human gut microbiome. *Nature
828   Microbiology 2021 6:7* **6,** 960–970 (2021). doi: 10.1038/s41564-021-00928-6
829   9.   Fan, Y., Wu, L. & Zhai, B. The mycobiome: interactions with host and implications in
830   diseases. *Curr Opin Microbiol* **75,** 102361 (2023). doi: 10.1016/J.MIB.2023.102361
831   10.   Chibani, C. M., Mahnert, A., Borrel, G., Almeida, A., Werner, A., Brugère, J.-F.,
832   Gribaldo, S., Finn, R. D., Schmitz, R. A. & Moissl-Eichinger, C. A catalogue of 1,167
833   genomes from the human gut archaeome. *Nat Microbiol* **7,** 48–61 (2022). doi:
834   10.1038/s41564-021-01020-9
835   11.   Kumpitsch, C., Fischmeister, F. P. S., Mahnert, A., Lackner, S., Wilding, M., Sturm,
836   C., Springer, A., Madl, T., Holasek, S. & Högenauer, C. Reduced B12 uptake and
837   increased gastrointestinal formate are associated with archaeome-mediated breath
838   methane emission in humans. *Microbiome* **9,** 1–18 (2021).
839   12.   Grice, E. A. & Segre, J. A. The skin microbiome. *Nat Rev Microbiol* **9,** 244–253
840   (2011).
841   13.   Kong, H. H. Skin microbiome: genomics-based insights into the diversity and role of
842   skin microbes. *Trends Mol Med* **17,** 320 (2011). doi: 10.1016/J.MOLMED.2011.01.013
843   14.   Oh, J., Byrd, A. L., Park, M., Kong, H. H. & Segre, J. A. Temporal Stability of the
844   Human Skin Microbiome. *Cell* **165,** 854–866 doi: 10.1016/j.cell.2016.04.008
845   15.   Byrd, A. L., Belkaid, Y. & Segre, J. A. The human skin microbiome. *Nat Rev Microbiol*
846   **16,** 143–155 (2018). doi: 10.1038/nrmicro.2017.157
847   16.   Probst, A. J., Auerbach, A. K. & Moissl-Eichinger, C. Archaea on Human Skin. *PLoS
848   One* **8,** (2013). doi: 10.1371/journal.pone.0065388

17. Moissl-Eichinger, C., Probst, A. J., Birarda, G., Auerbach, A., Koskinen, K., Wolf, P. &
Holman, H.-Y. N. Human age and skin physiology shape diversity and abundance of
Archaea on skin. *Sci Rep* **7,** (2017). doi: 10.1038/s41598-017-04197-4

18. Parks, D. H., Chuvochina, M., Rinke, C., Mussig, A. J., Chaumeil, P. A. & Hugenholtz,
P. GTDB: an ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based taxonomy.
Nucleic Acids Res **50,** D785–D794 (2022). doi: 10.1093/NAR/GKAB776

19. Rinke, C., Chuvochina, M., Mussig, A. J., Chaumeil, P.-A., Davín, A. A., Waite, D. W.,
Whitman, W. B., Parks, D. H. & Hugenholtz, P. A standardized archaeal taxonomy for
the Genome Taxonomy Database. *Nat Microbiol* 1–14 (2021).

20. Parks, D. H., Chuvochina, M., Chaumeil, P.-A., Rinke, C., Mussig, A. J. & Hugenholtz,
P. A complete domain-to-species taxonomy for Bacteria and Archaea. *Nat Biotechnol*
38, 1079–1086 (2020). doi: 10.1038/s41587-020-0501-8

21. Moissl-Eichinger, C. Archaea in artificial environments: their presence in global
spacecraft clean rooms and impact on planetary protection. *ISME J* **5,** 209–19 (2011).

22. Kozlowski, J. A., Kits, K. D. & Stein, L. Y. Comparison of Nitrogen Oxide Metabolism
among Diverse Ammonia-Oxidizing Bacteria. *Front Microbiol* **7,** (2016). doi:
10.3389/fmicb.2016.01090

23. Sun, Y., Liu, Y., Pan, J., Wang, F. & Li, M. Perspectives on Cultivation Strategies of
Archaea. *Microb Ecol* (2019). doi:10.1007/s00248-019-01422-7 doi: 10.1007/s00248-
019-01422-7

24. Umbach, A. K., Stegelmeier, A. A. & Neufeld, J. D. Archaea Are Rare and Uncommon
Members of the Mammalian Skin Microbiome. *mSystems* **6,** (2021). doi:
10.1128/MSYSTEMS.00642-21/ASSET/1CFB65AC-B6A0-4252-8E75-
629C6EA38AC1/ASSETS/IMAGES/LARGE/MSYSTEMS.00642-21-F004.JPG

25. Mahnert, A., Blohs, M., Pausan, M. R. & Moissl-Eichinger, C. The human archaeome:
methodological pitfalls and knowledge gaps. *Emerg Top Life Sci* **2.4,** 469–482 (2018).

26. Usuda, D., Higashikawa, T., Hotchi, Y., Usami, K., Shimozawa, S., Tokunaga, S.,
Osugi, I., Katou, R., Ito, S., Yoshizawa, T., Asako, S., Mishima, K., Kondo, A., Mizuno,
K., Takami, H., Komatsu, T., Oba, J., Nomura, T. & Sugita, M. Exophiala dermatitidis.
World J Clin Cases **9,** 7963–7972 (2021). doi: 10.12998/wjcc.v9.i27.7963

27. Untereiner, W. A. & Naveau, F. A. Molecular systematics of the Herpotrichiellaceae
with an assessment of the phylogenetic positions of Exophiala dermatitidis and
Phialophora americana. *Mycologia* **91,** 67–83 (1999). doi:
10.1080/00275514.1999.12060994

28. Tourna, M., Stieglmeier, M., Spang, A., Könneke, M., Schintlmeister, A., Urich, T.,
Engel, M., Schloter, M., Wagner, M. & Richter, A. Nitrososphaera viennensis, an
ammonia oxidizing archaeon from soil. *Proceedings of the National Academy of*
Sciences **108,** 8420–8425 (2011).

29. Abby, S. S., Melcher, M., Kerou, M., Krupovic, M., Stieglmeier, M., Rossel, C., Pfeifer,
K. & Schleper, C. Candidatus Nitrosocaldus cavascurensis, an ammonia oxidizing,
extremely thermophilic archaeon with a highly mobile genome. *Front Microbiol* **9,**
346006 (2018). doi: 10.3389/FMICB.2018.00028/BIBTEX

30. Alves, R. J. E., Kerou, M., Zappe, A., Bittner, R., Abby, S. S., Schmidt, H. A., Pfeifer,
K. & Schleper, C. Ammonia oxidation by the arctic terrestrial thaumarchaeote
candidatus nitrosocosmicus arcticus is stimulated by increasing temperatures. *Front*
Microbiol **10,** 1571 (2019). doi: 10.3389/FMICB.2019.01571/FULL

31. Lehtovirta-Morley, L. E., Ross, J., Hink, L., Weber, E. B., Gubry-Rangin, C., Thion, C.,
Prosser, J. I. & Nicol, G. W. Isolation of 'Candidatus Nitrosocosmicus franklandus', a
novel ureolytic soil archaeal ammonia oxidiser with tolerance to high ammonia
concentration. *FEMS Microbiol Ecol* **92,** 57 (2016). doi: 10.1093/FEMSEC/FIW057

32. Jung, M. Y., Kim, J. G., Sinninghe Damsté, J. S., Rijpstra, W. I. C., Madsen, E. L.,
Kim, S. J., Hong, H., Si, O. J., Kerou, M., Schleper, C. & Rhee, S. K. A hydrophobic
ammonia-oxidizing archaeon of the Nitrosocosmicus clade isolated from coal tar-
contaminated sediment. *Environ Microbiol Rep* **8,** 983–992 (2016). doi: 10.1111/1758-
2229.12477

33. Liu, L., Liu, M., Jiang, Y., Lin, W. & Luo, J. Production and Excretion of Polyamines To
Tolerate High Ammonia, a Case Study on Soil Ammonia-Oxidizing Archaeon '
Candidatus Nitrosocosmicus agrestis'. *mSystems* **6,** (2021). doi:
10.1128/MSYSTEMS.01003-20

34. Roller, B. R. K., Stoddard, S. F. & Schmidt, T. M. Exploiting rRNA operon copy
number to investigate bacterial reproductive strategies. *Nat Microbiol* **1,** 16160 (2016).
doi: 10.1038/nmicrobiol.2016.160

35. Klappenbach, J. A., Dunbar, J. M. & Schmidt, T. M. rRNA Operon Copy Number
Reflects Ecological Strategies of Bacteria. *Appl Environ Microbiol* **66,** 1328–1333
(2000). doi: 10.1128/AEM.66.4.1328-1333.2000

36. Konstantinidis, K. T., Rosselló-Móra, R. & Amann, R. Uncultivated microbes in need
of their own taxonomy. *ISME J* **11,** 2399–2406 (2017). doi: 10.1038/ismej.2017.113

37. Alves, R. J. E., Minh, B. Q., Urich, T., Haeseler, A. & Schleper, C. Unifying the global
phylogeny and environmental distribution of ammonia-oxidising archaea based on
amoA genes. *Nat Commun* **9,** 1517 (2018).

38. van Ulsen, P., Zinner, K. M., Jong, W. S. P. & Luirink, J. On display: autotransporter
secretion and application. *FEMS Microbiol Lett* **365,** (2018). doi:
10.1093/femsle/fny165

39. Kobe, B. The leucine-rich repeat as a protein recognition motif. *Curr Opin Struct Biol*
11, 725–732 (2001). doi: 10.1016/S0959-440X(01)00266-4

40. Alvarez, C. E. On the origins of arrestin and rhodopsin. *BMC Evol Biol* **8,** 222 (2008).
doi: 10.1186/1471-2148-8-222

41. Marullo, S. & Coureuil, M. Arrestins in host-pathogen interactions. *Handb Exp*
Pharmacol **219,** 361–74 (2014). doi: 10.1007/978-3-642-41199-1_18

42. Chan, J. M., Gori, A., Nobbs, A. H. & Heyderman, R. S. Streptococcal Serine-Rich
Repeat Proteins in Colonization and Disease. *Front Microbiol* **11,** (2020). doi:
10.3389/fmicb.2020.593356

43. Yeats, C., Bentley, S. & Bateman, A. New Knowledge from Old: In silico discovery of
novel protein domains in Streptomyces coelicolor. *BMC Microbiol* **3,** 3 (2003). doi:
10.1186/1471-2180-3-3

44. Gibbs, G. M., Roelants, K. & O'Bryan, M. K. The CAP Superfamily: Cysteine-Rich
Secretory Proteins, Antigen 5, and Pathogenesis-Related 1 Proteins—Roles in
Reproduction, Cancer, and Immune Defense. *Endocr Rev* **29,** 865–897 (2008). doi:
10.1210/er.2008-0032

45. Chen, C. K.-M., Chan, N.-L. & Wang, A. H.-J. The many blades of the β-propeller
proteins: conserved but versatile. *Trends Biochem Sci* **36,** 553–561 (2011). doi:
10.1016/j.tibs.2011.07.004

46. Brandwein, M., Steinberg, D. & Meshner, S. Microbial biofilms and the human skin
microbiome. *npj Biofilms and Microbiomes 2016 2:1* **2,** 1–6 (2016). doi:
10.1038/s41522-016-0004-z
47. Cinar, M. S., Niyas, A. & Avci, F. Y. Serine-rich repeat proteins: well-known yet little-
understood bacterial adhesins. *J Bacteriol* **206,** (2024). doi: 10.1128/jb.00241-23
48. Mondal, A. K., Lata, K., Singh, M., Chatterjee, S., Chauhan, A., Puravankara, S. &
Chattopadhyay, K. Cryo-EM elucidates mechanism of action of bacterial pore-forming
toxins. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1864,** 184013 (2022).
doi: 10.1016/j.bbamem.2022.184013
49. Garcia, C. J., Pericleous, A., Elsayed, M., Tran, M., Gupta, S., Callaghan, J. D.,
Stella, N. A., Franks, J. M., Thibodeau, P. H., Shanks, R. M. Q. & Kadouri, D. E.
Serralysin family metalloproteases protects Serratia marcescens from predation by
the predatory bacteria Micavibrio aeruginosavorus. *Sci Rep* **8,** 14025 (2018). doi:
10.1038/s41598-018-32330-4
50. Chua, W., Poh, S. E. & Li, H. Secretory Proteases of the Human Skin Microbiome.
Infect Immun **90,** (2022). doi: 10.1128/IAI.00397-21
51. Kumar, N. G., Contaifer, D., Wijesinghe, D. S. & Jefferson, K. K. Staphylococcus
aureus Lipase 3 (SAL3) is a surface-associated lipase that hydrolyzes short chain
fatty acids. *PLoS One* **16,** e0258106 (2021). doi: 10.1371/journal.pone.0258106
52. Chen, X. & Alonzo, F. Bacterial lipolysis of immune-activating ligands promotes
evasion of innate defenses. *Proceedings of the National Academy of Sciences* **116,**
3764–3773 (2019). doi: 10.1073/pnas.1817248116
53. Chen, Y.-L., Kuan, W.-H. & Liu, C.-L. Comparative Study of the Composition of Sweat
from Eccrine and Apocrine Sweat Glands during Exercise and in Heat. *Int J Environ
Res Public Health* **17,** 3377 (2020). doi: 10.3390/ijerph17103377
54. Cals-Grierson, M.-M. & Ormerod, A. D. Nitric oxide function in the skin. *Nitric Oxide*
10, 179–193 (2004). doi: 10.1016/j.niox.2004.04.005
55. Suschek, C. V., Opländer, C. & van Faassen, E. E. Non-enzymatic NO production in
human skin: Effect of UVA on cutaneous NO stores. *Nitric Oxide* **22,** 120–135 (2010).
doi: 10.1016/j.niox.2009.10.006
56. Wilson, M. *Microbial Inhabitants of Humans: Their Ecology and Role in Health and
Disease.* (Cambridge University Press, 2005).
57. Huang, C.-T., Chen, M.-L., Huang, L.-L. & Mao, I.-F. Uric acid and urea in human
sweat. *Chin J Physiol* **45,** 109–15 (2002).
58. Abby, S. S., Kerou, M. & Schleper, C. Ancestral reconstructions decipher major
adaptations of ammonia-oxidizing archaea upon radiation into moderate terrestrial
and marine environments. *mBio* **11,** 1–20 (2020). doi: 10.1128/MBIO.02371-
20/SUPPL_FILE/MBIO.02371-20-SF004.PDF
59. Sauder, L. A., Albertsen, M., Engel, K., Schwarz, J., Nielsen, P. H., Wagner, M. &
Neufeld, J. D. Cultivation and characterization of Candidatus Nitrosocosmicus
exaquare, an ammonia-oxidizing archaeon from a municipal wastewater treatment
system. *ISME J* **11,** 1142 (2017).
60. Lopes, E. G., Moreira, D. A., Gullón, P., Gullón, B., Cardelle-Cobas, A. & Tavaria, F.
K. Topical application of probiotics in skin: adhesion, antimicrobial and antibiofilm
in vitro assays. *J Appl Microbiol* **122,** 450–461 (2017). doi: 10.1111/jam.13349
61. Song, G. C., Im, H., Jung, J., Lee, S., Jung, M., Rhee, S. & Ryu, C. Plant growth-
promoting archaea trigger induced systemic resistance in Arabidopsis thaliana against

Pectobacterium carotovorum and Pseudomonas syringae. *Environ Microbiol* **21,** 940–
948 (2019).

62. Taffner, J., Cernava, T., Erlacher, A. & Berg, G. Novel insights into plant-associated
archaea and their functioning in arugula (Eruca sativa Mill.). *J Adv Res* (2019).

63. De La Cuesta-Zuluaga, J., Spector, T. D., Youngblut, N. D. & Ley, R. E. Genomic
Insights into Adaptations of Trimethylamine-Utilizing Methanogens to Diverse
Habitats, Including the Human Gut. *mSystems* **6,** e00939-20 (2021).

64. Cozannet, M., Borrel, G., Roussel, E., Moalic, Y., Allioux, M., Sanvoisin, A., Toffin, L.
& Alain, K. New Insights into the Ecology and Physiology of Methanomassiliicoccales
from Terrestrial and Aquatic Environments. *Microorganisms* **9,** 30 (2020). doi:
10.3390/microorganisms9010030

65. Soto, W. & Nishiguchi, M. K. Environmental Stress Selects for Innovations That Drive
Vibrio Symbiont Diversity. *Front Ecol Evol* **9,** (2021). doi: 10.3389/fevo.2021.616973

66. Brooke, J. S. Stenotrophomonas maltophilia: an Emerging Global Opportunistic
Pathogen. *Clin Microbiol Rev* **25,** 2–41 (2012). doi: 10.1128/CMR.00019-11

67. Ghazali, A.-K., Firdaus-Raih, M., Uthaya Kumar, A., Lee, W.-K., Hoh, C.-C. & Nathan,
S. Transitioning from Soil to Host: Comparative Transcriptome Analysis Reveals the
Burkholderia pseudomallei Response to Different Niches. *Microbiol Spectr* **11,** (2023).
doi: 10.1128/spectrum.03835-22

68. Rayan, G. M. & Flournoy, D. J. Microbiologic flora of human fingernails. *J Hand Surg*
Am **12,** 605–607 (1987). doi: 10.1016/S0363-5023(87)80217-4

69. Reyes, C., Hodgskiss, L. H., Baars, O., Kerou, M., Bayer, B., Schleper, C. & Kraemer,
S. M. Copper limiting threshold in the terrestrial ammonia oxidizing archaeon
Nitrososphaera viennensis. *Res Microbiol* **171,** 134–142 (2020). doi:
10.1016/J.RESMIC.2020.01.003

70. Meulenbroek, E. M., Peron Cane, C., Jala, I., Iwai, S., Moolenaar, G. F., Goosen, N. &
Pannu, N. S. UV damage endonuclease employs a novel dual-dinucleotide flipping
mechanism to recognize different DNA lesions. *Nucleic Acids Res* **41,** 1363–1371
(2013). doi: 10.1093/nar/gks1127

71. Großkopf, R., Janssen, P. H. & Liesack, W. Diversity and structure of the
methanogenic community in anoxic rice paddy soil microcosms as examined by
cultivation and direct 16S rRNA gene sequence retrieval. *Appl Environ Microbiol* **64,**
960–969 (1998).

72. Lane, D. in *Nucleic Acid Techniques in Bacterial Systematics* (eds. Stackebrandt, E. &
Goodfellow, M.) 115–175 (John Wiley & Sons, 1991).

73. Ochsenreiter, T., Selezi, D., Quaiser, A., Bonch-Osmolovskaya, L. & Schleper, C.
Diversity and abundance of Crenarchaeota in terrestrial habitats studied by 16S RNA
surveys and real time PCR. *Environ Microbiol* **5,** 787–797 (2003). doi: 10.1046/j.1462-
2920.2003.00476.x

74. Taylor, D. L. & Bruns, T. D. Community structure of ectomycorrhizal fungi in a *Pinus*
muricata forest: minimal overlap between the mature forest and resistant propagule
communities. *Mol Ecol* **8,** 1837–1850 (1999). doi: 10.1046/j.1365-294x.1999.00773.x

75. Tourna, M., Freitag, T. E., Nicol, G. W. & Prosser, J. I. Growth, activity and
temperature responses of ammonia-oxidizing archaea and bacteria in soil
microcosms. *Environ Microbiol* **10,** 1357–1364 (2008). doi: 10.1111/j.1462-
2920.2007.01563.x

76. Arce, M. I., von Schiller, D., Bengtsson, M. M., Hinze, C., Jung, H., Alves, R. J. E.,
Urich, T. & Singer, G. Drying and Rainfall Shape the Structure and Functioning of

1037   Nitrifying Microbial Communities in Riverbed Sediments. *Front Microbiol* **9,** (2018).
1038   doi: 10.3389/fmicb.2018.02794

1039   77.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
1040   sequence data. *Bioinformatics* **30,** 2114–2120 (2014). doi:
1041   10.1093/bioinformatics/btu170

1042   78.   De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M. & Van Broeckhoven, C.
1043   NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics* **34,**
1044   2666–2669 (2018). doi: 10.1093/bioinformatics/bty149

1045   79.   Kolmogorov, M., Bickhart, D. M., Behsaz, B., Gurevich, A., Rayko, M., Shin, S. B.,
1046   Kuhn, K., Yuan, J., Polevikov, E., Smith, T. P. L. & Pevzner, P. A. metaFlye: scalable
1047   long-read metagenome assembly using repeat graphs. *Nat Methods* **17,** 1103–1110
1048   (2020). doi: 10.1038/s41592-020-00971-x

1049   80.   Li, H. New strategies to improve minimap2 alignment accuracy. *Bioinformatics* **37,**
1050   4572–4574 (2021). doi: 10.1093/bioinformatics/btab705

1051   81.   Gremme, G., Steinbiss, S. & Kurtz, S. GenomeTools: A Comprehensive Software
1052   Library for Efficient Processing of Structured Genome Annotations. *IEEE/ACM Trans*
1053   *Comput Biol Bioinform* **10,** 645–656 (2013). doi: 10.1109/TCBB.2013.68

1054   82.   Chaumeil, P. A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: A toolkit to
1055   classify genomes with the genome taxonomy database. *Bioinformatics* **36,** 1925–1927
1056   (2020). doi: 10.1093/bioinformatics/btz848

1057   83.   Chklovski, A., Parks, D. H., Woodcroft, B. J. & Tyson, G. W. CheckM2: a rapid,
1058   scalable and accurate tool for assessing microbial genome quality using machine
1059   learning. *bioRxiv* 2022.07.11.499243 (2022). doi:10.1101/2022.07.11.499243 doi:
1060   10.1101/2022.07.11.499243

1061   84.   Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. DRep: A tool for fast and
1062   accurate genomic comparisons that enables improved genome recovery from
1063   metagenomes through de-replication. *ISME Journal* **11,** 2864–2868 (2017). doi:
1064   10.1038/ismej.2017.126

1065   85.   Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J.
1066   eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
1067   Prediction at the Metagenomic Scale. *Mol Biol Evol* **38,** 5825–5829 (2021). doi:
1068   10.1093/molbev/msab293

1069   86.   Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K.,
1070   Cook, H., Mende, D. R., Letunic, I., Rattei, T. & Jensen, L. J. eggNOG 5.0: a
1071   hierarchical, functionally and phylogenetically annotated orthology resource based on
1072   5090 organisms and 2502 viruses. *Nucleic Acids Res* **47,** D309–D314 (2019).

1073   87.   Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput Biol* **7,** e1002195
1074   (2011). doi: 10.1371/journal.pcbi.1002195

1075   88.   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using
1076   DIAMOND. *Nat Methods* **12,** 59 (2015). doi: 10.1038/nmeth.3176

1077   89.   Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching
1078   for the analysis of massive data sets. *Nat Biotechnol* **35,** 1026–1028 (2017). doi:
1079   10.1038/nbt.3988

1080   90.   Hyatt, D., Chen, G.-L., LoCascio, P. F., Land, M. L., Larimer, F. W. & Hauser, L. J.
1081   Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC*
1082   *Bioinformatics* **11,** 119 (2010). doi: 10.1186/1471-2105-11-119

1083   91.   Bengtsson-Palme, J., Ryberg, M., Hartmann, M., Branco, S., Wang, Z. & Godhe, A.
1084   Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS

sequences of fungi and other eukaryotes for analysis of environmental sequencing
data. *Meth Ecol Evol* **4,** (2013).

92. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.
Genome Biol **20,** 1–13 (2019).

93. Lu, J. & Salzberg, S. L. Ultrafast and accurate 16S rRNA microbial community
analysis using Kraken 2. *Microbiome* **8,** 124 (2020). doi: 10.1186/s40168-020-00900-
2

94. Vallenet, D., Calteau, A., Dubois, M., Amours, P., Bazin, A., Beuvin, M., Burlot, L.,
Bussell, X., Fouteau, S., Gautreau, G., Lajus, A., Langlois, J., Planel, R., Roche, D.,
Rollin, J., Rouy, Z., Sabatet, V. & Médigue, C. MicroScope: an integrated platform for
the annotation and exploration of microbial gene functions through genomic,
pangenomic and metabolic comparative analysis. *Nucleic Acids Res* **48,** D579–D589
(2020). doi: 10.1093/NAR/GKZ926

95. Makarova, K. S., Galperin, M. Y. & Koonin, E. V. Comparative genomic analysis of
evolutionarily conserved but functionally uncharacterized membrane proteins in
archaea: Prediction of novel components of secretion, membrane remodeling and
glycosylation systems. *Biochimie* **118,** 302–312 (2015). doi:
10.1016/j.biochi.2015.01.004

96. Zhang, H., Yohe, T., Huang, L., Entwistle, S., Wu, P., Yang, Z., Busk, P. K., Xu, Y. &
Yin, Y. dbCAN2: a meta server for automated carbohydrate-active enzyme
annotation. *Nucleic Acids Res* **46,** W95–W101 (2018). doi: 10.1093/nar/gky418

97. Saier, M. H., Reddy, V. S., Tsu, B. V., Ahmed, M. S., Li, C. & Moreno-Hagelsieb, G.
The Transporter Classification Database (TCDB): recent advances. *Nucleic Acids Res*
44, D372–D379 (2016). doi: 10.1093/nar/gkv1103

98. Dombrowski, N., Williams, T. A., Sun, J., Woodcroft, B. J., Lee, J.-H., Minh, B. Q.,
Rinke, C. & Spang, A. Undinarchaeota illuminate DPANN phylogeny and the impact
of gene transfer on archaeal evolution. *Nat Commun* **11,** 3939 (2020). doi:
10.1038/s41467-020-17408-w

99. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for
comparative genomics. *Genome Biol* **20,** 238 (2019). doi: 10.1186/s13059-019-1832-
y

100. Pausan, M. R., Csorba, C., Singer, G., Till, H., Schöpf, V., Santigli, E., Klug, B.,
Högenauer, C., Blohs, M. & Moissl-Eichinger, C. Exploring the Archaeome: Detection
of Archaeal Signatures in the Human Body. *Front Microbiol* **10,** (2019). doi:
10.3389/fmicb.2019.02796

101. Arce, M. I., von Schiller, D., Bengtsson, M. M., Hinze, C., Jung, H., Alves, R. J. E.,
Urich, T. & Singer, G. Drying and Rainfall Shape the Structure and Functioning of
Nitrifying Microbial Communities in Riverbed Sediments. *Front Microbiol* **9,** (2018).
doi: 10.3389/fmicb.2018.02794

102. Leininger, S., Urich, T., Schloter, M., Schwark, L., Qi, J., Nicol, G. W., Prosser, J. I.,
Schuster, S. C. & Schleper, C. Archaea predominate among ammonia-oxidizing
prokaryotes in soils. *Nature* **442,** 806–809 (2006).

103. Walters, W. A., Caporaso, J. G., Lauber, C. L., Berg-Lyons, D., Fierer, N. & Knight, R.
PrimerProspector: de novo design and taxonomic analysis of barcoded polymerase
chain reaction primers. *Bioinformatics* **27,** 1159–1161 (2011).

104. Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C., Al-Ghalith, G. A.,
Alexander, H., Alm, E. J., Arumugam, M. & Asnicar, F. *QIIME 2: Reproducible,*

1132		*interactive, scalable, and extensible microbiome data science.* (PeerJ Preprints,
1133		2018).
1134	105.	Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A. &
1135		Holmes, S. P. DADA2: High-resolution sample inference from Illumina amplicon data.
1136		*Nat Methods* **13,** 581–583 (2016). doi: doi:10.1038/nmeth.3869
1137	106.	Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A. & B.J, C. Simple statistical
1138		identification and removal of contaminant sequences in marker-gene and
1139		metagenomics data. *Microbiome* **6,** 226 (2018). doi: 10.1186/s40168-018-0605-2
1140	107.	Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. &
1141		Glockner, F. O. The SILVA ribosomal RNA gene database project: improved data
1142		processing and web-based tools. *Nucleic Acids Res* **41,** D590–D596 (2013). doi:
1143		10.1093/nar/gks1219
1144	108.	Robeson, M. S., O'Rourke, D. R., Kaehler, B. D., Ziemski, M., Dillon, M. R., Foster, J.
1145		T. & Bokulich, N. A. RESCRIPt: Reproducible sequence taxonomy reference
1146		database management. *PLoS Comput Biol* **17,** e1009581 (2021). doi:
1147		10.1371/journal.pcbi.1009581
1148	109.	Bokulich, N., Dillon, M., Bolyen, E., Kaehler, B., Huttley, G. & Caporaso, J. q2-
1149		sample-classifier: machine-learning tools for microbiome classification and regression.
1150		*J Open Source Softw* **3,** 934 (2018). doi: 10.21105/joss.00934
1151	110.	Lozupone, C. & Knight, R. UniFrac: a New Phylogenetic Method for Comparing
1152		Microbial Communities. *Appl Environ Microbiol* **71,** 8228–8235 (2005). doi:
1153		10.1128/AEM.71.12.8228-8235.2005
1154	111.	Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 - Approximately maximum-
1155		likelihood trees for large alignments. *PLoS One* **5,** (2010). doi:
1156		10.1371/journal.pone.0009490
1157	112.	Heidrich, V., Karlovsky, P. & Beule, L. 'SRS' R Package and 'q2-srs' QIIME 2 Plugin:
1158		Normalization of Microbiome Data Using Scaling with Ranked Subsampling (SRS).
1159		*Applied Sciences* **11,** 11473 (2021). doi: 10.3390/app112311473
1160	113.	Fernandes, A. D., Macklaim, J. M., Linn, T. G., Reid, G. & Gloor, G. B. ANOVA-Like
1161		Differential Expression (ALDEx) Analysis for Mixed Population RNA-Seq. *PLoS One*
1162		**8,** e67019 (2013). doi: 10.1371/journal.pone.0067019
1163	114.	Lin, H. & Peddada, S. Das. Analysis of compositions of microbiomes with bias
1164		correction. *Nat Commun* **11,** 3514 (2020). doi: 10.1038/s41467-020-17041-7
1165	115.	Mallick, H., Rahnavard, A., McIver, L. J., Ma, S., Zhang, Y., Nguyen, L. H., Tickle, T.
1166		L., Weingart, G., Ren, B. & Schwager, E. H. Multivariable association discovery in
1167		population-scale meta-omics studies. *Biorxiv* (2021).
1168	116.	Martino, C., Morton, J. T., Marotz, C. A., Thompson, L. R., Tripathi, A., Knight, R. &
1169		Zengler, K. A Novel Sparse Compositional Technique Reveals Microbial
1170		Perturbations. *mSystems* **4,** (2019). doi: 10.1128/mSystems.00016-19
1171	117.	Wang, H., Bagnoud, A., Ponce-Toledo, R. I., Kerou, M., Weil, M., Schleper, C. &
1172		Urich, T. Linking 16S rRNA Gene Classification to *amoA* Gene Taxonomy Reveals
1173		Environmental Distribution of Ammonia-Oxidizing Archaeal Clades in Peatland Soils.
1174		*mSystems* **6,** (2021). doi: 10.1128/mSystems.00546-21
1175	118.	Rinke, C., Schwientek, P., Sczyrba, A., Ivanova, N. N., Anderson, I. J., Cheng, J.-F.,
1176		Darling, A., Malfatti, S., Swan, B. K., Gies, E. A., Dodsworth, J. A., Hedlund, B. P.,
1177		Tsiamis, G., Sievert, S. M., Liu, W.-T., Eisen, J. A., Hallam, S. J., Kyrpides, N. C.,
1178		Stepanauskas, R., Rubin, E. M., Hugenholtz, P. & Woyke, T. Insights into the

phylogeny and coding potential of microbial dark matter. *Nature* **499,** 431–437 (2013).
doi: 10.1038/nature12352

119. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol Biol Evol* **30,** 772–780 (2013). doi:
10.1093/molbev/mst010

120. Criscuolo, A. & Gribaldo, S. BMGE (Block Mapping and Gathering with Entropy): a
new software for selection of phylogenetic informative regions from multiple sequence
alignments. *BMC Evol Biol* **10,** 210 (2010). doi: 10.1186/1471-2148-10-210

121. Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von
Haeseler, A. & Lanfear, R. IQ-TREE 2: New Models and Efficient Methods for
Phylogenetic Inference in the Genomic Era. *Mol Biol Evol* **37,** 1530–1534 (2020). doi:
10.1093/molbev/msaa015

122. Pruesse, E., Peplies, J. & Glöckner, F. O. SINA: accurate high-throughput multiple
sequence alignment of ribosomal RNA genes. *Bioinformatics* **28,** 1823–1829 (2012).

123. Larsson, A. AliView: a fast and lightweight alignment viewer and editor for large
datasets. *Bioinformatics* **30,** 3276–3278 (2014). doi: 10.1093/bioinformatics/btu531

124. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods*
14, 587–589 (2017). doi: 10.1038/nmeth.4285

125. Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W. & Gascuel, O.
New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies:
Assessing the Performance of PhyML 3.0. *Syst Biol* **59,** 307–321 (2010). doi:
10.1093/sysbio/syq010

1202

1203

1204