PDK-1/S6K and mTORC1 bypass systemic growth restrictions

to promote regeneration

Or how can some cells proliferate, when nobody else can ?

Ananthakrishnan Vijayakumar Maya [1,2,3], Liyne Nogay [1,2,3], Lara Heckmann [1,2,3], Isabelle Grass [1,2], Katrin

Kierdorf [4,5], Jörg Büscher [6], Anne-Kathrin Classen [1,2,5,7] *

* Corresponding author: anne.classen@biologie.uni-freiburg.de

1 Hilde-Mangold-Haus, University of Freiburg, 79104 Freiburg, Germany

2 Faculty of Biology, University of Freiburg, 79104 Freiburg, Germany

3 International Max Planck Research School for Immunobiology, Epigenetics, and Metabolism, Freiburg,

Germany

4 Institute of Neuropathology, Faculty of Medicine, University of Freiburg, Freiburg, Germany

5 CIBSS Centre for Integrative Biological Signalling Studies, University of Freiburg, Germany

6 Max Planck Institute of Immunobiology and Epigenetics, Freiburg, Germany.

7 BIOSS Centre for Biological Signalling Studies, University of Freiburg, Germany

Abstract

Tissue damage and inflammation trigger systemic signals that induce catabolic breakdown and nutrient release in distant organs, a process well-characterized in the context of tumor cachexia. While mechanisms allowing tumors to circumvent these systemic growth restrictions are known, the physiological processes that overcome inflammation-induced growth restrictions to support tissue repair and regeneration remain largely unexplored. In our study, we use a model of tissue inflammation and regeneration in developing *Drosophila* imaginal discs to dissect the key metabolic and signaling adaptations that help tissue overcome systemic growth restrictions. Our findings reveal a unique metabolic strategy used by rapidly proliferating cells in the regenerating domain. Instead of relying on the conventional Insulin-PI3K-Akt signaling pathway, these cells utilize a JAK/STAT-PDK1-S6K axis. This adaptation facilitates sustained protein synthesis and cellular growth despite the systemic catabolism associated with low insulin signaling. Specifically, we find that catabolic breakdown of the fat body is driven by the insulin-binding factor Impl2, which is expressed at the site of inflammatory damage. Notably, regenerative proliferation is also supported by mTORC1 activity and is associated with the upregulation of amino acid transporters in proliferating cells of the regenerating domain. These amino acid transporters align with a specific amino acid metabolite signature in the hemolymph, revealing a specialized metabolic program that meets the demands of fast-proliferating cells. Our work provides insight into how regenerating tissues rewire signaling pathways and adapt their metabolic growth to coordinate tissue repair with a conserved systemic nutrient provision response. These findings have important implications for understanding human diseases such as chronic wounds and cancer.

Introduction

Tissue damage and inflammation trigger a dynamic interplay between cellular signals and cell behaviors to promote repair and regeneration. Proper orchestration of these responses is crucial, as failure can lead to chronic wound healing pathologies, fibrosis, or diseases like cancer (Eming et al, 2017; Huang et al, 2022; MacCarthy-Morrogh & Martin, 2020; Pena & Martin, 2024; Rybinski et al, 2014). To better understand these diseases, a wide range of recent studies aim to dissect the pathological reprogramming of relevant metabolic circuits (Martinez-Reyes & Chandel, 2021; Stine et al, 2022; Wong & Verheyen, 2021). As cancers frequently adopt tissue repair pathways for their own advantage, these studies may also indirectly offer insights into the metabolism of tissue repair. However, a surprising gap exists in our knowledge about the precise metabolic adaptations employed by the normal physiological programs of tissue repair and regeneration (Eming et al, 2021; Eming et al, 2017; Meacham et al, 2022; Zhu & Thompson, 2019). With this study, we aim to provide insight into the local and systemic metabolic adaptations during physiological tissue repair and regeneration.

Two pathways central to metabolism and cellular growth in health and disease are the Insulin/PI3K/Akt and mTORC1 signaling pathways. Both are intimately linked by their shared effector ribosomal protein S6 kinase (S6K), which drives protein synthesis and cellular growth by activating protein translation. Insulin/PI3K/Akt is activated by binding of Insulin to its receptor (InR), thereby stimulating Phosphoinositide-3-kinase (PI3K) to produce Phosphatidylinositol (3,4,5)-trisphosphate (PIP3) at the plasma membrane. PIP3 recruits Phosphoinositide-dependent kinase 1 (PDK-1), but also activates Akt, which inhibits nuclear translocation of the transcription factor dFOXO and promotes survival. PDK-1 in turn phosphorylates and activates S6K, yet, for S6K to function optimally, at least two of its residues need to be phosphorylated. In addition to PDK-1, the mechanistic target of rapamycin complex 1 (mTORC1) and c-Jun N-terminal kinases (JNK) phosphorylate activating residues on S6K (Holz et al, 2005; Hopkins et al, 2020; Hoxhaj & Manning, 2020; Manning & Cantley, 2007; Wu et al, 2022). mTORC1 is central to anabolic growth by modulating cellular responses to nutrient availability and environmental cues (Saxton & Sabatini, 2017). While it is sensitive to amino acid availability, it can also be stimulated by Akt via crosstalk with Insulin/PI3K/Akt signaling. The interplay of Insulin/PI3K/Akt and mTORC1 signaling in metabolism is highly conserved from invertebrates to vertebrates. Previous studies demonstrated that Insulin signaling is active in *Drosophila* epidermal wounds, where it facilitates epithelial sealing by regulating actin dynamics immediately after damage (Kakanj et al, 2016; Nakamura et al, 2020). Recent evidence reveals an elevation of ribosomal S6K phosphorylation in mammalian skin wounds and correlating it with successful healing (Ring et al, 2023). Yet, the relevant molecular mechanisms and precise metabolic adaptations during repair responses have yet to be investigated.

Drosophila models have significantly advanced our understanding of tissue repair, regeneration and metabolism (Fox et al, 2020; Worley & Hariharan, 2022). Specifically, studies in developing wing imaginal

discs or the adult gut have highlighted the role of two key signaling pathways - JNK/AP1 and JAK/STAT

signaling - in epithelial repair and proliferation. The JNK/AP1 pathway is an early stress response pathway

which induces a senescence-like cell cycle arrest the site of tissue damage (Cosolo et al, 2019; Floc'hlay

et al, 2023; Jaiswal et al, 2023). The senescence-like state is associated with the production of

extracellular matrix and cytoskeletal remodelers, which promote the sealing of epithelial wounds

(Floc'hlay et al, 2023; Nakamura et al, 2020; Pena & Martin, 2024; Stevens & Page-McCaw, 2012).

Importantly, these cells also produce Unpaired (Upd) cytokines. Their paracrine activity activates

JAK/STAT in cells surrounding the site of inflammatory damage, where it promotes cell survival and

compensatory proliferation during tissue regeneration (Crucianelli et al, 2022; Floc'hlay et al, 2023;

Herrera & Bach, 2019; Jaiswal et al, 2023; La Fortezza et al, 2016; Santabarbara-Ruiz et al, 2015; Worley

et al, 2022). Compensatory proliferation itself is characterized by an accelerated cell cycle with

shortened S-phases and gap phases (**Fig S1A**) (Crucianelli et al, 2022; Herrera et al, 2013; Sustar et al,

2011). The distinct functional and proliferative demands of senescent and fast-proliferating cell

populations within the damaged tissue raises questions about how these distinct cell populations

metabolically adapt to successfully support tissue repair.

Tissue repair and tumor development share striking similarities; in fact, tumors have long been compared

to non-healing wounds (Dvorak, 1986). Accordingly, *Drosophila* tumor models activate JNK/AP1 and

JAK/STAT signaling, which promotes tumor progression (Enomoto & Igaki, 2022; Herrera & Bach, 2019; La

Marca & Richardson, 2020). To support their growth, tumors secrete signaling molecules that initiate

inter-organ signaling and systemic metabolic responses (Bilder et al, 2021; Hodgson et al, 2021; Liu et al,

2022). For instance, the Insulin-like peptide 8 (Dilp8), when secreted by tumors, disrupts hormone

balance by reducing Ecdysone and Insulin production through direct effects on the ring gland and Insulin-

producing cells (IPCs), with the effect of halting developmental progression of the tumor-bearing host

(Colombani et al, 2012; Garelli et al, 2012). Other cytokines, such as Upd2, Upd3, PDGF- and VEGF-

related factor 1 (Pvf1), Ecdysone-inducible gene L2 (Impl2), matrix metalloprotease 1 (MMP-1) or the

TNFα homologue Eiger (Egr), directly or indirectly trigger catabolic lipolysis and proteolysis to promote

nutrient release from muscles and the so-called fat body, an adipose tissue central for nutrient storage

and energy homeostasis (Agrawal et al, 2016; Ding et al, 2021; Figueroa-Clarevega & Bilder, 2015;

Hodgson et al, 2021; Lodge et al, 2021; Rajan & Perrimon, 2012; Romao et al, 2021; Song et al, 2019).

Amino acids or sugars are released in this manner and subsequently absorbed by tumors and facilitate

their anabolic growth (Cong et al, 2021; Khezri et al, 2021; Newton et al, 2020). This inter-organ signaling

network and metabolic state resembles cachexia, a clinical syndrome characterized by weight loss,

muscle atrophy and fatigue, typically observed in chronic inflammatory conditions, including cancer

(Argiles et al, 2014; Bilder et al, 2021; Hodgson et al, 2021; Liu et al, 2022; Setiawan et al, 2023). However,

the systemic and metabolic regulation of non-pathological tissue repair and regeneration remains poorly

understood. Previous studies suggest that cytokines may also be released upon physiological tissue

damage or starvation (Colombani et al, 2012; Garelli et al, 2012; Gontijo & Garelli, 2018; Hackney &

Cherbas, 2014; Halme et al, 2010; Sun & Poss, 2023). Further studies demonstrated the activation of fat body catabolic pathways with effects on Methionine, S-adenosylmethionine and Kynurenine metabolism promote effective imaginal disc regeneration (Kashio & Miura, 2020; Kashio et al, 2016). While these studies affirm that tumors may hijack tissue repair mechanisms, the underlying local and systemic metabolic adaptations that evolution developed to promote physiological processes of repair and regeneration remain little characterized. In our study we combine genetic analysis, quantitative imaging, untargeted metabolomics and information from single-cell RNA sequencing data to outline the local and systemic adaptations that selectively support fast-proliferating cells during regeneration through an Insulin-independent JAK/STAT-PDK1-S6K signaling axis.

Results

(1) The proliferating domain maintains high levels of protein synthesis during regeneration

Does tissue damage alter metabolic processes to drive regenerative proliferation within epithelial cells? To induce inflammation and regeneration, we expressed the *Drosophila* homologue of TNF-α, known as Eiger (Egr), for a 24 h period within the imaginal wing pouch using the *rn-GAL4* driver (Cosolo et al, 2019; Smith-Bolton et al, 2009). As previously observed, this led to significant cell death, accompanied by the activation of the JNK/AP-1 activity reporter TRE-RFP (**Fig 1A-C; Fig S1B,C**) (Chatterjee & Bohmann, 2012; Cosolo et al, 2019; Jaiswal et al, 2023; La Fortezza et al, 2016). JNK activation at the center of the pouch pauses the cell cycle in G2, as evidenced by FUCCI cell cycle analysis and the absence of EdU incorporation (**Fig 1A,B; Fig S1D-F**) (Cosolo et al, 2019; Floc'hlay et al, 2023; Jaiswal et al, 2023; Zielke et al, 2014). The JNK signaling domain exhibits characteristic features of cellular senescence, including the expression of senescence-associated-β-galactosidase (SA-β-gal) (**Fig S1G,H**), the matrix metalloprotease MMP-1 and mitogenic cytokines of the Upd family (Floc'hlay et al, 2023; Jaiswal et al, 2023). They induce compensatory proliferation in surrounding cells, which can be detected by rapid EdU incorporation (**Fig 1A,B**) (Crucianelli et al, 2022). We found that the majority of cells undergoing proliferation were located within 40 μm from the outer border of high JNK signaling cells (**Fig 1D**). To facilitate quantification of cell behaviors by TRE-RFP segmentation (see Materials and Methods) and account for disc size variation among the many experiments carried out in this study, we define a conservative 20μm band outside the high JNK signaling domain as the 'proliferative domain' (PD^{egr}) for the remainder of this study.

How can these rapidly proliferating cells be supported by biosynthesis and cellular growth? To answer this question, we monitored protein translation using O-propargyl-puromycin (OPP)-incorporation assays (Aviner, 2020). This assay revealed uniform protein synthesis levels throughout control wing imaginal discs. In the regenerating domain of *egr*-expressing discs protein translation proceeded at levels similar to control discs and was mildly reduced within the domain of high JNK-signaling, senescent cells (**Fig 1E-G**). Surprisingly, OPP incorporation significantly decreased in the notum of *egr*-expressing discs and

aligned with markedly low levels of EdU incorporation in these domains (**Fig 1H-M**). This observation suggests that protein synthesis rates as well as cell cycle entry and progression slow down significantly in cell populations at a distance from the site of tissue damage.

To test whether reduced proliferation in the notum of *egr*-expressing discs is connected to a senescent cell cycle arrest, we closely monitored JNK/AP-1 activation (TRE-RFP), cell cycle dynamics (FUCCI) and the senescence marker SA-β-GAL in the notum. However, we observed no elevation of JNK signaling nor SA-β-GAL activity, and FUCCI profiles were similar in control and *egr*-expressing disc (**Fig 1H,I; Fig S1G-J**). This suggests that the reduced proliferation in nota is distinct from JNK/AP-1-associated senescence responses in *egr*-expressing cells of the pouch. Hence, we conclude that the notum state may be more accurately described by the term quiescence.

Combining these observations, at least three distinct cell populations exist in the wing discs with inflammatory lesion, each characterized by different proliferative potentials and cellular programs: 1) a senescent cell population at the center of damage, exhibiting high protein synthesis, high JNK/AP-1 activation and a G2 cell cycle arrest; 2) an actively cycling cell population exhibiting high protein synthesis and low JNK/AP-1 activation; 3) a distal quiescent cell population displaying low rates of protein synthesis, low JNK/AP-1 activation and slow cycling activity (**Fig 1N,O**).

(2) Peripheral organs reduce growth

The lack of damage-associated signals in the notum may suggest that not local, but paracrine and even systemic changes reduce cell proliferation and biosynthesis. To investigate this, we first evaluated mitotic activity and protein synthesis in other peripheral tissues, such as the eye and leg imaginal disc. In larvae carrying *egr*-expressing wing discs, the mitotic wave anterior to the morphogenetic furrow in the developing eye was arrested, which also correlated with reduced protein synthesis (**Fig 2A-F**). Similarly, protein synthesis was diminished in leg discs, if compared to leg discs of wild type larvae (**Fig S2A,B**). Taken together, these findings reveal a widespread decline in cell proliferation and protein synthesis in *egr*-expressing larvae including distant tissues. These observations are consistent with previous reports on reduced eye disc and notum sizes after activation of stress responses in the wing pouch (Boulan et al, 2019).

We previously reported that *egr*-expressing larvae also exhibit a significant delay in initiating pupariation (La Fortezza et al, 2016). Accordingly, JNK activation in *egr*-expressing wing discs correlates with a pronounced increase in Dilp-8 expression likely underlying the delay in pupariation (**Fig 2G,H**) (Floc'hlay et al, 2023). Yet, previous studies also suggest, that Dilp8 and Egr itself may inhibit IPCs in the brain, indicating that *egr*-expressing larvae may experience low systemic Insulin signaling, limiting anabolic growth despite nutrient intake by feeding (Agrawal et al, 2016; Figueroa-Clarevega & Bilder, 2015; Ingaramo et al, 2020; Rajan & Perrimon, 2012; Vallejo et al, 2015). To understand if reduced protein

synthesis in distant tissues of *egr*-expressing larvae may indeed correspond to reduced systemic Insulin
signaling, we examined expression levels and nuclear localization of the dFOXO transcription factor, a
key mediator of Insulin signaling. Indeed, we observed elevated nuclear levels of FOXO in the peripheral
organs of the fat body dissected from *egr*-expressing larvae, if compared to wild-type controls (**Fig 2I-K**).
Combined, these observations strongly indicate that attenuation of Insulin signaling may occur within
distant non-inflamed tissues within 24 h of *egr*-expression, similar to systemic changes induced by
inflammatory tumors or related types of tissue damage (Ding et al, 2021; Figueroa-Clarevega & Bilder,
2015; Garelli et al, 2012; Hodgson et al, 2021; Jiang et al, 2022; Khezri et al, 2021; Lodge et al, 2021; Song
et al, 2019) .

(3) Amino acid importers, mTORC1 signaling and ribosomal S6 phosphorylation support
regenerative proliferation
How can the proliferative domain maintain protein translation, anabolic growth and proliferation a
systemic environment characterized by reduced protein synthesis and proliferation? To answer this
question, we analyzed regulatory inputs into protein translation and ribosome activity. We found that
transcript levels of S6K were upregulated in the wound-associated clusters of *egr*-expressing wing
imaginal discs in previously published scRNA-seq data sets (**Fig S3A,B**) (Floc'hlay et al, 2023), suggesting
that *S6K* may be regulated by the tissue repair program. Importantly, staining for phosphorylated
ribosomal S6 (p-S6) indeed revealed that the proliferating domain of *egr*-expressing discs exhibited
strongly elevated p-S6 levels (**Fig 3A-C**). This finding is in line with recent observations during wound
healing in mammalian skin (Ring et al, 2023). This suggests existence of a conserved mechanism that
elevates ribosomes activation in proliferating cells of the regenerating domain to specifically support their
growth.

S6K is activated by Insulin/PI3K/Akt and mTORC1 signaling (**Fig S3C**) (Liu & Sabatini, 2020; Wu et al, 2022).
Therefore, we assessed the activity of these two pathways in *egr*-expressing discs. To demonstrate that
the mTORC1 pathway is required to drive protein synthesis during wing disc regeneration, we inhibited
mTORC1 activity by feeding *egr*-expressing larvae the mTOR inhibitor Rapamycin for 24 h during *egr*-
expression (Li et al, 2014; Wu et al, 2022). Protein synthesis in control imaginal discs was only mildly
reduced by Rapamycin treatment. However, a pronounced reduction in protein synthesis could be
observed in the proliferating domain of *egr*-expressing discs upon Rapamycin treatment, demonstrating
that mTORC1 activity is specifically required to drive protein synthesis in the proliferating domain of *egr*-
expressing discs (**Fig 3D-H**).

mTORC1 activity itself is induced by uptake of nutrients and different amino acids (Jewell et al, 2013;
Lama-Sherpa et al, 2023; Liu & Sabatini, 2020; Shimobayashi & Hall, 2016; Yue et al, 2022). We thus
investigated if amino acids could be taken up preferentially by the proliferating cells in *egr*-expressing
discs. Indeed, transcripts of solute carrier group of membrane transport proteins (SLC) were upregulated

in *egr*-expressing discs, such as *CG15279*, a putative glycine and proline transporter of the SLC6 family,
path, a putative alanine and glycine transporter of the SLC 36 family, *mnd*, an amino acid/polyamine
transporter involved in leucine import and *CG5535*, a putative lysine, arginine, and ornithine transporter
of the SLC7 family (**Fig S3D**). Importantly, we could verify upregulation of CG5535 in the proliferating
domain by immunofluorescence (**Fig 3I-K**) and were similarly able to detect upregulation of CG1139
(Arcus), a putative alanine, glycine and proline transporter of the SLC36 family in *egr*-expressing discs (**Fig
S3E,F**). These observations suggest that amino acids could be preferentially imported by proliferating
cells. Furthermore, CG3168, a potential sugar transporter of the SLC 22 family and TRET-1, predicted to
transport the disaccharide trehalose into cells (**Fig S3G-I, Fig S3D**), were strongly upregulated, indicating
that additional energy sources can be imported preferentially by proliferating cells. Availability of energy
would promote mTORC1 activity, as ATP production serves as a positive regulator of mTORC1 activity by
inhibiting the mTORC1 antagonist AMPK (Jewell et al, 2013; Liu & Sabatini, 2020; Shimobayashi & Hall,
2016).
249
If the regeneration domain displays elevation of specific amino acid transporters, then what kind of
changes to amino acid composition can be observed in the hemolymph of *egr*-expressing larvae which
could be taken up by proliferating cells? We performed an untargeted metabolomic analysis of the
hemolymph from both control and *egr*-expressing larvae. Our results revealed an enrichment of many
amino acids in *egr*-expressing larval hemolymph, including those which have been reported to directly or
indirectly activate mTORC1 signaling (for example, Leucine, Arginine, Glutamine (Lama-Sherpa et al,
2023; Yue et al, 2022)). Surprisingly, many come enriched in form of mixed dipeptide species. Moreover,
elevated levels of dipeptide species and free amino acids associated with glutamate metabolism were
observed, including increased concentrations of glutamine, glutamate, and glutamyl dipeptides (**Fig 3L**).
These findings suggest that metabolic processes centered around glutamate conversion and
glutaminolysis may support regenerative metabolism (Altman et al, 2016). Notably, we also observed a
strong signature of urea cycle and arginine metabolism, characterized by high levels of ornithine, N-
acetyl-arginine, aspartate, but low levels of argininosuccinic acid, citrulline, and N-acetyl-ornithine. This
metabolite pattern together with the elevated expression of the ornithine transporter CG5535 in *egr*-
expressing discs could support an increased flux of the urea cycle from ornithine towards arginine to
provide a nitrogen acceptor for glutaminolysis, and potentially function to produce arginine, polyamines,
nitric oxide or non-essential amino acids (Keshet et al, 2018; Lieu et al, 2020; Marti & Reith, 2021). This
altered metabolite composition in the hemolymph of *egr*-expressing larvae may represent a highly flexible
supply of building blocks not only for protein synthesis, but also for energy and co-factor production or
glutathione redox metabolism. Combined, our data support a potential role for mTORC1 activation,
amino acid uptake and glutamate and arginine metabolism in supporting the rapid growth and
proliferation of the regenerating domain.
272
(4) Impl2 from wing discs induces a catabolic phenotype in the larval fat body

So how may these amino acids get enriched in the hemolymph of *egr*-expressing larvae? Previous studies

have shown that cytokines can induce the catabolic conversion and release of nutrients from distant

organs. Specifically, the break-down of fat body, muscle degradation and autophagy of imaginal discs

have been established as nutrient sources driving tumor growth (Ding et al, 2021; Figueroa-Clarevega &

Bilder, 2015; Katheder & Rusten, 2017; Khezri et al, 2021; Lodge et al, 2021; Newton et al, 2020; Song et

al, 2019). We wanted to better understand if similar catabolic changes may be induced by *egr*-expressing

larvae to facilitate regeneration. To this end, we employed Nile Red staining to visualize lipid droplets

within the fat body, performed F-actin staining to observe the morphology of body wall muscles, and

visualized the autophagy marker Ref(2)P in peripheral imaginal disc tissue. We did not observe changes

to muscle morphology nor imaginal disc autophagy in *egr*-expressing larvae, suggesting that these tissues

may not serve as a primary source for nutrients during early stages of tissue repair (**Fig S4A-D**). However,

in the fat body, we observed morphological changes consistent with nutrient mobilization, such as

increased droplet size and roundness, indicating that the observed hemolymph changes may be derived

from fat body catabolism (**Fig 4A-D**)

In addition to the cachexic cytokines Egr and Pvf1 (Agrawal et al, 2016; Song et al, 2019), expression of all

Upd family members, including Upd3, and the Insulin-growth factor binding protein Impl2, was strongly

elevated in *egr*-expressing discs (**Fig 4E-J, Fig S4E**). Impl2 has been previously described as contributing

to fat body break-down in adult flies with *Ras^{V12}, scrib* or Yki-dependent tumors (Ding et al, 2021; Figueroa-

Clarevega & Bilder, 2015; Kwon et al, 2015), while Upd3 induces JAK/STAT signaling and acts on fat bodies

during tumor growth in larval and adult hosts (Ding et al, 2021; Romao et al, 2021). Indeed, we observed

activation of the JAK/STAT reporter 10xSTAT-GFP in the fat body of *egr*-expressing larvae, suggesting that

egr-expressing discs may affect fat body signaling as well (**Fig 4K-M**). Moreover, when we ectopically

expressed Impl2 for 24 h using the wing-specific driver rn-GAL4, we found that this was sufficient to

induce morphological changes in the fat body characteristic of nutrient mobilization (**Fig 4N-Q**). These

observations indicate that the production of Impl2 and Upd cytokines from *egr*-expressing discs are likely

responsible to drive nutrient mobilization from the fat body, thereby enriching the hemolymph with

metabolites that support regenerative proliferation.

(5) PDK-1 acts independent of Insulin/PI3K/Akt signaling in S6K activation and is regulated by

JAK/STAT signaling

Nutrient availability, along with mTORC1 activation, are not sufficient to elevate levels of protein

synthesis in the proliferating domain. S6K requires activation by mTORC1, but also by the kinase PDK-1,

the latter being integral to Insulin/PI3K signaling (Hopkins et al, 2020; Hoxhaj & Manning, 2020; Wu et al,

2022). If *egr*-expressing discs induce low systemic Insulin signaling, could the proliferating domain

somehow activate Insulin/PI3K signaling to drive PDK-1 activity towards S6K?

To understand if Insulin/PI3K and downstream signaling effectors are active in proliferating cells, we
examined the nuclear localization of dFOXO, the downstream effector of Insulin/PI3K/Akt activity in the
proliferating domain. Strikingly, we observed that both cytoplasmic and nuclear levels of dFOXO were
elevated, suggesting that both dFOXO levels, as well as rates of nuclear translocation were upregulated
(**Fig 5A-C**). The high nuclear dFOXO levels in the proliferating domain strongly suggests that also locally,
Insulin/PI3K signaling is low in proliferating cells. This is consistent with a systemic environment of low
Insulin signaling. Our conclusion is further supported by experiments showing that InR levels remained
unchanged, and levels of Akt and phosphorylated Akt (P-Akt) were even reduced in the regenerating
domain (**Fig S5A-H**). Consequently, the proliferating domain has not evolved an Insulin-scavenging
mechanism to maintain high levels of protein synthesis in a growth restrictive environment with low
Insulin signaling.

Given this observation, the question arises as to how else S6K may become co-activated by PDK-1.
Strikingly, we observed that the expression of PDK-1 was strongly upregulated in the proliferating domain
(**Fig 5D-F**). Elevated PDK-1 protein levels may arise from post-translational mechanisms affecting PDK-1
stability, as we did not observe an increase in PDK-1 transcripts (Floc'hlay et al, 2023). We wanted to
identify the upstream regulator of PDK-1 elevation. In *egr*-expressing discs, JAK/STAT signaling is
activated in the proliferating domain to promote cell proliferation and survival (**Fig S5I,J**) (Crucianelli et
al, 2022; Jaiswal et al, 2023; La Fortezza et al, 2016). To test whether JAK/STAT signaling may regulate
PDK-1 levels, we expressed STAT92E using the *en-GAL4* driver in the posterior compartment of wing
imaginal discs, which drives JAK/STAT activation (Jaiswal et al, 2023). We found that JAK/STAT activation
increased the levels of PDK-1-GFP, phospho-S6 and protein synthesis (**Fig 5G-O**). This suggests that
JAK/STAT activation in the proliferating domain increases PDK-1 levels and thereby promotes ribosomal
S6 phosphorylation and protein synthesis. Combined, these observations reveal a novel function for
JAK/STAT in tissue repair and regeneration and outline a pathway whereby the combined action of PDK-1
and mTORC1 drives insulin-independent growth within the proliferating domain.

(6) PDK-1 is necessary and sufficient to drive Insulin-independent regenerative proliferation
To provide evidence that upregulation of PDK-1 levels plays a crucial role in the proliferating domain of
egr-expressing discs, we tested whether ectopic expression of PDK-1 alone was sufficient to drive protein
synthesis and promote growth of wing disc tissues. We therefore expressed either a wild-type form of
PDK-1 or a constitutively active form of PDK-1 for 24 hours in the wing pouch using the *rn-GAL4* driver.
Compared to control discs, we observed higher levels of protein synthesis and larger pouch sizes in both
PDK-1-expressing genotypes. The increased size of the pouch appeared to be supported by cell
proliferation, as EdU incorporation in PDK-1 expressing domains was clearly elevated (**Fig 6A-F, Fig S6A-**
D). These results align with previous studies (Radimerski et al, 2002; Rintelen et al, 2001) and
demonstrate that high PDK-1 levels are sufficient to support a metabolic program characteristic of
regenerative proliferation. Importantly, PDK-1 can auto-activate via transphosphorylation upon PIP3-

dependent recruitment to the plasma membrane (Levina et al, 2022). We therefore analyzed PIP3 levels

using the pleckstrin-homology domain reporter tGPH in *egr*-expressing discs (Britton et al, 2002). tGPH

levels at the membrane were consistently high, suggesting that PIP3 levels are not reduced in *egr*-

expressing discs and that the plasma membrane provides a functional substrate for PDK-1 recruitment

and autoactivation (**Fig S6E,F**).

To provide proof that PDK-1 function is also necessary to support the proliferative domain in *egr*-

expressing discs, we genetically reduced PDK-1 function by establishing heterozygosity for a null allele of

PDK-1 (Rintelen et al, 2001). Importantly, heterozygous control discs did not exhibit difference in protein

synthesis if compared to wild type discs, revealing that heterozygosity for PDK-1 does not limit the growth

during normal development. However, protein synthesis in the regenerating domain of *egr*-expressing

discs heterozygous for PDK-1 was significantly reduced (**Fig 6G-K**). These experiments demonstrate that

specifically the proliferating domain is sensitive to a reduction in PDK-1 levels, and that elevated PDK-1

levels are necessary for protein synthesis during regenerative proliferation. Notably, the fact that

overexpression of a wild-type PDK-1 alone is sufficient to promote proliferation and protein synthesis,

suggests that PDK-1 activity can be regulated via its protein levels alone, and as such potentially

independent of other canonical upstream signaling inputs, such as Insulin/PI3K signaling. Strikingly, PDK-

1 may also be a target of systemic regulation during growth restriction, as we found that PDK-1 levels were

lower in peripheral tissues of *egr*-expressing larvae, such as in the eye discs (**Fig 6L-N**). This indicates that

PDK-1 downregulation in peripheral tissues may be an effector of the systemic changes to reduce protein

synthesis and anabolic growth, further positioning PDK-1 as a central regulator and integrator of the

systemic and local metabolic reprogramming required for tissue repair.

Discussion

Our study provides insights into the local and systemic adaptations of metabolic processes during tissue

repair and regeneration (**Fig 6O,P**). Our findings distinguish adaptations in proliferating cells of the

regenerating domain from senescent or quiescent cells within damaged tissues. Here we show that the

behavior of quiescent-like cells distal to the site of inflammatory damage is characterized by reduced

protein synthesis and cell proliferation and correlates with a low insulin-signaling environment. Previous

work suggest that systemic growth restriction may prevent the progression and overgrowth of undamaged

tissues, thereby synchronizing regeneration with developmental progression (Cao et al, 2022; Colombani

et al, 2012; Garelli et al, 2012; Hackney & Cherbas, 2014; Halme et al, 2010).

Importantly, how the regenerating domain overcomes such a restrictive systemic environment was not

known. Here, we demonstrate that proliferating cells in *egr*-expressing wing discs display a distinct

upregulation of ribosomal S6 protein phosphorylation and maintain high levels of protein synthesis. A

domain characterized by ribosomal S6 protein phosphorylation was recently described in mouse and pig
skin wounds, implicating potential similarities in regenerative programs across species (Ring et al, 2023).
Besides mTORC1, PDK-1 and JNK can act as activators of S6K (Wu et al, 2022). Yet, JNK signaling in
proliferating cells of the regenerating domain is low. Our findings show that the PDK-1 kinase is essential
to support protein translation and growth during regeneration. PDK-1 is typically considered to be part of
the Insulin/AKT/FOXO; yet, in the regenerating domain, cells proliferate independent of Insulin/AKT/FOXO
activity. PDK-1 upregulation appears to act as a substitute compensating the lack of Insulin/AKT/FOXO
signaling and thus acts as an alternative driver of proliferative metabolism to support tissue repair. This
is supported by previous studies on AKT-independent but PDK-1-dependent growth during *Drosophila*
development (Radimerski et al, 2002; Rintelen et al, 2001). Given that PDK-1 may bypass a need for
Insulin/AKT/FOXO, it may not be surprising that PDK-1 upregulation is found in many cancers (Zheng et
al, 2023). In addition to S6K activity, Myc levels are elevated in the regenerating domain (Crucianelli et al,
2022; Smith-Bolton et al, 2009). The function of Myc as a positive regulator of ribosome biogenesis and
protein translation may thus further support growth and proliferation in this domain (Destefanis et al,
2020). In our study, we identify JAK/STAT signaling as an upstream regulator of PDK-1, phosphorylation of
ribosomal S6 protein, and protein synthesis. Our work thus uncovers a pathway whereby JAK/STAT
signaling mediates the known proliferation and survival-promoting function in imaginal disc
development, regeneration, and tumor growth (Herrera & Bach, 2019).
405
The increased expression of nutrient transporters in proliferating cells suggests that the regeneration
program allows for selective uptake of mobilized nutrients from the hemolymph environment. The
hemolymph from *egr*-expressing larvae exhibited enrichment of metabolites associated with glutamate
and arginine metabolism when compared to control larvae. Remarkably, the upregulation of both the urea
cycle metabolite ornithine in the hemolymph and the predicted ornithine-transporter CG5535 in the
regenerating domain suggest a role for urea cycle metabolism during regeneration. Prior research has
shown that dietary supplementation of ornithine significantly enhances wound healing parameters in
mice (Shi et al, 2002; Witte & Barbul, 2003). Earlier studies indicated elevated levels of ornithine,
glutamate, and glutamine in rat wound fluids (Albina et al, 1993). Our findings suggest that glutamate
metabolism may support regeneration, aligning with its described roles in tumor growth, inflammation,
and glutathione oxidative stress defenses (Altman et al, 2016; Ling et al, 2023; Zhu & Thompson, 2019).
By coupling glutaminolysis to the urea cycle, the proliferating cell would be able to channel nitrogen into
different metabolic processes such as non-essential amino acid (NEAA) synthesis via transamination,
thereby extending the advantages of urea cycle use beyond arginine, polyamines, or nitric oxide
production (Keshet et al, 2018).
421
422 Importantly, our study elucidates how wild type cells can proliferate and repair physiological
423 inflammation. The program systemically inhibits growth and proliferation and drives nutrient release form
424 energy-storing organs like the fat body. Undoubtedly, the systemic effects are part of an ancient and

conserved stress response and repair program designed to distribute nutrients from an energy store to the site of tissue damage. Evolutionarily, this program would guarantee a nutrient supply to the regenerating area and therefore prioritize repair, especially if feeding or foraging opportunities for the organism may be scarce or limited by injuries. If activated more long-term by inflammatory pathologies, like tumors, this program turns from a repair program to chronic cachexia and lethal organ wasting. While oncogenic mutations may facilitate nutrient use in such a cachexic environment (Lee et al, 2021; Newton et al, 2020; Santabarbara-Ruiz & Leopold, 2021; Song et al, 2019), our work reveals how normal tissue adapts within the repair program to overcome the systemic inhibition of growth. Our work highlights the JAK/STAT-PDK-1 and mTORC1 axis as crucial mediators of regenerative growth.

 Materials and Methods

Drosophila **maintenance**

All experiments were performed on *Drosophila melanogaster*. Fly strains (see **Table S1**) were maintained

on standard fly food (10L water, 74,5g agar, 243g dry yeast, 580g corn flour, 552ml molasses, 20.7g

Nipagin, 35ml propionic acid) at 18°C – 22°C. Larvae from experimental crosses were allowed to feed on

Bloomington formulation (175.7g Nutry-Fly,1100ml water 20g dry yeast, 1.45g Nipagin in 15ml Ethanol,

4.8ml Propionic acid) and raised at 18°C or 30°C to control GAL80ts-dependent induction of GAL4/UAS.

Our experimental design did not consider differences between sexes unless for genetic crossing

schemes.

Drosophila **genetics**

To induce expression of UAS-constructs, such as *UAS-egr*, under the control of rn-GAL4 in the wing

pouch, experiments were carried out as described in (Cosolo et al, 2019; La Fortezza et al, 2016; Smith-

Bolton et al, 2009) with minor modifications. Briefly, larvae of genotype *rn-GAL4, tub-GAL80^{ts}* and carrying

the desired *UAS*-transgenes were staged with a 6h egg collection and raised at 18°C at a density of 50

larvae/vial. Overexpression of transgenes was induced by shifting the temperature to 30°C for 24h on the

seventh day (D7) after egg deposition (AED) to relieve temperature-sensitive GAL80ts repression of GAL4.

Larvae were dissected after 24 h of *egr*-expression. Control genotypes were generated by crossing *rn-*

GAL4, tub-GAL80^{ts} into a wild type background.

Immunohistochemistry of wing imaginal discs

Wing discs from third instar larvae were dissected and fixed for 15 minutes at room temperature in 4%

paraformaldehyde in PBS. Washes were performed in PBS containing 0.1% TritonX-100 (PBT). The discs

were then incubated with primary antibodies (listed in **Table S2**) in 0.1% PBT, gently mixed overnight at

4°C. During incubation with cross-absorbed secondary antibodies coupled to Alexa Fluorophores at

room temperature for 2 hours, tissues were counterstained with DAPI (0.25 ng/µL, Sigma, D9542). Tissues

were mounted using SlowFade Gold Antifade (Invitrogen, S36936). To ensure comparability in staining

between different genotypes, experimental and control discs were processed in the same vial and

mounted on the same slides whenever possible. Images were acquired using the Leica TCS SP8

Microscope, using the same confocal settings for linked samples and processed using tools in Fiji.

Protein synthesis assays using OPP- Click-iT staining

OPP Assays were performed using Click-iT® Plus OPP Protein Synthesis Assay Kits (Invitrogen Molecular

Probe) according to manufacturer's instructions. Briefly, larvae were dissected and inverted cuticles

were incubated with a 1:1000 dilution of Component A in Schneider's medium for 15 minutes on a

nutator. Larval cuticles were fixed with 4% paraformaldehyde for 15 minutes, rinsed twice in 0.1% PBT,

and permeabilized with 0.5% PBT for 15 minutes. The cuticles were then stained with the Click-iT®

cocktail for 30 minutes at room temperature, protected from light. Further immunohistochemistry

analysis and sample mounting was performed as described above.

EdU labelling

EdU incorporation was performed using the Click-iT Plus EdU Alexa Fluor 647 Imaging Kit. Briefly, larval

cuticles were inverted in Schneider's medium and incubated with EdU (10µM final concentration) at RT

for 15 minutes. Cuticles were then fixed in 4% PFA/PBS for 15 minutes, washed for 30 minutes in PBT

0.5%. EdU-Click-iT labeling was performed according to manufacturer's guidelines. Further

immunohistochemistry analysis and sample mounting was performed as described above.

SA-β-Gal staining

Cell senescence detection kit from Invitrogen (C10850) was used to analyze senescence-associated β-

galactosidase activity. Briefly, larval cuticles were inverted in PBS, fixed with 4% PFA, washed with 1%

BSA (in PBS) and then incubated in working solution for 2h at 37°C, according to manufacturer's

instructions. Washing steps were performed in PBS and PBS containing 0.1% TritonX-100 (PBT). Further

immunohistochemistry analysis and sample mounting was performed as described above.

Fat body Nile Red staining

Early third instar larvae were collected in PBS and dissected, leaving the gut intact to prevent fat body

loss. Inverted cuticles were transferred to an Eppendorf tube and fixed with 4% paraformaldehyde/PBS

for 15 minutes. Samples were washed in 0.1% PBT. Cuticles were incubated with Nile Red (2 µg/mL in

PBS) for 1 hour, protected from light. Following incubation, samples were washed in PBS, the fat body

was dissected and mounted as described above.

Hemolymph sample preparation

Fifteen larvae were collected and thoroughly washed with Milli-Q water to remove any fly food particles.

Care was taken to ensure no food particles remained on the larvae surfaces, and they were dried using

Kim Tech paper wipes. Each larva was punctured in the center using forceps and transferred to a 0.5 mL

microcentrifuge tube with three 1 mm holes at the bottom. The 0.5 mL microcentrifuge tube was then

placed into a pre-cooled 1.5 mL microcentrifuge tube. One larva at a time was processed in this assembly

and centrifuged in a microfuge for 10 seconds. After each centrifugation, the larval carcass was removed

from the 0.5 mL microcentrifuge tube to prevent blockage of the holes. Hemolymph isolated from the 15

larvae was collected in the bottom 1.5 mL tube. From the total collected hemolymph, 8 µL was transferred

to a fresh 1.5 mL tube, and 10 µL of ultrapure Milli-Q water, 30 µL of methanol, an internal standard, and

50 µL of MTBE were added. The solution was mixed thoroughly and centrifuged at 1000g for 10 minutes at

4°C. After centrifugation, the organic and polar phases were collected separately in different tubes for

metabolite measurement.

Hemolymph metabolomic analyses

Non-targeted analysis of polar metabolites by LC-MS was carried out as described previously (Edwards-Hicks et al, 2020) using an Agilent 1290 Infinity II UHPLC in line with a Bruker Impact II QTOF-MS operating in negative and positive ion mode. Scan range was from 20 to 1050 Da and mass calibration was performed at the beginning of each run. LC separation was on a Waters Atlantis Premier BEH ZHILIC column (100 x 2.1 mm, 1.7 µm particles), buffer A was 20 mM ammonium carbonate and 5 µM medronic acid in milliQ H2O and buffer B was 90:10 acetonitrile:buffer A and the solvent gradient was from 95% to 55% buffer B over 14 minutes. Flow rate was 180 µL/min, column temperature was 35 °C, autosampler temperature was 5 °C and injection volume was 3 µL. Data processing including feature detection, feature deconvolution, and annotation of features was performed using MetaboScape (version 2023b). Quantile normalization was performed to minimize sample size effects and further statistical processing were performed with R. Metabolites with missing values were eliminated from the dataset. Both, the Shapiro-Wilk test and the Q-Q plot, showed that the data was not normally distributed. Therefore, a two-sided, unpaired Wilcoxon rank-sum test was performed on the quantile normalized data. Only metabolites with at least <0.75 and >1.5 fold change and a p-value < 0.19 and were selected for further analysis and were displayed in a heatmap with row-wise normalization (Fig 3L).

Image quantification and statistical analysis - General comments

For all quantifications, control and experimental samples were processed together and imaged in parallel, using the same confocal settings. Images were processed, analyzed and quantified using tools in Fiji (ImageJ 2.14) (Schindelin et al, 2012). Extreme care was taken to apply consistent methods to control and experimental samples (i.e., number of projected sections, thresholding methods, processing) for image analysis and quantifications. See **File S1** for macros used in FIJI during image segmentation and quantifications. Further details on segmentation and quantifications are provided below. Figure panels were assembled using Affinity Designer 2.4.0. Statistical analyses were performed in Graphpad Prism. Illustrations were done in Biorender.

To quantify fluorescence intensities inside the proliferative domain (PD^{egr}), a mask of the high JNK-signaling domain region was generated, followed by creating a 20µm band on the outside to mark the proliferative domain. Fluorescence intensities in proliferative domains was always compared to fluorescence intensities within the pouch domain of control discs, unless otherwise noted.

Quantification of cycling cells using EdU incorporation

(a) EdU quantification in notum: An xy-section with the maximum number of notum epithelial cells was selected, excluding any myoblast cells. A mask was created from the DAPI staining to represent nuclei and saved as ROI. A binary operation (AND) was used to compute a notum DAPI mask with the aid of a manually selected notum region (FileS1, Macro-1L(a)). This mask was then used to measure EdU intensity.

(b) EdU quantification within 10µm bands: In *egr*-expressing discs, TRE-RFP expression was used to generate a mask of the high JNK-signaling domain (FileS1, Macro-1D(a)). The boundary of this mask defined the regions inside and outside the high JNK region. Five 10µm bands were created outside the high JNK-signaling domain, and three 10µm bands inside of the JNK-signaling domain. Also applying a DAPI mask to each band (FileS1, Macro-1D(b)), the mean EdU intensity in each of the 10µm bands was measured. Consistent with previous studies (Cosolo et al, 2019; Jaiswal et al, 2023), the highest EdU intensities were observed in 3 bands outside the JNK-signaling domain, which was denoted as proliferative domain.

(c) EdU quantification in eye disc: For the eye disc, a DAPI mask (FileS1, Macro-2C(a)) was generated using the DAPI staining, which was then applied to measure EdU intensity in the region above the morphogenetic furrow. EdU intensity was also measured inside an EdU-based mask, which was generated on a single X-Y slice.

Quantification of Upd3-LacZ levels

A maximum projection of TRE-RFP expression was used to generate a mask of the high JNK-signaling domain in *egr*-expressing disc (FileS1, Macro-4J(a)). Upd3-LacZ intensity was measured within the TRE-RFP mask of the high JNK-signaling domain in *egr*-expressing disc, while for the control disc, Upd3-LacZ intensity was measured in the pouch.

Quantification of ImpL2-GFP levels

A pouch mask was generated using the maximum projection of an anti-Nubbin staining (FileS1, Macro-4G(a)) for both control and *egr*-expressing discs. ImpL2-GFP intensities were measured within the Nubbin mask on the sum projection of the ImpL2-GFP signals within the stack (seven slices in each disc).

Quantification of OPP levels in different regions of wing imaginal disc

In the control disc, mean OPP intensity was measured separately in the pouch, hinge, and notum regions. In *egr*-expressing discs, mean OPP intensity was measured in the high JNK signaling region (TRE-RFP mask), the proliferative region (20µm band around the TRE-RFP mask), and the notum region. In control discs, the pouch, hinge, and notum regions were selected using wing fold landmarks. In *egr*-expressing discs, a max projection of TRE-RFP intensities was used to create a mask of the high JNK signaling domain (FileS1, Macro-1G(a)), which was applied to measure OPP intensity in the high JNK region. A 20µm band outside the high JNK region was generated to mark the proliferative region, and OPP intensity in this domain was measured. For the notum, a manual selection along the first notum fold and outlining the edges of the notum was carried out.

Quantification of OPP levels in the eye disc

585   For the eye-antennal disc, the DAPI staining was used to generate a mask of the tissue outline (FileS1,
586   Macro-2F(a)) and combined with a manual selection of the eye disc to generate a region of interest (ROI)
587   within which we measured OPP intensity.
588

Quantification of nuclear dFOXO levels
In control discs, a slice with the maximum number of pouch cells was selected and a DAPI mask (FileS1,
Macro-5C(a)) was generated and combined with manual pouch selection to create a pouch nuclear
mask. This mask was used to measure mean dFOXO intensity in the pouch nucleus. In *egr*-expressing
discs, a single slice from the Z-stack with the maximum number of cells in the proliferative domain was
selected. A max projection of TRE-RFP intensities was used to create a mask of the high JNK signaling
domain, and a 20μm band was created to locate the proliferative region. A DAPI mask (FileS1, Macro-
5C(b)) was generated from the selected slice and combined with the 20μm band to create a new mask for
the proliferative cell nuclei, which was used to measure dFOXO intensity in the nucleus. Please note that
discs and fat body were treated with Leptomycin B (413nM) for 30 min before dissection to block nuclear
export.
600

Quantification of PDK-1 GFP, p-S6 and OPP levels in the en-GAL4 domain
The slice with the maximum number of pouch cells was selected, and masks of the anterior and posterior
pouch compartments were generated by segmenting the enGAL4, UAS-RFP expression domain. Levels of
p-S6 and Pdk1-GFP intensities were measured within these masks and the P/A ratio (en-GAL4 region/non-
en-GAL4 region) was calculated. Mean OPP intensity in the control and experimental pouch en-GAL4
domain was measured and reported.
607

608   **Quantification of 10xStat92E-GFP levels in the fat body**
609   A single slice from the Z-stack with the maximum number of nuclei was selected. A 200x200 pixel square
610   region was defined and used as a standard area for measurement of GFP intensities.
611

612   **Quantification of lipid droplet size and shape**
613   A single slice from the Z-stack through the most anterior fat body containing the maximum number of
614   nuclei was selected. A published STAR protocol was applied to measure lipid droplet area in both control
615   and experimental samples (https://doi.org/10.1016/j.xpro.2022.101230). Additionally, the roundness of
616   lipid droplets was measured using the same published workflow.
617

Author Contributions

Conceptualization AVM, AKC; Data curation AVM, LH, AKC; Formal analysis AVM, LN, LH, JB, AKC; Funding acquisition AKC; Investigation AVM, LN, AKC; Methodology AVM, LN, IG, JB, AKC; Project administration IG, AKC; Supervision KK, AKC; Validation AVM, LN, KK, AKC; Visualization AVM, LN, LH; Writing – original draft AVM, AKC; Writing – review & editing AVM, LN, KK, AKC

Acknowledgements

We thank the staff of the Life Imaging Center (LIC) in the Hilde Mangold House (HMH) of the Albert-Ludwigs-University of Freiburg for help with their confocal microscopy resources, and the excellent support in image recording. We specifically thank the DFG for supporting our imaging work through project number 414136422. We thank Susumi Hirobayashi, Hugo Stocker, Aurelio Teleman and Hong Xu, Iswar Hariharan, Dirk Bohmann for sharing reagents and the Bloomington *Drosophila* Stock Center (BDSC), the Vienna *Drosophila* Stock Collection (VDRC), the University of Zurich ORFeome Project (FlyORF), the Developmental Studies Hybridoma Bank (DSHB) and the Monoclonal Antibody Core Facility at the Helmholtz Zentrum Munich for providing fly stocks and antibodies. We thank the IMPRS-IE graduate school for supporting our doctoral researchers. We thank Olaf Gross and Valerie Hilgers for critical feed-back on the manuscript.

Funding

Funding for this work was provided by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germanyś Excellence Strategy (CIBSS – EXC-2189), the Heisenberg Program (CL490/3-1, CL490/4-1), and by the Boehringer Ingelheim Foundation (BIF Plus3 & Rise Up) to AKC. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Declaration of Interest

The authors declare no competing interests.

650 **Main Figures**

Figure 1. Spatial organization of cell proliferation and protein synthesis induced by inflammatory damage in wing imaginal disc.

A,B. Control wing disc (A) and a wing discs (B) after 24 hours of *egr*-expression within the wing pouch domain (see Fig 1C) under the control of the rn-GAL4 driver (rn^{ts} and rn^{ts}>egr, respectively). TRE-RFP visualizes JNK-pathway activity (magenta or grey) and EdU incorporation visualizes DNA replication (cyan or grey). Discs were stained with DAPI to visualize nuclei.

C. Schematic representation of a third instar wing imaginal disc, highlighting different regions (pouch, notum, and hinge) on the left. On the right, schematic representation of a wing imaginal disc after 24 hours of *egr*-expression in the pouch. The magenta area identifies the JNK-signaling domain representing inflammation and tissue damage.

D. Quantification of average EdU intensity in the DAPI area contained within 10μm bands segmented inward and outward from the edge of the high JNK-signaling domain in *egr*-expressing discs. The the high JNK-signaling domain is shaded in magenta, the proliferative domain marking the regenerating domain is shaded in cyan. Mean and 95% CI (confidence interval) are shown. Statistical significance was tested using a one-way ANOVA followed by Tukey's post-hoc test for multiple comparisons (n=7).

E,F. Protein synthesis visualized by O-proparyl-puromycin (OPP) incorporation into newly synthesized proteins in control wing disc (E) and *egr*-expressing discs (F). Discs were stained with DAPI to visualize nuclei.

G. Quantification of mean OPP intensity in three different regions of control and *egr*-expressing discs. In control disc, OPP intensity was measured in the pouch, hinge, and notum, while in *egr*-expressing discs, OPP intensity is measured in the high JNK signaling domain (approximating the central pouch), the proliferating region (approximating the peripheral pouch and hinge), and the notum. Mean and 95% CI are shown. Statistical significance was tested using a one-way ANOVA followed by Tukey's post-hoc test for multiple comparisons (control: n=8, *egr*-expressing disc: n=8).

H, I. Protein synthesis visualized by OPP incorporation in the notum of control (H) and *egr*-expressing discs (I).

J,K. EdU incorporation to visualize DNA replication in the notum of control (J) and *egr*-expressing discs (K).

L. EdU area per DAPI area quantified in the notum of control and *egr*-expressing discs, serving as a proxy for the number of cells undergoing DNA replication. Mean and 95% CI are shown. Statistical significance was tested using an Unpaired t-test (control: n=9, *egr*-expressing disc: n=8), with a p-value = 0.0001.

M. Mean EdU intensity per EdU-positive area quantified in the notum of control and *egr*-expressing discs, serving as a proxy for DNA replication speed. Mean and 95% CI are shown. Statistical significance was tested using an Unpaired t-test (control: n=9, *egr*-expressing disc: n=8), with a p-value = 0.0011.

N,O. Schematic representation of protein synthesis rates (blue shades) and mitotic activity (black dots) in control disc (left) and *egr*-expressing discs (right) (N). Based on activation of distinct signaling

689   pathways, or metabolic and mitotic activities in *egr*-expressing discs, three different cell populations

690   (senescent, proliferating and quiescent) can be distinguished (O).

692   Scale bars: 100 µm. Fluorescence intensities are reported as arbitrary units.

2

Figure 2. Inflammatory tissue damage induces a systemic reduction in growth and cell proliferation.

A,B. EdU incorporation to visualizes DNA replication in eye imaginal discs, dissected from larvae with

control (A) or *egr*-expressing (B) wing imaginal discs. The yellow arrowhead points to the expected

location of the mitotic wave just anterior to the morphogenetic furrow.

C. Mean EdU intensity per EdU-positive area serving as a proxy for DNA replication speed, quantified in

eye imaginal discs dissected from larvae with control or *egr*-expressing wing imaginal discs. Mean and

95% CI are shown. Statistical significance was tested using the Mann-Whitney test, p-value = 0.0286

(control: n=4, *egr*-expressing disc: n=4).

D,E. Protein synthesis visualized by OPP incorporation in the eye imaginal disc, dissected from larvae

with control (D) or *egr*-expressing (E) wing imaginal discs. TRE-RFP visualizes JNK-pathway activity.

F. Mean OPP intensity in eye imaginal discs, dissected from larvae with control or *egr*-expressing wing

imaginal discs. Mean and 95% CI are shown. Statistical significance was tested using an Unpaired t-test,

p-value = 0.0002 (control: n=5, *egr*-expressing disc: n=5).

G,H. Expression of Dilp8-GFP (cyan or grey) in control (G) and *egr*-expressing discs (H). TRE-RFP visualizes

JNK-pathway activity (magenta or grey). Discs were stained with DAPI to visualize nuclei.

I,J. Expression of dFOXO-GFP (cyan or grey) in fat body, dissected from larvae with control (I) or *egr*-

expressing (J) wing imaginal discs. Fat bodies were stained with DAPI to visualize nuclei (magenta or grey).

K. Mean nuclear dFOXO-GFP intensity in fat body, dissected from larvae with control or *egr*-expressing

wing imaginal discs. Statistical significance was tested using an unpaired t-test, p-value = 0.0003

(control: n=10, *egr*-expressing disc: n=9).

Scale bars: 100 μm. Fluorescence intensities are reported as arbitrary units.

3

Figure 3. mTORC signaling, nutrient transporters and metabolite changes in the hemolymph support regenerative proliferation

A,B. Immunostaining for phosphorylated-S6 (pS6, cyan or grey) in control (A) and *egr*-expressing discs (B). TRE-RFP visualizes JNK-pathway activity (magenta or grey). Discs were stained with DAPI to visualize nuclei.

C. Mean p-S6 intensity quantified in the pouch of control discs and the proliferative domain of *egr*-expressing discs (PD^egr). Mean and 95% CI are shown. Statistical significance was tested using the Mann-Whitney test, p-value = 0.0001 (control: n=9, *egr*-expressing discs: n=9).

D-G. Protein synthesis visualized by OPP incorporation in control (D,E) and *egr*-expressing discs (F,G) from larvae fed on food without (D,F) or with rapamycin (200 µM) (E,G) for 24 h during the 24 h period of *egr*-expression in wing discs.

H. Quantification of mean OPP intensity in the pouch of control discs and the proliferative domain of *egr*-expressing discs (PD^egr) from larvae fed on food without or with rapamycin (200 µM) for 24 hours during the 24 h period of *egr*-expression in wing discs. Mean and 95% CI are shown. Statistical significance was tested using a one-way ANOVA followed by Tukey's post-hoc test for multiple comparisons (control: non-fed:n=9, fed: n=10 ; *egr*-expressing disc: non-fed: n=10, fed: n=10).

I,J. Expression of CG5535-GFP in control (I) and *egr*-expressing discs (J). Discs were stained with DAPI to visualize nuclei.

K. Mean CG5535-GFP intensity quantified in the pouch of control discs and the proliferative domain of *egr*-expressing discs (PD^egr). Mean and 95% CI are shown. Statistical significance was tested using the Unpaired t-test, p-value = 0.0092 (control: n=4, *egr*-expressing discs: n=5).

L. Heat map representing relative concentration changes of metabolites identified in the larval hemolymph isolated from control larvae and *egr*-expressing larvae. The heat map was generated using quantile normalized data, and statistical analysis was performed using a two-sided, unpaired Wilcoxon rank-sum test (see Materials and Methods) and metabolites with at least <0.75 and >1.5 fold change were selected for further analysis. Metabolites in the heat map are ordered based on the log2 fold change between control and *egr*-expressing larval samples and Z-scores were visualized. Sample size was n=3 for both control and *egr*-expressing larvae.

Scale bars: 100 µm. Fluorescence intensities are reported as arbitrary units.

4

Figure 4. Impl2 signaling mobilizes nutrients in the fat body

A,B. Nile Red staining (cyan or grey) of fat body, dissected from larvae with control (A) or *egr*-expressing (B) wing imaginal discs. Fat bodies were stained with DAPI (magenta or grey) to visualize nuclei.

C, D. Quantification of area and roundness of lipid droplets in the fat body dissected from larvae with control (C) or *egr*-expressing (D) wing imaginal discs. Mean and 95% CI are shown. Statistical significance of was tested using an Unpaired t-test. Mean area, p-value = 0.0004. Mean roundness, p-value < 0.0001. (control: n=15, *egr*-expression in discs: n=18).

E,F. Expression of Impl2-GFP in control (A) and *egr*-expressing discs (B). Discs were stained with DAPI to visualize nuclei.

G,J. Impl2-GFP (G) and Upd3-lacZ (J) intensity quantified in the pouch of control wing discs and in the JNK-signaling domain of *egr*-expressing discs. Mean and 95% CI are shown and statistical significance was tested using Welch's t-test (Impl2-GFP: control: n=9, *egr*-expressing discs: n=11, p-value = 0.0001; Upd3-LacZ: control: n=8, *egr*-expressing disc: n=9, p-value < 0.0001).

H,I. Expression of Upd3-lacZ in control (C) and *egr*-expressing discs (D). Discs were stained with DAPI to visualize nuclei.

K,L. Activity of the JAK/STAT reporter *10xSTAT92E>GFP* (cyan or grey) in the fat body dissected from larvae with control (K) or *egr*-expressing (L) wing imaginal discs. TRE-RFP visualizes JNK-pathway activity (grey) and DAPI visualizes nuclei (magenta or grey).

M. Quantification of *10xSTAT92E>GFP* reporter intensity in the fat body dissected from larvae with control or *egr*-expressing wing imaginal discs. Mean and 95% CI are shown and statistical significance was tested using the Mann-Whitney test, p-value < 0.0001 (control: n=12, *egr*-expression in disc: n=14).

N,O. Nile Red staining of fat body, dissected from larvae with control (N) or Impl2-expressing (O) wing imaginal discs. Fat bodies were stained with DAPI to visualize nuclei.

P,Q. Quantification of area of lipid droplets and roundness of lipid droplets in the fat body dissected from larvae with either control discs or wing imaginal discs expressing ImpL2 for 24 h under the control of rn-GAL4. For lipid droplet area, statistical significance was tested using the Mann-Whitney test, p-value < 0.0001 (control: n=23, experiment: n=18). For lipid droplet roundness, statistical significance was tested using the Unpaired t-test, p-value=0.0088 (control: n=24, *egr*-expression in disc: n=18)

Scale bars: 100 µm. Fluorescence intensities are reported as arbitrary units.

5

Figure 5. STAT signaling regulates protein synthesis via regulation of Pdk1

A,B. Expression of dFOXO-GFP (cyan or grey) in control (A) and *egr*-expressing discs (B). TRE-RFP

visualizes JNK-pathway activity (magenta or grey) and DAPI visualizes nuclei (grey).

C. Mean nuclear dFOXO-GFP intensity quantified in the pouch of control discs or the proliferative domain

of *egr*-expressing discs (PD^{egr}). Mean and 95% CI are shown and statistical significance was tested using

the Mann-Whitney test, p-value = 0.0006 (control: n=7, *egr*-expressing disc: n=7).

D,E. Expression of Pdk1-GFP (cyan or grey) in control (D) and *egr*-expressing discs (E). TRE-RFP visualizes

JNK-pathway activity (magenta or grey) and DAPI visualizes nuclei (grey).

F. Quantification of Pdk1-GFP intensity in the pouch of control discs or the proliferative domain of *egr*-

expressing discs (PD^{egr}). Statistical significance was tested using a Welch's t-test, p-value = 0.0025

(control: n=10, *egr*-expressing disc: n=10).

G-N. Control wing imaginal discs from sibling larvae (G,J,M) and wing imaginal discs expressing UAS-

STAT92E in the posterior compartment under the control en-GAL4 which also drives expression of UAS-

RFP (H,K,N). Discs were assayed for expression of Pdk1-GFP (G,H), levels of p-S6 (J,K), and protein

synthesis using OPP incorporation assays (M,N). DAPI staining visualizes nuclei.

I,L,O. Quantifications of levels of Pdk1-GFP (I), p-S6 (L) and protein synthesis using OPP incorporation

assays (O) in control wing imaginal discs and wing imaginal discs expressing UAS-STAT92E in the

posterior compartment under the control en-GAL4. In the graph displaying Pdk1-GFP intensity (I),

statistical significance was tested using the Unpaired t-test, p-value < 0.0001 (control: n=5, experiment:

n=8). In the graph displaying p-S6 intensity (L), statistical significance was tested using Welch's t-test, p-

value = 0.0033 (control: n=5, experiment: n=8). In the graph displaying OPP intensity (O), statistical

significance was tested using the Unpaired t-test, p-value = 0.0004 (control: n=5, experiment: n=5). Mean

and 95% CI are shown.

Scale bars: 100 μm. Fluorescence intensities are reported as arbitrary units.

6

Figure 6. Pdk1 is a central regulator of regenerative metabolism

A,B. Protein synthesis visualized by OPP incorporation in control discs (A) and discs expressing UAS-PDK1 for 24 h under the control of rn-GAL4 (B). DAPI staining visualizes nuclei. The dotted white line indicates the boundary of the wing pouch based on landmark tissue folds, approximating the rn-GAL4 expression domain.

C,D. EdU incorporation in control (C) and discs expressing UAS-PDK1 for 24 h under the control of rn-GAL4 (D). DAPI staining visualizes nuclei. The dotted white line indicates the boundary of the wing pouch based on landmark tissue folds, approximating the rn-GAL4 expression domain.

E. Quantification of mean OPP intensity in the pouch of control or *Pdk1*-expressing wing discs. Mean and 95% CI are shown and statistical significance was tested using an Unpaired t-test, p-value < 0.0001 (control: n=6, *Pdk1*-expressing disc: n=6).

F. Quantification of Mean EdU intensity per EdU-positive area serving as a proxy for DNA replication speed, in the pouch of control or *Pdk1*-expressing wing discs. Mean and 95% CI are shown. Statistical significance was tested using the Welch's t-test, p-value = 0.0078 (control: n=8, *Pdk1*-expressing disc: n=8).

G-J. Protein synthesis visualized by OPP incorporation in control (G, H) and *egr*-expressing discs (I, J) that were either wild type (G, I) or heterozygous mutant for the *Pdk1^5* null allele (H, J). DAPI staining visualizes nuclei.

K. Quantification of mean OPP intensity in the pouch of control or the proliferative domain of *egr*-expressing discs (PD^{egr}) that were either wild type or heterozygous mutant for the *Pdk1^5* null allele. Statistical significance was tested using a one-way ANOVA followed by Tukey's post-hoc test for multiple comparisons. (control:n=6, control, *PDK1^5/+*: n=6 ; *egr*:n=8, *egr, PDK1^5/+*: n=7).

L, M. Pdk1-GFP expression in eye imaginal discs, dissected from larvae with control or *egr*-expressing wing imaginal discs. DAPI staining visualizes nuclei.

N. Quantifications of Pdk1-GFP expression in eye imaginal discs, dissected from larvae with control or *egr*-expressing wing imaginal discs. Mean and 95% CI are shown. Statistical significance was tested using an Unpaired t-test, p-value = 0.0031 (control: n=3, *egr*-expressing disc: n=3).

O, P. A model illustrating the local (O) and systemic (P) changes associated with anabolic growth and protein synthesis in normal tissues (O, left) and in the regenerating domain within an insulin-restricted environment. (O, right). Inflammatory cytokine signals (P) induce systemic changes that specifically support metabolic adaptations for rapid proliferation during regeneration.

Scale bars: 100 μm. Fluorescence intensities are reported as arbitrary units.

841 **Supplementary Figures and Tables**

S1

Figure S1. Cell cycle adaptations in response to inflammatory tissue damage

A. Schematic illustrating the spatial organization of cell signaling and cell behavior in damaged tissue following injury. The tissue repair region is divided into two spatially separated domains, indicated by different colors. Senescent cells are located in the high JNK/AP1 signaling domain (magenta), while regenerative proliferation occurs in the JAK/STAT signaling domain (cyan). See also (Cosolo et al, 2019; Crucianelli et al, 2022; Floc'hlay et al, 2023; Jaiswal et al, 2023) for details.

B, C. Control wing disc (B) and a wing discs (C) after 24 hours of *egr*-expression within the wing pouch domain under the control of the rn-GAL4 driver (rn^{ts} and rn^{ts}>egr, respectively). A co-expressed UAS-GFP construct visualizes the rn-GAL4 expression domain. Staining for cleaved Dcp-1 visualizes apoptosis. Discs were stained with DAPI to visualize nuclei. Please note the GFP-expression domain in (C) which does not stain for cDcp-1 and represents surviving cells with high levels of JNK activity (see Fig 1B). Levels of Dcp-1 were previously quantified (Cosolo et al, 2019; Jaiswal et al, 2023; La Fortezza et al, 2016).

D-F. Control (D) and *egr*-expressing wing disc (E), co-expressing the FUCCI cell cycle reporter system (F) consisting of mRFP-NLS.CycB.1-266 (red in overlay) and GFP-E2f1.1-230 (green in overlay). Discs were stained with DAPI to visualize nuclei. Cells in the Gap phase G1, express GFP; cells in early S-phase lack expression of both FUCCI constructs; cells in late S-phase express RFP; and cells in G2 express both reporters. Cell cycle changes were already previously quantified (Cosolo et al, 2019; Crucianelli et al, 2022; Jaiswal et al, 2023).

G,H. Analysis of senescence-associated β-galactosidase (SA-β-gal) activity (cyan or grey) in control (G) and *egr*-expressing wing disc (H). TRE-RFP visualizes JNK-pathway activity. Discs were stained with DAPI to visualize nuclei (magenta or grey). SA-β-gal activity was already previously reported and quantified (Floc'hlay et al, 2023; Jaiswal et al, 2023).

I,J. Nota of control (I) and wing disc expressing egr in the pouch (J), co-expressing the FUCCI cell cycle reporter system consisting of mRFP-NLS.CycB.1-266 (red in overlay) and GFP-E2f1.1-230 (green in overlay) (as described in Fig-S1F). Discs were stained with DAPI to visualize nuclei.

Scale bars: 100 µm.

S2

	Figure S2. Metabolic adaptation in peripheral tissues

875	**A, B.** Protein synthesis visualized by O-proparyl-puromycin (OPP) incorporation into newly synthesized

876	proteins in leg imaginal discs, dissected from larvae containing either control wing disc (A) or *egr-*

877	expressing wing discs (B). Discs were stained with DAPI to visualize nuclei.

879	Scale bar: 100 μm.

S3

Figure S3. Elevated levels of nutrient transporters and mTORC signaling support regenerative proliferation

A. UMAP plots generated using the Scope Wing Atlas created from control and *egr*-expressing discs analysed by single-cell RNA-Seq technology (Floc'hlay et al, 2022). Left: A 'wound' cluster appears in *egr*-expressing discs. Right: Further analysis revealed the identity of an alpha and beta cluster within *egr*-expressing discs with signatures characteristic of senescent cells (beta) and with characteristic of a cell and tissue damage response signature (alpha).

B. The scRNA UMAP plot displays the transcript levels of ribosomal protein S6 kinase (S6K). Grey indicates low expression, and red indicates high expression of transcripts.

C. A scheme illustrating InR/PI3K/AKT and mTORC signaling pathways involved in growth and protein synthesis in normal tissues.

D. scRNA UMAP plot displaying significantly elevated transcript levels detected for different nutrient transporters in alpha and beta clusters (Floc'hlay et al, 2023). Grey indicates low expression, and red indicates high expression of transcripts.

E,F. Staining for the putative alanine, glycine and proline amino acid transporter Arcus (CG1139, cyan or grey) in control (E) and *egr*-expressing discs (F). TRE-RFP visualizes JNK-pathway activation (magenta or grey). Discs were stained with DAPI to visualize nuclei.

G,H. Expression of Tret1.1-GFP (cyan or grey) in control (G) and *egr*-expressing discs (H). TRE-RFP visualizes JNK-pathway activation (magenta or grey). Discs were stained with DAPI to visualize nuclei.

I. Mean Tret1.1-GFP intensity quantified in the pouch of control discs and the proliferative domain of *egr*-expressing discs. Mean and 95% CI are shown. Statistical significance was tested using the Welch's t-test, p-value < 0.0001 (control: n=5, *egr*-expressing discs: n=9).

Scale bar: 100 μm. Fluorescence intensities are reported as arbitrary units.

Figure S4. Interorgan signals are induced by *egr*-driven inflammatory damage

A, B. *Drosophila* larval body wall muscles dissected from larvae containing either control (A) or *egr*-
expressing (B) wing discs. Actin was visualized using Phalloidin (cyan or grey) and TRE-RFP visualizes JNK-
pathway activation (magenta or grey). Muscles were stained with DAPI to visualize nuclei. Please note
that muscles do not activate JNK-signaling in response to inflammatory cues.

C,D. Nota of control (C) and *egr*-expressing wing disc (D) stained for the autophagy marker Ref(2)P. Discs
were stained with DAPI to visualize nuclei.

E. UMAP plots generated using the Scope Wing Atlas created from control and *egr*-expressing discs
analysed by single-cell RNA-Seq technology (Floc'hlay et al, 2022). Significantly elevated transcript levels
detected for different secreted paracrine ligands with known interorgan signaling roles in alpha and beta
clusters are shown. Grey indicates low expression, and red indicates high expression of transcripts.

Scale bar: 100 µm.

S5

Figure S5. Insulin/PI3K/Akt signaling is low in the proliferative domain

A-D. Control (A, C) and *egr*-expressing wing discs (B, D), expressing an Insulin receptor tagged with GFP
(InR-GFP) (A, B) or stained for Akt (C, D) (cyan or grey). TRE-RFP visualizes JNK-pathway activity (magenta
or grey). Discs were stained with DAPI to visualize nuclei.

E. Anti-phospho-Akt (S505) staining (cyan or grey) to test antibody specificity in UAS-Akt-overexpressing
clones marked by co-expression of UAS-GFP (magenta or grey) in the wing imaginal disc. Discs were
stained with DAPI to visualize nuclei.

F,G. Control (F) and *egr*-expressing wing discs (G), stained for phospho-Akt (S505) (cyan or grey). TRE-RFP
visualizes JNK-pathway activity (magenta or grey). Discs were stained with DAPI to visualize nuclei.

H. Quantification of mean p-Akt intensity in the control pouch and proliferative domain of *egr*-expressing
wing discs. Mean and 95% CI are shown and statistical significance was tested using an Unpaired t-test ,
p-value =0.0156. (Control; n=4, *egr*-expressing discs; n=5)

I, J. Control and *egr*-expressing discs stained for EdU to visualize DNA replication in discs also co-
expressing a 10xStat92E-GFP activity reporter (same disc is shown in Fig-1A). Discs were also stained
with DAPI to visualize nuclei. EdU intensity in 10xStat92E-GFP reporter-expressing domain was previously
quantified (Jaiswal et al, 2023).

Scale bar: 100 µm. Fluorescence intensities are reported as arbitrary units.

S6

Figures S6. Pdk1 is sufficient to drive elevated protein synthesis in wing imaginal disc

A,B. Protein synthesis visualized by OPP incorporation in control (A) and discs expressing a constitutively

active UAS-PDK1 (B) for 24 h under the control of rn-GAL4. Discs were stained with DAPI to visualize

nuclei. The dotted white line marks the border of the pouch using tissue fold landmarks.

C. Quantification of mean OPP intensity in the pouch of control and *Pdk1*^{CA}-expressing wing discs. Mean

and 95% CI are shown and statistical significance was tested using Mann-Whitney test. p-value = 0.0062.

(Control; n=8, *Pdk1*^{CA}-expressing discs; n=10).

D. Quantification of the pouch area in control and *Pdk1*^{CA}-expressing wing discs. Mean and 95% CI are

shown and statistical significance was tested using Unpaired t-test. p-value = 0.0003. (Control; n=9,

Pdk1^{CA}-expressing discs; n=8).

E,F. Control (E) and *egr*-expressing wing disc (F), also co-expressing the Pleckstrin Homology Domain

reporter for PIP3 levels (tGPH) (cyan or grey). TRE-RFP visualizes JNK-pathway activity (magenta or grey).

Discs were stained with DAPI to visualize nuclei.

Scale bar: 100 µm. Fluorescence intensities are reported as arbitrary units.

Table S1 Drosophila strains

Genotype	Source
Dilp8-GFP	BDSC: 33079
ImpL2-GFP	BDSC: 59778
UAS-ImpL2-3xHA	Fly-ORF F001712
Upd3.1-3 LacZ	Ishwar Hariharan lab
dFOXO-GFP	BDSC: 59766
en2.4-GAL4, UAS-mRPF.NLS/CyO	BDSC: 30557
Tret1-GFP	BDSC: 66367
Pdk1-GFP	BDSC: 59836
InR-eGFP	Hong Xu lab DOI: 10.7554/eLife.49309
tGPH-GFP	Yohanns Bellaiche
UAS-Akt1	BDSC: 8192
CG5535-GFP	BDSC: 64096
CycB-FUCCI': Ubi-mRFP1.NLS.CycB$^{1-266}$	Norman Zielke PMID: 24726363
E2F1-FUCCI, CycB-FUCCI'; Ubi-GFP.E2f1$^{1-230}$, Ubi-mRFP1.NLS.CycB$^{1-266}$	Norman Zielke PMID: 24726363
UAS-GFP^{S56T}	BDSC: 1521
rn$^{GAL4-DeltaS}$, tubGAL80^{ts}	BDSC: 8142, 7012
rn^{GAL4-5}, UAS-egr, tubP-GAL80^{ts}	Iswar Hariharan
10xStat92E-dGFP	Erica Bach PMID: 17008134
10xStat92E-GFP	Erica Bach PMID: 17008134
UAS-Stat92E.ORF.3xHA	Fly-ORF F000750 PMID: 23583758
'TRE-RFP': TRE-DsRed.T4	Dirk Bohmann PMID: 22509270
upd3.1-3-LacZ	Iswar Hariharan
UAS-dPdk1	Hugo Stocker
UAS-dPdk1(A467V)	Hugo Stocker https://doi.org/10.1073/pnas.011318098
dPdk1(5)ΔEP	Hugo Stocker lab https://doi.org/10.1073/pnas.011318098

964 **Table S2 Immunohistochemistry reagents**

Reagent	Source	
Mouse polyclonal anti-β-Galactosidase (1:1000)	Promega	Cat. #: Z3782
Mouse monoclonal anti-MMP-1 (1:30)	Developmental Studies Hybridoma Bank	Cat. #: 3A6B4
Mouse monoclonal anti-MMP-1 (1:30)	Developmental Studies Hybridoma Bank	Cat. #: 3B8D12
Mouse monoclonal anti-MMP-1 (1:30)	Developmental Studies Hybridoma Bank	Cat. #: 5H7B11
Mouse monoclonal anti-Nubbin (1:100)	Developmental Studies Hybridoma Bank	Cat. #: 2D4
Rabbit polyclonal anti-CG1139 (1:200)	Susumu Hirabayashi, Imperial College London Faculty of Medicine.	https://pubmed. ncbi.nlm.nih.gov /32938923/
Rabbit polyclonal anti-p-AKT (S505) (1:500)	Cell Signaling	Cat. #: 4084s
Rabbit polyclonal anti-AKT (1:500)	Cell Signaling	Cat. #: 9272s
Rat monoclonal anti-RFP (1:1000)	Chromotek	Cat. #: 5F8
Rabbit polyclonal anti-Drosophila p-S6 (1:200)	Aurelio Telemann, DKFZ	https://doi.org/1 0.1016/j.devcel. 2017.07.019
Rabbit polyclonal anti-Ref(2)p	Abcam	Cat. #: Ab178440
Rabbit anti-cleaved Dcp-1	Cell Signaling	Cat. #: 9578s
Rabbit monoclonal anti-GFP	Invitrogen	Cat. #: G10362
Goat anti-mouse IgG Alexa Flour 488	Invitrogen	Cat. #: A-11001
Goat anti-rabbit IgG Alexa Flour 488	Invitrogen	Cat. #: A-11008
Goat anti-mouse IgG Alexa Flour 555	Invitrogen	Cat. #: A-21422
Goat anti-rat IgG Alexa Flour 555	Invitrogen	Cat. #: A-21434
Goat anti-mouse Alexa Flour 647	Invitrogen	Cat. #: A-21235
Goat anti-rabbit Alexa Flour 647	Invitrogen	Cat. #: A-21244
Click-iT Plus EdU Alexa Fluor 647 Imaging Kit	Invitrogen	Cat. #: C10340
Phalloidin 488	Invitrogen	Cat #: A12379
Phalloidin 647	Invitrogen	Cat #: A22287
DAPI	Sigma Aldrich	Cat #: D9564
CellEvent Senescence Green Detection Kit	Invitrogen	Cat. #: C10850

Rapamycin	Thermo Fisher Scientific	Cat. #: J62473.MC
Click-iT Plus OPP Alexa Fluor 647 Imaging Kit	Invitrogen	Cat. #: C10458
Leptomycin	Sigma-Aldrich	Cat. #: L2913
Nile Red	Invitrogen	Cat. #: N1142

References

References

Agrawal, N., Delanoue, R., Mauri, A., Basco, D., Pasco, M., Thorens, B. & Leopold, P. (2016) The Drosophila TNF
Eiger Is an Adipokine that Acts on Insulin-Producing Cells to Mediate Nutrient Response. *Cell Metab*, 23(4), 675-
84.

Albina, J. E., Abate, J. A. & Mastrofrancesco, B. (1993) Role of ornithine as a proline precursor in healing wounds.
J Surg Res, 55(1), 97-102.

Altman, B. J., Stine, Z. E. & Dang, C. V. (2016) From Krebs to clinic: glutamine metabolism to cancer therapy.
Nat Rev Cancer, 16(10), 619-34.

Argiles, J. M., Busquets, S., Stemmler, B. & Lopez-Soriano, F. J. (2014) Cancer cachexia: understanding the
molecular basis. *Nat Rev Cancer*, 14(11), 754-62.

Aviner, R. (2020) The science of puromycin: From studies of ribosome function to applications in biotechnology.
Comput Struct Biotechnol J, 18, 1074-1083.

Bilder, D., Ong, K., Hsi, T. C., Adiga, K. & Kim, J. (2021) Tumour-host interactions through the lens of Drosophila.
Nat Rev Cancer, 21(11), 687-700.

Boulan, L., Andersen, D., Colombani, J., Boone, E. & Leopold, P. (2019) Inter-Organ Growth Coordination Is
Mediated by the Xrp1-Dilp8 Axis in Drosophila. *Dev Cell*, 49(5), 811-818 e4.

Britton, J. S., Lockwood, W. K., Li, L., Cohen, S. M. & Edgar, B. A. (2002) Drosophila's insulin/PI3-kinase pathway
coordinates cellular metabolism with nutritional conditions. *Dev Cell*, 2(2), 239-49.

Cao, X., Rojas, M. & Pastor-Pareja, J. C. (2022) Intrinsic and damage-induced JAK/STAT signaling regulate
developmental timing by the Drosophila prothoracic gland. *Dis Model Mech*, 15(1).

Chatterjee, N. & Bohmann, D. (2012) A versatile PhiC31 based reporter system for measuring AP-1 and Nrf2
signaling in Drosophila and in tissue culture. *PLoS One*, 7(4), e34063.

Colombani, J., Andersen, D. S. & Leopold, P. (2012) Secreted peptide Dilp8 coordinates Drosophila tissue growth
with developmental timing. *Science*, 336(6081), 582-5.

Cong, B., Nakamura, M., Sando, Y., Kondo, T., Ohsawa, S. & Igaki, T. (2021) JNK and Yorkie drive tumor
malignancy by inducing L-amino acid transporter 1 in Drosophila. *PLoS Genet*, 17(11), e1009893.

Cosolo, A., Jaiswal, J., Csordas, G., Grass, I., Uhlirova, M. & Classen, A. K. (2019) JNK-dependent cell cycle stalling
in G2 promotes survival and senescence-like phenotypes in tissue stress. *Elife*, 8.

Crucianelli, C., Jaiswal, J., Vijayakumar Maya, A., Nogay, L., Cosolo, A., Grass, I. & Classen, A. K. (2022) Distinct
signaling signatures drive compensatory proliferation via S-phase acceleration. *PLoS Genet*, 18(12), e1010516.

Destefanis, F., Manara, V. & Bellosta, P. (2020) Myc as a Regulator of Ribosome Biogenesis and Cell
Competition: A Link to Cancer. *Int J Mol Sci*, 21(11).

Ding, G., Xiang, X., Hu, Y., Xiao, G., Chen, Y., Binari, R., Comjean, A., Li, J., Rushworth, E., Fu, Z., Mohr, S. E., Perrimon, N. & Song, W. (2021) Coordination of tumor growth and host wasting by tumor-derived Upd3. *Cell Rep*, 36(7), 109553.

Dvorak, H. F. (1986) Tumors: wounds that do not heal. Similarities between tumor stroma generation and wound healing. *N Engl J Med*, 315(26), 1650-9.

Edwards-Hicks, J., Mitterer, M., Pearce, E. L. & Buescher, J. M. (2020) Metabolic Dynamics of In Vitro CD8+ T Cell Activation. *Metabolites*, 11(1).

Eming, S. A., Murray, P. J. & Pearce, E. J. (2021) Metabolic orchestration of the wound healing response. *Cell Metab*, 33(9), 1726-1743.

Eming, S. A., Wynn, T. A. & Martin, P. (2017) Inflammation and metabolism in tissue repair and regeneration. *Science*, 356(6342), 1026-1030.

Enomoto, M. & Igaki, T. (2022) Cell-cell interactions that drive tumorigenesis in Drosophila. *Fly (Austin)*, 16(1), 367-381.

Figueroa-Clarevega, A. & Bilder, D. (2015) Malignant Drosophila tumors interrupt insulin signaling to induce cachexia-like wasting. *Dev Cell*, 33(1), 47-55.

Floc'hlay, S., Balaji, R., Christiaens, V., González-Blas, C. B., De Winter, S., Hulselmans, G., De Waegeneer, M., Quan, X., Koldere, D., Atkins, M., Halder, G., Classen, A.-K. & Aerts, S. (2022) Shared enhancer gene regulatory networks between wound and oncogenic programs. *Biorxiv*.

Floc'hlay, S., Balaji, R., Stankovic, D., Christiaens, V. M., Bravo Gonzalez-Blas, C., De Winter, S., Hulselmans, G. J., De Waegeneer, M., Quan, X., Koldere, D., Atkins, M., Halder, G., Uhlirova, M., Classen, A. K. & Aerts, S. (2023) Shared enhancer gene regulatory networks between wound and oncogenic programs. *Elife*, 12.

Fox, D. T., Cohen, E. & Smith-Bolton, R. (2020) Model systems for regeneration: Drosophila. *Development*, 147(7).

Garelli, A., Gontijo, A. M., Miguela, V., Caparros, E. & Dominguez, M. (2012) Imaginal discs secrete insulin-like peptide 8 to mediate plasticity of growth and maturation. *Science*, 336(6081), 579-82.

Gontijo, A. M. & Garelli, A. (2018) The biology and evolution of the Dilp8-Lgr3 pathway: A relaxin-like pathway coupling tissue growth and developmental timing control. *Mechanisms of Development.* Elsevier Ireland Ltd.

Hackney, J. F. & Cherbas, P. (2014) Injury response checkpoint and developmental timing in insects. *Fly (Austin)*, 8(4), 226-31.

Halme, A., Cheng, M. & Hariharan, I. K. (2010) Retinoids regulate a developmental checkpoint for tissue regeneration in Drosophila. *Curr Biol*, 20(5), 458-63.

Herrera, S. C. & Bach, E. A. (2019) JAK/STAT signaling in stem cells and regeneration: from Drosophila to vertebrates. *Development*, 146(2).

Herrera, S. C., Martin, R. & Morata, G. (2013) Tissue homeostasis in the wing disc of Drosophila melanogaster:
immediate response to massive damage during development. *PLoS Genet*, 9(4), e1003446.

Hodgson, J. A., Parvy, J. P., Yu, Y., Vidal, M. & Cordero, J. B. (2021) Drosophila Larval Models of Invasive
Tumorigenesis for In Vivo Studies on Tumour/Peripheral Host Tissue Interactions during Cancer Cachexia. *Int J*
Mol Sci, 22(15).

Holz, M. K., Ballif, B. A., Gygi, S. P. & Blenis, J. (2005) mTOR and S6K1 mediate assembly of the translation
preinitiation complex through dynamic protein interchange and ordered phosphorylation events. *Cell*, 123(4),
569-80.

Hopkins, B. D., Goncalves, M. D. & Cantley, L. C. (2020) Insulin-PI3K signalling: an evolutionarily insulated
metabolic driver of cancer. *Nat Rev Endocrinol*, 16(5), 276-283.

Hoxhaj, G. & Manning, B. D. (2020) The PI3K-AKT network at the interface of oncogenic signalling and cancer
metabolism. *Nat Rev Cancer*, 20(2), 74-88.

Huang, W., Hickson, L. J., Eirin, A., Kirkland, J. L. & Lerman, L. O. (2022) Cellular senescence: the good, the bad
and the unknown. *Nat Rev Nephrol*, 18(10), 611-627.

Ingaramo, M. C., Sanchez, J. A., Perrimon, N. & Dekanty, A. (2020) Fat Body p53 Regulates Systemic Insulin
Signaling and Autophagy under Nutrient Stress via Drosophila Upd2 Repression. *Cell Rep*, 33(4), 108321.

Jaiswal, J., Egert, J., Engesser, R., Peyroton, A. A., Nogay, L., Weichselberger, V., Crucianelli, C., Grass, I., Kreutz,
C., Timmer, J. & Classen, A. K. (2023) Mutual repression between JNK/AP-1 and JAK/STAT stratifies senescent
and proliferative cell behaviors during tissue regeneration. *PLoS Biol*, 21(5), e3001665.

Jewell, J. L., Russell, R. C. & Guan, K. L. (2013) Amino acid signalling upstream of mTOR. *Nat Rev Mol Cell Biol*,
14(3), 133-9.

Jiang, H., Kimura, T., Hai, H., Yamamura, R. & Sonoshita, M. (2022) Drosophila as a toolkit to tackle cancer and
its metabolism. *Front Oncol*, 12, 982751.

Kakanj, P., Moussian, B., Gronke, S., Bustos, V., Eming, S. A., Partridge, L. & Leptin, M. (2016) Insulin and TOR
signal in parallel through FOXO and S6K to promote epithelial wound healing. *Nat Commun*, 7, 12972.

Kashio, S. & Miura, M. (2020) Kynurenine Metabolism in the Fat Body Non-autonomously Regulates Imaginal
Disc Repair in Drosophila. *iScience*, 23(12), 101738.

Kashio, S., Obata, F., Zhang, L., Katsuyama, T., Chihara, T. & Miura, M. (2016) Tissue nonautonomous effects of
fat body methionine metabolism on imaginal disc repair in Drosophila. *Proc Natl Acad Sci U S A*, 113(7), 1835-
40.

Katheder, N. S. & Rusten, T. E. (2017) Microenvironment and tumors-a nurturing relationship. *Autophagy*,
13(7), 1241-1243.

Keshet, R., Szlosarek, P., Carracedo, A. & Erez, A. (2018) Rewiring urea cycle metabolism in cancer to support
anabolism. *Nat Rev Cancer*, 18(10), 634-645.

Khezri, R., Holland, P., Schoborg, T. A., Abramovich, I., Takats, S., Dillard, C., Jain, A., O'Farrell, F., Schultz, S. W.,
Hagopian, W. M., Quintana, E. M., Ng, R., Katheder, N. S., Rahman, M. M., Teles Reis, J. G., Brech, A., Jasper, H.,
Rusan, N. M., Jahren, A. H., Gottlieb, E. & Rusten, T. E. (2021) Host autophagy mediates organ wasting and
nutrient mobilization for tumor growth. *EMBO J*, 40(18), e107336.

Kwon, Y., Song, W., Droujinine, I. A., Hu, Y., Asara, J. M. & Perrimon, N. (2015) Systemic organ wasting induced
by localized expression of the secreted insulin/IGF antagonist ImpL2. *Dev Cell*, 33(1), 36-46.

La Fortezza, M., Schenk, M., Cosolo, A., Kolybaba, A., Grass, I. & Classen, A. K. (2016) JAK/STAT signalling
mediates cell survival in response to tissue stress. *Development*, 143(16), 2907-19.

La Marca, J. E. & Richardson, H. E. (2020) Two-Faced: Roles of JNK Signalling During Tumourigenesis in the
Drosophila Model. *Front Cell Dev Biol*, 8, 42.

Lama-Sherpa, T. D., Jeong, M. H. & Jewell, J. L. (2023) Regulation of mTORC1 by the Rag GTPases. *Biochem Soc*
Trans, 51(2), 655-664.

Lee, J., Ng, K. G., Dombek, K. M., Eom, D. S. & Kwon, Y. V. (2021) Tumors overcome the action of the wasting
factor ImpL2 by locally elevating Wnt/Wingless. *Proc Natl Acad Sci U S A*, 118(23).

Levina, A., Fleming, K. D., Burke, J. E. & Leonard, T. A. (2022) Activation of the essential kinase PDK1 by
phosphoinositide-driven trans-autophosphorylation. *Nat Commun*, 13(1), 1874.

Li, J., Kim, S. G. & Blenis, J. (2014) Rapamycin: One drug, many effects. *Cell Metabolism.*

Lieu, E. L., Nguyen, T., Rhyne, S. & Kim, J. (2020) Amino acids in cancer. *Exp Mol Med*, 52(1), 15-30.

Ling, Z. N., Jiang, Y. F., Ru, J. N., Lu, J. H., Ding, B. & Wu, J. (2023) Amino acid metabolism in health and disease.
Signal Transduct Target Ther, 8(1), 345.

Liu, G. Y. & Sabatini, D. M. (2020) mTOR at the nexus of nutrition, growth, ageing and disease. *Nat Rev Mol Cell*
Biol, 21(4), 183-203.

Liu, Y., Saavedra, P. & Perrimon, N. (2022) Cancer cachexia: lessons from Drosophila. *Dis Model Mech*, 15(3).

Lodge, W., Zavortink, M., Golenkina, S., Froldi, F., Dark, C., Cheung, S., Parker, B. L., Blazev, R., Bakopoulos, D.,
Christie, E. L., Wimmer, V. C., Duckworth, B. C., Richardson, H. E. & Cheng, L. Y. (2021) Tumor-derived MMPs
regulate cachexia in a Drosophila cancer model. *Dev Cell*, 56(18), 2664-2680 e6.

MacCarthy-Morrogh, L. & Martin, P. (2020) *W O U N D H E A L I N G The hallmarks of cancer are also the*
hallmarks of wound healing.

Manning, B. D. & Cantley, L. C. (2007) AKT/PKB signaling: navigating downstream. *Cell*, 129(7), 1261-74.

Marti, I. L. A. A. & Reith, W. (2021) Arginine-dependent immune responses. *Cell Mol Life Sci*, 78(13), 5303-5324.

Martinez-Reyes, I. & Chandel, N. S. (2021) Cancer metabolism: looking forward. *Nat Rev Cancer*, 21(10), 669-
680.

Meacham, C. E., DeVilbiss, A. W. & Morrison, S. J. (2022) Metabolic regulation of somatic stem cells in vivo. *Nat*
Rev Mol Cell Biol, 23(6), 428-443.

Nakamura, M., Verboon, J. M., Allen, T. E., Abreu-Blanco, M. T., Liu, R., Dominguez, A. N. M., Delrow, J. J. &
Parkhurst, S. M. (2020) Autocrine insulin pathway signaling regulates actin dynamics in cell wound repair. *PLoS*
Genet, 16(12), e1009186.

Newton, H., Wang, Y. F., Camplese, L., Mokochinski, J. B., Kramer, H. B., Brown, A. E. X., Fets, L. & Hirabayashi,
S. (2020) Systemic muscle wasting and coordinated tumour response drive tumourigenesis. *Nat Commun*,
11(1), 4653.

Pena, O. A. & Martin, P. (2024) Cellular and molecular mechanisms of skin wound healing. *Nat Rev Mol Cell*
Biol.

Radimerski, T., Montagne, J., Rintelen, F., Stocker, H., van der Kaay, J., Downes, C. P., Hafen, E. & Thomas, G.
(2002) dS6K-regulated cell growth is dPKB/dPI(3)K-independent, but requires dPDK1. *Nat Cell Biol*, 4(3), 251-5.

Rajan, A. & Perrimon, N. (2012) Drosophila cytokine unpaired 2 regulates physiological homeostasis by
remotely controlling insulin secretion. *Cell*, 151(1), 123-37.

Ring, N. A. R., Dworak, H., Bachmann, B., Schadl, B., Valdivieso, K., Rozmaric, T., Heimel, P., Fischer, I., Klinaki,
E., Gutasi, A., Schuetzenberger, K., Leinfellner, G., Ferguson, J., Drechsler, S., Mildner, M., Schosserer, M.,
Slezak, P., Meyuhas, O., Gruber, F., Grillari, J., Redl, H. & Ogrodnik, M. (2023) The p-rpS6-zone delineates
wounding responses and the healing process. *Dev Cell*, 58(11), 981-992 e6.

Rintelen, F., Stocker, H., Thomas, G. & Hafen, E. (2001) PDK1 regulates growth through Akt and S6K in
Drosophila. *Proc Natl Acad Sci U S A*, 98(26), 15020-5.

Romao, D., Muzzopappa, M., Barrio, L. & Milan, M. (2021) The Upd3 cytokine couples inflammation to
maturation defects in Drosophila. *Curr Biol*, 31(8), 1780-1787 e6.

Rybinski, B., Franco-Barraza, J. & Cukierman, E. (2014) The wound healing, chronic fibrosis, and cancer
progression triad. *Physiol Genomics*, 46(7), 223-44.

Santabarbara-Ruiz, P. & Leopold, P. (2021) An Oatp transporter-mediated steroid sink promotes tumor-induced
cachexia in Drosophila. *Dev Cell*, 56(19), 2741-2751 e7.

Santabarbara-Ruiz, P., Lopez-Santillan, M., Martinez-Rodriguez, I., Binagui-Casas, A., Perez, L., Milan, M.,
Corominas, M. & Serras, F. (2015) ROS-Induced JNK and p38 Signaling Is Required for Unpaired Cytokine
Activation during Drosophila Regeneration. *PLoS Genet*, 11(10), e1005595.

Saxton, R. A. & Sabatini, D. M. (2017) mTOR Signaling in Growth, Metabolism, and Disease. *Cell*, 168(6), 960-
976.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C.,
Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P. & Cardona, A. (2012)
Fiji: an open-source platform for biological-image analysis. *Nat Methods*, 9(7), 676-82.

Setiawan, T., Sari, I. N., Wijaya, Y. T., Julianto, N. M., Muhammad, J. A., Lee, H., Chae, J. H. & Kwon, H. Y. (2023)
Cancer cachexia: molecular mechanisms and treatment strategies. *J Hematol Oncol*, 16(1), 54.

Shi, H. P., Fishel, R. S., Efron, D. T., Williams, J. Z., Fishel, M. H. & Barbul, A. (2002) Effect of supplemental
ornithine on wound healing. *J Surg Res*, 106(2), 299-302.

Shimobayashi, M. & Hall, M. N. (2016) Multiple amino acid sensing inputs to mTORC1. *Cell Research.* Nature
Publishing Group.

Smith-Bolton, R. K., Worley, M. I., Kanda, H. & Hariharan, I. K. (2009) Regenerative growth in Drosophila
imaginal discs is regulated by Wingless and Myc. *Dev Cell*, 16(6), 797-809.

Song, W., Kir, S., Hong, S., Hu, Y., Wang, X., Binari, R., Tang, H. W., Chung, V., Banks, A. S., Spiegelman, B. &
Perrimon, N. (2019) Tumor-Derived Ligands Trigger Tumor Growth and Host Wasting via Differential MEK
Activation. *Dev Cell*, 48(2), 277-286 e6.

Stevens, L. J. & Page-McCaw, A. (2012) A secreted MMP is required for reepithelialization during wound
healing. *Mol Biol Cell*, 23(6), 1068-79.

Stine, Z. E., Schug, Z. T., Salvino, J. M. & Dang, C. V. (2022) Targeting cancer metabolism in the era of precision
oncology. *Nat Rev Drug Discov*, 21(2), 141-162.

Sun, F. & Poss, K. D. (2023) Inter-organ communication during tissue regeneration. *Development*, 150(23).

Sustar, A., Bonvin, M., Schubiger, M. & Schubiger, G. (2011) Drosophila twin spot clones reveal cell division
dynamics in regenerating imaginal discs. *Dev Biol*, 356(2), 576-87.

Vallejo, D. M., Juarez-Carreno, S., Bolivar, J., Morante, J. & Dominguez, M. (2015) A brain circuit that
synchronizes growth and maturation revealed through Dilp8 binding to Lgr3. *Science*, 350(6262), aac6767.

Witte, M. B. & Barbul, A. (2003) Arginine physiology and its implication for wound healing. *Wound Repair*
Regen, 11(6), 419-23.

Wong, K. K. L. & Verheyen, E. M. (2021) Metabolic reprogramming in cancer: Mechanistic insights from
Drosophila. *DMM Disease Models and Mechanisms.* Company of Biologists Ltd.

Worley, M. I., Everetts, N. J., Yasutomi, R., Chang, R. J., Saretha, S., Yosef, N. & Hariharan, I. K. (2022) Ets21C
sustains a pro-regenerative transcriptional program in blastema cells of Drosophila imaginal discs. *Curr Biol*,
32(15), 3350-3364 e6.

Worley, M. I. & Hariharan, I. K. (2022) Imaginal Disc Regeneration: Something Old, Something New. *Cold Spring*
Harb Perspect Biol, 14(11).

Wu, X., Xie, W., Xie, W., Wei, W. & Guo, J. (2022) Beyond controlling cell size: functional analyses of S6K in
tumorigenesis. *Cell Death Dis*, 13(7), 646.

Yue, S., Li, G., He, S. & Li, T. (2022) The Central Role of mTORC1 in Amino Acid Sensing. *Cancer Res*, 82(17),
2964-2974.

Zheng, N., Wei, J., Wu, D., Xu, Y. & Guo, J. (2023) Master kinase PDK1 in tumorigenesis. *Biochim Biophys Acta*
Rev Cancer, 1878(6), 188971.

Zhu, J. & Thompson, C. B. (2019) Metabolic regulation of cell growth and proliferation. *Nat Rev Mol Cell Biol*,
20(7), 436-450.

Zielke, N., Korzelius, J., van Straaten, M., Bender, K., Schuhknecht, G. F. P., Dutta, D., Xiang, J. & Edgar, B. A.
(2014) Fly-FUCCI: A versatile tool for studying cell proliferation in complex tissues. *Cell Rep*, 7(2), 588-598.
1173

1174