Circadian clock disruption and growth of kidney cysts in autosomal dominant polycystic kidney disease

Abeda Jamadar[1,2], Christopher J. Ward[1,2], Viji Remadevi[1,2],

Meekha M Varghese[1,2], Pabla Navjot [3], Michelle L. Gumz [4], Reena Rao[1,2] *

[1]Kidney Institute, University of Kansas Medical Center, Kansas City, KS. USA

[2]Department of Medicine, Division of Nephrology, University of Kansas Medical Center, Kansas City, KS. USA

[3]Division of Pharmaceutics and Pharmacology, College of Pharmacy & Comprehensive Cancer Center, The Ohio State University, Columbus, Ohio, USA

[4]Department of Physiology and Aging, Department of Medicine, Division of Nephrology, Hypertension, and Renal Transplantation, University of Florida, Gainesville, FL, USA.

Correspondence:

* Reena Rao, PhD

Associate Professor of Medicine

5040 WHE

Jared Grantham Kidney Institute,

University of Kansas Medical Center,

3901 Rainbow Blvd, Kansas City,

Kansas 66160-3018, USA.

E-mail: rrao@kumc.edu

913 945 6849 - Phone

Running Title: Lack of BMAL1 causes rapid cyst growth in ADPKD.

Abstract: -250

Text: -3500

Keywords: Circadian clock, BMAL1, cell proliferation, cyst, lipogenesis, fatty-acid oxidation

Abstract

Background: Autosomal dominant polycystic kidney disease (ADPKD) is caused by mutations in the *PKD1* and *PKD2* genes, and often progresses to end stage renal disease (ESRD). ADPKD progression is not uniform among patients, suggesting that factors secondary to the *PKD1/2* gene mutation could regulate the rate of disease progression. Here we tested the effect of circadian clock disruption on ADPKD progression. Circadian rhythms are regulated by cell-autonomous circadian clocks composed of clock proteins. BMAL1 is a core constituent of the circadian clock.

Methods: To disrupt the circadian clock, we knocked out *Bmal1* gene in the renal collecting ducts of the *Pkd1*$^{RC/RC}$ (RC/RC) mouse model of ADPKD, to get *Pkd1*$^{RC/RC}$;*Bmal1*$^{f/f}$;*Pkhd1*^{cre} double knockout (DKO) mouse, and in *Pkd1* knockout mouse inner medullary collecting duct cells (*Pkd1*KO mIMCD3) to get *Pkd1Bmal1*KO cells.

Results: Human nephrectomy ADPKD kidneys and *Pkd1*KO mIMCD3 cells showed reduced *Bmal1* gene expression compared to normal controls. When compared to RC/RC kidneys, DKO kidneys showed increased cyst area, cyst number, kidney/body weight ratio, cell proliferation, apoptosis and renal fibrosis. Similarly, *in vitro Pkd1Bmal1*KO cells showed increased cell proliferation compared to *Pkd1KO* cells. Expression of clock genes was significantly altered, and lipogenesis and cholesterol synthesis-related genes was increased in DKO kidneys and *Pkd1Bmal1*KO cells compared to RC/RC kidneys and *Pkd1*KO cells respectively. Moreover, fatty-acid oxidation-related gene expression was reduced in *Pkd1Bmal1*KO cells compared to *Pkd1*KO cells.

Conclusion: Renal collecting duct specific *Bmal1* gene deletion disrupts the circadian clock and triggers accelerated ADPKD progression by altering lipid metabolism-related gene expression.

Key points:

- Lack of BMAL1, a circadian clock protein in renal collecting ducts disrupted the clock and increased cyst growth in an ADPKD mouse model.

- BMAL1 gene deletion increased cell proliferation, fibrosis, lipogenesis and reduced fatty acid oxidation related gene expression.

- Thus, circadian clock disruption could be a risk factor for accelerated disease progression in patients with ADPKD.

Introduction:

Autosomal dominant polycystic kidney disease (ADPKD) is the most common inherited kidney disease which is caused by mutations in the *PKD1* and *PKD2* genes. Progressive cyst growth in ADPKD kidneys is accompanied by inflammation and fibrosis, which often leads to loss of renal function and end stage renal disease (ESRD). However, disease progression to ESRD varies significantly among ADPKD patients, even within families with the same *PKD1* gene mutation [1]. Based on the Mayo-Irazabal classification for age-adjusted assessment of ADPKD progression, the predicted incidence of ESRD at 10 years ranges widely between 2.2 to 77.4% in older adults [2]. Thus, factors secondary to the *PKD1/2* gene mutation could regulate the rate of ADPKD progression to ESRD. Here we tested if disruption of the circadian clock is a trigger for accelerated ADPKD progression.

Circadian rhythms are intrinsic, cyclical ~24-hour oscillations in behavior and physiology that coordinate biological processes with the time of day. In mammals, circadian rhythms are regulated by cell-autonomous circadian clocks, which at the molecular level are comprised of clock proteins. The central clock, located in the suprachiasmatic nucleus in the hypothalamus, synchronizes peripheral clocks based on external timing cues. The canonical circadian clock involves heterodimerization of circadian proteins (transcription factors), CLOCK and BMAL1 (activators), which activate the gene expression of *Per1,2,3* and *Cry1,2* (repressors). PER and CRY proteins accumulate over time, repressing CLOCK and BMAL1 action, thereby inhibiting their own transcription. RORα/β and REV-ERBα/β function in an ancillary loop to mediate opposing action on BMAL1 transcription, thus driving rhythmic expression of BMAL1. These feedback loops result in ~24-hour oscillations of clock gene expression, and rhythmic expression of ~40% of protein-coding genes in mammals [3].

In mouse kidneys, ~13% of all genes are expressed in a circadian manner [4, 5]. Moreover, kidney functions including blood pressure rhythms, renal blood flow, glomerular filtration rate (GFR), and excretion of electrolytes such as sodium and potassium display circadian influence [6-9]. Chronic disruption of the circadian clock (chronodisruption) is a well-known risk factor for accelerated ageing, memory loss, metabolic defects, obesity, cancer, cardiovascular defects, and metabolic syndrome in humans [10, 11]. The role of chronodisruption in ADPKD pathogenesis is currently unclear.

BMAL1 is encoded by the *Arntl* (Aryl hydrocarbon receptor nuclear translocator-like protein-1) gene and is a core component of the molecular circadian clock. Together with other clock proteins, BMAL1 plays an essential role in cellular metabolic processes by regulating time-dependent utilization of nutrients and metabolites to support cell growth [12]. *Bmal1* is a master clock gene, and the only clock gene whose deletion alone leads to complete loss of circadian rhythms [13]. Here we examined the effect of collecting duct-specific gene deletion of *Bmal1* on disease progression in an ADPKD mouse model and *Pkd1* gene knockout cells.

Methods:

Human tissue samples: Tissues from de-identified human normal and APKD kidneys were "medical discard" nephrectomy samples from the Kansas PKD Research and Translational Center. IRB approval #5929 approval date 8/25/2020. The time of collection is unknown.

***Bmal1* gene knockout in WT and *Pkd1*KO mIMCD3 cells using Crispr-Cas12a:** Murine *Bmal1* (*Arntl* gene, ID 11865) was disrupted in mIMCD3 cells using a microhomology end joining-based strategy and Cas12a [14]. The **TTTG**GCGTATCTACCACAGGAACT sequence in exon 13 was targeted (PAM in bold). CRISPR sequence was, 5'/AltR1/rUrArArUrUrUrCrUrArCrUrCrUrUrGrUrArGrArUrGrCrGrUrArUrCrUrArCrCrArCrArGrGrArArCrU/AltR2/-3'. PCR products that showed deletion on PAGE were sequenced after alkaline phosphatase treatment (ExoSAP, # 78200, Applied Biosystems, Foster City, CA). *Bmal1*KO clone and *Pkd1Bmal1*KO had frame-shift deletions of 10bp and 16bp respectively.

Mouse model: RC/RC (p.R3277C) mice [15] were bred with *Bmal1*^{f/f} mice (B6.129S4(Cg)-*Bmal1tm1Weit/J*, carrying loxP sites flanking exon 8 of *Bmal1* gene, Strain#007668, The Jackson Laboratory), and mice carrying Pkhd1^{cre} [16]. The resultant *Pkd1*^{RC/RC};*Bmal1*^{f/f};*Pkhd1*^{cre} mice (RC/RC-*Bmal1*KO) were compared with *Bmal1*^{f/f} (wild type-WT), *Bmal1*^{f/f};*Pkhd1*^{cre} (*Bmal1*KO) and RC/RC littermates. Male mice were used and sacrificed at 8 months of age, between 11:30 AM and 12:30 PM. Studies were approved by our Univerisity's IACUC committee.

Recombination specificity in *Bmal1*KO and DKO mice:

Bmal1 recombination was confirmed by PCR on genomic DNA of whole papilla from mouse kidneys [17, 18]. PCR reaction products run on a 2% agarose gel yielded a 431-bp band for floxed *Bmal1* allele and 572 bp band for recombination product.

Statistics: Values expressed as mean±SEM for *in vivo* and mean±SD for *in vitro* studies. Data were analyzed by two-tailed unpaired T-test with Welch's correction; or one-way ANOVA followed by Tukey's multiple comparison test and two-way ANOVA followed by Bonferroni's multiple comparison test using GraphPad Prism software (Version 9). P≤0.05 was considered significant.

Supplementary methods: We provide detailed methods for *in vitro* BrdU incorporation [19, 20], quantification of cyst [21], Western blot [22, 23], TUNEL staining [24, 25], immunohistochemistry/immunofluorescence (IHC/IF) staining [26] and quantitative real time PCR (QRTPCR) [27] in supplementary methods, and primer list for QRTPCR in Supplemental Table 1.

Results:

1. Renal collecting duct specific *Bmal1* gene deletion in ADPKD mouse kidneys.

Examination of *BMAL1* (*ARNTL* gene) expression in human nephrectomy ADPKD kidneys showed over 2-fold decrease in *BMAL1* mRNA levels, compared to normal human kidneys (Fig 1A). Consistent with this, the kidney interactive transcriptomics database [28] of single cell RNA-sequencing showed reduced *Bmal1* expression in most renal cells of human ADPKD kidneys compared to normal control kidneys (Fig 1B). To examine the effect of *Pkd1* gene deletion on *Bmal1*, we examined its mRNA levels in control and *Pkd1$^{-/-}$* mouse renal inner medullary collecting duct cells (mIMCD3) [29] and mouse proximal tubular cells [16, 30]. Cells were synchronized using 100nM dexamethasone for 2h [31], and released for 12h before collection every 4h, for 48h. The control *Pkd1$^{+/+}$* mIMCD3 cells and *Pkd1$^{+/-}$* proximal tubule cells showed clear circadian oscillation of *Bmal1* mRNA, with ~20h interval between peaks, while *Bmal1* mRNA levels and circadian oscillation were significantly dampened in the corresponding *Pkd1$^{-/-}$* cells (Fig 1C, D).

To determine the role of BMAL1 in ADPKD progression, we chose the *Pkd1* gene hypomorphic *Pkd1$^{RC/RC}$* (RC/RC) mouse model of ADPKD. The RC/RC mice showed no significant difference in renal *Bmal1* levels at 4 or 8 months of age compared to wild type (WT) littermates (Supplemental Fig 1A,B,C,D,E). Renal collecting duct specific *Bmal1* gene knockout in WT mice- *Bmal1$^{f/f}$;Pkhd1^{cre}* (*Bmal1*KO) and RC/RC mice- RC/RC;*Bmal1$^{f/f}$;Pkhd1^{cre}* double knockout (DKO) mice were generated. To test the effect of *Bmal1* gene deletion, the DKO mice were compared with WT (*Bmal1$^{f/f}$*), RC/RC and *Bmal1*KO littermates. We detected recombination of exon-8 of the *Bmal1* gene (disrupted allele) in the *Bmal1*KO and DKO kidneys, while recombination was not detected in the WT or RC/RC kidneys (Figure 1E). BMAL1 expression was detected in the nuclei of tubular epithelial cells in WT and RC/RC mouse kidneys (Fig 1F).

In the *Bmal1*KO and DKO kidneys, BMAL1 was not detected in some tubular or cyst epithelial cells (Fig 1F). Most large cysts in DKO kidneys showed Dolichos biflorus agglutinin (DBA) staining, implying collecting ducts (Supplemental Fig 1F).

To determine the effect of Bmal1 gene deletion on the circadian clock, we measured clock gene expression. When compared to RC/RC kidneys, the DKO kidneys showed significantly increased mRNA levels of *Clock*, *Cry1*, *RevErba* and *Tef* (Fig 1G). RC/RC kidneys showed significant decreases in *Per1*, *Dbp* and *Tef*, and increases in *Clock* mRNA levels when compared to WT kidneys (Fig 1G). *Clock*, *Cry1* and *RevErba* genes were upregulated in both *Bmal1*KO and DKO kidneys compared to WT kidneys and RC/RC kidneys respectively (Fig 1G).

2. Renal collecting duct specific *Bmal1* gene deletion increased cyst growth in ADPKD mouse kidneys.

At 8 months of age, the DKO kidneys were larger and more cystic (Fig 2A, 2B), with significantly higher cyst number (Fig 2C), cystic index (Fig 2D), kidney weight (Fig 2E) and kidney to body weight ratio (Fig 2F) compared to RC/RC kidneys. No significant changes were observed in the *Bmal1*KO kidneys compared to WT kidneys (Fig 2A,E,F). The results show that gene deletion of *Bmal1* in collecting ducts accelerates renal cyst growth in an ADPKD kidney but has no overt effect on WT kidneys.

3. *Bmal1* gene deletion increased cell turnover and fibrosis in ADPKD mouse kidneys.

To examine the effect of *Bmal1* gene deletion on cell proliferation, kidney tissue sections were immunostained for Ki67 and quantified. DKO mouse kidneys showed significantly increased Ki67 stained nuclei compared to RC/RC kidneys (Fig 3A,B). Immunoblot of kidney tissue lysate showed increased cyclin D1 and reduced P53 protein levels in DKO kidneys compared to

RC/RC kidneys (Fig 3C,D,E). In DKO kidneys, pERK/ERK, pCREB/CREB, and pS6/S6 ratios were significantly increased, suggesting higher activity compared to RC/RC kidneys (Fig 3C,F,G,H). No change was observed in c-Myc levels (Supplemental Fig 2A). No significant difference was observed between *Bmal1*KO and WT kidneys in Ki67 staining (Supplemental Fig 2B, C), or in cyclin D1, pERK/ERK or pCREB/CREB protein levels (Supplemental Fig 2D,E). TUNEL staining of kidney tissue sections (Fig 4A,B) and mRNA levels of renal injury markers, *Kim1* and *Ngal1* (Fig 4C,D) were significantly increased in DKO kidneys compared to RC/RC kidneys. The increased cell proliferation and apoptosis indicate high turnover of cells in the DKO kidneys.

DKO mouse kidneys also showed increased renal fibrosis. Immunostaining for the myofibroblast marker αSMA and the extracellular matrix (ECM) protein, collagen1 were increased in the DKO kidneys compared to RC/RC kidneys (Fig 4A). Moreover, mRNA levels of ECM proteins, *collagen1a*, *collagen3a1* and *fibronectin* were significantly higher in the DKO kidneys compared to RC/RC kidneys (Fig 4E, F, G, H).

4. *Bmal1* gene knockout altered the expression of circadian clock genes and gene targets of BMAL1 in mIMCD3 cells.

To better understand the mechanism by which *Bmal1* gene knockout in the renal collecting ducts accelerates disease progression in RC/RC mice, we deleted *Bmal1* gene in WT and *Pkd1* gene knockout mIMCD3 cell lines using Crispr-Cas12a (Fig 5A, Supplemental Fig 3). WT, *Bmal1*KO, *Pkd1*KO, and *Pkd1Bmal1*KO cells were compared. Consistent with results in mouse kidneys, cell proliferation was significantly increased in *Pkd1Bmal1*KO cells compared to *Pkd1*KO cells, while *Bmal1*KO cells showed no significant difference compared to WT cells (Fig 5B). Examination of circadian clock gene expression showed significantly reduced *Per1*, *Per2*,

RevErba and *Rora*, and increased *Cry1* mRNA levels in *Pkd1Bmal1*KO cells when compared to *Pkd1*KO cells (Fig 5C). The *Pkd1*KO cells showed significant increase in *Per1, Per2, RevErba, Rora* and *Hlf* mRNA levels compared to WT cells (Fig 5C). Significant difference in *Per2, Cry1* and *RevErba*, members of the repressor arm of the clock were common to both *Bmal1*KO and *Pkd1Bmal1*KO cells when compared to WT and *Pkd1*KO cells respectively (Fig 5C). *Rorb* and *Dbp* were not detected.

We also examined BMAL1's prospective gene targets. BMAL1, a transcription factor binds to E-box elements on gene promoters [32]. The *Pkd1* gene promoter contains an E-box element [33], although it is unknown if *Bmal1* is a transcriptional regulator of *Pkd1*. Contrary to our expectation, the *Bmal1*KO cells had significantly increased *Pkd1* mRNA levels when compared to WT cells (Supplemental Fig 4A). We also examined gene expression of *Ets1*, *Sp1*, *Rxra*, *Cebpa*, *Hnf4a*, *Fli1*, *Hnf1a*, *Hnf1b*, *Hnf4a*, *Pax8* and *Tead4*, based on a previous Chip-seq analysis [34] that identified these genes as targets of BMAL1 in WT mouse kidneys [9]. Of the genes detected, *Cebpa, Ets1* and *Hnf1b* were significantly reduced in DKO kidneys and *Pkd1Bmal1*KO cells when compared to RC/RC kidneys and *Pkd1*KO cells respectively (Supplemental Fig 4A,B).

5. *Bmal1* gene knockout altered the lipid metabolism-related gene expression in *Pkd1Bmal1*KO mIMCD3 cells and DKO mouse kidneys.

Reduced fatty-acid oxidation and increased fatty acid synthesis enhance cyst growth in ADPKD [35-37]. Since *Bmal1* is known to regulate fatty-acid oxidation, lipogenesis and cholesterol metabolism [38-42], we examined genes regulating these processes. In *Pkd1Bmal1*KO cells, mRNA levels of fatty-acid oxidation-related genes, *Acox1, Acox2, Acadsb, Acadvl, Cpt1a, Cypt2, Fabp3*, and *Ppara* showed significant decreases, while *Pgc1a* was increased compared

to *Pkd1*KO cells (Fig 5D). *Pkd1*KO cells showed significantly reduced *Acox2*, *Ppara* and *Pgc1a* mRNA levels compared to WT cells (Fig 5D). *Acox2*, *Acadsb*, *Acadvl* and *Cpt1a* are common genes that were significantly decreased in both *Bmal1*KO and *Pkd1Bmal1*KO cells compared to WT and Pkd1KO cells respectively (Fig 5D).

Lipogenesis associated genes, *Acaca*, *Acss2*, *Dagla*, *Daglb*, *Fads1*, *Fads3*, *Scd1*, and Srebp1c were significantly increased in *Pkd1Bmal1*KO cells compared to *Pkd1*KO cells (Fig 5E). *Acss2*, *Fads3* and *Srebp1c* are common genes that were upregulated in *Pkd1Bmal1*KO cells, but downregulated in the *Bmal1*KO cells compared to WT and *Pkd1*KO cells respectively (Fig 5E). Compared to WT cells, *Pkd1*KO cells showed significantly increased *Srebp1c*, and reduced *Dagla* mRNA levels (Fig 5E). Genes related to cholesterol synthesis including *Hmgcs1*, *Hmgcr*, *Mvk* and *Mvd* were significantly increased in *Pkd1Bmal1*KO cells when compared to *Pkd1*KO cells (Fig 5F). When compared to WT cells, the *Pkd1*KO cells showed significantly reduced levels of *Hmgcs1*, *Hmgcsr*, *Mvk*, *Pmvk*, *Fdps* and *Mvd* (Fig 5F). Other fatty acid oxidation and lipogenesis-related genes that remained unchanged are shown in Supplemental Fig 5A,B.

Examination of lipogenesis-related genes in our mouse study showed significantly increased mRNA levels of *Acaca*, *Dagla*, *Fads1*, *Scd1*, *Srebp1c*, *Mag1* and *Lpcat* genes in DKO mouse kidneys compared to RC/RC kidneys (Fig 6A-H). Similarly, cholesterol metabolism-related *Mvk*, *Pmvk* and *Fdps6* genes were significantly increased in DKO kidneys compared to RC/RC kidneys (Fig 6I, J, K). Fatty-acid oxidation-related genes were not significantly different between DKO and RC/RC kidneys. However, RC/RC kidneys showed significantly reduced expression of *Acox1*, *Acadsb*, *Ppara*, *Acat1*, *Acat2* and *Acads* compared to WT kidneys (Supplemental Fig 6 A-N). Unchanged lipogenesis and cholesterol synthesis-related genes are shown in Supplemental Fig 7A-I.

Discussion

Our studies show that disruption of the circadian clock stimulates accelerated cyst growth in ADPKD. We found reduced *Bmal*1 gene expression in human nephrectomy ADPKD kidneys, and in *Pkd1* gene knockout mIMCD3 and proximal tubule cells, but not in *Pkd1* gene hypomorphic RC/RC mouse kidneys. Renal collecting duct specific gene deletion of *Bmal1* in RC/RC mice increased renal cell proliferation, apoptosis, fibrosis, and cyst growth. *Bmal1* gene deletion significantly altered the expression of other clock genes, reduced the expression of BMAL1 target genes and increased the expression of lipogenesis and cholesterol synthesis-related genes in *Pkd1*KO cells *in vitro*, and in RC/RC mouse kidneys. *Bmal1* gene deletion also reduced the expression of fatty-acid oxidation-related gene expression in *Pkd1*KO mIMCD3 cells.

Our finding that *Bmal1* gene knockout in the renal collecting ducts increased cyst growth in ADPKD mice is significant because it shows for the first time that chronodisruption accelerates ADPKD progression. Chronodisruption is disease inducing, and known to directly drive metabolic dysfunction, cause type-2 diabetes, obesity, various cancers and neurodegenerative conditions, and adversely affect fetal development [43-47]. Environmental or behavioral factors such as mistimed feeding, shift work or jet lag can induce chronodisruption. Shift work for instance, reduces the estimated GFR in humans and is a risk factor for CKD [48-50], cardiovascular disease [51], multiple cancers [52] and metabolic disorders [53]. Since chronodisruption by alterations in behavior or the environment are systemic effects, we used renal collecting duct specific *Bmal1* gene deletion as a model for genetic disruption of the molecular clock so as to focus on the kidney. In our study, *Bmal1* gene knockout altered the expression of clock genes including *Cry1*, *Per1*, *Per2*, and *RevErba* that are in the negative arm of the clock, and are known to be regulated by *Bmal1*.

We found that collecting duct-specific gene deletion of *Bmal1* dramatically increased cyst growth, cell proliferation and fibrosis in the RC/RC mouse model of ADPKD. In prior studies, *Bmal1* gene knockout reduced locomotor activity, and induced tendon calcification, sarcopenia, hypersensitivity to insulin-shock, fragmented sleep, and rapid ageing in mice [54, 55]. In the kidneys, nephron-specific *Bmal1* gene deletion caused Na^+ retention in response to a K^+-restricted diet [56], lowered blood pressure [56-58] and altered the urinary metabolome in mice [54]. However, the effect of altering *Bmal1* gene expression on kidney disease progression is context dependent, as shown in experimental mouse models [59]. In a mouse model of AKI, shRNA mediated *Bmal1* gene knockout alleviated AKI, while adenoviral overexpression of *Bmal1* accelerated renal injury [60]. Similarly in unilateral ureteral obstruction model of CKD, global gene knockout of *Bmal1* reduced fibrosis [61] or had no effect on fibrosis or disease progression [62]. In contrast to the above mentioned studies, and supporting our results, renal tubule specific *Bmal1* gene knockout worsened hyperglycemia, and accelerated renal hypertrophy in mice with streptozotocin-induced type-1 diabetes, compared to similarly treated WT mice [63]. Similarly, global or proximal tubule specific gene knockout of *Bmal1* worsened disease progression in the adenine induced model of CKD in mice [64,65]. We found that in the presence of *Pkd1* gene mutation, *Bmal1* gene knockout in renal collecting ducts increased cell proliferation indicated by increase in Ki-67 immunostaining, and pERK/ERK1/2, pCREB/CREB, pS6/S6 and CyclinD1 levels, all factors known to increase cell proliferation in ADPKD kidneys [24]. Our data is consistent with previous studies where *Bmal1* gene deletion or inactivation increases the proliferation rate of cancer cells [66, 67].

BMAL1 binds to E-box elements on promoters of rhythmically and non-rhythmically expressed genes [32]. Over 7 million E-box elements exist in the mouse genome, ~ 0.7% of which

are bound by CLOCK-BMAL1 or NPAS2-BMAL1 dimers in mouse peripheral tissues [34, 68].

Based on this published data, we examined 12 prospective gene targets of BMAL1 and

identified *Ets1*, *Hnf1b* and *Cebpa* to be significantly reduced in *Bmal1* deficient *Pkd1*KO cells

and RC/RC mouse kidneys. C/EBPα plays an essential role in embryonic development and

supresses cell proliferation by stabilizing p21 and inhibiting *cdk2*, *cdk4* and *E2F* [69]. Inhibiting

C/EBPα contributes to tumor progression in liver, pancreatic, breast and lung cancers [69].

Similarly, ETS1 and HNF1 transcription factors have oncogenic or tumor supressor roles in a

context-dependent manner in human cancers [70, 71]. HNF1β and ETS1 are also thought to

regulate *Pkd1* gene expression [33, 72, 73].

Fatty-acid metabolism is impaired in ADPKD and is thought to enhance renal cyst growth

[35-37]. Previous studies in RC/RC mice have shown reduced renal mRNA levels of fatty-acid

oxidation-related genes compared to WT mice, and treatment with fenofibrate, a PPARα agonist

inhibited cyst growth in the RC/RC mice [35]. Similarly, pups born from female mice fed with a high

fat diet had a significantly higher cystic index at P21 as compared to those fed with low fat diet

[36]. *In vitro* studies have also shown that *Pkd1* gene mutation leads to defects in fatty-acid

oxidation [36, 37]. Consistent with these studies, we found fatty-acid oxidation-related genes to be

significantly reduced in *Pkd1*KO cells compared to WT cells, and in RC/RC mouse kidneys

compared to WT mouse kidneys.

BMAL1 is known to regulate fatty-acid oxidation [38-42]. Bignon *et al* [42] showed that renal

tubule specific *Bmal1* gene deletion in WT mice significantly reduced mRNA and protein

expression of ACADS, ACADM, ACADL, and ACADVL which catalyze the rate-limiting step of β-

oxidation, suggesting disrupted fatty-acid oxidation. However in our study, renal collecting duct

specific *Bmal1* gene deletion in the *Bmal1*KO mice or DKO mice did not alter fatty-acid

oxidation-related genes when compared to WT or RC/RC mouse kidneys. The reason for this disparity is presently unclear. Our results from the mIMCD3 cells were different, with 4 of the 14 fatty acid oxidation-related genes showing significant decreases in the *Bmal1*KO and *Pkd1Bmal1*KO cells compared to WT and *Pkd1*KO cells respectively.

Unlike fatty-acid oxidation, our *in vitro* and *in vivo* results consistently showed increased lipogenesis and cholesterol metabolism-related genes in *Bmal1* gene deleted cells and mouse kidneys. The mRNA levels of 7 out of the 14 lipogenesis-related genes tested in *Pkd1Bmal1*KO cells and in DKO kidneys showed significant increase compared to WT cells and RC/RC kidneys. Of the lipogenesis-related genes, *Acaca*, *Fads1*, *Scd1* and *Srebp1c* were upregulated in both DKO mouse kidneys and in *Pkd1Bmal1*KO cells, suggesting regulation by *Bmal1*. The *Pkd1Bmal1*KO cells and DKO kidneys showed significant increases in several cholesterol synthesis-related genes. Mvk was a frequently upregulated genes in the cells and kidneys. Renal collecting duct specific *Bmal1* gene deletion in WT mice (*Bmal1*KO mice) did not alter any lipogenesis, cholesterol synthesis or fatty-acid oxidation-related genes when compared to WT mouse kidneys, suggesting that *Bmal1* gene knockout in the ADPKD kidneys is important for these changes.

Although *Bmal1* gene deletion altered the expression of multiple clock genes and BMAL1 regulated genes and lipid metabolism-related genes in the mIMCD3 cells and mouse kidneys, their changes were not always similar in the cells and kidneys. This could be because *Bmal1* is deleted only in the collecting duct cells of the mouse kidneys, while the IMCD3 cells had complete gene deletion of *Bmal1*. Moreover, the mice were all sacrificed around 12 PM, while the mIMCD3 cells were collected 24h after synchronization *in vitro*.

While patients with ADPKD often develop non-dipping hypertension, polyuria, polydipsia, and nocturnal polyuria [74-76], it had not previously been shown whether chronodisruption accelerates ADPKD. Our results show that *Bmal1* gene knockout in the renal collecting ducts disrupts the circadian clock and expression of key elements of cell proliferation, fibrosis and lipid metabolism, thereby accelerating cyst growth in our ADPKD models. It is possible, therefore, that chronodisruption in patients with ADPKD might be a cause of, or contribute to more rapid disease progression.

Disclosure Statement:

The authors have declared that no conflict of interest exists.

Funding:

This study was supported by, DOD Discovery Award W81XWH2310011, NIHR56DK128962-01 and NIH1R01DK135308-01 grants to RR.

Acknowledgments:

We thank Dr. Peter Harris and Dr. Katharina Hopp for providing *Pkd1*RC/RC mice; Dr. Stefan Somlo and Dr. Yiqiang Cai (George M. O'Brien Kidney Center, Yale University, New Haven, CT) for *Pkd1*KO and control proximal tubular and inner medullary collecting ducts cells; and Dr. Darren Wallace for human kidney tissues (NIDDK PKD-Research Resource Consortium and the Kansas PKD Research and Translational Core Center-U54 DK126126).

Author contributions:

RR conceptualized and designed studies, analyzed the results, and wrote the paper. AJ performed experiments, analyzed results and wrote parts of the paper. VR, CJW and MMS performed some studies, analyzed results and edited the paper. MLG conceptualized studies and reviewed the paper. All authors read, edited and approved the paper.

Data sharing statements:

No proteomics, transcriptomics or GWAS data was generated in this paper. All in vivo studies using mouse models were done, and tissues analyzed at the University of

Kansas Medical Center. All data that was generated are provided in the figures in the main text or in Supplementary Material. The datasets used to examine Bmal1 expression levels in human samples are openly available on the kidney interactive transcriptomics database (https://humphreyslab.com/SingleCell/) and permission was obtained to use it. No proteomics, transcriptomics or GWAS data was generated in this paper. If any added information is required to re-examine or analyze the data shown in this paper, they will be readily available from the lead author. All requests for data will be processed based on institutional policies for noncommercial research purposes. Data sharing could require a data transfer agreement as determined by the University of Kansas Medical Center's legal department.

REFERENCE

1. Rossetti S, Kubly VJ, Consugar MB*, et al.* Incompletely penetrant PKD1 alleles suggest a role for gene dosage in cyst initiation in polycystic kidney disease. *Kidney Int* 2009; **75:** 848-855.

2. Irazabal MV, Rangel LJ, Bergstralh EJ*, et al.* Imaging classification of autosomal dominant polycystic kidney disease: a simple model for selecting patients for clinical trials. *J Am Soc Nephrol* 2015; **26:** 160-172.

3. Xie Y, Tang Q, Chen G*, et al.* New Insights Into the Circadian Rhythm and Its Related Diseases. *Front Physiol* 2019; **10:** 682.

4. Wolff CA, Gutierrez-Monreal MA, Meng L*, et al.* Defining the age-dependent and tissue-specific circadian transcriptome in male mice. *Cell Rep* 2023; **42:** 111982.

5. Zhang R, Lahens NF, Ballance HI*, et al.* A circadian gene expression atlas in mammals: implications for biology and medicine. *Proc Natl Acad Sci U S A* 2014; **111:** 16219-16224.

6. Bronner F, Lemaire R. Comparison of calcium kinetics in man and the rat. *Calcif Tissue Res* 1969; **3:** 238-248.

7. Firsov D, Bonny O. Circadian rhythms and the kidney. *Nat Rev Nephrol* 2018; **14:** 626-635.

8. Gumz ML. Molecular basis of circadian rhythmicity in renal physiology and pathophysiology. *Exp Physiol* 2016; **101:** 1025-1029.

9. Johnston JG, Pollock DM. Circadian regulation of renal function. *Free Radic Biol Med* 2018; **119:** 93-107.

10. Mohandas R, Douma LG, Scindia Y*, et al.* Circadian rhythms and renal pathophysiology. *J Clin Invest* 2022; **132**.

11. Rivera AS, Akanbi M, O'Dwyer LC*, et al.* Shift work and long work hours and their association with chronic health conditions: A systematic review of systematic reviews with meta-analyses. *PLoS One* 2020; **15:** e0231037.

12. Kinouchi K, Sassone-Corsi P. Metabolic rivalry: circadian homeostasis and tumorigenesis. *Nat Rev Cancer* 2020; **20:** 645-661.

13. Bunger MK, Wilsbacher LD, Moran SM*, et al.* Mop3 is an essential component of the master circadian pacemaker in mammals. *Cell* 2000; **103:** 1009-1017.

14. Ata H, Ekstrom TL, Martinez-Galvez G*, et al.* Robust activation of microhomology-mediated end joining for precision gene editing applications. *PLoS Genet* 2018; **14:** e1007652.

15. Hopp K, Ward CJ, Hommerding CJ*, et al.* Functional polycystin-1 dosage governs autosomal dominant polycystic kidney disease severity. *J Clin Invest* 2012; **122:** 4257-4273.

16. Kakade VR, Tao S, Rajagopal M*, et al.* A cAMP and CREB-mediated feed-forward mechanism regulates GSK3beta in polycystic kidney disease. *J Mol Cell Biol* 2016; **8:** 464-476.

17. Crislip GR, Wohlgemuth SE, Wolff CA*, et al.* Apparent Absence of BMAL1-Dependent Skeletal Muscle-Kidney Cross Talk in Mice. *Biomolecules* 2022; **12**.

18. Rao R, Patel S, Hao C*, et al.* GSK3beta mediates renal response to vasopressin by modulating adenylate cyclase activity. *J Am Soc Nephrol* 2010; **21:** 428-437.

19. Jamadar A, Dwivedi N, Mathew S*, et al.* Vasopressin Receptor Type-2 Mediated Signaling in Renal Cell Carcinoma Stimulates Stromal Fibroblast Activation. *International Journal of Molecular Sciences* 2022; **23:** 7601.

20. Jamadar A, Suma SM, Mathew S*, et al.* The tyrosine-kinase inhibitor Nintedanib ameliorates autosomal-dominant polycystic kidney disease. *Cell Death Dis* 2021; **12:** 947.

21. Tao S, Kakade VR, Woodgett JR*, et al.* Glycogen synthase kinase-3beta promotes cyst expansion in polycystic kidney disease. *Kidney Int* 2015; **87:** 1164-1175.

22. Norregaard R, Tao S, Nilsson L*, et al.* Glycogen synthase kinase 3alpha regulates urine concentrating mechanism in mice. *Am J Physiol Renal Physiol* 2015; **308:** F650-660.

23. Sinha S, Dwivedi N, Woodgett J*, et al.* Glycogen synthase kinase-3beta inhibits tubular regeneration in acute kidney injury by a FoxM1-dependent mechanism. *FASEB J* 2020; **34:** 13597-13608.

24. Dwivedi N, Tao S, Jamadar A*, et al.* Epithelial Vasopressin Type-2 Receptors Regulate Myofibroblasts by a YAP-CCN2-Dependent Mechanism in Polycystic Kidney Disease. *J Am Soc Nephrol* 2020; **31:** 1697-1710.

25. Sinha S, Dwivedi N, Tao S*, et al.* Targeting the vasopressin type-2 receptor for renal cell carcinoma therapy. *Oncogene* 2020; **39:** 1231-1245.

26. Singh SP, Tao S, Fields TA*, et al.* Glycogen synthase kinase-3 inhibition attenuates fibroblast activation and development of fibrosis following renal ischemia-reperfusion in mice. *Dis Model Mech* 2015; **8:** 931-940.

27. Dwivedi N, Jamadar A, Mathew S*, et al.* Myofibroblast depletion reduces kidney cyst growth and fibrosis in autosomal dominant polycystic kidney disease. *Kidney Int* 2023; **103:** 144-155.

28. Muto Y, Dixon EE, Yoshimura Y*, et al.* Defining cellular complexity in human autosomal dominant polycystic kidney disease by multimodal single cell analysis. *Nat Commun* 2022; **13:** 6497.

29.   Krappitz M, Bhardwaj R, Dong K, *et al.* XBP1 Activation Reduces Severity of Polycystic Kidney Disease due to a Nontruncating Polycystin-1 Mutation in Mice. *J Am Soc Nephrol* 2023; **34:** 110-121.

30.   Shibazaki S, Yu Z, Nishio S, *et al.* Cyst formation and activation of the extracellular regulated kinase pathway after kidney specific inactivation of Pkd1. *Hum Mol Genet* 2008; **17:** 1505-1516.

31.   Lee JY, Jung E, Yeo H, *et al.* The Natural Janus Kinase Inhibitor Agerarin Downregulates Interleukin-4-Induced PER2 Expression in HaCaT Keratinocytes. *Molecules* 2022; **27**.

32.   Rey G, Cesbron F, Rougemont J, *et al.* Genome-wide and phase-specific DNA-binding rhythms of BMAL1 control circadian output functions in mouse liver. *PLoS Biol* 2011; **9:** e1000595.

33.   Lantinga-van Leeuwen IS, Leonhard WN, Dauwerse H, *et al.* Common regulatory elements in the polycystic kidney disease 1 and 2 promoter regions. *Eur J Hum Genet* 2005; **13:** 649-659.

34.   Beytebiere JR, Trott AJ, Greenwell BJ, *et al.* Tissue-specific BMAL1 cistromes reveal that rhythmic transcription is associated with rhythmic enhancer-enhancer interactions. *Genes Dev* 2019; **33:** 294-309.

35.   Lakhia R, Yheskel M, Flaten A, *et al.* PPARalpha agonist fenofibrate enhances fatty acid beta-oxidation and attenuates polycystic kidney and liver disease in mice. *Am J Physiol Renal Physiol* 2018; **314:** F122-F131.

36.   Menezes LF, Lin CC, Zhou F, *et al.* Fatty Acid Oxidation is Impaired in An Orthologous Mouse Model of Autosomal Dominant Polycystic Kidney Disease. *EBioMedicine* 2016; **5:** 183-192.

37.   Podrini C, Rowe I, Pagliarini R, *et al.* Dissection of metabolic reprogramming in polycystic kidney disease reveals coordinated rewiring of bioenergetic pathways. *Commun Biol* 2018; **1:** 194.

38.   Zhang D, Tong X, Nelson BB, *et al.* The hepatic BMAL1/AKT/lipogenesis axis protects against alcoholic liver disease in mice via promoting PPARalpha pathway. *Hepatology* 2018; **68:** 883-896.

39.   Zhang D, Tong X, Arthurs B, *et al.* Liver clock protein BMAL1 promotes de novo lipogenesis through insulin-mTORC2-AKT signaling. *J Biol Chem* 2014; **289:** 25925-25935.

40.   Pan X, Mota S, Zhang B. Circadian Clock Regulation on Lipid Metabolism and Metabolic Diseases. *Adv Exp Med Biol* 2020; **1276:** 53-66.

41.   Pan X, Bradfield CA, Hussain MM. Global and hepatocyte-specific ablation of Bmal1 induces hyperlipidaemia and enhances atherosclerosis. *Nat Commun* 2016; **7:** 13011.

42.   Bignon Y, Wigger L, Ansermet C, *et al.* Multiomics reveals multilevel control of renal and systemic metabolism by the renal tubular circadian clock. *J Clin Invest* 2023; **133**.

43.   Huang C, Zhang C, Cao Y, *et al.* Major roles of the circadian clock in cancer. *Cancer Biol Med* 2023; **20:** 1-24.

44.   Mendez N, Corvalan F, Halabi D*, et al.* From gestational chronodisruption to noncommunicable diseases: Pathophysiological mechanisms of programming of adult diseases, and the potential therapeutic role of melatonin. *J Pineal Res* 2023; **75:** e12908.

45.   Parameswaran G, Ray DW. Sleep, circadian rhythms, and type 2 diabetes mellitus. *Clin Endocrinol (Oxf)* 2022; **96:** 12-20.

46.   Paschos GK. Diurnal rhythms and obesity. *Curr Opin Clin Nutr Metab Care* 2021; **24:** 333-338.

47.   Verma AK, Singh S, Rizvi SI. Aging, circadian disruption and neurodegeneration: Interesting interplay. *Exp Gerontol* 2023; **172:** 112076.

48.   Sasaki S, Yoshioka E, Saijo Y*, et al.* Short sleep duration increases the risk of chronic kidney disease in shift workers. *J Occup Environ Med* 2014; **56:** 1243-1248.

49.   Ko BS, Shin SY, Hong JE*, et al.* Effect of night shift work on the reduction of glomerular filtration rate using data from Korea Medical Institute (2016-2020). *Ann Occup Environ Med* 2023; **35:** e22.

50.   Charles LE, Gu JK, Fekedulegn D*, et al.* Association between shiftwork and glomerular filtration rate in police officers. *J Occup Environ Med* 2013; **55:** 1323-1328.

51.   Dochi M, Suwazono Y, Sakata K*, et al.* Shift work is a risk factor for increased total cholesterol level: a 14-year prospective cohort study in 6886 male workers. *Occup Environ Med* 2009; **66:** 592-597.

52.   Kaakour D, Fortin B, Masri S*, et al.* Circadian Clock Dysregulation and Prostate Cancer: A Molecular and Clinical Overview. *Clin Med Insights Oncol* 2023; **17:** 11795549231211521.

53.   Hermansson J, Hallqvist J, Karlsson B*, et al.* Shift work, parental cardiovascular disease and myocardial infarction in males. *Occup Med (Lond)* 2018; **68:** 120-125.

54.   Nikolaeva S, Ansermet C, Centeno G*, et al.* Nephron-Specific Deletion of Circadian Clock Gene Bmal1 Alters the Plasma and Renal Metabolome and Impairs Drug Disposition. *J Am Soc Nephrol* 2016; **27:** 2997-3004.

55.   Takahashi JS, Hong HK, Ko CH*, et al.* The genetics of mammalian circadian order and disorder: implications for physiology and disease. *Nat Rev Genet* 2008; **9:** 764-775.

56.   Crislip GR, Douma LG, Masten SH*, et al.* Differences in renal BMAL1 contribution to Na(+) homeostasis and blood pressure control in male and female mice. *Am J Physiol Renal Physiol* 2020; **318:** F1463-F1477.

57.   Tokonami N, Mordasini D, Pradervand S*, et al.* Local renal circadian clocks control fluid-electrolyte homeostasis and BP. *J Am Soc Nephrol* 2014; **25:** 1430-1439.

58.   Zhang D, Jin C, Obi IE, *et al.* Loss of circadian gene Bmal1 in the collecting duct lowers blood pressure in male, but not female, mice. *Am J Physiol Renal Physiol* 2020; **318:** F710-F719.

59.   Juffre A, Gumz ML. Recent advances in understanding the kidney circadian clock mechanism. *Am J Physiol Renal Physiol* 2024; **326:** F382-F393.

60.   Zha M, Tian T, Xu W, *et al.* The circadian clock gene Bmal1 facilitates cisplatin-induced renal injury and hepatization. *Cell Death Dis* 2020; **11:** 446.

61.   Zhang J, Liu C, Liang Q, *et al.* Postnatal deletion of Bmal1 in mice protects against obstructive renal fibrosis via suppressing Gli2 transcription. *FASEB J* 2021; **35:** e21530.

62.   Rey-Serra C, Tituana J, Lin T, *et al.* Reciprocal regulation between the molecular clock and kidney injury. *Life Sci Alliance* 2023; **6**.

63.   Ansermet C, Centeno G, Bignon Y, *et al.* Dysfunction of the circadian clock in the kidney tubule leads to enhanced kidney gluconeogenesis and exacerbated hyperglycemia in diabetes. *Kidney Int* 2022; **101:** 563-573.

64.   Fang Y, Jo SK, Park SJ, *et al.* Role of the Circadian Clock and Effect of Time-Restricted Feeding in Adenine-Induced Chronic Kidney Disease. *Lab Invest* 2023; **103:** 100008.

65.   Liu C, Li S, Ji S, *et al.* Proximal tubular Bmal1 protects against chronic kidney injury and renal fibrosis by maintaining of cellular metabolic homeostasis. *Biochim Biophys Acta Mol Basis Dis* 2023; **1869:** 166572.

66.   Gwon DH, Lee WY, Shin N, *et al.* BMAL1 Suppresses Proliferation, Migration, and Invasion of U87MG Cells by Downregulating Cyclin B1, Phospho-AKT, and Metalloproteinase-9. *Int J Mol Sci* 2020; **21**.

67.   Elshazley M, Sato M, Hase T, *et al.* The circadian clock gene BMAL1 is a novel therapeutic target for malignant pleural mesothelioma. *Int J Cancer* 2012; **131:** 2820-2831.

68.   Marri D, Filipovic D, Kana O, *et al.* Prediction of mammalian tissue-specific CLOCK-BMAL1 binding to E-box DNA motifs. *Sci Rep* 2023; **13:** 7742.

69.   Lourenco AR, Coffer PJ. A tumor suppressor role for C/EBPalpha in solid tumors: more than fat and blood. *Oncogene* 2017; **36:** 5221-5230.

70.   Kim GC, Lee CG, Verma R, *et al.* ETS1 Suppresses Tumorigenesis of Human Breast Cancer via Trans-Activation of Canonical Tumor Suppressor Genes. *Front Oncol* 2020; **10:** 642.

71.   Chandra S, Srinivasan S, Batra J. Hepatocyte nuclear factor 1 beta: A perspective in cancer. *Cancer Med* 2021; **10:** 1791-1804.

72.   Puri S, Rodova M, Islam MR, *et al.* Ets factors regulate the polycystic kidney disease-1 promoter. *Biochem Biophys Res Commun* 2006; **342:** 1005-1013.

73.   Bergmann C, von Bothmer J, Ortiz Bruchle N, *et al.* Mutations in multiple PKD genes may explain early and severe polycystic kidney disease. *J Am Soc Nephrol* 2011; **22:** 2047-2056.

74.   Sans-Atxer L, Torra R, Fernandez-Llama P. Hypertension in autosomal-dominant polycystic kidney disease (ADPKD). *Clin Kidney J* 2013; **6:** 457-463.

75.   Seeman T, Dusek J, Vondrak K*, et al.* Renal concentrating capacity is linked to blood pressure in children with autosomal dominant polycystic kidney disease. *Physiol Res* 2004; **53:** 629-634.

76.   Zittema D, Boertien WE, van Beek AP*, et al.* Vasopressin, copeptin, and renal concentrating capacity in patients with autosomal dominant polycystic kidney disease without renal impairment. *Clin J Am Soc Nephrol* 2012; **7:** 906-913.

Figure Legends:

Figure1. Renal collecting duct specific *Bmal1* gene knockout in WT and RC/RC mice.

(A) *Bmal1* mRNA levels in normal human kidneys (NHK) and ADPKD kidneys measured by QRTPCR. ***P<0.001 by T-test (B) *Bmal1* gene expression in human normal control and ADPKD kidneys using kidney interactive transcriptomics database of single cell RNA-seq data. (C) Time course showing fold change in *Bmal1* mRNA relative to 18S mRNA in mouse inner medullary collecting duct cells (mIMCD3) (n=4), and (D) proximal tubule cells (n=3). ***P<0.001 by 2-way ANOVA for C and D. Cells were synchronized using dexamethasone (100nM, for 2h) and released into media with 5% FBS. After 12h, cell lysates were collected every 4 hours for 48h. (E) Agarose gel showing recombination PCR products of genomic DNA obtained from mouse kidney. The 431 bp band denotes the floxed *Bmal1* allele and the 572 bp band represents the recombination product. A single 431 bp band indicates absence of recombination. (F) Immunostaining for BMAL1 (brown) in kidney tissue sections of WT, *Bmal1*KO, RC/RC and RC/RC;*Bmal1*KO (DKO) mice (male, 8 months old) (Scale bar = 100µm). (G) QRTPCR analysis of mRNA levels of clock genes in whole kidney tissue lysates. *=*vs* WT, # =*vs Bmal1*KO and $=*vs* RC/RC. *P<0.05, **P<0.01, ***P<0.001 by One-way ANOVA. n=5 in WT and *Bmal1*KO, and n=7 in RC/RC and DKO.

Figure 2. Renal collecting duct specific *Bmal1* gene knockout increased cyst growth in RC/RC mice. (A) Images of H&E staining of kidney sections and (B) magnified images. Scale bar = 1 mm for A and 400µM for B. (C) Cyst number, (D) cystic index, (E) total kidney weight, and (F) two kidney to body-weight ratio. **P<0.01, ***P<0.001 by T-test for C and D, and One-way ANOVA for E and F.

Fig 3. Renal collecting duct specific *Bmal1* gene knockout increased cell proliferation in RC/RC mouse kidneys. (A) Immunostaining for Ki-67 in mouse kidney tissue sections (brown nuclear staining denotes Ki-67). Scale bar = 100 µm. (B) Quantitation of Ki-67-stained nuclei per kidney section. (C) Immunoblot on kidney tissue lysate and quantitation of band density for (D) Cyclin D1, (E) P53, (F) pERK/ERK ratio, (G) pCREB/CREB ratio, (H) pS6/S6 ratio. $*P<0.05$, $**P<0.01$, $***P<0.001$ by T-test.

Fig 4. Renal collecting duct specific *Bmal1* gene knockout increased apoptosis and fibrosis in RC/RC mouse kidneys. (A) TUNEL staining (green) (Scale bar 400µm), and immunostaining for αSMA (red), and Collagen1 (green) (Scale bar 100µm) in mouse kidney tissue sections. (B) Quantitation of TUNEL staining per kidney tissue section. (C) QRTPCR for *Kim1*, (D) *Ngal1* (E) *αSMA*, (F) *Collagen 1a1*, (G) *Collagen 3a1,* and (H) *Fibronectin* in kidney tissue lysates. $*P<0.05$, $**P<0.01$, $***P<0.001$ by T-test in B and One-way ANOVA in C, D, E, F, G and H.

Fig 5. *Bmal1* gene knockout increased cell proliferation and altered clock genes and altered lipid metabolism-related gene expression in *Pkd1*KO mIMCD3 cells. A) Western blot shows WT and *Bmal1*KO clones in WT and *Pkd1* gene knockout mIMCD3 cells. WT-*Bmal1*KO clone #4 and *Pkd1Bmal1*KO clone# 3 were used for further studies. (B) Cell proliferation assessed by BrdU incorporation assay. $***P<0.001$ by One-way ANOVA. (C) QRTPCR analysis for mRNA levels of clock genes in cells. (D) mRNA levels of genes that regulate fatty acid oxidation, (E) lipogenesis, and (F) cholesterol synthesis. * = *vs* WT, # = *vs Bmal1*KO and $ = *vs Pkd1*KO. $*P<0.05$, $**P<0.01$, $***P<0.001$ by One-way ANOVA. n=4 in C,D,E and F.

Fig 6. Renal collecting duct specific *Bmal1* gene knockout increased lipogenesis and cholesterol metabolism-related gene expression in RC/RC mouse kidneys. The mRNA levels of genes that regulate lipogenesis, (A) *Acaca*, (B) *Dagla*, (C) *Fads1*, (D) *Scd1*, (E) *Srebp1c*, (F) *Magl*, (G) *Lpcat*, (H) *Elovl6*; and genes that regulate cholesterol metabolism, (I) *Mvk*, (J) *Pmvk*, and (K) *Fdps*. *P<0.05, **P<0.01, ***P<0.001 by One-way ANOVA.

Supplemental Figure legends:

Supplemental 1. *Bmal1* expression in WT and RC/RC mouse kidneys: (A) *Bmal1* mRNA expression measured by QRTPCR in WT and RC/RC kidneys at 4 and 8 months of age. (B) Western blot for BMAL1 and (C) densitometry for 4 month old mouse kidneys (WT, n=2, RC/RC, n=4), and (D) Western blot and (E) densitometry for 8 month old mouse kidneys (WT, n=6, RC/RC, n=7). (F) Immunostaining for DBA (red) and DAPI (blue).

Supplemental 2. Cell proliferation in mouse kidneys: (A) Western blot for c-Myc. (B) Ki67 stained kidney section (Scale bar 100µm) and (C) Ki-67 quantitation (n=6) (D) Western blot and (E) densitometry for pro-proliferative proteins in WT and *Bmal1*KO mouse kidneys (n=6)..

Supplement 3. *Bmal1* gene knockout clones of WT and *Pkd1*KO mIMCD3 cells: (A) Gene knockout of *Bmal1* by CRISPR/Cas12a in WT and *Pkd1* gene knockout mIMCD3 cells. Sequences of wild-type mIMCD3 (WT), and the selected clones #4 of WT and clone #3 of *Pkd1*KO mIMCD3 cells, validating the deletion of base pairs in exon 2.

Supplemental 4. BMAL1 regulated genes: (A) QRTPCR for mRNA levels of BMAL1 target genes in mIMCD3 cell lysates and (B) in mouse kidney tissue lysates. * = *vs* WT, # = *vs Bmal1*KO and $ = *vs Pkd1*KO. *P<0.05, **P<0.01, ***P<0.001 by by One-way ANOVA.

Supplemental 5. Fatty acid oxidation and lipogenesis-related gene expression in *Pkd1*KO mIMCD3 cells. (A) QRTPCR analysis for mRNA levels of genes related to fatty acid oxidation (n=4) and (C) lipogenesis (n=4). *P<0.05, **P<0.01, ***P<0.001 by by One-way ANOVA.

Supplemental 6. Fatty-acid oxidation-related gene expression in mouse kidneys. The mRNA levels of genes that regulate fatty-acid oxidation. *P<0.05, **P<0.01, ***P<0.001 by One-way ANOVA.

Supplemental 7. Lipogenesis and cholesterol metabolism-related gene expression in mouse kidneys. QRTPCR analysis for mRNA levels of genes that regulate lipogenesis and cholesterol metabolism in whole kidney lysates.

(A) RC/RC (Pkd1$^{RC/RC}$)

DKO (Pkd1$^{RC/RC}$;Bmal1$^{f/f}$;Pkhd1Cre)

WT

Bmal1KO (Bmal1$^{f/f}$;Pkhd1Cre)

(B) RC/RC DKO

Lipogenesis

Cholesterol synthesis