Assessing the effects of female protections on size structure and spawning potential in two

clawed lobster fisheries subject to varying exploitation levels

Theberge, Kaitlyn[1], Sørdalen, Tonje K. [2], Pugh, Tracy L.[3], Kindsvater, Holly K. [1,*]

[1]Department of Fish and Wildlife Conservation, Virginia Tech, 310 West Campus Dr. Ste 11,

Blacksburg, VA 24061

[2]Centre for Coastal Research, Department of Natural Sciences, University of Agder, 4630

Kristiansand, Norway

[3]Massachusetts Division of Marine Fisheries, 836 South Rodney French Blvd., New Bedford,

MA 02744

*Corresponding author, hkindsvater@vt.edu, Department of Fish and Wildlife Conservation,

Virginia Tech, 310 West Campus Dr. Ste 11, Blacksburg, VA 24061

Keywords,

fisheries, size-selectivity, v-notching, fishery management, demographic modelling, crustacean,

population dynamics

ABSTRACT

Current fishery management practices in both the U.S. Gulf of Maine and Norwegian

clawed lobster fisheries primarily focus on conserving mature females to maximize egg

production. While abundance of adult American lobsters (*Homarus americanus*) in the Gulf of

Maine remains high, declines appear to be on the horizon. Similarly, the European lobster

(*Homarus gammarus*), is facing its lowest recorded population size in southern Norway.

Understanding how management strategies and fishing practices impact lobster size structure

and spawning potential could inform management to improve resiliency to climate-induced

changes. In the Gulf of Maine fishery, egg-bearing (ovigerous) female lobsters are not only

protected from harvest, but also v-notched which offers additional protection up to several

years. Norway, however, protects egg-bearing females without v-notching. Comparing these

fisheries allows us to test the effect of the different management practices and how they interact

with key vital rates, including growth and natural mortality rates. We used deterministic size- and

age-structured models and empirically estimated growth and molt functions to simulate relative

changes in abundance, size structure, egg production, and sex ratios in response to these two

female protection strategies. Our findings suggest that in all scenarios, controlling total fishing

effort to low or moderate levels - relative to the $F > 1$ that has been estimated for American

lobster - is most important for the effectiveness of size-based restrictions on harvest of larger

individuals. Both forms of female protection enhance overall egg production in both species

across levels of fishing intensity, but also result in a skewed sex ratio in favor of females and a

more pronounced size disparity between female and male lobsters. Moreover, our results

suggest that American and European lobster populations exhibit differential responses to the

management strategies, likely due to variations in estimates of natural mortality rates and

growth rates. Our results highlight the sensitivity of management effectiveness to assumptions

regarding the underlying biology, but also provide a clear message that current intense fishing

practices have likely depleted the ability of both species to compensate for fishing mortality in

the long term.

1. INTRODUCTION

Practices to improve the sustainable management of marine fisheries include regulations

for the size and sex of legal catch. While the intention of these regulations is often to protect

older mature females, they can also have unintended consequences for various aspects of

population dynamics, including size structure, sex ratio, and mating behavior (Hines *et al.* 2003;

Rowe and Hutchings 2003; Sato *et al.* 2005; Baker *et al.* 2022). These downstream effects can

impact recruitment, ultimately either supporting or undermining population productivity.

Consequently, discerning how populations respond demographically to selective harvesting

remains essential for forecasting sustainability and designing optimal management strategies

(Rochet and Marty 2016). Crustacean fisheries worldwide use a diverse range of management

regulations, in part due to the distinctive characteristics of lobsters and crabs, which allow non-

invasive sex identification and because females carry their developing eggs externally. For

instance, lobster fisheries often implement bans on the harvesting of females within the

harvestable size range if the female is carrying eggs, with the goal of enhancing overall egg

production as a countermeasure against overfishing.

American lobsters (*Homarus americanus*) are widely distributed in the northwestern

Atlantic Ocean from Newfoundland, Canada to North Carolina, USA; however most are found

north of Rhode Island, USA (Caputi *et al.* 2013). European lobsters (*Homarus gammarus*) are

found in the northeastern Atlantic from the Norwegian Arctic Circle to as far south as Morocco

and parts of the Mediterranean (Wahle *et al.* 2013). American and European lobsters are close

relatives, and share many biological similarities, but the fisheries for each species are managed

differently to account for fishery scale and local cultural values. Contrasting the American

lobster fishery in the U.S. Gulf of Maine to the European lobster fishery in southern Norway

provides an opportunity to more closely examine management strategies' effects on

demography. This includes exploring how life history traits like size at maturity or natural

mortality influence the consequences of management on sex-specific size structure, sex ratio,

and annual egg production.

In the U. S. Gulf of Maine American lobster fishery, the harvesting of egg-bearing

females is prohibited. Fishers are also required to make a V-shaped incision in the egg-bearing

females' tail, a practice known as "v-notching," before releasing them back into the water (Code

of Federal Regulations 2020); this protects the female from future harvest even while not

carrying eggs. In the nearshore Gulf of Maine, where the majority of US landings originate, v-

notching affords protection for up to four years (see Mazur *et al.* 2019). V-notching is not a

common practice in Europe, and it is mainly found in Irish fisheries (Department of the Marine,

Dublin 1994). In the Norwegian fishery for European lobster, harvesting of egg-bearing females

is also prohibited, although it does not involve a v-notching requirement (Kleiven *et al.* 2011).

Since the Norwegian fishery is open only for a two-month-long season (October and

November), we assume this regulation provides only a one-year protection period for the egg-

bearing females.

Protection of mature females coupled with elevated fishing pressure on males could

affect adult sex ratios and sex-specific size structure. Past work on clawed lobsters suggests

that differential fishing selectivity for males and females may lead to suboptimal mating

conditions including skewed sex ratios (Daniel *et al.* 1989; Pugh *et al.* 2013; Jury *et al.* 2019),

reduced mating success (Pugh 2014; Goldstein *et al.* 2014), and sperm limitation (Gosselin *et*

al. 2005). It may also affect the prevalence of sexually-selected traits like claw size in males

(Sørdalen *et al.* 2020). American lobster sex ratios (female:male) in coastal Maine have

reportedly ranged from 1:1 in survey work to 1.4:1 in commercial traps in the 1970s and 1980s

(Krouse 1973; Daniel *et al*. 1989). More recently, work in northern Massachusetts documented

roughly equal sex ratios in smaller lobsters from commercial traps, but above harvestable sizes,

the majority (60-90%) were female (Pugh *et al.* 2013). Differential protection of females,

resulting in intense fishing pressure on males, has been suggested as at least partially

responsible for female-skewed sex ratios in these studies as well as in other portions of the U.S.

fishery (Jury *et al*. 2019).

The aim of this study is to identify how fishing intensity and types of female protections

affect sex ratio, size structure, size disparity between females and males, and egg production

for both American and European lobsters. We use demographic models of population dynamics

to incorporate life history traits to predict population-level trends in size structure and

productivity across a range of fishing pressures and management scenarios. Comparing both

American and European lobsters in this study provides an opportunity to examine how two

forms of restrictions on female harvest (a ban on harvest of egg-bearing females and v-

notching, corresponding to one or four years of protection), interact with species life histories to

impact population demographics.

2. METHODS

2.1 Model Overview

For both American and European lobster, we constructed a deterministic sex-specific

population dynamics model using the open-source statistical software R version 4.0.2 (R Core

Team 2020). This model was based on the methods for a deterministic model of age-structured

population dynamics with annual time steps developed in Mangel (2006) and Kindsvater *et al.*

(2020). We extended the model to incorporate the unique growth patterns of clawed lobsters in

order to estimate average size at age, allowing us to address the size-selectivity of each fishery

(Table 1), including variations on female lobster protections with slot limits (minimum and

maximum sizes) on harvest. This approach made it possible to compare relative trends and

changes in size structure, sex ratio, and total egg production for different scenarios of female

protection in the American and European lobster populations. These attributes were chosen

because they are quantifiable traits that are important for reproductive success. Furthermore,

our approach provides an opportunity to examine relative trends in egg production and size

structure for a range of fishing intensities, without the need for indices of abundance or landings

data. The model was run for three scenarios of differing female protection (none, 1-year, 4-

year), each at varying levels of fishing intensity (F) (see Figure 1). We included the southern

Norwegian practice of banning harvest of ovigerous females which results in protection from

harvest for one year (1-year protection scenario in the model), as well as the Gulf of Maine

fishery requirement of v-notching egg-bearing females, assuming it protects them for four years,

even if they are not carrying eggs (4-year protection scenario in the model). By comparing

management strategies across a range of fishing pressures, and accounting for the unique

patterns of growth and natural mortality that have been estimated for each species, we can

predict the relative impacts of female protection on the size structure of the adult population, as

well as resulting size disparities between the sexes for both clawed lobster fisheries.

2.2 Growth

Lobsters and other crustaceans have a discontinuous growth pattern where they

undergo ecdysis, or molting. Lobsters molt more frequently when they are smaller and growing

fast, and less frequently as they get older and larger (Phillips *et al.* 1980; Huntsberger *et al.*

2020). While recent advances have been made in aging crustaceans, the loss at each molt of

the hard body parts that may track age makes it difficult to accurately assign age (Kilada *et al.*

2012; Wahle *et al.* 2013; Huntsberger *et al.* 2020; Fairfield *et al.* 2021). Thus lobster size-at-age

models are based on data for molting increments (the difference between pre- and post-molt

size) and the duration of time between molts (the intermolt period), which are typically collected

from mark-recapture studies of wild individuals or laboratory growth experiments (Ennis 1978;

Krouse 1981; Campbell 1983; Lawton *et al.* 1984; ASMFC 2020; Sørdalen *et al.* 2022).

Our estimates of size-at-age were based on previously compiled data for annual size-

based molt frequency and molt increments gained from mark-recapture studies of both

American (ASMFC 2006) and European lobster (Sørdalen *et al.* 2022). This allowed for a

projection of the mean probabilistic relationship between size and age for each species. The

model used estimates of mean annual growth to calculate egg production based on the mean

size of spawning females. It also calculated mean change in size for males and females across

scenarios. The growth equations for European lobster were parameterized using data from a

tag-recapture survey from both protected areas and fishing areas in southern Norway (Sørdalen

et al. 2022). Data on molt frequency for the largest individuals was unfortunately limited for both

species.

Annual growth in terms of carapace length was the product of annual molting probability

and molt increment. The annual probability of molting P_{molt} and the molt increment MI_g were

calculated as a function of sex and pre-molt carapace length *CL*. Due to differences in the

reporting practices for each fishery for each species (total length vs. carapace length), we

converted total length to carapace length for European lobster, using the equations in Table 2.

The frequency of molting and the increment of growth vary with individual size, temperature and

between fished and unfished populations, which likely differ in density. Unfortunately, sufficient

data to characterize these potential differences are not available.

For American lobster, annual probability of molting was given by a logistic equation, and

molt increment was a linear function until it reaches an inflection point, after which molt

increments were constant (Table 2). Note that *f* represented an inflection point where the length

at which 10% of individuals were mature and presumably beginning to allocate resources to

reproduction instead of growth (ASMFC 2009, 2020). Annual probability of molting was then

multiplied by the molt increment for each size class to calculate the mean annual growth for a

given sex and pre-molt size. Molt increment for European lobster females and males were linear

functions of pre-molt carapace length (Table 2). The relationship between molting probability

and pre-molt carapace length for European lobster was predicted from the relationship in

Sørdalen *et al.* (2022) (reported in Supplemental Table 1). Therefore, for all growth calculations

for European lobster, we combined the data on molt increment and molt frequency both in the

protected area and the fished area to ensure our length-molt relationship estimates represented

the widest range of size classes available.

2.3 Maximum Age

We assumed that individuals of both species do not live beyond 40 years of age.

Historically, American lobsters have been observed to be as large as 217 mm carapace length

or as much as 19 kg in weight, suggesting they can live for more than four decades (Lawton and

Lavalli 1995). However, individuals of these larger sizes are extremely rare and difficult to

capture (Cooper and Uzmann 1980; Sheehy *et al.* 1999). We used a maximum age of 40 years,

given assumed rates of natural mortality and high fishing mortality in both American and

Norwegian fisheries (ASMFC 2020; Fernández-Chacón *et al.* 2021).

2.4 Reproduction

Size-at-maturity and growth rate are dependent on sex in clawed lobsters. For both

species, the cumulative probability a female was mature at a given carapace length was

represented by logistic equations with parameters for size at 50% maturity (L_{mat}) and shape

parameters (α and β). Values for each species are reported in Table 2. The functions used for

this relationship, known as the maturation ogive for each species, were taken from the primary

literature and differed slightly for each species. The equation for American lobster (ASMFC

2020) was:

$$p_{mat} = 1 - \frac{1}{1+e^{(\alpha+\beta\times CL)}}$$ (1)

The other parameters are estimated from logistic regressions, which have been reported in

slightly different functional forms. In American lobster, size at 50% maturity has shifted over

time (Waller *et al.* 2019, 2021). Thus the most recent stock-wide estimates (and parameters) for

Gulf of Maine female maturity were extracted from the 2020 American lobster stock assessment

(see Appendix 1 in ASMFC 2020).

Since the size at 50% maturity (L_{mat}) for the European lobster population in Southern

Norway has not been determined, we relied on recent estimates of parameters for the size-

specific probability of maturation obtained from Orkney in Scotland, which is similar in latitude to

southern Norway (Coleman *et al.* 2023). The equation for European lobster (Walker 2005;

Coleman *et al.* 2023) was:

$$p_a = \frac{1}{1+e^{-(\alpha+\beta\times CL)}}$$ (2)

In American lobsters, males tend to reach physiological maturity (the ability to produce sperm

and extrude spermatophores) at smaller sizes than females (Krouse 1973; Pugh *et al.* 2015).

However, it is likely that males only participate in mating activities if they are at least the same

size or larger than the mature females since females typically prefer larger males, and the male

must be able to physically manipulate the female into mating position (Templeman 1933; Waddy

and Aiken 1991; Sørdalen *et al.* 2018). Therefore, the functions used for female size-at-maturity

were also used to approximate male functional maturity.

For both species, egg production in each year was calculated based on the proportion of

mature females in the population in all age groups at a given time, their fecundity-at-size

(estimated using functions in Table 2, with values reported in Supplemental Table 2), and their

size-based spawning frequency. In year *t,* total egg production E_t is calculated as the summed

egg production over all age classes *a*:

$$E_t = \sum_a^{40} N_{a,t,f} \times p_a \times Y_a \times SF_a \tag{3}$$

The total number of mature females in each age class in each year is represented by $N_{a,t,f}$,

where *f* signifies that we are only considering females. The probability a given female of age *a* is

mature is p_a (given Eq. 1 and Eq. 2); this is multiplied by both the fecundity at size or age *Y*

(here, given as a function of age, Y_a) and the spawning frequency for a given age or size (SF_a).

This product is the number of eggs produced by a single female in each age class in a given

year, which is then multiplied by the total number of females in that age class $N_{a,t,f}$. The

functions and parameters used to calculate *Y* and *SF* for each species are derived from

empirical data and are given in Table 2. For American lobsters, we assumed female spawning

frequency was every other year, as females alternate with molting years, until they are large

enough to spawn twice per three years (Waddy and Aiken 1986; Waddy *et al.* 1995; ASMFC

2020). European lobsters tend to spawn every other year regardless of size, with some

variability in the larger size classes (Sørdalen 2019).

The population in the model reaches a stable distribution of individuals given annual

birth and death rates, sex, and average individual growth before fishing begins and the

population reaches a new steady state. Recruitment was deterministic and was assumed to

follow a density-dependent relationship between recruitment and spawning stock biomass. For

both species, there is considerable uncertainty in stock recruitment relationships. for lobster

(Sundelöf et al. 2015). Our approach to modeling stock dynamics is focused on the effects of

fishing on adult age structure, not on recruitment overfishing, so we made the most conservative

decision possible when selecting recruitment functions. For American lobster, a Ricker function

is sometimes supported over a Beverton-Holt relationship (ASMFC 2020). However, preliminary

analyses suggested our results were less sensitive to the choice of a Beverton-Holt recruitment

function (Beverton and Holt 1957). The number of female recruits R in a year is given by

$$R = \frac{cE}{1+d} \qquad\qquad (4)$$

Total egg production E is calculated in Eq. 3. Parameter values (*c* and *d*) were arbitrary

and are given in Table 2; we chose these parameters to have minimal influence on our results

and conclusions after preliminary sensitivity analyses.

For European lobster, a Beverton-Holt stock-recruitment relationship has also been

estimated (Bannister and Addison 1986). That relationship is given by

$$R = \frac{gE}{1+(E/h)} \qquad\qquad (5)$$

Again, specific parameter values (*g* and *h*) are given in Table 2. Individuals from both

species are recruited to the model at 56 mm carapace length, which was assumed to

correspond to roughly five years of age (ASMFC 2020), we only tracked age classes from five to

40, past which chances of survival were assumed to be negligible. Sex ratio at recruitment was

assumed to be constant at 1:1.

2.5 Fishing Dynamics

As individuals recruit to the model, the simulated population will eventually reach an

equilibrium size, depending on species- and sex- specific recruitment, growth, maturity,

reproduction, and mortality rates (Table 2). The population dynamics of each age class over

time are given by

263
$$N_{a,t+1} = \begin{cases} R & if\ a = 5 \\ N_{a,t}S & if\ a > 5 \end{cases}$$
(6)

In Eq. 6, $N_{a,t}$ is the number of individuals of in each age class in year t. Survival (S) to the next

year is calculated according to the natural mortality (M) and, for age/size classes vulnerable to

fishery, fishing mortality (F) (sex-specific values given in Table 2). Both mortality rates are

instantaneous, such that $S = e^{-(M+F)}$. Our simulated population grows from an arbitrary starting

size at $t = 0$ until it reaches the unfished equilibrium, or steady state (Mangel 2006). After 100

years, we simulated size-selective fishing, with pressures ranging from lightly fished populations

experiencing an annual harvest rate of approximately 25% (corresponding to an instantaneous

fishing mortality rate (F) of 0.3) to intensely fished populations experiencing an annual harvest

rate of approximately 60% (corresponding to an F of 0.9). We simulated the fished population

for another 100 years, so that each simulation was for 200 years.

For each species, two types of ovigerous female protections were simulated to reflect

actual conservation practices in the fisheries. A scenario without any female protections was

also simulated for comparison. The first scenario of female protection was a 1-year protection

for caught females, which represents a ban on harvest of egg-bearing females, consistent with

Norwegian fishing practices. The second scenario of protection was a 4-year protection period

representing the v-notch given to captured egg-bearing females in the Gulf of Maine. For both

protection methods in the model, once caught and returned, females were removed from the

fishable population for the relevant duration of time where they grew, reproduced, and survived

until rejoining the fishable population. We assumed 100% fisher compliance with protection

measures for egg-bearing females and size limits. For *H. gammarus*, sex-specific natural

mortality was converted from the mean survival of lobsters in three protected areas in Norway

for large and small individuals (Fernández-Chacón *et al.* 2021). Protected females and

individuals outside of the species-specific slot limit therefore only experienced natural mortality

and did not experience fishing mortality in our simulation.

Both American and European lobster fisheries have experienced high rates of harvest in

the past and present; in the US lobster fishery, past stock assessments estimated F at greater

than 1.0 (an annual harvest rate of 63%) in some years (ASMFC 2006, 2009; Kleiven *et al.*

2022). Simulating results for a wide range of fishing intensities provides insight on how

exploitation interacts with different scenarios of female protection. For each species' fishery,

selectivity of the fishery is defined by a slot limit. Those individuals within the harvestable size

range were subjected to fishing mortality in the model. Slot limits were different for each species

and reflect current management practices (Table 1).

The model predicted the number of individuals of each sex at equilibrium, as well as

their size-at-age with and without fishing. These values were used to quantify population size

structure, size disparity between males and females, sex ratio, and egg production for each

fishing intensity and management scenario. Size structure was calculated by summing the

number of individuals by sex per size/age group for each scenario. Size disparity was calculated

by subtracting the mean size of mature males from the mean size of mature females for each

scenario. Sex ratio, or proportion female, was calculated by dividing the total mature female

population by the total mature population of males and females. We also calculated annual egg

production of all spawning females (year class strength) as the sum of all eggs that are

produced by all spawning females in a given year under each fishing scenario, based on

available fecundity-at-size estimates from the literature (Estrella and Cadrin 1995; Agnalt 2008).

3. RESULTS

3.1 Size structure

For both American and European lobsters, we used sex-specific growth functions to

estimate size structure from age structure. In each species, males tended to grow faster than

females (Figure 2). Although growth is continuous, for plotting purposes, we group individuals in

10 mm carapace length size bins. Males move through these bins more quickly than females of

the same cohort (Figure 3). Differences between the two species, especially in the larger size

classes, can be attributed to differences in the estimated rates of natural mortality and growth of

each sex (Figure 3 A and C). To clearly see how fishing pressure and female protection

measures affect population size structure, we compared two levels of fishing intensity

representing moderate (F = 0.3) and intense (F = 0.9) fishing pressure (Figures 3-5). The

predicted stable age distributions corresponding to all panels in Figures 3-5 can be seen in

Supplemental Figures 1 and 2. Note the "intense" fishing scenario is still less than the level of F

historically estimated for American lobsters (ASMFC 2006). The clearest effect of fishing is, as

expected, reducing population abundance (numbers). Because our model includes strong

assumptions about density-dependent recruitment (for which we have little empirical

information) we are focused on relative trends and changes in demography, rather than

predicting yield. The abundance of mature American lobster decreased by more than 60% with

moderate fishing intensity (F = 0.3) in our model, depending on the female protection scenario.

However, this level of fishing allowed for the persistence of both sexes into the size classes

above the slot and even into the largest sizes of 156-165 mm carapace length (Figure 4).

Under intense fishing (F = 0.9), the abundance of mature lobsters decreased by more than

88%, depending on the female protection scenario, and resulted in an extremely truncated size

distribution for both sexes, with few adults persisting in the population above the slot limit.

Management protecting females increased abundance for larger females under both levels of

fishing, again more so in the 4-year protection scenario than the 1-year protection scenario

(Figure 4).

Trends from our model for European lobster showed that moderate fishing pressure (F =

0.3) reduced abundance for all size classes above the minimum legal size, but relative declines

in abundance were less dramatic than for American lobster, ranging from 36-39% depending on

the scenario of female protection. Moderate fishing allowed for adults of both sexes to survive

long enough to outgrow the slot and reach the protected larger size classes (Figure 5). In

model scenarios with female protections, large females above the slot limit were slightly more

abundant (discussed below). Under intense fishing pressure (F = 0.9), numbers of mature

lobsters decreased by 60-63% across all female protection scenarios. The size structure was

more truncated, especially for males, in the largest harvestable size classes, with very few

individuals of either sex surviving to grow larger than the slot. Even with the 4-year protection, in

our model of European lobster, there were very few females in the size bin just above the slot

limit compared to the unfished population (Figure 5).

3.2 Size Disparity Between Sexes

The size distribution of populations of American lobsters responded differently than

European lobsters to the same scenarios of female protections and range of fishing mortality,

due in part to inherent differences in the sizes of mature individuals evident in the unfished

scenario (Figure 3). This resulted in species differences in the size disparity between sexes

(Figure 6). Without fishing, American lobster mature males were on average 4 mm carapace

length (CL), or about 3%, larger than mature females (Figure 6). Without specific measures

protecting females during fishing, males remained larger than females, by an average of 3 to 7

mm, across fishing levels. With one year of protection, females were up to 5 mm (CL) smaller

than males until a fishing level of 0.6, where males and females were on average the same size.

Above F = 0.6, the mean difference increased so that females were up to 5 mm (CL) larger than

males at F = 0.9. Management measures protecting females for four years led to the most

extreme size disparity, where females were on average 2.5 mm CL larger than males at F = 0.3,

but as much as 8 to 16 mm larger than males at $F \geq 0.5$. In other words, at $F \geq 0.5$ the average

male was 7 to 14% smaller than the average female. This predicted pattern is consistent with

the size disparity observed in the Georges Bank sub-stock of American lobster (ASMFC 2020;

T. Pugh, *personal observation*).

In all scenarios, the predicted average size of European lobster females was larger than

males, because the low survival of males led to few in the largest size classes, even in the

unfished population. Therefore, without fishing, the size difference between European lobster

males and females was predicted to be opposite to American lobster, with mature females on

average 5.5 mm (CL) or 4% larger than males (Figure 6). This size disparity in favor of females

decreased slightly with low to moderate fishing pressure (F = 0.1 – 0.4) in all female protection

scenarios. With F > 0.4, females remained larger than males, particularly so in the 4-year

protection scenario. Only in the scenario without female protection measures at very high fishing

pressure (F > 0.7), did females and males show very little size disparity (less than 2% difference

in CL). The greatest disparity occurred in the 4-year protection scenario with intense fishing

pressure (F > 0.8) where females were up to 4% larger than males (4.5 mm in CL).

3.3 Sex Ratio

For American lobster*,* sex ratios were fairly even (near 1:1) without female protections,

but became more biased toward females as protection measures increased (from 1 to 4 years)

and as fishing intensity increased (Figure 7). In the 4-year protection scenario, the bias of the

sex ratio toward females reached 65% female at very high fishing pressure ($F = 0.9$). For

European lobster, as with the pattern of size disparity, the unfished population was biased

toward females. This female bias in sex ratio persisted across levels of fishing in every female

protection scenario and fishing pressure, with very little difference arising from female protection

scenarios. The model predicted female European lobsters consistently comprised between 56-

70% of the mature population, such that 70% of the mature population was female without

fishing, which is consistent with patterns observed in unfished areas (Fernández-Chacón et al.

2021). At low fishing pressure ($F = 0.1 – 0.3$) the proportion of females in the population for both

the 1-year and 4-year protection scenarios remained very similar to the scenario without female

protections. As fishing intensity increased, the proportion female in the 1-year protection

scenario remained within 0.03% of those in the 4-year protection scenario. At $F ≥ 0.5$, females

without any protections decreased to between 56 – 59% of the total population.

3.4 Egg Production

As egg production is a function of the number of spawning females in the population,

even the lowest fishing pressure had a negative effect on total egg production for both species

(Figure 8). Across all scenarios for both species, increasing fishing intensity reduced egg

production compared to unfished production levels, with the largest declines occurring as F

increased from 0 to around 0.4. Of the two female protection scenarios, the 4-year protection

resulted in the least decline in egg production, especially at the highest levels of fishing

pressure. The effectiveness of the two measures of female protection in increasing egg

production were most evident for moderate to intense levels of fishing, particularly for American

lobsters.

Overall, the egg production in American lobster was more sensitive to changes in fishing

pressure compared to the European lobster, meaning that the *relative* impact of fishing on egg

production was greater in American lobster. While American lobster population numbers were

overall much higher than the European lobster, egg production declined to a larger extent with

fishing in American lobster (Figure 8). With increasing levels of fishing pressure, egg production

declined from 71% to 4% of unfished levels for American lobster. When fishing pressure

exceeded $F = 0.6$ without any female protections, egg production was extremely low. Although

the 1-year female protection improved egg production slightly, the 4-year protection consistently

led to up to 13% higher egg production at all levels of fishing pressure.

In European lobster, the sensitivity of egg production to changes in fishing pressure was

slightly lower compared to American lobster. Across all management scenarios, egg production

with increasing fishing pressure declined from 70% to 12% of unfished production levels. At low

fishing pressure, both forms of female protections had minimal impact on egg production, but

their efficacy in improving annual egg production became more apparent as fishing pressure

approached moderate levels (Figure 8), with protection of females for four years being more

effective than one year of protection.

4. DISCUSSION

The aim of this study was to assess the effects of different management regulations,

particularly two types of protection applied to reproductive females, on population size structure,

sex ratio, and egg production in response to varying levels of fishing intensity for both American

and European lobster fisheries. As expected, fishing led to large declines in population size and

commensurate decreases in egg production, which occurred even with slot limits in place under

intense fishing. Implementing one year of protection for egg-bearing females allowed some

females to reach larger sizes, resulting in improved egg production in both species, particularly

at higher fishing pressure, consistent with other research (Sundelöf *et al*. 2015). The 4-year

protection scenario further enhanced egg production, most notably for American lobsters.

However, particularly for American lobsters, female protections also led to skewed sex ratios

and increased size differences between females and males, which could potentially have

unintended consequences for population productivity if the disparity between males and females

affects fertilization success. Differences in the size distributions of American and European

lobster under different fishing scenarios in Figures 4 and 5 are due to the high rates of both

growth and mortality of males estimated from experimentally protected and fished populations of

European lobster (Sørdalen *et al.* 2022). These differences in life history traits led to less

impactful effects of female protection measures, especially 4-year protection, on our modeled

populations of European lobster.

Effects of fishing on productivity

The nuanced implications of fishing mortality on population size are often overlooked in

discussions surrounding resource management focused on yields. We show that realistically

intense levels of fishing pressure led to a substantial reduction in abundance, and that even low

levels of fishing had serious effects on annual egg production (spawning potential) in both

species. Our results suggest that management will have to drastically reduce F in order to see

signs of recovery. Low population biomass and spawning potential can profoundly affect a

population's ability to recover from overfishing or to maintain resilience to natural or

anthropogenic environmental variation. Depleted populations may reach a critical threshold of

density, where decreased fitness of the remaining individuals may negatively affect population

growth rate, a phenomenon known as depensation (Allee 1938; Hutchings 2014). One

mechanism contributing to depensation is when individuals are unable to reproduce

successfully because they are unable to locate a viable mate (Errington 1940; Gascoigne et al.

2009). Variable reproductive success in the Caribbean spotted spiny lobster (*P. guttatus*) in

isolated patch reefs has been attributed to lack of mating opportunities and subsequently

population-level depensation (Robertson and Butler 2009). Depleted populations also yield less

over longer timescales, resulting in a lose-lose situation both from a perspective of ecosystem

function and from a sustainable management view. An additional factor undermining population

productivity comes from the fact that fisheries consistently remove the largest individuals, which

can inadvertently lead to selection for slower growing individuals, smaller body size, and smaller

size at maturity (Hutchings and Rowe 2008). In other words, lower population sizes not only

diminish yield to the fisheries at present but can also further diminish returns by eroding genetic

variation in growth that may be important for long-term sustainable yield (Conover and Munch

2002). Both clawed lobster species studied here show both temporal and spatial variation in the

size at maturation, which could reflect the evolutionary effects of harvest, in addition to recent

increases in temperature, which is known to lead to earlier maturation (Le Bris et al. 2017;

Waller et al. 2021).

Effects of fishing on population size structure

We were able to link population age structure to population size structure by calculating

the expected growth trajectory of the average individual in a deterministic environment for each

species. We incorporated discontinuous growth by calculating mean size at age given annual

growth probabilities, which opens new possibilities for crustacean modelling. For simplicity, we

did not incorporate environmental variability or density effects on growth, instead relying on

parameters and equations that have been estimated previously through field and lab studies.

Our results illustrate how fishing reduces population abundance, erodes size structure and

spawning potential, and may lead to the evolution of faster growth and smaller sizes at maturity.

These effects could explain the worrisome status of the European lobster populations in

Norway, which have declined by 92% over the past 90 years (Kleiven *et al.* 2022). While

American lobster in the Gulf of Maine is still at very high abundance, the Southern New England

US lobster stock is significantly depleted and may be at or near critical thresholds (ASMFC

2020; Pugh *et al*. 2023).

Removal of large females from the population can negatively affect overall productivity

and population resilience as large females in aquatic environments are not only more fecund,

but can exhibit different spawning behavior to maximize reproductive success (a bet-hedging

strategy) (Hixon *et al*. 2014). Age-truncated populations have also been found to track

environmental fluctuations more closely, affecting the resilience of the populations, and

consequently destabilizing the fisheries (Wright and Trippel 2009; Rouyer *et al*. 2011). This

effect of fishing on population demography is exaggerated at higher intensities of fishing, which

was clearly demonstrated in our results for both species (Figures 4 and 5), even though both

simulated fisheries have a slot limit (minimum and maximum harvest size). Management

measures taken to protect reproductive females are designed to preserve size-based fecundity,

thereby promoting the survival of larger, more fecund females to support overall population

growth. However, our model suggested protecting only females within the harvestable slot

resulted in females surviving longer and thus outgrowing their male counterparts.

The species trends in size-disparity across fishing levels arise (Figure 6) from these

differences in estimates of growth and natural mortality. The disparity in average size of

American lobsters was biased toward larger males, but this relationship reversed with fishing,

especially in scenarios with protection of mature females (Figure 6). For European lobster, in all

model scenarios, on average, females were larger than males (even though males grow much

faster; Figure 2) because of the low survival of males. This disparity initially decreases as fishing

pressure increases, as both large females and males are removed from the population.

However, the female-biased disparity between the sexes increased slightly at higher fishing in

scenarios with female protections (Figure 6). This is somewhat in line with observations from

southern Norway where females on average were larger than males in no-take areas, (where

fishing mortality is close to zero), a size disparity that was slightly amplified in fished areas

(Sørdalen *et al*. 2020, Table 1). However, these trap-based studies have also been found to

under-sample large males, potentially biasing the growth and survival parameters used here

(Helms 2023).

Effects of fishing on the size disparity between sexes

The truncation of the adult size distribution and changes to sex ratios in response to

fishing and female protections (Figure 7) can be expected to influence the mating system of

both American and European lobsters, considering their similar mating behavior. Both depletion

of large individuals and skewed sex ratios in larger size classes can limit the opportunity to find

mates and affect the opportunity for and the strength of sexual selection (Kokko *et al*. 2012).

Female lobsters typically select a male with large body size and large crusher claw, which

indicate dominance status (Talbot and Helluy 1995; Atema and Steinbach 2007; Sørdalen *et al.*

2018). In European lobster, size-selective fishing appears to disrupt these sexually selected

traits, suggesting that fishing can alter mating behavior and weaken sexual selection on male

traits (Sørdalen *et al.* 2018, 2020). Although females may mate with smaller males when faced

with scarcity of choice, experimental studies suggest that smaller males may not provide

sufficient sperm to fertilize the clutch of larger females, potentially leading to sperm limitation

(Gosselin *et al.* 2003; Pugh 2014; Pugh *et al.* 2015; Tang *et al.* 2019). While some evidence

from other exploited crustacean populations suggests that changes in population sex ratio and

size structure may result in sperm limitation (Pardo *et al*. 2015; Baker *et al*. 2022), this

relationship still needs further investigation for clawed lobsters. The results of our model are

sensitive to sex- and size-based mortality, and do not address potential variation in mating

success or fertilization rates resulting from predicted size-disparities and sex-ratios. Further

understanding of these relationships in clawed lobsters is needed to fully understand the

potential for sperm limitation to limit their population productivity beyond direct effects of

reduced egg protection.

Effects of fishing on spawning potential

The spawning potential (egg production) of each species declined dramatically with

fishing intensity, but was affected differently by each measure of female protection (Figure 8). In

this study, European lobsters did not experience the same degree of improvement in egg

production or size disparity as American lobsters with female protections. The model outputs

show that v-notching as a form of 4-year protection is an effective means of boosting egg

production if the female can survive long enough to benefit from this. Because of the assumed

low natural mortality of American lobster, females were able to survive through the 4 years of

protection to continue growing and reproducing. In species with higher estimates of natural

mortality rates, including the European lobster, v-notching (represented here as 4-year

protection) is predicted to only marginally enhance egg producing potential compared to a ban

on harvest of egg-bearing females alone (Sundelöf et al. 2015). Future studies could test the

hypothesis that European lobster would benefit from a narrower slot limit (a lower maximum

legal size) to improve the chances both sexes outgrow the size classes vulnerable to harvest.

However, even with female protections, our results show that fishing at high intensities may

lower spawning potential enough to reduce the overall population's resilience to fishing or poor

environmental conditions. Intensively fishing a population to a spawning potential ratio of less

than 0.4 has been suggested to be a critical threshold past which populations do not have a

sufficient spawning potential to buffer natural variability in recruitment (Wiedenmann et al. 2006;

Zhou et al. 2020; Conrad and Kindsvater 2024). In Figure 8, our model predicts this threshold

occurs at low to moderate levels of fishing, depending on the species and female protection

scenario.

Survival and growth rates

The literature-based estimates of survival and growth rates that we used contribute to

our results, including the effects of slot limits and protections such as v-notching on size

structure and egg production, because these rates determine the time an individual spends in

the size classes vulnerable to fishing and the probability of surviving (Chong-Montenegro and

Kindsvater 2022). We based survival on mortality estimates for European lobster calculated

from a trap-based mark-recapture study (Fernández-Chacón *et al*. 2021), and like any study

conducted in this manner, estimates are dependent on season, sex and size-dependent

catchability, trap design (Miller 1990), and assumptions around immigration/emigration (Skerritt

et al. 2023). By contrast, the natural mortality parameter used for American lobster was the

value used by recent stock assessments for Gulf of Maine lobsters; it does not differentiate by

sex and is not empirically derived (ASMFC 2020). If this parameter is an underestimate of the

natural mortality of American lobster, then we may be overestimating the effectiveness of v-

notching and results may be more similar to the modeled European lobster results currently

shown (and vice versa). Future studies to simulate both possibilities would clarify this

discrepancy, and improved empirical data on natural mortality is clearly desirable.

The growth parameters used in this study for European lobster are based on recent tag-

recapture data that estimated molting probability and molt increment (Sørdalen *et al.* 2022). We

combined data from protected and fished areas to maximize the size range used in the

parameter estimates. This choice may have impacted some of the modeled differences between

male and female growth since lobsters from fished areas do not grow as fast as those subjected

to protection, an effect particularly evident in females (Sørdalen et al. 2022). The plateauing of

male growth (see Figure 2) is likely due to the combined effect of insufficient empirical data to

estimate molt parameters at the larger sizes and the relatively high natural mortality estimate

used for males. The growth parameters for American lobster are mainly from older data that

have been recycled over a number of stock assessments (ASMFC 2020). This could be an

indication that updated estimates of natural mortality, growth, and fecundity are necessary to

accurately represent American lobster population dynamics into the future, especially since it

has already been shown that their size-at-maturity has been changing over time (Waller *et al.*

2019).

The comparative nature of our work demonstrates the importance of the estimated

natural mortality and other life history parameters to modeled results, and shows that accurate

knowledge of species life histories is needed for improved modelling and for truly understanding

the effect of fishing on population demography and dynamics. Yet intense fishing activities in the

past decades have left few areas unaffected by fishing, thereby limiting the availability of

reference areas for collecting data on life history parameters and natural behavior. No-take

areas have been adopted as management tool in recent years, with 60 such areas already

implemented across southern Norway, aimed at aiding the recovery of local populations of

European lobster (Moland *et al*. 2021; Knutsen *et al*. 2022). Remarkably, these Norwegian

protected areas have demonstrated that they need not be large to elicit population-level

responses, likely in part due to the coastal fjord landscape and the high site fidelity of individual

European lobsters (Moland et al. 2021). The observed changes in demographic structures,

phenotypic composition, mating behavior, and reproductive output within reserves in Norway

and other locations have provided valuable insights into the effects of fishing and the impact of

release from fishing pressure (Bertelsen and Matthews 2001; Goni *et al*. 2003; Sørdalen *et al*.

2018, 2020, 2022; Fernández-Chacón *et al*. 2021). Our simulation results clearly showed

differences in the demographics between the unfished population (comparable to a reserve),

very low levels of F, and higher levels of fishing intensity. Whether no-take areas as

implemented are supporting overall population growth and the fishery through spillover is yet to

be determined.

5. CONCLUSION

The sensitivity of overall population size and size structure to fishing pressure as shown in our

results for both species emphasizes the need for implementing targeted measures of controlling

fishing mortality, to prevent further depletion and facilitate recovery of lobster populations. Such

measures could range from narrowing the harvest window of minimum and maximum size

limits, or more directly controlling F by implementing catch quotas. Understanding how

management decisions influence population structure is crucial, particularly in light of the

predicted decline in American lobster recruitment and that there are still no signs of recovery for

European lobster (Sundelöf et al. 2015; Le Bris *et al*. 2018; Oppenheim *et al*. 2019; Kleiven *et*

al. 2022). As managers explore options for addressing the challenges faced by these two

species, the development and refinement of models like the one we have presented here can

serve as critical tools for evaluating management strategies and forecasting possible outcomes.

Author Contributions:

Conceptualization: K.T., T.K.S., T.L.P, H.K.K., Methodology and Software: K.T., H.K.K.,

Writing original draft: K.T., Reviewing and Editing: K.T., T.K.S., T.L.P., H.K.K.

Acknowledgements:

K.T. and H.K.K. were supported by the Department of Fish and Wildlife Conservation at Virginia

Tech. This study was completed as part of a K.T.'s Master of Science thesis, and she would like

to thank her committee, particularly Leandro Castello. We would also like to thank Kim

Halvorsen for assistance with growth data and for insightful conversations about *H. gammarus*.

K.T. also thanks Hailey Conrad for assistance troubleshooting R code.

REFERENCES

Agnalt, A.-L. 2008. Fecundity of the European lobster (*Homarus gammarus*) off southwestern Norway after stock enhancement: do cultured females produce as many eggs as wild females? ICES Journal of Marine Science 65(2):164–170.

Alden, R. 2023, November 22. FISHERIES FROM 1973 TO 2023 — How We Got Here. https://www.islandinstitute.org/2023/11/22/fisheries-from-1973-to-2023-how-we-got-here/.

Allee, W. C. 1938. The social life of animals, by W.C. Allee ... W.W. Norton & Company, inc., New York,.

ASMFC. 2006. Stock Assessment Report No. 06-03 of the Atlantic States Marine Fisheries Commission American Lobster Stock Assessment for Peer Review. Atlantic States Marine Fisheries Commission.

ASMFC. 2009. 2009 American Lobster Stock Assessment Report for Peer Review. Atlantic States Marine Fisheries Commission.

ASMFC. 2020. 2020 American Lobster Benchmark Stock Assessment and Peer Review Report. Atlantic States Marine Fisheries Commission, NA15NMF4740069.

Atema, J., and M. A. Steinbach. 2007. Chemical Communication and Social Behavior of the Lobster *Homarus americanus* and Other Decapod Crustacea. Pages 115–144 *in* J. E. Duffy and M. Thiel, editors. Evolutionary Ecology of Social and Sexual Systems, 1st edition. Oxford University PressNew York.

Atema, J., and R. Voigt. 1995. Behavior and Sensory Biology. Pages 313–348 Biology of the Lobster. Elsevier.

Baker, K. D., D. R. J. Mullowney, and B. Sainte-Marie. 2022. Large males matter: Low sperm reserves in female snow crab (*Chionoecetes opilio*) off Newfoundland, Canada. Fisheries Research 253:106385.

Bannister, R. C. A., and J. T. Addison. 1986. Effect of Assumptions about the Stock–Recruitment Relationship on a Lobster (*Homarus gammarus*) Stock Assessment. Canadian Journal of Fisheries and Aquatic Sciences 43(11):2353–2359.

Bertelsen, R. D., and T. R. Matthews. 2001. Fecundity dynamics of female spiny lobster (*Panulirus argus*) in a south Florida fishery and Dry Tortugas National Park lobster sanctuary. Marine and Freshwater Research 52(8):1559.

Beverton, R. J. H., and S. J. Holt. 1957. On the dynamics of exploited fish populations. Fish. Invest. Ser., UK.

Campbell, A. 1983. Growth of Tagged American Lobsters, *Homarus americanus*, in the Bay of Fundy. Canadian Journal of Fisheries and Aquatic Sciences 40(10):1667–1675.

Caputi, N., S. de Lestang, S. Frusher, and R. A. Wahle. 2013. The impact of climate change on exploited lobster stocks. Pages 84–112 Lobsters: Biology, Management, Aquaculture & Fisheries: Second Edition.

Chong-Montenegro, C., and H. K. Kindsvater. 2022. Demographic Consequences of Small-Scale Fisheries for Two Sex-Changing Groupers of the Tropical Eastern Pacific. Frontiers in Ecology and Evolution 10:850006.

Code of Federal Regulations. 2020. 50 CFR Part 697 Size, harvesting and landing requirements. Pages 1077–1126.

Coleman, M. T., M. Garratt, N. Hold, I. S. M. Bloor, S. R. Jenkins, J. S. Porter, O. Tully, and M. C. Bell. 2023. A standardized assessment of geographic variation in size at maturity of European lobster (*Homarus gammarus* L.) in the North East Atlantic. ICES Journal of Marine Science 80(4):911–922.

Conover, D. O., and S. B. Munch. 2002. Sustaining Fisheries Yields Over Evolutionary Time Scales. Science 297(5578):94–96.

Conrad, H. M., and H. K. Kindsvater. 2024. Modeling potential consequences of climate-driven sex
reversal on Southern and Summer Flounder population dynamics. Marine and Coastal Fisheries
16(2):e10289.
Cooper, R. A., and J. R. Uzmann. 1980. Ecology of juvenile and adult Homarus. Page 97-141 *in* J. S. Cobb
and R. Phillips, editors. The biology and management of lobster. Academic Press, New York.
Cowan, D. F., and J. Atema. 1990. Moult staggering and serial monogamy in American lobsters, *Homarus*
americanus. Animal Behaviour 39(6):1199–1206.
Daniel, P. C., R. C. Bayer, and C. Waltz. 1989. Egg Production of V-Notched American Lobsters (*Homarus*
americanus) along Coastal Maine. Journal of Crustacean Biology 9(1):77–82.
Department of the Marine, Dublin. 1994. Lobster (Conservation of Stocks) Order. Page S.I. No. 304 of
1994.
Ennis, G. P. 1978. Growth curves for Newfoundland lobsters from data on molt increment and
proportion molting. Fisheries and Marine Service, Fisheries and Environment Canada, St. John's,
Newfoundland.
Errington, P. L. 1940. Natural Restocking of Muskrat-Vacant Habitats. The Journal of Wildlife
Management 4(2):173.
Estrella, B. T., and S. X. Cadrin. 1995. Fecundity of the American lobster (*Homarus americanus*) in
Massachusetts coastal waters. ICES Marine Science Symposium 199:61–72.
Fairfield, E. A., D. S. Richardson, C. L. Daniels, C. L. Butler, E. Bell, and M. I. Taylor. 2021. Ageing European
lobsters (*Homarus gammarus*) using DNA methylation of evolutionarily conserved ribosomal
DNA. Evolutionary Applications 14(9):2305–2318.
Fernández-Chacón, A., L. Buttay, E. Moland, H. Knutsen, and E. M. Olsen. 2021. Demographic responses
to protection from harvesting in a long-lived marine species. Biological Conservation
257:109094.
Gascoigne, J., L. Berec, S. Gregory, and F. Courchamp. 2009. Dangerously few liaisons: a review of mate-
finding Allee effects. Population Ecology 51(3):355–372.
Goldstein, J. S., T. L. Pugh, E. A. Dubofsky, K. L. Lavalli, M. Clancy, and W. H. Watson. 2014. A Noninvasive
Method For In situ Determination of Mating Success in Female American Lobsters (*Homarus*
americanus). Journal of Visualized Experiments (84):50498.
Goni, R., A. Quetglas, and O. Renones. 2003. Size at maturity, fecundity and reproductive potential of a
protected population of the spiny lobster *Palinurus elephas* (Fabricius, 1787) from the western
Mediterranean. Marine Biology 143(3):583–592.
Gosselin, T., B. Sainte-Marie, and L. Bernatchez. 2003. Patterns of sexual cohabitation and female
ejaculate storage in the American lobster (*Homarus americanus*). Behavioral Ecology and
Sociobiology 55(2):151–160.
Gosselin, T., B. Sainte-Marie, and L. Bernatchez. 2005. Geographic variation of multiple paternity in the
American lobster, *Homarus americanus*. Molecular Ecology 14(5):1517–1525.
Helms, I. M. 2023. Size selectivity of standardized lobster traps used in long-term monitoring and their
ability to catch large European lobsters (*Homarus gammarus*). University of Agdar, Kristiansand,
Norway.
Hines, A. H., P. R. Jivoff, P. J. Bushmann, J. van Montfrans, S. A. Reed, D. L. Wolcott, and T. G. Wolcott.
2003. Evidence for Sperm Limitation in the Blue Crab,. Bulletin of Marine Science 72(2):24.
Hixon, M. A., D. W. Johnson, and S. M. Sogard. 2014. BOFFFFs: on the importance of conserving old-
growth age structure in fishery populations. ICES Journal of Marine Science 71(8):2171–2185.
Huntsberger, C. J., R. Kilada, W. G. Ambrose, and R. A. Wahle. 2020. Age-at-size relationships of the
American lobster (*Homarus americanus*) from three contrasting thermal regimes using gastric
mill band counts as a direct aging technique. Canadian Journal of Fisheries and Aquatic Sciences
77(10):1733–1740.

Hutchings, J. A. 2014. Renaissance of a caveat: Allee effects in marine fish. ICES Journal of Marine Science 71(8):2152–2157.

Hutchings, J. A., and S. Rowe. 2008. Consequences of sexual selection for fisheries-induced evolution: an exploratory analysis: Consequences of sexual selection for fisheries-induced evolution. Evolutionary Applications 1(1):129–136.

Jury, S., T. Pugh, H. Henninger, J. Carloni, and W. Watson. 2019. Patterns and possible causes of skewed sex ratios in American lobster (*Homarus americanus*) populations. Invertebrate Reproduction & Development 63(3):189–199.

Kilada, R., B. Sainte-Marie, R. Rochette, N. Davis, C. Vanier, and S. Campana. 2012. Direct determination of age in shrimps, crabs, and lobsters. Canadian Journal of Fisheries and Aquatic Sciences 69(11):1728–1733.

Kindsvater, H. K., K. T. Halvorsen, T. K. Sørdalen, and S. H. Alonzo. 2020. The consequences of size-selective fishing mortality for larval production and sustainable yield in species with obligate male care. Fish and Fisheries 21(6):1135–1149.

Kleiven, A. R., S. H. Espeland, S. Stiansen, K. Ono, F. Zimmermann, and E. M. Olsen. 2022. Technological creep masks continued decline in a lobster (*Homarus gammarus*) fishery over a century. Scientific Reports 12(1):3318.

Kleiven, A. R., E. M. Olsen, and J. H. Vølstad. 2011. Estimating Recreational and Commercial Fishing Effort for European Lobster *Homarus gammarus* by Strip Transect Sampling. Marine and Coastal Fisheries 3(1):383–393.

Knutsen, J. A., A. R. Kleiven, E. M. Olsen, H. Knutsen, S. H. Espeland, T. K. Sørdalen, S. H. Thorbjørnsen, J. A. Hutchings, A. Fernández-Chacón, M. Huserbråten, D. Villegas-Ríos, K. T. Halvorsen, P. J. Nillos Kleiven, T. K. Langeland, and E. Moland. 2022. Lobster reserves as a management tool in coastal waters: Two decades of experience in Norway. Marine Policy 136:104908.

Kokko, H., H. Klug, and M. D. Jennions. 2012. Unifying cornerstones of sexual selection: operational sex ratio, Bateman gradient and the scope for competitive investment. Ecology Letters 15(11):1340–1351.

Krouse, J. S. 1973. Maturity, Sex Ratio, and Size Composition of the Natural Population of American lobster *Homarus americanus*, along the Maine Coast. FISHERY BULLETIN 71:9.

Krouse, J. S. 1981. Movement, Growth, and Mortality of American Lobsters, *Homarus americanus*, Tagged Along the Coast of Maine. Page 17. U.S. Department of Commerce; National Oceanic and Atmospheric Administration; National Marine Fisheries Service, NOAA Technical Report NMFS SSRF-747.

Lawton, P., and K. L. Lavalli. 1995. Postlarval, Juvenile, Adolescent, and Adult Ecology. Pages 47–88 *in* J. R. Factor, editor. Biology of the lobster *Homarus americanus*. Academic Press, San Diego.

Lawton, R. P., P. Brady, C. Sheehan, W. Sides, E. Kouloheras, M. Borgatti, and V. Malkoski. 1984. Growth and Movement of Tagged Lobsters released in Western Cape Cod Bay, 1970-1977. Pages 119–129 *in* J. D. Davis and D. Merriman, editors. Observations on the Ecology and Biology of Western Cape Cod Bay, Massachusetts. Springer US, New York, NY.

Le Bris, A., K. E. Mills, R. A. Wahle, Y. Chen, M. A. Alexander, A. J. Allyn, J. G. Schuetz, J. D. Scott, and A. J. Pershing. 2018. Climate vulnerability and resilience in the most valuable North American fishery. Proceedings of the National Academy of Sciences of the United States of America 115(8):1831–1836.

Le Bris, A., A. J. Pershing, J. Gaudette, T. L. Pugh, and K. M. Reardon. 2017. Multi-scale quantification of the effects of temperature on size at maturity in the American lobster (*Homarus americanus*). Fisheries Research 186:397–406.

Mangel, M. 2006. The Theoretical Biologist's Toolbox: Quantitative Methods for Ecology and Evolutionary Biology, 1st edition. Cambridge University Press.

774 Mazur, M., B. Li, J. Chang, and Y. Chen. 2019. Contributions of a conservation measure that protects the
775  spawning stock to drastic increases in the Gulf of Maine American lobster fishery. Marine
776  Ecology Progress Series 631:127–139.
777 Miller, R. J. 1990. Effectiveness of Crab and Lobster Traps. Canadian Journal of Fisheries and Aquatic
778  Sciences 47(6):1228–1251.
779 Moland, E., A. Fernández-Chacón, T. K. Sørdalen, D. Villegas-Ríos, S. H. Thorbjørnsen, K. T. Halvorsen, M.
780  Huserbråten, E. M. Olsen, P. J. Nillos Kleiven, A. R. Kleiven, H. Knutsen, S. H. Espeland, C. Freitas,
781  and J. A. Knutsen. 2021. Restoration of Abundance and Dynamics of Coastal Fish and Lobster
782  Within Northern Marine Protected Areas Across Two Decades. Frontiers in Marine Science
783  8:674756.
784 Oppenheim, N. G., R. A. Wahle, D. C. Brady, A. G. Goode, and A. J. Pershing. 2019. The cresting wave:
785  larval settlement and ocean temperatures predict change in the American lobster harvest.
786  Ecological Applications 29(8).
787 Pardo, L. M., Y. Rosas, J. P. Fuentes, M. P. Riveros, and O. R. Chaparro. 2015. Fishery Induces Sperm
788  Depletion and Reduction in Male Reproductive Potential for Crab Species under Male-Biased
789  Harvest Strategy. PLOS ONE 10(3):e0115525.
790 Phillips, B. F., J. S. Cobb, and R. W. George. 1980. General Biology. Pages 1–82 *in* J. S. Cobb and B. F.
791  Phillips, editors. The Biology and management of lobsters. Academic Press, New York.
792 Pugh, T., R. Glenn, and W. Watson. 2023. Spatial differences in female lobster mating success and
793  population characteristics in southern Massachusetts. Marine Ecology Progress Series 718:69–
794  84.
795 Pugh, T. L. 2014. The Potential for Sperm Limitation in American Lobsters (*Homarus americanus*) as
796  Indicated by Female Mating Activity and Male Reproductive Capacity. University of New
797  Hampshire.
798 Pugh, T. L., M. Comeau, W. H. Watson, and K. Benhalima. 2015. Variation in the size and composition of
799  ejaculates produced by male American lobsters, *Homarus americanus*. Milne Edwards, 1837
800  (Decapoda: Nephropidae). Journal of Crustacean Biology 35(5):593–604.
801 Pugh, T. L., J. S. Goldstein, K. L. Lavalli, M. Clancy, and W. H. Watson. 2013. At-sea determination of
802  female American lobster (*Homarus americanus*) mating activity: Patterns vs. expectations.
803  Fisheries Research 147:327–337.
804 R Core Team. 2020. R: A language and environment for   statistical computing. R Foundation for
805  Statistical Computing, Vienna, Austria.
806 Robertson, D. N., and M. J. Butler. 2009. Variable reproductive success in fragmented populations.
807  Journal of Experimental Marine Biology and Ecology 377(2):84–92.
808 Rochet, M., and L. Marty. 2016. Effects of Fishing on the Population. Pages 188–225 *in* T. Jakobsen, M. J.
809  Fogarty, B. A. Megrey, and E. Moksness, editors. Fish Reproductive Biology, 1st edition. Wiley.
810 Rouyer, T., G. Ottersen, J. M. Durant, M. Hidalgo, D. Ø. Hjermann, J. Persson, L. C. Stige, and N. C.
811  Stenseth. 2011. Shifting dynamic forces in fish stock fluctuations triggered by age truncation?
812  Global Change Biology 17(10):3046–3057.
813 Rowe, S., and J. A. Hutchings. 2003. Mating systems and the conservation of commercially exploited
814  marine fish. Trends in Ecology & Evolution 18(11):567–572.
815 Sato, T., S. Goshima, and M. Ashidate. 2005. Negative Effects of Delayed Mating on the Reproductive
816  Success of Female Spiny King Crab, *Paralithodes Brevipes*. Journal of Crustacean Biology
817  25(1):105–109.
818 Sheehy, M., R. Bannister, J. F. Wickins, and P. Shelton. 1999. New perspectives on the growth and
819  longevity of the European lobster (*Homarus gammarus*). Canadian Journal of Fisheries and
820  Aquatic Sciences 56(10):1904–1915.

Skerritt, D., M. Bell, K. Lees, A. Mill, N. Polunin, and C. Fitzsimmons. 2023. Estimating catchability and
density of the European lobster *Homarus gammarus* from continuous, short-term mark-
recapture data. Marine Ecology Progress Series 715:79–89.
Sørdalen, T. K. 2019. Marine reserves and selective fishing shape mating behaviour, secondary sexual
trait and growth in European lobster. PhD Thesis, Unpublished.
Sørdalen, T. K., K. T. Halvorsen, H. B. Harrison, C. D. Ellis, L. A. Vøllestad, H. Knutsen, E. Moland, and E. M.
Olsen. 2018. Harvesting changes mating behaviour in European lobster. Evolutionary
Applications 11(6):963–977.
Sørdalen, T. K., K. T. Halvorsen, and E. M. Olsen. 2022. Protection from fishing improves body growth of
an exploited species. Proceedings of the Royal Society B: Biological Sciences
289(1987):20221718.
Sørdalen, T. K., K. T. Halvorsen, L. A. Vøllestad, E. Moland, and E. M. Olsen. 2020. Marine protected areas
rescue a sexually selected trait in European lobster. Evolutionary Applications 13(9):2222–2233.
Sundelöf, A., Grimm, V., Ulmestrand, M. and Fiksen, Ø., 2015. Modelling harvesting strategies for the
lobster fishery in northern Europe: the importance of protecting egg-bearing
females. *Population Ecology*, *57*, pp.237-251.
Talbot, P., and S. Helluy. 1995. Reproduction and Embryonic Development. Pages 177–216 *in* J. R. Factor,
editor. Biology of the lobster *Homarus americanus*. Academic Press, San Diego.
Templeman, W. 1933. Mating in the American Lobster. Contributions to Canadian Biology and Fisheries
8(1):421–432.
Thorbjørnsen, S., E. Moland, M. Huserbråten, J. Knutsen, H. Knutsen, and E. Olsen. 2018. Replicated
marine protected areas (MPAs) support movement of larger, but not more, European lobsters to
neighbouring fished areas. Marine Ecology Progress Series 595:123–133.
Waddy, S. L., and D. E. Aiken. 1986. Multiple Fertilization and Consecutive Spawning in Large American
Lobsters, *Homarus americanus*. Canadian Journal of Fisheries and Aquatic Sciences 43(11):2291–
2294.
Waddy, S. L., and D. E. Aiken. 1991. Mating and Insemination in the American Lobster, *Homarus
americanus*. Pages 126–144 *in* R. T. Bauer and J. W. Martin, editors. Crustacean Sexual Biology.
Columbia University Press, New York.
Waddy, S. L., D. E. Aiken, and D. P. V. De Kleijn. 1995. Control of Growth and Reproduction. Pages 217–
266 Biology of the Lobster. Elsevier.
Wahle, R. A., C. Brown, and K. Hovel. 2013. The Geography and Body-Size Dependence of Top-Down
Forcing in New England's Lobster-Groundfish Interaction. Bulletin of Marine Science 89(1):189–
212.
Walker, T. I. 2005. Reproduction in Fisheries Science. Pages 81–127 *in* W. C. Hamlett, editor.
Reproductive Biology and Phylogeny of Chondrichthyes: Sharks, Batoids and Chimaeras. Science
Publishers, Enfield, NH.
Waller, J. D., K. M. Reardon, S. E. Caron, B. P. Jenner, E. L. Summers, and C. J. Wilson. 2021. A
comparison of the size at maturity of female American lobsters (*Homarus americanus*) over
three decades and across coastal areas of the Gulf of Maine using ovarian staging. ICES Journal
of Marine Science 78(4):1267–1277.
Waller, J. D., K. M. Reardon, S. E. Caron, H. M. Masters, E. L. Summers, and C. J. Wilson. 2019. Decrease
in size at maturity of female American lobsters *Homarus americanus* (H. Milne Edwards, 1837)
(Decapoda: Astacidea: Nephropidae) over a 50-year period in Maine, USA. Journal of Crustacean
Biology 39(4):509–515.
Wiedenmann, J., M. Americas, and M. Mangel. 2006. A Review of Rebuilding Plans for Overfished Stocks
in the United States. Lenfest Ocean Program, Washington, DC.

Wright, P. J., and E. A. Trippel. 2009. Fishery-induced demographic changes in the timing of spawning:
consequences for reproductive success. Fish and Fisheries 10(3):283–304.
Zhou, S., A. E. Punt, Y. Lei, R. A. Deng, and S. D. Hoyle. 2020. Identifying spawner biomass per-recruit
reference points from life-history parameters. Fish and Fisheries 21(4):760–773.
872
873
874

876   Figure 1. Summary of the order of events that the model simulates in each time step.
           Items with an asterisk are variable depending on the scenario.

878   Figure 2. Predicted growth of American (*H. americanus*) and European (*H.
           gammarus*) lobsters over time. Age here is relative since these curves were
879   generated using annual size-based probabilities of molting and molting increment for
           females and males of American (ASMFC 2009, 2020) and European (Sørdalen *et al.*
           2022) lobsters.

Figure 3. Size structure plots of the total population of American (*H. americanus*) in Panel A and European (*H. gammarus*) lobsters in Panel C over time without fishing. Size bins are in millimeters carapace length. Lighter versions of the colors for females and males indicate immature individuals, and darker colors are mature. Panels B and D show only the mature population for American and European lobsters, respectively. The size bins in between the vertical red lines contain the harvestable size limits for reference. The American lobster slot limit is 83-127 mm carapace length. The slot limits for European lobster are 86.9-114.76 mm (females) and 88.6-117.09 mm (males) carapace length.

Figure 4. Size structures of American lobsters with and without female protections during fishing compared to the unfished size structure. Size bins are in millimeters carapace length. Note that these are for mature individuals only at *F* = 0.3 and *F* = 0.9. The first row shows size structures after fishing without any female protections. The second row shows size structures after fishing with 1-year protections for ovigerous females. The third row shows after fishing with 4-year protections for ovigerous females. The size bins in between the vertical red lines contain the harvestable size limits of 83-127 mm carapace length for American lobster.

Figure 5. Size structures of European lobsters with and without female protections during fishing compared to the unfished size structure. Size bins are in millimeters carapace length. Note that these are for mature individuals only at F = 0.3 and F = 0.9. The first row shows size structures after fishing without any female protections. The second row shows size structures after fishing with 1-year protections for ovigerous females. The third row shows after fishing with 4-year protections for ovigerous females. The slot limits for European lobster are 86.9-114.76 mm (female) and 88.6-117.09 mm (males) carapace length.

891 ·

894 Figure 6. Mean size disparity by sex calculated by subtracting mean mature male carapace length from mean mature female carapace length for American (*H. americanus*) and European (*H. gammarus*) lobsters. The red horizontal lines indicate the threshold (zero) below which males are larger than females on average, or above which females are on average larger than males.

899 Figure 7. Proportion female for American (*H. americanus*) and European (*H. gammarus*) lobsters over in three alternative female protection scenarios and varied levels of fishing pressure.

Figure 8. Egg production during fishing scaled to the unfished level of egg production for American (*H. americanus*) and European (*H. gammarus*) lobsters in three alternative female protection

904   scenarios and varied levels of fishing pressure.

907 **Table 1.** Harvest regulations for American (ASMFC 2020) and European lobster fisheries
908 (Kleiven et al. 2011; Sørdalen et al. 2018).

Regulation	Gulf of Maine American lobster (*H. americanus*)	Southern Norway European lobster (*H. gammarus*)
Minimum harvestable size	83 mm CL	**Females**: 86.9 mm CL **Males**: 88.6 mm CL (Converted from 250 mm total length for each sex)
Maximum harvestable size	127 mm CL	**Females**: 114.76 mm CL **Males**: 117.09 mm CL (Converted from 320 mm total length)
Harvest season	January 1 – December 31 (all year)	October 1 – November 30 (Thorbjørnsen *et al.* 2018)
Trap limit	800 traps per fisher, limited to 600 traps in Zone E (Alden 2023)	Commercial fishers may use up to 100 traps per day, whereas recreational fishers may use 10 traps per day
Egg-bearing female protection	V-notching, no harvest of egg-bearing females	No harvest of egg-bearing females

911 **Table 2.** Species-specific relationships used to calculate average growth, maturation, and
912 recruitment. Average growth per year is calculated by converting Total Length to Carapace
913 Length (for European lobster), and then estimating the product of the probability of molting and
914 the molt increment at each size.

Parameter / Equation Name	Gulf of Maine American lobster (*H. americanus*)	Southern Norway European lobster (*H. gammarus*)
Total Length (TL) to Carapace Length (CL) conversion	n/a	Female: $CL = 0.3842 \times TL - 8.45$ Male: $CL = 0.3987 \times TL - 10.42$ (Sørdalen *et al.* 2022)
Probability of molting (P_{molt})	$p_{molt\,a} = \dfrac{1}{1 + e^{(a + b \times CL)}}$ a = Female: -6.571 Male: -6.834 b = Female: 0.05901 Male: 0.06046 (ASMFC 2009, 2020)	*See Supplemental Table 1*
Molt Increment (MI)	$MI_a = \begin{bmatrix} m_a + n_a \times CL & for\ CL < f \\ m_a + n_a f & for\ CL \geq f \end{bmatrix}$ m_a = Female: -3.90 Male: -4.51 n_a = Female: 0.22 Male: 0.23 f = Female: 76 Male: 86 (ASMFC 2009, 2020)	$MI_g = m_g + n_g \times CL$ m_g = Female: 13.31382 Male: 9.2874 n_g = Female: -0.05861 Male: 0.02223 (Sørdalen *et al.* 2022)
Maturation ogive parameters (Eqs 1 and 2)	α = -17.14056 β = 0.19664 (ASMFC 2020)	α = -40.59 β = 0.43 (Coleman *et al.* 2023)
L_{mat}	87 mm (ASMFC 2020)	92.5 mm (Coleman *et al.* 2023)
Spawning frequency (SF)	$for\ L_{mat} > CL > 120\quad freq = 0.5$ $for\ CL > 120\qquad freq = 0.66$ (Waddy and Aiken 1986; Waddy *et al.* 1995; ASMFC 2020)	$freq = 0.50$ (Sørdalen 2019)
Fecundity at size (Y)	$Y = 1.09886(0.000919833[CL^{3.580220}])$ (Estrella and Cadrin 1995)	$Y = 468.17(CL - 33004)$ (Agnalt 2008)
Stock-Recruitment relationship parameters	$c = 1$ $d = 9E\text{-}10$	g = 100,000 recruits per 10^8 eggs h = 0.851 eggs (x10^{-8}) (Bannister and Addison 1986)
Instantaneous Natural Mortality *M*	0.15 (ASMFC 2020)	**Females**: *Small:* 0.24208 *Large:* 0.33281 **Males**: *Small:* 0.38074 *Large:* 0.63534 (Fernández-Chacón *et al.* 2021)
Size cutoff value for small and large natural mortality	n/a	**Females:** 88.16 mm CL **Males:** 89.09 mm CL (Converted from 250 mm TL) (Fernández-Chacón *et al.* 2021)

Supplemental Table 1. Probabilities of molting given pre-molt carapace length (CL) for females and males of European lobster (*H. gammarus*). Data from Sørdalen *et al.* (2022) on lobsters in fished and protected areas were combined, then converted from total length to carapace length for this study.

Pre-molt CL	Probability Molting	
	Female	**Male**
56	0.954438	0.998032
57	0.951875	0.997854
58	0.949056	0.997675
59	0.945928	0.997459
60	0.942785	0.99723
61	0.93953	0.996999
62	0.9361	0.996749
63	0.932807	0.99645
64	0.929357	0.996128
65	0.925385	0.995805
66	0.921234	0.995418
67	0.916776	0.995005
68	0.911843	0.994589
69	0.906924	0.994089
70	0.902283	0.993558
71	0.896988	0.993023
72	0.891413	0.992377
73	0.886097	0.99163
74	0.879958	0.990856
75	0.873675	0.990035
76	0.866951	0.989122
77	0.859425	0.988222
78	0.852112	0.987164
79	0.845206	0.985996
80	0.837312	0.984842
81	0.829186	0.983594
82	0.821436	0.982141
83	0.812478	0.98051
84	0.802719	0.978914
85	0.793065	0.977048
86	0.784099	0.974972
87	0.774081	0.972935
88	0.763528	0.970546
89	0.753681	0.967912
90	0.742565	0.965321
91	0.731203	0.962273
92	0.720504	0.958945
93	0.70734	0.955662
94	0.693841	0.951792
95	0.6823	0.947606
96	0.669446	0.943471
97	0.656279	0.939029
98	0.644049	0.933856
99	0.630305	0.928179
100	0.616654	0.922635
101	0.602488	0.91617
102	0.587108	0.90915
103	0.572675	0.902291
104	0.559467	0.894289
105	0.546175	0.885699
106	0.531627	0.87731
107	0.516899	0.867523
108	0.503462	0.857146
109	0.488632	0.847024
110	0.47273	0.835228
111	0.457912	0.822887
112	0.443497	0.810875
113	0.428896	0.796905
114	0.416458	0.782497
115	0.401268	0.768517
116	0.386254	0.753918
117	0.374399	0.737259
118	0.359801	0.71995
119	0.34544	0.703471
120	0.334156	0.684753
121	0.320325	0.665627
122	0.306783	0.64964
123	0.296191	0.626156
124	0.283265	0.604382
125	0.270663	0.586642
126	0.260849	0.564143
127	0.251267	0.541395
128	0.239636	0.52306
129	0.228355	0.500058
130	0.219613	0.477057
131	0.20904	0.458721
132	0.198821	0.440496
133	0.190929	0.417955
134	0.181417	0.395729
135	0.172254	0.378258
136	0.165199	0.356878
137	0.156721	0.336017
138	0.148578	0.319787
139	0.142326	0.300124
140	0.134833	0.281124
141	0.127654	0.266481
142	0.122157	0.255807

Supplemental Table 2. Fecundity-at-size estimates for American lobster (*H. americanus*; Estrella and Cadrin 1995) and European lobster (*H. gammarus*; Agnalt 2008). Functions and parameters are in Table 2.

H. americanus		H. gammarus	
Carapace Length (mm)	Average fecundity	Carapace Length (mm)	Average fecundity
73.72	295	74.30	1781
84.84	8117	82.13	5445
95.44	12374	89.03	8679
104.66	17213	95.04	11493
112.32	22166	100.23	13920
118.54	26891	104.71	16018
123.61	31241	108.63	17853
127.80	35198	112.02	19443
131.32	38796	114.92	20798
134.34	42083	117.56	22033
136.97	45103	119.87	23115
139.28	47895	121.97	24099
141.35	50491	123.86	24985
143.22	52917	125.58	25788
144.91	55196	127.13	26515
146.47	57346	128.54	27172
147.90	59381	129.86	27790
149.23	61314	131.11	28377
150.47	63156	132.23	28901
151.63	64916	133.29	29398
152.72	66601	134.29	29865
153.75	68218	135.23	30304
154.72	69773	136.12	30721
155.64	71272	136.95	31113
156.51	72717	137.78	31500
157.35	74115	138.56	31865
158.15	75467	139.30	32211
158.91	76777	139.99	32536
159.64	78049	140.68	32858
160.34	79284	141.33	33161
161.01	80484	141.94	33449
161.66	81653	142.55	33735
162.29	82791		

Supplemental Figure 1. Relative abundance of American lobster individuals at each age in the model, starting at age five. The unfished abundance is shown twice, once in each column for comparison to the different types of female protections under *F* = 0.3 and *F* = 0.9.

Supplemental Figure 2. Relative abundance of European lobster individuals at each age in the model, starting at age five. The unfished abundance is shown twice, once in each column for comparison to the different types of female protections under *F* = 0.3 and *F* = 0.9.