Title:

Inhibitory basal ganglia nuclei differentially innervate pedunculopontine nucleus subpopulations and evoke differential motor and valence behaviors.

Author list and affiliations:

[1,2,3]Michel Fallah

[2]Kenea C Udobi

[2]Aleksandra E Swiatek

[2]Chelsea B Scott

[1,2,4]Rebekah C Evans*

[1]Interdisciplinary Program in Neuroscience, Georgetown University, Washington, DC, USA 20007

[2]Department of Neuroscience, Georgetown University Medical Center, Washington, DC, USA 20007

[3]Senior author

[4]Lead contact

Lead Contact footnote:

*Rebekah Evans, Ph.D.

re285@georgetown.edu

Department of Neuroscience

Georgetown University

3970 Reservoir Rd. NW, NRB EG31

Washington DC

Email addresses:

re285@georgetown.edu

Graphical Abstract:

Summary:
The canonical basal ganglia model predicts that the substantia nigra *pars reticulata* (SNr) and the globus pallidus *externa* (GPe) will have specific effects on locomotion: the SNr inhibiting locomotion and the GPe enhancing it. In this study, we use *in vivo* optogenetics to show that the GPe exerts non-canonical effects on locomotion while the SNr has no gross motor impact through inhibition of the PPN. We also show that these structures mediate opposing effects on reward. To understand how these structures differentially modulate the PPN, we use *ex vivo* whole-cell recording with optogenetics to comprehensively dissect the SNr and GPe synaptic connections to regionally- and molecularly-defined populations of PPN neurons. The SNr inhibits all PPN subtypes, but most strongly inhibits caudal glutamatergic neurons. The GPe selectively inhibits caudal glutamatergic and GABAergic neurons, avoiding both cholinergic and rostral cells. This circuit characterization reveals non-canonical basal ganglia pathways for locomotion and valence.

Key words: basal ganglia; brainstem; locomotion; reward; mesencephalic locomotor region; inhibition; electrophysiology; optogenetics; substantia nigra; pedunculopontine nucleus

Introduction:
The pedunculopontine nucleus (PPN) is a brainstem structure heavily interconnected with subcortical structures, such as the basal ganglia, thalamus, and spinal cord. The PPN has been implicated in locomotor control and valence processing[1–4]. However, understanding PPN circuitry is complicated by its anatomical and molecular heterogeneity. It has three molecularly-defined cell types, distinguished by their major neurotransmitter[5] and distinct rostral and caudal topography. While cholinergic and glutamatergic neurons are more densely packed in the caudal PPN, GABAergic neurons are more densely packed in the rostral PPN[5,6]. These molecularly-defined populations display very minimal overlap[5,7,8]. This unique topography has been conserved across species[9–11], but it is unknown how the basal ganglia connects to each of the distinct cell types in the rostral and caudal PPN regions to modulate locomotor and valence outputs.

While the PPN interacts with all basal ganglia nuclei[3,9], the inhibitory inputs from the substantia nigra *pars reticulata* (SNr) and the globus pallidus *externus* (GPe) to the PPN are of particular interest because of their roles in the canonical motor output pathways for 'stop' and 'go,' respectively. Both the SNr and GPe are known to send axonal projections to the PPN[12–18]. Previous studies have identified SNr and GPe monosynaptic connections to the PPN cholinergic and glutamatergic neurons using rabies tracing[13–17]. However, inputs to the GABAergic neurons have only been described in terms of the entire mesencephalic locomotor region (MLR), which extends beyond the PPN[14]. While SNr projections to MLR GABAergic neurons were identified but cannot be isolated to the PPN, GPe projections to the MLR GABAergic neurons were not detected[14]. Of note, these previous studies consider the PPN as a whole rather than separating its rostral and caudal subregions. While viral tracing can provide insight into neuroanatomical connections, limitations of these techniques[19–21] and evidence that the rostral and caudal PPN may have distinct functions[22–25] prompt a

need for electrophysiological studies to determine whether the SNr and GPe inputs
differentially innervate regionally-defined PPN neurons.
Optogenetic manipulations of the molecularly-defined PPN cell types have resulted in
variable, and sometimes contradictory, locomotor and valence behaviors[7,14,16,17,22–33].
More recent studies have refined our understanding of these opposing behavioral
effects by selectively stimulating the rostral or caudal GABAergic and glutamatergic
PPN neurons[22–25]. In the basal ganglia, direct stimulation of molecularly-defined SNr
and GPe subpopulations can also have distinct effects on locomotion and valence[34–42].
Despite the potential for the basal ganglia to modify behavior through the motor
brainstem[14], it is not clear how the SNr and GPe influence the rostral and caudal PPN to
modify locomotion and valence.
Here, we use *ex vivo* whole-cell patch clamp electrophysiology paired with
optogenetics to selectively stimulate the SNr or GPe axons while recording inhibitory
inputs to the cholinergic, GABAergic, and glutamatergic PPN neurons to
comprehensively characterize the strength and pattern of inhibition across the
rostrocaudal PPN axis. We find that the GPe preferentially inhibits caudal GABAergic
and glutamatergic PPN neurons whereas the SNr most strongly inhibits a caudal
'hotspot' of PPN glutamatergic neurons. Stimulating SNr or GPe axons over the PPN *in*
vivo evokes opposing valence processing outcomes with place aversion during SNr
stimulation and place preference during GPe stimulation. Surprisingly, and counter to
the predictions of the canonical basal ganglia model, the GPe decreases locomotion
through its projections to the PPN. While both the SNr and GPe inhibit the PPN, our
results show that each nucleus differentially modulates the activity of specific cell types
in the rostral and caudal PPN and are implicated in non-canonical basal ganglia circuits
for modulating locomotion and valence processing.
104
Results:
SNr and GPe axons display distinct distribution patterns across the rostral and
caudal PPN.
To characterize the axon distribution pattern from the SNr and GPe throughout the
rostral and caudal extent of the PPN, we stereotaxically injected a channelrhodopsin
virus with a fluorescent marker (AAV1-hSyn-ChR2-eYFP) into either the SNr or GPe of
ChAT-Cre/Ai9-tdTomato mice (**Figure 1A&C**). We sliced 200 μm sagittal slices from
mouse brains three weeks after viral injection, and cleared the brain slices using the
CUBIC clearing method[43]. Using the red fluorescent ChAT-positive neurons to define
the borders of the PPN, we found that SNr axons appear in both the rostral and caudal
PPN. The SNr axons fill the rostral PPN evenly, but appear more distinctly clustered in
specific areas of the caudal PPN (**Figure 1B**), whereas GPe axonal projections more
densely fill the caudal region and strikingly avoid the rostral region (**Figure 1D**). These
anatomical findings predict that the SNr will inhibit both rostral and caudal PPN neurons,
while the GPe will selectively inhibit the caudal PPN. However, this axon imaging
method does not reveal whether these axons form functional synapses with the neurons
in the PPN and does not provide information on the strength of inhibition from each
structure onto specific PPN cell types. Therefore, to evaluate the functional connectivity

and synaptic characteristics from
these inhibitory basal ganglia output
structures to regionally- and
molecularly-defined PPN neural
subpopulations, we used *ex vivo*
whole-cell electrophysiology to
systematically record the inhibitory
input to the rostral and caudal PPN
from the SNr and GPe in three
genetic mouse lines (ChAT-cre,
Vgat-cre, and Vglut2-cre) which label
cholinergic, GABAergic, and
glutamatergic PPN subpopulations,
respectively (**Figure S1**).
137

Figure 1. SNr and GPe axons display distinct distribution patterns across the rostral and caudal PPN. (A,C) Stereotaxic injection of AAV1 delivering *hSyn-ChR2-eYFP* to the SNr or GPe of ChAT-Cre/Ai9-tdTomato mice, respectively. (B, D) Confocal images of EYFP-filled SNr or GPe axons across the PPN, respectively. *CNII: cranial nerve II, scp: superior cerebellar peduncle; PB: parabrachial nucleus*

The SNr inhibits cholinergic
neurons in both the rostral and
caudal PPN to a similar extent, but
evokes stronger rebound in
rostral neurons.
To evaluate the functional
connection between the SNr and cholinergic PPN neurons, we performed *ex vivo*
whole-cell patch clamp electrophysiology paired with optogenetics to record synaptic
currents in ChAT-Cre/Ai9-tdTomato mice injected with ChR2 in the SNr three weeks
prior (**Figure 2A**). Neurons were targeted for whole-cell patch clamp in both the rostral
and caudal PPN across three sagittal slices to evaluate the medial to lateral extent of
the PPN (**Figure 2B**). To measure optically-evoked inhibitory post-synaptic currents
(oIPSCs), cholinergic PPN neurons were held at -50 mV in whole-cell voltage clamp
and blue (470 nm) light was applied (2 ms pulse duration, 13 mW) to activate the ChR2-
infected SNr axons. In connected neurons, inhibitory currents were recorded in the
presence of glutamatergic receptor blockers (AP5, NBQX, CNQX) during stimulation
(**Figure 2C**). In a subset of neurons, a GABA-a receptor blocker, GABAzine, was
applied to confirm the oIPSCs were GABA-mediated (**Figure 2C**). GABAzine eliminated
SNr-evoked inhibitory currents in most cells, but a small residual current remained in
some cells (see *Methods* for details). We found that all rostral (n=19/19 cells, N=6 mice)
and essentially all caudal (n=26/27, N=6) recorded cholinergic PPN neurons responded
to SNr axon activation with an observable oIPSC (**Figure 2D**).
To determine whether the SNr input to the rostral and caudal PPN cholinergic
neurons differed in synaptic strength, we measured the amplitude of each oIPSC in a 20
Hz train of 2 ms optical stimulations applied for 2 seconds. There was no difference in
amplitude of the first oIPSC in the stimulus train between rostral vs caudal cholinergic
neurons (mean ± SEM; first oIPSC n=15 Rostral: 92.6 ± 18 pA, n=20 Caudal: 115.2 ±
17 pA; unpaired t-test, t=0.9041, df=33, p=0.3725; **Figure 2E&F**). Going forward, the
first oIPSC amplitude of the train will be referred to as the first oIPSC.
While the rostral and caudal anatomical separation of PPN neurons is commonly
used, it is a relatively coarse division. Therefore, we carefully mapped the location of

each recorded cholinergic PPN neuron by matching slices to their bregma reference in
the Paxinos and Franklin's mouse brain atlas and aligning the midline of cholinergic
neuron distribution which separates the loosely spread rostral neurons and densely
packed caudal neurons (See *Methods* for details, **Figure 2G**). The strength of the SNr
inhibitory connection is depicted by a color scale representative of the first oIPSC
(**Figure 2G, *right***). Throughout both the rostral and caudal PPN, cholinergic neurons
receiving larger inputs from the SNr are intermixed with neurons receiving smaller
inputs.
To compare short-term synaptic plasticity characteristics between rostral and caudal
cholinergic PPN neurons, we normalized the amplitude of each current in the train to the
first current. The oIPSC amplitude remained relatively constant with each subsequent
pulse in both rostral and caudal cholinergic neurons (**Figure 2H**). In addition, the paired-
pulse ratio (PPR) of the peak of the first two currents in the stimulation train did not
differ (mean ± SEM; PPR n=15 Rostral: 0.96 ± 0.04, n=20 Caudal: 0.97 ± 0.05;
unpaired t-test, t=0.1082, df=33, p=0.9145; **Figure 2I**). SNr synaptic inputs onto
cholinergic neurons do not undergo significant short-term synaptic plasticity in either the
rostral or caudal PPN.
Although oIPSC amplitude is indicative of the strength of a synapse, current size
does not always correlate with functional impact on a neuron's action potential
output[20,44,45]. Because cholinergic PPN neurons display spontaneous firing[46–50], we
characterized the impact of SNr-mediated inhibition on action potential firing rate. While
optically stimulating SNr axons with the same 20 Hz blue light protocol as above, we
observed a decrease in firing rate in both rostral and caudal cholinergic PPN neurons
Figure 2J-K). However, there was no significant difference between the percent
frequency change in rostral and caudal neurons during SNr axon stimulation (mean ±
SEM; %Frequency n=14 Rostral: During Stimulation 35.62 ± 8.87 %, Post Stimulation
136.1 ± 7.803 %; n=23 Caudal: During Stimulation 52.11 ± 8.43 %, Post Stimulation
119.1 ± 4.614 %; 2way ANOVA, Stimulation$_{F(1, 35)=81.9}$ p<0.0001, Region$_{F(1, 35)=0.002052}$
p=0.9641, Interaction$_{F(1, 35)=3.273}$ p=0.0790; **Figure 2K**) or in the absolute decrease in
firing frequency during optical stimulation [median (IQR); ΔFrq During Opto n=14
Rostral: -2.47 Hz (-3.37 to -1.44 Hz), n=23 Caudal: -2.10 Hz (-3.11 to -0.84 Hz); Mann
Whitney, U=131, p=0.3602; **Figure 2L**]. These data show that the SNr equally inhibits
action potential firing in rostral and caudal PPN cholinergic neurons.
Post-inhibitory rebound firing has been observed in the PPN as well as other
brainstem structures[49,51–57] and is associated with greater excitability, temporal
encoding, and generation of oscillatory activity[58]. Indeed, we found that the rostral and
caudal cholinergic PPN neurons exhibited an increase in action potential firing after
inhibition (**Figure 2K**). This is in agreement with previous literature showing rebound
spikes following inhibitory postsynaptic potentials recorded in PPN neurons of male rats
during electrical stimulation of the SNr[51,52]. However, rostral cholinergic neurons had
greater rebound activity compared with caudal cholinergic neurons as measured by the
absolute change in rebound frequency [median (IQR); ΔRebFrq n=14 Rostral: 1.64 Hz
(1.00 to 2.17 Hz), n=22 Caudal: 0.83 Hz (0.42 to 1.14 Hz); Mann Whitney, U=79,
p=0.0142; **Figure 2M**]. This aligns with previous findings showing that rostral cholinergic
neurons tend to have stronger intrinsic rebound activity compared to caudal neurons[49].

Importantly, we show that such rebound can be evoked by synaptic inhibition from the
SNr.

Figure 2. SNr inhibition of rostral and caudal ChAT+ PPN neurons. (A) Experimental set up to identify red ChAT+ PPN neurons for whole-cell patch clamp while stimulating ChR2-filled SNr axons [N=6]. (B) White arrowheads pointing to neurobiotin-filled patched neurons within the PPN across three 200µm slices. (C) Example trace of the first five oIPSCs [blue] in the 2-second 20 Hz train inhibited by GABA-a receptor blocker, GABAzine [green], while holding the cell at -50mV. (D) Percent connected among patched neurons in the rostral and caudal regions. (E) Average oIPSC amplitude at each of 40 optogenetic light pulses in n=15 rostral neurons and n=20 caudal neurons. (F) *Left*, Individual cell data for the first oIPSC amplitude and, *right*, example current traces. (G) Cell mapping of patched neuron locations with the first oIPSC amplitude represented by the color scale. (H) Normalized current amplitudes in E. (I) *Left*, Individual cell data for the PPR between the first two oIPSC amplitudes in the train and, *right*, example current traces. (J) Example voltage traces of action potential firing during a 2-second 20 Hz train stimulation in rostral (left) and caudal (right) neurons. (K) Percent of pre-optical stimulation firing frequency during stimulation and rebound in n=14 rostral vs n=23 caudal neurons. (L) Individual cell data for the absolute change in frequency during optical stimulation [ΔFrq During Opto]. (M) Individual cell data for the absolute change in rebound frequency post-stimulation [ΔRebFrq]; rostral vs. caudal p=0.0142. (N) Correlation analysis, color scale representing Spearman r [-1,1] and size representing p-value [1,0]. (O) Negative correlation between the absolute change in frequency during stimulation and post-stimulation rebound; r=-0.372, p=0.039. * p<0.05; bar graph data represent mean ± SEM; box plots show median line with boxes showing IQR and whiskers showing 9th and 91st percentiles.

To detect trends among SNr connectivity characteristics and intrinsic
electrophysiological properties of cholinergic PPN neurons, we performed a correlation
analysis to determine the strength and directionality of the relationship among five
parameters [first oIPSC, PPR, ΔFrq During Opto, pre-optical stimulation frequency (Pre-
Opto Frq), and ΔRebFrq; **Figure 2N**]. We found a weak but significant negative
correlation between the frequency decrease during SNr axon stimulation and the
rebound frequency increase (Spearman r=-0.372, p=0.039, **Figure 2N&O**). This finding
is expected since strong inhibition is more effective at producing rebound activity than
weak inhibition. However, rostral PPN cholinergic neurons do not receive significantly
stronger inhibition from the SNr compared with caudal cholinergic neurons (**Figure
2F&L**). Therefore, larger rebound frequency in rostral neurons is likely due to
differences in intrinsic cellular properties rather than synaptic properties[49]. Put together,
these findings show that the SNr inhibits rostral and caudal cholinergic PPN neurons to
a similar extent, but evokes stronger rebound firing in the rostral neurons.
230
**The SNr inhibits GABAergic neurons in both the rostral and caudal PPN to a
similar extent.**
While there are several rabies tracing studies suggesting SNr axons project to the
cholinergic and glutamatergic PPN neurons[13–17], SNr projections to the GABAergic PPN
neurons are inferred from a study of the entire mesencephalic locomotor region[14] and a
characterization of their synaptic strength has not been conducted. Therefore, we
evaluated the synaptic characteristics between the SNr and GABAergic neurons in the
rostral and caudal PPN. In Vgat-Cre/Ai9-tdTomato mice, we injected ChR2 into the SNr
to optically stimulate axons over the PPN while patching Vgat+ GABAergic PPN
neurons (**Figure 3A**). We post-hoc stained the slices for ChAT+ neurons to verify the
location of all patched GABAergic neurons within the cholinergic PPN. We found that all
rostral (n=12/12, N=6) and most caudal (n=20/22, N=6) recorded GABAergic PPN
neurons received inhibitory input from the SNr (**Figure 3B**).
Although there was no major difference in the proportion of GABAergic neurons
receiving inhibitory input from the SNr in the rostral vs caudal PPN, we compared the
current amplitudes to characterize the synaptic strength and pattern of this connection.
The oIPSC amplitudes across the 2-second stimulation as well as the first oIPSC
showed no difference between rostral and caudal GABAergic neurons [median (IQR);
first oIPSC n=9 Rostral: 76.5 pA (27.0 to 209.9 pA), n=13 Caudal: 59.7 pA (37.7 to
269.7 pA); Mann Whitney, U=50, p=0.6005; **Figure 3C&D**]. We also did not observe
significant differences in short-term synaptic plasticity between rostral and caudal
GABAergic neurons [median (IQR); PPR n=9 Rostral: 1.01 (0.88 to 1.16), n=13 Caudal:
1.06 (0.96 to 1.15); Mann Whitney, U=48, p=0.5123; **Figure 3 F&G**]. Defining short-
term depression as a PPR<0.95 and short-term potentiation as PPR>1.05, we found 3/9
rostral and 7/13 caudal GABAergic PPN neurons showed short-term potentiation, while
3/9 rostral and 3/13 caudal GABAergic PPN neurons showed short-term depression.
These data suggest additional heterogeneity within the GABAergic PPN subpopulation.
Similar to the SNr connection with cholinergic PPN neurons, SNr input to the
GABAergic PPN does not result in significant short-term synaptic plasticity and does not
differ in amplitude or characteristics between rostral and caudal regions.

Figure 3. SNr inhibition of rostral and caudal Vgat+ PPN neurons. (A) Experimental set up to identify red Vgat+ PPN neurons for whole-cell patch clamp while stimulating ChR2-filled SNr axons [N=6]. (B) Percent connected among patched neurons in the rostral and caudal regions. (C) Average oIPSC amplitude at each of 40 optogenetic light pulses in n=9 rostral neurons and n=13 caudal neurons. (D) *Left*, Individual cell data for the first oIPSC amplitude and, *right*, example current traces. (E) Cell mapping of patched neuron locations with the first oIPSC amplitude represented by the color scale. (F) Normalized current amplitudes in C. (G) *Left*, Individual cell data for the PPR between the first two oIPSC amplitudes in the train and, *right*, example current traces. (H) Percent of pre-optical stimulation firing frequency [% Pre-Opto Frq] during stimulation and and rebound in n=7 rostral and n=19 caudal neurons. (I) Individual cell data for the absolute change in frequency during optical stimulation [ΔFrq During Opto]. (J) Correlation analysis, color scale representing Spearman r [-1,1] and size representing p-value [1,0]. (K) Negative correlation between the absolute change in frequency during stimulation and first oIPSC amplitude; r=-0.755, p=0.001. (L) Negative correlation between the pre-optical stimulation firing frequency and the PPR; r =-0.706, p=0.002. Box plots show median line with boxes showing IQR and whiskers showing 9th and 91st percentiles.

Because GABAergic PPN neurons display spontaneous firing[50], we recorded action
potential firing in current clamp while optically stimulating the SNr axons. The percent
inhibition was not significantly different between rostral and caudal neurons (mean ±
SEM; %Frequency n=7 Rostral: During Stimulation 37.91 ± 15.91 %, Post Stimulation
100.0 ± 14.57 %; n=19 Caudal: During Stimulation 30.68 ± 8.03 %, Post Stimulation
113.2 ± 5.187 %; 2way ANOVA, Stimulation$_{F(1, 24)=44.63}$ p<0.0001, Region$_{F(1, 24)=0.09642}$
p=0.7588, Interaction$_{F(1, 24)=0.8841}$ p=0.3565; **Figure 3H**). There was also no difference in
the absolute frequency change during optical stimulation between rostral and caudal
neurons [median (IQR); ΔFrq During Opto n=7 Rostral: -5.03 Hz (-5.55 to -1.34 Hz),
n=19 Caudal: -4.29 Hz (-11.02 to -2.94 Hz); Mann Whitney, U=55, p=0.5336; **Figure 3I**].
These data show that the SNr equally inhibits action potential firing in rostral and caudal
GABAergic PPN neurons and does not evoke significant rebound firing in either neural
population.

Overall, we found that the SNr inhibits the rostral and caudal GABAergic PPN neurons to a similar extent, but observed greater variability in synaptic strength and impact on neuronal action potential firing among caudal GABAergic neurons. One caudal subgroup responded with very small (<60 pA) amplitude currents and another subgroup responded with currents more than triple that amplitude, with one neuron responding with a 2.2 nA inhibitory current. We mapped the location of each recorded GABAergic neuron, as described above, and found no specific location corresponding with the neurons receiving stronger SNr inhibition across the PPN landscape (**Figure 3E**). This strong variability suggests that GABAergic PPN neurons may be divided into functionally distinct neural types that do not correspond to their rostrocaudal anatomical location. To determine if the neurons receiving larger inhibitory currents have distinct characteristics, we performed a correlation analysis (**Figure 3J**). As expected, we found that the absolute frequency reduction during inhibition is correlated with the first oIPSC amplitude recorded in a cell (Spearman r=-0.755, p=0.001; **Figure 3K**). Surprisingly, however, we found a significant negative correlation between PPR and the pre-optical stimulation frequency (Pre-Opto Frq, **Figure 3L**). Specifically, input to slower-firing neurons displayed greater short-term synaptic facilitation. This finding supports the idea that multiple functional neural subpopulations may be present within the GABAergic PPN population. Together, these results show that the SNr functionally inhibits GABAergic neurons across the rostrocaudal extent of the PPN, but suggests the presence of additional heterogeneity within the GABAergic PPN population.

The SNr differentially inhibits the rostral and caudal glutamatergic PPN neurons.
Previous rabies tracing anatomical studies have identified SNr inputs to glutamatergic PPN neurons[14,16,17], but a full characterization of their synaptic strength and rostral and caudal connectivity has not been conducted. In Vglut2-Cre/Ai9-tdTomato mice, we injected ChR2 into the SNr to optically stimulate axons over the PPN while patching Vglut2+ glutamatergic neurons in the PPN (**Figure 4A**). While holding cells at -50mV and applying 470 nm blue light stimulation, neurons were identified as connected if oIPSCs were observed. Similar to our findings in cholinergic and GABAergic neurons, we found that all rostral (n=19/19, N=6) and all caudal (n=28/28, N=6) recorded glutamatergic neurons received inhibitory input from the SNr (**Figure 4B**).
Comparing the oIPSC amplitudes across the 20 Hz train, we discovered that the caudal glutamatergic neurons receive larger inhibitory currents than rostral glutamatergic neurons (**Figure 4C**). The median first oIPSC measured in caudal glutamatergic neurons was also significantly larger than those recorded in rostral neurons [median (IQR); first oIPSC n=13 Rostral: 61.6 pA (19.2 to 113.9 pA), n=13 Caudal: 313.5 pA (143.8 to 729.1 pA); Mann Whitney, U=29, p=0.0035; **Figure 4D**]. After post-hoc staining for the cholinergic PPN and mapping the location of each recorded glutamatergic PPN neuron as described above, we identified a 'hotspot' of strong SNr inhibition in a medial-caudal group of glutamatergic neurons (**Figure 4E**). This finding shows that a subset of caudal PPN neurons preferentially receive exceptionally strong SNr input.
To determine whether the SNr input to the rostral and caudal PPN show differential short-term synaptic plasticity, we compared the normalized oIPSC amplitudes. We found that inhibitory inputs to the rostral glutamatergic neurons display short-term

synaptic facilitation with amplitudes increasing in subsequent stimulations (**Figure 4F**).
However, strong facilitation occurred in only a few neurons that drove the mean upward
and there was no significant difference between the PPR observed in rostral and caudal
glutamatergic neurons when activating SNr axons [median (IQR); PPR n=13 Rostral:
1.07 (0.86 to 1.38), n=13 Caudal: 0.92 (0.89 to 1.02); Mann Whitney, U=77, p=0.7241;
Figure 4G].
Because glutamatergic PPN neurons spontaneously fire[16,50], we wanted to determine
if larger inhibitory currents mediated larger decreases in the firing rate of caudal PPN
neurons. Therefore, we recorded tonic action potential firing in glutamatergic PPN
neurons while optically stimulating SNr axons (**Figure 4H**). Although inhibition
measured as the percent of pre-optical stimulation firing frequency was not significantly
different between rostral and caudal neurons (mean ± SEM; %Frequency n=13 Rostral:
During Stimulation 32.62 ± 10.21 %, Post Stimulation 100.3 ± 6.113 %; n=17 Caudal:
During Stimulation 24.71 ± 7.30 %, Post Stimulation 97.48 ± 2.228 %; 2way ANOVA,
Stimulation$_{F(1, 28)=133.6}$ p<0.0001, Region$_{F(1, 28)=0.5214}$ p=0.4762, Interaction$_{F(1, 28)=0.1765}$
p=0.6776; **Figure 4I**), we found that the absolute frequency decrease was significantly
larger in caudal glutamatergic neurons compared to rostral neurons during SNr
stimulation [median (IQR); ΔFrq During Opto n=13 Rostral: -3.21 Hz (-12.21 to -1.95
Hz), n=17 Caudal: -14.08 Hz (-20.04 to -6.07 Hz); Mann Whitney, U=55, p=0.0197;
Figure 4J]. To examine the discrepancy between the percent decrease and the
absolute decrease in firing frequency, we compared the spontaneous firing frequency
between rostral and caudal glutamatergic neurons in cells that had no holding current
applied. We found that caudal glutamatergic neurons fire faster than rostral
glutamatergic neurons [median (IQR); Spontaneous Frequency n=11 Rostral: 6.8 Hz
(5.4 to 17.2 Hz), n=16 Caudal: 20.6 Hz (11.2 to 37.2 Hz); Mann Whitney, U=45,
p=0.0343; **Figure 4K**]. Together, these findings show that fast-firing caudal
glutamatergic PPN neurons receive larger SNr-mediated IPSCs and display greater
decreases in their action potential firing.
Since we found that the caudal glutamatergic neurons receive larger inhibitory
currents and display greater decreases in firing rate during SNr axon stimulation, we
performed a correlation analysis to evaluate the strength of the relationship among the
synaptic and firing characteristics (**Figure 4L**). Interestingly, there was a weak but
significant positive correlation between the PPR and frequency change during
stimulation showing that input displaying short-term synaptic depression occurs in
neurons with greater decreases in firing frequency (Spearman r=0.486, p=0.030; **Figure**
4M). As expected, SNr-mediated oIPSCs with larger amplitudes are correlated with
greater decreases in firing frequency during stimulation (Spearman r=-0.841,
p<0.00001; **Figure 4N**). Greater pre-optical stimulation firing frequency is correlated
with larger frequency decreases during stimulation (Spearman r=-0.791, p<0.0001;
Figure 4O). This finding is expected because the fast-firing neurons can be more
strongly inhibited in terms of absolute decrease in action potential frequency (i.e. a floor
effect). Surprisingly, however, pre-optical stimulation firing frequency is correlated with
the first oIPSC amplitude (Spearman r=0.818, p<0.0001; **Figure 4P**), indicating that the
SNr is more strongly connected to glutamatergic PPN neurons with faster firing rates.
These correlations highlight strong selectivity in SNr projections to PPN glutamatergic
neurons. Overall, these findings support that the SNr selectively targets and strongly

inhibits fast-firing caudal glutamatergic PPN neurons compared to rostral neurons, and
that there is a medial-caudal hotspot where SNr inhibition of PPN glutamatergic neurons
is particularly strong.

Figure 4. SNr inhibition of rostral and caudal Vglut2+ PPN neurons. (A) Experimental set up to identify red Vglut2+ PPN neurons for whole-cell patch clamp while stimulating ChR2-filled SNr axons [N=6]. (B) Percent connected among patched neurons in the rostral and caudal regions. (C) Average oIPSC amplitude at each of 40 optogenetic light pulses in n=13 rostral neurons and n=13 caudal neurons. (D) *Left*, Individual cell data for the first oIPSC amplitude and, *right*, example current traces; p=0.0035. (E) Cell mapping of patched locations with the first oIPSC amplitude represented by the color scale. (F) Normalized current amplitudes in C. (G) *Left*, Individual cell data for the PPR between the first two oIPSC amplitudes in the train and, *right*, example current traces. (H) Example voltage traces of action potential firing during a 2-second 20 Hz train stimulation in rostral and caudal neurons, top to bottom. (I) Percent of pre-optical stimulation firing frequency [% Pre-Opto Frq] during stimulation and rebound in n=13 rostral and n=17 caudal neurons. (J) Individual cell data for the absolute change in frequency during optical stimulation [ΔFrq During Opto]; p=0.0197. (K) Spontaneous frequency in n=11 rostral and n=16 caudal neurons; p=0.0343. (L) Correlation analysis, color scale representing Spearman r [-1,1] and size representing p-value [1,0]. (M) Positive correlation between the absolute change in frequency during stimulation and PPR, r=0.486, p=0.030. (N) Negative correlation between the absolute change in frequency during stimulation and first oIPSC amplitude; r=-0.841, p<0.00001. (O) Negative correlation between the absolute change in frequency during stimulation and pre-optical stimulation frequency; r=-0.791, p<0.0001. (P) Positive correlation between the first oIPSC amplitude and pre-optical stimulation frequency; r=0.818, p<0.0001. * p<0.05, ** p<0.01; box plots show median line with boxes showing IQR and whiskers showing 9th and 91st percentiles.

369
The SNr most strongly inhibits caudal glutamatergic PPN neurons.

Most of the previous PPN circuit work has been done with rabies tracing. Aside from technical limitations, rabies tracing can only indicate the presence of synaptic connections. Our electrophysiological findings comprehensively characterize the strength and characteristics of the SNr input to each of the PPN cell types in both the rostral and caudal regions. Using this comprehensive dataset, we compared SNr input across the different PPN cell types in each region under the same electrophysiological conditions. We found that the median amplitude of the first SNr-mediated oIPSC was the same among rostral PPN neurons irrespective of cell type [median (IQR); first oIPSC n=15 ChAT+: 74.7 pA (44.5 to 136.5 pA), n=9 Vgat+: 76.5 pA (27.0 to 209.9 pA), n=13 Vglut2+: 61.6 pA (19.2 to 113.9 pA); Kruskal-Wallis, $H_{(2,34)}$=0.7002 p=0.7046; **Figure 5A**]. The impact of these inhibitory inputs on neuronal firing did not differ with cell type as shown by the absolute change in frequency during SNr stimulation [median (IQR); ΔFrq During Opto n=14 ChAT+: -2.47 Hz (-3.37 to -1.44 Hz), n=7 Vgat+: -5.03 Hz (-5.54 to -1.34 Hz), n=13 Vglut2+: -3.21 Hz (-12.21 to -1.95 Hz); Kruskal-Wallis, $H_{(2,31)}$=2.389 p=0.3028; **Figure 5B**].

We found that the SNr-mediated oIPSCs in caudal glutamatergic neurons were significantly larger than in cholinergic and GABAergic caudal PPN neurons [median (IQR); first oIPSC n=20 ChAT+: 109.3 pA (58.5 to 168.6 pA), n=12 Vgat+: 59.7 pA (37.7 to 269.7 pA), n=13 Vglut2+: 313.5 pA (143.8 to 729.1 pA); Kruskal-Wallis, $H_{(2,43)}$=8.717 p=0.0128; Dunn's test, ChAT+ vs Vgat+ p=0.6409, ChAT+ vs Vglut2+ p=0.0042, Vgat+ vs Vglut2+ p=0.0295; **Figure 5C**]. Likewise, the impact of SNr inhibition on neuron firing rate was strongest in caudal glutamatergic neurons. The absolute frequency decrease during stimulation in caudal glutamatergic neurons was larger than both cholinergic and GABAergic neurons [median (IQR); ΔFrq During Opto n=23 ChAT+: -2.10 Hz (-3.11 to -0.84 Hz),

Figure 5. The SNr most strongly inhibits caudal glutamatergic PPN neurons. (A,C) Individual cell data for the first oIPSC amplitude recorded in each cell type for rostral and caudal PPN neurons, respectively. (B,D) Individual cell data for the absolute change in frequency during stimulation in each cell type for rostral and caudal PPN neurons, respectively. (E) Graphical depiction of SNr stimulation results. * p<0.05, ** p<0.01, **** p<0.0001; box plots show median line with boxes showing IQR and whiskers showing 9th and 91st percentiles.

n=19 Vgat+: -4.29 Hz (-11.02 to -2.94 Hz), n=17 Vglut2+: -14.08 Hz (-20.04 to -6.07
Hz); Kruskal-Wallis, $H_{(2,56)}$=24.99 p<0.0001; Dunn's test, ChAT+ vs Vgat+ p=0.0049,
ChAT+ vs Vglut2+ p<0.0001, Vgat+ vs Vglut2+ p=0.0332; **Figure 5D**]. While we found
no significant difference between SNr-mediated oIPSCs recorded in GABAergic and
cholinergic neurons (**Figure 5C**), the GABAergic neuron firing frequency was more
strongly inhibited than cholinergic (**Figure 5D**). The greater reduction in firing frequency
may be driven by the caudal GABAergic subgroup receiving particularly large oIPSCs
(**Figure 3D**) and could be explained if GABAergic PPN neurons have higher input
resistance than cholinergic neurons, as has been previously suggested[59]. Altogether,
our findings show that the SNr most strongly inhibits the glutamatergic neurons followed
by GABAergic neurons in the caudal PPN and more weakly inhibits rostral PPN neurons
without any cell type bias (**Figure 5E**).
427
The GPe preferentially, but weakly, inhibits a subset of caudal GABAergic and
glutamatergic PPN neurons.
GPe projections to the cholinergic and glutamatergic PPN neurons have been shown
in previous rabies tracing studies[14–17], but projections to the GABAergic MLR neurons
was not detected[14]. To evaluate the synaptic strength of these connections and
determine regional connectivity, we repeated the previous whole-cell patch clamp
experiments paired with optogenetics, but this time injecting ChR2 into the GPe of
ChAT-Cre/Ai9-tdTomato, Vgat-Cre/Ai9-tdTomato, and Vglut2-Cre/tdTomato mice
(**Figure 6A**). If oIPSCs were observed in response to optical stimulation, the neuron
was identified as connected. In a subset of neurons, a GABA-a receptor blocker,
GABAzine, was applied to confirm the oIPSCs were GABA-mediated (See *Methods* for
details; **Figure 6B**). We recorded GPe-mediated oIPSCs in all three cell types of the
PPN. As predicted by the low density of GPe axons present in the rostral PPN (**Figure**
1D), very few rostral PPN neurons received synaptic input from the GPe. This was true
within each molecularly-defined PPN cell type [caudal vs rostral % connected; ChAT+:
30% vs 0% (n=9/30 vs 0/7), Vgat+: 68% vs 38% (n=21/31 vs 6/16), Vglut2+: 69% vs
26% (n=24/35 vs 5/19); **Figure 6Ci-iii**]. These findings show that the GPe preferentially
targets caudal PPN neurons compared to rostral neurons. Since so few rostral PPN
neurons received GPe inhibition, we focused our characterization of GPe inhibition to
the caudal PPN.
The percentage of connected caudal neurons was greater than the percentage of
connected rostral neurons in both GABAergic and glutamatergic neurons (one-sided
Fisher's exact test; Vgat: OR=0.2857, CI=0.09190 to 0.9391, p=0.047; Vglut2:
OR=0.1637, CI=0.05069 to 0.5767, p=0.0033; **Figure 6Cii&iii**), but we found no
statistical difference between the rostral and caudal connectivity in cholinergic neurons
(OR=0, CI=0 to 1.301, p=0.1150; **Figure 6Ci**). The number of connected caudal
cholinergic neurons was not significantly greater than its 0% rostral connectivity (**Figure**
6Ci). In addition, the average oIPSC amplitude across the train and the first oIPSC
measured in the connected cholinergic neurons was small [median (IQR); first oIPSC
n=6 ChAT+: 22.6 pA (9.75 to 50.20 pA); **Figure 6D&E**]. Among recorded caudal
cholinergic PPN neurons, both connected and non-connected, these small inhibitory
currents displayed negligible effects on the neuronal firing frequency of the cholinergic
PPN population in both the percent of pre-optical stimulation frequency and absolute

frequency change [median (IQR); n=29 ChAT+: %Frequency During Stimulation 96.67
% (85.1 to 100.7 %); ΔFrq During Opto -0.16 Hz (-0.90 to 0.03 Hz); **Figure 6H&I**].
These findings indicate that the GPe has essentially no direct effect on the cholinergic
PPN neural population. For this reason, we exclude GPe inhibition of cholinergic PPN
neurons in the synaptic characterizations shown in Figures 6F&H. Together, we show
that the GPe preferentially inhibits caudal GABAergic and glutamatergic neurons while
largely avoiding rostral and cholinergic PPN neurons.
To determine whether the synaptic strength and characteristics from the GPe to the
PPN differed between GABAergic and glutamatergic PPN subtypes, we evaluated the
amplitude of the GPe-mediated oIPSCs during the 2-second 20 Hz optical stimulation.
Both GABAergic and glutamatergic neurons responded to GPe stimulation with oIPSCs
ranging from ones to hundreds of picoamperes (**Figure 6D**). While there was no
significant difference in the first oIPSC measured in GABAergic and glutamatergic
neurons [median (IQR); first oIPSC n=19 Vgat+: 24.77 pA (6.72 to 54.17 pA), n=15
Vglut2+: 65.71 pA (14.50 to 133.4 pA); Kruskal-Wallis, $H_{(2,37)}$=3.050, p=0.2176; **Figure
6E**], the normalized amplitudes show that GPe input to GABAergic neurons facilitates
with subsequent stimulations, while the input to glutamatergic neurons depresses
(**Figure 6F**). The GPe-mediated inhibitory input to glutamatergic neurons displayed a
significantly lower PPR than to GABAergic neurons [median (IQR); PPR n=19 Vgat+:
0.97 (0.76 to 1.2), n=15 Vglut2+: 0.81 (0.55 to 0.93); Mann-Whitney, U=76, p=0.0206;
Figure 6G]. While short-term synaptic plasticity patterns were variable among
GABAergic neurons with some depressing and others facilitating, GPe input to
glutamatergic neurons generally depresses.
Short-term synaptic depression can dampen the impact that repeated IPSCs have on
action potential output. Therefore, we compared the firing frequency change in each cell
type during optogenetic stimulation of GPe axons. The PPN GABAergic neuron
population firing rate was decreased to about 58% of their pre-stimulation firing with a
median frequency decrease of 1.92 Hz during stimulation (**Figure 6H&I**). Similarly, the
glutamatergic neuron population firing rate decreased to about 64% of their pre-
stimulation firing with a median frequency decrease of 0.89 Hz during stimulation
(**Figure 6H&I**). We found no statistical difference between GPe inhibition of firing
frequency in caudal GABAergic and glutamatergic PPN neurons (mean ± SEM;
%Frequency n=29 ChAT+: During Stimulation 92.39 ± 7.59 %, Post Stimulation 104.10
± 5.23 %; n=18 Vgat+: During Stimulation 57.70 ± 10.11 %, Post Stimulation 91.81 ±
6.98 %; n=24 Vglut2+: During Stimulation 64.15 ± 7.63 %, Post Stimulation 104.7 ± 5.39
%; 2way ANOVA, Stimulation$_{F(1, 69)=25.46}$ p<0.0001, Cell type$_{F(1, 69)=4.444}$ p=0.0153,
Interaction$_{F(1, 69)=2.805}$ p=0.0674; Fisher's LSD, df=138, %Frequency During Stimulation:
ChAT+ vs. Vglut2+ p=0.0035, ChAT+ vs. Vgat+ p=0.0016, Vglut2+ vs. Vgat+ p=0.5378,
%Frequency Post Stimulation: ChAT+ vs. Vglut2+ p=0.8883, ChAT+ vs. Vgat+
p=0.2838, Vglut2+ vs. Vgat+ p=0.2184; median (IQR); ΔFrq During Opto n=25 ChAT+: -
0.16 Hz (-0.90 to 0.03 Hz), n=18 Vgat+: -1.92 Hz (-3.85 to -0.60 Hz), n=29 Vglut2+: -
0.89 Hz (-3.28 to -0.43 Hz); Kruskall-Wallis, $H_{(2,69)}$=17.21, p=0.0002; Dunn's test,
ChAT+ vs Vgat+ p=0.0002, ChAT+ vs Vglut2+ p=0.0009, Vgat vs Vglut2+ p=0.4020;
Figure 6H&I). These findings show that the GPe inhibits GABAergic and glutamatergic
neural populations to a similar extent.

Figure 6. GPe inhibition of the three PPN cell types. (A) Experimental set up to identify red ChAT+, Vgat+, and Vglut2+ PPN neurons for whole-cell patch clamp while stimulating ChR2-filled GPe axons [N=6]. (B) Example trace of the first five oIPSCs in the 2-second 20 Hz train [blue] inhibited by GABA-a receptor blocker, GABAzine [green], while holding the cell at -50mV. (C) *Left*, Percent connected among patched neurons in the rostral and caudal regions and, *right*, cell mapping of patched locations with the first oIPSC amplitude represented by the color scale. *Top to bottom*, i. ChAT+, ii. Vgat+, and iii. Vglut2+ datasets. (D) Average oIPSC amplitude at each of 40 optogenetic light pulses in n=6 ChAT+, n=19 Vgat+, and n=15 Vglut2+ caudal PPN neurons. (E) *Left*, Individual cell data for the first oIPSC amplitude and, *right*, example current traces. (F) Normalized current amplitudes in C. (G) *Left*, Individual cell data for the PPR between the first two oIPSC amplitudes in the train; p=0.0206. R*ight, top*, example current trace of short-term synaptic facilitation in VgAT+ neurons. *Right, bottom*, example current traces of short-term synaptic depression in Vgat+ and Vglut2+ neurons. (H) Percent of pre-optical stimulation firing frequency [%Pre-Opto Frq] during and post-stimulation in n=25 ChAT+, n=18 Vgat+, and n=29 Vglut2+ caudal PPN neurons. (I) Individual cell data for the absolute change in frequency during stimulation [ΔFrq During Opto].

(J) Correlation analysis for Vgat+ neurons, color scale representing Spearman r [-1,1] and size representing p-value [1,0]. (K) Negative correlation between the absolute change in frequency during stimulation and first oIPSC amplitude; r=-0.627, p=0.044. (L) Correlation analysis for Vglut2+ neurons. (M) Negative correlation between the absolute change in frequency during stimulation and the pre-stimulation firing frequency; r=-0.648, p=0.014. *p<0.05, **p<0.01; box plots show median line with boxes showing IQR and whiskers showing 9th and 91st percentiles.

To determine if there are significant relationships among the synaptic and firing
characteristics, we performed a correlation analysis. Among GABAergic neurons
receiving GPe inhibition, we found that the amplitude of the first oIPSC is significantly
correlated with the absolute change in frequency during stimulation (Spearman r=-
0.627, p=0.044, **Figure 6K**). Although GPe input to GABAergic neurons exhibited
various PPRs, the correlation analyses revealed no significant relationship between
PPR and absolute change in frequency during stimulation (**Figure 6J**). These findings
suggest that the initial oIPSC amplitude of GPe input to GABAergic PPN neurons has a
stronger relationship with firing rate than short-term synaptic plasticity does.
As nearly all GPe inputs to glutamatergic neurons exhibited short-term synaptic
depression, we expected the correlation analysis to reveal a significant relationship
between PPR and the change in firing frequency. However, similar to the GPe input to
GABAergic neurons, PPR was not predictive of impact on neuronal activity (**Figure 6L**).
GPe inhibition of the glutamatergic neurons showed a significant correlation between
the absolute change in frequency during stimulation and pre-stimulation firing frequency
(Spearman r=-0.648, p=0.014, **Figure 6M**). However, because the inhibitory synaptic
current amplitude did not correlate with firing frequency, this correlation is likely due to a
floor effect in which faster firing neurons have a greater capacity for reductions in firing
frequency. Together, these findings show that the GPe preferentially inhibits caudal
GABAergic and glutamatergic PPN neurons.
527
***In vivo* activation of GPe and SNr axons in the PPN show differential effects on**
locomotion and valence.
Once we established that the SNr and GPe inhibit different profiles of regionally- and
molecularly-defined subpopulations of the PPN using *ex vivo* optogenetic
electrophysiology, we wanted to evaluate the behavioral consequences of selectively
stimulating these inputs *in vivo*. Direct optogenetic activation of PPN subpopulations
have been previously shown to either promote or inhibit motion, sometimes with
contradictory results[14,16,17,22–28]. To determine whether differential inhibitory inputs onto
the PPN influenced locomotor activity, we bilaterally injected wild-type mice with ChR2
in either the SNr or GPe and implanted an optical fiber over the PPN (**Figure 7A&B**).
Control mice were injected with an eGFP virus and implanted. After three weeks, we
placed each mouse in the open field to measure gross locomotor behavior with and
without optical stimulation of the SNr or GPe axons. We tracked the distance traveled
for each mouse at baseline and during bilateral stimulation of either the SNr or GPe by
applying blue light (473 nm, 4-4.5 mW) at 20 Hz with 2 ms pulses (4% duty cycle), for 1

Figure 7. *In vivo* activation of GPe and SNr axons in the PPN show differential effects on locomotion and opposite effects on valence. (A) Experimental set up to stimulate ChR2-filled SNr or GPe axons over the PPN *in vivo.* (B) Representative image of optical fiber tract overlaid with the approximate optical fiber placement for SNr- [green] and GPe- [orange] injected mice. (C) Distance traveled over time in an open field with 1 minute 20 Hz 4.25 mW optical stimulations over the PPN in N=9 control (Ctrl) mice (black circles), N=8 mice injected with ChR2 in the SNr (green diamonds), and N=9 mice injected with ChR2 in the GPe (orange hexagons); vertical blue lines represent periods of optical stimulation. (D) Average distance traveled for each mouse across the six 1-minute optical stimulations in the high (4.25 mW) and low laser power (0.25 mW) conditions (E) Representative mouse track tracings during real time place preference task in a three-chamber box and continuously stimulating EGFP- or ChR2-filled axons over the PPN at 20 Hz in SNr- and GPe-injected mice when the mice are in the stimulation zone. (F) Percent time spent in the stimulation zone in N=16 control mice, N=9 mice injected with ChR2 in the SNr, and N=10 mice injected with ChR2 in the GPe with 4.25 or 0.25 mW laser power. (G) Percent time spent in the stimulation zone during the first minute reintroduced to the RTPP box on day 2 of RTPP with the laser off (no chamber is stimulated) Black line = median. * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001; box plots show median line with boxes showing IQR and whiskers showing 9th and 91st percentiles. See related supplemental Figures S2-S4 and Videos S1-S4.

minute at a time (**Figure 7C**). We found that stimulation of these two basal ganglia
inputs to the PPN resulted in differential motor behaviors. Specifically, stimulation of
SNr axons in the PPN increased distance traveled, while stimulation of GPe axons in
the PPN decreased distance traveled [mean ± SEM; N=9 (3M, 6F) Control (Ctrl): 2.22 ±
0.24 m, N=8 (5M, 3F) SNr: 4.11 ± 0.51 m, N=9 (4M, 5F) GPe: 0.57 ± 0.18 m; Welch
ANOVA p<0.0001; Dunnett's test, Ctrl$_{4.25mW}$ vs SNr$_{4.25mW}$ p=0.0263, Ctrl$_{4.25mW}$ vs
GPe$_{4.25mW}$ p=0.0002; **Figure 7D & Figure S2**, see Video S1&2].
The PPN has been implicated in reward processing and stimulation of the PPN can
be reinforcing[7,28,29,31,35,60]. Therefore, we evaluated the effect of GPe or SNr axon

stimulation in the PPN on valence in the real-time place preference (RTPP) task. In a three-chamber apparatus, the mouse could freely move between chambers for ten minutes. Optical stimulation of the SNr or GPe axons in the PPN was applied when the mouse entered the stimulated chamber and remained on at 20 Hz (4% duty cycle) until the mouse exited that chamber (**Figure 7E**). To reduce the motor effects of optical stimulation, this experiment was conducted with unilateral stimulation only. Interestingly, we again found opposite effects when stimulating the two basal ganglia inputs to the PPN. Mice avoided the stimulated chamber when the SNr axons were stimulated, but preferred the stimulated chamber when the GPe axons were stimulated (mean ± SEM; N=16 (6M, 10F) Ctrl: 40.76 ± 2.0 %, N=9 (5M, 4F) SNr: 23.99 ± 5.2 %, N=10 (5M, 5F) GPe: 76.11 ± 3.7 %; Welch ANOVA, p<0.0001; Dunnett's test, $Ctrl_{4.25mW}$ vs $SNr_{4.25mW}$ p=0.0437, $Ctrl_{4.25mW}$ vs $GPe_{4.25mW}$ p<0.0001, $Ctrl_{4.25mW}$ vs $SNr_{0.25mW}$ p=0.0695, $Ctrl_{4.25mW}$ vs $GPe_{0.25mW}$ p=0.0467; **Figure 7F**, see Video S3&4). To further control for potential motion confounds, we recorded the location preference of the mice in the absence of optical stimulation the day after the stimulated RTPP assay (regardless of whether they received low or high power on day 1). The mice were placed in the central, neutral chamber and their location was tracked for 10 minutes. During the first minute of this assay, 8 out of 9 mice who had SNr optical stimulation on the first day continued avoiding the stimulated chamber while 9 out of 10 mice which had GPe optical stimulation continued to prefer the previously stimulated chamber compared to the median percent time spent in the stimulated chamber for mice in the control condition [median (IQR); N=16 (6M, 10F) Ctrl: 47% (14.79 to 69.24 %), N=9 (5M, 4F) SNr: 1.833% (0 to 41.75 %), N=10 (5M, 5F) GPe: 92.83% (58 to 99.21 %); Kruskall-Wallis, p=0.0008; Dunn's test, Ctrl vs SNr p=0.1433, Ctrl vs GPe 0.0312; **Figure 7G**).

In addition to sending projections to the PPN, the GPe and SNr send axon collaterals to multiple other brain areas[12,37,61,62]. Therefore, it is possible that stimulating these axons in the PPN could cause antidromic stimulation of the GPe and SNr cell bodies, resulting in the inhibition of non-PPN brain regions. However, it has recently been shown that 0.25 mW laser power stimulation of axons can prevent antidromic activation of cell bodies[42]. Therefore, we repeated these optogenetic behavioral experiments using 0.25 mW laser power. We found that the locomotion effect of SNr stimulation over the PPN was no longer present during low power stimulation, but GPe stimulation still significantly decreased locomotion (mean ± SEM; distance traveled N=9 (3M, 6F) Ctrl: 2.22 ± 0.24 m, N=6 (3M, 3F) SNr: 2.24 ± 0.50 m, N=8 (4M, 4F) GPe: 1.09 ± 0.22 m; Welch ANOVA p<0.0001; Dunnett's test, $Ctrl_{4.25mW}$ vs $SNr_{0.25mW}$ p>0.999, $Ctrl_{4.25mW}$ vs $GPe_{0.25mW}$ p=0.0131; **Figure 7D**). In the RTPP task, SNr axon stimulation continued to evoke place aversion at the trend level and GPe axon stimulation continued to significantly evoke place preference (mean ± SEM; % time in stimulation zone N=16 (6M, 10F) Ctrl: 40.76 ± 2.1 %, N=6 (3M, 3F) SNr: 27.72 ± 4.0 %, N=8 (4M, 4F) GPe: 68.41 ± 8.5 %; Welch ANOVA, p<0.0001; Dunnett's test, $Ctrl_{4.25mW}$ vs $SNr_{0.25mW}$ p=0.0695, $Ctrl_{4.25mW}$ vs $GPe_{0.25mW}$ p=0.0467; **Figure 7F**). These findings show that decreased locomotion and place preference are mediated by GPe inhibition of the PPN while aversion is mediated by SNr inhibition of the PPN.

The increase in locomotion observed when stimulating the SNr axons in the PPN with high laser power is unexpected given that previous studies directly stimulating SNr neurons report decreases in movement[34,36]. Because injecting constitutive AAVs into

the SNr can infect both GABAergic neurons and adjacent dopaminergic neurons of the
substantia nigra pars compacta (SNc), we wanted to determine whether stimulating
dopaminergic axons in the PPN enhanced locomotion and evoked aversion. To test
this, we injected a cre-dependent ChR2 into the substantia nigra of DAT-cre mice (3M
heterozygotes and 3F homozygotes) and ran open field and RTPP behavioral assays.
We found that selective stimulation of dopaminergic axons in the PPN increased
locomotion at high laser power (**Figure S3A**) and had no effect on locomotion at low
power (**Figure S3C**). This finding suggests that dopaminergic neurons in the SNc send
collateral projections to both the PPN and other locomotor-related structures (e.g. the
striatum) such that stimulation of PPN-projecting SNc neurons increases locomotion.
The effect of dopaminergic axon stimulation did not differ from whole substantia nigra
stimulation in either high (mean ± SEM; N=8 (5M, 3F) $SNr_{4.25mW}$: 4.107 ± 0.51 m, N=6
(3M, 3F) $DAT-Cre_{4.25mW}$: 3.778 ± 0.28 m; unpaired t-test, t=0.5159, df=12, p=0.6153;
Figure S3B) or low laser power (mean ± SEM; N=6 (3M, 3F) $SNr_{0.25mW}$: 2.238 ± 0.58 m,
N=9 (3M, 3F) $DAT-Cre_{0.25mW}$: 1.984 ± 0.30 m; unpaired t-test, t=0.4350, df=10,
p=0.6728; **Figure S3C**). On the other hand, we found that stimulation of dopaminergic
axons in the PPN did not cause place aversion, but rather induced real-time place
preference at both high and low laser power (**Figure S3D-F**). Selectively stimulating
dopaminergic axons induced place preference behavior significantly different from and
opposite to whole substantia nigra stimulation at high (mean ± SEM; N=9 (5M, 4F)
$SNr_{4.25mW}$: 23.99 ± 5.18 %, N=6 (3M, S3F) $DAT-Cre_{4.25mW}$: 70.83 ± 10.06 %; unpaired t-
test, t=4.545, df=13, p=0.0005; **Figure S3E**) and low laser power (mean ± SEM; N=6
(3M, 3F) $SNr_{0.25mW}$: 27.72 ± 4.0 %, N=6 (3M, 3F) $DAT-Cre_{0.25mW}$: 50.82 ± 4.8 %;
unpaired t-test, t=3.697, df=10, p=0.0041; **Figure S3F**). Together, these findings show
that the the nigral increase in locomotion is not mediated through the PPN, and that
dopaminergic input from the SNc to the PPN is rewarding while GABAergic input from
the SNr to the PPN is aversive.
625
Discussion:
In this study, we comprehensively characterize the synaptic strength and impact on
neuron action potential firing in molecularly- and regionally-defined PPN subpopulations
while stimulating the SNr or GPe using *ex vivo* electrophysiology and optogenetics. We
also stimulated the SNr or GPe axons in the PPN *in vivo* which revealed that the SNr
and GPe evoke differential locomotion and valence processing behaviors in non-
canonical basal ganglia circuits. Together, these findings show that distinct and
selective inhibition of PPN subpopulations by the SNr and GPe can alter behavioral
output.
635
Region- and cell type-specific inputs to the PPN
The SNr and GPe have been previously shown to form monosynaptic synapses with
PPN neurons using rabies viral tracing[12–17]. While SNr input to cholinergic and
glutamatergic PPN neurons has been reproduced across different studies[13–17], SNr
inputs to GABAergic PPN neurons has only been inferred from studies of the entire
mesencephalic locomotor region (MLR), consisting of both the PPN and cuneiform
nucleus[14]. Several studies have used slice electrophysiology to show that the SNr
inhibits PPN neurons generally[32] or to compare SNr input to cholinergic and non-

cholinergic PPN neurons[48,51,52]. However, a full characterization of the SNr input to
regionally- and molecularly-defined PPN neurons in adult animals has been lacking.
Using *ex vivo* whole-cell patch recordings, we were able to measure inhibitory currents
in each of the PPN cell types during SNr stimulation showing that the SNr functionally
inhibits each cell type within the rostral PPN to a similar extent while most strongly
inhibiting a medial-caudal 'hotspot' of glutamatergic neurons.
While the GPe has been shown to project to the cholinergic and glutamatergic PPN
neurons[15–17], our electrophysiological results show that GPe inhibition of the cholinergic
neurons is weak to non-existent and its inhibition of the glutamatergic PPN neurons is
caudally biased. In contrast to a previous study showing that the GPe does not project
to GABAergic MLR neurons[14], we find that the GPe also preferentially inhibits a caudal
subgroup of GABAergic PPN neurons. While the GPe selectively inhibits a caudal
subpopulation of GABAergic and glutamatergic PPN neurons, its inhibition is much
weaker than SNr inhibition of the PPN in terms of the proportion of connected PPN
neurons, the inhibitory current amplitudes, and the inhibition of neuronal firing. Overall,
our findings utilize a systematic approach to characterize the synaptic strength between
these inhibitory basal ganglia nuclei and the PPN across its rostrocaudal axis for each
cell type allowing us to compare SNr and GPe to PPN circuitry with regional-level
granularity. These results also encourage future work involving the PPN to consider the
individual influence of its rostral and caudal regions.
664
Noncanonical GPe motor circuits
We found that stimulation of GPe axons over the PPN decreased locomotion in both
low (putatively local) and high (putatively generating antidromic activity) laser power
stimulations. This behavioral outcome appears counter to the canonical model of basal
ganglia movement pathways, in which GPe activation increases movement. Although
there is strong support for this canonical model, recent *in vivo* recordings have found
subpopulations of GPe neurons that show activity patterns counter to this model
(neurons that increase activity during immobility)[63,64]. Similarly, a subset of GPe neurons
paradoxically increase activity upon indirect pathway striatal projection neuron
stimulation[41]. While one explanation for these heterogeneous responses could be local
inhibition within the GPe[40,41,65,66], another possibility is that distinct subpopulations within
the GPe differentially modulate locomotor behavior[39–41,61,67]. Our findings support non-
canonical basal ganglia motor pathways involving GPe neurons that project to the PPN.
Our characterization of region-specific inhibitory inputs to the PPN helps us
understand how GPe inhibition of the PPN can mediate decreases in locomotion. Our
electrophysiology experiments show that the GPe exerts minimal influence on the
rostral PPN neurons which appear to decrease locomotion[22,24,25] and preferentially
inhibits caudal GABAergic and glutamatergic PPN neurons. Because stimulating the
caudal GABAergic and glutamatergic PPN neurons increases locomotion[23], GPe
selective inhibition of these caudal neurons could explain our behavioral experiments
showing that stimulation of GPe axons over the PPN decreases locomotion. Future
studies are needed to determine if GPe inhibition of both cell types is necessary for
mediating decreased locomotion. Aligned with our findings, selectively stimulating the
Npas1+ or FoxP2+ subpopulations in the GPe decreases locomotion[39,41]. Some groups
have found that direct stimulation of the PV+ GPe neurons increases locomotion[39,62];

however, one study shows that inhibition of PV+ GPe neurons can increase
locomotion[42]. While we show that GPe projections to the PPN decrease locomotion,
future studies are needed to determine which GPe subpopulations project to the PPN to
modulate locomotion.
694
GABAergic vs dopaminergic effects of nigral projections to the PPN
While we found that stimulating nigral axons in the PPN with high laser power
increased locomotion, we observed no locomotor effect when using low laser power
stimulation. The simultaneous inhibition of caudal PPN neurons that promote
movement[23] and rostral PPN neurons that suppress movement[22,24,25] may explain a net
zero effect on locomotion during low power optical stimulation of SNr axons in the PPN.
Indeed, low laser power stimulation of nigral axons in the most rostral implant location
resulted in increased locomotion, while the most caudal implant location resulted in
decreased locomotion (**Figure S4**). Interestingly, the enhanced locomotion observed
during high laser power stimulation suggests that a subset of nigral neurons projecting
to the PPN promote movement. However, the SNr canonically suppresses movement
and the direct somatic stimulation of PV+ and GAD2+ SNr neurons that send axon
collaterals to the PPN[12,68] decreases gross locomotor movement[34,68]. Since injecting
constitutive AAVs into the substantia nigra can infect both GABAergic neurons of the
SNr and adjacent dopaminergic neurons of the SNc, it is possible that stimulation of
dopaminergic activity may mask the impact of stimulating the GABAergic SNr alone.
Both the increased locomotion with high laser power and the lack of change with low
laser power was recapitulated when we ran these experiments while selectively
stimulating dopaminergic projections from the SNc to the PPN (**Figure S3**). Therefore,
our findings suggest that activation of the PPN-projecting dopaminergic SNc neurons
promotes locomotion, but not through direct actions at the PPN. This is consistent with
previous literature showing that SNc dopaminergic neurons send axons to both the PPN
and striatum[69] and that direct stimulation of nigrostriatal dopaminergic neurons can
increase movement initiation and vigor[70]. Therefore, antidromic stimulation of the SNc
may evoke dopamine release into the striatum to increase movement in our high laser
power experiments. Future studies are needed to evaluate the motor behavioral impact
of the different SNr cell types when inhibiting the entire PPN versus the rostral or caudal
PPN subregions.
By contrast, we find that both high and low power non-selective SNr axon stimulation
in the PPN is aversive (**Figure 7E-G**), while both high and low power selective
stimulation of dopaminergic SNc axons in the PPN is rewarding (**Figure S3**). These
findings are particularly interesting because they indicate that the dopaminergic vs
GABAergic inputs from the substantia nigra to the PPN serve opposite functions
regarding valence signaling. Importantly, these findings indicate that GABAergic input
from the SNr is highly aversive and that this aversive signaling occurs through the PPN.
Interestingly, these findings also show that dopaminergic SNc neurons do not require
antidromic stimulation to signal reward. This means that dopaminergic signaling within
the PPN may serve to reinforce behavior. Future work is needed to parse out specific
reward and aversion pathways involving neural subpopulations within the SNr, SNc, and
PPN.
735

Differential PPN inhibition by the SNr and GPe mediate opposing valence processing outcomes

We found that stimulation of SNr axons over the PPN was aversive in real time place preference (RTPP), while stimulation of GPe axons over the PPN led to a preference for the stimulated chamber. Previous studies directly stimulating SNr and GPe subpopulations have implicated these structures in reward and aversion processing. Inhibition of Vgat+ SNr neurons has been shown to increase place preference[36] while its direct stimulation has no effect[34,36]. However, selective stimulation of the PV+ SNr neurons induces aversion in RTPP[34]. Although the GPe has been predominantly studied in the context of movement, recent work has begun to reveal a role for the GPe in valence processing[42,71–73]. One recent study shows that inhibition of PV+ GPe neurons induced aversion while inhibition of Npas1+ GPe neurons induced placed preference in RTPP[42]. Because activation of PV+ SNr neurons is aversive[34] and inhibition of PV+ GPe neurons is aversive[42], it is likely that our results are due to activation of the PV+ SNr and GPe axons in the PPN. While we show that activation of the PPN-projecting SNr and GPe subpopulations strongly affect valence processing, future studies are needed to determine whether these effects are specifically mediated by PV+ SNr and GPe projections to the PPN.

Neurons in the PPN have been implicated in reward and aversive-related behaviors[7,28,29,31–33,35,60]. The cholinergic PPN neurons are implicated in positive valence with place preference and increased lever pressing through their projections to the ventral tegmental area (VTA)[28,29]. Aligned with our findings, direct inhibition of the cholinergic PPN neurons evokes place aversion[28]. Of particular significance, we show that the SNr inhibits the cholinergic PPN neurons while the GPe axons avoid cholinergic PPN neurons. Therefore, it is possible that the difference between the SNr and GPe effects on valence is due to the difference in their inhibition of PPN cholinergic neurons.

Direct stimulation of the glutamatergic PPN neurons has also been implicated in positive valence with increased reinforcement behavior[7]. Both the cholinergic and glutamatergic PPN neurons directly influence dopamine release in the striatum and dopaminergic neuron activity in the SNc and VTA[7,28,29,74–78]. Specifically, the caudal PPN neurons are thought to innervate the medial part of the SNc and the VTA[9,78], two major reward pathway hubs. Our data show that the SNr most strongly inhibits the caudal glutamatergic PPN neurons. Therefore, strong SNr inhibition of these neurons would remove excitatory drive from the reward-related dopaminergic neurons. This pathway could also contribute to the aversive effect of SNr axon stimulation in the PPN. Future experiments are needed to test whether direct inhibition of the caudal glutamatergic PPN neurons receiving SNr input is sufficient to evoke aversion.

The pathways underlying the GPe-mediated rewarding effect are less clear and may indicate functional heterogeneity among caudal PPN neurons. We find that the GPe only inhibits a subset of caudal GABAergic and glutamatergic PPN neurons. It is possible that the subset of GABAergic PPN neurons targeted by the GPe disinhibits local cholinergic and glutamatergic neurons to increase excitation of the rewarding dopaminergic neuron pathways. Another possibility is that the specific GPe-inhibited PPN population may have particularly aversive-properties. For example, PPN neurons that excite amygdala nuclei evoke place avoidance[35,60]. In this case, GPe inhibition of these neurons could be rewarding by removing excitation of amygdala nuclei involved in

negative valence processing. To understand the PPN circuitry involved in GPe-mediated place preference, future experimental work must determine whether inhibiting GPe-targeted GABAergic or glutamatergic PPN subpopulations alone is sufficient to evoke place preference.

Conclusions

We have systematically characterized SNr and GPe inputs across the rostrocaudal axis of the PPN for each cell type – cholinergic, GABAergic, and glutamatergic. We show that the SNr inhibits nearly every PPN cell recorded with differential strength determined by cell type and anatomical biases while identifying a medial-caudal 'hotspot' of glutamatergic neurons most strongly inhibited by the SNr. In contrast, the GPe strikingly avoids the cholinergic PPN neurons and weakly, but selectively, inhibits a subpopulation of caudal GABAergic and glutamatergic neurons. We proposed that the differential inhibition of regionally-biased, mixed-cell type PPN subpopulations can alter behavioral outputs. By stimulating these inhibitory basal ganglia axons over the PPN, we show that the SNr evokes place aversion while the GPe evokes place preference and decreases locomotion. Together, our findings show that the SNr and GPe mediate opposing valence processing outcomes through the PPN and support a non-canonical basal ganglia motor pathway in which the GPe decreases locomotion through its projections to the PPN.

Acknowledgements:
Funding for this research was provided by an NIH BRAIN Initiative grant R00NS112417 (to RCE), NIH NINDS grant 1F30NS132399 (to MF), Michael J Fox Foundation Research Grant MJFF-022863 (to RCE), and Parkinson's Foundation Summer Student Fellowship PF-SSF-941810 (to AES). This research was supported by the Microscopy and Imaging Shared Resource (MISR, S10RR025661) and the Lombardi Comprehensive Cancer Center grant (P30-CA051008). We would like to thank the Division of Comparative Medicine for their animal care and management and Dr. Stefano Vicini, Dr. Aryn Gittis, and members of the Evans lab for their feedback on earlier versions of this manuscript.

Author Contributions:
M.F. and R.C.E. conceptualized the experiments; M.F. performed stereotaxic injections, conducted the electrophysiological experiments, and analyzed the data; A.E.S. and K.C.U. performed stereotaxic injections, conducted the behavioral experiments, and analyzed the data; C.B.S. performed stereotaxic injections for pilot experiments; M.F. wrote the initial draft of the manuscript; M.F. and R.C.E. edited and revised the manuscript.

Declaration of Interests
The authors declare no competing interests.

828 **Star Methods text**
829 **KEY RESOURCES TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
goat anti-choline acetyltransferase (ChAT)	Millipore	Cat#: AB144P, RRID: AB_2079751
Streptavidin, Cy5	Invitrogen	Cat#: SA1011
Streptavidin, DyLight™ 405	Invitrogen	Cat#: 21831
Donkey anti-goat IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 647	Invitrogen	Cat#: A-21447, RRID: AB_2535864
Sheep anti-tyrosine hydroxylase (TH)	Novus Biologicals	Cat# NB300-110, RRID: AB_10002491
DyLight™ 405 AffiniPure™ Donkey Anti-Sheep IgG (H+L)	Jackson ImmunoResearch Laboratories	Cat# 713-475-147, RRID: AB_2340740
Bacterial and virus strains		
AAV1-hSyn-hChR2(H134R)-EYFP	Addgene (Deisseroth Lab)	Cat#: 26973
AAV1-Ef1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA	Addgene (Deisseroth Lab)	Cat#: 20298
AAV1-hSyn-EGFP	Addgene (Roth Lab)	Cat#: 50465
Chemicals, peptides, and recombinant proteins		
D-AP5	Tocris and HelloBio	Cat#: 0106 and HB0225
NBQX	Tocris and HelloBio	Cat#: 1044 and HB0443
CNQX disodium salt	HelloBio	Cat#: HB0205
SR 95531 hydrobromide (GABAzine)	Tocris and HelloBio	Cat#: 1262 and HB0901
Experimental models: Organisms/strains		
ChAT-cre mice: B6.129S-Chat$^{tm1(cre)Lowl}$/MwarJ	Jackson	RRID: IMSR_JAX:031661
Vgat-cre mice: B6J.129S6(FVB)-Slc32a1$^{tm2(cre)Lowl}$/MwarJ	Jackson	RRID: IMSR_JAX:028862
Vglut2-cre mice: B6J.129S6(FVB)-Slc17a6$^{tm2(cre)Lowl}$/MwarJ	Jackson	RRID: IMSR_JAX:028863
Ai9-tdTomato mice: B6.Cg-Gt(ROSA)26Sor$^{tm9(CAG-tdTomato)Hze}$/J	Jackson	RRID: IMSR_JAX:007909
DAT-cre mice [B6.SJL-Slc6a3$^{tm1.1(cre)Bkmn}$/J	Jackson	RRID: IMSR JAX:006660
C57BL/6J	Jackson	RRID: IMSR_JAX:000664
Software and algorithms		
Igor Pro	WaveMetrics	Version 9.00
GraphPad Prism	Dotmatics	Version 9.5.0
Fiji software	Open source on GitHub	Version 1.51n
ANY-maze® software	Stoelting Company, Wood Dale, IL.	Version 7.36
Clampex software	Molecular Devices	Version 11.2
Other		

Multiclamp™ 700B amplifier		
Axon™ Digidata® 1550B		
Leica SP8AOBS++		
Zeiss Axio Imager Z2		
Sutter Instrument Model P-97		

830
831 **RESOURCE AVAILABILITY**
832
833 **Lead contact**
834 Further information and requests for resources and reagents should be directed to and will be fulfilled by
835 the lead contact, Dr. Rebekah C. Evans (re285@georgetown.edu).
836
837 **Materials availability**
838 This study did not generate new unique reagents.
839
840 **Data and code availability**
841 • All data and code supporting the current study will be shared by the lead contact upon request.
842 • No original code was generated in this paper.
843 • Any additional information required to reanalyze the data reported in this paper is available from
844 the lead contact upon request.
845
846 **EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS**
847
848 Animal Welfare
849 All animal procedures were approved by the Georgetown University Medical Center Institutional Animal
850 Care and Use Committee (IACUC). Measures were taken to ensure minimal animal suffering and
851 discomfort, and protocols were designed to minimize the number of animals used.
852
Animal Subjects
Homozygous Ai9-TdTomato mice [B6.Cg-Gt(ROSA)26Sor[tm9(CAG-tdTomato)Hze]/J; JAX# 007909] were bred
with homozygous ChAT-Cre mice [B6.129S-Chat[tm1(cre)Lowl]/MwarJ; JAX# 031661], homozygous Vgat-Cre
mice [B6J.129S6(FVB)-*Slc32a1*[tm2(cre)Lowl]/MwarJ; JAX# 028862], and Vglut2-Cre mice [B6J.129S6(FVB)-
Slc17a6[tm2(cre)Lowl]/MwarJ; JAX# 028863] to obtain ChAT-Cre/Ai9-tdTomato, Vgat-Cre/Ai9-tdTomato, and
Vglut2-Cre/Ai9-tdTomato in-house, respectively. C57BL/6J wildtype mice (Jax# 000664) and DAT-cre
mice [B6.SJL-Slc6a3[tm1.1(cre)Bkmn]/J; JAX#006660] were obtained from Jackson for behavioral experiments.
Animals were housed under a 12:12 light-dark cycle (lights on at 06:00 AM) with food and water *ad*
libitum.
862
Experimental Groups
For electrophysiological studies, ChAT-Cre/Ai9-tdTomato, Vgat-Cre/Ai9-tdTomato, and Vglut2-Cre/Ai9-
tdTomato mice of both sexes (3 males and 3 females in each group) aged 2-5 months were used. For
behavioral studies, C57BL/6J wildtype mice of both sexes, with age-matched littermates randomly
assigned to control or experimental groups. DAT-cre mice of both sexes were used for behavioral
experiments where specified. Behavioral assessments were exclusively conducted during the light phase,
and all testing chambers were cleaned with 70% ethanol between mice to mitigate potential olfactory
influences.
871
872 **METHOD DETAILS**
873
Viral injections and optical fiber implantation surgeries
Mice at least 7 weeks old were briefly anesthetized with inhaled 5% isoflurane using an anesthetic
chamber and placed onto a heated pad within the stereotaxic frame (Stoelting 51730UD). The skull was
stabilized with evenly positioned ear bars and nose cone properly positioned to deliver continuous 1-3%
isoflurane and oxygen at a steady flow of 1L/min throughout the surgery duration. Bupivacaine (5 mg/kg)
and carprofen (5mg/kg) were administered as local anesthetic and analgesic, respectively. A small

incision was made on the scalp to visualize bregma and lambda, which were used as references to level
the skull. Bilateral holes were drilled at either the coordinates of the SNr (AP -3.1 mm, ML ± 1.4 mm, DV -
4.7 mm relative to bregma) or GPe (AP -0.3 mm, ML ± 1.9mm, DV -3.9 mm relative to bregma). The 5µL
Hamilton microsyringe was positioned and 250 nL virus was injected at a rate of 0.2 µL/min. The syringe
was raised and rested 0.5 mm above the injection site for 10 minutes. For *ex vivo* optogenetic
experiments, AAV1-hSyn-hChR2(H134R)-EYFP (2.3 x 10^13 particles per milliliter, Addgene, Cat# 26973)
was injected into either the SNr or GPe. For *in vivo* optogenetic experiments, AAV1-hSyn-hChR2(H134R)-
EYFP 2.0 x 10^13 particles per milliliter, Addgene, Cat #26973), AAV1-hSyn-EGFP (1.1 x 10^13 particles
per milliliter, Addgene, Cat# 50465), or AAV1-Ef1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA
(2.3 x 10^13 particles per milliliter, Addgene, Cat#20298) was injected into either the SNr or GPe for
experimental and control mice, respectively. For mice used in the *in vivo* optogenetic experiments, an
optical fiber (200 µm core, 0.22 NA, 3.7 mm length) was implanted over the middle-to-rostral PPN (AP -
4.5, ML ± 1.1 mm, DV -3.5 relative to bregma). The skin on either side of the incision site was joined,
glued together using VetBond tissue adhesive, and fastened with two wound clips or the optical fiber was
fixed in position with dental cement. (C&B-Metabond Quick! Adhesive Luting Cement by Parkell Products
Inc, Patterson# 553-3484/CAT# S380; Jet Denture Repair Package by LANG, CAT# 1223F2). Post-
surgery, buprenorphine SR (1.5 mg/kg) was administered for long-acting analgesia. Mice were allowed to
recover on a heating pad until fully awake and were monitored daily for signs of distress or infection.
898
Electrophysiological solutions
Acute brain slices were prepared in modified artificial cerebrospinal fluid (aCSF) solutions with final
osmolarities ~300-310 mOsm and ~7.4 pH. The slicing solution contained (in mM) 198 glycerol, 2.5 KCl,
1.2 NaH_2PO_4, 20 HEPES, 25 $NaHCO_3$, 10 glucose, 10 $MgCl_2$, and 0.5 $CaCl_2$. The holding solution
contained (in mM) 92 NaCl, 2.5 KCl, 1.2 NaH_2PO_4, 30 $NaHCO_3$, 20 HEPES, 2 $MgCl_2$, 2 $CaCl_2$, 35
glucose, 5 sodium ascorbate, 3 sodium pyruvate, and 2 thiourea. Recording aCSF was made up of (in
mM): 125 NaCl, 25 $NaHCO_3$, 3.5 KCl, 1.25 NaH_2PO_4, 10 glucose, 1 $MgCl_2$, 2 $CaCl_2$. Whole-cell patch
clamp recordings used a potassium methane sulfonate ($KMeSO_3$)-based internal solution containing (in
mM) 122 methanesulfonic acid, 9 NaCl, 9 HEPES, 1.8 $MgCl_2$, 4 Mg-ATP, 0.3 Tris-GTP, and 14
phosphocreatine for a final osmolarity between 290 and 305 mOsm. The internal solution contained
neurobiotin (0.1-0.3%) for post-hoc staining. These reagents enable reliable electrophysiology recordings
in adult brainstem neurons[46,79].
911
Slicing and Electrophysiology
Animals ages 2-5 months old were anesthetized with inhaled isoflurane and transcardially perfused with
ice-cold slicing solution that had been bubbled with 95% O_2 and 5% CO_2. Mice were decapitated and
brains were quickly extracted from the skull, keeping the cerebellum intact. For sagittal slices, the brain
hemispheres were separated and the medial side was glued onto a 3% agar block fixed to the stage of a
semi-automatic Leica VT1200 microtome. 200 µm-thick slices were obtained and incubated at 34°C in
holding solution for 30 minutes then kept at room temperature. In all steps, the modified aCSF solutions
are bubbled with 95% O_2 and 5% CO_2. Slices in recording chamber during whole-cell patch clamp
experiments were continuously perfused with oxygenated recording aCSF kept at 28-34°C using a water
bath and in-line Warner heater. Cells were visualized using an Olympus OpenStand upright microscope
and 565nm ThorLabs LED light. Recording pipettes with resistance between 1.5-4 MΩ were prepared
using borosilicate glass capillaries (World Precision Instruments, Inc. 1B150F-4, 4-inch length, 1.5 mm
OD, 0.84 mm ID) with a micropipette puller (Sutter Instrument Model P-97). Recordings were obtained
using a Multiclamp™ 700B amplifier and Axon™ Digidata® 1550B controlled by Clampex 11.2. Voltage-
clamp signals were low-pass filtered at 2kHz and sampled at 10 kHz. Current-clamp signals were low-
pass filtered at 10kHz and sampled at 10 kHz. At the end of recording, the cell was sealed by moving the
recording pipette slightly above the soma and an image was taken of the PPN and pipette tip.
929
Ex vivo **optogenetic activation**
Whole-field optogenetic activation of channelrhodopsin-infected axons in brain slice was achieved with
a blue (470 nm) ThorLabs LED light sent to the tissue via a silver mirror. Light intensity measured at the
objective back aperture was 13 mW. Light activation was applied at 20 Hz with 2 ms pulse intervals for 2
seconds. All recordings were conducted in the presence of glutamatergic receptor blockers – 50µM D-

AP5 (Tocris Cat#0106 and HelloBio Cat#H0225), and 5µM NBQX (Tocris Cat#1044 and HelloBio
Cat#HB0443) alone or in combination with 20µM CNQX (HelloBio Cat#HB0205).
In voltage-clamp, cells were determined to be connected if observable optically evoked inhibitory post
synaptic currents (oIPSCs) were measured while holding the cell at -50 mV. In a subset of neurons, 10µM
GABAzine (Tocris Cat#1262 and HelloBio Cat#HB0901) was applied to ensure the oIPSCs were GABA-a
receptor mediated. In a few cells (17/54), small inhibitory currents remained with amplitudes averaging
about 24% of the first oIPSC measured. Any cells that required greater than 200pA to be held at -50mV or
that had access resistance that exceeded 30 MΩ at the start of recording were excluded from oIPSC
quantitative analyses as they were likely unhealthy and too leaky; however, the first oIPSC amplitude was
used as an approximate measure of input strength in the cell maps (SNr: 21/46; GPe: 11/40). oIPSC
amplitudes were measured from the baseline to the peak of each current. The paired pulse ratio (PPR)
was calculated from the amplitudes of the first two currents evoked in the 20 Hz train.
For current-clamp recordings, most patched PPN neurons spontaneously fired, but a few required a
small amount of hyperpolarizing or depolarizing current to fire [(R)ostral, (C)audal; SNr-PPN$_{ChAT}$ R: 3/14,
C: 4/23; SNr-PPN$_{Vgat}$ R: 2/7, C: 9/19; SNr-PPN$_{Vglut2}$ R: 2/13, C: 1/17; GPe-PPN$_{ChAT}$ R: 2/7, C: 9/25; GPe-
PPN$_{Vgat}$ R: 2/7, C: 8/18; GPe-PPN$_{Vglut2}$ R: 6/12, C: 7/29 required current). Neurons injected with more than
60 pA were excluded. In this manuscript, we use "pre-stimulation firing frequency" to refer to the time
period right before optical stimulation. The pre-stimulation firing frequency was measured using the
number of action potentials in the one-second epoch before the start of stimulation. The firing frequency
during stimulation was measured using the middle one-second of the 2-second stimulation. The post-
stimulation firing frequency was measured in the one-second epoch immediately following the end of the
stimulation. The absolute frequency change is the difference between the pre-simulation and during
stimulation frequencies. The rebound frequency change is the difference between the post-stimulation
and pre-stimulation frequencies. These values include cells with and without holding current applied. By
contrast, the spontaneous firing rate of the cell includes only cells with no holding current applied.
960
***In vivo* optogenetic activation**
Two behavioral tests, the open field (OF) test and real-time place preference (RTPP), were conducted
sequentially to assess gross locomotion and place preference in animals of at least 11 weeks old (mean
age ± SEM; SNr-injected mice: 132 ± 23.4 days old; GPe-injected mice: 110 ± 7.4 days old). Optical
stimulation was achieved by the application of blue light (473 nm) at 20 Hz with 2 ms pulses (4% duty
cycle). Behavioral testing was conducted under both high laser power (4.0-4.5 mW) and low laser power
(0.2-0.25 mW) conditions. Low power testing was employed to control for potential behavioral effects
related to antidromic stimulation. On day 1, mice underwent OF testing and were randomly assigned to
high or low power stimulation conditions. On day 2 mice were tested in the open field arena for a second
time under opposite power conditions. On day 3 mice were tested in the RTPP arena for 10 minutes and
were again randomly assigned to high or low power conditions. On day 4, mice were tested in the RTPP
arena for 10 minutes without any stimulation to assess retention of place memory. Finally, on day 5, mice
underwent RTPP under opposite power conditions (low vs high) and opposite chamber as stimulation
zone (stripes vs spots) to those on Day 3.
975
Open Field Test
OF test was conducted in an opaque arena measuring 40.64 x 40.64 x 40.64 cm. Animals were placed in
the center of the arena and allowed to explore freely for 10 minutes. This was followed by 5 minutes of
discontinuous bilateral photostimulation at either high or low power, 5 minutes of recovery, another 5
minutes of discontinuous photostimulation at 20 Hz (at either high or low power), and a final 5 minutes of
recovery. The discontinuous photostimulation consisted of three 1-minute photostimulation periods
interspersed with 1-minute intervals of no photostimulation. Movement during the OF test were recorded
and analyzed using ANY-maze® software (Stoelting Company, Wood Dale, IL).
984
Real-Time Place Preference
The real-time place preference (RTPP) apparatus consisted of a rectangular behavioral arena with three
chambers, each measuring 24.60 x 27.94 x 27.94 cm, with a non-reflective grey background. The
adjacent chambers had distinct visual cues: one chamber featured stripes and the other spots. These
striped and spotted zones were randomly assigned as stimulation zones between animals. The center
chamber and one of the adjacent chambers served as neutral zones, while the other adjacent chamber

served as a stimulation zone. Animals received unilateral optical stimulation when in the assigned
stimulation zone. Laser-on side was randomly chosen and counter-balanced between mice. Mice were
also randomly assigned to have low laser power RTPP first or high laser power RTPP first. In each case,
mice were given an unstimulated 10-minute trial on the day between the first and second RTPP
experiment to 'unlearn' which side was stimulated and the second RTPP experiment stimulated the
opposite chamber compared to the first RTPP experiment. For example, one mouse would have high
power stimulation on the striped side on day 1, no stimulation on day 2, and low power stimulation on the
spotted side on day 3. To mitigate locomotor effects with bilateral stimulation, unilateral optical stimulation
was used in RTPP. Individual subjects were placed in the RTPP apparatus and allowed 10 minutes to
explore both compartments. The time spent in each compartment was recorded in real-time using ANY-
maze® software (Stoelting Company, Wood Dale, IL).
1002
Immunohistochemistry and confocal imaging
After electrophysiological experiments, brain slices were fixed overnight in a 4% w/v paraformaldehyde
(PFA) solution in phosphate buffer (PB) solution, pH 7.6 at 4°C. The fixed brain slices were then stored in
phosphate buffer (PB) solution until immunostaining.
Brain slices were also collected at the conclusion of behavioral experiments. Mice were deeply
anesthetized and perfused with phosphate buffer (PB), followed by a fixative solution containing 4% w/v
paraformaldehyde (PFA) in PB, pH 7.6, at 4°C. Whole brains were extracted and fixed overnight in the
same PFA solution. After fixation, brains were stored in PB solution until further processing. For cubic
processing, 200 μm-thick sagittal slices were obtained from post-behavioral whole mouse brains using
the PELCO easiSlicer™ Vibratory Tissue Slicer.
A CUBIC tissue clearing protocol [43] was combined with immunofluorescence staining as in[79] for all
fixed brain slices from both electrophysiological and behavioral experiments. All steps are performed at
room temperature on a shaker. Slices were placed in CUBIC reagent 1 for 1-2 days; then washed in PB
3x for 1 hour each; placed in blocking solution (0.5% fish gelatin in PB) for 3 hours; placed in primary
antibodies for 2-3 days; washed in PB 3x for 2 hours each; placed in secondary antibodies for 2-3 days;
washed in PB 3x for 2 hours each; and placed in CUBIC reagent 2 for 2 hours before mounting onto
slides (Fisherbrand 12-550-403) in reagent 2 and sealed with frame-seal incubation chambers (Thermo
Scientific AB-0577) and a coverslip (Corning 2845-18).
Neurobiotin-filled patched neurons were stained with streptavidin antibody (Cy5, Invitrogen
Cat#SA1011, 1:1000 or DyLight™ 405, Invitrogen Cat#21831, 1:1000). Goat anti-ChAT primary antibody
(Sigma Cat#AB144P, 1:200) and donkey anti-goat Alexa Fluor™ 647 secondary antibody (Invitrogen
Cat#A-21447, 1:333) was used to identify the borders of the PPN. On an example slice to show the SNr
injection site, sheep anti-TH primary antibody (Novus Biologicals Cat#NB300-110, 1:1000) and donkey
anti-sheep (1:100) was used to delineate the GABAergic SNr from the dopamine-rich TH+ SNc neurons.
To identify the approximate location of patched cells for cell mapping, slices were imaged as tiled z-
stacks using a Leica SP8AOBS++ at the Microscopy and Imaging Shared Resource core facility at
Georgetown University. Additionally, to verify the injection site and optical implant placement in the brain
slices from mice that underwent behavioral testing, tiled z-stacks were obtained using either the
aforementioned Leica SP8AOBS++ or a Zeiss Axio Imager Z2 at the Georgetown Department of
Neuroscience core imaging facility.
1033
Approximate Cell Location
An image of the PPN with the pipette tip still in position was taken at the end of each cell recording and
confocal images of the slices after CUBIC clearing were obtained as described above. Confocal images
were overlapped with the pipette images using the streptavidin-stained patched cells (which aligned with
the pipette tip images) and autofluorescence of the superior cerebellar peduncle (scp) white fibers as
landmarks. The perimeter of the PPN was determined by the presence of ChAT+ neurons and generally
extended from the edge of the SNr to approximately 2/3 up the scp fibers. ChAT+ neurons were identified
by tdTomato expression in the ChAT-Cre mouse line and anti-ChAT staining in the Vgat-Cre and Vglut2-
Cre mouse lines. Since there is variability introduced during the slicing process, we matched slices to
three sagittal templates representing 1.00, 1.25, and 1.50 mm lateral to the midline in the Paxinos and
Franklin's Mouse Brain Atlas 4th edition. The most medial slice (1.00 mm lateral to the midline)
characteristically had few to no fibers of the superior cerebellar peduncle at the edge of the slice where
the cerebellum would tether. All neurons on this slice were considered 'caudal' following the typical shape

of the PPN. Slices matching 1.25 and 1.50 mm from the midline had a region of densely clustered
cholinergic neurons and a region of loosely dispersed cholinergic neurons. An oval ROI was created
following the length and orientation of ChAT+ neuron spread. The midline of the oval demarcated the
rostral and caudal regions. Overlapping the midlines allowed us to evaluate rostral-caudal differences
across mice with a small margin of error, particularly along the midline. For this reason, we also mapped
the location of patched cells which shows the distance from the midline and can reveal medial-lateral
differences. The XY coordinates were determined using Fiji software and plotted in Igor. The color scale
applied to the points represented the amplitude of the first oIPSC and ranged from 0 to 250pA so that
amplitudes greater than 250pA were shown in the same maximal bright color.
1056

Approximate Implant Location
To verify the location of the optical implant, anatomical landmarks for locating the PPN were identified by
immunofluorescent labeled PPN cholinergic neurons. Sections containing visible optical fiber scarring
were stained with anti-ChAT immunofluorescent labeling. These histological images were then visually
matched to the corresponding sagittal planes of the stereotaxic atlas, based on the identified anatomical
landmarks. The approximate position of the optical cannula was localized in the histological images and
was in the middle-to-rostral part of the PPN.
1064

1065   **QUANTIFICATION AND STATISTICAL ANALYSIS**
1066

Electrophysiological traces were processed in Igor Pro 9 (Wavemetrics) and all statistical analyses were
performed in GraphPad Prism 9.5. All parametric data in text is reported as mean $\pm$ standard error of the
mean (SEM). Two-tailed unpaired t-tests were used to compare between two groups and one-way
ANOVA followed by Dunnett's post hoc test was used to compare three or more groups. If the equality of
variances was violated (i.e. the ratio of the largest and smallest standard deviations was greater than 2
and Bartlett test was statistically significant), Welch's ANOVA test was alternatively used. Two-way
ANOVA followed by Fisher's LSD post hoc test was performed for percent of pre-optical stimulation
frequency as each comparison group was predetermined. Non-parametric data in text is reported as
median [interquartile range (IQR)]. Boxplots show medians, 25th and 75th percentile as first and third
quartile box edges, and 9th and 91st percentiles as whiskers. We used Mann-Whitney rank-sum test for
comparing two groups and Kruskal-Wallis for multiple comparisons. As each comparison group within the
Kruskal-Wallis tests was planned and stand alone, uncorrected Dunn's post hoc test was performed to
determine significance between groups. To determine if caudal GPe-connectivity is significantly greater
than rostral GPe-connectivity, as expected from axon projection patterns, a one-tailed Fisher's exact test
and odds ratio effect size was performed. For correlation matrix analyses, non-parametric Spearman r
was computed. Statistical details of experiments can be found in the text, figure legends, and
Supplemental Table 1. Biological replicates are individual cells (n) from 6 separate mice (N=3 males and
3 females) in the electrophysiological experiments and are individual mice of both sexes in the behavioral
experiments (numbers described in results text). Same sex littermates were randomly allocated to control
or experimental groups and experimenters were not blinded during experiments. Statistical significance
was evaluated as $p < 0.05$.
1088
1089
1090

Main figure titles and legends
Figure 1. SNr and GPe axons display distinct distribution patterns across the
rostral and caudal PPN. (A,C) Stereotaxic injection of AAV1 delivering *hSyn-ChR2-*
eYFP to the SNr or GPe of ChAT-Cre/Ai9-tdTomato mice, respectively. (B, D) Confocal
images of EYFP-filled SNr or GPe axons across the PPN, respectively. *CNII: cranial*
nerve II, scp: superior cerebellar peduncle; PB: parabrachial nucleus
1097

Figure 2. SNr inhibition of rostral and caudal ChAT+ PPN neurons. (A)
Experimental set up to identify red ChAT+ PPN neurons for whole-cell patch clamp
while stimulating ChR2-filled SNr axons [N=6]. (B) White arrowheads pointing to

neurobiotin-filled patched neurons within the PPN across three 200μm slices. (C)
Example trace of the first five oIPSCs [blue] in the 2-second 20 Hz train inhibited by
GABA-a receptor blocker, GABAzine [green], while holding the cell at -50mV. (D)
Percent connected among patched neurons in the rostral and caudal regions. (E)
Average oIPSC amplitude at each of 40 optogenetic light pulses in n=15 rostral neurons
and n=20 caudal neurons. (F) *Left*, Individual cell data for the first oIPSC amplitude and,
right, example current traces. (G) Cell mapping of patched neuron locations with the first
oIPSC amplitude represented by the color scale. (H) Normalized current amplitudes in
E. (I) *Left*, Individual cell data for the PPR between the first two oIPSC amplitudes in the
train and, *right*, example current traces. (J) Example voltage traces of action potential
firing during a 2-second 20 Hz train stimulation in rostral (left) and caudal (right)
neurons. (K) Percent of pre-optical stimulation firing frequency during stimulation and
rebound in n=14 rostral vs n=23 caudal neurons. (L) Individual cell data for the absolute
change in frequency during optical stimulation [ΔFrq During Opto]. (M) Individual cell
data for the absolute change in rebound frequency post-stimulation [ΔRebFrq]; rostral
vs. caudal p=0.0142. (N) Correlation analysis, color scale representing Spearman r [-
1,1] and size representing p-value [1,0]. (O) Negative correlation between the absolute
change in frequency during stimulation and post-stimulation rebound; r=-0.372,
p=0.039. * p<0.05; bar graph data represent mean ± SEM; box plots show median line
with boxes showing IQR and whiskers showing 9th and 91st percentiles.
1121
Figure 3. SNr inhibition of rostral and caudal Vgat+ PPN neurons. (A) Experimental
set up to identify red Vgat+ PPN neurons for whole-cell patch clamp while stimulating
ChR2-filled SNr axons [N=6]. (B) Percent connected among patched neurons in the
rostral and caudal regions. (C) Average oIPSC amplitude at each of 40 optogenetic light
pulses in n=9 rostral neurons and n=13 caudal neurons. (D) *Left*, Individual cell data for
the first oIPSC amplitude and, *right*, example current traces. (E) Cell mapping of
patched neuron locations with the first oIPSC amplitude represented by the color scale.
(F) Normalized current amplitudes in C. (G) *Left*, Individual cell data for the PPR
between the first two oIPSC amplitudes in the train and, *right*, example current traces.
(H) Percent of pre-optical stimulation firing frequency [% Pre-Opto Frq] during
stimulation and and rebound in n=7 rostral and n=19 caudal neurons. (I) Individual cell
data for the absolute change in frequency during optical stimulation [ΔFrq During Opto].
(J) Correlation analysis, color scale representing Spearman r [-1,1] and size
representing p-value [1,0]. (K) Negative correlation between the absolute change in
frequency during stimulation and first oIPSC amplitude; r=-0.755, p=0.001. (L) Negative
correlation between the pre-optical stimulation firing frequency and the PPR; r =-0.706,
p=0.002. Box plots show median line with boxes showing IQR and whiskers showing
9th and 91st percentiles.
1140
Figure 4. SNr inhibition of rostral and caudal Vglut2+ PPN neurons. (A)
Experimental set up to identify red Vglut2+ PPN neurons for whole-cell patch clamp
while stimulating ChR2-filled SNr axons [N=6]. (B) Percent connected among patched
neurons in the rostral and caudal regions. (C) Average oIPSC amplitude at each of 40
optogenetic light pulses in n=13 rostral neurons and n=13 caudal neurons. (D) *Left*,
Individual cell data for the first oIPSC amplitude and, *right*, example current traces;

p=0.0035. (E) Cell mapping of patched locations with the first oIPSC amplitude
represented by the color scale. (F) Normalized current amplitudes in C. (G) *Left*,
Individual cell data for the PPR between the first two oIPSC amplitudes in the train and,
right, example current traces. (H) Example voltage traces of action potential firing during
a 2-second 20 Hz train stimulation in rostral and caudal neurons, top to bottom. (I)
Percent of pre-optical stimulation firing frequency [% Pre-Opto Frq] during stimulation
and rebound in n=13 rostral and n=17 caudal neurons. (J) Individual cell data for the
absolute change in frequency during optical stimulation [ΔFrq During Opto]; p=0.0197.
(K) Spontaneous frequency in n=11 rostral and n=16 caudal neurons; p=0.0343. (L)
Correlation analysis, color scale representing Spearman r [-1,1] and size representing
p-value [1,0]. (M) Positive correlation between the absolute change in frequency during
stimulation and PPR, r=0.486, p=0.030. (N) Negative correlation between the absolute
change in frequency during stimulation and first oIPSC amplitude; r=-0.841, p<0.00001.
(O) Negative correlation between the absolute change in frequency during stimulation
and pre-optical stimulation frequency; r=-0.791, p<0.0001. (P) Positive correlation
between the first oIPSC amplitude and pre-optical stimulation frequency; r=0.818,
p<0.0001. * p<0.05, ** p<0.01; box plots show median line with boxes showing IQR and
whiskers showing 9th and 91st percentiles.
1165
Figure 5. The SNr most strongly inhibits caudal glutamatergic PPN neurons.
(A,C) Individual cell data for the first oIPSC amplitude recorded in each cell type for
rostral and caudal PPN neurons, respectively. (B,D) Individual cell data for the absolute
change in frequency during stimulation in each cell type for rostral and caudal PPN
neurons, respectively. (E) Graphical depiction of SNr stimulation results. * p<0.05, **
p<0.01, **** p<0.0001; box plots show median line with boxes showing IQR and
whiskers showing 9th and 91st percentiles.
1173
Figure 6. GPe inhibition of the three PPN cell types. (A) Experimental set up to
identify red ChAT+, Vgat+, and Vglut2+ PPN neurons for whole-cell patch clamp while
stimulating ChR2-filled GPe axons [N=6]. (B) Example trace of the first five oIPSCs in
the 2-second 20 Hz train [blue] inhibited by GABA-a receptor blocker, GABAzine
[green], while holding the cell at -50mV. (C) *Left*, Percent connected among patched
neurons in the rostral and caudal regions and, *right*, cell mapping of patched locations
with the first oIPSC amplitude represented by the color scale. *Top to bottom*, i. ChAT+,
ii. Vgat+, and iii. Vglut2+ datasets. (D) Average oIPSC amplitude at each of 40
optogenetic light pulses in n=6 ChAT+, n=19 Vgat+, and n=15 Vglut2+ caudal PPN
neurons. (E) *Left*, Individual cell data for the first oIPSC amplitude and, *right*, example
current traces. (F) Normalized current amplitudes in C. (G) *Left*, Individual cell data for
the PPR between the first two oIPSC amplitudes in the train; p=0.0206. R*ight, top*,
example current trace of short-term synaptic facilitation in VgAT+ neurons. *Right,*
bottom, example current traces of short-term synaptic depression in Vgat+ and Vglut2+
neurons. (H) Percent of pre-optical stimulation firing frequency [%Pre-Opto Frq] during
and post-stimulation in n=25 ChAT+, n=18 Vgat+, and n=29 Vglut2+ caudal PPN
neurons. (I) Individual cell data for the absolute change in frequency during stimulation
[ΔFrq During Opto]. (J) Correlation analysis for Vgat+ neurons, color scale representing
Spearman r [-1,1] and size representing p-value [1,0]. (K) Negative correlation between

the absolute change in frequency during stimulation and first oIPSC amplitude; r=-
0.627, p=0.044. (L) Correlation analysis for Vglut2+ neurons. (M) Negative correlation
between the absolute change in frequency during stimulation and the pre-stimulation
firing frequency; r=-0.648, p=0.014. *p<0.05, **p<0.01; box plots show median line with
boxes showing IQR and whiskers showing 9th and 91st percentiles.
1198
Figure 7. *In vivo* activation of GPe and SNr axons in the PPN show differential
effects on locomotion and opposite effects on valence. (A) Experimental set up to
stimulate ChR2-filled SNr or GPe axons over the PPN *in vivo*. (B) Representative image
of optical fiber tract overlaid with the approximate optical fiber placement for SNr-
[green] and GPe- [orange] injected mice. (C) Distance traveled over time in an open
field with 1 minute 20 Hz 4.25 mW optical stimulations over the PPN in N=9 control
(Ctrl) mice (black circles), N=8 mice injected with ChR2 in the SNr (green diamonds),
and N=9 mice injected with ChR2 in the GPe (orange hexagons); vertical blue lines
represent periods of optical stimulation. (D) Average distance traveled for each mouse
across the six 1-minute optical stimulations in the high (4.25 mW) and low laser power
(0.25 mW) conditions (E) Representative mouse track tracings during real time place
preference task in a three-chamber box and continuously stimulating EGFP- or ChR2-
filled axons over the PPN at 20 Hz in SNr- and GPe-injected mice when the mice are in
the stimulation zone. (F) Percent time spent in the stimulation zone in N=16 control
mice, N=9 mice injected with ChR2 in the SNr, and N=10 mice injected with ChR2 in the
GPe with 4.25 or 0.25 mW laser power. (G) Percent time spent in the stimulation zone
during the first minute reintroduced to the RTPP box on day 2 of RTPP with the laser off
(no chamber is stimulated) Black line = median. * p<0.05, ** p<0.01, *** p<0.001, ****
p<0.0001; box plots show median line with boxes showing IQR and whiskers showing
9th and 91st percentiles. See related supplemental Figures S2-S4 and Videos S1-S4.
Supplemental information
Document S1. Figure S1, related to Figures 2, 3, 4, & 6; Figure S2, S3, S4 related to Figure 7.
Document S2. Table S1, related to Figures 2, 3, 4, 5, & 6.
Video S1. Example video of stimulating SNr axons over the PPN in an open field, related to Figure 7
Video S2. Example video of stimulating GPe axons over the PPN in an open field, related to Figure 7
Video S3. Example video of stimulating SNr axons over the PPN in the striped zone, related to Figure 7
Video S4. Example video of stimulating GPe axons over the PPN in the striped zone, related to Figure 7

A.

B.

C. ChAT-Cre/Ai9-TdTomato

D. Vgat-Cre/Ai9-TdTomato

E. Vglut2-Cre/Ai9-TdTomato

1231

Figure S1. (A) Representative image of SNr injected with ChR2-EYFP. Cell bodies and axons filled with
ChR2 are represented in green. Post hoc staining of tyrosine hydroxylase (TH) to label dopamine
neurons in the substantia nigra *pars compacta*. TH+ neurons are represented in magenta. **(B)**
Representative image of GPe injected with ChR2-EYFP in a Vgat-Cre/Ai9-TdTomato mouse. Cell bodies
and axons filled with ChR2 are represented in green. Vgat+ neurons and their axonal projections are
represented in red. **(C-E)** Graphical depiction of virus spread in N=6 SNr and N=6 GPe injected mice in
each mouse line: **(C)** ChAT-Cre/Ai9-TdTomato, **(D)** Vgat-Cre/Ai9-TdTomato, and **(E)** Vglut2-Cre/Ai9-
TdTomato mice.
1240

Figure S2. Speed measurements across age and sex. (A) Speed of SNr-ChR2 mice during optical
stimulation in the open field. Individual mouse traces are shown as green broken lines, with the group
average represented by a solid green line with diamond markers. SNr-ChR2 mice exhibited increased
speed during stimulation. (B) Speed of GPe-ChR2 mice during six 1-minute optical stimulations in the
open field. Individual mouse traces are shown as orange broken lines, with the group average
represented by a solid orange line with hexagon markers. GPe-ChR2 mice exhibited a reduction in speed
during stimulation. Data are presented as mean ± SEM.(C) Distance traveled in the open field as a
function of age in SNr-ChR2 and control mice. Older SNr-ChR2 mice tend to travel further, whereas older
control mice travel less. Individual mouse traces are shown. (D) Comparison of locomotion between male
and female SNr-ChR2 mice in the open field, showing a trend toward increased movement in female mice
with SNr stimulation.

Figure S3. Opposite place preference despite similar motor responses to nigral axon stimulation in the PPN of DAT-Cre and wildtype mice.
(A) Distance traveled over time in an open field during 1-minute optical stimulations (20 Hz, 4.25 mW) over the pedunculopontine nucleus (PPN). Data are shown for **N = 6** DAT-Cre mice and wildtype mice injected with ChR2 in the substantia nigra. SNr data from Figure 7. Data are presented as mean ± SEM. (B) Average distance traveled for each mouse during stimulation across six 1-minute optical stimulations under high laser power (4.25 mW). (C) Average distance traveled for each mouse during stimulation across six 1-minute optical stimulations under low laser power (0.25 mW). (D) Representative movement traces from individual mice during a real-time place preference task in a three-chamber box. Mice received continuous optical stimulation (20 Hz) of EGFP- or ChR2-expressing axons over the PPN when in the stimulation zone. (E) Percentage of time spent in the stimulation zone under high laser power (4.25 mW) for **N = 9** wildtype and **N = 6** DAT-Cre mice injected with ChR2 in the substantia nigra. (F) Percentage of time spent in the stimulation zone under low laser power (0.25 mW) for the same groups.

Figure S4. Rostral vs caudal implant site alters SNr axon stimulation effects. (A) Implant location (same as Figure 7B) with most rostral (purple) and most caudal (red) bilateral implant pairs circled for SNr axon stimulaton. (B) Distance traveled in open field in mouse with rostral implants (purple) and mouse with caudal implants (red).

1313

1314 **Table 1. Statistical details of exact p-values and critical values for SNr input to ChAT+ PPN neurons**

| Parameter | Statistical Test | p-value | ChAT+ | |
			Rostral	Caudal
oIPSC Amplitude	unpaired t-test, t=0.9041, df=33	0.3725	n=15, 92.6 ± 18 pA	n=20, 115.2 ± 17 pA
PPR	unpaired t-test, t=0.1082, df=33	0.9145	n=15, 0.96 ± 0.04	n=20, 0.97 ± 0.05
Avg membrane potential	Mann Whitney, U=87	0.4706	n=11, -54.24 (-56.94 to -50.20) mV	n=19, -53.00 (-55.38 to -48.84) mV
% Pre-Opto Frq	Repeated Measures 2way ANOVA Stimulation: F(1, 35)=81.9 Region: F(1, 35)=0.002052 Interaction: F(1, 35)=3.273	<0.0001 0.9641 0.0790	n=14 During: 35.62 ± 8.869 % Post: 136.1 ± 7.803 %	n=23 During: 52.11 ± 8.432 % Post: 119.1 ± 4.614 %
ΔFrq During Opto	Mann Whitney, U=131	0.3602	n=14, -2.47 (-3.37 to -1.44) Hz	n=23, -2.10 (-3.11 to -0.84) Hz
ΔRebound Frq	Mann Whitney, U=79	0.0142	n=14, 1.64 (1.00 to 2.17) Hz	n=22, 0.83 (0.42 to 1.14) Hz

1315
1316

1317

1318

1319

1320 **Table 2. Statistical details of exact p-values and critical values for SNr input to Vgat+ PPN neurons**

| Parameter | Statistical Test | p-value | Vgat+ | |
			Rostral	Caudal
oIPSC Amplitude	Mann Whitney, U=50	0.6005	n=9, 76.5 (27.0 to 209.9) pA	n=13, 59.7 (37.7 to 269.7) pA
PPR	Mann Whitney, U=48	0.5123	n=9, 1.01 (0.88 to 1.16)	n=13, 1.06 (0.96 to 1.15);
Avg membrane potential	unpaired t-test, t=2.468, df=13	0.0282	n=5, -57.91 ± 1.630 mV	n=10, -52.00 ± 1.472 mV
% Pre-Opto Frq	Repeated Measures 2way ANOVA Stimulation: F(1, 24)=44.63 Region: F(1, 24)=0.09642 Interaction: F(1, 24)=0.8841	<0.0001 0.7588 0.3565	n=7 During: 37.91 ± 15.91 % Post: 100.0 ± 14.57 %	n=19 During: 35.62 ± 8.869 % Post: 113.2 ± 5.187 %
ΔFrq During Opto	Mann Whitney, U=55	0.5336	n=7, -5.03 (-5.55 to -1.34) Hz	n=19, -4.29 (-11.02 to -2.94) Hz
ΔRebound Frq	Mann Whitney, U=63	>0.9999	n=7, 0.80 (-1.31 to 2.33) Hz	n=18, 0.59 (-2.35 to 1.47) Hz

1321

Table 3. Statistical details of exact p-values and critical values for SNr input to Vglut2+ PPN neurons

Parameter	Statistical Test	p-value	Vglut2+	
			Rostral	Caudal
oIPSC Amplitude	Mann Whitney, U=29	0.0035	n=13, 61.6 (19.2 to 113.9) pA	n=13, 313.5 (143.8 to 729.1) pA
PPR	Mann Whitney, U=77	0.7241	n=13, 1.07 (0.86 to 1.38)	n=13, 0.92 (0.89 to 1.02)
Avg membrane potential	unpaired t-test, t=0.3838, df=25	0.7043	n=11, -53.03 ± 1.815	n=16, -52.09 ± 1.601
% Pre-Opto Frq	Repeated Measures 2way ANOVA Stimulation: F(1,28)=133.6 Region: F(1, 28)=0.5214 Interaction: F(1,28)=0.1765	 <0.0001 0.4762 0.6776	n=13 During: 32.62 ± 10.21 % Post: 100.3 ± 6.113 %	n=17 During: 24.71 ± 7.297 % Post: 97.48 ± 2.228 %
ΔFrq During Opto	Mann Whitney, U=55	0.0343	n=13, -3.21 (-12.21 to -1.95) Hz	n=17, -14.08 (-20.04 to -6.07) Hz
ΔRebound Frq	Mann Whitney, U=88	0.3627	n=13, 0.37 (-0.027 to 0.48) Hz	n=17, -0.23 (-0.82 to 1.08) Hz

Table 4. Statistical details of exact p-values and critical values for GPe input to ChAT+, Vgat+, and Vglut2+ PPN neurons

Parameter	Statistical Test	p-value	ChAT+	Vgat+	Vglut2+
oIPSC Amplitude	Kruskal-Wallis, H(2,37)=3.050	0.2176	n=6, 22.6 (9.75 to 50.20) pA	n=19, 24.77 (6.72 to 54.17) pA	n=15, 65.71 (14.50 to 133.4) pA
PPR	Mann Whitney, U=76	0.0206		n=19, 0.97 (0.76 to 1.2)	n=15, 0.81 (0.55 to 0.93
Avg membrane potential	Kruskal-Wallis, H(2,69)=2.963	0.2273	n=25, -54.16 (-58.05 to -51.10) mV	n=18, -54.28 (-55.47 to -48.77) mV	n=29, -55.32 (-57.97 to -52.28) mV
% Pre-Opto Frq	Repeated Measures 2way ANOVA Stimulation: F(1, 69)=25.46 Cell type: F(1, 69)=4.444 Interaction: F(1, 69)=2.805	 <0.0001 0.0153 0.0674	n=29 During: 92.39 ± 7.59 % Post: 104.10 ± 5.23 %	n=18 During: 57.70 ± 10.11 % Post: 91.81 ± 6.98 %	n=24 During: 64.15 ± 7.63 % Post: 104.7 ± 5.39 %
	Uncorrected Fisher's LSD, df=138 t=2.971 t=3.223 t=0.6177 t=0.1407 t=1.076 t=1.236	 0.0035 0.0016 0.5378 0.8883 0.2838 0.2184	During: ChAT+ vs. Vglut2+: 28.24 (CI 9.446 to 47.03) ChAT+ vs. Vgat+: 34.69 (CI 13.41 to 55.97) Vglut2+ vs. Vgat+: 6.454 (CI -14.21 to 27.11) Post: ChAT+ vs. Vglut2+: -1.337 (CI -20.13 to 17.45) ChAT+ vs. Vgat+: 11.58 (CI -9.702 to 32.86) Vglut2+ vs. Vgat+: 12.92 (CI -7.741 to 33.58)		
ΔFrq During Opto	Kruskal-Wallis, H(2,69)=17.21	0.0002	n=25, -0.16 (-0.90 to 0.03) Hz	n=18, -1.92 (-3.85 to -0.60) Hz	n=29, -0.89 (-3.28 to -0.43) Hz
	Uncorrected Dunn's test	0.0009 0.0002 0.4020	ChAT+ vs. Vglut2+: Z=3.332 ChAT+ vs. Vgat+: Z=3.755 Vglut2+ vs. Vgat+: Z=0.8380		
ΔRebound Frq	Kruskal-Wallis, H(2,69)=17.21	0.6772	n=8, 0.1041 (-0.52 to 0.41) Hz	n=19, 0.07082 (-0.36 to 0.57) Hz	n=7, -0.1421 (-2.58 to 0.71) Hz

References

1. Ryczko, D. (2022). The Mesencephalic Locomotor Region: Multiple Cell Types, Multiple Behavioral Roles, and Multiple Implications for Disease. Neuroscientist, 10738584221139136. https://doi.org/10.1177/10738584221139136.

2. Bastos-Gonçalves, R., Coimbra, B., and Rodrigues, A.J. (2024). The mesopontine tegmentum in reward and aversion: from cellular heterogeneity to behaviour. Neuroscience & Biobehavioral Reviews, 105702. https://doi.org/10.1016/j.neubiorev.2024.105702.

3. Mena-Segovia, J., and Bolam, J.P. (2017). Rethinking the Pedunculopontine Nucleus: From Cellular Organization to Function. Neuron *94*, 7–18. https://doi.org/10.1016/j.neuron.2017.02.027.

4. French, I.T., and Muthusamy, K.A. (2018). A Review of the Pedunculopontine Nucleus in Parkinson's Disease. Frontiers in Aging Neuroscience *10*, 99. https://doi.org/10.3389/fnagi.2018.00099.

5. Wang, H.-L., and Morales, M. (2009). Pedunculopontine and laterodorsal tegmental nuclei contain distinct populations of cholinergic, glutamatergic and GABAergic neurons in the rat. Eur J Neurosci *29*, 10.1111/j.1460-9568.2008.06576.x. https://doi.org/10.1111/j.1460-9568.2008.06576.x.

6. Mena-Segovia, J., Micklem, B. r., Nair-Roberts, R. g., Ungless, M. a., and Bolam, J. p. (2009). GABAergic neuron distribution in the pedunculopontine nucleus defines functional subterritories. Journal of Comparative Neurology *515*, 397–408. https://doi.org/10.1002/cne.22065.

7. Yoo, J.H., Zell, V., Wu, J., Punta, C., Ramajayam, N., Shen, X., Faget, L., Lilascharoen, V., Lim, B.K., and Hnasko, T.S. (2017). Activation of Pedunculopontine Glutamate Neurons Is Reinforcing. J Neurosci *37*, 38–46. https://doi.org/10.1523/JNEUROSCI.3082-16.2016.

8. Steinkellner, T., Yoo, J.H., and Hnasko, T.S. (2019). Differential Expression of VGLUT2 in Mouse Mesopontine Cholinergic Neurons. eNeuro *6*. https://doi.org/10.1523/ENEURO.0161-19.2019.

9. Martinez-Gonzalez, C., Bolam, J., and Mena-Segovia, J. (2011). Topographical Organization of the Pedunculopontine Nucleus. Frontiers in Neuroanatomy *5*, 22. https://doi.org/10.3389/fnana.2011.00022.

10. Alam, M., Schwabe, K., and Krauss, J.K. (2011). The pedunculopontine nucleus area: critical evaluation of interspecies differences relevant for its use as a target for deep brain stimulation. Brain *134*, 11–23. https://doi.org/10.1093/brain/awq322.

11. Marín, O., Smeets, W.J.A.J., and González, A. (1998). Evolution of the basal ganglia in tetrapods: a new perspective based on recent studies in amphibians.

Trends in Neurosciences *21*, 487–494. https://doi.org/10.1016/S0166-
2236(98)01297-1.

12. McElvain, L.E., Chen, Y., Moore, J.D., Brigidi, G.S., Bloodgood, B.L., Lim, B.K.,
Costa, R.M., and Kleinfeld, D. (2021). Specific populations of basal ganglia output
neurons target distinct brain stem areas while collateralizing throughout the
diencephalon. Neuron *109*, 1721-1738.e4.
https://doi.org/10.1016/j.neuron.2021.03.017.

13. Zhao, P., Wang, H., Li, A., Sun, Q., Jiang, T., Li, X., and Gong, H. (2022). The
Mesoscopic Connectome of the Cholinergic Pontomesencephalic Tegmentum.
Frontiers in Neuroanatomy *16*.

14. Roseberry, T.K., Lee, A.M., Lalive, A.L., Wilbrecht, L., Bonci, A., and Kreitzer, A.C.
(2016). Cell-Type-Specific Control of Brainstem Locomotor Circuits by Basal
Ganglia. Cell *164*, 526–537. https://doi.org/10.1016/j.cell.2015.12.037.

15. Huerta-Ocampo, I., Dautan, D., Gut, N.K., Khan, B., and Mena-Segovia, J. (2021).
Whole-brain mapping of monosynaptic inputs to midbrain cholinergic neurons. Sci
Rep *11*, 9055. https://doi.org/10.1038/s41598-021-88374-6.

16. Dautan, D., Kovács, A., Bayasgalan, T., Diaz-Acevedo, M.A., Pal, B., and Mena-
Segovia, J. (2021). Modulation of motor behavior by the mesencephalic locomotor
region. Cell Reports *36*. https://doi.org/10.1016/j.celrep.2021.109594.

17. Caggiano, V., Leiras, R., Goñi-Erro, H., Masini, D., Bellardita, C., Bouvier, J.,
Caldeira, V., Fisone, G., and Kiehn, O. (2018). Midbrain circuits that set locomotor
speed and gait selection. Nature *553*, 455–460.
https://doi.org/10.1038/nature25448.

18. Henrich, M.T., Geibl, F.F., Lakshminarasimhan, H., Stegmann, A., Giasson, B.I.,
Mao, X., Dawson, V.L., Dawson, T.M., Oertel, W.H., and Surmeier, D.J. (2020).
Determinants of seeding and spreading of α-synuclein pathology in the brain. Sci
Adv *6*. https://doi.org/10.1126/sciadv.abc2487.

19. Ugolini, G. (2011). Chapter 10 - Rabies Virus as a Transneuronal Tracer of
Neuronal Connections. In Advances in Virus Research Research Advances in
Rabies., A. C. Jackson, ed. (Academic Press), pp. 165–202.
https://doi.org/10.1016/B978-0-12-387040-7.00010-X.

20. Linders, L.E., Supiot, Laura.F., Du, W., D'Angelo, R., Adan, R.A.H., Riga, D., and
Meye, F.J. (2022). Studying Synaptic Connectivity and Strength with Optogenetics
and Patch-Clamp Electrophysiology. Int J Mol Sci *23*, 11612.
https://doi.org/10.3390/ijms231911612.

21. Glasgow, S.D., McPhedrain, R., Madranges, J.F., Kennedy, T.E., and Ruthazer,
E.S. (2019). Approaches and Limitations in the Investigation of Synaptic

Transmission and Plasticity. Front. Synaptic Neurosci. *11*.
https://doi.org/10.3389/fnsyn.2019.00020.

22. Goñi-Erro, H., Selvan, R., Leiras, R., and Kiehn, O. (2023). Pedunculopontine
Chx10+ neurons control global motor arrest in mice. Nat Neurosci, 1–13.
https://doi.org/10.1038/s41593-023-01396-3.

23. Masini, D., and Kiehn, O. (2022). Targeted activation of midbrain neurons restores
locomotor function in mouse models of parkinsonism. Nat Commun *13*, 504.
https://doi.org/10.1038/s41467-022-28075-4.

24. Gut, N.K., Yilmaz, D., Kondabolu, K., Huerta-Ocampo, I., and Mena-Segovia, J.
(2022). Selective inhibition of goal-directed actions in the mesencephalic locomotor
region (Neuroscience) https://doi.org/10.1101/2022.01.18.476772.

25. Huang, Y., Wang, S., Wang, Q., Zheng, C., Yang, F., Wei, L., Zhou, X., and Wang,
Z. (2024). Glutamatergic Circuits in the Pedunculopontine Nucleus Modulate
Multiple Motor Functions. Neurosci. Bull. https://doi.org/10.1007/s12264-024-01314-
y.

26. Josset, N., Roussel, M., Lemieux, M., Lafrance-Zoubga, D., Rastqar, A., and
Bretzner, F. (2018). Distinct Contributions of Mesencephalic Locomotor Region
Nuclei to Locomotor Control in the Freely Behaving Mouse. Current Biology *28*, 884-
901.e3. https://doi.org/10.1016/j.cub.2018.02.007.

27. Tello, A.J., Zouwen, C.I. van der, Dejas, L., Duque-Yate, J., Boutin, J., Medina-Ortiz,
K., Suresh, J.S., Swiegers, J., Sarret, P., and Ryczko, D. (2024). Dopamine-
sensitive neurons in the mesencephalic locomotor region control locomotion
initiation, stop, and turns. Cell Reports *43*.
https://doi.org/10.1016/j.celrep.2024.114187.

28. Xiao, C., Cho, J.R., Zhou, C., Treweek, J.B., Chan, K., McKinney, S.L., Yang, B.,
and Gradinaru, V. (2016). Cholinergic Mesopontine Signals Govern Locomotion and
Reward through Dissociable Midbrain Pathways. Neuron *90*, 333–347.
https://doi.org/10.1016/j.neuron.2016.03.028.

29. Dautan, D., Souza, A.S., Huerta-Ocampo, I., Valencia, M., Assous, M., Witten, I.B.,
Deisseroth, K., Tepper, J.M., Bolam, J.P., Gerdjikov, T.V., et al. (2016). Segregated
cholinergic transmission modulates dopamine neurons integrated in distinct
functional circuits. Nat Neurosci *19*, 1025–1033. https://doi.org/10.1038/nn.4335.

30. Ferreira-Pinto, M.J., Kanodia, H., Falasconi, A., Sigrist, M., Esposito, M.S., and
Arber, S. (2021). Functional diversity for body actions in the mesencephalic
locomotor region. Cell *184*, 4564-4578.e18.
https://doi.org/10.1016/j.cell.2021.07.002.

31. Zhang, S., Mena-Segovia, J., and Gut, N.K. (2024). Inhibitory Pedunculopontine
Neurons Gate Dopamine-Mediated Motor Actions of Unsigned Valence. Curr

Neuropharmacol *22*, 1540–1550.
https://doi.org/10.2174/1570159X21666230911103520.

32. Hormigo, S., Vega-Flores, G., Rovira, V., and Castro-Alamancos, M.A. (2019).
Circuits That Mediate Expression of Signaled Active Avoidance Converge in the
Pedunculopontine Tegmentum. J. Neurosci. *39*, 4576–4594.
https://doi.org/10.1523/JNEUROSCI.0049-19.2019.

33. Hormigo, S., Shanmugasundaram, B., Zhou, J., and Castro-Alamancos, M.A.
(2021). A Signaled Locomotor Avoidance Action Is Fully Represented in the Neural
Activity of the Midbrain Tegmentum. J. Neurosci. *41*, 4262–4275.
https://doi.org/10.1523/JNEUROSCI.0027-21.2021.

34. Rizzi, G., and Tan, K.R. (2019). Synergistic Nigral Output Pathways Shape
Movement. Cell Reports *27*, 2184-2198.e4.
https://doi.org/10.1016/j.celrep.2019.04.068.

35. Liu, A., Cheng, Y., and Huang, J. (2023). Neurons innervating both the central
amygdala and the ventral tegmental area encode different emotional valences. Front
Neurosci *17*, 1178693. https://doi.org/10.3389/fnins.2023.1178693.

36. Galaj, E., Han, X., Shen, H., Jordan, C.J., He, Y., Humburg, B., Bi, G.-H., and Xi, Z.-
X. (2020). Dissecting the Role of GABA Neurons in the VTA versus SNr in Opioid
Reward. J Neurosci *40*, 8853–8869. https://doi.org/10.1523/JNEUROSCI.0988-
20.2020.

37. Mastro, K.J., Bouchard, R.S., Holt, H.A.K., and Gittis, A.H. (2014). Transgenic
Mouse Lines Subdivide External Segment of the Globus Pallidus (GPe) Neurons
and Reveal Distinct GPe Output Pathways. J. Neurosci. *34*, 2087–2099.
https://doi.org/10.1523/JNEUROSCI.4646-13.2014.

38. Tian, J., Yan, Y., Xi, W., Zhou, R., Lou, H., Duan, S., Chen, J.F., and Zhang, B.
(2018). Optogenetic Stimulation of GABAergic Neurons in the Globus Pallidus
Produces Hyperkinesia. Front Behav Neurosci *12*, 185.
https://doi.org/10.3389/fnbeh.2018.00185.

39. Pamukcu, A., Cui, Q., Xenias, H.S., Berceau, B.L., Augustine, E.C., Fan, I.,
Chalasani, S., Hantman, A.W., Lerner, T.N., Boca, S.M., et al. (2020). Parvalbumin+
and Npas1+ Pallidal Neurons Have Distinct Circuit Topology and Function. J
Neurosci *40*, 7855–7876. https://doi.org/10.1523/JNEUROSCI.0361-20.2020.

40. Cui, Q., Pamukcu, A., Cherian, S., Chang, I.Y.M., Berceau, B.L., Xenias, H.S.,
Higgs, M.H., Rajamanickam, S., Chen, Y., Du, X., et al. (2021). Dissociable Roles of
Pallidal Neuron Subtypes in Regulating Motor Patterns. J. Neurosci. *41*, 4036–4059.
https://doi.org/10.1523/JNEUROSCI.2210-20.2021.

41. Aristieta, A., Barresi, M., Azizpour Lindi, S., Barrière, G., Courtand, G., de la
Crompe, B., Guilhemsang, L., Gauthier, S., Fioramonti, S., Baufreton, J., et al.

(2021). A Disynaptic Circuit in the Globus Pallidus Controls Locomotion Inhibition.
Current Biology *31*, 707-721.e7. https://doi.org/10.1016/j.cub.2020.11.019.

42. Isett, B.R., Nguyen, K.P., Schwenk, J.C., Yurek, J.R., Snyder, C.N., Vounatsos,
M.V., Adegbesan, K.A., Ziausyte, U., and Gittis, A.H. (2023). The indirect pathway of
the basal ganglia promotes transient punishment but not motor suppression.
Neuron. https://doi.org/10.1016/j.neuron.2023.04.017.

43. Susaki, E.A., Tainaka, K., Perrin, D., Yukinaga, H., Kuno, A., and Ueda, H.R.
(2015). Advanced CUBIC protocols for whole-brain and whole-body clearing and
imaging. Nat Protoc *10*, 1709–1727. https://doi.org/10.1038/nprot.2015.085.

44. Ambrosi, P., and Lerner, T.N. (2022). Striatonigrostriatal circuit architecture for
disinhibition of dopamine signaling. Cell Reports *40*.
https://doi.org/10.1016/j.celrep.2022.111228.

45. Buchholz, M.O., Guilabert, A.G., Ehret, B., and Schuhknecht, G.F.P. (2023). How
synaptic strength, short-term plasticity, and input synchrony contribute to neuronal
spike output. PLOS Computational Biology *19*, e1011046.
https://doi.org/10.1371/journal.pcbi.1011046.

46. Chen, R.Y.-T., and Evans, R.C. (2024). Comparing tonic and phasic dendritic
calcium in cholinergic pedunculopontine neurons and dopaminergic substantia nigra
neurons. European Journal of Neuroscience *59*, 1638–1656.
https://doi.org/10.1111/ejn.16281.

47. Tubert, C., Zampese, E., Pancani, T., Tkatch, T., and Surmeier, D.J. (2023). Feed-
forward metabotropic signaling by Cav1 Ca2+ channels supports pacemaking in
pedunculopontine cholinergic neurons. Neurobiology of Disease *188*, 106328.
https://doi.org/10.1016/j.nbd.2023.106328.

48. Takakusaki, K., Shiroyama, T., and Kitai, S.T. (1997). Two types of cholinergic
neurons in the rat tegmental pedunculopontine nucleus: electrophysiological and
morphological characterization. Neuroscience *79*, 1089–1109.
https://doi.org/10.1016/S0306-4522(97)00019-5.

49. Baksa, B., Kovács, A., Bayasgalan, T., Szentesi, P., Kőszeghy, Á., Szücs, P., and
Pál, B. (2019). Characterization of functional subgroups among genetically identified
cholinergic neurons in the pedunculopontine nucleus. Cell. Mol. Life Sci. *76*, 2799–
2815. https://doi.org/10.1007/s00018-019-03025-4.

50. Kroeger, D., Ferrari, L.L., Petit, G., Mahoney, C.E., Fuller, P.M., Arrigoni, E., and
Scammell, T.E. (2017). Cholinergic, Glutamatergic, and GABAergic Neurons of the
Pedunculopontine Tegmental Nucleus Have Distinct Effects on Sleep/Wake
Behavior in Mice. J Neurosci *37*, 1352–1366.
https://doi.org/10.1523/JNEUROSCI.1405-16.2016.

51. Granata, A.R., and Kitai, S.T. (1991). Inhibitory substantia nigra inputs to the
pedunculopontine neurons. Exp Brain Res *86*, 459–466.
https://doi.org/10.1007/BF00230520.

52. Kang, Y., and Kitai, S.T. (1990). Electrophysiological properties of pedunculopontine
neurons and their postsynaptic responses following stimulation of substantia nigra
reticulata. Brain Research *535*, 79–95. https://doi.org/10.1016/0006-8993(90)91826-
3.

53. Evans, R.C., Zhu, M., and Khaliq, Z.M. (2017). Dopamine Inhibition Differentially
Controls Excitability of Substantia Nigra Dopamine Neuron Subpopulations through
T-Type Calcium Channels. J. Neurosci. *37*, 3704–3720.
https://doi.org/10.1523/JNEUROSCI.0117-17.2017.

54. Villalobos, C.A., and Basso, M.A. (2022). Optogenetic activation of the inhibitory
nigro-collicular circuit evokes contralateral orienting movements in mice. Cell Rep
39, 110699. https://doi.org/10.1016/j.celrep.2022.110699.

55. Rajaram, E., Kaltenbach, C., Fischl, M.J., Mrowka, L., Alexandrova, O., Grothe, B.,
Hennig, M.H., and Kopp-Scheinpflug, C. (2019). Slow NMDA-Mediated Excitation
Accelerates Offset-Response Latencies Generated via a Post-Inhibitory Rebound
Mechanism. eNeuro *6*. https://doi.org/10.1523/ENEURO.0106-19.2019.

56. Zhu, T., Wei, S., and Wang, Y. (2022). Post-Inhibitory Rebound Firing of Dorsal
Root Ganglia Neurons. J Pain Res *15*, 2029–2040.
https://doi.org/10.2147/JPR.S370335.

57. Beaver, M.L., and Evans, R.C. (2025). Muscarinic receptor activation preferentially
inhibits rebound in vulnerable dopaminergic neurons. J. Neurosci.
https://doi.org/10.1523/JNEUROSCI.1443-24.2025.

58. Sivaramakrishnan, S., and Lynch, W.P. (2017). Rebound from Inhibition: Self-
Correction against Neurodegeneration? J Clin Cell Immunol *8*, 492.
https://doi.org/10.4172/2155-9899.1000492.

59. Bordas, C., Kovacs, A., and Pal, B. (2015). The M-current contributes to high
threshold membrane potential oscillations in a cell type-specific way in the
pedunculopontine nucleus of mice. Frontiers in Cellular Neuroscience *9*.

60. Aitta-aho, T., Hay, Y.A., Phillips, B.U., Saksida, L.M., Bussey, T.J., Paulsen, O., and
Apergis-Schoute, J. (2018). Basal Forebrain and Brainstem Cholinergic Neurons
Differentially Impact Amygdala Circuits and Learning-Related Behavior. Current
Biology *28*, 2557-2569.e4. https://doi.org/10.1016/j.cub.2018.06.064.

61. Dong, J., Hawes, S., Wu, J., Le, W., and Cai, H. (2021). Connectivity and
Functionality of the Globus Pallidus Externa Under Normal Conditions and
Parkinson's Disease. Frontiers in Neural Circuits *15*.

62. Lilascharoen, V., Wang, E.H.-J., Do, N., Pate, S.C., Tran, A.N., Yoon, C.D., Choi, J.-
H., Wang, X.-Y., Pribiag, H., Park, Y.-G., et al. (2021). Divergent pallidal pathways
underlying distinct Parkinsonian behavioral deficits. Nat Neurosci *24*, 504–515.
https://doi.org/10.1038/s41593-021-00810-y.

63. Dodson, P.D., Larvin, J.T., Duffell, J.M., Garas, F.N., Doig, N.M., Kessaris, N.,
Duguid, I.C., Bogacz, R., Butt, S.J.B., and Magill, P.J. (2015). Distinct
developmental origins manifest in the specialized encoding of movement by adult
neurons of the external globus pallidus. Neuron *86*, 501–513.
https://doi.org/10.1016/j.neuron.2015.03.007.

64. Turner, R.S., and Anderson, M.E. (1997). Pallidal Discharge Related to the
Kinematics of Reaching Movements in Two Dimensions. Journal of Neurophysiology
77, 1051–1074. https://doi.org/10.1152/jn.1997.77.3.1051.

65. Kita, H., and Kita, S.T. (1994). The morphology of globus pallidus projection neurons
in the rat: an intracellular staining study. Brain Research *636*, 308–319.
https://doi.org/10.1016/0006-8993(94)91030-8.

66. Sadek, A.R., Magill, P.J., and Bolam, J.P. (2007). A Single-Cell Analysis of Intrinsic
Connectivity in the Rat Globus Pallidus. J. Neurosci. *27*, 6352–6362.
https://doi.org/10.1523/JNEUROSCI.0953-07.2007.

67. Mastro, K.J., Zitelli, K.T., Willard, A.M., Leblanc, K.H., Kravitz, A.V., and Gittis, A.H.
(2017). Cell-specific pallidal intervention induces long-lasting motor recovery in
dopamine-depleted mice. Nat Neurosci *20*, 815–823.
https://doi.org/10.1038/nn.4559.

68. Liu, D., Li, W., Ma, C., Zheng, W., Yao, Y., Tso, C.F., Zhong, P., Chen, X., Song,
J.H., Choi, W., et al. (2020). A common hub for sleep and motor control in the
substantia nigra. Science *367*, 440–445. https://doi.org/10.1126/science.aaz0956.

69. Ryczko, D., Cone, J.J., Alpert, M.H., and Dubuc, R. (2016). A descending dopamine
pathway conserved from basal vertebrates to mammals. PNAS *113*.
https://doi.org/10.1073/pnas.1600684113.

70. da Silva, J.A., Tecuapetla, F., Paixão, V., and Costa, R.M. (2018). Dopamine neuron
activity before action initiation gates and invigorates future movements. Nature *554*,
244–248. https://doi.org/10.1038/nature25457.

71. Baker, M., Kang, S., Hong, S.-I., Song, M., Yang, M.A., Peyton, L., Essa, H., Lee,
S.W., and Choi, D.-S. (2023). External globus pallidus input to the dorsal striatum
regulates habitual seeking behavior in male mice. Nat Commun *14*, 4085.
https://doi.org/10.1038/s41467-023-39545-8.

72. Farries, M.A., Faust, T.W., Mohebi, A., and Berke, J.D. (2023). Selective encoding
of reward predictions and prediction errors by globus pallidus subpopulations.
Current Biology *0*. https://doi.org/10.1016/j.cub.2023.08.042.

73. Arkadir, D., Morris, G., Vaadia, E., and Bergman, H. (2004). Independent Coding of
Movement Direction and Reward Prediction by Single Pallidal Neurons. J. Neurosci.
24, 10047–10056. https://doi.org/10.1523/JNEUROSCI.2583-04.2004.

74. Forster, G.L., and Blaha, C.D. (2003). Pedunculopontine tegmental stimulation
evokes striatal dopamine efflux by activation of acetylcholine and glutamate
receptors in the midbrain and pons of the rat. Eur J Neurosci *17*, 751–762.
https://doi.org/10.1046/j.1460-9568.2003.02511.x.

75. Estakhr, J., Abazari, D., Frisby, K., McIntosh, J.M., and Nashmi, R. (2017).
Differential Control of Dopaminergic Excitability and Locomotion by Cholinergic
Inputs in Mouse Substantia Nigra. Current Biology *27*, 1900-1914.e4.
https://doi.org/10.1016/j.cub.2017.05.084.

76. Galtieri, D.J., Estep, C.M., Wokosin, D.L., Traynelis, S., and Surmeier, D.J. (2017).
Pedunculopontine glutamatergic neurons control spike patterning in substantia nigra
dopaminergic neurons. eLife *6*, e30352. https://doi.org/10.7554/eLife.30352.

77. Futami, T., Takakusaki, K., and Kitai, S.T. (1995). Glutamatergic and cholinergic
inputs from the pedunculopontine tegmental nucleus to dopamine neurons in the
substantia nigra pars compacta. Neuroscience Research *21*, 331–342.
https://doi.org/10.1016/0168-0102(94)00869-H.

78. Good, C.H., and Lupica, C.R. (2009). Properties of distinct ventral tegmental area
synapses activated via pedunculopontine or ventral tegmental area stimulation in
vitro. J Physiol *587*, 1233–1247. https://doi.org/10.1113/jphysiol.2008.164194.

79. Evans, R.C., Twedell, E.L., Zhu, M., Ascencio, J., Zhang, R., and Khaliq, Z.M.
(2020). Functional Dissection of Basal Ganglia Inhibitory Inputs onto Substantia
Nigra Dopaminergic Neurons. Cell Reports *32*, 108156.
https://doi.org/10.1016/j.celrep.2020.108156.

1658