Decoding critical targets and signaling pathways in EBV-mediated diseases

using large language models

Jingwen Yu[1], Yaohao Wang[1], Haidong Wang[1], Zhi Wei[2], Yonggang Pei[1, *]

[1] School of Public Health and Emergency Management, Southern University of

Science and Technology, Shenzhen, Guangdong, China

[2] Department of Computer Science, New Jersey Institute of Technology, Newark, NJ,

USA

Running title: Utilizing large language models to analyze EBV-mediated diseases

Keywords: EBV, Antigens, Pathways, LLMs

Corresponding author:

Yonggang Pei Ph.D.

Assistant professor

School of Public Health and Emergency Management

Southern University of Science and Technology

Email: peiyg@sustech.edu.cn

Tel: 86-0755-88015615

Abstract

Epstein-Barr virus (EBV), a member of the gamma-herpesvirus, is the first identified human oncovirus and is associated with various malignancies. Understanding the intricate interactions between EBV antigens and cellular pathways is crucial to unravel the molecular mechanisms in EBV-mediated diseases. However, fully elucidating EBV-host interactions and the associated pathogenesis remains a significant challenge. In this study, we utilized LLMs to screen 36,105 EBV-relevant scientific publications, presenting a descriptive overview of the interactions between EBV antigens and host cellular pathways through comprehensive interaction networks. We described the critical roles of EBV antigens by constructing functional subsets of host proteins targeted by EBV antigens and illustrated the interactions using protein-protein interaction (PPI) networks. Furthermore, we developed an antigen-pathway network that highlights the connections with EBV-associated diseases, including DLBCL, BL, NPC, gastric cancer (GC), and post-transplant lymphoproliferative disorders (PTLD). Utilizing our dataset and public dataset, we validated the efficacy of using BLLF3-targeted TLR2-associated factors as marker genes for DLBCL. Next, we confirmed the co-expression of calcium pathway factors induced by LMP1 over-expression in BL. Finally, based on shared results suggesting that LMP1 actively regulates the glycolysis pathway, we identified and validated the correlation and co-expression of LMP1-induced PARP1, HIF1A, HOXB3, and key transcription-related factors, revealing the complete picture of LMP1's influence on the glycolysis pathway. Our study presents a comprehensive functional encyclopedia of the interactions

between EBV antigens and host signaling pathways across various EBV-associated disease contexts, providing valuable insights for the development of therapeutic strategies.

Introduction

Epstein-Barr virus (EBV) was discovered in 1964 and has infected over 90% of the world's population[1]. The oncovirus contributes to various human diseases, such as Burkitt lymphoma (BL), nasopharyngeal carcinoma (NPC), Hodgkin lymphoma (HL), diffuse large B cell lymphoma (DLBCL), multiple sclerosis (MS), and so on. In the 60 years since its discovery, the intricate roles of EBV in regulating interconnected host cellular pathways have provided valuable insights into the mechanisms underlying EBV-related disease progression. EBV infection in human cells triggers a series of changes in cellular biology including cell cycle transition, cell apoptosis, DNA mismatch repair, and multiple cellular signaling pathways. Such modulations are mediated by various EBV antigens, promoting viral latency and oncogenesis. For instance, EBV-encoded Latent Membrane Protein 1 (LMP1) mimics CD40 signaling via its two C-terminal activating regions, namely CTAR1 and CTAR2, which are shown to activate both the canonical and noncanonical NF-κB pathways[2–5]. EBV nuclear antigen 1 (EBNA1) inhibits the canonical NF-κB pathway by inhibiting the phosphorylation of IKKα/β[6]. Although the NF-κB pathway serves as a key factor in EBV latent infection and oncogenesis, the intricate and intertwined network of EBV antigens and NF-κB factors presents a significant challenge for the development of

anti-EBV therapeutic strategies. Therefore, it is imperative to comprehensively

examine the landscape of current studies on EBV antigens and their interactions with

cellular signaling pathways to further explore the EBV-mediated pathogenesis.

Traditional literature mapping techniques, including systematic reviews and

meta-analyses, often fall short of capturing the full breadth of EBV literature.

However, the rapid development in machine learning offers promising tools for

analyzing potential key factors in EBV-mediated pathogenesis from vast amounts of

literature. A recent study characterizes the biomedical literature into a 2D map[7]. The

embedding contains 21 million papers, presenting the unlimited possibilities of

LLMs-assisted literature study. Another work introduced a knowledge graph of the

m6A regulatory pathway utilizing GPT-4 to elucidate m6A's regulatory mechanisms

in cancer phenotypes across various cancers[8]. This burgeoning technology has

catalyzed a proliferation of studies exploring its capabilities and applications in the

fields of biological and medical research, driving innovation and prompting ongoing

discourse on its potential impacts and future developments.

Our study aims to utilize LLMs for the first time to analyze decades of research

on how EBV antigens regulate cellular factors and signaling pathways. We present a

comprehensive framework of EBV antigens, their interacting host targets, and

associated pathways, constructing an extensive network that delineates virus-host

interaction. We then mapped the diverse background of EBV-related diseases for

antigen-pathway relationships and revealed the critical interactions between EBV

antigens and cellular pathways in EBV-mediated diseases. Collectively, this study

offers a unique overview of current EBV-host interactions and provides valuable

resources for the development of anti-EBV therapeutic strategies in the future.

Methods

Data sources

The data in this study were obtained from the PubMed database

(https://pubmed.ncbi.nlm.nih.gov/) using EDirect. The initial data collection was

performed on February 28, 2024. The final dataset comprises 36,105 EBV-related

articles, including only original articles and reviews published in English. The RNA-

seq dataset employed to investigate differentially expressed genes induced by LMP1

overexpression was obtained from the Gene Expression Omnibus (GEO) database

under accession number GSE247495. Differential gene expression analysis was

conducted using the R package DESeq2 (1.40.2). Genes with an adjusted p-value

(padj) less than 0.05 and a log2 fold change greater than log2(2) were considered

significantly differentially expressed.

ChatGPT-4 prompt engineering

To ensure accuracy, we provided ChatGPT with the lists of EBV antigens and

EBV-related diseases through system messages. The prompt for antigen-target-

pathway summary is as follows: "If the abstract concludes that EBV-encoded proteins

affect another protein or pathway, show me only the EBV protein, affected protein

and pathway. If not, return 'Error' and nothing else. Use a temperature of 0. All EBV

proteins were listed before. The abstract: XXX". The prompt for EBV-related disease

mapping: "Return which specific EBV-related disease does the abstract attached in the

end investigates. If unknown, return 'Error' and nothing else. The abstract: XXX".

To minimize costs, we initially utilized the free version GPT-3.5 Turbo via the

OpenAI API. Items not resulting in "Error" were then processed using GPT-4. Next,

we employed synonym substitution to standardize the naming of various aliases. For

instance, "EBV nuclear antigen 1", "EBNA1" and "EBNA-1" are all annotated as

"EBNA1"; "EB2", "Mta", "SM" and "BS-MLF1" are all annotated as "BMLF1".

(Table S1)

GPT-4 generated data evaluation

To evaluate the accuracy of the GPT-generated results, we randomly selected a

total of 115 entries for manual inspection. The inspection focused on the EBV

antigens mentioned in the corresponding articles, the host factors associated with

these antigens, and the annotated KEGG pathways. The criteria for a "correct result"

are as follows: (1) the article clearly describes the interaction between the EBV

antigen and its corresponding host factors; (2) the article identifies the biological

changes influenced by this interaction and their relevance to the annotated KEGG

results.

Annotation from GPT summarized pathways to KEGG database

The results generated by LLMs are highly variable. To establish a unique

standard for readability, we annotated the LLM-generated pathway outputs to the

KEGG pathway database due to its simplicity, accessibility, and widespread use

within the academic community. This mapping was performed based on our

customized R script and empirical manual substitution. A Sankey diagram shows a

complete mapping of the substitution (Table S21).

Gene enrichment analysis

LLM-acquired target host protein annotation is performed using the R package

STRINGdb (2.12.1). Annotated IDs are submitted to UniProt for symbol conversion

(https://www.uniprot.org/). The generated list of genes of interest (GOIs) was

subjected to gene enrichment analysis using three primary tools: the KEGG database[9],

Reactome (https://reactome.org/), and STRINGdb.

Omics analysis

PCA reduction and cohort analyses are performed using UCSC Xena[10]

(https://xena.ucsc.edu/) and GEPIA2 (http://gepia2.cancer-pku.cn/)[11]. The dataset

comprised TCGA Diffuse Large B-Cell Lymphoma, GTEx Spleen and GTEx Blood.

LMP1 overexpression RNA-seq data is normalized and analyzed using DESeq2.

Visualization

The figures were created with BioRender (https://biorender.com), the R package

wordcloud2 (0.2.1), circlize (0.4.16), or pheatmap (1.0.12) with customized scripts.

PPI network is visualized in Cytoscape (3.9.1) and colored based on protein family

types. Other visualizations are made by a combination of R packages including

ggplot2 (3.5.0), ggalluvial (0.12.5), and enrichplot (1.20.3).

Results

LLMs-assisted data mining reveals the key insights in EBV research

To create a literature pool for information extraction, our initial dataset comprised

36,105 EBV-related research articles available in the PubMed database as of the start

of this project (February 28, 2024). These publications span a temporal range of 57

years, from 1968 to 2024 (Figure 1a). Overall, there has been a steady increase in the

number of research articles related to EBV over these years, with an average of

approximately 1000 publications annually.

Next, to better characterize the current EBV studies, we summarized the available

articles into a table of PubMed ID, EBV antigens, target proteins, and the related

signaling pathways using ChatGPT-4. The returned dataset contained information

from 2,827 publications related to EBV antigen-pathway interactions and was

subsequently subjected to rigorous data reduction and filtering (Figure 1b). The final

dataset includes 2,289 documented entries of EBV antigens and their associated

pathways, derived from 1,839 publications (Figure 1b, Table S3). To validate the

accuracy of the information extracted using ChatGPT, we manually compared a

random subset of 115 entries (~5%) of the GPT-4 generated summaries with the

corresponding original abstracts. The results indicate that our prompts achieved an

average accuracy of 93.04% (Figure 1c, Table S4), demonstrating that our GPT-4-

based in-house pipeline can effectively analyze additional literature.

Analysis of EBV antigens and their impacts on cellular pathways reveals that

LMP1 is the most frequently studied antigen, followed by EBNA1, BZLF1, LMP2A,

and EBNA2, which are primarily latent or lytic proteins (Figure 1d). Throughout the

2010s, the significance of EBNA3C has increased, making it the fourth most studied

antigen (Figure 1e). The most studied pathways focus on the mechanisms underlying

EBV infection (Figure 1f). Pathways involving EBV-mediated cell apoptosis are the

next most thoroughly studied areas. Abundant studies concentrate on the oncogenic

mechanism of EBV infection, mainly on how EBV antigens regulate cell cycle and

replication. The most extensively studied mechanisms include the NF-κB signaling

pathway, MAPK signaling pathway, TLR signaling pathway, PI3K/Akt signaling

pathway, TNF signaling pathway, and TGF-β signaling pathway (Figure 1f). On the

temporal scale, research on cell apoptosis, NF-κB signaling pathway, cell cycle, and

MAPK signaling pathway have always been the four unyielding topics (Figure 1g),

suggesting their critical roles in EBV-mediated diseases.

Pathway mapping reveals distinct signatures of EBV antigens

To clarify the critical pathways that EBV antigens regulate, we manually

annotated the GPT-generated pathways to the KEGG pathway entries (Table S2). The

chord diagram denotes the frequency of top 33 most studied EBV antigens and

correlated pathways, indicating an intricate and interconnected network of cellular

pathways induced by EBV infection (Figure 2a). To clearly illustrate the concentrated

interactions between EBV antigens and their related pathways, we present a chord

diagram as a heatmap (Figure 2b). In this figure, the pathways are clustered using the

maximum linkage clustering method, and EBV antigens are arranged empirically

based on their roles in EBV latent or lytic infection. We observed that EBV latent

membrane proteins, LMP1 and LMP2A, exhibit similar interaction patterns across

different host cellular pathways, although research on LMP1 has been more thorough

as previously discussed (Figure 1d). The resemblance of these two EBV antigens lies

in the shared role in their ability to reprogram host immune response through the

MAPK, PI3K/Akt, or TNF pathways, and their oncogenic functions in regulating the

p53 or PD1/PD-L1 checkpoint pathway. Despite the significant scarcity of LMP2A-

related publications compared to LMP1, the studies collectively indicate that LMP2A

contributes more to the BCR signaling pathway and the Wnt pathway than LMP1

(Figure 2b).

EBV nuclear antigens (EBNAs) contribute to the establishment and maintenance

of viral latency, especially EBNA1, which is present in all EBV latent programs

except for latency 0. EBNA1 directly binds to the host genome and is essential for the

replication and maintenance of EBV episome[1]. It also serves as a transcription factor

regulating multiple gene expression. Our analysis shows that EBNA1 differs from

other EBNAs in terms of regulating DNA replication, and oxidative phosphorylation.

EBNA2 plays a crucial role in the initial stage of EBV-mediated B-cell transformation.

Interestingly, EBNA2 has been confirmed to bind to super-enhancer sites that are

critical for the maintenance of EBV latency[12,13]. Our findings highlight the distinctive

role of EBNA2 in cAMP signaling pathways, fatty acid biosynthesis, and cell

adhesion lipid molecules, indicating its potential to facilitate B-cell transformation

and proliferation through lipid metabolism. Similar to EBNA1, EBNA3A and

EBNA3C are shown to interfere with the host genome and promote immune evasion,

cell cycle control, and inhibition of apoptosis as transcription factors. EBNALP binds

to EBNA2 and functions as a coactivator to promote the transformation of B-cells

during EBV infection[1]. For immediate-early lytic genes that are responsible for EBV

reactivation, both BZLF1 and BRLF1 are well-studied. BZLF1 activates lytic gene

expression, initiating the lytic cycle, while BRLF1 synergizes with BZLF1 to promote

lytic gene transcription, leading to viral replication and release. Studies on BZLF1

seem more diverse on multiple cellular oncogenic pathways.

Other EBV antigens have received relatively less attention compared to the

previously mentioned latent and lytic antigens (Table S5). EBV glycoproteins play

roles in infection and entry into host cells. Glycoprotein L (gL) cooperates with gH

and gB to form the infection complex, mediating EBV-cell fusion and entry.

Gene enrichment uncover defining characteristics of EBV-induced cellular

modulations

To further investigate the associated diseases or pathways from the level of target

proteins, we annotated the EBV research literature on a higher resolution after

enrichment of GPT-generated target proteins using the KEGG database and the

Reactome database. Disease-wise, KEGG enrichment revealed that EBV infection

may be associated with traits typical of a wide range of diseases, including bladder

cancer, pancreatic cancer, colorectal cancer, small cell lung cancer, and others (Figure

3a). The results suggest a possible correlation between these malignant diseases and

EBV infection. However, it's important to note that the KEGG database does not

include EBV-related diseases like Burkitt lymphoma. Consequently, the diseases

highlighted are primarily autoimmune-related and various forms of cancer (Figure 3a).

The pathways enriched solely from target proteins and those retrieved from

literature exhibit notable similarities, with extensive studies on pathways such as

apoptosis, VEGF signaling pathway, TNF signaling pathway, Toll-like receptor

signaling pathway, and IL-17 signaling pathway (Figure 3b). This indicates the

coherence between GPT-generated pathways and the original articles. The last three

pathways highlight the importance of studying how innate immunity is affected by

viral antigens. The IL-17 signaling pathway triggers downstream NF-κB signaling

pathway and is known to be a key pathogenic factor of autoimmune diseases[14].

However, the relationship between the IL-17 pathway and EBV infection remains

unexplored. The high expression of IL-17 in multiple sclerosis patients could present

a novel area of EBV research and provide druggable candidates for certain

autoimmune diseases[15].

To provide a more comprehensive overview of the functions of EBV-associated

target proteins within their host, we used the Reactome database for annotation and

identified 375 biological processes ($p<0.05$) related to EBV infection (Figure 3c).

This network shows that EBV-targeted host proteins predominantly form clusters in

approximately 10 functional pathway groups. The largest clusters include innate

immunity pathways, represented by the TNF signaling pathway, Toll-like receptor

signaling cascades, and NF-κB signaling pathways. These are followed by cell growth

and differentiation pathways, exemplified by the PI3K/AKT signaling pathways and

related factors such as interleukins. Subsequently, DNA replication and mismatch

repair, along with cell cycle-related pathways, are prominently featured, particularly

those regulated by TP53. Moreover, the NOTCH signaling pathway and its regulation

of the TGF-β signaling pathways mediated by the SMAD proteins are also critical for

cell proliferation, differentiation, transitions, and apoptosis in our model of cellular

signaling crosstalk[16].

Intriguingly, we have identified some active mediators bridging the

interconnected cellular signaling crosstalk (Figure 3c). These active members include

deubiquitination and sumoylation that regulate protein modification and proteostasis;

TP53-mediated cellular senescence on DNA replication and cell cycle; PTEN and

downstream AKT signaling activation mediated by PIP3; BCR signaling for multiple

pathways spanning the immune response to cell proliferation; Fc epsilon receptor

(FCERI) signaling for the same function. Targets related to SARS-CoV-2 infection are

enriched alongside the immune signaling pathways, suggesting a shared immune

response elicited by EBV and SARS-CoV-2. This indicates the potential for

developing broad-spectrum antivirals for both viruses by targeting shared oncogenic

factors. Additionally, more discrete, smaller clusters were identified, including the

Wnt signaling pathway, mRNA editing, as well as the unfolded protein response (UPR)

essential for maintaining cellular homeostasis and survival under conditions of protein

misfolding and ER stress.

Analysis of host protein-protein interactions identifies key mediators in EBV-

mediated diseases

To further identify critical cellular factors involved in EBV-associated diseases,

we focused on EBV-targeted host proteins independent of pathway enrichment, which

may provide insights into key target proteins. Bibliographic analysis of publications

over time reveals trends in EBV-related host proteins (Figure 4a). p53 and c-Myc are

the two most studied proteins across all periods. Genomic profiling has identified

frequent TP53[17–19] and MYC[20,21] mutations or rearrangements in EBV-positive

malignancies. CD23 and CD40 are two critical factors for EBV infection on the B-cell

membrane. Earlier studies identified that EBNA2 specifically induces CD23

expression vital for B-cell transformation[22] and verified the role of LMP1 in viral

latency maintenance via mimicry of the CD40 signaling pathways[23]. As previously

discussed in Figure 2, multiple EBV antigens including BHRF1 and LMP1 are

associated with Bcl-2 related apoptosis in lymphoma and immortalized B-cells[19,24,25].

To better characterize the pathways and networks of EBV-related host factors

described above, we annotated all the target proteins using the STRING database and

performed protein-protein interaction (PPI) analysis. The top 3 enriched clusters are

shown (Figure 4). The first PPI cluster centers around TP53 genes, apoptotic

pathways, mTOR signaling, and the MAPK pathway cascades along with gene

silencing mediated by DICER1 and m6A methylation (Figure 4b). This cluster reflects

the role of EBV in suppressing the host apoptotic pathways and related signaling

transduction pathways to promote viral latency and oncogenesis. Clinical trials and

genomic profiling studies have identified the downregulation and mutation of TP53 in

multiple EBV-associated cancers[26–29]. EBV-encoded microRNAs (EBERs) are shown

to help maintain Burkitt's lymphomas by targeting caspase 3 to inhibit apoptosis and

aid the transformation of naïve B-cells[30]. Such inhibition usually works in harmony

with other factors like Bcl-2. Besides triggering classic apoptotic pathways, EBV also

bypasses the host antiviral systems by decreasing host Dicer protein via EBNA1-

mediated microRNAs degradation[31]. Recent studies have also uncovered the role of

circular RNAs in promoting the progression of EBV-associated gastric carcinoma

through the transactivation of METTL3[32]. Along this pathway, caspases have been

found to inhibit the m6A RNA modification process, thereby promoting EBV

replication[33]. Additionally, EBV infection is known to induce TCL1 gene expression

in Burkitt's lymphoma, which also serves as an activator for the Akt pathway[34].

The second cluster highlights EBV's roles in transcriptional regulation and

epigenetic modification, particularly through transcriptional factors (TFs) (Figure 4c).

MYC is a key TF associated with multiple cancers including EBV-associated

lymphomas[35]. Moreover, MYC abundance represses EBV reactivation and controls

the lytic switch in Burkitt lymphoma[36]. 3D genome mapping of EBV-positive

lymphoblastoid cell lines (LCLs) revealed distinctive EBV super-enhancer (SE)

targets like MCL1, IRF4, and EBF[37]. Bioinformatic analysis identified that H3K27ac

SEs overlap with STAT5 and NFAT targets, indicating their roles in promoting EBV

latency[13]. On a larger scale, broad screening of TFs integrated over 700 high-

throughput sequencing datasets and provides an atlas of EBV-related transcriptomic

and epigenetic host-virus regulatory interactions[38]. These EBV-specific TFs, such as

SPI1, EBF1, and PAX5, collectively contribute to EBV-induced oncogenesis and

latency[13].

The last cluster primarily comprises factors related to the innate immune response,

including cellular receptors, cytokines, and chemokines that regulate immune cell

activation, inflammatory response, and the tumor microenvironment (TME) (Figure

4d). EBV orchestrates immune evasion by cellular receptors like CD40, CD80, IL4,

IL6, IL10 and their ligands[39,40]. For example, LMP1 mimics CD40 to activate

downstream NF-κB pathways to induce B-cell proliferation[23]. After infection, EBV

hijacks host signaling transduction factors like STATs[41,42] and IRFs[43] to promote cell

survival and proliferation. LMP1 and LMP2A then activate the JAK/STAT pathway,

to facilitate cell growth factors and anti-apoptotic signaling. Meanwhile, EBV and

other herpesviruses induce the PI3K/Akt pathway to aid viral infection, latency, and

reactivation[44]. This cluster is also rich in enzymes, multiple cellular growth factors

(CGFs), and extracellular matrix (ECM) proteins, which represent the invasiveness of

EBV-infected cells. Early research discovered that viral LMP1 and BZLF1 proteins

up-regulate the expression and activity of matrix metalloproteinase 1 (MMP1), and

thereby confer the invasive properties of the cells[45]. Besides common growth factors,

we also identified Angiotensin-Converting Enzyme 2 (ACE2) in this cluster. Study

indicates that the lytic replication of EBV induces ACE2 expression in human

epithelial cells, which enhances the entry of SARS-CoV-2[46].

Exploring the critical relationships in EBV-mediated diseases

Due to the complex background of EBV pathogenesis, we next seek to elucidate

the influence of EBV antigens on host pathways in the context of multiple EBV-

associated malignancies (Figure 5a-b). In this section we demonstrate how our dataset

and generated PPI networks can be utilized to assist EBV-host interaction studies

using basic omics analyses. We plot the connections identified for EBV antigens and

host pathways onto circular connection hubs for DLBCL, BL, NPC, GC, and PTLD

(Figure 5c, Figure 6a and Figure S1).

Approximately 9% of DLBCLs are EBV positive with type II lantency[1]. In

DLBCL-centered hubs, we noticed the special linkage of EBV encoded dUTPase

BLLF3 and the TNF signaling pathways (Figure 5c). This indicates that BLLF3

induce DLBCL through the TNF pathways. dUTPase is crucial for maintaining the

balance of intracellular nucleotides, particularly during DNA replication[47]. Our

dataset pointed to study exploring BLLF3 and its interaction with host factor TLR2[48].

Our PPI networks show several known key factors closely interacting with TLR2 in

the background of EBV infection, such as IRAK1, HMGB1, TLR9, TNF, NFKB1,

IL1B and IL10. Next, we conducted pan-cancer analysis to examine the utility of

these proteins in distinguishing between DLBCL tumor and normal samples (Figure

5d). The results indicated clear difference between the tumor and samples from

normal tissues, suggesting their potential as marker genes for DLBCL diagnosis. As

DLBCL does not have corresponding gene expression data for normal tissue in TCGA,

we used whole blood and spleen, both of which are immune cells, as normal samples

for comparison. Taken apart, we noticed the high expression of HMGB1, TNF,

NFKB1 and IL10 in DLBCL tumor samples which function in TNF pathway (Figure

5e). At last, we examined the correlation of TNF expression and DLBCL patients'

survival probability (Figure 5f). The results show the survival possibility of patients

with high TNF expression dropped to 75% in approximately four years while samples

with low expression survived 100%. These findings above show the potential in

studying BLLF3 mediated TNF pathway activation in the carcinogenesis of DLBCL.

LMP1 is the most studied EBV antigens in our literature pool (Figure 5b). In BL,

literature indicates that LMP1 activates the calcium signaling pathway by

overexpressing the host factor CAMK4[49] (Figure 6a, Table S3). Our PPI network

shows that HDAC4 and CREB1 directly interact with CAMK4, and the secondary

factors that interact with these two, such as MEF2D, RUNX3, RUNX2, NFATC2,

JUND, FOS, RELA, TP53, TAF4, MECP2, CTNNB1, ATF4, and CRTC2 (Figure 4c).

We observed that although a dox-induced overexpression of LMP1 in the Akata cells

did not result in a significant increase in CAMK4, its interacting proteins listed above

exhibited a co-expression pattern and consistent upregulation (Figure 6b, c). This

suggests that the calcium signaling pathway may play a role in the oncogenesis of BL

by a LMP1-dependent manner.

Finally, we locate studies working on similar pathways associated with the same

antigens across cell lines from different EBV-related malignancies to compensate the

utility. For instance, studies on BL and NPC both demonstrated that LMP1 is

associated with glycolysis (Figure 6a, Figure S1). EBV typically establishes type I

latency in BL. Overexpressing LMP1 in BL cell lines suggests that LMP1 activates

PARP1 to induce glycolysis and promote oncogenesis in a HIF-1α-dependent

manner[45]. Study in NPC cell lines demonstrates that LMP1 represses Hox genes via

stalling the key mediators of DNA transcription like RNA pol II to modulate

glycolysis[51]. These studies in cell lines with different EBV latencies demonstrate the

role of LMP1 in regulating glycolysis. We next explore the transcriptomic correlation

for PARP1, HIF1A, HOXB3, and key transcription-related factors (Figure 6d-e).

Result show high correlation, which suggests a more complete mechanism that LMP1

activates PARP1, which subsequently binds to HIF-1α and RNA pol II and activates

HoxB3, finally modulate glycolysis.

Discussion

This study, for the first time, presents a mapping of EBV antigens, host factors,

and cellular pathways from available EBV-related articles using LLMs. We have

summarized the major pathways associated to each EBV antigens, and provided a

descriptive overview of core PPI clusters. At last, we provided examples of ustilizing

our dataset for marker gene analyses.

However, certain limitations in our research need to be addressed. While our

initial data pool contained 36,105 EBV-related publications, our strict prompts and

data filtering reduced this to 2,289 entries, encompassing 1,839 unique publications

(~5.1%) for downstream analyses. If loosening the criteria, we will be able to obtain

more entries for deeper analyses, although the accuracy may decrease. Secondly, our

literature study was based solely on conclusions drawn from abstracts of the available

publications which has its intrinsic limitation in the study of network biology.

Although we used the frequency of topics as a proxy for their importance in the EBV

antigen-pathway mapping, the sheer number of publications may not fully represent

actual biological significance. Occasionally, a higher citation rate may influence their

proportions, potentially translating non-experimental results. Future efforts may focus

on using supervised machine learning algorithms to explore the transcriptomic

networks of EBV-induced cellular changes for enhanced utility and usability. In the

last part, we showcase through basic omics analyses that our EBV literature analysis

bears the potential to inspire researchers to engage with former published work of

scientists worldwide for novel effort in illustrating the mechanism of EBV-host

interaction. Finally, EBV latency types are critical in understanding EBV gene

expression and virus-host interaction. However, such information is largely

unattainable in abstracts. Future research based on more powerful tools can extract the

main text for related information.

This study represents a pioneering effort in the application of LLMs in EBV-

related research, highlighting the potential of LLMs in advancing our understanding

of viral interactions and cellular pathways. Unlike broad-scale context generation, our

study clarifies the relationships between EBV antigens and various cellular pathways

based on the manual annotation of GPT-generated summaries, presenting networks of

EBV-related host proteins for future research. Beyond EBV, the big data era and

advancements in various LLMs underscore the urgent need for advanced, digitized

platforms in all areas of viral research. Such tools are pivotal for accelerating research

progress, advancing basic virology, and fostering the development of novel antivirals.

References

1.   Damania, B., Kenney, S. C. & Raab-Traub, N. Epstein-Barr virus: Biology and

clinical disease. *Cell* S0092867422011217 (2022) doi:10.1016/j.cell.2022.08.026.

2.   Hatzivassiliou, E., Miller, W. E., Raab-Traub, N., Kieff, E. & Mosialos, G. A fusion of

the EBV latent membrane protein-1 (LMP1) transmembrane domains to the CD40

cytoplasmic domain is similar to LMP1 in constitutive activation of epidermal growth

factor receptor expression, nuclear factor-kappa B, and stress-activated protein kinase. *J*

Immunol **160**, 1116–1121 (1998).

3.   Kaykas, A. & Sugden, B. The amino-terminus and membrane-spanning domains of

LMP-1 inhibit cell proliferation. *Oncogene* **19**, 1400–1410 (2000).

4.   Gewurz, B. E. *et al.* Genome-wide siRNA screen for mediators of NF-κB activation.

Proc. Natl. Acad. Sci. U.S.A. **109**, 2467–2472 (2012).

5.   Luftig, M. *et al.* Epstein–Barr virus latent membrane protein 1 activation of NF-κB

through IRAK1 and TRAF6. *Proc. Natl. Acad. Sci. U.S.A.* **100**, 15595–15600 (2003).

6.   Valentine, R. *et al.* Epstein-Barr virus-encoded EBNA1 inhibits the canonical NF-

kappaB pathway in carcinoma cells by inhibiting IKK phosphorylation. *Mol Cancer* **9**, 1

(2010).

7. González-Márquez, R., Schmidt, L., Schmidt, B. M., Berens, P. & Kobak, D. The

landscape   of   biomedical   research.   *Patterns*   100968   (2024)

doi:10.1016/j.patter.2024.100968.

8. Wu, X. *et al.* reguloGPT: Harnessing GPT for Knowledge Graph Construction of

Molecular Regulatory Pathways. Preprint at https://doi.org/10.1101/2024.01.27.577521

(2024).

9. Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic*

Acids Res **28**, 27–30 (2000).

10. Goldman, M. J. *et al.* Visualizing and interpreting cancer genomics data via the Xena

platform. (2020).

11. Tang, Z., Kang, B., Li, C., Chen, T. & Zhang, Z. GEPIA2: an enhanced web server

for large-scale expression profiling and interactive analysis.

12. Bal, E. *et al.* Super-enhancer hypermutation alters oncogene expression in B cell

lymphoma. *Nature* **607**, 808–815 (2022).

13. Zhou, H. *et al.* Epstein-Barr Virus Oncoprotein Super-enhancers Control B Cell

Growth. *Cell Host & Microbe* **17**, 205–216 (2015).

14. Mills, K. H. G. IL-17 and IL-17-producing cells in protection versus pathology. *Nat*

Rev Immunol **23**, 38–54 (2023).

15. Luchtman, D. W., Ellwardt, E., Larochelle, C. & Zipp, F. IL-17 and related cytokines

involved in the pathology and immunotherapy of multiple sclerosis: Current and future

developments. *Cytokine Growth Factor Rev* **25**, 403–413 (2014).

16. K, L. Signaling Cross Talk between TGF-β/Smad and Other Signaling Pathways.

Cold Spring Harbor perspectives in biology **9**, (2017).

17. Menter, T. *et al.* Mutational landscape of B-cell post-transplant lymphoproliferative

disorders. *Br J Haematol* **178**, 48–56 (2017).

18. Bruce, J. P. *et al.* Whole-genome profiling of nasopharyngeal carcinoma reveals

viral-host co-operation in inflammatory NF-κB activation and immune escape. *Nat*

Commun **12**, 4193 (2021).

19. Frontzek, F. *et al.* Molecular profiling of EBV associated diffuse large B-cell

lymphoma. *Leukemia* **37**, 670–679 (2023).

20. Panea, R. I. *et al.* The whole-genome landscape of Burkitt lymphoma subtypes.

Blood **134**, 1598–1607 (2019).

21. Garcia-Reyero, J. *et al.* Genetic lesions in MYC and STAT3 drive oncogenic

transcription factor overexpression in plasmablastic lymphoma. *Haematologica* **106**,

1120–1128 (2021).

22. Wang, F. *et al.* Epstein-Barr virus nuclear antigen 2 specifically induces expression

of the B-cell activation antigen CD23. *Proc Natl Acad Sci U S A* **84**, 3452–3456 (1987).

23. Uchida, J. *et al.* Mimicry of CD40 signals by Epstein-Barr virus LMP1 in B

lymphocyte responses. *Science* **286**, 300–303 (1999).

24. Oudejans, J. J. *et al.* BHRF1, the Epstein-Barr virus (EBV) homologue of the BCL-2

protooncogene, is transcribed in EBV-associated B-cell lymphomas and in reactive

lymphocytes. *Blood* **86**, 1893–1902 (1995).

25. Kenney, J. L., Guinness, M. E., Curiel, T. & Lacy, J. Antisense to the epstein-barr

virus (EBV)-encoded latent membrane protein 1 (LMP-1) suppresses LMP-1 and bcl-2

expression and promotes apoptosis in EBV-immortalized B cells. *Blood* **92**, 1721–1727

(1998).

26. Ribeiro, J. *et al.* P53 deregulation in Epstein-Barr virus-associated gastric cancer.

Cancer Lett **404**, 37–43 (2017).

27. Sarkozy, C. *et al.* Mutational landscape of gray zone lymphoma. *Blood* **137**, 1765–

1776 (2021).

28. Lee, J. E. *et al.* Clinicopathologic and genomic characteristics of mucinous gastric

adenocarcinoma. *Gastric Cancer* **25**, 697–711 (2022).

29. Lee, H. S., Chang, M. S., Yang, H.-K., Lee, B. L. & Kim, W. H. Epstein-barr virus-

positive gastric carcinoma has a distinct protein expression profile in comparison with

epstein-barr virus-negative carcinoma. *Clin Cancer Res* **10**, 1698–1705 (2004).

30. Vereide, D. T. *et al.* Epstein-Barr virus maintains lymphomas via its miRNAs.

Oncogene **33**, 1258–1264 (2014).

31. Mansouri, S., Pan, Q., Blencowe, B. J., Claycomb, J. M. & Frappier, L. Epstein-Barr

virus EBNA1 protein regulates viral latency through effects on let-7 microRNA and dicer.

J Virol **88**, 11166–11177 (2014).

32. Zhang, J.-Y. *et al.* ebv-circRPMS1 promotes the progression of EBV-associated

gastric carcinoma via Sam68-dependent activation of METTL3. *Cancer Lett* **535**, 215646

(2022).

33. Zhang, K. *et al.* Caspases Switch off the m^{6}A RNA Modification Pathway to Foster

the Replication of a Ubiquitous Human Tumor Virus. *mBio* **12**, e01706-21 (2021).

34. Bell, A. & Rickinson, A. B. Epstein-Barr virus, the TCL-1 oncogene and Burkitt's

lymphoma. *Trends Microbiol* **11**, 495–497 (2003).

35. Dhanasekaran, R. *et al.* The MYC oncogene - the grand orchestrator of cancer

growth and immune evasion. *Nat Rev Clin Oncol* **19**, 23–36 (2022).

36. Guo, R. *et al.* MYC Controls the Epstein-Barr Virus Lytic Switch. *Mol Cell* **78**, 653-

669.e8 (2020).

37. Jiang, S. *et al.* The Epstein-Barr Virus Regulome in Lymphoblastoid Cells. *Cell Host*

Microbe **22**, 561-573.e4 (2017).

38. Arvey, A. *et al.* An Atlas of the Epstein-Barr Virus Transcriptome and Epigenome

Reveals Host-Virus Regulatory Interactions. *Cell Host & Microbe* **12**, 233–245 (2012).

39. Park, G. B. *et al.* CD80 (B7.1) and CD86 (B7.2) induce EBV-transformed B cell

apoptosis through the Fas/FasL pathway. *Int J Oncol* **43**, 1531–1540 (2013).

40. Kis, L. L., Takahara, M., Nagy, N., Klein, G. & Klein, E. IL-10 can induce the

expression of EBV-encoded latent membrane protein-1 (LMP-1) in the absence of EBNA-

2 in B lymphocytes and in Burkitt lymphoma- and NK lymphoma-derived cell lines. *Blood*

107, 2928–2935 (2006).

41. Weber-Nordt, R. M. *et al.* Constitutive activation of STAT proteins in primary

lymphoid and myeloid leukemia cells and in Epstein-Barr virus (EBV)-related lymphoma

cell lines. *Blood* **88**, 809–816 (1996).

42. Hsiao, J.-R. *et al.* Constitutive activation of STAT3 and STAT5 is present in the

majority of nasopharyngeal carcinoma and correlates with better prognosis. *Br J Cancer*

89, 344–349 (2003).

43. Wang, L. *et al.* LMP1 signaling pathway activates IRF4 in latent EBV infection and a

positive circuit between PI3K and Src is required. *Oncogene* **36**, 2265–2274 (2017).

44. Cohen, J. I. Herpesviruses in the Activated Phosphatidylinositol-3-Kinase-δ

Syndrome. *Front Immunol* **9**, 237 (2018).

45. Lu, J., Chua, H.-H., Chen, S.-Y., Chen, J.-Y. & Tsai, C.-H. Regulation of matrix

metalloproteinase-1 by Epstein-Barr virus proteins. *Cancer Res* **63**, 256–262 (2003).

46. Verma, D., Church, T. M. & Swaminathan, S. Epstein-Barr Virus Lytic Replication

Induces ACE2 Expression and Enhances SARS-CoV-2 Pseudotyped Virus Entry in

Epithelial Cells. *J Virol* **95**, e0019221 (2021).

47. Fleischmann, J., Kremmer, E., Greenspan, J. S., Grässer, F. A. & Niedobitek, G.

Expression of viral and human dUTPase in Epstein-Barr virus-associated diseases.

Journal of Medical Virology **68**, 568–573 (2002).

48. Williams, M. & Ariza, M. E. EBV Positive Diffuse Large B Cell Lymphoma and

Chronic Lymphocytic Leukemia Patients Exhibit Increased Anti-dUTPase Antibodies.

Cancers (Basel) **10**, 129 (2018).

49. Mosialos, G. *et al.* A Ca2+/calmodulin-dependent protein kinase, CaM kinase-Gr,

expressed after transformation of primary human B lymphocytes by Epstein-Barr virus

(EBV) is induced by the EBV oncogene LMP1. *J Virol* **68**, 1697–1705 (1994).

50. Hulse, M. *et al.* Poly(ADP-ribose) polymerase 1 is necessary for coactivating

hypoxia-inducible factor-1-dependent gene expression by Epstein-Barr virus latent

membrane protein 1. *PLoS Pathog* **14**, e1007394 (2018).

51. Jiang, Y. *et al.* Repression of Hox genes by LMP1 in nasopharyngeal carcinoma and

modulation of glycolytic pathway genes by HoxC8. *Oncogene* **34**, 6079–6091 (2015).

Acknowledgments: We thank all members of our laboratory for helpful advice and

assistance.

Funding: This work was supported by the Science, Technology and Innovation

Commission of Shenzhen Municipality (grant no. JCYJ20230807093208017 to Y.P.),

the Guangdong Natural Science Foundation (grant no. 2024A1515013126 to Y.P.).

Author contributions: Y.P. conceived the application of LLM in the EBV-related

literature analysis. J.Y. designed and carried out the data mining and visualization.

Y.W., H.W., and Z.W. provided expertise in technical support and data analysis. J.Y.

wrote the manuscript and Y.P. revised the manuscript.

Competing interests: The authors declare no potential conflicts of interest.

Figure legends

Figure 1. LLMs-assisted data mining reveals the key insights in EBV research. (**a**)

The trend of EBV-related publications in different years. (**b**) The pipeline in this study

is illustrated as a flowgraph. (**c**) Evaluation of GPT-generated output is conducted by

examining 102 entries recorded for EBNA3C and LMP2. (**d**) The relative frequency

of EBV antigens (log2 transformed) in our analysis is illustrated in the word cloud. (**e**)

Statistical analysis highlights the research focus on various EBV antigens over time,

showcasing only the four most studied antigens for each decade. (**f**) The relative

frequency of EBV-related pathways was shown in the word cloud. (**g**) Statistical

analysis highlights the research focus on various EBV-related pathways over time,

showcasing only the four most pathways for each decade.

Figure 2. Pathway mapping reveals the distinct signatures of EBV antigens. (**a**) The

chord diagram shows the linkage between the top 33 EBV antigens and their

corresponding pathways. (**b**) Heatmap view of the linkage between EBV latent or

lytic antigens and the associated pathways.

Figure 3. Gene enrichment and pathway analysis uncover defining characteristics of

EBV-induced cellular modulations. (a-b) KEGG enrichment shows the top 10 EBV-

related diseases or pathogens (a) and signaling pathways (b). (c) Reactome

enrichment maps the landscape of EBV-targeted pathways and their correlations based

on the shared host proteins.

Figure 4. Analysis of host protein-protein interactions identifies key mediators in

EBV-mediated diseases. (a) Statistical analysis reveals the focus of study every

decade on EBV-targeted host proteins. TF, transcription factor; GPCR, G Protein-

Coupled Receptor. (b-d) Top 3 PPI networks describe the basic functional clusters of

EBV-targeted host proteins.

Figure 5. Revealing the key relationships in EBV-mediated DLBCLs. (a) The bar plot

shows the numbers of total articles of the top 12 most studied EBV-related diseases.

(b) The bar plot shows the numbers of total articles of the top 7 most studied EBV

antigens. (c) The circular hub graph elucidates the DLBCL-related antigen-pathway

relations. (d) PCA analysis highlights distinct DLBCL tumor samples using TLR2-

related marker gene expressions. (e) The TLR2-related marker gene expressions

between normal tissue and tumor tissue. (f) Survival plot shows high TNF expression

leads to less survival probability (red indicates samples with normalized TNF counts >

7.907 and blue indicates less).

Figure 6. Exploring the functions of EBV antigens in BL. (a) The circular hub graph

elucidates the BL-related antigen-pathway relations. (b-c) The CAMK4-related

factors are co-expressed by LMP1 overexpression. (d-e) The PARP1-related factors

are co-expressed by LMP1 overexpression.

Supplementary materials

Figure S1: Determining the key relationships in EBV-mediated NPC, PTLD, and GC.

(a-c) The circular hub graph elucidates the NPC-related, PTLD-related and GC-

related antigen-pathway relations. NPC, nasopharyngeal carcinoma; PTLD,

Posttransplant lymphoproliferative disorders; GC, gastric cancer.;

Table S1: The formal names and aliases used in GPT-generated term substitution;

Table S2: The correspondence between GPT-4 generated pathways and manually

annotated pathway to KEGG entries;

Table S3: The final dataset includes 2,289 entries of EBV antigens and their

associated pathways;

Table S4: Validation for GPT-generated results;

Table S5: Ranking of most studied EBV antigens (with >10 related articles);

a

b

log2(associated publications)

Latent antigens   Lytic antigens

TP53-associated apoptotic signaling pathways

EBV-associated TFs and TF-related factors

Node family
- Enzyme (Enzyme)
- Epigenetic (Epigenetic)
- GPCR (GPCR)
- Ion Channel (Ion Channel)
- Kinase (Kinase)
- Nuclear Receptor (Nuclear Receptor)
- TF (TF)
- TF-Epigenetic (TF-Epigenetic)
- Transporter (Transporter)
- Others (Others)

Immune response and related factors