Title: Meta-analysis of public RNA-sequencing data of drought and salt stresses in different phenotypes of *Oryza sativa*

Mitsuo Shintani 1 and Hidemasa Bono 1, 2, *

1 Graduate School of Integrated Sciences for Life, Hiroshima University, Higashi-Hiroshima, Japan
2 Genome Editing Innovation Center, Hiroshima University, Higashi-Hiroshima, Japan

*Correspondence: Hidemasa Bono
3-10-23 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-0046, Japan
Email: bonohu@hiroshima-u.ac.jp
Tel: +81-82-424-4013

Abstract:

Environmental stresses, such as drought and salt, adversely affect plant growth and crop productivity. While many studies have focused on established components of stress signaling pathways, research on unknown elements remains limited. In this study, we collected RNA sequencing (RNA-Seq) data from *Oryza sativa* subsp. *indica* and *Oryza sativa* subsp. *japonica* registered in public databases and conducted a meta-analysis integrating multiple studies. Focusing on two types of stress conditions (salt and drought), we aimed to identify novel stress-responsive genes in *Oryza sativa* by comparing RNA-Seq data from stress-resistant and stress-susceptible cultivars. We analyzed 105 paired datasets with different phenotypes under drought and salt stress conditions to identify genes with common expression changes across multiple studies. A meta-analysis identified 10 genes specifically upregulated in resistant cultivars and 12 specifically upregulated in susceptible cultivars under both drought and salt stress conditions. Furthermore, by comparing previously identified stress-responsive genes in *Arabidopsis thaliana*, we explored genes potentially involved in stress response mechanisms that are conserved across plant species. The genes identified in this data-driven study that potentially determine plant stress resistance or susceptibility phenotypes may serve as research targets for elucidating novel plant stress mechanisms and candidates for genome editing.

Keywords:

Oryza sativa, Salt stress, Drought stress, RNA-sequencing, Meta-analysis

1. Introduction

Rice (*Oryza sativa*) is an important food in East Asia, South Asia, the Middle East, Latin America, and the West Indies, and it is estimated to account for more than one-fifth of the calories consumed by humans worldwide [1]. Rice is a crucial crop because it is high in calories and contains more abundant amounts of essential vitamins and minerals than other grains [2]. However, rice production faces salinity and drought stress, which hinders its cultivation.

Salinity and drought are major abiotic stresses that inhibit plant growth and productivity [3–7]. Rice is susceptible to the negative effects of drought and salt damage, which lead to serious issues of reduced yields. Twenty percent of agricultural land currently used worldwide is affected by salt stress, and this percentage increases every year owing to anthropogenic and natural factors [8]. High Na+ levels induce K+ and Ca2+ efflux from the cytoplasm, causing intracellular homeostatic imbalance, nutrient deficiency, oxidative stress, growth retardation, and cell death [9]. In addition, high salt concentrations reduce photosynthesis through stomatal limitations, such as stomatal closure [10]. Salt stress also has a negative effect on photosynthesis because of non-stomatal limitations, such as chlorophyll dysfunction, deficiency of photosynthetic enzyme proteins and membranes, and disruption of the ultrastructure of the chloroplast [11–13].

Drought represents a state of soil water deficit in which insufficient water is available for plants to fully grow and complete their life cycles. As soil water evaporates, salinity is concentrated in the soil solution, causing drought and salinity simultaneously [14].

Across Asia, 20% of all rice-producing areas are affected by drought each year [15]. Several studies focusing on plant drought have shown that drought delays the flowering time of rice plants and consequently suppresses plant growth, resulting in reduced ear number and fertile fruit number, ultimately reducing yield [16–19]. Drought stress causes changes in the antioxidant and osmotic regulatory systems of rice, leading to the accumulation of antioxidants and osmotic regulators [20,21].

Previous studies identified several genes that contribute to salinity and drought stress in rice [8,22,23]. For example, the transcription factors *OsMYBc*, *OsbZIP71*, and *OsNF-YC13* improve salt tolerance by activating K+/Na+ transporters and increasing the K+/Na+ ratio [24–26].

In addition, *OsP5CS1* and *OsP5CR* play crucial roles in synthesizing compatible solute proline, increasing proline levels, and thereby enhancing salt tolerance [27]. *OsCPK4* and *OsCPK12*, which encode calcium-dependent protein kinases, are involved in scavenging reactive oxygen species (ROS), thereby enhancing salt tolerance [28,29]. Similarly, the transcription factors *OsZFP179* and *OsZFP182* also improve salt tolerance by enhancing ROS scavenging ability [30,31].

Regarding drought tolerance, the expression of *OsCPK9* promotes stomatal closure and improves osmotic regulation, thereby enhancing plant drought tolerance [32]. The transcription factors *OsNAC5* and *OsNAC10* increase grain yield and root development under drought conditions [33,34]. *OsDREB2A* enhances the survival of transgenic plants under drought stress and *OsDREB2B* increases root length and number [35,36].

However, current research has only revealed a small part of the overall stress response mechanism in rice. In particular, the causes of

large differences in stress sensitivity among rice varieties are not fully understood. Therefore, identifying the factors that determine the differences in stress sensitivity is important for elucidating the entire stress response mechanism in rice.

In this study, we focused on the differences in drought and salt stress responses among rice varieties and attempted to identify genes specifically involved in stress sensitivity (susceptible), stress tolerance (resistant), and each phenotype. Through the integrated analysis of a large amount of RNA sequencing (RNA-Seq) data retrieved from public databases, we explored novel responsive genes that have not been reported to be associated with salt or drought stress. By comparing differentially expressed genes identified between cultivars with different susceptibilities to stress, we narrowed down candidate genes involved in the phenotype.

In a previous study, we performed a meta-analysis of stress conditions involving abscisic acid (ABA), salt, and dehydration in *Arabidopsis thaliana* [37]. By comparing the stress-responsive genes identified in *Oryza sativa* from this study with those from the meta-analysis of *Arabidopsis thaliana*, we identified candidate genes that contribute to stress response mechanisms conserved across plant species.

The findings of this study provide a new perspective for basic research on the stress response mechanisms in rice. Moreover, the identified genes have the potential for use as target candidates for genome editing to develop stress-resistant rice.

2. Results

2.1 Overview of this study

Initially, we present an overview of the entire study. Through comprehensive data analysis, this research aimed to achieve two primary objectives:

(1) Identification of genes suggested to contribute to stress tolerance phenotypes in *Oryza sativa.*

(2) Identification of genes implicated by the data to be involved in common stress response mechanisms across plant species, specifically in *Oryza sativa* and *Arabidopsis thaliana.*

This study consisted of six steps as shown in Figure 1.

Figure 1. Overview of the six steps of this study. Step 1: RNA-Seq data for *Oryza sativa* were retrieved from public databases (National Center for Biotechnology Information Gene Expression Omnibus (NCBI GEO) and European Bioinformatics Institute BioStudies (EBI BioStudies). Step 2: Samples were categorized according to stress type (salt/drought) and phenotype (resistant/susceptible). Differentially expressed genes (DEGs) were identified

using the TN-score method. Step 3: Enrichment analysis was performed on the DEGs for each phenotype and stress condition to evaluate the validity of the analysis. Step 4: Genes involved in the phenotypic differences between *Oryza sativa* cultivars were selected under drought and salt stress conditions. Step 5: Common genes among different stress conditions were selected. Step 6: Genes involved in stress response mechanisms conserved across different plant species were selected.

2.2 Retrieval and curation of RNA-Seq data from public databases

RNA-Seq data were retrieved from the National Center for Biotechnology Information Gene Expression Omnibus (NCBI GEO) and European Bioinformatics Institute BioStudies (EBI BioStudies) public databases. Compared with microarray data, RNA-Seq data were primarily derived from the Illumina platform, making them more suitable for comparative analyses of studies from different research groups. Therefore, only RNA-Seq data were used in this study, while microarray data were excluded. In this study, based on the description in the papers and metadata of databases, the phenotype of cultivars reported to present strong resistance was defined as "resistant" and the cultivars reported to present relatively weak resistance were defined as "susceptible." A total of 202 samples were collected from the 11 projects, resulting in 105 paired datasets of stress-treated and untreated control samples. Because of manual curation, the collected data were divided into four categories based on the stress type and phenotype: (1) salt-resistant, (2) salt-susceptible, (3) drought-resistant, and (4) drought-susceptible. The number of paired datasets for each category was as follows: 24 for salt-resistant, 36 for salt-susceptible, 31 for drought-resistant, and 14 for drought-susceptible. To systematically organize the data used in the analysis, a table including the stress type, phenotype, subspecies, research project ID, and experimental pair number was prepared for each rice variety (Table 1). When the number of collected sample pairs was compared among the tissues, it was revealed that samples derived from "leaf" contained the most data (Figure 2). Details on all other metadata are presented in Supplementary Table 1.

Table 1. Summary of *Oryza sativa* sample metadata utilized in the meta-analysis

Stress type	Phenotype	Cultivar	Sub species	Study project ID	Number of sample pairs
Salt	Resistant	Pokkali	*indica*	ERP003982	15
Salt	Resistant	Baldo	*japonica*	SRP131107	6
Salt	Resistant	Tainung 67	*japonica*	SRP397488	3
Salt	Susceptible	Nipponbare	*japonica*	SRP200410	6
Salt	Susceptible	Nipponbare	*japonica*	ERP002527	9
Salt	Susceptible	IR29	*indica*	ERP003983	15
Salt	Susceptible	Vialone Nano	*japonica*	SRP131107	6
Drought	Resistant	Nagina 22	*indica*	SRP253184	2

Drought	Resistant	P28	*indica*	SRP042233	4
Drought	Resistant	H471	*indica*	SRP042233	4
Drought	Resistant	PR403	*indica*	SRP286181	6
Drought	Resistant	Azucena	*japonica*	SRP065945	15
Drought	Susceptible	Nipponbare	*japonica*	SRP065476	2
Drought	Susceptible	IR64	*indica*	SRP253184	2
Drought	Susceptible	HHZ	*indica*	SRP042233	4
Drought	Susceptible	PY6	*indica*	SRP286181	6

Figure 2. Tissue distribution of *Oryza sativa* samples utilized in the meta-analysis

Visualization of the distribution of RNA-seq sample tissue counts from different projects of *Oryza sativa*.

2.3 Analysis of collected data and selection of differentially expressed genes in Oryza sativa

The collected samples were subjected to quality control and expression quantification using Salmon[38], which calculated the transcripts per million (TPM) values for each gene in each sample (Supplementary Table 2). Stress-treated and non-treated samples were paired for each gene, and the expression ratios (TN ratios) and TN scores were calculated. The TN ratio was calculated as follows:

TN ratio = (stress-treated TPM + 1) / (non-treated TPM + 1).

If the TN ratio was higher than the threshold, the gene was considered upregulated; whereas if it was lower than the reciprocal of the threshold, it was considered downregulated. Otherwise, it was considered unchanged. To classify the upregulated and downregulated genes, we evaluated 1.5-fold, 2-fold, 5-fold, and 10-fold thresholds and finally chose the 2-fold threshold. Therefore, genes with a TN ratio higher than 2 were classified as upregulated, while genes with a TN ratio lower than 0.5 were classified as downregulated. The TN score of each gene was determined by subtracting the number of downregulated experiments from the number of upregulated experiments to assess changes in gene expression under stress conditions across the entire experiment.

The collected samples were classified into the four aforementioned categories based on the phenotype and type of stress treatment, and the TN scores for each category were calculated. After considering multiple expression ratio thresholds, we adopted a 2-fold threshold (TN2). This threshold was slightly lower to provide a comprehensive analysis. More severe scores for the 5-fold (TN5) and 10-fold (TN10) thresholds were also calculated and are listed in the Supplementary Table. The TN ratios and TN scores for all the genes are available online (Supplementary Tables 3 and 4).

Additionally, the distribution of TN scores for the genes was visualized using scatter plots (Supplementary Figure 1). Focusing on points where scores changed significantly on the scatter plot, the top and bottom genes based on TN2 scores were selected as differentially expressed genes (DEGs). This accounted for the top or bottom 1% of all reference genes. Abrupt score changes indicated that the expression of these genes was remarkably variable in the dataset used in this study. The number of identified genes and range of TN2 scores are summarized in Table 2. Lists of the upregulated and downregulated genes are available online (Supplementary Tables 5 and 6, respectively). For the DEGs identified in rice, the corresponding orthologs in *Arabidopsis thaliana* were determined using sequence similarity searches with BLASTP. The list of these genes is available online (Supplementary Table 7).

Table 2. Distribution of DEG numbers and TN2 scores in *Oryza sativa* under different stress conditions and phenotypes abased on the meta-analysis

Treatment type	Phenotype	Regulation	All pairs	TN Score (S)	Number of *Oryza sativa* genes	Number of ortholog *Arabidopsis* genes
Salt	Resistant	upregulated	24	$6 \leqq S \leqq 15$	397	307
Drought	Resistant	upregulated	31	$13 \leqq S \leqq 26$	569	421
Salt	Susceptible	upregulated	36	$10 \leqq S \leqq 22$	456	312
Drought	Susceptible	upregulated	14	$9 \leqq S \leqq 13$	441	264

Salt	Resistant	downregulated	24	-17 ≤ S ≤ -5	473	379
Drought	Resistant	downregulated	31	-22 ≤ S ≤ -11	415	314
Salt	Susceptible	downregulated	36	-16 ≤ S ≤ -6	489	302
Drought	Susceptible	downregulated	14	-14 ≤ S ≤ -7	556	282

2.4 Enrichment analysis to evaluate the characteristics of DEGs

To evaluate the characteristics of the identified DEGs, we performed enrichment analyses of individual DEG groups using ShinyGO. The results are shown in Supplementary Figure 2 and Supplementary Table8. The top two GO terms for each DEG are listed in Table 3.

Table 3. Top two GO terms from enrichment analyses of individual DEGs of *Oryza sativa* under different stress conditions and phenotypes

Treatment type	Phenotype	Regulation	GO term ID	GO name	Fold Enrichment	Enrichment FDR
Salt	Resistant	upregulated	GO:0006568	Tryptophan metabolic proc.	25.9507772	1.67E-06
Salt	Resistant	upregulated	GO:0006586	Indolalkylamine metabolic proc.	25.9507772	1.67E-06
Salt	Resistant	downregulated	GO:0010227	Floral organ abscission	27.1640091	0.00693101
Salt	Resistant	downregulated	GO:0009838	Abscission	24.4476082	0.00929488
Salt	Susceptible	upregulated	GO:0009631	Cold acclimation	40.5612245	1.41E-08
Salt	Susceptible	upregulated	GO:0042542	Response to hydrogen peroxide	15.2104592	0.00013353
Salt	Susceptible	downregulated	GO:0030417	Nicotianamine metabolic proc.	79.3237251	8.73E-05

Salt	Susceptible	downregulated	GO:0030418	Nicotianamine biosynthetic proc.	79.3237251	8.73E-05
Drought	Resistant	upregulated	GO:0009631	Cold acclimation	27.7756211	3.58E-06
Drought	Resistant	upregulated	GO:0009845	Seed germination	11.9386442	0.00010894
Drought	Resistant	downregulated	GO:0016540	Protein autoprocessing	58.1707317	0.01989467
Drought	Resistant	downregulated	GO:0009068	Aspartate family amino acid catabolic proc.	43.6280488	0.00360792
Drought	Susceptible	upregulated	GO:0006097	Glyoxylate cycle	82.4308756	9.19E-05
Drought	Susceptible	upregulated	GO:0046487	Glyoxylate metabolic proc.	82.4308756	9.19E-05
Drought	Susceptible	downregulated	GO:0009068	Aspartate family amino acid catabolic proc.	32.8211009	0.00430687
Drought	Susceptible	downregulated	GO:0006546	Glycine catabolic proc.	28.1323722	0.00567738

Enrichment analysis revealed distinct patterns of gene regulation under different stress conditions and phenotypes. These results suggest that diverse stress response mechanisms may exist depending on the type of stress and phenotypic traits.

Furthermore, we focused on DEGs that showed similar expression patterns (upregulation or downregulation) across different phenotypes and stress conditions and considered them as a single group. We performed enrichment analyses for two types of combined DEGs. The first group consisted of all genes whose expression was upregulated under at least one condition (Figure 3a), and the second group consisted of all genes whose expression was downregulated under at least one condition (Figure 3b).

(a) All DEG of *Oryza sativa* genes list upregulated

(b) All DEG of *Oryza sativa* genes list downregulated

Figure 3. Enrichment analysis of the combined DEGs with similar expression patterns across phenotypes and stress conditions (a) All upregulated genes with at least one DEG (b) All downregulated genes with at least one DEG

For upregulated DEGs (Figure 3a), the most significantly enriched terms were "diterpenoid metabolic processes" and "water deprivation." Other enriched terms included responses to various abiotic stressors, such as heat and salt stress, and hormone responses, especially abscisic acid (ABA). These results support the validity of the selection of DEGs responsive to salt and drought stress by the meta-analysis.

For the downregulated genes (Figure 3b), the analysis revealed a different set of enriched terms. The most significantly enriched terms were related to nitrogen metabolism, including "nitrate metabolic processes," "nitrate assimilation," and "reactive nitrogen species metabolic processes." Interestingly, several terms involved in oxidative stress response and detoxification were also significantly enriched, such

as "hydrogen peroxide catabolic process" and "reactive oxygen species metabolic process." Additionally, photosynthesis-related processes and transmembrane transport were among the enriched pathways for downregulated genes. Detailed results of the enrichment analysis for individual DEGs and all combined DEGs are provided in Supplementary Table 8.

2.5 Selection of genes involved in phenotypic differences between Oryza sativa cultivars under drought and salt stress conditions

We compared the DEGs between the stress-resistant and stress-susceptible phenotypes for both salt and drought stress to identify genes specific to each condition (Supplementary Figure 3). The numbers of common and unique genes in each group are summarized in Table 4.

Table 4. Number of common and unique DEGs between resistant and susceptible cultivars under salt and drought stress

Stress type	Regulation	Common to both phenotypes	Specific to Resistant phenotype	Specific to Susceptible phenotype
Salt	upregulated	106	291	350
Salt	downregulated	114	359	375
Drought	upregulated	298	271	143
Drought	downregulated	249	166	307

We conducted an enrichment analysis of these individual gene sets, and the results are shown in Supplementary Figure 3. However, as no hits were found in gene set (j), only those for the "All available gene sets" setting were used. The top GO terms for each DEG are summarized in Table 5.

Table 5. Top enriched terms from enrichment analysis of individual DEG groups associated with phenotypic differences in salt and drought stress responses in *Oryza sativa*

Treatment type	Unique or common	Phenotype	Regulation	GO term ID	GO name	Fold Enrichment	Enrichment FDR
Salt	unique	Resistant	up	GO:0051501	Diterpene phytoalexin metabolic proc.	76.6607143	1.50E-04
Salt	unique	Susceptible	up	GO:0006572	Tyrosine catabolic proc.	71.1940299	0.01518192

Salt	common	Resistant and Susceptible	up	GO:0009631	Cold acclimation	96.4285714	3.86E-06
Salt	unique	Resistant	down	GO:0010227	Floral organ abscission	36.0271903	0.03397284
Salt	unique	Susceptible	down	GO:0030417	Nicotianamine metabolic proc.	104.300292	0.00019586
Salt	common	Resistant and Susceptible	down	GO:0006826	Iron ion transport	50.9615385	4.00E-05
Drought	unique	Resistant	up	GO:0055129	L-proline biosynthetic proc.	69.0637066	0.04002621
Drought	unique	Susceptible	up	GO:0008610	Lipid biosynthetic proc.	4.65547531	0.03500824
Drought	common	Resistant and Susceptible	up	GO:0009631	Cold acclimation	43.6067772	5.47E-06
Drought	unique	Resistant	down	KEGG: osa00904	Diterpenoid biosynthesis	31.162892	0.00535764
Drought	unique	Susceptible	down	GO:0009768	Photosynthesis, light harvesting in photosystem I	23.9297659	0.03759248
Drought	common	Resistant and Susceptible	down	GO:0016540	Protein autoprocessing	96.9512195	0.00545431

Interestingly, the enrichment analysis showed that the most significantly enriched term for both the upregulated genes in the salt susceptible and salt resistant groups as well as the drought susceptible and drought resistant groups was "GO:0009631 Cold acclimation." This suggests that the importance of stress mechanisms of cold acclimation may be similar to that of both salinity and drought stress, regardless of the *Oryza sativa* phenotype.

In addition, in Figure 3(a), the most significantly enriched term in the enrichment analysis of the upregulated combine DEGs was "diterpenoid metabolic processes." In contrast, the most significantly

enriched term in the genes specifically downregulated in the drought-resistant phenotype was "diterpenoid biosynthesis (KEGG: osa00904)." This suggests that the pathways involving diterpenoids may regulate stress responses by upregulating or downregulating the expression of related genes. All gene lists, Venn diagrams visualizing overlapping genes, and enrichment results are shown in Supplementary Tables (Supplementary Table 9 and 10).

2.6 Selection of genes common among different stress conditions

The overlap of genes that were commonly upregulated or downregulated across the four categories based on the two stress conditions and two phenotypes each was visualized using UpSet plots (Figure 4). All overlapping genes are listed in Supplementary Table 11. The number of genes whose expression specifically increased or decreased in each phenotype under both stress conditions is summarized in Table 6. Among these genes, those whose expression were upregulated in the resistant and susceptible varieties are summarized in Table 7. In total, 10 genes were consistently upregulated in the resistant phenotype under both stress conditions. These included genes involved in various cellular processes, such as transcription regulation (*Oshox25* and *PHR3*), secondary metabolism (*OsOSC4* and *OsPSY*), and peroxidase (*prx130*).

Figure 4. UpSet plots showing the overlap and specificity of DEGs in *Oryza sativa* under various stress conditions and phenotypes. UpSet plots illustrating the overlap of commonly or specifically regulated genes across the four categories combining salt and drought stress conditions with resistant and susceptible phenotypes. (a) Upregulated genes; (b) downregulated genes

All overlapping genes are listed in Supplementary Table 11. The number of genes whose expression specifically increased or decreased in each phenotype under both stress conditions is summarized in Table 6.

Table 6. *Oryza sativa* genes that were specifically regulated in the resistant or susceptible phenotype under both salt and drought stress conditions

Phenotype	Regulation	Number of Genes
Resistant	Upregulated	10
Resistant	Downregulated	6
Susceptible	Upregulated	12
Susceptible	Downregulated	34

Among these genes, those whose expression were upregulated in the resistant and susceptible varieties are summarized in Table 7.

Table 7. List of *Oryza sativa* genes that were specifically upregulated in the resistant or susceptible phenotype under both salt and drought stress conditions

Overlap	Total number of genes	Os-Gene stable ID	Os-Gene name	Os-Gene description
Drought_Resistant _up Salt_Resistant_up	10	Os12g0 594000	OsW D40- 198	WD40 repeat-like domain containing protein
		Os09g0 379600	Osho x25	Similar to Homeobox-leucine zipper protein HOX25
		Os03g0 709300	OsUC L9	Similar to Chemocyanin precursor (Basic blue protein) (Plantacyanin)
		Os02g0 140400	OsOS C4	Similar to Beta-amyrin synthase
		Os02g0 139000	PHR3	Transcription factor, Regulation of Pi signaling and homeostasis, Tolerance to low-Pi stress
		Os08g0 290700	-	Winged helix repressor DNA-binding domain containing protein
		Os11g0 112400	prx13 0	Peroxidase (EC 1.11.1.7)
		Os01g0 767600	-	Conserved hypothetical protein
		Os09g0 555500	OsPS Y	Similar to Chloroplast phytoene synthase 3

		Os01g0 155800	-	Conserved hypothetical protein
Drought_Susceptible_up Salt_Susceptible_up	12	Os01g0 159800	OsbHLH035	Similar to DNA binding protein
		Os05g0 119100	-	Hypothetical protein
		Os02g0 209400	-	Conserved hypothetical protein
		Os10g0 450900	GRP	Similar to Glycine-rich cell wall structural protein 2 precursor
		Os10g0 517500	RRJ1	Cys/Met metabolism, pyridoxal phosphate-dependent enzyme domain containing protein
		Os12g0 510750	-	Conserved hypothetical protein
		Os09g0 278000	-	Hypothetical conserved gene
		Os07g0 673900	HIGD2	Hypoxia induced protein, Early stage of hypoxia signaling
		Os11g0 533400	-	Conserved hypothetical protein
		Os11g0 700900	C10923	Glycoside hydrolase, subgroup, catalytic core domain containing protein
		Os03g0 305100	-	Similar to AMP-binding protein
		Os03g0 184550	-	Similar to Dihydroflavonol-4-reductase

In total, 10 genes were consistently upregulated in the resistant phenotype under both stress conditions. These included genes involved in various cellular processes, such as transcription regulation (*Oshox25* and *PHR3*), secondary metabolism (*OsOSC4* and *OsPSY*), and peroxidase (*prx130*).

In contrast, 12 genes were consistently upregulated in the susceptible phenotype under both salt and drought stress conditions. These genes have different functional profiles, including a bHLH transcription factor (*OsbHLH035*), glycine-rich cell wall structural protein (*GRP*), hypoxia-

induced gene (*HIGD2*), and enzymes involved in various metabolic processes (*RRJ1* and *C10923*).

2.7 Selection of genes involved in stress response mechanisms conserved across different plant species

  In a previous study, we performed a meta-analysis of *Arabidopsis thaliana* under ABA, salt, and dehydration conditions and identified genes that were upregulated or downregulated under each condition [37]. Here, we used *Arabidopsis thaliana* orthologous genes corresponding to the *Oryza sativa* genes identified in this study to determine whether any genes shared common. The overlap in expression of commonly regulated genes was visualized using UpSet plots (Supplementary Figure 4). Satisfying the above criteria, 11 genes were upregulated (At_ABA_up, At_Dehydration_up, At_Salt_up, Os_Drought_Resistant_up, Os_Drought_Susceptible_up, Os_Salt_Resistant_up, and Os_Salt_Susceptible_up) and 1 gene was downregulated (At_ABA_down, At_Dehydration_down, At_Salt_down, Os_Drought_Resistant_down, Os_Drought_Susceptible_down, Os_Salt_Resistant_down, and Os_Salt_Susceptible_down) under salt and drought conditions (Table 8). These genes may be involved in the stress response mechanisms that are common across different plant species.

Table 8. Comparison of ABA, salt, and drought stress-responsive genes conserved among different plant species

Overlap	Total number of genes	At-Gene stable ID	At-Gene Name	At-Description
At_ABA_up At_Dehydration_up At_Salt_up Os_Drought_Resistant_up Os_Drought_Susceptible_up Os_Salt_Resistant_up Os_Salt_Susceptible_up	11	AT1G52690	LEA7	Late embryogenesis abundant protein (LEA) family protein
		AT3G03341	-	cold-regulated protein
		AT1G07430	HAI2	highly ABA-induced PP2C protein 2
		AT2G35300	LEA18	Late embryogenesis abundant protein, group 1 protein

		AT3G48510	-	AtIII18x5-like protein
		AT2G39050	EULS3	hydroxyproline-rich glycoprotein family protein
		AT3G24520	HSFC1	heat shock transcription factor C1
		AT2G46270	GBF3	G-box binding factor 3
		AT5G66400	RAB18	Dehydrin family protein
		AT2G47770	TSPO	TSPO(outer membrane tryptophan-rich sensory protein)-like protein
		AT2G46680	HB-7	homeobox 7
At_ABA_down At_Dehydration_down At_Salt_down Os_Drought_Resistant_down Os_Drought_Susceptible_down Os_Salt_Resistant_down Os_Salt_Susceptible_down	1	AT5G05440	PYL5	Polyketide cyclase/dehydrase and lipid transport superfamily protein
At_ABA_up At_Dehydration_up At_Salt_up Os_Drought_Resistant_up Os_Salt_Resistant_up	6	AT4G19230	CYP707A1	cytochrome P450, family 707, subfamily A, polypeptide 1
		AT1G60190	PUB19	ARM repeat superfamily protein
		AT1G45249	ABF2	abscisic acid responsive elements-binding factor 2
		AT3G63060	EDL3	EID1-like 3

				senescence regulator (Protein of unknown function, DUF584)
		AT1G29640	-	
		AT1G48000	MYB112	myb domain protein 112
At_ABA_down At_Dehydration_down At_Salt_down Os_Drought_Resistant_down Os_Salt_Resistant_down	1	AT1G80050	APT2	adenine phosphoribosyl transferase 2

For the *Oryza sativa* genes in the 'resistant' phenotype category used in this study, six genes were upregulated (At_ABA_up, At_Dehydration_up, At_Salt_up, Os_Drought_Resistant_up, and Os_Salt_Resistant_up) and one gene was downregulated (At_ABA_down, At_Dehydration_down, At_Salt_down, Os_Drought_Resistant_down, and Os_Salt_Resistant_down) was downregulated under salt and drought conditions (Table 8). These genes may be involved in stress responses that are common across plant species, particularly in mechanisms important for stress tolerance.

The list of *Oryza sativa* gene IDs identified in this study was converted to *Arabidopsis thaliana* IDs and compared with previous studies. All gene lists are available online (Supplementary Table 12).

3. Discussion

This study aimed to identify novel genes and potential pathways involved in drought and salt stress responses in *Oryza sativa* using a meta-analysis of publicly available RNA-Seq data. By integrating datasets from multiple studies covering different *Oryza sativa* cultivars and stress conditions, we aimed to reveal robust and consistent gene expression patterns associated with stress resistance or susceptibility. Analysis of the tissue types of the collected samples revealed that while salt and drought stress are expected to have significant effects on the roots, the majority of the affected samples in this study consisted of "leaf" and "shoot" tissues, with a limited number of "root-"specific samples. Although this study focused on phenotypes and stress types rather than tissue specificity, tissue-specific analyses will likely become possible in the future as the amount of publicly available RNA-Seq data increases.

This study focused on several *Oryza sativa* cultivars that exhibited different phenotypic responses to salt and drought stress. The phenotypic classifications of "resistant" and "susceptible" used in this study are based solely on existing research reports and may not fully capture the nuanced stress tolerance levels of each cultivar. In addition, differences in stress treatment methods among the studies should be considered.

To minimize the impact of these differences and provide a robust comparative analysis of multiple datasets, we adopted the following approach. First, we directly compared the gene expression ratios between stressed and control samples in the same study. Subsequently, we calculated the TN score, which facilitated integration and comparative analysis of the data across multiple studies.

Based on the TN score, DEGs were identified for each dataset and categorized by stress type and phenotype. Enrichment analysis of each DEG set revealed the distinct and characteristic terms associated with each DEG group. Furthermore, we performed an enrichment analysis of genes that were upregulated or downregulated, irrespective of the phenotypic classification or stress type. The most significantly enriched term for upregulated genes was "diterpenoid metabolic process," with 13 genes included in this category (Figure 3(a)). Diterpenoid phytoalexins have been shown to be potentially involved in the resistance of cereal plants to environmental stresses, such as drought [39]. The genes are listed in Supplementary Table 13.

These genes included those whose functions in stress mechanisms have been elucidated in previous studies. *OsCPS4* encodes a key enzyme that contributes to the biosynthesis of labdane-related diterpenoid (LRD) phytoalexins in *Oryza sativa*. Mutant analysis showed that the loss of function of *OsCPS4* significantly increased sensitivity to drought stress, suggesting that LRD biosynthesis involving *OsCPS4* contributes to the regulation of stomatal closure, independent of the ABA pathway [40]. The genes with unknown functions listed in this study may include those that contribute to the stress response through their involvement in LRD metabolism. Interestingly, in addition to the upregulated DEGs, the term enriched in genes specifically downregulated in the drought stress susceptible samples was "diterpenoid biosynthesis (KEGG: osa00904)" (Supplementary Figure 3 (d), Supplementary Table 13). This contrasting expression pattern suggests a complex regulatory mechanism in the diterpenoid pathway. In other words, the genes in this pathway may show diverse expression patterns depending on the type of stress, plant phenotype, and cultivar.

We also focused on a group of genes whose expression was upregulated in both the resistant and susceptible cultivars under the two stresses. Some of the genes commonly upregulated under salt and drought stress in the resistant phenotype included (*OsPSY3* and *Os09g0555500*). *PSY* is a rate-limiting enzyme in carotenoid synthesis and represents a precursor required for ABA synthesis in various plants [41]. *OsPSY* has been shown to be upregulated by salt and drought treatments [42–44]. Therefore, the selection results were consistent with those of existing studies. Other upregulated genes included transcription factor *OsPHR3* (*Os02g0139000*), whose expression is induced by inorganic phosphate (Pi) deficiency and plays a role in improving tolerance to Pi deficiency as well as regulating nitrogen (N) homeostasis [45–47]. Deficiency of Pi or N induces various morphophysiological adaptive responses and gene expression regulation in plants [48–51]. Although several studies have suggested a relationship between Pi and N deficiency and the ABA pathway and its biosynthesis, many aspects remain unclear and further research is needed [52–56].

Finally, a comparison with previous *Arabidopsis* meta-analysis data revealed several genes whose expression was altered in *Oryza sativa* and *Arabidopsis thaliana* under both salt and drought stress conditions (Table 8). To focus on genes involved in mechanisms commonly conserved between plant species regardless of the stress phenotype, we focused on genes shared by seven DEGs, with three in *Arabidopsis thaliana* and four in *Oryza sativa*. For example, *HSFC1* (*AT3G24520*) represented a gene commonly upregulated in both species, regardless of phenotype.

Previous studies on *Arabidopsis thaliana* have suggested that the expression of *HSFC1* is involved in cold responses via a pathway independent of the C-repeat binding factor (CBF), a well-known transcription factor that controls cold acclimation in plants [57,58].

The ortholog of *Arabidopsis HSFC1* gene in *Oryza sativa* was *HSfC2b* (*Os06g0553100*). However, the functions and roles of these genes under salinity and drought conditions remain unclear. Heat shock factors (HSFs) that contribute to environmental stress tolerance occur in both *Arabidopsis thaliana* and *Oryza sativa* [59,60]. Given the consistent upregulation of *HSFC1* and its ortholog *HSfC2b* across different stress conditions and plant species, these genes are promising candidates for further investigation.

This study had some limitations. First, because the identification of DEGs was based on the selection of TN scores rather than statistical tests, the results should be interpreted with caution. Second, the RNA-Seq data used in this study were collected under different growth and experimental conditions. Third, experimental validation using plant samples exposed to specific stress environments was not performed, indicating that further research is required to elucidate the functions of these genes. These limitations should be considered when interpreting our results and planning future studies.

4. Materials and methods

4.1 Manual curation of gene expression data from public databases

RNA-Seq data were collected from the National Center for Biotechnology Information Gene Expression Omnibus (NCBI GEO) (https://www.ncbi.nlm.nih.gov/geo/) [61] and European Bioinformatics Institute BioStudies (EBI BioStudies) (https://www.ebi.ac.uk/biostudies/) [62]. In the NCBI GEO, a comprehensive search was performed using the following query: ("drought"[All Fields] OR "water deficit"[All Fields] OR "salt"[All Fields] OR "NaCl"[All Fields] OR "salinity"[All Fields]) AND "*Oryza sativa*"[porgn] AND "Expression profiling by high throughput sequencing"[Filter].

In the EBI BioStudies, projects related to stress treatments were selected by searching with the following parameters: "Keywords: 'RNA-Seq' AND '*Oryza sativa*' AND ('salt' OR 'drought')," "Collection: 'Array Express'."

Manual curation was performed based on the search results. The manual curation process included collecting metadata from the database, such as stress treatment conditions and sample tissues, along with the raw RNA-Seq data. Based on the descriptions in the papers and database metadata, the phenotype of each cultivar was classified as either "resistant" or "susceptible" to stress. Another criterion was the presence of paired stress-treated and stress-untreated control samples in the same project. This process resulted in the creation of 105 paired datasets from 202 samples from 11 projects.

4.2 Gene expression quantification

FASTQ-formatted files for each RNA-Seq run accession number were obtained using the prefetch and fastq-dump commands of the SRA Toolkit (v3.0.0) [https://github.com/ncbi/sra-tools]. Quality control of the raw reads was performed using Trim Galore (v0.6.7) [https://github.com/FelixKrueger/TrimGalore]. Transcript quantification was performed using Salmon (v1.8.0) [https://github.com/COMBINE-lab/salmon] [38], and the cDNA sequences downloaded from Ensembl Plants (release 56) (Oryza_sativa.IRGSP-1.0. cdna.all.fa) were used as reference. To ensure data consistency and enable accurate gene expression comparisons, we used the *Oryza sativa japonica* reference transcriptome for expression quantification in both *japonica* and *indica* datasets. This approach enhances comparability and unifies annotations. We chose the *japonica* reference because of its more comprehensive and well-curated gene annotations, which provided a rich foundation for functional analysis across both subspecies. As a result, the quantitative RNA-Seq data were calculated as the TPM. Transcript-level TPM values were summarized at the gene level using tximport (v1.26.1) [https://bioconductor.org/packages/release/bioc/html/tximport.html].

4.3 TN ratio and score calculation

Gene expression data from different experiments were normalized by calculating the TN ratio, which represents the ratio of gene expression between stress-treated (T) and non-treated (N) samples. If the TN ratio was higher than the threshold, the gene was considered to be upregulated; whereas if it was less than the reciprocal of the threshold, the gene was considered downregulated. Otherwise, the gene was considered unchanged. To classify the upregulated and downregulated genes, 1.5-fold, 2-fold, 5-fold, and 10-fold thresholds were evaluated, and a 2-fold threshold was adopted. Therefore, genes with a TN ratio higher

than 2 were classified as upregulated, while genes with a TN ratio lower than 0.5 were classified as downregulated. The TN score of each gene was determined by subtracting the number of downregulated experiments from the number of upregulated experiments to assess changes in gene expression under stress conditions across multiple experiments. The TN ratio and TN score were computed using code that had been utilized in a previous study [63].

4.4 Gene set enrichment analysis

Gene set enrichment analysis was performed using the web tool ShinyGO (v0.77) [http://bioinformatics.sdstate.edu/go77/] [64] to analyze the DEG sets. For the *Oryza sativa* analysis, the species was set to "*Oryza sativa Japonica* Group," and the pathway database was set to "GO Biological Process." All other parameters were set to the default settings.

4.5 Visualization

A web-based Venn diagram tool [https://bioinformatics.psb.ugent.be/webtools/Venn/] and the UpSet plot tool [https://asntech.shinyapps.io/intervene/] [65] were used to search for and visualize overlapping genes.

4.6 Functional annotation of Oryza sativa genes

To functionally annotate the *Oryza sativa* genes, we retrieved the "Gene ID," "Gene name," and "Gene description" for each gene from the Ensembl Plants (release 56, *Oryza sativa Japonica* Group genes (IRGSP-1.0)). We then identified the putative orthologs of these genes in *Arabidopsis thaliana* by performing a BLASTP (v2.12.0) search using the *Oryza sativa* protein sequences (*Oryza_sativa*.IRGSP-1.0.pep.all.fa.gz) as queries against the *Arabidopsis thaliana* protein database obtained from Ensembl Plants (release 56, *Arabidopsis_thaliana*.TAIR10.pep.all.fa.gz). The BLASTP search was conducted with an E-value cutoff of 1e-10. For each *Oryza sativa* gene, the top hit in the *Arabidopsis thaliana* protein database was considered the putative ortholog. The corresponding "Gene ID," "Gene name," and "Gene description" for the *Arabidopsis thaliana* orthologs were obtained from the Ensembl Plants (release 56, *Arabidopsis thaliana* genes (TAIR10)).

5. Conclusion

Through a meta-analysis of publicly available RNA-Seq data, we obtained a list of potential target genes related to drought and salt stress responses in *Oryza sativa* and identified candidate genes that may influence stress-related phenotypes. These findings provide a valuable resource for future studies aimed at understanding the molecular mechanisms underlying stress resistance in rice and developing new strategies to improve crop productivity under adverse environmental conditions.

Supplementary Materials: The Supplementary Material for this article can be found online at https://doi.org/10.6084/m9.figshare.25547083.v1.

Author Contributions: Conceptualization, M.S. and H.B.; Methodology, M.S. and H.B.; Software, M.S. and H.B.; Validation, M.S. and H.B.; Formal Analysis, M.S. and H.B.; Investigation, M.S.; Resources, M.S. and H.B.; Data Curation, M.S.; Writing — Original Draft Preparation, M.S.; Writing — Review and Editing, M.S. and H.B.; Visualization, M.S.; Supervision, H.B.; Project Administration, H.B.; Funding Acquisition, H.B. All authors have read and agreed to the published version of the manuscript.

Funding: The authors declare that they received financial support for the research, authorship, and publication of this article. This work was supported by the Center of Innovation for Bio-Digital Transformation (BioDX), an open innovation platform for industry–academia cocreation (COI-NEXT) and the Japan Science and Technology Agency (JST, COI-NEXT, JPMJPF2010).

Data Availability Statement: The data presented in this study are publicly available at Figshare (https://doi.org/10.6084/m9.figshare.25547083.v1).

Acknowledgments: Computations were performed at the Hiroshima University Genome Editing Innovation Center.

Conflicts of Interest: The authors declare that this study was conducted in the absence of any commercial or financial relationships that could be construed as potential conflicts of interest.

References

1. Sharif, M.K.; Butt, M.S.; Anjum, F.M.; Khan, S.H. Rice Bran: A Novel Functional Ingredient. *Crit. Rev. Food Sci. Nutr.* **2014**, *54*, 807–816.

2. Mohidem, N.A.; Hashim, N.; Shamsudin, R.; Che Man, H. Rice for Food Security: Revisiting Its Production, Diversity, Rice Milling Process and Nutrient Content. *Collect. FAO Agric.* **2022**, *12*, 741.

3. Billah, M.; Aktar, S.; Brestic, M.; Zivcak, M.; Khaldun, A.B.M.; Uddin, M.S.; Bagum, S.A.; Yang, X.; Skalicky, M.; Mehari, T.G.; et al. Progressive Genomic Approaches to Explore Drought- and Salt-Induced Oxidative Stress Responses in Plants under Changing Climate. *Plants* **2021**, *10*, doi:10.3390/plants10091910.

4. Ma, Y.; Dias, M.C.; Freitas, H. Drought and Salinity Stress Responses and Microbe-Induced Tolerance in Plants. *Front. Plant Sci.* **2020**, *11*, 591911.

5. Angon, P.B.; Tahjib-Ul-Arif, M.; Samin, S.I.; Habiba, U.; Hossain, M.A.; Brestic, M. How Do Plants Respond to Combined Drought and Salinity Stress?-A Systematic Review. *Plants* **2022**, *11*, doi:10.3390/plants11212884.

6. Yang, Y.; Guo, Y. Elucidating the Molecular Mechanisms Mediating Plant Salt-Stress Responses. *New Phytol.* **2018**, *217*, 523–539.

7. Kamran, M.; Parveen, A.; Ahmar, S.; Malik, Z.; Hussain, S.; Chattha, M.S.; Saleem, M.H.; Adil, M.; Heidari, P.; Chen, J.-T. An Overview of Hazardous Impacts of Soil Salinity in Crops, Tolerance Mechanisms, and Amelioration through Selenium Supplementation. *Int. J. Mol. Sci.* **2019**, *21*, doi:10.3390/ijms21010148.

8. Liu, C.; Mao, B.; Yuan, D.; Chu, C.; Duan, M. Salt Tolerance in Rice: Physiological Responses and Molecular Mechanisms. *The Crop Journal* **2022**, *10*, 13–25.

9. Ahanger, M.A.; Agarwal, R.M. Salinity Stress Induced Alterations in Antioxidant Metabolism and Nitrogen Assimilation in Wheat (Triticum Aestivum L) as Influenced by Potassium Supplementation. *Plant Physiol. Biochem.* **2017**, *115*, 449–460.

10. Munns, R.; Tester, M. Mechanisms of Salinity Tolerance. *Annu. Rev. Plant Biol.* **2008**, *59*, 651–681.

11. Jiang, C.; Zheng, Q.; Liu, Z.; Xu, W.; Liu, L.; Zhao, G.; Long, X. Overexpression of Arabidopsis Thaliana Na+/H+ Antiporter Gene Enhanced Salt Resistance in Transgenic Poplar (Populus × Euramericana 'Neva'). *Trees* **2012**, *26*, 685–694.

12. Mittal, S.; Kumari, N.; Sharma, V. Differential Response of Salt Stress on Brassica Juncea: Photosynthetic Performance, Pigment, Proline, D1 and Antioxidant Enzymes. *Plant Physiol. Biochem.* **2012**, *54*, 17–26.

13. Gengmao, Z.; Shihui, L.; Xing, S.; Yizhou, W.; Zipan, C. The Role of Silicon in Physiology of the Medicinal Plant (Lonicera Japonica L.) under Salt Stress. *Sci. Rep.* **2015**, *5*, 12696.

14. Impacts of Soil Salinity/Sodicity on Soil-Water Relations and Plant Growth in Dry Land Areas: A Review. *Journal of Natural Sciences Research* **2021**, doi:10.7176/jnsr/12-3-01.

15. Gowda, V.R.P.; Henry, A.; Yamauchi, A.; Shashidhar, H.E.; Serraj, R. Root Biology and Genetic Improvement for Drought Avoidance in Rice. *Field Crops Res.* **2011**, *122*, 1–13.

16. Yue, B.; Xiong, L.; Xue, W.; Xing, Y.; Luo, L.; Xu, C. Genetic Analysis for Drought Resistance of Rice at Reproductive Stage in Field with Different Types of Soil. *Theor. Appl. Genet.* **2005**, *111*, 1127–1136.

17. Zou, G.-H.; Liu, H.-Y.; Mei, H.-W.; Liu, G.-L.; Yu, X.-Q.; Li, M.-S.; Wu, J.-H.; Chen, L.; Luo, L.-J. Screening for Drought Resistance of Rice Recombinant Inbred Populations in the Field. *J. Integr. Plant Biol.* **2007**, *49*, 1508–1516.

18. Liu, G.; Mei, H.; Liu, H.; Yu, X.; Zou, G.; Luo, L. Sensitivities of Rice Grain Yield and Other Panicle Characters to Late-Stage Drought Stress Revealed by Phenotypic Correlation and QTL Analysis. *Mol. Breed.* **2010**, *25*, 603–613.

19. Ekanayake, I.J.; Datta, S.K.D.E.; Steponkus, P.L. Spikelet Sterility and Flowering Response of Rice to Water Stress at Anthesis. *Ann. Bot.* **1989**, *63*, 257–264.

20. Ouyang, S.-Q.; Liu, Y.-F.; Liu, P.; Lei, G.; He, S.-J.; Ma, B.; Zhang, W.-K.; Zhang, J.-S.; Chen, S.-Y. Receptor-like Kinase OsSIK1 Improves Drought and Salt Stress Tolerance in Rice (Oryza Sativa) Plants. *Plant J.* **2010**, *62*, 316–329.

21. Wang, X.; Liu, H.; Yu, F.; Hu, B.; Jia, Y.; Sha, H.; Zhao, H. Differential Activity of the Antioxidant Defence System and Alterations in the Accumulation of Osmolyte and Reactive Oxygen Species under Drought Stress and Recovery in Rice (Oryza Sativa L.) Tillering. *Sci. Rep.* **2019**, *9*, 8543.

22. Panda, D.; Mishra, S.S.; Behera, P.K. Drought Tolerance in Rice: Focus on Recent Mechanisms and Approaches. *Rice Sci.* **2021**, *28*, 119–132.

23. Hassan, M.A.; Dahu, N.; Hongning, T.; Qian, Z.; Yueming, Y.; Yiru, L.; Shimei, W. Drought Stress in Rice: Morpho-Physiological and Molecular Responses and Marker-Assisted Breeding. *Front. Plant Sci.* **2023**, *14*, 1215371.

24. Wang, R.; Jing, W.; Xiao, L.; Jin, Y.; Shen, L.; Zhang, W. The Rice High-Affinity Potassium Transporter1;1 Is Involved in Salt Tolerance and Regulated by an MYB-Type Transcription Factor. *Plant Physiol.* **2015**, *168*, 1076–1090.

25. Liu, C.; Mao, B.; Ou, S.; Wang, W.; Liu, L.; Wu, Y.; Chu, C.; Wang, X. OsbZIP71, a BZIP Transcription Factor, Confers Salinity and Drought Tolerance in Rice. *Plant Mol. Biol.* **2014**, *84*, 19–36.

26. Manimaran, P.; Venkata Reddy, S.; Moin, M.; Raghurami Reddy, M.; Yugandhar, P.; Mohanraj, S.S.; Balachandran, S.M.; Kirti, P.B. Activation-Tagging in Indica Rice Identifies a Novel Transcription Factor Subunit, NF-YC13 Associated with Salt Tolerance. *Sci. Rep.* **2017**, *7*, 9341.

27. Sripinyowanich, S.; Klomsakul, P.; Boonburapong, B.; Bangyeekhun, T.; Asami, T.; Gu, H.; Buaboocha, T.; Chadchawan, S. Exogenous ABA Induces Salt Tolerance in Indica Rice (Oryza Sativa L.): The Role of OsP5CS1 and OsP5CR Gene Expression during Salt Stress. *Environ. Exp. Bot.* **2013**, *86*, 94–105.

28. Campo, S.; Baldrich, P.; Messeguer, J.; Lalanne, E.; Coca, M.; San Segundo, B. Overexpression of a Calcium-Dependent Protein Kinase Confers Salt and Drought Tolerance in Rice by Preventing Membrane Lipid Peroxidation. *Plant Physiol.* **2014**, *165*, 688–704.

29. Asano, T.; Hayashi, N.; Kobayashi, M.; Aoki, N.; Miyao, A.; Mitsuhara, I.; Ichikawa, H.; Komatsu, S.; Hirochika, H.; Kikuchi, S.; et al. A Rice Calcium-Dependent Protein Kinase OsCPK12 Oppositely Modulates Salt-Stress Tolerance and Blast Disease Resistance. *Plant J.* **2012**, *69*, 26–36.

30. Sun, S.-J.; Guo, S.-Q.; Yang, X.; Bao, Y.-M.; Tang, H.-J.; Sun, H.; Huang, J.; Zhang, H.-S. Functional Analysis of a Novel Cys2/His2-Type Zinc Finger Protein Involved in Salt Tolerance in Rice. *J. Exp. Bot.* **2010**, *61*, 2807–2818.

31. Huang, J.; Sun, S.; Xu, D.; Lan, H.; Sun, H.; Wang, Z.; Bao, Y.; Wang, J.; Tang, H.; Zhang, H. A TFIIIA-Type Zinc Finger Protein Confers Multiple Abiotic Stress Tolerances in Transgenic Rice (Oryza Sativa L.). *Plant Mol. Biol.* **2012**, *80*, 337–350.

32. Wei, H.; Chen, C.; Ma, X.; Zhang, Y.; Han, J.; Mei, H.; Yu, S. Comparative Analysis of Expression Profiles of Panicle Development among Tolerant and Sensitive Rice in Response to Drought Stress. *Front. Plant Sci.* **2017**, *8*, 437.

33. Jeong, J.S.; Kim, Y.S.; Baek, K.H.; Jung, H.; Ha, S.-H.; Do Choi, Y.; Kim, M.; Reuzeau, C.; Kim, J.-K. Root-Specific Expression of OsNAC10 Improves Drought Tolerance and Grain Yield in Rice under Field Drought Conditions. *Plant Physiol.* **2010**, *153*, 185–197.

34. Jeong, J.S.; Kim, Y.S.; Redillas, M.C.F.R.; Jang, G.; Jung, H.; Bang, S.W.; Choi, Y.D.; Ha, S.-H.; Reuzeau, C.; Kim, J.-K. OsNAC5 Overexpression Enlarges Root Diameter in Rice Plants Leading to Enhanced Drought Tolerance and Increased Grain Yield in the Field. *Plant Biotechnol. J.* **2013**, *11*, 101–114.

35. Cui, M.; Zhang, W.; Zhang, Q.; Xu, Z.; Zhu, Z.; Duan, F.; Wu, R. Induced Over-Expression of the Transcription Factor OsDREB2A Improves Drought Tolerance in Rice. *Plant Physiol. Biochem.* **2011**, *49*, 1384–1391.

36. Matsukura, S.; Mizoi, J.; Yoshida, T.; Todaka, D.; Ito, Y.; Maruyama, K.; Shinozaki, K.; Yamaguchi-Shinozaki, K. Comprehensive Analysis of Rice DREB2-Type Genes That Encode Transcription Factors Involved in the Expression of Abiotic Stress-Responsive Genes. *Mol. Genet. Genomics* **2010**, *283*, 185–196.

37. Shintani, M.; Tamura, K.; Bono, H. Meta-Analysis of Public RNA Sequencing Data of Abscisic Acid-Related Abiotic Stresses in Arabidopsis Thaliana. *Front. Plant Sci.* **2024**, *15*, 1343787.

38. Patro, R.; Duggal, G.; Love, M.I.; Irizarry, R.A.; Kingsford, C. Salmon Provides Fast and Bias-Aware Quantification of Transcript Expression. *Nat. Methods* **2017**, *14*, 417–419.

39. Murphy, K.M.; Zerbe, P. Specialized Diterpenoid Metabolism in Monocot Crops: Biosynthesis and Chemical Diversity. *Phytochemistry* **2020**, *172*, 112289.

40. Zhang, J.; Li, R.; Xu, M.; Hoffmann, R.I.; Zhang, Y.; Liu, B.; Zhang, M.; Yang, B.; Li, Z.; Peters, R.J. A (Conditional) Role for Labdane-Related Diterpenoid Natural Products in Rice Stomatal Closure. *New Phytol.* **2021**, *230*, 698–709.

41. Zhou, X.; Rao, S.; Wrightstone, E.; Sun, T.; Lui, A.C.W.; Welsch, R.; Li, L. Phytoene Synthase: The Key Rate-Limiting Enzyme of Carotenoid Biosynthesis in Plants. *Front. Plant Sci.* **2022**, *13*, 884720.

42. Welsch, R.; Wüst, F.; Bär, C.; Al-Babili, S.; Beyer, P. A Third Phytoene Synthase Is Devoted to Abiotic Stress-Induced Abscisic Acid Formation in Rice and Defines Functional Diversification of Phytoene Synthase Genes. *Plant Physiol.* **2008**, *147*, 367–380.

43. Li, F.; Vallabhaneni, R.; Wurtzel, E.T. PSY3, a New Member of the Phytoene Synthase Gene Family Conserved in the Poaceae and Regulator of Abiotic Stress-Induced Root Carotenogenesis. *Plant Physiol.* **2008**, *146*, 1333–1345.

44. Du, H.; Chang, Y.; Huang, F.; Xiong, L. GID1 Modulates Stomatal Response and Submergence Tolerance Involving Abscisic Acid and Gibberellic Acid Signaling in Rice. *J. Integr. Plant Biol.* **2015**, *57*, 954–968.

45. Guo, M.; Ruan, W.; Li, C.; Huang, F.; Zeng, M.; Liu, Y.; Yu, Y.; Ding, X.; Wu, Y.; Wu, Z.; et al. Integrative Comparison of the Role of the PHOSPHATE RESPONSE1 Subfamily in Phosphate Signaling and Homeostasis in Rice. *Plant Physiol.* **2015**, *168*, 1762–1776.

46. Ruan, W.; Guo, M.; Wu, P.; Yi, K. Phosphate Starvation Induced OsPHR4 Mediates Pi-Signaling and Homeostasis in Rice. *Plant Mol. Biol.* **2017**, *93*, 327–340.

47. Sun, Y.; Luo, W.; Jain, A.; Liu, L.; Ai, H.; Liu, X.; Feng, B.; Zhang, L.; Zhang, Z.; Guohua, X.; et al. OsPHR3 Affects the Traits Governing Nitrogen Homeostasis in Rice. *BMC Plant Biol.* **2018**, *18*, 241.

48. Bechtaoui, N.; Rabiu, M.K.; Raklami, A.; Oufdou, K.; Hafidi, M.; Jemo, M. Phosphate-Dependent Regulation of Growth and Stresses Management in Plants. *Front. Plant Sci.* **2021**, *12*, 679916.

49. Khan, F.; Siddique, A.B.; Shabala, S.; Zhou, M.; Zhao, C. Phosphorus Plays Key Roles in Regulating Plants' Physiological Responses to Abiotic Stresses. *Plants* **2023**, *12*, 2861.

50. Sakuraba, Y. Molecular Basis of Nitrogen Starvation-Induced Leaf Senescence. *Front. Plant Sci.* **2022**, *13*, 1013304.

51. Xing, J.; Cao, X.; Zhang, M.; Wei, X.; Zhang, J.; Wan, X. Plant Nitrogen Availability and Crosstalk with Phytohormones Signallings and Their Biotechnology Breeding Application in Crops. *Plant Biotechnol. J.* **2023**, *21*, 1320–1342.

52. Fang Zhu, X.; Sheng Zhao, X.; Wu, Q.; Fang Shen, R. Abscisic Acid Is Involved in Root Cell Wall Phosphorus Remobilization Independent of Nitric Oxide and Ethylene in Rice (Oryza Sativa). *Ann. Bot.* **2018**, *121*, 1361–1368.

53. Zhang, Y.; Li, T.-T.; Wang, L.-F.; Guo, J.-X.; Lu, K.-K.; Song, R.-F.; Zuo, J.-X.; Chen, H.-H.; Liu, W.-C. Abscisic Acid Facilitates Phosphate Acquisition through the Transcription Factor ABA INSENSITIVE5 in Arabidopsis. *Plant J.* **2022**, *111*, 269–281.

54. Jaskolowski, A.; Poirier, Y. Phosphate Deficiency Increases Plant Susceptibility to Botrytis Cinerea Infection by Inducing the Abscisic Acid Pathway. *Plant J.* **2024**, *119*, 828–843.

55. Hsieh, P.-H.; Kan, C.-C.; Wu, H.-Y.; Yang, H.-C.; Hsieh, M.-H. Early Molecular Events Associated with Nitrogen Deficiency in Rice Seedling Roots. *Sci. Rep.* **2018**, *8*, 12207.

56. Zakari, S.A.; Asad, M.-A.-U.; Han, Z.; Zhao, Q.; Cheng, F. Relationship of Nitrogen Deficiency-Induced Leaf Senescence with ROS Generation and ABA Concentration in Rice Flag Leaves. *J. Plant Growth Regul.* **2020**, *39*, 1503–1517.

57. Zhao, C.; Zhang, Z.; Xie, S.; Si, T.; Li, Y.; Zhu, J.-K. Mutational Evidence for the Critical Role of CBF Transcription Factors in Cold Acclimation in Arabidopsis. *Plant Physiol.* **2016**, *171*, 2744–2759.

58. Jia, Y.; Ding, Y.; Shi, Y.; Zhang, X.; Gong, Z.; Yang, S. The Cbfs Triple Mutants Reveal the Essential Functions of CBFs in Cold Acclimation and Allow the Definition of CBF Regulons in Arabidopsis. *New Phytol.* **2016**, *212*, 345–353.

59. Wenjing, W.; Chen, Q.; Singh, P.K.; Huang, Y.; Pei, D. CRISPR/Cas9 Edited HSFA6a and HSFA6b of Arabidopsis Thaliana Offers ABA and Osmotic Stress Insensitivity by Modulation of ROS Homeostasis. *Plant Signal. Behav.* **2020**, *15*, 1816321.

60. Zhang, Y.; Wang, C.; Wang, C.; Yun, L.; Song, L.; Idrees, M.; Liu, H.; Zhang, Q.; Yang, J.; Zheng, X.; et al. OsHsfB4b Confers Enhanced Drought Tolerance in Transgenic Arabidopsis and Rice. *Int. J. Mol. Sci.* **2022**, *23*, doi:10.3390/ijms231810830.

61. Barrett, T.; Wilhite, S.E.; Ledoux, P.; Evangelista, C.; Kim, I.F.; Tomashevsky, M.; Marshall, K.A.; Phillippy, K.H.; Sherman, P.M.; Holko, M.; et al. NCBI GEO: Archive for Functional Genomics Data Sets--Update. *Nucleic Acids Res.* **2013**, *41*, D991-5.

62. Sarkans, U.; Gostev, M.; Athar, A.; Behrangi, E.; Melnichuk, O.; Ali, A.; Minguet, J.; Rada, J.C.; Snow, C.; Tikhonov, A.; et al. The BioStudies Database-One Stop Shop for All Data Supporting a Life Sciences Study. *Nucleic Acids Res.* **2018**, *46*, D1266–D1270.

63. Ono, Y.; Bono, H. Multi-Omic Meta-Analysis of Transcriptomes and the Bibliome Uncovers Novel Hypoxia-Inducible Genes. *Biomedicines* **2021**, *9*, doi:10.3390/biomedicines9050582.

64. Ge, S.X.; Jung, D.; Yao, R. ShinyGO: A Graphical Gene-Set Enrichment Tool for Animals and Plants. *Bioinformatics* **2020**, *36*, 2628–2629.

65. Khan, A.; Mathelier, A. Intervene: A Tool for Intersection and Visualization of Multiple Gene or Genomic Region Sets. *BMC Bioinformatics* **2017**, *18*, 287.