Synaptic cell adhesion molecule *Cdh6* identifies a class of sensory neurons with novel functions in colonic motility

Julieta Gomez-Frittelli[1,2], Gabrielle Devienne[2,3], Lee Travis[4], Melinda A. Kyloh[4], Xin Duan[5], Tim J. Hibberd[4], Nick J. Spencer[4], John R. Huguenard[2,3], Julia A. Kaltschmidt[2,6]

[1]Department of Chemical Engineering, Stanford University; Stanford, CA, USA.

[2]Wu Tsai Neurosciences Institute, Stanford University; Stanford, CA, USA.

[3]Department of Neurology & Neurological Sciences, Stanford University; Stanford, CA, USA.

[4]College of Medicine and Public Health, Flinders University; Adelaide, Australia.

[5]Department of Ophthalmology, School of Medicine, University of California San Francisco; San Francisco, CA, USA.

[6]Department of Neurosurgery, Stanford University School of Medicine; Stanford, CA, USA.

*Corresponding author. Email: jukalts@stanford.edu

Abstract:

Intrinsic sensory neurons are an essential part of the enteric nervous system (ENS) and play a crucial role in gastrointestinal tract motility and digestion. Neuronal subtypes in the ENS have been distinguished by their electrophysiological properties, morphology, and expression of characteristic markers, notably neurotransmitters and neuropeptides. Here we investigated synaptic cell adhesion molecules as novel cell type markers in the ENS. Our work identifies two Type II classic cadherins, *Cdh6* and *Cdh8,* specific to sensory neurons in the mouse colon. We show that *Cdh6+* neurons demonstrate all other distinguishing classifications of enteric sensory neurons including marker expression of *Calcb* and *Nmu*, Dogiel type II morphology and AH-type electrophysiology and I_H current. Optogenetic activation of *Cdh6+* sensory neurons in distal colon evokes retrograde colonic motor complexes (CMCs), while pharmacologic blockade of rhythmicity-associated current I_H disrupts the spontaneous generation of CMCs. These findings provide the first demonstration of selective activation of a single neurochemical and functional class of enteric neurons, and demonstrate a functional and critical role for sensory neurons in the generation of CMCs.

One-Sentence Summary:

Intrinsic sensory neurons of the enteric nervous system in the mouse distal colon exclusively express synaptic cell adhesion molecules *Cdh6* and *Cdh8,* evoke retrograde colonic motor complexes (CMCs) when stimulated, and possess rhythmicity-associated I_H current, which is necessary for the production of spontaneous CMCs.

Main Text:

Introduction

Sensory signaling within the gastrointestinal (GI) tract plays a critical role in the autonomous regulation of digestion. The GI tract is the only internal organ system containing its own sensory neurons. Intrinsic primary afferent neurons (IPANs) detect relevant stimuli through chemo- and mechano-sensation and direct appropriate GI functions via downstream components of the enteric nervous system (ENS), including ascending and descending interneurons, and excitatory and inhibitory motor neurons (*1*). These neuronal subtypes have begun to be distinguished morphologically, electrophysiologically, and by marker expression, classically especially of neurotransmitters (*2*, *3*), and together this information has provided an opening to characterize individual neuron subtype function within the GI tract.

Synaptic cell adhesion molecules define neuronal subtype connectivity within many regions of the CNS. Type II Cadherins are a family of synaptic cell adhesion molecules with combinatorial expression in multiple neural circuits of the CNS, including retina, limbic, olivonuclear, and auditory projection systems (*4–6*). Type II Cadherins bind homophilically by expression of the same cadherin at both the pre- and post-synapse, which stabilizes developing synapses between correct partners while incorrect synapses are pruned away (*7*, *8*). Recent RNA-Seq studies of human and mouse ENS have identified synaptic cell adhesion molecules, including Type II Cadherins, expressed in enteric neuronal subtypes (*9–11*). However, the specificity of their expression has yet to be harnessed to assess neuronal subtype-specific function in the ENS.

Here we identify Type II Cadherin, *Cdh6*, as a novel marker for IPANs of the colonic ENS. We demonstrate the sensory identity of *Cdh6* neurons by immunohistochemical, morphological and neurophysiological classification. *Cdh6* neurons express IPAN markers *Calcb* and *Nmu*. Sparse labeling of individual IPANs reveals they project mainly circumferentially and branch extensively in myenteric ganglia. Whole cell patch clamp recordings of sensory neurons *in situ* reveal action potential (AP) slow after-hyperpolarizations characteristic of IPANs, and hyperpolarization-activated cationic current (I_H), a rhythmicity indicator in thalamocortical and other systems (*12*). Using a *Cdh6* genetic mouse model, we show that optogenetic activation of distal colon IPANs is sufficient to evoke retrograde CMCs, while pharmacologic block of I_H in IPANs disrupts colonic rhythmicity and reversibly abolishes spontaneous CMCs.

Results

Expression of the type II classic cadherin *Cdh6* in colonic IPANs

To identify Cadherins expressed in enteric neuronal subtypes in mouse, we screened recently published RNA-Seq data (*10*, *11*) for Classic Type II cadherin expression. *Cdh6* and *Cdh8* appeared to be restricted to IPAN subsets in both small intestine and colon (*10*, *11*). *Cdh9* was previously identified in a separate population of IPANs in the small intestine (*9–11*). We validated *Cdh6* and *Cdh8* expression by RNAscope *in situ* hybridization in the myenteric plexus, which contains the enteric motility circuitry. *Cdh6* mRNA was expressed in $14.7 \pm 0.8\%$ of myenteric neurons in small intestine (jejunum) and in $6.8 \pm 0.3\%$ of myenteric neurons in distal colon (mean $\pm$ SEM) [Fig.1A-C]. *Cdh8* was almost exclusively co-expressed in *Cdh6+* neurons, although at a much lower level of detection [Fig.1D-H]. We therefore focused our further analysis on *Cdh6+* neurons.

To confirm IPAN identity of *Cdh6+* neurons, we first established the differential
expression of two putative and broadly used markers of IPANs, *Calcb* and *Nmu* (*3, 9, 11*). We
found that all *Nmu+* neurons co-express *Calcb* in both jejunum and distal colon [Fig.1I-K]. In
contrast, only about half of *Calcb+* neurons in the jejunum and two-thirds in the distal colon co-
express *Nmu* [Fig.1L].

We next assessed co-expression of *Cdh6* with *Calcb* and *Nmu.* In the jejunum, we found
that nearly all *Nmu+* neurons and *Calcb+* neurons express *Cdh6* [Fig.1P, U], though only about
three-quarters of all *Cdh6+* neurons express *Calcb* and only about half express *Nmu* [Fig.1O, T].
In contrast, in the distal colon, while *Cdh6* is only expressed in about two-thirds of all *Calcb+*
neurons [Fig.1U], nearly all *Cdh6+* neurons express *Nmu* and *Calcb* [Fig.1O, T]. Taken together,
our data show that in the myenteric plexus, *Cdh6* is expressed exclusively in *Calcb+/Nmu+*
IPANs in the mouse distal colon [Fig.1W, X].

Mouse colonic IPANs display AH-type electrophysiology and I_H current

We next assessed the electrophysiological properties of *Cdh6+* IPANs. We focused our
analysis on the colon, and for ease of neuron tracing, took advantage of a genetic strategy to
sparsely label *Cdh6+* neurons. Previous studies of Hb9:GFP transgenic mice have shown that
due to the inserted transgene's proximity to *Cdh6*, *Cdh6+* neurons can express eGFP (*13*).
Hb9:GFP+ neurons were rare and projected extensively throughout the myenteric plexus
[Fig.2A-C]. *In situ* hybridization confirmed *eGFP* expression was limited to a small fraction (3.5
± 0.8%) of *Cdh6+* colonic neurons [Fig.2D, F-H], and all *eGFP+* neurons expressed *Cdh6*
[Fig.2E].

We developed a protocol to perform whole-cell patch-clamp recordings (*14*) in
Hb9:GFP+ colonic neurons. Membrane capacitance reflecting overall size of these cells was 32 ±
8.7 pF [Fig.3D]; their resting membrane potential (RMP) was -49.4 ± 2.9 mV [Fig.3C]. The
input resistance (R_{in}) was 393 ± 54.7 MΩ [Fig.3E] as computed from the slope of the voltage-
current (V-I) relationship. All patched neurons had large-amplitude action potentials (AP; 72 ±
2.5 mV, [Fig.3F, E]) with threshold of -26.4 ± 0.9 mV [Fig.3L] and a half width of 1.2 ± 0.1 ms
[Fig.3H] elicited at rheobase (20 ± 4.5 pA, [Fig.3G]), each followed by an afterhyperpolarization
(AHP= -67.7 ± 0.9 mV, [Fig.3K]). In addition, the first derivative of membrane voltage during
the action potential (dV/dt, [Fig.3J]) exhibited an inflection during the repolarization phase,
suggesting the presence of fast and slow AP repolarization mechanisms [Fig.3F]. Patched
Hb9:GFP+ neurons generally could not sustain repetitive APs in response to long depolarizing
current pulses. All patched neurons responded to hyperpolarizing current pulses with a time-
dependent membrane potential sag (-6.3 ± 1.2 mV, [Fig.3B, M]), and a rebound depolarization
following the hyperpolarization [Fig.3B, N].

Consistent with these findings of sag and rebound, voltage clamp step hyperpolarizations
revealed I_H (hyperpolarization-activated current, [Fig 3.N]) and, upon repolarization, I_T (transient
inward presumed Ca^{2+} current) (*15*) in terms of its kinetics [Fig 3.O] and steady-state
inactivation [Fig.3O, P]. Taken together, these results show that Hb9:GFP+ / *Cdh6+* colonic
neurons have AP after-hyperpolarizing (AH)-type electrophysiology typical of IPANs, including
rhythm generating currents I_H and I_T (*16–18*).

Colonic IPANs have Dogiel type II morphology and abundant projections throughout the
myenteric plexus

To visualize the morphology and projections of individual patched Hb9:GFP+ neurons,
we included biocytin in the internal solution for post-fixation single-cell tracing [Fig.4]. Patched
Hb9:GFP+ neurons displayed Dogiel type II morphology (*19*), with large smooth cell somas and
multiple branching neurites. Projections were mainly circumferential and extensively branched
within myenteric ganglia. Thus, Hb9:GFP+ neurons display morphological features
characteristic of IPANs (*2*, *19*).

To further visualize the full extent of IPAN circuitry in the myenteric plexus, we
intercrossed Cdh6[CreER] (*20*) and LSL-tdTomato (Ai14) (*21*) mice and induced Cre expression at
5-8 weeks of age [Fig.5A, B]. We confirmed tdTomato+ labeling in myenteric *Cdh6+* neurons
[Fig.5E, F], representing about 5% of the total neuronal population [Fig.5G]. tdTomato+ neurons
had large cell somas (major axis, 27.8 ± 0.7 μm; minor axis, 15.9 ± 0.4 μm) [Fig.5D]. All
ganglia of the myenteric plexus were densely innervated by tdTomato+ fibers [Fig.5A, B], which
also projected into the circular muscle. We noted additional tdTomato+ labeling of some putative
longitudinal and circular muscle cells [Fig.5A, B]. Taken together, our data reveal an IPAN array
that spans the entire motility circuitry of the colonic myenteric plexus.

Optogenetic activation of distal colon IPANs evokes CMCs

In thalamocortical relay neurons, I_H contributes to intrinsic slow rhythmic burst firing at
1-2Hz (*22*). During colonic motor complexes (CMCs), large regions of the ENS oscillate in
synchrony at 1-2Hz to generate traveling contractions along the colon (*23*). Recent calcium
imaging studies have shown that IPANs participate, along with all other subtypes of enteric
neurons, in this synchronized oscillatory firing (*24*). Furthermore, our electrophysiological
studies confirm the presence of I_H in mouse colonic IPANs [Fig.3N]. However, the role of
IPANs in spontaneous CMCs is not well understood.

To interrogate the functional role of *Cdh6+* IPANs in CMCs, we performed *ex vivo*
colonic contraction force recordings in conjunction with optogenetic activation. To express
ChR2-eYFP in *Cdh6+* cells, we intercrossed Cdh6[CreER] and LSL-ChR2-eYFP (Ai32) (*25*) mice
and induced Cre expression at 5-8 weeks of age [Fig.6A-C]. In Cdh6[CreER]+;ChR2-eYFP+ colon
preparations we observed spontaneous CMCs at regular intervals of about 3-5 minutes. Blue
light stimulation of *Cdh6+* IPANs in distal colon 60-90 seconds after a spontaneous CMC
("control" CMC) resulted in an evoked, premature CMC that began during stimulation (N = 17
stimulations; n = 5 mice)[Fig.6D, E]. Evoked CMCs traveled retrogradely from the distal to the
proximal colon. They were similar to spontaneous CMCs in peak amplitude, area under the
curve, and duration, though the contractile force (peak amplitude and AUC) was slightly weaker
in the proximal colon [Fig.6F-H]. Blue light stimulation in proximal or mid colon failed to
generate CMCs (n = 5/5, data not shown). In comparison, stimulation in control Cdh6[CreER]-
negative;ChR2-eYFP+ colons failed to evoke any CMCs (N = 28 stimulations; n = 7
mice)[Fig.S1].

CMCs have previously been shown to depend on nicotinic cholinergic transmission (*24*).
We performed optogenetic stimulation in the presence of hexamethonium, a blocker of nicotinic
cholinergic transmission. Spontaneous CMCs were abolished in hexamethonium, and CMCs
could not be evoked by optogenetic stimulation [Fig.6I]. We conclude that activation of *Cdh6+*
distal colon IPANs evokes retrograde-traveling but otherwise characteristic and hexamethonium-
sensitive CMCs.

Blockade of I_H current in colonic IPANs disrupts CMC production

To determine whether I_H in IPANs contributes to oscillatory firing driving CMCs, we
measured colonic contraction force on a tethered pellet in the presence of I_H blockers ZD7288 or
CsCl (*26*, *27*). Spontaneous CMCs were recorded in all preparations of both proximal and distal
colon prior to drug application [Fig.7]. Addition of 10 μM ZD7288 to the recording chamber
abolished spontaneous CMCs and washout of ZD7288 recovered spontaneous CMC activity
[Fig.6J, Fig.S2]. Addition of 2mM CsCl also impaired or altered spontaneous rhythmic
production of typical CMCs [Fig.6K, Fig.S3]; rhythmic contractions increased in frequency
(proximal colon, n = 5/6; distal colon, 6/6), decreased in amplitude (proximal colon, n = 5/6;
distal colon, n = 4/6, p = 0.0606), or in some cases even included significant retrograde force
components (proximal colon, n = 2/6; distal colon, n = 2/6). We conclude that pharmacologic
blockade of I_H in IPANs impairs the production of CMCs in the mouse colon.

Discussion

Our study shows that in the myenteric plexus, *Cdh6* is expressed exclusively in
Calcb+/Nmu+ IPANs located in the mouse colon, while in the small intestine, *Cdh6* is also
expressed in some *Calcb+/Nmu-* and *Calcb-/Nmu-* neurons. We confirm the IPAN identity of
Cdh6+ colonic neurons by electrophysiological recordings revealing AH-type signature, and
single neuron tracings showing Dogiel type II morphology. Finally, we demonstrate that
activation of IPANs in the distal colon evokes retrograde CMCs, while pharmacologic blockade
of I_H, a rhythmicity associated current we show also present in mouse colonic IPANs, disrupts
spontaneous CMC generation.

Together with *Cdh8*, which we show to be co-expressed with *Cdh6,* our study validates
two new adhesion molecules specific to IPANs in the colon. Notably, these markers show a
different expression pattern than another cadherin, *Cdh9*,which is exclusively expressed in the
small intestine in mouse, not in the colon (*9*). *Cdh9* is expressed in a subset of IPANs not
expressing either *Calcb* or *Nmu* (*3*, *9*, *11*). It is possible that in the small intestine *Cdh6* and *Cdh9*
mark some of the same neurons. However, RNA-Seq data from two separate studies suggest this
is unlikely (*10*, *11*). We conclude that *Cdh6/Cdh8* IPANs are a separate population from *Cdh9*
IPANs.

Though IPANs are positioned to initiate motility by activating other subtypes in enteric
circuitry, their role in spontaneous CMCs has been debated, as CMCs can occur in the absence of
luminal contents, without any apparent stimulus for IPANs to sense, or upon stimulation of
nitrergic populations (*2*, *28*). Only recently has evidence emerged to suggest that IPANs
participate in oscillatory rhythmic firing of the ENS during CMCs, and that activation of
neuronal subtypes expressing calretinin, including IPANs, together can evoke CMCs (*24*, *29*).
Our results demonstrate that excitement of IPANs alone in the distal colon is capable of
producing retrograde CMCs. The exact mechanism causing CMC generation at an established
and controlled frequency of once every few minutes, without evident stimulus, remains to be
determined.

A major observation was that optogenetic stimulation of IPANs readily evoked CMCs
from the distal colon, but never from the proximal colon. Failure of proximal colonic stimulation
to evoke CMCs may reflect inhibition by IPAN recruitment of descending pathways; or
conversely, high efficacy of stimulation from distal colon may reflect bias toward activation of
ascending excitatory cholinergic pathways. In addition, our stimulus paradigm in the proximal
colon may not have activated enough IPANs based on light density and illumination. In previous
studies, optogenetic stimulation of calretinin-expressing neurons or choline acetyl transferase

(ChAT)-expressing neurons (both of which include IPANs, interneurons and motor neurons) elicited anterograde CMCs regardless of stimulus location, including proximal colon (*29, 30*). Proximal colon stimulation of nitric oxide synthase-expressing neurons also elicited CMCs (*28*). How broad and non-specific activation of such disparate neuron classes readily evokes CMCs remains unclear.

Spontaneous and evoked CMCs were abolished in hexamethonium, confirming that CMC synchronous firing is dependent on nicotinic cholinergic transmission (*24*). This reinforces that nicotinic cholinergic transmission is required for greater activation and synchrony of the entire ENS motility network to generate CMCs.

Through our electrophysiological investigation of IPANs in the distal colon using voltage clamp, our work reveals the presence of two voltage gated ion conductances and their underlying currents, I_T and I_H. Slow AHP, I_H and I_T have been previously identified as distinguishing characteristics of IPANs in rat and guinea pig (*16–18*). Although prior studies noted I_H and proposed I_T in guinea pig Dogiel type II neurons (*17*), they have not previously been reported in studies of intact mouse distal colon myenteric plexus due to the conventional reliance on sharp electrode recordings in which voltage clamp is not possible (*31*). The presence of these two currents in IPANs is similar to thalamocortical relay neurons and other cell types, in which intrinsic rhythmicity is supported (*22, 32*), and may similarly promote autonomous rhythmic activity in colonic IPANs.

I_H is conducted through hyperpolarization-activated cyclic nucleotide gated (HCN) channels (*33*). HCN channel family members HCN1 and HCN2 have been shown to be present in mouse distal colonic Dogiel type II neurons (*16*), and RNA-Seq ENS screens similarly indicate their expression in IPANs (*10, 11*). Knockout of HCN2 in mouse leads to a severe growth restriction phenotype due to malnutrition and GI dysmotility (*34*). Here we demonstrate that pharmacologic blockade of I_H, which we show to be present in IPANs, with two distinct HCN channel blockers ZD7288 and CsCl abolishes CMCs, an otherwise persistent and ongoing pattern of motor activity in the mouse colon. We speculate that I_H in colonic IPANs, as in thalamocortical neurons, plays a role in promoting either rhythmic oscillatory single neuron activity, or network burst firing, or both (*22, 32*). Blocking I_H may impair individual IPANs' ability to fire rhythmically, or the ability of IPANs to synchronize into a network burst firing mode. Failure of IPANs to fully activate and synchronize could prevent generation of both (1) synchronized rhythmic myenteric network activation of motility circuits, and (2) the resulting synchronized contractions that sum to much larger contractile forces during CMCs.

Finally, we note that Type II cadherins are homophilic synaptic cell adhesion molecules (*7, 8*). Restricted expression of two Type II cadherins, *Cdh6* and *Cdh8*, to mouse colonic IPANs raises the possibility of these cadherins supporting IPAN-IPAN synaptic connections. While broadly speaking, IPANs are not known to receive synaptic input and in fact have been characterized electrophysiologically by their lack thereof (*35*), some work has in fact suggested that AH-AH neuron interconnected pairs may exist (*36*). Immunohistochemical and electron microscopy investigation of synapses on enteric neurons further showed that calbindin-positive neurons, presumed IPANs, do receive synapses, though fewer than non-calbindin neurons, and some of those synapses were also calbindin-reactive (*37*). These observations informed a proposed "IPAN driver circuit" theory, in which IPANs form an interconnected network of positive feedback to synchronize and amplify sensory signaling and thus activate large swaths of enteric circuitry (*38*). In contrast, more recently, activation of large regions of the ENS has been suggested to be driven by interneuronal networks (*39*). Further investigation may determine

258 whether synaptic adhesion molecules, such as *Cdh6* and *Cdh8*, may in fact support IPAN-IPAN
259 synapses underlying an "IPAN driver circuit."
260
261

References and Notes

1. C. Fung, P. Vanden Berghe, Functional circuits and signal processing in the enteric nervous system. *Cell. Mol. Life Sci.* **77**, 4505–4522 (2020).

2. K. Nurgali, M. J. Stebbing, J. B. Furness, Correlation of electrophysiological and morphological characteristics of enteric neurons in the mouse colon. *J. Comp. Neurol.* **468**, 112–124 (2004).

3. Z.-D. Qu, M. Thacker, P. Castelucci, M. Bagyánszki, M. L. Epstein, J. B. Furness, Immunohistochemical analysis of neuron types in the mouse small intestine. *Cell Tissue Res.* **334**, 147–161 (2008).

4. S. C. Suzuki, T. Inoue, Y. Kimura, T. Tanaka, M. Takeichi, Neuronal Circuits Are Subdivided by Differential Expression of Type-II Classic Cadherins in Postnatal Mouse Brains. *Mol. Cell. Neurosci.* **9**, 433–447 (1997).

5. X. Duan, A. Krishnaswamy, M. A. Laboulaye, J. Liu, Y.-R. Peng, M. Yamagata, K. Toma, J. R. Sanes, Cadherin Combinations Recruit Dendrites of Distinct Retinal Neurons to a Shared Interneuronal Scaffold. *Neuron* **99**, 1145-1154.e6 (2018).

6. M. Honjo, H. Tanihara, S. Suzuki, T. Tanaka, Y. Honda, M. Takeichi, Differential Expression of Cadherin Adhesion Receptors in Neural Retina of the Postnatal Mouse. *Invest. Ophthalmol. Vis. Sci.* **41**, 546–551 (2000).

7. M. Yamagata, X. Duan, J. R. Sanes, Cadherins Interact With Synaptic Organizers to Promote Synaptic Differentiation. *Front. Mol. Neurosci.* **11** (2018).

8. R. Basu, M. R. Taylor, M. E. Williams, The classic cadherins in synaptic specificity. *Cell Adhes. Migr.* **9**, 193–201 (2015).

9. A. A. May-Zhang, E. Tycksen, A. N. Southard-Smith, K. K. Deal, J. T. Benthal, D. P. Buehler, M. Adam, A. J. Simmons, J. R. Monaghan, B. K. Matlock, D. K. Flaherty, S. S. Potter, K. S. Lau, E. M. Southard-Smith, Combinatorial Transcriptional Profiling of Mouse and Human Enteric Neurons Identifies Shared and Disparate Subtypes In Situ. *Gastroenterology* **160**, 755-770.e26 (2021).

10. E. Drokhlyansky, C. S. Smillie, N. V. Wittenberghe, M. Ericsson, G. K. Griffin, G. Eraslan, D. Dionne, M. S. Cuoco, M. N. Goder-Reiser, T. Sharova, O. Kuksenko, A. J. Aguirre, G. M. Boland, D. Graham, O. Rozenblatt-Rosen, R. J. Xavier, A. Regev, The Human and Mouse Enteric Nervous System at Single-Cell Resolution. *Cell* **182**, 1606-1622.e23 (2020).

11. K. Morarach, A. Mikhailova, V. Knoflach, F. Memic, R. Kumar, W. Li, P. Ernfors, U. Marklund, Diversification of molecularly defined myenteric neuron classes revealed by single-cell RNA sequencing. *Nat. Neurosci.* **24**, 34–46 (2021).

12. C. Wahl-Schott, M. Biel, HCN channels: Structure, cellular regulation and physiological function. *Cell. Mol. Life Sci.* **66**, 470 (2008).

13. M. A. Laboulaye, X. Duan, M. Qiao, I. E. Whitney, J. R. Sanes, Mapping Transgene Insertion Sites Reveals Complex Interactions Between Mouse Transgenes and Neighboring Endogenous Genes. *Front. Mol. Neurosci.* **11** (2018).

14. N. Osorio, P. Delmas, Patch clamp recording from enteric neurons in situ. *Nat. Protoc.* **6**, 15–27 (2011).

15. J. R. Huguenard, Low-threshold calcium currents in central nervous system neurons. *Annu. Rev. Physiol.* **58**, 329–348 (1996).

16. J. Xiao, T. V. Nguyen, K. Ngui, P. J. L. M. Strijbos, I.-S. Selmer, C. B. Neylon, J. B. Furness, Molecular and functional analysis of hyperpolarisation-activated nucleotide-gated (HCN) channels in the enteric nervous system. *Neuroscience* **129**, 603–614 (2004).

17. K. Nurgali, T. V. Nguyen, H. Matsuyama, M. Thacker, H. L. Robbins, J. B. Furness, Phenotypic changes of morphologically identified guinea-pig myenteric neurons following intestinal inflammation. *J. Physiol.* **583**, 593–609 (2007).

18. Y. Mao, B. Wang, W. Kunze, Characterization of Myenteric Sensory Neurons in the Mouse Small Intestine. *J. Neurophysiol.* **96**, 998–1010 (2006).

19. J. B. Furness, H. L. Robbins, J. Xiao, M. J. Stebbing, K. Nurgali, Projections and chemistry of Dogiel type II neurons in the mouse colon. *Cell Tissue Res.* **317**, 1–12 (2004).

20. J. N. Kay, I. D. la Huerta, I.-J. Kim, Y. Zhang, M. Yamagata, M. W. Chu, M. Meister, J. R. Sanes, Retinal Ganglion Cells with Distinct Directional Preferences Differ in Molecular Identity, Structure, and Central Projections. *J. Neurosci.* **31**, 7753–7762 (2011).

21. L. Madisen, T. A. Zwingman, S. M. Sunkin, S. W. Oh, H. A. Zariwala, H. Gu, L. L. Ng, R. D. Palmiter, M. J. Hawrylycz, A. R. Jones, E. S. Lein, H. Zeng, A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nat. Neurosci.* **13**, 133–140 (2010).

22. D. A. McCormick, H. C. Pape, Properties of a hyperpolarization-activated cation current and its role in rhythmic oscillation in thalamic relay neurones. *J. Physiol.* **431**, 291–318 (1990).

23. N. J. Spencer, M. Costa, T. J. Hibberd, J. D. Wood, Advances in colonic motor complexes in mice. *Am. J. Physiol.-Gastrointest. Liver Physiol.* **320**, G12–G29 (2021).

24. T. J. Hibberd, L. Travis, L. Wiklendt, M. Costa, S. J. H. Brookes, H. Hu, D. J. Keating, N. J. Spencer, Synaptic activation of putative sensory neurons by hexamethonium-sensitive nerve pathways in mouse colon. *Am. J. Physiol. Gastrointest. Liver Physiol.* **314**, G53–G64 (2018).

25. L. Madisen, T. Mao, H. Koch, J. Zhuo, A. Berenyi, S. Fujisawa, Y.-W. A. Hsu, A. J. Garcia, X. Gu, S. Zanella, J. Kidney, H. Gu, Y. Mao, B. M. Hooks, E. S. Boyden, G. Buzsáki, J. M. Ramirez, A. R. Jones, K. Svoboda, X. Han, E. E. Turner, H. Zeng, A toolbox

of Cre-dependent optogenetic transgenic mice for light-induced activation and silencing.
Nat. Neurosci. **15**, 793–802 (2012).

26. N. C. Harris, A. Constanti, Mechanism of block by ZD 7288 of the hyperpolarization-
activated inward rectifying current in guinea pig substantia nigra neurons in vitro. *J.*
Neurophysiol. **74**, 2366–2378 (1995).

27. J. J. Galligan, H. Tatsumi, K. Z. Shen, A. Surprenant, R. A. North, Cation current activated
by hyperpolarization (IH) in guinea pig enteric neurons. *Am. J. Physiol.-Gastrointest. Liver*
Physiol. **259**, G966–G972 (1990).

28. S. D. Koh, B. T. Drumm, H. Lu, H. J. Kim, S.-B. Ryoo, H.-U. Kim, J. Y. Lee, P.-L. Rhee,
Q. Wang, T. W. Gould, D. Heredia, B. A. Perrino, S. J. Hwang, S. M. Ward, K. M.
Sanders, Propulsive colonic contractions are mediated by inhibition-driven poststimulus
responses that originate in interstitial cells of Cajal. *Proc. Natl. Acad. Sci.* **119**,
e2123020119 (2022).

29. T. J. Hibberd, J. Feng, J. Luo, P. Yang, V. K. Samineni, R. W. Gereau, N. Kelley, H. Hu, N.
J. Spencer, Optogenetic Induction of Colonic Motility in Mice. *Gastroenterology* **155**, 514-
528.e6 (2018).

30. A. Efimov, T. Hibberd, Y. Wang, M. Wu, K. Zhang, K. Ting, S. Madhvapathy, M. Lee, J.
Kim, J. Kang, M. Riahi, H. Zhang, L. Travis, E. J. Govier, L. Yang, N. Kelly, Y. Huang, A.
Vázquez-Guardado, N. J. Spencer, J. A. Rogers, Remote optogenetic control of the enteric
nervous system and brain-gut axis in freely-behaving mice enabled by a wireless, battery-
free optoelectronic device. *Biosens. Bioelectron.*, 116298 (2024).

31. Z. Li, W. Boesmans, Y. Kazwiny, M. M. Hao, P. Vanden Berghe, Simultaneous whole-cell
patch-clamp and calcium imaging on myenteric neurons. *Am. J. Physiol.-Gastrointest.*
Liver Physiol. **323**, G341–G347 (2022).

32. H. C. Pape, D. A. McCormick, Noradrenaline and serotonin selectively modulate thalamic
burst firing by enhancing a hyperpolarization-activated cation current. *Nature* **340**, 715–718
(1989).

33. E. E. Benarroch, HCN channels. *Neurology* **80**, 304–310 (2013).

34. D. W. Fisher, P. Luu, N. Agarwal, J. E. Kurz, D. M. Chetkovich, Loss of HCN2 leads to
delayed gastrointestinal motility and reduced energy intake in mice. *PloS One* **13**, e0193012
(2018).

35. G. D. S. Hirst, M. E. Holman, I. Spence, Two types of neurones in the myenteric plexus of
duodenum in the guinea-pig. *J. Physiol.* **236**, 303–326 (1974).

36. W. A. A. Kunze, J. B. Furness, J. C. Bornstein, Simultaneous intracellular recordings from
enteric neurons reveal that myenteric ah neurons transmit via slow excitatory postsynaptic
potentials. *Neuroscience* **55**, 685–694 (1993).

37. S. Pompolo, J. B. Furness, Ultrastructure and synaptic relationships of calbindin-reactive,
Dogiel type II neurons, in myenteric ganglia of guinea-pig small intestine. *J. Neurocytol.*
17, 771–782 (1988).

38. J. D. Wood, "Chapter 21 - Cellular Neurophysiology of Enteric Neurons" in *Physiology of*
the Gastrointestinal Tract (Fifth Edition), L. R. Johnson, F. K. Ghishan, J. D. Kaunitz, J. L.
Merchant, H. M. Said, J. D. Wood, Eds. (Academic Press, Boston, 2012;
https://www.sciencedirect.com/science/article/pii/B978012382026600021X), pp. 629–669.

39. B. B. Barth, N. J. Spencer, W. M. Grill, "Activation of ENS Circuits in Mouse Colon:
Coordination in the Mouse Colonic Motor Complex as a Robust, Distributed Control
System" in *The Enteric Nervous System II*, N. J. Spencer, M. Costa, S. M. Brierley, Eds.
(Springer International Publishing, Cham, 2022), pp. 113–123.

40. J. Gomez-Frittelli, R. Hamnett, J. A. Kaltschmidt, Comparison of wholemount dissection
methods for neuronal subtype marker expression in the mouse myenteric plexus. bioRxiv
[Preprint] (2023). https://doi.org/10.1101/2023.01.17.524014.

41. R. Hamnett, L. B. Dershowitz, V. Sampathkumar, Z. Wang, J. Gomez-Frittelli, V. De
Andrade, N. Kasthuri, S. Druckmann, J. A. Kaltschmidt, Regional cytoarchitecture of the
adult and developing mouse enteric nervous system. *Curr. Biol.* **32**, 4483-4492.e5 (2022).

42. R. A. Guyer, R. Stavely, K. Robertson, S. Bhave, J. L. Mueller, N. M. Picard, R. Hotta, J.
A. Kaltschmidt, A. M. Goldstein, Single-cell multiome sequencing clarifies enteric glial
diversity and identifies an intraganglionic population poised for neurogenesis. *Cell Rep.* **42**,
112194 (2023).

43. J. R. Huguenard, D. A. Prince, A novel T-type current underlies prolonged Ca(2+)-
dependent burst firing in GABAergic neurons of rat thalamic reticular nucleus. *J. Neurosci.*
12, 3804–3817 (1992).

44. M. Ishizawa, Propulsive Activity of the Isolated Segmental Colon of Guinea-Pig. *Jpn. J.*
Smooth Muscle Res. **20**, 401–406 (1984).

45. T. J. Hibberd, M. Costa, D. J. Smolilo, L. J. Keightley, S. J. Brookes, P. G. Dinning, N. J.
Spencer, Mechanisms underlying initiation of propulsion in guinea pig distal colon. *Am. J.*
Physiol.-Gastrointest. Liver Physiol. **323**, G71–G87 (2022).

Acknowledgments: We thank the members of the Kaltschmidt laboratory for experimental
advice and discussions, Vanda Lennon (Mayo Clinic) for the HuC/D primary antibody and
Beatriz G. Robinson for 3D printing of organ chambers for electrophysiology.

Funding:

National Institute of General Medical Sciences of the National Institutes of Health Award
T32GM120007 (JGF)

National Institutes of Health Grant R01 EY030138 (XD)

National Health and Medical Research Council (NHMRC) project grant 1156416 (NJS)

Australian Research Council (ARC) Discovery Project grant DP220100070 (NJS)

NINDS 5RO1NS34774 (JRH)

Wu Tsai Neurosciences Institute, Stanford University (JAK)

Department of Neurosurgery, Stanford University (JAK)

National Institutes of Health Grant R21 HD110950 (JAK)

The Firmenich Foundation (JAK)

The Carol and Eugene Ludwig Family Foundation (JAK)

Stanford ADRC Developmental Project Grant (National Institutes of Health Grant P30AG066515) (JAK)

Author contributions:

Conceptualization: JGF, JAK

Resources: NJS, JRH, JAK

Methodology: JGF, GD, XD, TJH

Investigation: JGF, GD, LT, MAK, TJH

Visualization: JGF, GD, TJH

Funding acquisition: NJS, JRH, JAK

Supervision: NJS, JRH, JAK

Writing – original draft: JGF, GD

Writing – review & editing: JGF, NJS, JRH, JAK

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Fig. 1. *Cdh6* **expression overlaps with IPAN markers** *Calcb* **and** *Nmu*. (**A, B**) Representative images of jejunum (A) and distal colon (B) myenteric plexus labeled with HuC/D (IHC) (blue) and *Cdh6* (RNA) (red). (**C**) Proportion of total HuC/D neurons positive for *Cdh6* (jejunum, n = 9; distal colon, n = 9). (**D, E**) As in (A, B) for HuC/D (IHC) (blue), *Cdh6* (RNA) (red), and *Cdh8* (RNA) (green). (**F**) Proportion of *Cdh6+* neurons positive for *Cdh8* (jejunum, n = 8; distal colon, n = 8). (**G**) Proportion of *Cdh8+* neurons positive for *Cdh6* (jejunum, n = 8; distal colon, n = 8). (**H**) Proportion of total HuC/D neurons positive for *Cdh8* (jejunum, n = 8; distal colon, n = 8). (**I, J**) As in (A, B) for HuC/D (IHC) (blue), *Calcb* (RNA) (red), and *Nmu* (RNA) (green). (**K**) Proportion of *Nmu+* neurons positive for *Calcb* (jejunum, n =5 ; distal colon, n = 5). (**L**) Proportion of *Calcb+* neurons positive for *Nmu* (jejunum, n = 5; distal colon, n = 5). (**M, N**) As in (A, B) for HuC/D (IHC) (blue), *Cdh6* (RNA) (red), and *Nmu* (RNA) (green). (**O**) Proportion of *Cdh6+* neurons positive for *Nmu* (jejunum, n = 7; distal colon, n = 8). (**P**) Proportion of *Nmu+* neurons positive for *Cdh6* (jejunum, n = 7; distal colon, n = 8). (**Q**) Proportion of total HuC/D neurons positive for *Nmu* (jejunum, n = 7; distal colon, n = 8). (**R, S**) As in (A, B) for HuC/D (IHC) (blue), *Cdh6* (RNA) (red), and *Calcb* (RNA) (green). (**T**) Proportion of *Cdh6+* neurons positive for *Calcb* (jejunum, n = 9; distal colon, n = 7). (**U**) Proportion of *Calcb+* neurons positive for *Cdh6* (jejunum, n = 9; distal colon, n = 7). (**V**) Proportion of total HuC/D neurons positive for *Calcb* (jejunum, n = 9; distal colon, n = 9). (**W, X**) Schematic of marker overlap in jejunum (W) and distal colon (X). Scale bar represents 100 μm for (A, B, I, J, M, N, R, S), 50 μm for (D, E). All charts (mean ± SEM). * p<0.05; ** p<0.01; *** p<0.001; **** p<0.0001.

Fig. 2. Hb9:GFP+ is expressed in a small proportion of *Cdh6*+ colon myenteric neurons.
(**A, B**) Representative images of Hb9:GFP+ colon myenteric plexus labeled with HuC/D (IHC)
(magenta) and GFP (green). (**C**) Proportion of total HuC/D neurons positive for GFP (n = 3). (**D**)
Proportion of *Cdh6*+ neurons positive for *eGFP* (n = 3). (**E**) Proportion of *eGFP*+ neurons
positive for *Cdh6* (n=3). (**F-H**) Representative images of Hb9:GFP+ colon myenteric plexus
labeled with HuC/D (IHC) (blue), *Cdh6* (RNA) (red), and e*GFP* (RNA) (green). Scale bar
represents 200 μm for (A, B), 50 μm for (E-G). All charts (mean ± SEM).

464
465
466

Fig. 3. Hb9:GFP+ neurons have AH electrophysiological characteristics. (**A**) IR
videomicroscopy image of an Hb9:GFP neuron that presented a large soma located in a ganglion
(scale bar, 10 μm). (**B**) Current-clamp recordings of the same neuron in (A) obtained in response
to application of current pulse (bottom traces) of −50 pA, and + 10 pA. Note the presence of a
sag and a post-hyperpolarization rebound depolarization. (**C-E**) Box-and-whisker plots of
cellular properties of recorded neurons. (C) resting membrane potential (RMP), (D) capacitance
(Cm), and (E) membrane resistance (Rm) (N = 5). (**F**) Averaged traces of the first spike
(rheobase action potential) after a depolarization step of 1s. (**J**) Averaged derivative traces of the
first spike (rheobase action potential). An inflection on the repolarizing phase is observed in the
first derivative (arrow). (**G-I, K-N**) Box-and-whisker plots of electrophysiological properties of
recorded neurons; rheobase action potential (AP, G-I) (G) current threshold, (H) half-width, (I)
amplitude, (K) afterhyperpolarization (AHP), and (L) threshold. (M, N) non-AP properties sag
(mV) and rebound (mV). (**O**) H and T currents in recorded neurons. Top: example of currents
obtained from voltage protocol. Bottom: 500 ms hyperpolarizations ranging from -90 to -45 for
500 ms followed by depolarizing to -40 mV. Hyperpolarizations evoked slowly activating
inward current (H-Current, arrow), followed by a transient inward current upon post-
conditioning step to -30mV (T- current, arrow) Largest T and H currents were obtained with the
most hyperpolarized potentials (red trace). (**P**) Normalized peak I_T plotted versus holding
potential to obtain the I/Imax curve (N = 3/3). Scale bar represents 10 μm for (A).

Fig. 4. Hb9:GFP+ neurons have circumferential branching projections. (**A-C**)
Representative images of Hb9:GFP+ colon myenteric plexus labeled with HuC/D (blue),
streptavidin (red), and GFP (green). (**D, E**) Tracings of Hb9:GFP+ neurons filled with biocytin
during whole cell patch clamp recording. (**F**) Image of patched and filled Hb9:GFP+ neuron
traced in (E). (**G, H**) Inset of (F). Scale bar represents 40 µm for (A-C), 200 µm for (D-F), 100
µm for (G, H).

Fig. 5. Cdh6^{CreER}+/ tdTomato+ neurons have Dogiel type II morphology. (**A, B**)
Representative images of Cdh6^{CreER}+;tdTomato+ distal colon myenteric plexus labeled with
HuC/D (IHC) (magenta) and tdTomato (IHC) (green). (**C**) Inset of (A). (**D**) Dimensions of
tdTomato+ neurons (major and minor axes) (N = 73; n = 3). (**E**) Representative image of
Cdh6^{CreER}+;tdTomato+ distal colon myenteric plexus labeled with HuC/D (IHC) (blue),
tdTomato (IHC) (red), and *Cdh6* (RNA) (green). Arrowheads indicate *Cdh6*+ / tdTomato+ cells;
arrowhead, *Cdh6*+ / tdTomato-negative cell. (**F**) Proportion of *Cdh6*+ neurons positive for
tdTomato (n = 6). (**G**) Proportion of total HuC/D neurons positive for tdTomato (proximal colon,
n = 5; mid colon, n = 5; distal colon, n = 10). Scale bar represents 100 μm for all images. All
charts (mean ± SEM).

Fig. 6. Optogenetic stimulation of colonic *Cdh6*+ neurons evokes CMCs, while pharmacologic blockade of I_H abolishes spontaneous CMCs (**A, B**) Representative images of Cdh6^{CreER}+;ChR2-eYFP+ distal colon myenteric plexus labeled with HuC/D (magenta) and GFP (green). (**C**) Proportion of total HuC/D neurons positive for ChR2-eYFP (n = 6). (**D**) Representative force traces. Blue bars indicate timing of light stimulation. LEDs placed distal to distal hook. (**E**) CMC intervals recorded from force traces. Evoked (blue) intervals represent the time from the prior spontaneous CMC before stimulation to the evoked CMC following stimulation. Control (grey) intervals represent the time between the spontaneous CMC prior to stimulation and the previous spontaneous CMC (n = 5). Paired t test, one tailed. (**F**) CMC peak amplitude recorded from force traces. Evoked (blue) indicates the evoked CMC following stimulation. Grey (control) indicates the spontaneous CMC prior to stimulation (n = 5). Paired t test, two tailed. (**G**) CMC AUC (area under the curve). Evoked (blue) and control (grey) as in (F) (n = 5). Paired t test, two tailed. (**H**) CMC duration. Evoked (blue) and control (grey) as in (F) (n = 5). Paired t test, two tailed. (**I**) Representative force traces. Hex indicates addition of 300 µM hexamethonium. Blue bars indicate timing of light stimulation. LEDs placed distal to distal hook (n = 5/5). (**J**) Representative force traces on tethered pellets. First arrowhead indicates addition of 10 µM ZD7288. Second arrowhead indicates washout in Krebs. Third arrowhead indicates addition of 1 µM TTX. ZD7288 abolished CMCs in both proximal and distal colon (n = 6/6, p = 0.0022, Fisher's exact test). Washout in Krebs restored CMCs in both proximal and distal colon (n = 6/6, p = 0.0022, Fisher's exact test). (**K**) As in (J). First arrowhead indicates addition of 2 mM CsCl. Second arrowhead indicates washout in Krebs. Third arrowhead indicates addition of 1 µM TTX. Typical CMC production was impaired or altered by CsCl (proximal colon, n = 5/6,

532   p = 0.0152; distal colon, n = 6/6, p = 0.0022, Fisher's exact test): increased frequency (proximal
533   colon, n = 5/6, p = 0.0152; distal colon, n = 6/6, p = 0.0022, Fisher's exact test), decreased in
534   amplitude (proximal colon, n = 5/6, p = 0.0152; distal colon, n = 4/6, p = 0.0606, Fisher's exact
535   test); retrograde force (proximal colon, n = 2/6; distal colon, n = 2/6). Scale bar represents 100
536   μm for (A, B). * p<0.05; ** p<0.01; *** p<0.001.

538
539

Materials and Methods

Mice

  All procedures conformed to the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals and were approved by the Stanford University Administrative Panel on Laboratory Animal Care. Mice were group housed up to a maximum of five adults per cage. Food and water were provided ad libitum and mice were maintained on a 12:12 LD cycle. Male and female mice were used in all experiments.

  Wild-type C57BL/6J mice (#000664), Hb9:GFP mice (#005029), Ai14 (#007908), and Ai32 (#024109) (*21*) mice were obtained from the Jackson Laboratory and from the Animal Resource Center (ARC) in Western Australia, with JAX heritage. Cdh6^{CreER} mice (#029428) (*20*) were provided by Xin Duan (UCSF). Cdh6^{CreER} mice were crossed to Ai14 mice and Ai32 mice to generate mice heterozygous for each allele, termed Cdh6^{CreER};Ai14 and Cdh6^{CreER};Ai32. Tamoxifen (20 mg/mL in corn oil) was administered via oral gavage to a final dose of 2.5mg/10g mouse for five consecutive days beginning at 5-8 weeks of age. Induced mice were group housed for at least four weeks prior to experiments.

  Adult male and female mice (Cdh6^{CreER+};Ai32 and Cdh6^{CreER-};Ai32) aged 16 to 19 weeks were euthanised by isoflurane inhalation overdose in accordance with Flinders Animal Welfare Committee guidelines (ethics approval #4004). The protocol for animal euthanasia is approved by the National Health and Medical Research Council (NHMRC) Australian code for the care and use of animal for scientific purposes (8thedition, 2013) and recommendations from the NHMRC Guidelines to promote the wellbeing of animals used for scientific purposes (2008).

Dissection

  Mice were culled by CO_2 and cervical dislocation. Small intestine and colon were removed and flushed with ice cold PBS, then placed in a Sylgard-lined Petri dish with ice cold PBS for further dissection.

Wholemount preparations

  Intestinal segments were prepared and fixed as in Gomez-Frittelli *et al* (*40*). Briefly, each intestinal segment was opened along the mesentery border, pinned flat under light tension serosa-side up, and fixed in 4% PFA in PBS at 4°C with gentle rocking for 90 minutes. Segments were washed three times in PBS at 4°C for at least 10 minutes with gentle rocking. Muscularis was separated from the mucosa at one end of the segment with fine forceps for 2-3 mm, then pinned mucosa-side up in the dish. The mucosa was peeled away from the muscularis with fine forceps while the muscularis was gently held down in the dish with a cotton swab. For immunohistochemistry, segments were processed immediately or stored in PBS with 0.1% NaN$_3$ at 4°C until use. For RNAscope, muscularis segments were post-fixed in 4% PFA in PBS at 4°C with gentle rocking overnight, then washed three times in PBS at 4°C with gentle rocking for at least 10 minutes each wash before use.

Immunohistochemistry

Immunohistochemistry was performed as described previously (*41*). Briefly, muscularis wholemount tissue segments about 7mm x 7mm were incubated with primary antibodies in PBT (PBS with 1% BSA and 0.3% Triton-X 100) at 4°C with gentle rocking overnight, then washed three times in PBT for at least ten minutes each at room temperature with gentle shaking. Tissues were incubated in secondary antibodies in PBT for 2 hours with gentle shaking, washed twice in PBT, and twice in PBS, then mounted on Superfrost Plus slides with Fluoromount G medium (Southern Biotech). Primary antibodies: human anti-HuC/D (1:75k) (gift from Vanda Lennon); sheep anti-GFP (1:1k) (Biogenesis); rabbit anti-RFP (1:1k) (Rockland); rabbit anti-PGP9.5 (1:4k) (Abcam). Secondary antibodies: donkey anti-human AlexaFluor (AF)-647 (1:500); donkey anti-sheep AF-488 (1:1k); donkey anti-rabbit AF-488 (1:1k); streptavidin AF-546 (1:500).

RNAscope

In situ hybridization in combination with IHC was performed on muscularis wholemount tissues using the RNAscope Multiplex Fluorescent V2 Assay kit with RNA-Protein Co-detection Ancillary Kit [ACD], according to the manufacturer's instructions with modifications as previously described (*42*). Probes used were *Cdh6* (#519541), *Cdh8* (#485461), *Nmu* (#446831), *Calcb* (#425511), and *eGFP* (#400281).

Confocal imaging

Images were acquired on a Leica SP8 confocal microscope using a 20x (NA 0.75) oil objective at 1024 x 1024 pixel resolution. Tiled images (24-30 tiles) of Z-stacks (2.5µm between planes) were acquired and stitched together using the Navigator mode within LASX (Leica). Imaged regions were located away from the mesenteric border.

Image analysis and quantification

Image analysis was performed using ImageJ/FIJI (NIH, Bethesda, MD), as described previously (*41*). HuC/D images (z-stack individual planes) were blurred and thresholded; then maximally projected and total neurons counted using the Analyze Particles function. $Cdh6^{CreER}$;Ai32 expression and RNAscope *in situ* hybridization and $Cdh6^{CreER}$;Ai14 expression were counted manually. Cell tracing was performed in Imaris using Filament Tracer (Bitplane, Oxford Instruments).

Electrophysiological recordings

Whole cell patch clamp electrophysiological recordings of Hb9:GFP+ neurons were performed according to Osorio & Delmas (*14*), with modifications for recording from the colon. The protocol is described in brief below.

Tissue dissection and preparation

Mice aged 8-10 weeks were culled by CO_2 and cervical dislocation. The colon was removed and flushed with ice cold oxygenated Krebs solution [118 mM NaCl, 4.8 mM KCl, 1 mM NaH2PO4, 25 mM NaHCO3, 1.2 mM MgCl2, 2.5 mM CaCl2 and 11 mM glucose, supplemented with scopolamine (2 M) and nicardipine (6 µM)], then placed in a Sylgard-lined Petri dish with ice cold oxygenated Krebs solution for further dissection. Krebs solution was changed out for fresh oxygenated solution every 5 minutes. Under a dissection microscope, the

colon was pinned and the mucosa peeled away using fine forceps, leaving a few millimeters of
mucosa along the edges of the tissue for pinning stability. The muscularis was then flipped over
and re-pinned, serosa side up, and the longitudinal muscle carefully peeled away. The tissue was
transferred to a custom 3D-printed recording chamber lined with a thin layer of clear Sylgard,
and re-pinned under light tension, with the myenteric plexus facing up. The tissue was kept at
32°C and was continuously perfused with oxygenated Krebs solution. Hb9:GFP+ neurons were
visually identified within a ganglion under epifluorescence illumination with a 455nM LED
(Thorlabs, M455L2) and a 470 (excitation)/525 (emission) nm wavelength filter set. A local
perfusion of protease XIV (0.2% in Krebs) (Sigma, P5417) was applied on top of the targeted
cell to digest any muscle fiber residue. A 1-2MΩ pipet with a trimmed arm hair glued to the tip
was used to brush and clean the surface of the ganglion. Further cleaning with 1mg/mL
collagenase (Worthington, CLS-4) 4mg/mL dispase (Sigma, D4693) in Krebs solution was also
performed to expose the GFP neuron for patching.
648
649 *Patching and recording*
650
Patch pipettes (4–6 MΩ) pulled from borosilicate glass were filled with internal solution
containing in mM: 144 K-gluconate, 3 MgCl2, 0.5 EGTA, 10 HEPES, pH 7.2 (285/295 mOsm)
and 2% biocytin (Millipore Sigma, B4261-100MG). Patch clamp recordings were collected with
a Multiclamp 700A (Molecular Devices) amplifier, a Digidata 1440 digitizer and pClamp10.7
(Molecular Devices). Recordings were sampled and filtered at 10kHz. Passive properties
analysis was performed using pClamp10.7. Analysis of action potentials was performed using a
custom MATLAB (MathWorks) software. All recordings were performed at 32°C. Membrane
potentials were not corrected for liquid junction potential. Immediately after whole cell
configuration, the cell was maintained at -70mV and a short voltage clamp membrane test
protocol consisting of 20 times 600ms, 10mV depolarization steps was performed to assess cell
health and recording conditions. Recordings were performed in Hb9:GFP+ colonic neurons with
an access resistance less than 30MΩ (16.69 ± 2.63 MΩ). Next, the current clamp mode was used
to measure resting membrane potential (RMP), input resistance (Rin) and APs stimulated.
Membrane potential was not adjusted from resting potential, and cells were depolarized by 1
second current pulses in 10pA increments until APs were triggered (rheobase). Finally, if the seal
was still stable, a voltage clamp steady-state inactivation of T-current protocol was performed as
previously described (*43*). In brief, a sequence of depolarization from -90 to -45mV for 500ms
quickly followed by a depolarization to -40mV for 200ms. Tissues were then fixed and
immunostained according to *Wholemount preparations* and <u>Immunohistochemistry</u> sections
above.
671
<u>Mechanical recordings & optical stimulation</u>
Optogenetic stimulation experiments were performed as previously described (*29*). A 2.5
mm stainless steel rod was inserted through the lumen of the colon and mounted in an organ bath
(120 * 40 * 12 mm; L*W*H) located on a heated base. Krebs solution (35.5-36°C) superfused
the bath (~5 mL/min). Smooth muscle force was recorded via 4 evenly spaced hooks in the
colonic muscularis externa, each linked to an isometric force transducer (Grass FT03C) by suture
thread. Initial base resting tension was set between 0.5 - 1.0 g. Preamplified signals (Biomedical
Engineering, Flinders University) were digitized by a PowerLab 16/35 (ADInstruments, Bella
Vista, NSW, Australia) and recorded using LabChart 7 software (ADInstruments) on iMac
computer. Post hoc analysis of the mechanical recordings was done using LabChart 8 software
on PC.

683
For optical stimulation during mechanical recordings *in vitro*, two LEDs (emitting 470
nmλ photons; C470DA2432, Cree Inc., NC, USA) were used, driven by a variable power supply.
The area of light emission from each LED was 240 μm x 320 μm (0.0768 mm^2). To characterise
LED function, light power density across a range of currents was measured 5 mm from the LED
using a standard photodiode power sensor (S120C; Thorlabs, NJ, USA) and a power meter
(Thorlabs, PM100USB). The stimulator panel within LabChart software was used to set
parameters and manually trigger LED pulse trains via the 10V analogue output of the PowerLab
and an ILD1 opto-isolator.
692
<u>Intraluminal pellet CMC recordings</u>
To record proximal and distal colon CMCs separately (*44*, *45*), full length colon was
bisected halfway between the caeco-colonic junction and terminal rectum, creating equal length
proximal and distal colon preparations. Each preparation was suspended vertically on a stainless-
steel holder inside a glass, water jacketed organ bath containing Krebs solution (Fig.6J, K). A 2.7
mm diameter synthetic pellet (polymethyl methacrylate, "Perspex") was placed inside the gut
lumen and linked by stainless steel rod to a force transducer (MLT0420, ADInstruments),
allowing measurement of both anterograde and retrograde propulsive forces on the pellet.
Signals were amplified by bridge amplifier (FE224, ADInstruments), digitized at 1kHz (PLCF1,
ADInstruments) and recorded using LabChart 8 software.
703
704   ZD7288 (73777, Sigma-Aldrich) was dissolved in water as stock solution at 10 mM.
705   Caesium Chloride (C4036, Sigma-Aldrich) was dissolved in water as stock solution at 200 mM.
706   Tetrodotoxin citrate (T-550, Alomone Labs) was dissolved in water as stock solution at 3 mM.
707   Control, ZD7288, CsCl and washout periods were at least 30 mins; TTX was applied for at least
708   10 minutes.
709
710   <u>Statistical analysis</u>
711   Statistical tests and graphical representation of data were performed using Prism 9
712   software (GraphPad). Statistical comparisons were performed using paired t tests (one-tailed,
713   CMC intervals; two-tailed, peak amplitude, AUC, duration) and Welch's t test (marker
714   colocalizations). Asterisks indicate significant differences.
715
716
717

A   Cdh6^{CreER}-;ChR2-eYFP+

719 **Fig. S1.**

720 **Optogenetic stimulation control.** (**A**) Representative force traces of control Cdh6^{CreER}-
721 ;Chr2eYFP+ colon (n = 5). Blue bars indicate timing of light stimulation. LEDs placed distal to
722 distal hook.

723
724

ZD7288

A Proximal colon

Force

B Distal colon

Force

Fig. S2. Pharmacologic blockade of I_H with ZD7288 abolishes spontaneous CMCs. (A, B) Representative force traces from tethered pellet in proximal half (A) or distal half (B) of colon. Addition of 10 µM ZD7288 (first arrowhead), followed by washout in Krebs (second arrowhead), and addition of 1 µM TTX (third arrowhead). Scale bars represent 2 g force (vertical bars) and 10 minutes (horizontal bars) for all traces.

CsCl

A Proximal colon

B Distal colon

734 **Fig. S3. Pharmacologic blockade of I_H with CsCl impairs generation of CMCs. (A, B)**
735 Representative force traces from tethered pellet in proximal half (A) or distal half (B) of colon.
736 Addition of 2 mM CsCl (first arrowhead), followed by washout in Krebs (second arrowhead),
737 and addition of 1 μM TTX (third arrowhead). Scale bars represent 2 g force (vertical bars) and
738 10 minutes (horizontal bars) for all traces.

742