Repeated activation of preoptic area-recipient neurons in posterior paraventricular nucleus mediates chronic heat-induced negative emotional and hyperarousal states

Zhiping Cao[1], Wing-Ho Yung[2]*, Ya Ke[1]*

[1] School of Biomedical Sciences, Faculty of Medicine, The Chinese University of Hong Kong, Shatin, Hong Kong, China.

[2] Department of Neuroscience, City University of Hong Kong, Kowloon Tong, Hong Kong, China

* **Corresponding author**:

Wing-Ho Yung, PhD; Department of Neuroscience, City University of Hong Kong, Tat Chee Avenue, Kowloon, Kowloon, Hong Kong. Email: whyung@cityu.edu.hk

Ya Ke, PhD; Lo Kwee Seong Integrated Biomedical Sciences Building, Tai Po Road, Sha Tin District, New Territories, Hong Kong. Email: yake@cuhk.edu.hk

Abstract

Mental and behavioral disorders are associated with extended period of hot weather as found in heatwaves, but the underlying neural circuit mechanism is poorly known. The posterior paraventricular thalamus (pPVT) is a hub for emotional processing and receives inputs from the hypothalamic preoptic area (POA), the well-recognized thermoregulation center. The present study was designed to explore whether chronic heat exposure leads to aberrant activities in POA recipient pPVT neurons and subsequent changes in emotional states. By devising an air heating paradigm mimicking the condition of heatwaves and utilizing emotion-related behavioral tests, viral track tracing, *in vivo* calcium recordings, optogenetic manipulations and electrophysiological recordings, we found that chronic heat exposure for 3 weeks led to negative emotional and hyperarousal states in mice. The pPVT neurons receive monosynaptic excitatory and inhibitory innervations from the POA. These neurons exhibited a persistent increase in neural activity following chronic heat exposure, which was essential for chronic heat-induced emotional changes. Notably, these neurons were also prone to display stronger neuronal activities associated with anxiety responses to stressful situations. Furthermore, we observed saturated neuroplasticity in the POA-pPVT excitatory pathway after chronic heat exposure that occluded further potentiation. Taken together, long-term aberration in the POA to pPVT pathway offers a neurobiological mechanism of emotional and behavioral changes seen in extended period of hot weather like heatwaves.

Introduction

Against a backdrop of global warming, heatwaves that are characterized by abnormally hot weather for extended periods have become more frequent and intense. This is highlighted by that the July 2023 was the hottest month ever recorded by human (Tollefson, 2023). Previous studies have established a correlation between heatwaves and impaired physical health leading to increased mortality (Barriopedro et al., 2011; Garcia-Herrera et al., 2010; Mitchell et al., Mora et al., 2017; 2016; Robine et al., 2008). In addition, elevated temperatures could impact mental health by triggering feelings of anger, stress, aggression, and depression (Beecher et al., 2016). In fact, epidemiological research conducted across various regions extensively demonstrated a positive association between chronic heat exposure and higher hospital admissions for mental and behavioral disorders (Amr et al., 2012; Basu et al., 2018; Dominiak et al., 2015; Hansen et al., 2008; Lee et al., 2002; Mulder et al., 1990; Trang et al., 2015; Whitney et al., 1999). Nevertheless, our understanding of how the brain regulates emotional changes following prolonged heat exposure remains limited.

Current understanding about heat-induced emotional changes mainly focuses on the impact of heat stress on the hypothalamic-pituitary-adrenal (HPA) axis, which leads to abnormal plasma concentrations of the hormone cortisol and neurotransmitters such as 5-hydroxytryptamine (5-HT), noradrenaline and adrenaline, all of which play a significant role in emotional states (McMorris et al., 2006; Wang et al., 2015). However, the relationship between chronic heat exposure and emotional changes is likely to be more complex. Multifaceted

factors, such as the levels of physiological stress (Sampath et al., 2023), changes of neuroendocrine system (Podstawski et al., 2021), sleep disturbances (Altena et al., 2022), adaptation process (Oppermann et al., 2021), even the age and individual health status (Kenny et al., 2010; Malmquist et al., 2022), could interact in a various way to directly and indirectly contribute to emotional changes in response to chronic heat exposure. Thus, gaining insights into the mechanisms behind emotional changes caused by chronic heat exposure is crucial for a more comprehensive understanding of the neurobiological foundations of emotional regulation.

Emerging evidence has suggested that the paraventricular thalamus (PVT) serves as an important integrative node that detects aversive sensory and homeostatic challenges and regulates emotional responses and adaptive behaviors. As such, PVT neurons receive inputs from diverse nuclei in the hypothalamus, midbrain, and hindbrain, and has been implicated in a variety of behavioral responses including arousal (Ren et al., 2018), pain (Jurik et al., 2015), anxiety (Heilbronner et al., 2004; Li et al., 2010; Spencer et al., 2004), fear and fear memory (Do-Monte et al., 2015; Penzo et al., 2021). Therefore, PVT is now considered as an essential component of the emotional processing system in the brain (Penzo et al., 2021).

Among the various inputs from different brain regions to the PVT, the hypothalamus preoptic area (POA) is a well-known thermoregulatory center of the brain (Yu et al., 2016; Zhang et al., 2011; Zhao et al., 2016). Anatomically, the posterior PVT (pPVT) has been reported to receive projections from POA warm-sensitive neurons (Tan et al., 2016). Here, we hypothesize that the direct

projection from POA to pPVT contributes to chronic heat-induced emotional

disturbances. Based on the murine model, we dissected the synaptic

connection from POA to pPVT, tracked its activities and interrogated its

involvement in behavioral responses of the animals under acute and chronic

heat exposure paradigms. Our results uncovered long-term aberration in the

activity of this pathway and pPVT neurons mediating chronic heat-induced

changes in emotional and hyperarousal states, providing a neurobiological

basis of how chronic heat affects mental states.

Results

Chronic heat produces negative emotional and hyperarousal states but

not depression-like behaviors in mice.

c57BL/6 mice were exposed to chronic heat by putting them into a pre-heated

(38 ± 2℃) chamber daily for 21 days with free access of food and water (details

in **Materials and Methods**). In our preliminary experiments, various durations

of daily heat exposure were tested. We settled to 90 minutes due to its

robustness in inducing emotional state-related behavioral changes without

causing collapse of the animals. Throughout the entire process of chronic heat

exposure, we monitored the physiological states of mice daily. The chronic

treatment did not induce abnormal changes in body temperature (Figure 1-

figure supplement 1A and B), but affected normal gain in body weight (Figure

1-figure supplement 1C and D) and reduced food consumption (Figure 1-figure

supplement 1E and F), compared with control mice. On the other hand, mice

subjected to chronic heat displayed a higher prevalence of negative emotional behaviors in various behavioral tests. Specifically, in the elevated plus maze test, chronic heat-exposed mice spent less time in the open arms (Figure 1B and C), indicating increased stress levels. Moreover, in the three-chamber test, these mice displayed reduced interests in exploring the unfamiliar male mouse compared to the inanimate object (Figure 1D and E), suggesting decreased sociability. Similarly, during the female encounter test, these mice performed less engagement with the unfamiliar female mouse compared with the unfamiliar male mouse (Figure 1F and G), indicating decreased innate motivation. Interestingly, the chronic heat-exposed mice displayed decreased latencies for the first attack but increased attack durations in the resident-intruder test (Figure 1H and I), indicating elevated aggression levels.

We also observed that after chronic heat exposure, the mice displayed heightened alertness, suggesting a state of hyperarousal. To confirm this, we quantified the acoustic startle response of the animals by digital video capture of their rapid movements (Pantoni et al., 2020). Chronic heat-exposed mice displayed amplified acoustic startle response characterized by increased body fluctuations within 200 ms of the delivery of a 105 dB sound stimulus (Figure 1J-M and Figure 1-video supplement 1). However, these mice did not exhibit clear hyper-locomotion during the three-chamber test, the female encounter test, or the open field test (Figure 1-figure supplement 1G-I). Collectively, our data demonstrate that chronic heat exposure triggers increased negative emotional states as well as increased arousal in mice.

The fact that the observed changes in emotional and arousal states are the

results of long-term rather than short-term heat exposure is supported by the

lack of effect following one-time acute heat exposure for 90 mins (Figure 1-

figure supplement 2). Furthermore, we did not observe signs of depression in

chronic heat-exposed mice, as there were no significant increases in immobile

time during forced swimming (Figure 1N) and the tail suspension tests (Figure

1O), no obvious decrease in sucrose intake (Figure 1P), and no sign of

decreased exploration as there was no change in the time spent by these

animals in the center zone of the open field (Figure 1Q).

Involvement of the hypothalamic preoptic area to posterior

paraventricular thalamus projection.

To explore the involvement of POA and pPVT in chronic heat exposure induced

behavioral changes, we first studied the effect of heat treatment *per se* on their

activities via c-Fos staining. We found that acute heat treatment (for 90 mins)

significantly increased c-Fos expressions of POA and pPVT neurons (Figure 2-

figure supplement 1A-C), supporting not only the notion that the POA serves as

a thermoregulation center (Morrison et al., 2011) but also the recruitment of

pPVT following heat treatment. Next, to dissect the anatomical relationship

between POA and pPVT especially in the context of heat exposure, we injected

viruses (AAV1-hSyn-cre, AAV-FLEX-TVA-mCherry, AAV-FLEX-RG and

subsequently SAD△G (EnvA) virus) of a monosynaptic retrograde tracing rabies

virus system (details in **Materials and Methods**) into pPVT, which resulted in

clear expression of retrogradely-labeled neurons within POA, its upstream area

(Figure 2A-C). Moreover, a significant overlap (84 ± 5%, 3 mice) was observed

between the retrogradely-labeled and c-Fos expressed neurons in POA

following heat exposure (Figure 2D). Furthermore, almost all the POA recipient

pPVT neurons identified by a Cre-dependent anterograde labelling strategy

((97 ± 3%, 3 mice)) were strongly activated by heat exposure (Figure 2-figure

supplement 2). Together, our results indicate that most of the pPVT-projecting

POA neurons were neurons that respond to heat treatment which would then

recruit their downstream neurons in pPVT by exerting a net excitatory influence.

To functionally characterize the synaptic connection between POA and pPVT,

we administered AAV9-hSyn-ChR2-mCherry into the POA for optogenetic

manipulation of its neurons and their terminals in pPVT. By patch-clamp

recordings in brain slices prepared from these animals, we first confirmed

robust firing of ChR2-mCherry-expressing neurons in POA upon light

stimulation at various frequencies (Figure 2E). When we applied 473 nm light

stimulation to POA terminals in pPVT, in some neurons, a clear inhibitory

outward postsynaptic current was recorded in pPVT neurons held at +10 mV,

which was sensitive to picrotoxin (Figure 2F). In some other neurons, an inward

excitatory postsynaptic current sensitive to CNQX was recorded when the

neurons were held at -70 mV (Figure 2G). Quantification of the presence of

these light-evoked postsynaptic currents revealed the presence of both

excitatory and inhibitory currents in 34.9% of neurons recorded, only inhibitory

currents in 52.4% of neurons and only excitatory currents in 12.7% (Figure 2H).

Since these currents could be completely blocked by tetrodotoxin (TTX) but

restored with the addition of 4-aminopyridine (4-AP), it indicated the

monosynaptic connection, (an example shown in Figure 2I-K). Collectively, our

findings demonstrate that heat responsive POA neurons project directly to

pPVT via both excitatory and inhibitory innervations.

Activity changes of POA recipient pPVT neurons throughout chronic heat

exposure.

We speculated that repeated activation of POA recipient pPVT neurons

underlie the emotional state changes observed following chronic heat exposure.

To gain support for this hypothesis, we first monitored the activity changes of

these neurons during the course of chronic heat exposure by *in vivo* fibre

photometry of calcium signals (Figure 3A and B). We injected AAV1-hSyn-Cre-

EGFP into POA and Cre-dependent AAV9-hSyn-Flex-jGCaMP8F into pPVT

and implanted optical fibers into pPVT neurons after enough expression of

GCaMP. The fibre photometry recordings revealed fluctuations in signals

composed of spontaneous events (details in **Materials and Methods**)

superimposed on the baseline. The detected spontaneous calcium events

presumably represent brief trains or bursts of neuronal firing (Ali et al., 2020).

We first confirmed that the signals were stable for extended period (at least 21

days) in a group of mice not undergoing any treatment (Figure 3-figure

supplement 1). As illustrated in Figure 3C, prior to subjecting the mice to chronic

heat exposure, we recorded the neuronal activities of POA recipient pPVT

neurons for each mouse as the pre-heat control. We then recorded the calcium

events one day after acute heat exposure on day 1 and after chronic heat

exposure on day 21. Interestingly, when compared with the pre-heat condition

of day 1, both the frequency and amplitude during heat exposure on day 1 and

day 21 were significantly increased (Figure 3D and F). However, such increases in calcium events disappeared on the day following heat exposure on day 1 indicating its transient nature (Figure 3E). In contrast, after 21 days of heat exposure, such increases in the frequency of calcium events persisted (Figure 3G). It is worth noting that there were no significant differences in both the frequency and amplitude of calcium events during heat exposure on day 1 and day 21 (Figure 3H).

Based on the increased neuronal activities of POA recipient pPVT neurons after chronic heat exposure, we investigated the impact of activating the pathway from the POA to pPVT on emotional states and arousal levels of mice (Fig. S6A). Utilizing the real-time place preference behavioral paradigm, we noticed that when the blue light was turned on, mice that had already gone through the habituation process quickly entered the chamber with the light turned off (Figure 3-supplement figure 2B and C). This finding suggests that the activation of the POA to pPVT circuit induces an aversive emotional valence. Notably, optogenetic activation did not affect the movement activity of the mice during the experiment (Figure 3-supplement figure 2D). On the other hand, as pupil size is widely recognized as the gold standard for reflecting arousal levels in rodent models (Privitera et al., 2020), we further studied the effect of optogenetic activation of POA excitatory terminals in pPVT in mice under head fixation. The photos (Figure 3-supplement figure 2E) and video (Figure 3-video supplement 1) clearly demonstrated the enlargement of both the pupil (Figure 3-supplement figure 2F and G) and eye sizes (Figure 3-supplement figure 2H and I) of the mice during light stimulation. Collectively, our results demonstrate that POA recipient pPVT neurons exhibited heightened activities after chronic

heat exposure. Moreover, the activation of the POA to pPVT circuit is associated with negative emotional states and higher levels of arousal.

POA recipient pPVT neurons are sufficient and necessary for chronic heat exposure-induced negative emotional and hyperarousal states in mice.

To confirm the direct relationship between repeated activation of POA recipient pPVT neurons and the negative emotional and hyperarousal states in mice, we conducted chronic activation of POA excitatory terminals within pPVT by injecting CaMKII-promotor encoded ChR2 into the POA and implanting an optical fiber in pPVT. We applied a chronic activation protocol, described by Sidor et al. (Sidor et al., 2014), which involved a cycle of 2 minutes on and 2 minutes off, for a total of 20 minutes per day, over a period of up to 21 days. Following chronic optogenetic activation, we observed several behavioral changes in mice that were similar to those seen in mice exposed to chronic heat. Notably, in the elevated plus maze test, mice subjected to chronic optogenetic activation spent less time in the open arms (Figure 4B and C), representing increased stress levels. In the three-chamber test, these mice displayed reduced interests in exploring the unfamiliar male mouse compared to an inanimate object (Figure 4D and E), suggesting decreased sociability. During the female encounter test, these mice engaged less with the unfamiliar female mouse compared with the unfamiliar male mouse (Figure 4F and G), indicating decreased innate motivation. In the resident-intruder aggression test, the chronic heat-exposed mice exhibited both the decreased latencies in initiating the first attack and increased attack durations (Figure 4H and I),

suggesting elevated aggression levels. Additionally, compared to the control

group, mice after chronic optogenetic activation exhibited more pronounced

body fluctuations when captured by a high-resolution camera in response to a

105 dB sound stimulus, indicating the heightened arousal levels (Figure 4J-M).

Our results directly demonstrated that chronic optogenetic activation of POA

excitatory terminals within pPVT circuit could induce negative emotional and

hyperarousal states in mice.

To establish the necessity of POA recipient pPVT neurons for chronic heat

exposure-induced emotional changes, we optogenetically inhibited the POA

recipient pPVT neurons during chronic heat exposure by employing a cycle

stimulation strategy of 3 minutes ON followed by 3 minutes OFF. Only mice

with sufficient virus infection and accurate fiber implantation were included in

the data analysis (Figure 4O and P). The optogenetic inhibition prevented

chronic heat-induced behavioral changes, including anxiety levels (Figure 4Q

and R). sociability (Figure 4S), innate motivation (Figure 4T), aggression levels

(Figure 4U and V) and arousal state (Figure 4W and X). These findings further

supported the essential role of POA recipient pPVT neurons in mediating

chronic heat exposure-induced negative emotional and hyperarousal states.

Chronically activated POA recipient pPVT neurons exhibit exaggerated

response to stressful situations.

To gain further insight into the roles of POA recipient pPVT neurons in stress-

related behaviors, we elucidated the relationship of their activities with specific

behaviors indicative of emotional transitions and hyperarousal states. In the

elevated plus maze test in which the chronic heat-exposed mice spent less time

in the open arms (Figure 1B and C), there were also reduced number of entries

into the open arms (Figure 5B). More detailed analysis revealed that, compared

with untreated mice, these animals more frequently paused at the center of the

maze and then returned to the closed arms (Figure 5C). Furthermore, mice after

chronic heat exposure exhibited a higher chance of running, rather than walking,

back to the closed arms (Figure 5D) and their running speeds significantly

increased compared to their pre-heat condition (Figure 5E), presumably

reflecting a state of increased anxiety. Interestingly, POA recipient pPVT

neurons consistently exhibited a peak in activity when they were about to run

back to the closed arm, while chronic heat-exposed mice exhibited a

significantly larger peak (Figure 5F-H). Notably, no comparable increases of

activities in the POA recipient pPVT neurons were observed when chronic heat-

exposed mice paused at the center and finally walked out to the open arms

(Figure 5-supplement figure 1A-C) or walked to the closed arms (Figure 5-

supplement figure 1D-F). Similar phenomena were found with respect to the

transitions from the open arm to the closed arm, not only in terms of number of

transitions (Figure 5I and J), walking and running behavior (Figure 5K), and

speed of running (Figure 5L), but also a larger calcium peak response aligned

with the running behavior (Figure 5M-O).

In the chronic heat-treated mice, we also observed a context-dependent

change in behavior reminiscent of the hyperarousal states when we introduced

them back into the heat chamber. The mice exhibited apparent locomotion

hyperactivity (Figure 5P), as evidenced by a notable increase in motion speeds

exceeding 400 mm/s (Figure 5Q) and a higher frequency of fast running (Figure

5S). A clear time correlation was found between the heightened calcium activities in POA recipient pPVT neurons and the main component of locomotion hyperactivity, namely fast running under this condition (Figure 5T-V). In contrast, when being placed into a different chamber not associated with heat exposure, no obvious changes were observed among these chronic heat-exposed mice in their total distance traveled (Figure 5-supplement figure 1G), motion speed (Figure 5-supplement figure 1H), number of running exceeding 400 mm/s (Figure 5-supplement figure 1I), the frequency of fast running (Figure 5-supplement figure 1J), or the neuronal activities (Figure 5-supplement figure 1K-M). Collectively, our data suggest the heightened or exaggerated activities of POA recipient pPVT neurons after chronic heat exposure could serve as a driving force for behaviors manifesting negative emotional and hyperarousal states in response to stressful conditions.

Increased pre- and post-synaptic excitability of pPVT neurons but saturated circuitry neuroplasticity capacity following chronic heat exposure.

To explore the potential mechanisms underlying the heightened activities of POA recipient pPVT neurons after chronic heat exposure, we conducted *in vitro* slice recordings on pPVT neurons obtained from mice exposed to room temperature, acute heat, and chronic heat, respectively (Figure 6A and B). Both acute and chronic heat exposure significantly increased the amplitude of miniature inhibitory postsynaptic currents (mIPSCs) in pPVT neurons but there was no difference in mIPSCs amplitude between the acute and chronic heat

groups (Figure 6C). On the other hand, we observed a significant increase in
the frequency of miniature excitatory postsynaptic currents (mEPSCs) specially
in pPVT neurons after chronic but not acute heat exposure (Figure 6F). These
results suggest that while heat exposure could modulate both inhibitory and
excitatory synaptic inputs onto pPVT neurons, there was a differential increase
in presynaptic excitability of the excitatory pathway after chronic heat exposure
but an increase in postsynaptic response to inhibitory input. Furthermore, when
we examined the excitability of pPVT neurons by injecting a sequence of inward
currents up 100 pA to pPVT neurons, pPVT neurons from chronic heat-exposed
mice exhibited significant increases in the number of action potentials from 30
to 70 pA (Figure 6G and H). Further analysis unveiled that these neurons
exhibited a reduced rheobase for action potential generation (Figure 6I).
However, no apparent alterations were observed in the other parameters
(Figure 6-supplement figure 1).

The increase in presynaptic excitability of the POA to pPVT excitatory pathway
suggested plastic changes induced by the chronic heat treatment. While it
would be difficult to follow the excitability of this pathway in vivo during the
chronic heat treatment, we sought to examine the synaptic plasticity capacity
of this pathway before and after chronic heat treatment to shed light on its
involvement. We injected CaMKII-promotor encoded ChR2 into the POA of
mice and divided them into two groups: a control group exposed to room
temperature and a chronic heat group. After 21 days of heat exposure, sagittal
slices that largely preserved the POA to pPVT pathway were prepared from
both the control and chronic heat groups to induce long-term potentiation (LTP)
(Figure 6J). Local optogenetic stimulation vertically (470 nm) through the

objective was applied to elicit light-evoked EPSCs in POA recipient pPVT

neurons and baseline of 10 minutes was recorded. High frequency stimulation

(HFS) is an effective induction protocol for eliciting LTP at excitatory synapses

(Zhu et al., 2016; Grover et al., 2009; Hwong et al., 2008). Therefore, by

delivering blue light at various frequencies (10, 30, 50, 70, 90 Hz) to activate

the POA excitatory terminals within pPVT, we developed an optogenetic HFS

protocol (HFS$_{opto}$: light pulses at 30 Hz, repeated 3 times at 20 s interval)

effective in eliciting potentiated EPSCs on POA recipient pPVT neurons after

baseline recording. HFS$_{opto}$ protocol could successfully induce LTP at POA-

pPVT synapses. The amplitude of light-evoked EPSCs in the control group

following HFS$_{opto}$, at the onset, showed an obvious increase and gradually

became stable potentiation, lasting at least 30 min (150 % ± 18 of baseline, 14

neurons from 5 mice) (Figure 6K and L). However, when we applied the same

protocol to the POA recipient pPVT neurons of mice after chronic heat exposure,

typically no LTP could be induced (Figure 6K-M), suggesting that the pathway

in these mice were already saturated during chronic heat exposure.

Taken together, our findings suggest enhanced excitatory inputs to pPVT

neurons and increased membrane excitability of these neurons may underlie

the behavioral changes observed in mice following chronic heat exposure.

Discussion

In this study, we aimed to uncover the neurobiological basis of how extended

periods of high temperature such as heatwaves impact on emotional

processing in the mammalian brain. We demonstrated that chronic heat

exposure led to negative emotional and hyperarousal states in mice, but not

depression-like behavior. We discovered that hypothalamic POA recipient

pPVT neurons in the thalamus mediate such emotional state changes via

increased baseline activities and excitability in response to stressful conditions.

Presumably, this is attributed to that repeated activation of the POA to pPVT

circuit modulates both pre- and post-synaptic processes leading to their long-

term changes.

Different from utilizing heating panels (Venkatachalam et al., 2007) to

investigate heat-induced thermoregulation, our study employed an air heating

strategy. This choice was motivated by the aim to replicate the realistic

conditions induced by heatwaves. To ensure the well-being of the animals and

prevent unnecessary physiological harm, we maintained temperatures around

38°C, as temperatures exceeding 40℃ can lead to adverse effects on the body

(Deuis et al., 2017). Our findings also revealed that the protocol of chronic heat

exposure at 90 min per day for 21 days did not impact the thermoregulatory

function in mice. In the various emotional state-related behavioral tests we

conducted, including the open field, three-chamber, and female encounter test,

we did not observe locomotion hyperactivity in mice. However, a distinct

acoustic startle response was noted following chronic heat exposure. This may

initially appear contradictory, as hyperarousal states are typically associated

with hyperlocomotion (Zhang et al., 2020). However, POA recipient pPVT

neurons exhibited heightened responses to stressful and aversive situations,

which suggested that chronic heat exposure might primarily induce a state of

increased susceptibility towards anxiety and hyperarousal rather than

persistent hyperarousal states in mice.

There are a number of studies revealing the connections from the POA to pPVT.

Some previous reports have already shown that both excitatory and inhibitory

neurons within POA contribute to thermoregulatory function (Morrison et al.,

2011). Augustine et al. on the other hand demonstrated that POA neurons

expressing nNOS send both excitatory and inhibitory projections to PVT, with

an emphasis on their roles in thirst regulation (Augustine et al., 2018). In

contrast, Allen et al. (Allen et al., 2017) previously described those thirst-related

POA neurons sending projections to PVT mediated reinforcing behavior,

highlighting their independence from POA heat-responsive neurons. The

findings of the present study enriched our understanding on that heat-

responsive neurons in the POA could send both excitatory and inhibitory

projections to pPVT and uncovered the role of the POA recipient pPVT neurons

in chronic heat exposure-induced emotional changes.

Notably, we observed that POA recipient pPVT neurons exhibited higher

baseline activities and displayed heightened response to stress or aversive

stimuli following chronic heat exposure. Interestingly, prior studies have

reported that PVT neurons exhibited stronger activation when mice were

subjected to novel stimuli after repeated restraint stress (Vertes et al., 2015;

Bhatnagar et al., 1998; Heydendael et al., 2011). This effectively supports our

theory of heightened susceptibility to anxiety and our results provide a rationale

for this widely observed phenomenon. Meanwhile, the specific neuronal signals

correlated with actions indicative of heightened anxiety such as fast running to

closed arms and fast running in chronic-heat exposed chamber were not

induced by the motion *per se*. Our recording was conducted when mice were

spontaneously behaving without any pre-set cues, excluding the influence of

salient stimuli on the activity of POA recipient pPVT neurons (Penzo et al., 2021).

In addition to the enhanced excitatory connections from POA to pPVT after chronic heat exposure, we observed an increased amplitude of inhibitory postsynaptic response after both acute and chronic heat exposure. Increased inhibition could act to compensate for the initial over-excitation at critical time, to ensure that neurons briefly activated by stress can be effectively and prevent the occurrence of the excessive stress-related behaviors (Perica et al., 2023). pPVT neurons are anatomically innervated by strong inhibitory projections from different brian nuclei (Penzo et al., 2021). Acute stress-mediated disinhibition of pPVT neurons, as reported by Beas et al. (Beas et al., 2018), can rapidly induce an anxiety-like state in mice. Other studies also showed that inhibition of activated neurons induced by stressful stimuli is effective to avoid abnormal affective behavioral performances (Cruz et al., 2024; Ghasemi et al., 2022). Our results showed that long-term optogenetic inhibition of POA recipient pPVT neurons during chronic heat exposure can effectively prevent changes of emotional and hyperarousal states in mice. Therefore, the increase in mIPSC amplitude observed under both acute and chronic heat conditions in this study suggests the potential presence of negative feedback regulation in response to the over-excitation. However, under chronic heat condition, further persistent significant increase in excitatory synaptic transmission may surpass this effect, leading to a final net activation in the pPVT neurons. However, the potential mechanisms by which increased excitatory synaptic transmission and enhanced postsynaptic inhibitory responses interact to achieve activation or inhibition of neurons themselves still require further study.

Current hypothesis regarding the mechanism of pPVT in mediating anxiety-like

behavior primarily focused on the dopamine-mediated disinhibition of

hypothalamic inhibitory terminals within pPVT during acute stress (Beas et al.,

2018). Our study distinctly revealed that the enhanced excitatory inputs to pPVT

and increased excitability following chronic heat exposure are responsible for

the ensuing negative emotional and hyperarousal states. Interestingly, different

from previous studies that reported a decrease in the amplitude of mIPSC in

pPVT-expressing dopamine receptor 2 neurons 24 hours after acute restraint

stress (Beas et al., 2018), our slice recording from pPVT neurons 24 hours after

acute heat exposure exhibited an increase in mIPSCs amplitude. The reasons

behind these differences could be attributed to the treatment applied and the

specific neuronal types under investigation. Both the enhanced presynaptic

excitability and increased intrinsic postsynaptic excitability following chronic

heat exposure suggested the involvement of long-term neuroplasticity. The fact

that LTP of the POA to pPVT pathway was no longer inducible after chronic

heat exposure strongly suggests that this pathway was potentiated during the

heat treatment, which account for the increased excitability of POA terminals to

pPVT and their increased neuronal activities. Overall, both the enhanced

excitatory inputs to pPVT and increased excitability may play an indispensable

role in chronic heat exposure-induced emotional changes in mice.

There are certain limitations in the scope of the present study. First, our study

was conducted solely on male mice. Further investigation is required to

determine whether chronic heat exposure can elicit similar behavioral

phenotypes in female mice. Also, the pPVT is known to express various peptide

receptors (Curtis et al., 2021). However, the specific roles of these peptide

receptors in chronic heat exposure-induced emotional changes remain unexplored. Some possible mechanisms may contribute to heightened activities of POA recipient pPVT neurons, such as the increased sensitization, the increased activity of hypothalamus-pituitary-adrenal axis, the transition from chronic adaptation to pathological condition (Herman, 2013), and even the chronic effect of dopamine-induced disinhibition of inhibitory terminals within pPVT, all of which required further investigations. Furthermore, POA and pPVT were both previously reported to be activated by stimuli other than heat (Penzo et al., 2021; Zhang et al., 2021). Additional studies are needed to explore the specificity of POA recipient pPVT neurons on chronic heat-induced emotional changes.

In conclusion, our findings provide compelling evidence that the heightened activities of POA recipient pPVT neurons are critically involved in chronic heat-exposure-mediated negative emotional and hyperarousal states, and render mice to enter anxiety-like state more readily, as summarized in Fig. 6N. These results shed a new light on the mechanisms underlying heatwaves-induced emotional changes.

Materials and Methods

Animals

Adult c57BL/6 male mice (25-45g) were used in this study. Animals were bred and maintained by the Laboratory Animal Service Centre of The Chinese University of Hong Kong (CUHK). All experiments were performed following the

CUHK guideline approved by the Animal Experimentations and Ethics Committee.

Stereotaxic surgery and viruses

Mice were anesthetized with ketamine and xylazine and placed gently in a stereotaxic frame (Narashige, Tokyo). A Hamilton syringe (33-gauge) filled with AAV viruses was placed into target brain areas according to the corresponding coordinates: POA (+0.62 mm A/P, 0 mm M/L, 5.83 mm D/V), pPVT (-1.47 mm A/P, 0 mm M/L, 3.13 mm D/V) from brain skull. 0.10-0.35 µl viruses were injected at 10 nl/min speed. After injection, the needle was left in the targeted brain area for additional 10 minutes before retraction. The following viruses were used: AAV1-hSyn-Cre (Addgene, retrograde tracing), AAV1-hSyn-Cre-EGFP (Addgene, optogenetics, photometry recording), Rabies virus (BrainVTA, retrograde tracing), AAV9-hSyn-ChR2-EGFP (Addgene, functional connectivity), AAV9-hSyn-Flex-jGCaMP8F-WPRE (Addgene, photometry recording), AAV9-CaMKII-ChR2-mCherry (Addgene, optogenetic activation), AAV9-hSyn-Dio-eNpHR3.0-EGFP (Addgene, optogenetic inhibition), AAV9-hSyn-Dio-EGFP (Addgene, control experiments). All AAV virus titers were > 5 x 10^{12} particles per ml.

Protocols for chronic heat exposure

The temperature (38 ± 2℃) was adopted based on other reports (Bridges et al., 2012; Cassuto, 1968; Cure, 1989; Jin et a., 2011; Rousset et al., 1984). A

home-made chamber was constructed for chronic heat exposure (80 cm [L] ×

40 cm [W] × 60 cm [H]) which was composed of two built-in components

(heating system and an environmental monitoring sensor). For the heating

system, a heater with (1) adjustable buttons that could control the heating

power and (2) a "dual heating" setting that was able to stop heating when the

temperature exceeded 40℃ in the chamber and restarted heating when the

temperature in the chamber decreased to lower than 36℃. After the air

temperature of the chamber was gradually heated to the targeted temperature

ranges and was monitored to be stable for at least 30 minutes, mice would be

gently put into the chamber for 90 minutes of heat exposure per day for 21 days.

During heat exposure, mice were freely accessible to drinking water and

consuming food. The time point within each daytime to start heat exposure was

random. For the environmental monitoring sensor, it was used to monitor the

chamber temperature value and humidity level (Peng He Dian Zi, Shanghai,

China).

The method for measuring the body temperature of mice was based on a

previous report (Kawakami et al., 2018). Briefly, prior to exposing them to heat,

the mice were gently held by their tails to expose their lower abdomen. An

infrared thermometer sensor was positioned beneath the lower abdomen and

the temperature was measured. This measurement process was repeated

three times for each mouse, and the stable values obtained were then averaged

to determine the specific mouse's body temperature.

Behavioral experiments

A variety of emotion-related behavioral tests were conducted on mice after

acute and chronic heat exposure.

Elevated plus maze test. The self-made elevated plus maze (EPM) apparatus

was made up of two open arms (25 cm [L] × 5 cm [W] × 0.5 cm [H]) and two

closed arms (25 cm [L] × 5 cm [W] × 16 cm [H]) arranged in a "plus" shape and

elevated 50 cm above the floor. The mice were gently placed at the junction of

the open and closed arms while facing toward one of the open arms to initiate

the test. Mice were allowed for a 6-min exploration, and their behavior was

videotaped and then analyzed and quantified by the ANY-Maze tracking

software (Version 4.7, Stoelting CO).

Three-chamber test & Female encounter test. The chambers for both the three-

chamber test and female encounter test were self-made by using an open-field

arena (50 cm [L] x 40 cm [W] x 30 cm [H]) with two identical transparent

chambers (40 cm [L] x 1 cm [W] x 20 cm [H]) with holes in the middle surface,

the stranger male and female mouse were handled for 3 min and then

habituated in a wire cage placed in the 3-chamber apparatus for 5-10 min for

four consecutive days. The tested mouse was habituated to the 3-chamber

apparatus for 10 min with all chambers unobstructed when the handling was

over. The tested mouse was then guided to the center chamber when the

habituation was over, and bidirectional exits were blocked. A wired cup with a

stranger mouse and an empty cup was introduced into the other two chambers,

respectively, and then all chambers were opened for a sociability test. For the

female encounter test, a wired cup with a strange male mouse and a wired cup

with a strange female mouse was introduced into the left and right chambers,

respectively, and all chambers were then opened for a motivation test. Their behavior was videotaped and then analyzed and quantified by the ANY-Maze tracking software (Version 4.7, Stoelting CO).

Resident-intruder aggression test. Mice were fed in a single cage, in which a female mouse, separated by a wire mesh screen, was put in the corner of each cage for seven days to facilitate each tested mouse to build their territory. After chronic heat exposure, the unfamiliar intruder mouse with a color label on the back was introduced to the cage of each tested mouse, and the video was recorded for 15 minutes. Each tested mouse's first attack latency and total attack durations were calculated and counted manually and double-blind by another lab mate.

Acoustic startle response test. The acoustic startle response (ASR) test was conducted based on a recent study (Pantoni et al., 2020). In our study, a sound-proof box was purchased, in which a trumpet, a high-speed camera (200 frames per second, Plexon), and a box made from acrylic plate (10 cm [L] x 5 cm [W] x 25 cm [H]) were prepared. Mice exposed to sound at 105 dB measured by a decibel meter exhibited obvious startle response while 75 dB evoked no response. 105 dB was thus set as the sound decibel of pulse-only stimulation with 0.2 s duration, while 75 dB was set as pre-pulse stimulation with 0.2 s duration. Before the ASR test, mice were placed into an acrylic box for three days of environmental habituation and then for the test. Pre-pulse inhibition (PPI) test started with at least 20 min baseline to make the tested mice quiet and immobile, and then the habituation phase, which consisted of 10 pulses, each 20 s apart, and finally presented with a PPI phase, in which 20 trials: 5

pulse-only trials and 15 pre-pulse & pulse trials were delivered. In the pre-pulse & pulse trials, the pre-pulse was presented about 500 ms before the pulse stimulus. The mice's startle response was captured by a high-speed camera and then analyzed with the machine learning method: DeepLabCut. The brief details were as follows: the mice's body parts were manually labeled and labeled data were input into the google co-lab for the machine learning process until the learning error was lower than 0.5. The successful training data were exported as a .csv file. The startle amplitude of mice was calculated as the average value of all the labeled body parts' vibration amplitude which was shown as a standard deviation within 200 ms after the sound was broadcasted. Each trial was manually checked, and startle response affected by the mice's random moving would not be collected for further analysis. The startle response was visualized with custom MATLAB scripts.

The formula for calculating the PPI ratio was:

PPI ratio = amplitude of pre-pulse & pulse trial / amplitude of pulse-only trial.

Open field test (OFT). Mice were gently placed in the center of the OFT chamber (60 cm [L] × 60 cm [W] × 40 cm [H]) and allowed for free exploration for 10 mins. Mice's behaviors were analyzed and quantified by the ANY-Maze tracking software (Version 4.7, Stoelting CO).

Forced swim test (FST). The apparatus of the forced swim test was a cylindrical tank that was 30 cm high and 20 cm in diameter. And the water was 15 cm tall at room temperature (23-25℃). Tested mice were gently put into water, and the

video was recorded for 6 min. Mice's immobile time in the water was analyzed and quantified by the ANY-Maze tracking software (Version 4.7, Stoelting CO).

Sucrose preference test (SPT). Mice from different groups were habituated for two days for two bottles (one is a sucrose solution bottle, the other is a water bottle, with left and right random placement) in their home cages. After habituation, the sucrose preference of mice from both groups was tested. The intake volume of water and 2% sucrose solution was measured within 2 hours. Sucrose preference was determined as the ratio of the intake volume of sucrose solution to total water consumption.

Tail-suspension test (TST). The tested mice were gently grabbed with the tail and invertedly suspended in front of the recording camera by fixing their tail to the adhesive tape for 6 minutes of recording. Mice immobile time during the suspension was analyzed and quantified by the ANY-Maze tracking software (Version 4.7, Stoelting CO).

Immunohistochemistry

Immunostaining for c-Fos was conducted on four distinct groups of mice: those exposed to acute heat, chronic heat, those injected with rabies virus into the pPVT and exposed to heat, and those with virus-labelled POA recipient pPVT neurons and exposed to heat.

Retrograde tracing. The rabies virus system was adopted for retrograde tracing, as previously reported (Watabe-Uchida et al., 2012). On day 1, 50 nl of AAV1-hSyn-cre, 50nl of AAV-FLEX-TVA-mCherry, and 50 nl of AAV-FLEX-RG were

injected into pPVT. Subsequently, 80 nl of SAD△G virus was injected into

pPVT on day 10. Mice were then allowed seven days for full expression before

being sacrificed.

Tissue preparation. For mice from different groups, brain samples were

prepared as follows: 40 to 60 minutes after exposure to heat, mice were

anesthetized with i.p. injection of a ketamine-xylazine cocktail and

transcardially perfused with 1X phosphate-buffered saline (PBS, Invitrogen)

and then 4% paraformaldehyde (PFA, Sigma-Aldrich). The brain was then

extracted and postfixed overnight in 4% PFA and finally dehydrated with 30%

glucose solution for 48 h. The well-prepared brain samples were embedded in

OCT (ThermoFisher) and sliced into 30-μm coronal sections by a cryostat

(Leica) and stored in 1X PBS before immunohistochemistry.

Immunofluorescent staining. Brain sections were blocked in 5% normal goat

serum (ThermoFisher) in PBS with 0.3% Triton X-100 (Sigma Aldrich) for 40

min and then incubated with primary rabbit anti-c-Fos antibody (1:2000, Cell

Signaling Technology) at 4°C overnight. After several thorough washing pieces

in PBS, the sections were incubated with goat anti-rabbit IgG-Alexa Fluor 488

secondary antibody (1:1000, Invitrogen) in a blocking solution for 2 hours. After

that, the sections were rinsed in PBS again and finally mounted onto glass

slides. Microscopic images were taken under a confocal laser scanning

microscope (C1, Nikon).

Neuronal counting. As for the statistical analysis of c-Fos positive neurons,

profile counting based on the rectangular square drawn in each picture was

performed on image J. For every counted brain area, c-Fos positive neurons

on three consecutive brain slices were selected and measured. And the value

averaged from profile counting on three consecutive brain slices was regarded

as the c-Fos positive neuronal number of specific brain nuclei for one mouse.

Optogenetic experiments

For activation: 100 nL of AAV9-CaMKII-ChR2-mCherry was injected into POA,

optical fibers in pPVT. For inhibition: 100 nL of AAV1-hSyn-Cre-EGFP was

injected into POA, 200 nL of AAV9-hSyn-Dio-eNpHR3.0-EGFP or AAV9-hSyn-

Dio-EGFP in pPVT.

Real-time place preference test. Mice were placed into a two-chamber box for

10 minutes of habituation to make sure that mice had no preference for either

side of the chamber. Another 10 minutes' recording was started. The left

chamber was set as the light ON compartment while the other was the light

OFF chamber. Once mice entered light ON chamber, 473-nm light with 10 ms,

10 Hz (unless otherwise indicated, Newdoon Technology) was delivered via an

optic cable (200-μm core, 0.37 NA, Doric Lens) while the light was turned off

when mice entered light OFF chamber. Laser power was 5 mW measured at

the tip of the fiber, which was implanted 0.1-0.2 mm above the targeted nucleus.

Elevated plus maze test, three-chamber test, and female encounter test. To

explore the effect of optogenetic activation of POA excitatory terminals within

pPVT on the levels of anxiety, sociability, and motivation in mice, mice in these

three behavioral tests were recorded with 3 minutes as the baseline, followed

by another 3 minutes' delivery of 473-nm light stimulation via an optic cable

(200-μm core, 0.37 NA, Doric Lens).

Measurement of pupil size. To directly reflect the effect of POA excitatory

terminals within pPVT on hyperarousal levels, mice were head-fixed and 473-

nm light was delivered with 10 Hz, 10 ms at the mode of 10 s light off, 10 s light

on, and 10 s light off to measure the changes of pupil size. And pupil size of

mice during the whole process was synchronously captured through a high-

speed camera at 30 frames per second. All pictures were finally integrated into

video format through the FFmpeg program. The changes in pupil size among

established videos were further analyzed by the DeepLabCut method (Mathis

et al., 2018) for precise tracking and plotting. The time point for statistical

analysis of normalized pupil size was at 5s for Pre, 15s for Light, and 25s for

Post, respectively.

Chronic optogenetic stimulation. Mice are subjected to 473-nm light stimulation

with 10 ms, 10 Hz (unless otherwise indicated, Newdoon Technology) via an

optic cable (200-μm core, 0.37 NA, Doric Lens) 2 mins-on, 2 mins-off for 20 min

per day for up to 21 days (Sidor et al., 2014). Laser power does not exceed 5

mW measured at the tip of the fiber.

Optogenetic inhibition. Mice exposed to chronic heat were synchronously

illuminated with a 589-nm light laser (Newdoon Technology) to inhibit POA

recipient pPVT neurons specifically. The protocol was set at 0.0001 Hz, 9999

ms, 20 mW with a cyclical mode consisting of 3 minutes' light on and 3 minutes'

light off for the whole process of chronic heat exposure. The emotion-related

behaviors and hyperarousal states were measured as formerly mentioned and

analyzed by Any-maze software.

Fiber photometry

100 nL of AAV1-hSyn-Cre-EGFP in POA, 200 nL of AAV9-hSyn-Flex-

jGCaMP8F-WPRE in pPVT, optical fibers in pPVT. Calcium signals were

recorded for 5 minutes for each mouse before, during heat exposure on day 1,

the second day after heat exposure on 1st day, and after chronic heat exposure

on day 22. The frequency and amplitude of calcium events were collected for

data analysis.

Before and after chronic heat exposure, the performance of each mouse in the

elevated plus maze and the chronic heat-exposed chamber was in combination

with calcium recording for 10 minutes, respectively. Targeting of event-related

time points and calculation of speed were achieved through the SLEAP

machine learning method (Pereira et al., 2022).

Fiber photometry data were collected with a TDT system at a sampling

frequency of 1017 Hz. The LED power at the tip of the patch cord was less than

20 µW. The 405- and 465- signals were simultaneously recorded. The

isosbestic 405 nm control signal was filtered using a polyfit regression to limit

the influence of fluorescence decay during the session. And the fitted control

signal was then subtracted from the 465-signal to remove artifacts from the

intracellular calcium-dependent GCaMP fluorescence. The calculation for the

dynamics of fluorescence in chronic calcium recording was performed using the

formula $\triangle F/F=(F_{465}-F_{405}) / F405$. As corroborated by previous studies (Shao et

al., 2022; Xia et al., 2021), a calcium signal wave exceeding $\mu + 3\sigma$ was

regarded as a fluorescence transient. Here, μ and σ respectively denote the

average and the standard deviation of the baseline signal of each mouse. The

baseline signal for each mouse was randomly selected, with a duration ranging

from 5 to 10 seconds. As for the activity changes of POA recipient pPVT

neurons during emotional transitions and hyperarousal states, the $\triangle F/F$ was

then analyzed using a z-Score relative to the mean and standard deviation of

the session. Data were visualized and analyzed with custom MATLAB scripts.

Electrophysiological recordings

Brain slice preparation and whole-cell recording. As for the confirmation of

functional connectivity, coronal mice brain slices were prepared as follows.

Mice quickly anesthetized by isoflurane were then transcardially perfused with

pH7.4 NMDG artificial cerebrospinal fluid (aCSF) in which contains (mM):

NMDG 92, KCl 2.5, NaH2PO4 1.25, NaHCO3 20, HEPES 10, Glucose 25, Na-

ascorbate 5, thiourea 2, Na-pyruvate 3, MgSO4 10, CaCl2 0.5, N-acetyl-L-

cysteine 12. Once completion of perfusion, mice's brain was quickly and

smoothly extracted and placed into cutting NMDG solution for frozen 30 s. And

coronal sections at 270-μm were prepared with a vibratome (Campden

5100MZ-PLUS Vibrotome) and then transferred to a chamber with 34 °C NMDG

aCSF for the first 15 minutes of recovery. And subsequently, brain slices were

further gently transferred into another chamber holding aCSF at room

temperature containing HEPES, pH 7.4, which contains (mM): NaCl 92, KCl 2.5,

NaH2PO4 1.25, NaHCO3 20, HEPES 10, Glucose 25, Na-ascorbate 5, thiourea

2, Na-pyruvate 3, MgSO4 10, CaCl2 0.5, N-acetyl-L-cysteine 12. After 40 to 60

minutes of recovery, the brain slices were placed into a recording chamber with

normal aCSF, pH 7.4, which contained (mM): NaCl 125, KCl 2.5, Glucose 11,

NaHCO3 26, NaH2PO4 1.25, CaCl2 2, and MgCl2 2 for neuronal recording.

Functional connectivity. To record optically evoked postsynaptic currents, we

used an internal solution containing (mM): K-gluconate 130, KCl 10, HEPES 10,

EGTA 1, MgCl2 2, Na2-ATP 2, Na3-GTP 0.4. Light pulses at 470 nm were

delivered through a light stimulator (Polygon400 DSI-E-0470-0590-NK1

Dynamic Spatial Illuminator) to activate POA ChR2-EGFP-expressing terminals

while patched pPVT neurons were held at -70mV. The neuron with access

resistance <20 MΩ and leak current <100 pA was included for data collection.

Measurements of synaptic activity and intrinsic properties. To measure the

changes in synaptic transmission of pPVT neurons after chronic heat exposure

and acute heat exposure, coronal mice's brain slices were prepared as

mentioned above. The internal solution for both miniature inhibitory (sIPSCs,

held at +10 mV) and excitatory (sEPSCs, held at -70 mV) post synaptic currents

recording was Cesium-gluconate based solution with pH 7.3-7.4 which contains

(mM): CsMSF 130, HEPES 10, EGTA 1, MgCl2-6H2O 2, Na2ATP 2, NaGTP

0.4, QX-314-Cl 5, TEA-OH 5. And Tetrodotoxin (0.5 mM) was present for both

miniature EPSCs (mEPSCs) and miniature IPSCs (mIPSCs) recording. Once

completion of recording, brain slices were fixed overnight with 4% PFA and the

correct position of recorded pPVT neurons was identified by staining with

streptavidin-conjugated Alex 405 (ThermoFisher). For measuring the firing

frequency under current clamp recording, steady state current was injected in

+10 pA increments from 0 to 100 pA. All action potential properties and

excitability recordings in Figure S8 were performed in the presence of 10 µM

CNQX and 100 µM picrotoxin. As mentioned in the previous paper (Knowland

et al., 2017), the firing rate evoked by current injections and the half-width were

calculated in Clampfit software (Molecular Devices) by using peak detection

function and full-width at half max amplitude, respectively. The threshold was

measured as the change in the voltage from rest at which the slope =20 V/s.

Membrane resistance was calculated from the change in voltage elicited after

a 50 ms 5 mV hyperpolarizing step from -70 mV (the last 10 ms of the step from

baseline was taken as $\triangle$V). Capacitance and membrane time constant were

calculated by Clampfit during the first minute after breaking in to cell.

Long-term potentiation. To measure the zneuroplasticity of the POA to pPVT

pathway, two batches of mice with enough expression of CaMKII-ChR2-

mCherry within the POA were prepared: One serving as the control group and

the other as the chronic heat group. Sagittal brain slices from the mice were

prepared, and a low-chloride internal solution was used to record pPVT neurons.

EPSCs evoked by focal optogenetic stimulation through the objective

(approximately 2-3 mm above the surface of brain slice) in a top-down vertical

direction were elicited using a Polygon400 Multiwavelength Dynamic Patterned

Illuminator (Mightex). A circular area (roughly 500 µm diameter) covering the

recorded cell soma was illuminated with brief pulses of blue light (470 nm,

duration: 2 ms). LTP induced by optogenetic stimulation of pPVT neurons was

elicited by HFS consisting of three episodes of 30 Hz blue light pulses at 20 s

intervals.

All above signals were collected using a MultiClamp 700B amplifier controlled

by Clampfix 10.4 software via a Digital data 1550 interface (Molecular Devices).

Electrical signals were filtered at 3 kHz, digitized at 10 kHz, and further

analyzed using Clampfit 10.7 (Molecular Devices).

Statistical analysis

Statistical analysis was performed using GraphPad Prism 8.0. Values were

shown as mean ± standard error of the mean (SEM). Student's t-test (paired or

unpaired, parametric or nonparametric), one-way ANOVA with Tukey post-hoc

test, or two-way ANOVA followed by Sidak post-hoc test, were conducted for

statistical analysis. The value for statistical significance was $p < 0.05$. For each

experiment, the detailed statistics are described in the corresponding figure

legend.

Data availability

All data are contained in the main text, extended data or the supplementary

materials are available from the author upon reasonable request. The source

data underlying Figs 1C, E, G-I, L, M-Q, 2H, K, 3D-H, 4C, E, G-I, L, M, P-W,

5B-E, H-L, O-S, V, 6C-F, H, I, L, M and Figure 1-figure supplement 1A-I, Figure

1-figure supplement 2B, D, F, G, H, K, L, Figure 2-figure supplement 1C, Figure

3-figure supplement 1C, D, Figure 3-figure supplement 2C, D, G, I, Figure 5-

figure supplement 1C, F, G-J, M, Figure 6-figure supplement 1A-F are provided

as a Source Data file.

Acknowledgements: This work was funded by following funding sources:

HKRGC C4012-22G & HMRF 09203236. We thank Dr. Xu Tao, Romeo Goh,

Zhuyun and Rong Kanglin for their technical assistance.

Conflict of interest: The authors declare no competing financial interests.

Author contributions

Y Ke and WH Yung perceived the project, designed the experiments, and

analyzed the data. ZP Cao designed the experiments, collected, and analyzed

the data. All authors discussed the findings and contributed to the writing of the

manuscript.

Author ORCIDs

Ya Ke https://orcid.org/0000-0001-8166-6653

Zhiping Cao https://orcid.org/0000-0001-5560-7385

Ethics

All experimental procedures were approved by the Animal Experimentations

and Ethics Committee (AEEC number: 18-232-MIS-5-C) at the Chinese

University of Hong Kong (Animal license number: (23-120) in DH/HT&A/8/2/1

Pt.45). All surgery was performed under deep surgical anesthesia and every

effort was made to minimize animal suffering.

References

Ali F, Kwan AC (2020) **Interpreting in vivo calcium signals from neuronal**

cell bodies, axons, and dendrites: a review *Neurophotonics* **7**:011402.

Allen WE, DeNardo LA, Chen MZ, Liu CD, Loh KM, Fenno LE, Ramakrishnan

C, Deisseroth K, Luo LQ (2017) **Thirst-associated preoptic neurons encode**

an aversive motivational drive *Science* **357**:1149-1155.

Altena E, Baglioni C, Sanz-Arigita E, Cajochen C, Riemann D (2023) **How to**

deal with sleep problems during heatwaves: practical recommendations

from the European Insomnia Network *J Sleep Res* **32**:e13704.

Amr M, Volpe FM (2012) **Seasonal influences on admissions for mood**

disorders and schizophrenia in a teaching psychiatric hospital in Egypt *J*

Affect Disord **137**:56-60.

Augustine V, Gokce SK, Lee S, Wang B, J. Davidson T, Reimann F, Gribble F,

Deisseroth K, Lois C, Oka Y (2018) **Hierarchical neural architecture**

underlying thirst regulation *Nature* **555**:204-209.

Barriopedro D, Fischer EM, Luterbacher J, Trigo RM, Garcia-Herrera R (2011) **The hot summer of 2010: redrawing the temperature record map of Europe** *Science* **332**:220-224.

Basu R, Gavin L, Pearson D, Ebisu K, Malig B (2018) **Examining the association between apparent temperature and mental health related emergency room visits in California** *Am J Epidemiol* **187**:726-735.

Beas BS, Wright BJ, Skirzewski M, Leng Y, Hyun JH, Koita O, Ringelberg N, Kwon HB, Buonanno A, Penzo MA (2018) **The locus coeruleus drives disinhibition in the midline thalamus via a dopaminergic mechanism** *Nat Neurosci* **21**:963-973.

Beecher ME, Eggett D, Erekson D, Rees LB, Bingham J, Klundt J, Bailey RJ, Ripplinger C, Kirchhoefer J, Gibson R, Griner D, Cox JC, Boardman RD (2016) **Sunshine on my shoulders: weather, pollution, and emotional distress** *J Affect Disord* **205**:234-238.

Bhatnagar S, Dallman M (1998) **Neuroanatomical basis for facilitation of hypothalamic–pituitary–adrenal responses to a novel stressor after chronic stress** *Neuroscience* **84**:1025-1039.

Bridges MT, Tulapurkar ME, Shah NG, Singh IS, Hasday JD (2012) **Tolerance for chronic heat exposure is greater in female than male mice** *Int J Hyperthermia* **28**:747-755.

Cassuto Y (1968) **Metabolic adaptations to chronic heat exposure in the golden hamster** *Am J Physiol* **214**:1147-1151.

Cruz B, Vozella V, Borgonetti V, Bullard R, Bianchi PC, Kirson D, Bertotto LB,

Bajo M, Vlkolinsky R, Messing RO, Zorrilla EP, Roberto M (2024)

Chemogenetic inhibition of central amygdala CRF-expressing neurons

decreases alcohol intake but not trauma-related behaviors in a rat model

of post-traumatic stress and alcohol use disorder *Mol Psychiatry* Online

ahead of print 21 March 2024; doi: 10.1038/s41380-024-02514-8.

Cure M (1989) **Plasma corticosterone response in continuous versus**

discontinuous chronic heat exposure in rat *Physiol Behav* **45**:1117-1122.

Curtis GR, Oakes K, Barson JR (2021) E**xpression and Distribution of**

Neuropeptide-Expressing Cells Throughout the Rodent Paraventricular

Nucleus of the Thalamus *Frontiers in Behavioral Neuroscience* **14**:634163.

Deuis JR, Dvorakova LS, Vetter I (2017) **Methods Used to Evaluate Pain**

Behaviors in Rodents *Front Mol Neurosci* **10**:284.

Dominiak M, Swiecicki L, Rybakowski J (2015) **Psychiatric hospitalizations**

for affective disorders in Warsaw, Portland: Effects of season and

intensity of sunlight *Psychiatry Res* **229**:287-294.

Do-Monte FH, Quiñones-Laracuente K, Quirk GJ (2015) **A temporal shift in**

the circuits mediating retrieval of fear memory *Nature* **519**:460-463.

García-Herrera R, Díaz J, Trigo RM, Fischer EM (2010) **A review of the**

European summer heat wave of 2003 *Crit Rev Environ Sci Technol* **40**:267-

306.

Ghasemi M, Navidhamidi M, Rezaei F, Azizikia A, Mehranfard N (2022)

Anxiety and hippocampal neuronal activity: Relationship and potential

mechanisms *Cog Affect Behav Neurosci* **22**:431-449.

Grover LM, Kim E, Cooke JD, Holmes WR (2009) **LTP in hippocampal area**

CA1 is induced by burst stimulation over a broad frequency range

centered around delta *Learn Mem* **16**:69-81.

Hansen A, Bi P, Nitscheke M, Ryan P, Pisaniello D, Tucker G (2008) **The effect**

of heat waves on mental health in a temperate Australian city *Environ*

Health Perspect **116**:1369-1375.

Heilbronner U, Kampen MV, Flüggeg G (2004) **The Alpha-2B Adrenoceptor**

in the Paraventricular Thalamic Nucleus is Persistently Upregulated by

Chronic Psychosocial Stress *Cell Mol Neurobiol* **24**:815-831.

Herman JP (2013) **Neural control of chronic stress adaptation** *Front Behav*

Neurosci **7**: 61.

Heydendael W, Sharma K, Iyer V, Luz S, Piel D, Beck S, Bhatnagar S (2011)

Orexins/hypocretins act in the posterior paraventricular thalamic nucleus

during repeated stress to regulate facilitation to novel stress

Endocrinology **152**:4738-4752.

Hwong, Udaka Y, Kauer JA (2008) **High-Frequency Afferent Stimulation**

Induces Long-Term Potentiation of Field Potentials in the Ventral

Tegmental Area *Neuropsychopharmacology* **33**:1704-1712.

Jin Y, Hu YX, Han DP, Wang M (2011) **Chronic heat stress weakened the**

innate immunity and increased the virulence of highly pathogenic avian

influenza virus H5N1 in mice *J Biomed Biotechnol* **2011**:367846.

Jurik A, Auffenberg E, Klein S, Deussing JM, Schmid RM, Wotjak CT,

Thoeringer CK (2015) **Roles of prefrontal cortex and paraventricular**

thalamus in affective and mechanical components of visceral nociception.

Pain **156**:2479-2491.

Kawakami Y, Sielski R, and Kawakami T (2018) **Mouse Body Temperature**

Measurement Using Infrared Thermometer During Passive Systemic

Anaphylaxis and Food Allergy Evaluation *J Vis Exp* **139**:58391.

Kenny GP, Yardley J, Brown C, Sigal RJ, Jay O (2010) **Heat stress in older**

individuals and patients with common chronic diseases *CMAJ* **182**:1053-

1060.

Knowland D, Lilascharoen V, Pacia CP, Shin S, Wang EHJ, Lim BK (2017)

Distinct Ventral Pallidal Neural Populations Mediate Separate Symptoms

of Depression *Cell* **170**:284-297.

Lee HJ, Kim L, Joe SH, Suh KY (2002) **Effects of season and climate on the**

first manic episode of bipolar affective disorder in Korea *Psychiatry Res*

113:151-159.

Li YH, Li S, Wei CG, Wang HY, Sui N, Kirouac GJ (2010) **Orexins in the**

paraventricular nucleus of the thalamus mediate anxiety-like responses

in rats *Psychopharmacology (Berl)* **212**:251-265.

Malmquist A, Hjerpe M, Glaas E, Karlsson H, Lassi T (2002) **Elderly People's**

Perceptions of Heat Stress and Adaptation to Heat: An Interview Study *Int*

J Environ Res Public Health **19**: 3775.

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M

(2018) **DeepLabCut: markerless pose estimation of user-defined body**

parts with deep learning *Nat Neuroscience* **21**:1281-11289.

McMorris T, Swain J, Smith M, Corbett J, Delves S, Sale C, Harris RC, Potter

J (2006) **Heat stress, plasma concentrations of adrenaline, noradrenaline,**

5-hydroxytryptamine and cortisol, mood state and cognitive performance.

Int J Psychophysiol **61**:204-215.

Mitchell D, Heaviside C, Vardoulakis S, Huntingford C, Masato G, Guillod PB,

Frumhoff P, Bowery A, Wallom D and Allen M (2016) **Attributing human**

mortality during extreme heat waves to anthropogenic climate change.

Environ Res Lett **11**:74006.

Mora C, Dousset B, Caldwell IR, Powell FE, Geronimo RC, Bielecki CR,

Counsell CWW, Dietrich BS, Johnston ET, Louis LV, Lucas MP, McKenzie MM,

Shea AG, Tseng Han, Giambelluca TW, Leon LR, Hawkins E & Trauernicht C

(2017) **Global risk of deadly heat** *Nat Clim Change* **7**:501-506.

Morrison SF, Nakamura K (2011) **Central neural pathways for**

thermoregulation *Front Biosci* **16**:74-104.

Mulder RT, Cosgriff JP, Smith AM, Joyce PR (1990) **Seasonality of mania in**

New Zealand *Aust N Z J Psychiatry* **24**:187-190.

Oppermann E, Kjellstrom T, Lemke B, Otto M, Lee JKW (2021) **Establishing**

intensifying chronic exposure to extreme heat as a slow onset event with

implications for health, wellbeing, productivity, society and economy *Curr*

Opin Environ Sustain **50**:225-235.

Pantoni MM, Herrera GM, Van Alstyne KR, Anagnostaras SG (2020)

Quantifying the Acoustic Startle Response in Mice Using Standard Digital

Video *Front Behav Neurosci* **14**:83.

Penzo MA, Gao Claire (2021) **The paraventricular nucleus of the thalamus:**

an integrative node underlying homeostatic behavior *Trends Neurosci*

44:538-549.

Penzo MA, Robert V, Tucciarone J, De Bundel D, Wang MH, Van Aelst L,

Darvas M, Parada LF, Palmiter R, He M, Huang ZJ, Li B (2015) **The**

paraventricular thalamus controls a central amygdala fear circuit *Nature*

519:455-459.

Pereira TD, Tabris N, Matsliah A, Turner D, Li JY, Ravindranath S,

Papadoyannis ES, Normand E, Deutsch D, Wang Z, McKenzie-Smith GC,

Mitelut CC, Castro MD, D'Uva J, Kislin M, Sanes DH, Kocher SD, Wang SSH,

Falkner AL, Shaervitz JW, Murthy M (2022) **SLEAP: A deep learning system**

for multi-animal pose tracking *Nat Method* **19**:486-495.

Perica MI, Luna B (2023) **Impact of stress on excitatory and inhibitory**

markers of adolescent cognitive critical period plasticity *Neurosci*

Biobehav Rev **153**:105378.

Podstawski R, Boryslawski K, Pomianowski A, Krystkiewicz W, Żurek P (2021) **Endocrine Effects of Repeated Hot Thermal Stress and Cold Water Immersion in Young Adult Men** *Am J Mens Health* **15**:15579883211008339.

Privitera M, Ferrari KD, von Ziegler LM, Sturman O, Duss SN, Floriou-Servou A, Germain PL, Vermeiren Y, Wyss MT, P De Deyn P, Weber B, Bohacek J (2020) **A complete pupillometry toolbox for real-time monitoring of locus coeruleus activity in rodents** *Nat Protoc* **15**:2301-2320.

Ren SC, Wang YL, Yue FG, Cheng XF, Dang RZ, Qiao QC, Sun XQ, Li X, Jiang Q, Yao JW, Qin H, Wang GZ, Liao X, Gao D, Xia JX, Zhang J, Hu B, Yan JA, Wang YJ, Xu M, Han YY, Tang XD, Chan XW, He C, Hu ZA (2018) **The paraventricular thalamus is a critical thalamic area for wakefulness** *Science* **362**:429-434.

Robine JM, Cheung SLK, Le Roy S, Van Oyen H, Griffiths C, Michel JP, Herrmann FR (2008) **Death toll exceeded 70 000 in Europe during the summer of 2003** *C R Biol* **331**:171-178.

Rousset B, Cure M, Jordan D, Kervran A, Bornet H, Mornex R (1984) **Metabolic alterations induced by chronic heat exposure in the rat: the involvement of thyroid function** *Pflugers Arch* **401**:64-70.

Sampath V, Shalakhti O, Veidis E, Efobi JAI, Shamji MH, Agache I, Skevaki C, Renz H, Nadeau KC (2023) **Acute and chronic impacts of heat stress on planetary health** *Allergy* **78**:2109-2120.

Shao J, Gao DS, Liu YH, Chen SP, Liu NA, Zhang L, Zhou XY, Xiao Q, Wang LP, Hu HL, Wang F (2022) **Cav3.1-driven bursting firing in ventromedial**

hypothalamic neurons exerts dual control of anxiety-like behavior and energy expenditure *Mol Psychiatry* **27**:2901-2913.

Sidor MM, McClung C (2014) **Timing matters: using optogenetics to chronically manipulate neural circuitry and rhythms** *Front Behav Neurosci* **8**:41.

Spencer SJ, Fox JC, Day TA (2004) **Thalamic paraventricular nucleus lesions facilitate central amygdala neuronal responses to acute psychological stress** *Brain Res* **997**:234-237.

Tan CL, Cooke EK, Leib DE, Lin YC, Daly GE, Zimmerman CA, Knight ZA (2016) **Warm-sensitive neurons that control body temperature** *Cell* **167**:47-59.

Tollefson J (2023) **Earth's hottest month: these charts show what happened in July and what comes next** *Nature* **620**:703-704.

Trang PM, Rocklöv J, Giang KB, Van Minh H, Tinh LT, Nilsson M (2015) **Weather variations and hospital admissions for depressive disorders: a case study in Hanoi** *Ann Psychiatry Ment Health* **5**:1095.

Venkatachalam K, Montell C. TRP channels (2007) **Annu Rev Biochem** 76:387-417.

Vertes RP, Linley SB, Hoover WB (2015) **Limbic circuitry of the midline thalamus** *Neurosci Biobehav Rev* **54**:89-107.

Wang L, Liu FD, Luo Y, Zhu LQ, Li GH (2015) **Effect of acute heat stress on adrenocorticotropic hormone, cortisol, interleukin-2, interleukin-12 and apoptosis gene expression in rats** *Biomed Rep* **3**:425-429.

Watabe-Uchida M, Zhu L, Ogawa SK, Vamanrao A, Uchida N (2012) **Whole-Brain Mapping of Direct Inputs to Midbrain Dopamine Neurons** *Neuron* **74**:858-873.

Whitney DK, Sharma V, Kueneman K (1999) **Seasonality of manic-depressive illness in Canada** *J Affect Disord* **55**:99-105.

Xia G, Han Y, Meng F, He Y, Srisai D, Farias M, Dang MH, Palmiter RD, Xu Y, Wu Q (2021) **Reciprocal control of obesity and anxiety-depressive disorder via a GABA and serotonin neural circuit** *Mol Psychiatry* **26**:2837-2853.

Yu S, Qualls-Creekmore E, Rezai-Zadeh K, Jiang YY, Berthoud HR, Morrison CD, Derbenev AV, Zsombok A, Münzberg H (2016) **Glutamatergic preoptic area neurons that express leptin receptors drive temperature-dependent body weight homeostasis** *J Neurosci* **36**:5034-5046.

Zhang GW, Shen L, Tao C, Jung AH, Peng B, Li Z, Zhang LI, Whit Tao HZ (2021) **Medial preoptic area antagonistically mediates stress-induced anxiety and parental behavior** *Nat Neurosci* **24**:516-528.

Zhang QH, Hao JW, Li GL, Ji XJ, Zhou M, Yao YM (2020) **Long-lasting neurobehavioral alterations in burn-injured mice resembling post-traumatic stress disorder in humans** *Exp Neurol* **323**:113084.

Zhang Y, Kerman IA, Laque A, Nguyen P, Faouzi M, Louis GW, Jones JC, Rhodes C, Münzberg H (2011) **Leptin-receptor-expressing neurons in the dorsomedial hypothalamus and median preoptic area regulate sympathetic brown adipose tissue circuits** *J Neurosci* **31**:1873-1884.

Zhao ZD, Yang WZ, Gao CC, Fu X, Zhang W, Zhou Q, Chen WP, Ni XY, Lin

JK, Yang J, Xu XH, Shen WL (2016) **A hypothalamus circuit that controls**

body temperature *Proc Natl Acad Sci USA* **114**:2042-2047.

Zhu YT, Wang RB, Wang YH (2016) **A Comparative Study of the Impact of**

Theta-Burst and High-Frequency Stimulation on Memory Performance

Front Hum Neurosci **10**:19.

1079

Figure legends

Figure 1. Chronic heat exposure produces negative emotional and

hyperarousal states but not depression-like behaviors. (A) Experimental

schematics. Mice (n=10 in each group) were divided into Control and Heat

groups and conducted with chronic exposure to room temperature and heat

conditions, respectively, followed by behavioral tests. **(B, C)** The heatmap of

representative tracking trace examples in EPM test and the time spent in the

open arms of EPM (Mann-Whitney unpaired two-tailed U test; U=0, ***p<0.001).

(D, E) The heatmap of representative tracking trace examples in TCT and the

interaction time with an unfamiliar male mouse relative to the inanimate object

in TCT (Mann-Whitney unpaired two-tailed U test; U=13, ***p=0.0039). **(F, G)**

The heatmap of representative tracking trace examples in FET and the time

surrounding the unfamiliar female mouse compared to an unfamiliar male

mouse in FET (Mann-Whitney unpaired two-tailed U test; U=10, **p=0.0015).

(H, I) The 1st-time attack latency (Mann-Whitney unpaired two-tailed U test;

U=21, *p=0.0288) and the attack durations (Mann-Whitney unpaired two-tailed

U test; U=23, *p=0.0433) in the aggression test. **(J, K)** Visualized and

representative acoustic startle response example (left panel) and corresponding labeled body parts' skeletons (right panel) in ASR test from Control and Chronic Heat group.**(L, M)** The startle amplitude (Mann-Whitney unpaired two-tailed U test; U=6, ***p=0.0003) and the pre-pulse / pulse ratio (Mann-Whitney unpaired two-tailed U test; U=26, p=0.0753) in ASR test. **(N-Q)** The immobile time in the forced swim test (Mann-Whitney unpaired two-tailed U test; U=35, p=0.2799) and in the tail suspension test (Mann-Whitney unpaired two-tailed U test; U=26, p=0.0753; The percentage of sucrose intake in the sucrose preference test (Mann-Whitney unpaired two-tailed U test; U=49.5, p=0.4887); The time spent in the center of the open field test (Mann-Whitney unpaired two-tailed U test; U=35.5, p=0.2888). *p<0.05, **p<0.01, ***p<0.001, N.S.: not significant.

Figure 2. Involvement of the hypothalamic preoptic area to posterior paraventricular thalamus projections. (A) The strategy of virus injection followed by heat exposure-induced c-Fos staining (n=3 mice). **(B)** The representative microphotograph showed the expression of rabies virus in the pPVT regions. **(C)** Rabies virus retro-labeled POA neurons. Upper: magnification: x4, scale bar: 400 μm; Lower: magnification: x10, scale bar: 40 μm; The top right corner of the lower picture: magnification: x60, scale bar: 5 μm. **(D)** The representative microphotograph showed that most of pPVT rabies virus retro-labeled POA neurons co-stained with heat exposure-induced c-Fos. Upper: magnification: x10, scale bar: 40 μm; Lower: magnification: x60, scale bar: 10 μm. **(E)** The strategy of virus injection (n=5 mice) and patch-clamp

recording was performed on POA expressing ChR2-mCherry neurons and

pPVT neurons using potassium and cesium, low chloride internal solutions,

respectively. Representative traces showed that POA expressing ChR2-

mCherry neurons exhibited robust firing in response to optical stimulation at

different frequencies (n=15 neurons from 5 mice). **(F**, **G)** Representative traces

showed that blue light stimulation evoked either excitatory postsynaptic current

(EPSC) which could be blocked by cyanquixaline (CNQX, 10 μM) or inhibitory

postsynaptic current (IPSC) which could be blocked by picrotoxin (PTX, 100

nM). **(H)** Pie chart showed the projection types recorded on pPVT neurons

(n=63 neurons from 10 mice). **(I)** The representative recorded pPVT neuron

was visualized by biocytin staining and was found being surrounded by POA

expressing ChR2-mCherry terminals. Magnification: x60, scale bar: 20 μm. **(J**,

K) The representative trace showed that the application of tetrodotoxin (TTX)

eliminated the oPSC held at -70 mV while the addition of 4-amino-pyridine (4-

AP, 1mM) recovered it and the quantification (n=24 neurons from 10 mice).

1136

Figure 3. Activity changes of POA recipient pPVT neurons throughout

chronic heat exposure. (A) Experimental schematics. Mice (n=5) were

stereotaxically injected with Cre-dependent GCaMP into the pPVT, followed by

the implantation of optical fiber and chronic calcium recording. **(B)**

Representative microphotographs showed the virus expression in the POA and

pPVT regions, scale bar: all 40 μm. **(C)** From the top to the bottom: the

representative calcium events (on the left panel) and the calcium events from

different mice (on the right panel) on different days. Each vertical stripe in the

heatmap represents one calcium event for each mouse. Statistical analysis of the frequency and amplitude of POA recipient pPVT neurons' calcium events compared for **(D)** Pre and During heat exposure on day 1 (Paired, parametric, two tailed t-test; Frequency (*t=6.278, df=4, P=0.0033*); Amplitude (*t=4.344, df=4, *p=0.0122*)); **(E)** Pre heat on day 1 and After heat on day 2 (Paired, parametric, two tailed t-test; Frequency (*t=0.2787, df=4, p=0.7943*); Amplitude (*t=1.726, df=4, p=0.1595*)); **(F)** Pre heat on day 1 and During heat exposure on day 21 (Paired, parametric, two tailed t-test; Frequency (*t=6.124, df=4, **p=0.0036*); Amplitude (*t=4.704, df=4, **p=0.0093*)); **(G)** Pre heat on day 1 and after Chronic heat on day 22 (Paired, parametric, two tailed t-test; Frequency (*t=6.216, df=4, **p=0.0034*); Amplitude (*t=1.36, df=4, p=0.2454*)); **(H)** During heat exposure on day 1 and day 21 (Paired, parametric, two tailed t-test; Frequency (*t=1.242, df=4, p=0.2821*); Amplitude (*t=0.9424, df=4, p=0.3993*)). * *p*<0.05, ** *p*<0.01, N.S.: not significant.

Figure 4. POA recipient pPVT neurons are sufficient and necessary for chronic heat exposure-induced negative emotional and hyperarousal states in mice. (A) Experimental schematics. Mice (n=10 in each group) were stereotaxically injected with either AAV1-hSyn-EGFP or AAV9-CaMKII-ChR2-mCherry into the POA, followed by the implantation of optical fiber into pPVT, chronic optogenetic activation, and behavioral tests. **(B, C)** The heatmap of representative tracking trace examples in EPM test and the time spent in the open arms (Mann-Whitney unpaired two-tailed U test; U=2, ***p*<0.001). **(D, E)** The heatmap of representative tracking trace examples in TCT and the

interaction time with an unfamiliar male mouse relative to the inanimate object (Mann-Whitney unpaired two-tailed U test, U=8; ***p=0.0007). **(F, G)** The heatmap of representative tracking trace examples in FET and the time surrounding the unfamiliar female mouse compared to an unfamiliar male mouse (Mann-Whitney unpaired two-tailed U test; U=13, **p=0.0039). **(H, I)** The 1st-time attack latency (Mann-Whitney unpaired two-tailed U test; U=8, p=0.007) and the attack durations (Mann-Whitney unpaired two-tailed U test; U=19, *p=0.0185) in the aggression test. **(J, K)** Visualized acoustic startle response examples (upper panel) and corresponding labeled body parts' skeletons (lower panel) from hSyn-EGFP group and ChR2-mCherry group. **(L, M)** The startle amplitude (Mann-Whitney unpaired two-tailed U test; U=17, *p=0.0115) and the pre-pulse / pulse ratio (Mann-Whitney unpaired two-tailed U test; U=48, p=0.9118) in ASR test. **(N)** Experimental schematics. Mice (n⩾6 in each group) were stereotaxically injected with AAV1-hSyn-Cre-EGFP in the POA and either AAV9-Ef1a-Dio-eNpHR3.0-EGFP or AAV8-hSyn-Dio-EGFP into the pPVT, followed by the implantation of optical fiber and behavioral tests. **(O)** Representative microphotographs showed the expression of AAV1-hSyn-Cre-EGFP and AAV9-hSyn-Dio-eNpHR3.0 within the POA and pPVT, respectively. Magnification: x4, scale bar: 100 µm (upper panel) and 250 µm (lower panel). **(P, Q)** The time (One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 27)}$=83.03, ***p<0.001; Control vs. Chronic heat, ***p<0.001; Chronic heat vs. Chronic heat + inhibition, ***p<0.001) and the entry numbers (One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 20.92)}$=27.96, ***p<0.001; Control vs. Chronic heat, ***p<0.001; Chronic heat vs. Chronic heat + inhibition, ***p<0.001) into the open arms of EPM. **(R)** The time spent with an

unfamiliar male mouse compared to an inanimate object in TCT (One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 27)}=5.381$, *$p=0.0108$; Control vs. Chronic heat, *$p=0.0292$; Chronic heat vs. Chronic heat + Inhibition, *$p=0.0175$). **(S)** The ratio of time with an unfamiliar female mouse compared to an unfamiliar male mouse in FET (One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 27)}=10.94$, ***$p=0.0003$; Control vs. Chronic Heat, ***$p=0.0002$; Chronic Heat vs. Chronic Heat + Inhibition, *$p=0.0335$). **(T, U)** The 1st attack latency (One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 17)}=7.426$, *$p=0.0048$; Control vs. Chronic Heat, *$p=0.0157$; Chronic Heat vs. Chronic Heat + Inhibition, *$p=0.0063$) and the attack durations (One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 17)}=13.38$, ***$p=0.0003$; Control vs. Chronic Heat, ***$p=0.0003$; Chronic Heat vs. Chronic Heat + Inhibition, **$p=0.0051$) in the aggression test. **(V, W)** The startle amplitude (One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 21)}=21.03$, ****$p<0.0001$; Control vs. Chronic Heat, ***$p<0.001$; Chronic Heat vs. Chronic Heat + Inhibition, ***$p<0.001$) and the pre-pulse / pulse ratio (One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 21)}=3.802$, *$p=0.039$; Control vs. Chronic Heat, $p=0.2616$; Chronic Heat vs. Chronic Heat + Inhibition, $p=0.2246$) in ASR test. *$p<0.05$, **$p<0.01$, ***$p<0.001$, N.S.: not significant.

Figure 5. Chronically activated POA recipient pPVT neurons exhibited exaggerated response to stressful situations. (A) Experimental schematics. Mice (n=5) were stereotaxically injected with AAV1-hSyn-Cre-EGFP in the POA

and AAV9-hSyn-Flex-jGCaMP8F-WPRE in the pPVT, followed by the

implantation of optical fiber and calcium recording. **(B)** The number of entries

into the open arms of EPM (Paired, parametric, two tailed t-test; t=6.901, df=4,

p*=0.0023). **(C) The number of instances where the mice paused at the center

area followed by back to closed arms of EPM (Paired, parametric, two tailed t-

test; t=9, df=4, ***p*=0.0008). **(D)** Pie chart showed the changes of the

percentage of pause and then running back to closed arms. **(E)** The changes

of running speed in EPM (Mann-Whitney unpaired two-tailed U test; U=139,

p*=0.0028). **(F-H) Heatmaps showed the calcium activities of POA recipient

pPVT neurons when mice performed the pause-and-run-back-to-closed-arms

behavior for pre (n=13 trials from 5 mice) and post conditions (n=47 trials from

5 mice), Z-Score calcium signals, and statistical comparison (Mann-Whitney

unpaired two-tailed U test; U=194, *p*=0.0455). **(I)** The time spent in open arms

of EPM (Paired, parametric, two tailed t-test; t=3.649, df=4, *p*=0.0218). **(J)** The

total number of the behavioral event: back from open to closed arms (Paired,

parametric, two tailed t-test; t=6.901, df=4, **p*=0.0023). **(K)** Pie chart showed

the changes of number of running episodes from open to closed arms for pre

and post conditions. **(L)** The changes of running speed in EPM (Mann-Whitney

unpaired two-tailed U test; U=23, ***p*=0.0005). **(M-O)** Heatmaps showed the

calcium activities of POA recipient pPVT neurons when mice performed fast

running from open to closed arms for pre (n=14 trials from 5 mice) and post

conditions (n=13 trials from 5 mice), Z-Score calcium signals, and statistical

comparison (Mann-Whitney unpaired two-tailed U test; U=42, *p*=0.0168). **(P)**

The total distances of mice travelled in the chamber previously subjected to

chronic heat (Paired, parametric, two tailed t-test; t=6.876, df=4, **p*=0.0023).

(Q) The changes of motion speed for pre and post conditions from one representative mouse. **(R)** The number of instances with motion speed exceeding 400 mm/s (Paired, parametric, two tailed t-test; t=5.100, df=4, ***p=0.007). **(S)** The number of fast running episodes for pre and post conditions. **(T-V)** Heatmaps showed the calcium activities of POA recipient pPVT neurons when the mice performed fast running in the previous chronic heat-exposed chamber (n=90 trials from 5 mice) compared to pre-heat condition (n=38 trials from 5 mice), Z-Score calcium signals, and statistical comparison (Mann-Whitney unpaired two-tailed U test; U=1218, *p=0.01). *p<0.05, **p<0.01, ***p<0.001.

Figure 6. Increased pre- and post-synaptic excitability of pPVT neurons but saturated circuitry neuroplasticity capacity following chronic heat exposure. (A) Experimental schematics. *In vitro* brain slice recording was performed on mice from three groups (n=16 neurons from 3 mice in the Room temperature group, n=20 neurons from 4 mice in the Acute heat group, and n=20 neurons from 4 mice in the Chronic heat group). **(B)** Representative traces of miniature postsynaptic currents from three groups. Duration: 12 s. Scale bar: 10 pA. **(C, D)** The changes of mIPSCs' amplitude (One-way repeated measures ANOVA with Tukey post-hoc test; F(2, 54)=8.226, ***p=0.0008, Room temperature vs. Acute heat, ***p=0.0005; Room temperature vs. Chronic heat, *p=0.0268) and frequency (One-way repeated measures ANOVA with Tukey post-hoc test; F(2, 53)=1.346, *p*=0.269, Room temperature vs. Acute heat, *p*=0.4327; Room temperature vs. Chronic heat, *p*=0.9787) of pPVT neurons.

(E, F) The changes of mEPSCs' amplitude (One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 52.98)}=0.1995$, $p=0.8198$, Room temperature vs. Acute heat, $p=0.8433$; Room temperature vs. Chronic heat, $p=0.8433$) and frequency (One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 50.1)}=2.981$, $p=0.0598$, Room temperature vs. Acute heat, *p=0.2931*; Room temperature vs. Chronic heat, **p=0.0473*) of pPVT neurons. **(G)** The representative traces of action potential of pPVT neurons upon 100 pA current injection. **(H)** The changes of excitability of pPVT neurons when different currents were injected to the patched pPVT neurons (n=11 neurons from 3 mice in each groups; Mann-Whitney unpaired two-tailed U test; 30 pA: U=22.5, ***p=0.0091*; 40 pA: U=31, **p=0.048*; 50 pA: U=29.5, **p=0.0412*; 60 pA: U=25, **p=0.0167*; 70 pA: U=29.5, **p=0.0396*). **(I)** The changes of rheobase of action potential (Mann-Whitney unpaired two-tailed U test; U=28.5, **p=0.0345*). **(J)** Experimental schematics. Mice (n=5 mice in each group) stereotaxically injected with AAV9-CaMKII-ChR2-mCherry into the POA were then divided into the Room temperature and Chronic heat groups. Sagittal slices of mice were prepared for LTP induction and recording. **(K)** The representative traces showed pPVT neurons from mice exposed to room temperature and chronic heat exhibited different amplitude of oEPSCs after blue light-mediated high frequency stimulation. **(L)** The LTP induction and recording of POA to pPVT pathway from slices of mice exposed to room temperature and chronic heat conditions after blue light stimulation at 30 Hz (n=14 neurons from control group and 18 neurons from chronic heat group). **(M)** Statistical comparison of the amplitude of oEPSCs at different time points (Two-way repeated measures ANOVA with Sidak post-hoc test; Interaction: $F_{(3, 120)}=0.3486$, *p=0.7902*;

Optical stimulation main effect: $F_{(3, 120)}=76.2$, ***$p<0.001$; Time points effect:

$F_{(3, 120)}=0.8215$, $p=0.4844$; 1st min: Room temperature vs. Chronic heat:

***$p<0.001$; 10th min: Room temperature vs. Chronic heat: $p=0.0001$; 20th min:

Room temperature vs. Chronic heat: ***$p<0.001$; 30th min: Room temperature

vs. Chronic heat: **$p=0.0024$). *$p<0.05$, **$p<0.01$, ***$p<0.001$, N.S.: not

significant.

1298

Figure 7. A working model of neural circuit mechanisms underlying POA

recipient pPVT neurons-mediated chronic heat exposure-induced

negative emotional and hyperarousal states. Different from acute heat

exposure, chronic heat exposure-induced enhancement in excitatory inputs to

pPVT and saturated neuroplasticity contributed to the increased membrane

excitability underlie the heightened activities of POA recipient pPVT neurons,

rendering mice become more susceptible to stressful situations manifested as

negative emotions and hyperarousal states.

1307