Repeated activation of preoptic area-recipient neurons in posterior paraventricular nucleus mediates chronic heat-induced negative emotional valence and hyperarousal states

Zhiping Cao[1], Wing-Ho Yung[2]*, Ya Ke[1]*

[1] School of Biomedical Sciences, Faculty of Medicine, The Chinese University of Hong Kong, Shatin, Hong Kong, China.

[2] Department of Neuroscience, City University of Hong Kong, Kowloon Tong, Hong Kong, China

* **Corresponding author**:

Wing-Ho Yung, PhD; Department of Neuroscience, City University of Hong Kong, Tat Chee Avenue, Kowloon, Kowloon, Hong Kong. Email: whyung@cityu.edu.hk

Ya Ke, PhD; Lo Kwee Seong Integrated Biomedical Sciences Building, Tai Po Road, Sha Tin District, New Territories, Hong Kong. Email: yake@cuhk.edu.hk

Abstract

Mental and behavioral disorders are associated with extended period of hot weather as found in heatwaves, but the underlying neural circuit mechanism is poorly known. The posterior paraventricular thalamus (pPVT) is a hub for emotional processing and receives inputs from the hypothalamic preoptic area (POA), the well-recognized thermoregulation center. The present study was designed to explore whether chronic heat exposure leads to aberrant activities in POA recipient pPVT neurons and subsequent changes in emotional states. By devising an air heating paradigm mimicking the condition of heatwaves and utilizing emotion-related behavioral tests, viral track tracing, *in vivo* calcium recordings, optogenetic manipulations and electrophysiological recordings, we found that chronic heat exposure for 3 weeks led to negative emotional valence and hyperarousal states in mice. The pPVT neurons receive monosynaptic excitatory and inhibitory innervations from the POA. These neurons exhibited a persistent increase in neural activity following chronic heat exposure, which was essential for chronic heat-induced emotional changes. Notably, these neurons were also prone to display stronger neuronal activities associated with anxiety responses to stressful situations. Furthermore, we observed saturated neuroplasticity in the POA-pPVT excitatory pathway after chronic heat exposure that occluded further potentiation. Taken together, long-term aberration in the POA to pPVT pathway offers a neurobiological mechanism of emotional and behavioral changes seen in extended periods of hot weather like heatwaves.

Introduction

Against a backdrop of global warming, heatwaves that are characterized by abnormally hot weather for extended periods have become more frequent and intense. This is highlighted by that the July 2023 was the hottest month ever recorded by human (Tollefson, 2023). Traditional views support that repetitive heat exposure and the related heat acclimation have some beneficial effects in allowing the animals and humans to gain heat tolerance, such as strengthening the cardiovascular system, reducing energy metabolism and weight (Murray KO, et al., 2022; Podstawski R, et al., 2021; Périard et al., 2016; Fausnacht DW, et al., 2021). However, many studies have also established a correlation between heatwaves and impaired physical health leading to increased mortality (Barriopedro et al., 2011; Garcia-Herrera et al., 2010; Mitchell et al., Mora et al., 2017; 2016; Robine et al., 2008). In addition, elevated temperatures could impact mental health by triggering feelings of anger, stress, aggression, and depression (Beecher et al., 2016). In fact, epidemiological research conducted across various regions also extensively demonstrated a positive association between chronic heat exposure and higher hospital admissions for mental and behavioral disorders (Amr et al., 2012; Basu et al., 2018; Dominiak et al., 2015; Hansen et al., 2008; Lee et al., 2002; Mulder et al., 1990; Trang et al., 2015; Whitney et al., 1999). Nevertheless, our understanding of how the brain regulates emotional changes following prolonged heat exposure remains limited.

Current understanding about heat-induced emotional changes mainly focuses

on the impact of heat stress on the hypothalamic-pituitary-adrenal (HPA) axis,

which leads to abnormal plasma concentrations of the hormone cortisol and

neurotransmitters such as 5-hydroxytryptamine (5-HT), noradrenaline and

adrenaline, all of which play a significant role in modulating emotional states

(McMorris et al., 2006; Wang et al., 2015). However, how heat affects

emotions from the perspective of neural circuits has not been substantiated.

Moreover, the relationship between chronic heat exposure and emotional

changes is likely to be more complex. Multifaceted factors, such as the levels

of physiological stress (Sampath et al., 2023), changes of neuroendocrine

system (Podstawski et al., 2021), sleep disturbances (Altena et al., 2022),

adaptation process (Oppermann et al., 2021), even the age and individual

health status (Kenny et al., 2010; Malmquist et al., 2022), could interact in a

various way to directly and indirectly contribute to emotional changes in

response to chronic heat exposure. Thus, gaining insights into the circuitry

mechanisms behind emotional changes caused by chronic heat exposure is

crucial for a more comprehensive understanding of the neurobiological

foundations of emotional regulation.

Emerging evidence has suggested that the paraventricular thalamus (PVT)

serves as an important integrative node that detects aversive sensory and

homeostatic challenges and regulates emotional responses and adaptive

behaviors. As such, PVT neurons receive inputs from diverse nuclei in the

hypothalamus, midbrain, and hindbrain, and has been implicated in a variety

of behavioral responses including arousal (Ren et al., 2018), pain (Jurik et al.,

2015), anxiety (Heilbronner et al., 2004; Li et al., 2010; Spencer et al., 2004),

fear and fear memory (Do-Monte et al., 2015; Penzo et al., 2021). Therefore, PVT is now considered as an essential component of the emotional processing system in the brain (Penzo et al., 2021).

Among the various inputs from different brain regions to the PVT, the hypothalamus preoptic area (POA) is a well-known thermoregulatory center of the brain (Yu et al., 2016; Zhang et al., 2011; Zhao et al., 2016). Anatomically, the posterior PVT (pPVT) has been reported to receive projections from POA warm-sensitive neurons (Tan et al., 2016). Here, we hypothesize that the direct projection from POA to pPVT contributes to chronic heat-induced emotional disturbances. Based on the murine model, we dissected the synaptic connection from POA to pPVT, tracked its activities and interrogated its involvement in behavioral responses of the animals under acute and chronic heat exposure paradigms. Our results uncovered long-term aberration in the activity of this pathway and pPVT neurons mediating chronic heat-induced changes in emotional and hyperarousal states, providing a neurobiological basis of how chronic heat affects mental states.

Results

Chronic heat produces negative emotional valence and hyperarousal states but not depression-like behaviors in mice.

c57BL/6 mice were exposed to chronic heat by putting them into a pre-heated (38 ± 2℃) chamber daily for 21 days with free access of food and water (details in **Materials and Methods**). In our preliminary experiments, various

durations of daily heat exposure were tested. We settled to 90 minutes due to its robustness in inducing emotional state-related behavioral changes without causing the collapse of the animals. Throughout the entire process of chronic heat exposure, we monitored the physiological states of mice daily. We did not observe abnormal changes in body temperature the day after chronic heat exposure (Figure 1-figure supplement 1A and B) but affected normal gain in body weight (Figure 1-figure supplement 1C and D) and reduced food consumption (Figure 1-figure supplement 1E and F), compared with control mice. On the other hand, mice subjected to chronic heat displayed a higher prevalence of stress responses in various behavioral tests. Specifically, in the elevated plus maze test, chronic heat-exposed mice spent less time in the open arms (Figure 1B and C), indicating increased stress levels. Moreover, in the three-chamber test, these mice displayed reduced interests in exploring the unfamiliar male mouse compared to the inanimate object (Figure 1D and E), suggesting decreased sociability. Similarly, during the female encounter test, these mice performed less engagement with the unfamiliar female mouse compared with the unfamiliar male mouse (Figure 1F and G), indicating decreased innate motivation. Interestingly, the chronic heat-exposed mice displayed decreased latencies for the first attack but increased attack durations in the resident-intruder test (Figure 1H and I), indicating elevated aggression levels.

We also observed that after chronic heat exposure, the mice displayed heightened alertness, suggesting a state of hyperarousal. To confirm this, we quantified the acoustic startle response of the animals by digital video capture of their rapid movements (Pantoni et al., 2020). Chronic heat-exposed mice

displayed amplified acoustic startle response characterized by increased body

fluctuations within 200 ms of the delivery of a 105 dB sound stimulus (Figure

1J-M and Figure 1-video supplement 1). However, these mice did not exhibit

clear hyper-locomotion during the three-chamber test, the female encounter

test, or the open field test (Figure 1-figure supplement 1G-I), suggesting

increased agitation in mice. Collectively, our data demonstrate that chronic

heat exposure triggers increased stress responses as well as increased

arousal in mice.

The fact that the observed changes in emotional and arousal states are the

results of long-term rather than short-term heat exposure is supported by the

lack of effect measured one day following one-time acute heat exposure for

90 mins (Figure 1-figure supplement 2). Furthermore, we did not observe

signs of depression in chronic heat-exposed mice, as there were no

significant increases in immobile time during forced swimming (Figure 1N)

and the tail suspension tests (Figure 1O), no obvious decrease in sucrose

intake (Figure 1P), and no sign of decreased exploration as there was no

change in the time spent by these animals in the center zone of the open field

(Figure 1Q).

Involvement of the hypothalamic preoptic area to posterior

paraventricular thalamus projection.

To explore the involvement of POA and pPVT in chronic heat exposure

induced behavioral changes, we first studied the effect of heat treatment *per*

se on their activities via c-Fos staining. We found that acute heat treatment

(for 90 mins) significantly increased c-Fos expressions of POA and pPVT

neurons (Figure 2-figure supplement 1A-C), supporting not only the notion

that the POA serves as a thermoregulation center (Morrison et al., 2011) but

also the recruitment of pPVT following heat treatment. Next, to dissect the

anatomical relationship between POA and pPVT especially in the context of

heat exposure, we injected viruses (AAV1-hSyn-cre, AAV-FLEX-TVA-

mCherry, AAV-FLEX-RG and subsequently SAD△G (EnvA) virus) of a

monosynaptic retrograde tracing rabies virus system (details in **Materials and**

Methods) into pPVT, which resulted in clear expression of retrogradely-

labeled neurons within POA, its upstream area (Figure 2A-C). Moreover, a

significant overlap (84 ± 5%, 3 mice) was observed between the retrogradely-

labeled and c-Fos expressed neurons in POA following heat exposure (Figure

2D). Furthermore, almost all the POA recipient pPVT neurons identified by a

Cre-dependent anterograde labelling strategy (93 ± 2%, 3 mice) were strongly

activated by heat exposure (Figure 2-figure supplement 2). Together, our

results indicate that most of the pPVT-projecting POA neurons were neurons

that respond to heat treatment which would then recruit their downstream

neurons in pPVT by exerting a net excitatory influence.

To functionally characterize the synaptic connection between POA and pPVT,

we administered AAV9-hSyn-ChR2-mCherry into the POA for optogenetic

manipulation of its neurons and their terminals in pPVT. By patch-clamp

recordings in brain slices prepared from these animals, we first confirmed

robust firing of ChR2-mCherry-expressing neurons in POA upon light

stimulation at various frequencies (Figure 2E). When we applied 473 nm light

stimulation to POA terminals in pPVT, in some neurons, a clear inhibitory outward postsynaptic current was recorded in pPVT neurons held at +10 mV, which was sensitive to picrotoxin (Figure 2F). In some other neurons, an inward excitatory postsynaptic current sensitive to CNQX was recorded when the neurons were held at -70 mV (Figure 2G). Quantification of the presence of these light-evoked postsynaptic currents revealed the presence of both excitatory and inhibitory currents in 34.9% of neurons recorded, only inhibitory currents in 52.4% of neurons and only excitatory currents in 12.7% (Figure 2H). Since these currents could be completely blocked by tetrodotoxin (TTX) but restored with the addition of 4-aminopyridine (4-AP), it indicated the monosynaptic connection, (an example shown in Figure 2I-K). Collectively, our findings demonstrate that heat responsive POA neurons project directly to pPVT via both excitatory and inhibitory innervations.

Activity changes of POA recipient pPVT neurons throughout chronic heat exposure.

We speculated that repeated activation of POA recipient pPVT neurons underline the emotional valence changes observed following chronic heat exposure. To gain support for this hypothesis, we first monitored the activity changes of these neurons during the course of chronic heat exposure by *in vivo* fibre photometry of calcium signals (Figure 3A and B). We injected AAV1-hSyn-Cre-EGFP into POA and Cre-dependent AAV9-hSyn-Flex-jGCaMP8F into pPVT and implanted optical fibers into pPVT neurons after enough expression of GCaMP. The fibre photometry recordings revealed fluctuations

in signals composed of spontaneous events (details in **Materials and Methods**) superimposed on the baseline. The detected spontaneous calcium events presumably represent brief trains or bursts of neuronal firing (Ali et al., 2020). We first confirmed that the signals were stable for extended period (at least 21 days) in a group of mice not undergoing any treatment (Figure 3-figure supplement 1). As illustrated in Figure 3C, prior to subjecting the mice to chronic heat exposure, we recorded the neuronal activities of POA recipient pPVT neurons for each mouse as the pre-heat control. We then recorded the calcium events one day after acute heat exposure on day 1 and after chronic heat exposure on day 21. Interestingly, when compared with the pre-heat condition of day 1, both the frequency and amplitude during heat exposure on day 1 and day 21 were significantly increased (Figure 3D and F). However, such increases in calcium events disappeared on the day following heat exposure on day 1 indicating its transient nature (Figure 3E). In contrast, after 21 days of heat exposure, such increases in the frequency of calcium events persisted (Figure 3G). It is worth noting that there were no significant differences in both the frequency and amplitude of calcium events during heat exposure on day 1 and day 21 (Figure 3H).

Based on the increased neuronal activities of POA recipient pPVT neurons after chronic heat exposure, we investigated the impact of activating the pathway from the POA to pPVT on emotional valence and arousal levels of mice (Figure 3-supplement figure 3A). Utilizing the real-time place preference behavioral paradigm, we noticed that when the blue light was turned on, mice that had already gone through the habituation process quickly entered the chamber with the light turned off (Figure 3-supplement figure 2B and C). This

finding suggests that the activation of the POA to pPVT circuit induces an

aversive emotional valence. Notably, optogenetic activation did not affect the

movement activity of the mice during the experiment (Figure 3-supplement

figure 2D). And we did not observe any change in core body temperature

when stimulating the POA-pPVT circuit. On the other hand, as pupil size is

widely recognized as the gold standard for reflecting arousal levels in rodent

models (Privitera et al., 2020), we further studied the effect of optogenetic

activation of POA excitatory terminals in pPVT in mice under head fixation.

The photos (Figure 3-supplement figure 2E) and video (Figure 3-video

supplement 1) clearly demonstrated the enlargement of both the pupil (Figure

3-supplement figure 2F and G) and eye sizes (Figure 3-supplement figure 2H

and I) of the mice during light stimulation. Collectively, our results

demonstrate that POA recipient pPVT neurons exhibited heightened activities

after chronic heat exposure. Moreover, the activation of the POA to pPVT

circuit is associated with negative emotional states and higher levels of

arousal.

POA recipient pPVT neurons are sufficient and necessary for chronic

heat exposure-induced negative emotional valence and hyperarousal

states in mice.

To confirm the direct relationship between repeated activation of POA

recipient pPVT neurons and the negative emotional and hyperarousal states

in mice, we conducted chronic activation of POA excitatory terminals within

pPVT by injecting CaMKII-promotor encoded ChR2 into the POA and

implanting an optical fiber in pPVT. We applied a chronic activation protocol,

described by Sidor et al. (Sidor et al., 2014), which involved a cycle of 2

minutes on and 2 minutes off, for a total of 20 minutes per day, over a period

of up to 21 days. Following chronic optogenetic activation, we observed

several behavioral changes in mice that were similar to those seen in mice

exposed to chronic heat. Notably, in the elevated plus maze test, mice

subjected to chronic optogenetic activation spent less time in the open arms

(Figure 4B and C), representing increased stress levels. In the three-chamber

test, these mice displayed reduced interests in exploring the unfamiliar male

mouse compared to an inanimate object (Figure 4D and E), suggesting

decreased sociability. During the female encounter test, these mice engaged

less with the unfamiliar female mouse compared with the unfamiliar male

mouse (Figure 4F and G), indicating decreased innate motivation. In the

resident-intruder aggression test, the chronic heat-exposed mice exhibited

both the decreased latencies in initiating the first attack and increased attack

durations (Figure 4H and I), suggesting elevated aggression levels.

Additionally, compared to the control group, mice after chronic optogenetic

activation exhibited more pronounced body fluctuations when captured by a

high-resolution camera in response to a 105 dB sound stimulus, indicating the

heightened arousal levels (Figure 4J-M). Our results directly demonstrated

that chronic optogenetic activation of POA excitatory terminals within pPVT

circuit could induce negative emotional and hyperarousal states in mice.

To establish the necessity of POA recipient pPVT neurons for chronic heat

exposure-induced emotional changes, we optogenetically inhibited the POA

recipient pPVT neurons during chronic heat exposure by employing a cycle

stimulation strategy of 3 minutes ON followed by 3 minutes OFF. Only mice with sufficient virus infection and accurate fiber implantation were included in the data analysis (Figure 4O and P). The optogenetic inhibition prevented chronic heat-induced behavioral changes, including anxiety levels (Figure 4Q and R). sociability (Figure 4S), innate motivation (Figure 4T), aggression levels (Figure 4U and V) and arousal state (Figure 4W and X). These findings further supported the essential role of POA recipient pPVT neurons in mediating chronic heat exposure-induced negative emotional and hyperarousal states.

Chronically activated POA recipient pPVT neurons exhibit exaggerated response to stressful situations.

To gain further insight into the roles of POA recipient pPVT neurons in stress-related behaviors, we elucidated the relationship of their activities with specific behaviors indicative of emotional transitions and hyperarousal states. In the elevated plus maze test in which the chronic heat-exposed mice spent less time in the open arms (Figure 1B and C), there were also reduced number of entries into the open arms (Figure 5B). More detailed analysis revealed that, compared with untreated mice, these animals more frequently paused at the center of the maze and then returned to the closed arms (Figure 5C). Furthermore, mice after chronic heat exposure exhibited a higher chance of running, rather than walking, back to the closed arms (Figure 5D) and their running speed significantly increased compared to their pre-heat condition (Figure 5E), presumably reflecting a state of increased anxiety. Interestingly,

POA recipient pPVT neurons consistently exhibited a peak in activity when they were about to run back to the closed arm, while chronic heat-exposed mice exhibited a significantly larger peak (Figure 5F-H). Notably, no comparable increases of activities in the POA recipient pPVT neurons were observed when chronic heat-exposed mice paused at the center and finally walked out to the open arms (Figure 5-supplement figure 1A-C) or walked to the closed arms (Figure 5-supplement figure 1D-F). Similar phenomena were found with respect to the transitions from the open arm to the closed arm, not only in terms of number of transitions (Figure 5I and J), walking and running behavior (Figure 5K), and speed of running (Figure 5L), but also a larger calcium peak response aligned with the running behavior (Figure 5M-O).

In the chronic heat-treated mice, we also observed a context-dependent change in behavior reminiscent of the hyperarousal states when we introduced them back into the heat chamber. The mice exhibited apparent locomotion hyperactivity (Figure 5P), as evidenced by a notable increase in motion speeds exceeding 400 mm/s (Figure 5Q) and a higher frequency of fast running (Figure 5S). A clear time correlation was found between the heightened calcium activities in POA recipient pPVT neurons and the main component of locomotion hyperactivity, namely fast running under this condition (Figure 5T-V). In contrast, when being placed into a different chamber not associated with heat exposure, no obvious changes were observed among these chronic heat-exposed mice in their total distance traveled (Figure 5-supplement figure 1G), motion speed (Figure 5-supplement figure 1H), number of running exceeding 400 mm/s (Figure 5-supplement figure 1I), the frequency of fast running (Figure 5-supplement figure 1J), or the

neuronal activities (Figure 5-supplement figure 1K-M). Collectively, our data

suggest the heightened or exaggerated activities of POA recipient pPVT

neurons after chronic heat exposure could serve as a driving force for

behaviors manifesting negative emotional and hyperarousal states in

response to stressful conditions.

Increased pre- and post-synaptic excitability of pPVT neurons but

saturated circuitry neuroplasticity capacity following chronic heat

exposure.

To explore the potential mechanisms underlying the heightened activities of

POA recipient pPVT neurons after chronic heat exposure, we conducted *in*

vitro slice recordings on pPVT neurons obtained from mice exposed to room

temperature, acute heat, and chronic heat, respectively (Figure 6A and B).

Both acute and chronic heat exposure significantly increased the amplitude of

miniature inhibitory postsynaptic currents (mIPSCs) in pPVT neurons but

there was no difference in mIPSCs amplitude between the acute and chronic

heat groups (Figure 6C). On the other hand, we observed a significant

increase in the frequency of miniature excitatory postsynaptic currents

(mEPSCs) specially in pPVT neurons after chronic but not acute heat

exposure (Figure 6F). These results suggest that while heat exposure could

modulate both inhibitory and excitatory synaptic inputs onto pPVT neurons,

there was a differential increase in presynaptic excitability of the excitatory

pathway after chronic heat exposure but an increase in postsynaptic response

to inhibitory input. Furthermore, when we examined the excitability of pPVT

neurons by injecting a sequence of inward currents up to 100 pA to pPVT

neurons, pPVT neurons from chronic heat-exposed mice exhibited significant

increases in the number of action potentials from 30 to 70 pA (Figure 6G and

H). Further analysis unveiled that these neurons exhibited a reduced

rheobase for action potential generation (Figure 6I). However, no apparent

alterations were observed in the other parameters (Figure 6-supplement

figure 1).

The increase in presynaptic excitability of the POA to pPVT excitatory

pathway suggested plastic changes induced by the chronic heat treatment.

While it would be difficult to follow the excitability of this pathway in vivo during

the chronic heat treatment, we sought to examine the synaptic plasticity

capacity of this pathway before and after chronic heat treatment to shed light

on its involvement. We injected CaMKII-promotor encoded ChR2 into the

POA of mice and divided them into two groups: a control group exposed to

room temperature and a chronic heat group. After 21 days of heat exposure,

sagittal slices that largely preserved the POA to pPVT pathway were prepared

from both the control and chronic heat groups to induce long-term potentiation

(LTP) (Figure 6J). Local optogenetic stimulation vertically (470 nm) through

the objective was applied to elicit light-evoked EPSCs in POA recipient pPVT

neurons and baseline of 10 minutes was recorded. High frequency stimulation

(HFS) is an effective induction protocol for eliciting LTP at excitatory synapses

(Zhu et al., 2016; Grover et al., 2009; Hwong et al., 2008). Therefore, by

delivering blue light at various frequencies (10, 30, 50, 70, 90 Hz) to activate

the POA excitatory terminals within pPVT, we developed an optogenetic HFS

protocol (HFS$_{opto}$: light pulses at 30 Hz, repeated 3 times at 20 s interval)

effective in eliciting potentiated EPSCs on POA recipient pPVT neurons after

baseline recording. HFS$_{opto}$ protocol could successfully induce LTP at POA-

pPVT synapses. The amplitude of light-evoked EPSCs in the control group

following HFS$_{opto}$, at the onset, showed an obvious increase and gradually

became stable potentiation, lasting at least 30 min (150 % ± 18 of baseline,

14 neurons from 5 mice) (Figure 6K and L). However, when we applied the

same protocol to the POA recipient pPVT neurons of mice after chronic heat

exposure, typically no LTP could be induced (Figure 6K-M), suggesting that

the pathway in these mice were already saturated during chronic heat

exposure.

Taken together, our findings suggest enhanced excitatory inputs to pPVT

neurons and increased membrane excitability of these neurons may underlie

the behavioral changes observed in mice following chronic heat exposure.

Discussion

In this study, we aimed to uncover the neurobiological basis of how extended

periods of high temperature such as heatwaves impact on emotional

processing in the mammalian brain. We demonstrated that chronic heat

exposure led to negative emotional valence and hyperarousal states in mice,

but not depression-like behavior. We discovered that hypothalamic POA

recipient pPVT neurons in the thalamus mediate such emotional state

changes via increased baseline activities and excitability in response to

stressful conditions. Presumably, this is attributed to that repeated activation

of the POA to pPVT circuit modulates both pre- and post-synaptic processes

leading to their long-term changes.

Different from utilizing heating panels (Venkatachalam et al., 2007) to

investigate heat-induced thermoregulation, our study employed an air heating

strategy. This choice was motivated by the aim of replicating the realistic

conditions induced by heatwaves. To ensure the well-being of the animals

and prevent unnecessary physiological harm, we maintained temperatures

around 38°C, as temperatures exceeding 40°C can lead to adverse effects on

the body (Deuis et al., 2017). It is noted that our study differs from many

previous studies that applied less high temperature but continuously over

several days and weeks, which could have beneficial effects related to

cardiovascular fitness, energy metabolism and even cognitive functions

(Murray et al., 2022). Our findings revealed that the protocol of chronic heat

exposure at 90 min per day for 21 days did not impact the thermoregulatory

function in mice. In the various emotional state-related behavioral tests we

conducted, including the open field, three-chamber, and female encounter test,

we did not observe locomotion hyperactivity in mice. However, a distinct

acoustic startle response was noted following chronic heat exposure. This

may initially appear contradictory, as hyperarousal states are typically

associated with hyperlocomotion (Zhang et al., 2020). However, POA

recipient pPVT neurons exhibited heightened responses to stressful and

aversive situations, which suggested that chronic heat exposure might

primarily induce a state of increased susceptibility towards anxiety and

hyperarousal rather than persistent hyperarousal states in mice.

There are a number of studies revealing the connections from the POA to pPVT. Some previous reports have already shown that both excitatory and inhibitory neurons within POA contribute to thermoregulatory function (Morrison et al., 2011). Augustine et al. on the other hand demonstrated that POA neurons expressing nNOS send both excitatory and inhibitory projections to PVT, with an emphasis on their roles in thirst regulation (Augustine et al., 2018). In contrast, Allen et al. (Allen et al., 2017) previously described those thirst-related POA neurons sending projections to PVT mediated reinforcing behavior, highlighting their independence from POA heat-responsive neurons. The findings of the present study enriched our understanding that heat-responsive neurons in the POA could send both excitatory and inhibitory projections to pPVT, and uncovered the role of excitatory projections from POA to pPVT in chronic heat exposure-induced emotional changes. While the role of the excitatory inputs in mediating the negative emotional valence and hyperarousal state is unveiled in this study, the role of the inhibitory inputs awaits further investigation.

Notably, we observed that POA recipient pPVT neurons exhibited higher baseline activities and displayed heightened response to stress or aversive stimuli following chronic heat exposure. Consistently, a recent study also found that long-term heat exposure rendered POA neurons to become tonically active (Ambroziak et al., 2024). Interestingly, prior studies have reported that PVT neurons exhibited stronger activation when mice were subjected to novel stimuli after repeated restraint stress (Vertes et al., 2015; Bhatnagar et al., 1998; Heydendael et al., 2011). This also effectively supports our theory of heightened susceptibility to anxiety and our results

provide a rationale for this widely observed phenomenon. Meanwhile, the

specific neuronal signals correlated with actions indicative of heightened

anxiety such as fast running to closed arms and fast running in chronic-heat

exposed chamber were not induced by the motion *per se*. Our recording was

conducted when mice were spontaneously behaving without any pre-set cues,

excluding the influence of salient stimuli on the activity of POA recipient pPVT

neurons (Penzo et al., 2021).

In addition to the enhanced excitatory connections from POA to pPVT after

chronic heat exposure, we observed an increased amplitude of inhibitory

postsynaptic response after both acute and chronic heat exposure. Increased

inhibition could act to compensate for the initial over-excitation at critical time,

to ensure that neurons briefly activated by stress can be effectively and

prevent the occurrence of excessive stress-related behaviors (Perica et al.,

2023). pPVT neurons are anatomically innervated by strong inhibitory

projections from different brian nuclei (Penzo et al., 2021). Acute stress-

mediated disinhibition of pPVT neurons, as reported by Beas et al. (Beas et

al., 2018), can rapidly induce an anxiety-like state in mice. Other studies also

showed that inhibition of activated neurons induced by stressful stimuli is

effective to avoid abnormal affective behavioral performances (Cruz et al.,

2024; Ghasemi et al., 2022). Our results showed that long-term optogenetic

inhibition of POA recipient pPVT neurons during chronic heat exposure can

effectively prevent changes of emotional and hyperarousal states in mice.

Therefore, the increase in mIPSC amplitude observed under both acute and

chronic heat conditions in this study suggests the potential presence of

negative feedback regulation in response to the over-excitation. However,

under chronic heat exposure, further persistent significant increase in

excitatory synaptic transmission may surpass this effect, leading to a final net

activation in the pPVT neurons. However, the potential mechanisms by which

increased excitatory synaptic transmission and enhanced postsynaptic

inhibitory responses interact to achieve activation or inhibition of neurons

themselves still require further study.

Current hypothesis regarding the mechanism of pPVT in mediating anxiety-

like behavior primarily focused on the dopamine-mediated disinhibition of

hypothalamic inhibitory terminals within pPVT during acute stress (Beas et al.,

2018). Our study distinctly revealed that the enhanced excitatory inputs to

pPVT and increased excitability following chronic heat exposure are

responsible for the ensuing negative emotional and hyperarousal states.

Interestingly, different from previous studies that reported a decrease in the

amplitude of mIPSC in pPVT-expressing dopamine receptor 2 neurons 24

hours after acute restraint stress (Beas et al., 2018), our slice recording from

pPVT neurons 24 hours after acute heat exposure exhibited an increase in

mIPSCs amplitude. The reasons behind these differences could be attributed

to the treatment applied and the specific neuronal types under investigation.

Both the enhanced presynaptic excitability and increased intrinsic

postsynaptic excitability following chronic heat exposure suggested the

involvement of long-term neuroplasticity. The fact that LTP of the POA to

pPVT pathway was no longer inducible after chronic heat exposure strongly

suggested that this pathway was potentiated during the heat treatment, which

accounted for the increased excitability of POA terminals to pPVT and their

increased neuronal activities. Overall, both the enhanced excitatory inputs to

pPVT and increased excitability may play an indispensable role in chronic heat exposure-induced emotional changes in mice.

There are certain limitations in the scope of the present study. First, our study was conducted solely on male mice. Further investigation is required to determine whether chronic heat exposure can elicit similar behavioral phenotypes in female mice. Also, the pPVT is known to express various peptide receptors (Curtis et al., 2021). However, the specific roles of these peptide receptors in chronic heat exposure-induced emotional changes remain unexplored. Some possible mechanisms may contribute to heightened activities of POA recipient pPVT neurons, such as the increased sensitization, the increased activity of hypothalamus-pituitary-adrenal axis, the transition from chronic adaptation to pathological condition (Herman, 2013), and even the chronic effect of dopamine-induced disinhibition of inhibitory terminals within pPVT, all of which required further investigations. Moreover, POA and pPVT were both previously reported to be activated by stimuli other than heat (Penzo et al., 2021; Zhang et al., 2021). Additional studies are needed to explore the specificity of POA recipient pPVT neurons on chronic heat-induced emotional changes.

In conclusion, our findings provide compelling evidence that the heightened activities of POA recipient pPVT neurons are critically involved in chronic heat-exposure-mediated negative emotional valence and hyperarousal states, and render mice to enter anxiety-like state more readily, as summarized in Figure 7. These results shed a new light on the mechanisms underlying heatwaves-induced emotional changes.

Materials and Methods

Animals

Adult c57BL/6 male mice (25-45g) were used in this study. Animals were bred and maintained by the Laboratory Animal Service Centre of The Chinese University of Hong Kong (CUHK). All experiments were performed following the CUHK guideline approved by the Animal Experimentations and Ethics Committee.

Stereotaxic surgery and viruses

Mice were anesthetized with ketamine and xylazine and placed gently in a stereotaxic frame (Narashige, Tokyo). A Hamilton syringe (33-gauge) filled with AAV viruses was placed into target brain areas according to the corresponding coordinates: POA (+0.62 mm A/P, 0 mm M/L, 5.83 mm D/V), pPVT (-1.47 mm A/P, 0 mm M/L, 3.13 mm D/V) from brain skull. 0.10-0.35 µl viruses were injected at 10 nl/min speed. After injection, the needle was left in the targeted brain area for an additional 10 minutes before retraction. The following viruses were used: AAV1-hSyn-Cre (Addgene, retrograde tracing), AAV1-hSyn-Cre-EGFP (Addgene, optogenetics, photometry recording), Rabies virus (BrainVTA, retrograde tracing), AAV9-hSyn-ChR2-EGFP (Addgene, functional connectivity), AAV9-hSyn-Flex-jGCaMP8F-WPRE (Addgene, photometry recording), AAV9-CaMKII-ChR2-mCherry (Addgene, optogenetic activation), AAV9-hSyn-Dio-eNpHR3.0-EGFP (Addgene,

optogenetic inhibition), AAV9-hSyn-Dio-EGFP (Addgene, control experiments).

All AAV virus titers were > 5 x 10^{12} particles per ml.

Protocols for chronic heat exposure

The temperature (38 ± 2□) was adopted based on other reports (Bridges et

al., 2012; Cassuto, 1968; Cure, 1989; Jin et a., 2011; Rousset et al., 1984). A

home-made chamber was constructed for chronic heat exposure (80 cm [L] ×

40 cm [W] × 60 cm [H]) which was composed of two built-in components

(heating system and an environmental monitoring sensor). For the heating

system, a heater with (1) adjustable buttons that could control the heating

power and (2) a "dual heating" setting that was able to stop heating when the

temperature exceeded 40□ in the chamber and restarted heating when the

temperature in the chamber decreased to lower than 36□. After the air

temperature of the chamber was gradually heated to the targeted temperature

ranges and was monitored to be stable for at least 30 minutes, mice would be

gently put into the chamber for 90 minutes of heat exposure per day for 21

days. During heat exposure, mice were freely accessible to drinking water and

consuming food. The time point within each day to start heat exposure was

random. For the environmental monitoring sensor, it was used to monitor the

chamber temperature value and humidity level (Peng He Dian Zi, Shanghai,

China). And the behavioral tests were conducted the following day after 21

days' heat exposure.

The method for measuring the body temperature of mice was based on a

previous report (Kawakami et al., 2018). Briefly, prior to exposing them to heat,

the mice were gently held by their tails to expose their lower abdomen. An

infrared thermometer sensor was positioned beneath the lower abdomen and

the temperature was measured. This measurement process was repeated

three times for each mouse, and the stable values obtained were then

averaged to determine the specific mouse's body temperature.

Behavioral experiments

A variety of emotion-related behavioral tests were conducted on mice after

acute and chronic heat exposure.

Elevated plus maze test. The self-made elevated plus maze (EPM) apparatus

was made up of two open arms (25 cm [L] $\times$ 5 cm [W] $\times$ 0.5 cm [H]) and two

closed arms (25 cm [L] $\times$ 5 cm [W] $\times$ 16 cm [H]) arranged in a "plus" shape

and elevated 50 cm above the floor. The mice were gently placed at the

junction of the open and closed arms while facing toward one of the open

arms to initiate the test. Mice were allowed for a 6-min exploration, and their

behavior was videotaped and then analyzed and quantified by the ANY-Maze

tracking software (Version 4.7, Stoelting CO).

Three-chamber test & Female encounter test. The chambers for both the

three-chamber test and female encounter test were self-made by using an

open-field arena (50 cm [L] x 40 cm [W] x 30 cm [H]) with two identical

transparent chambers (40 cm [L] x 1 cm [W] x 20 cm [H]) with holes in the

middle surface, the stranger male and female mouse were handled for 3 min

and then habituated in a wire cage placed in the 3-chamber apparatus for 5-

10 min for four consecutive days. The tested mouse was habituated to the 3-

chamber apparatus for 10 min with all chambers unobstructed when the

handling was over. The tested mouse was then guided to the center chamber

when the habituation was over, and bidirectional exits were blocked. A wired

cup (of size 0.5 cm [L] x 0.5 cm [W], which prevents large body contact and

mating behavior but only induces innate sex-motivated exploratory movement

around the cup) with a stranger mouse and an empty cup was introduced into

the other two chambers, respectively, and then all chambers were opened for

a sociability test. For the female encounter test, a wired cup with a strange

male mouse and a wired cup with a strange female mouse was introduced

into the left and right chambers, respectively, and all chambers were then

opened for a motivation test. Their behavior was videotaped and then

analyzed and quantified by the ANY-Maze tracking software (Version 4.7,

Stoelting CO).

Resident-intruder aggression test. Mice were fed in a single cage, in which a

female mouse, separated by a wire mesh screen, was put in the corner of

each cage for seven days to facilitate each tested mouse to build their territory.

After chronic heat exposure, the unfamiliar intruder mouse with a color label

on the back was introduced to the cage of each tested mouse, and the video

was recorded for 15 minutes. Each tested mouse's first attack latency and

total attack durations were calculated and counted manually and double-blind

by another lab mate.

Acoustic startle response test. The acoustic startle response (ASR) test was

conducted based on a recent study (Pantoni et al., 2020). In our study, a

sound-proof box was purchased, in which a trumpet, a high-speed camera (200 frames per second, Plexon), and a box made from acrylic plate (10 cm [L] x 5 cm [W] x 25 cm [H]) were prepared. Mice exposed to sound at 105 dB measured by a decibel meter exhibited obvious startle response while 75 dB evoked no response. 105 dB was thus set as the sound decibel of pulse-only stimulation with 0.2 s duration, while 75 dB was set as pre-pulse stimulation with 0.2 s duration. Before the ASR test, mice were placed into an acrylic box for three days of environmental habituation and then for the test. Pre-pulse inhibition (PPI) test started with at least 20 min baseline to make the tested mice quiet and immobile, and then the habituation phase, which consisted of 10 pulses, each 20 s apart, and finally presented with a PPI phase, in which 20 trials: 5 pulse-only trials and 15 pre-pulse & pulse trials were delivered. In the pre-pulse & pulse trials, the pre-pulse was presented about 500 ms before the pulse stimulus. The mice's startle response was captured by a high-speed camera and then analyzed with the machine learning method: DeepLabCut. The brief details were as follows: the mice's body parts were manually labeled and labeled data were input into the google co-lab for the machine learning process until the learning error was lower than 0.5. The successful training data were exported as a .csv file. The startle amplitude of mice was calculated as the average value of all the labeled body parts' vibration amplitude within 200 ms after the sound was broadcast. Each trial was manually checked, and startle response affected by the mice's random moving would not be collected for further analysis. The startle response was visualized with custom MATLAB scripts.

The formula for calculating the PPI ratio was:

PPI ratio = amplitude of pre-pulse & pulse trial / amplitude of pulse-only trial.

Open field test (OFT). Mice were gently placed in the center of the OFT

chamber (60 cm [L] $\times$ 60 cm [W] $\times$ 40 cm [H]) and allowed for free exploration

for 10 mins. Mice's behaviors were analyzed and quantified by the ANY-Maze

tracking software (Version 4.7, Stoelting CO).

Forced swim test (FST). The apparatus of the forced swim test was a

cylindrical tank that was 30 cm high and 20 cm in diameter. And the water

was 15 cm tall at room temperature (23-25℃). Tested mice were gently put

into water, and the video was recorded for 6 min. Mice's immobile time in the

water was analyzed and quantified by the ANY-Maze tracking software

(Version 4.7, Stoelting CO).

Sucrose preference test (SPT). Mice from different groups were habituated for

two days for two bottles (one is a sucrose solution bottle, the other is a water

bottle, with left and right random placement) in their home cages. After

habituation, the sucrose preference of mice from both groups was tested. The

intake volume of water and 2% sucrose solution was measured within 2 hours.

Sucrose preference was determined as the ratio of the intake volume of

sucrose solution to total water consumption.

Tail-suspension test (TST). The tested mice were gently grabbed with the tail

and invertedly suspended in front of the recording camera by fixing their tail to

the adhesive tape for 6 minutes of recording. Mice immobile time during the

suspension was analyzed and quantified by the ANY-Maze tracking software

(Version 4.7, Stoelting CO).

Immunohistochemistry

Immunostaining for c-Fos was conducted on four distinct groups of mice: those exposed to acute heat, chronic heat, those injected with rabies virus into the pPVT and exposed to heat, and those with virus-labelled POA recipient pPVT neurons and exposed to heat.

Retrograde tracing. The rabies virus system was adopted for retrograde tracing, as previously reported (Watabe-Uchida et al., 2012). On day 1, 50 nl of AAV1-hSyn-cre, 50nl of AAV-FLEX-TVA-mCherry, and 50 nl of AAV-FLEX-RG were injected into pPVT. Subsequently, 80 nl of SAD△G virus was injected into pPVT on day 10. Mice were then allowed seven days for full expression before being sacrificed.

Tissue preparation. For mice from different groups, brain samples were prepared as follows: 40 to 60 minutes after exposure to heat, mice were anesthetized with i.p. injection of a ketamine-xylazine cocktail and transcardially perfused with 1X phosphate-buffered saline (PBS, Invitrogen) and then 4% paraformaldehyde (PFA, Sigma-Aldrich). The brain was then extracted and postfixed overnight in 4% PFA and finally dehydrated with 30% glucose solution for 48 h. The well-prepared brain samples were embedded in OCT (ThermoFisher) and sliced into 30-μm coronal sections by a cryostat (Leica) and stored in 1X PBS before immunohistochemistry.

Immunofluorescent staining. Brain sections were blocked in 5% normal goat serum (ThermoFisher) in PBS with 0.3% Triton X-100 (Sigma Aldrich) for 40

min and then incubated with primary rabbit anti-c-Fos antibody (1:2000, Cell

Signaling Technology) at 4°C overnight. After several thorough washing

pieces in PBS, the sections were incubated with goat anti-rabbit IgG-Alexa

Fluor 488 secondary antibody (1:1000, Invitrogen) in a blocking solution for 2

hours. After that, the sections were rinsed in PBS again and finally mounted

onto glass slides. Microscopic images were taken under a confocal laser

scanning microscope (C1, Nikon).

Neuronal counting. As for the statistical analysis of c-Fos positive neurons,

profile counting based on the rectangular square drawn in each picture was

performed on image J. For every counted brain area, c-Fos positive neurons

on three consecutive brain slices were selected and measured. And the value

averaged from profile counting on three consecutive brain slices was

regarded as the c-Fos positive neuronal number of specific brain nuclei for

one mouse.

Optogenetic experiments

For activation: 100 nL of AAV9-CaMKII-ChR2-mCherry was injected into POA,

optical fibers in pPVT. For inhibition: 100 nL of AAV1-hSyn-Cre-EGFP was

injected into POA, 200 nL of AAV9-hSyn-Dio-eNpHR3.0-EGFP or AAV9-

hSyn-Dio-EGFP in pPVT.

Real-time place preference test. Mice were placed into a two-chamber box for

10 minutes of habituation to make sure that mice had no preference for either

side of the chamber. Another 10 minutes' recording was started. The left

chamber was set as the light ON compartment while the other was the light

OFF chamber. Once mice entered light ON chamber, 473-nm light with 10 ms,

10 Hz (unless otherwise indicated, Newdoon Technology) was delivered via

an optic cable (200-µm core, 0.37 NA, Doric Lens) while the light was turned

off when mice entered light OFF chamber. Laser power was 5 mW measured

at the tip of the fiber, which was implanted 0.1-0.2 mm above the targeted

nucleus.

Elevated plus maze test, three-chamber test, and female encounter test. To

explore the effect of optogenetic activation of POA excitatory terminals within

pPVT on the levels of anxiety, sociability, and motivation in mice, mice in

these three behavioral tests were recorded with 3 minutes as the baseline,

followed by another 3 minutes' delivery of 473-nm light stimulation via an optic

cable (200-µm core, 0.37 NA, Doric Lens).

Measurement of pupil size. To directly reflect the effect of POA excitatory

terminals within pPVT on hyperarousal levels, mice were head-fixed and 473-

nm light was delivered with 10 Hz, 10 ms at the mode of 10 s light off, 10 s

light on, and 10 s light off to measure the changes of pupil size. And pupil size

of mice during the whole process was synchronously captured through a high-

speed camera at 30 frames per second. All pictures were finally integrated

into video format through the FFmpeg program. The changes in pupil size

among established videos were further analyzed by the DeepLabCut method

(Mathis et al., 2018) for precise tracking and plotting. The time point for

statistical analysis of normalized pupil size was at 5s for Pre, 15s for Light,

and 25s for Post, respectively.

Chronic optogenetic stimulation. Mice are subjected to 473-nm light stimulation with 10 ms, 10 Hz (unless otherwise indicated, Newdoon Technology) via an optic cable (200-μm core, 0.37 NA, Doric Lens) 2 mins-on, 2 mins-off for 20 min per day for up to 21 days (Sidor et al., 2014). Laser power does not exceed 5 mW measured at the tip of the fiber.

Optogenetic inhibition. Mice exposed to chronic heat were synchronously illuminated with a 589-nm light laser (Newdoon Technology) to inhibit POA recipient pPVT neurons specifically. The stimulus was 20 mW with a cyclical mode consisting of 3 minutes' light continuous on and 3 minutes' light off, which was conducted for 21 days only during the heat exposure period (90 minutes per day) for each mouse. The emotion-related behaviors and hyperarousal states were measured as formerly mentioned and analyzed by Any-maze software.

Fiber photometry

100 nL of AAV1-hSyn-Cre-EGFP in POA, 200 nL of AAV9-hSyn-Flex-jGCaMP8F-WPRE in pPVT, optical fibers in pPVT. Calcium signals were recorded for 1 minute for each mouse before, during heat exposure on day 1, the second day after heat exposure on 1st day, and after chronic heat exposure on day 22. The frequency and amplitude of calcium events were collected for data analysis.

Before and after chronic heat exposure, the performance of each mouse in the elevated plus maze and the chronic heat-exposed chamber was in

combination with calcium recording for 10 minutes, respectively. Targeting of

event-related time points and calculation of speed were achieved through the

SLEAP machine learning method (Pereira et al., 2022).

Fiber photometry data were collected with a TDT system at a sampling

frequency of 1017 Hz. The LED power at the tip of the patch cord was less

than 20 μW. The 405- and 465- signals were simultaneously recorded. The

isosbestic 405 nm control signal was filtered using a polyfit regression to limit

the influence of fluorescence decay during the session. And the fitted control

signal was then subtracted from the 465-signal to remove artifacts from the

intracellular calcium-dependent GCaMP fluorescence. The calculation for the

dynamics of fluorescence in chronic calcium recording was performed using

the formula $\triangle F/F = (F_{465} - F_{405}) / F405$. As corroborated by previous studies

(Shao et al., 2022; Xia et al., 2021), a calcium signal wave exceeding $\mu + 3\sigma$

was regarded as a fluorescence transient. Here, μ and σ respectively denote

the average and the standard deviation of the baseline signal of each mouse.

The baseline signal for each mouse was randomly selected, with a duration

ranging from 5 to 10 seconds. As for the activity changes of POA recipient

pPVT neurons during emotional transitions and hyperarousal states, the $\triangle F/F$

was then analyzed using a z-Score relative to the mean and standard

deviation of the session. Data were visualized and analyzed with custom

MATLAB scripts.

Electrophysiological recordings

Brain slice preparation and whole-cell recording. As for the confirmation of

functional connectivity, coronal mice brain slices were prepared as follows.

Mice quickly anesthetized by isoflurane were then transcardially perfused with

pH7.4 NMDG artificial cerebrospinal fluid (aCSF) in which contains (mM):

NMDG 92, KCl 2.5, NaH_2PO_4 1.25, $NaHCO_3$ 20, HEPES 10, Glucose 25, Na-

ascorbate 5, thiourea 2, Na-pyruvate 3, $MgSO_4$ 10, $CaCl_2$ 0.5, N-acetyl-L-

cysteine 12. Once completion of perfusion, mice's brain was quickly and

smoothly extracted and placed into cutting NMDG solution for frozen 30 s.

And coronal sections at 270-μm were prepared with a vibratome (Campden

5100MZ-PLUS Vibrotome) and then transferred to a chamber with 34□°C

NMDG aCSF for the first 15 minutes of recovery. And subsequently, brain

slices were further gently transferred into another chamber holding aCSF at

room temperature containing HEPES, pH 7.4, which contains (mM): NaCl 92,

KCl 2.5, NaH_2PO_4 1.25, $NaHCO_3$ 20, HEPES 10, Glucose 25, Na-ascorbate 5,

thiourea 2, Na-pyruvate 3, MgSO4 10, $CaCl_2$ 0.5, N-acetyl-L-cysteine 12.

After 40 to 60 minutes of recovery, the brain slices were placed into a

recording chamber with normal aCSF, pH 7.4, which contained (mM): NaCl

125, KCl 2.5, Glucose 11, $NaHCO_3$ 26, NaH_2PO_4 1.25, $CaCl_2$ 2, and $MgCl_2$ 2

for neuronal recording.

Functional connectivity. To record optically evoked postsynaptic currents, we

used an internal solution containing (mM): K-gluconate 130, KCl 10, HEPES

10, EGTA 1, $MgCl_2$ 2, Na_2-ATP 2, Na_3-GTP 0.4. Light pulses at 470 nm were

delivered through a light stimulator (Polygon400 DSI-E-0470-0590-NK1

Dynamic Spatial Illuminator) to activate POA ChR2-EGFP-expressing

terminals while patched pPVT neurons were held at -70mV. The neuron with

access resistance <20 MΩ and leak current <100 pA was included for data

collection.

Measurements of synaptic activity and intrinsic properties. To measure the

changes in synaptic transmission of pPVT neurons after chronic heat

exposure and acute heat exposure, coronal mice's brain slices were prepared

as mentioned above. The internal solution for both miniature inhibitory

(sIPSCs, held at +10 mV) and excitatory (sEPSCs, held at -70 mV) post

synaptic currents recording was Cesium-gluconate based solution with pH

7.3-7.4 which contains (mM): CsMSF 130, HEPES 10, EGTA 1, $MgCl_2$-$6H_2O$

2, Na_2ATP 2, NaGTP 0.4, QX-314-Cl 5, TEA-OH 5. And Tetrodotoxin (0.5 mM)

was present for both miniature EPSCs (mEPSCs) and miniature IPSCs

(mIPSCs) recording. Once completion of recording, brain slices were fixed

overnight with 4% PFA and the correct position of recorded pPVT neurons

was identified by staining with streptavidin-conjugated Alex 405

(ThermoFisher). For measuring the firing frequency under current clamp

recording, steady state current was injected in +10 pA increments from 0 to

100 pA. All action potential properties and excitability recordings in Figure S8

were performed in the presence of 10 µM CNQX and 100 µM picrotoxin. As

mentioned in the previous paper (Knowland et al., 2017), the firing rate

evoked by current injections and the half-width were calculated in Clampfit

software (Molecular Devices) by using peak detection function and full-width

at half max amplitude, respectively. The threshold was measured as the

change in the voltage from rest at which the slope =20 V/s. Membrane

resistance was calculated from the change in voltage elicited after a 50 ms 5

mV hyperpolarizing step from -70 mV (the last 10 ms of the step from baseline

was taken as $\triangle$V). Capacitance and membrane time constant were calculated

by Clampfit during the first minute after breaking in to cell.

Long-term potentiation. To measure the neuroplasticity of the POA to pPVT

pathway, two batches of mice with enough expression of CaMKII-ChR2-

mCherry within the POA were prepared: One serving as the control group and

the other as the chronic heat group. Sagittal brain slices from the mice were

prepared, and a low-chloride internal solution was used to record pPVT

neurons. EPSCs evoked by focal optogenetic stimulation through the

objective (approximately 2-3 mm above the surface of brain slice) in a top-

down vertical direction were elicited using a Polygon400 Multiwavelength

Dynamic Patterned Illuminator (Mightex). A circular area (roughly 500 μm

diameter) covering the recorded cell soma was illuminated with brief pulses of

blue light (470 nm, duration: 2 ms). LTP induced by optogenetic stimulation of

pPVT neurons was elicited by HFS consisting of three episodes of 30 Hz blue

light pulses at 20 s intervals.

All above signals were collected using a MultiClamp 700B amplifier controlled

by Clampfix 10.4 software via a Digital data 1550 interface (Molecular

Devices). Electrical signals were filtered at 3 kHz, digitized at 10 kHz, and

further analyzed using Clampfit 10.7 (Molecular Devices).

Statistical analysis

Statistical analysis was performed using GraphPad Prism 8.0. Values were

shown as mean $\pm$ standard error of the mean (SEM). Student's t-test

(paired or unpaired, parametric or nonparametric), one-way ANOVA with

Tukey post-hoc test, or two-way ANOVA followed by Sidak post-hoc test,

were conducted for statistical analysis. The value for statistical significance

was $p < 0.05$. For each experiment, the detailed statistics are described in

the corresponding figure legend.

Data availability

All data are contained in the main text, extended data or the supplementary

materials are available from the author upon reasonable request. The source

data underlying Figs 1C, E, G-I, L, M-Q, 2H, K, 3D-H, 4C, E, G-I, L, M, P-W,

5B-E, H-L, O-S, V, 6C-F, H, I, L, M and Figure 1-figure supplement 1A-I,

Figure 1-figure supplement 2B, D, F, G, H, K, L, Figure 2-figure supplement

1C, Figure 3-figure supplement 1C, D, Figure 3-figure supplement 2C, D, G, I,

Figure 5-figure supplement 1C, F, G-J, M, Figure 6-figure supplement 1A-F

are provided as a Source Data file.

Acknowledgements: This work was funded by following funding sources:

HKRGC C4012-22G & HMRF 09203236. We thank Dr. Xu Tao, Romeo Goh,

Zhuyun and Rong Kanglin for their technical assistance.

Conflict of interest: The authors declare no competing financial interests.

Author contributions

Y Ke and WH Yung perceived the project, designed the experiments, and analyzed the data. ZP Cao designed the experiments, collected, and analyzed the data. All authors discussed the findings and contributed to the writing of the manuscript.

Author ORCIDs

Ya Ke https://orcid.org/0000-0001-8166-6653

Zhiping Cao https://orcid.org/0000-0001-5560-7385

Ethics

All experimental procedures were approved by the Animal Experimentations and Ethics Committee (AEEC number: 18-232-MIS-5-C) at the Chinese University of Hong Kong (Animal license number: (23-120) in DH/HT&A/8/2/1 Pt.45). All surgery was performed under deep surgical anesthesia and every effort was made to minimize animal suffering.

References

Ali F, Kwan AC (2020) **Interpreting in vivo calcium signals from neuronal cell bodies, axons, and dendrites: a review** *Neurophotonics* **7**:011402.

Allen WE, DeNardo LA, Chen MZ, Liu CD, Loh KM, Fenno LE, Ramakrishnan

C, Deisseroth K, Luo LQ (2017) **Thirst-associated preoptic neurons**

encode an aversive motivational drive *Science* **357**:1149-1155.

Altena E, Baglioni C, Sanz-Arigita E, Cajochen C, Riemann D (2023) **How to**

deal with sleep problems during heatwaves: practical recommendations

from the European Insomnia Network *J Sleep Res* **32**:e13704.

Ambroziak W, Nencini S, Pohle J, Zuza K, Pino G, Lundh S, Araujo-Sosa C,

Goetz LIL, Schrenk-Siemens K, Manoj G, Herrera MA, Acuna C, Siemens J

(2024) **Thermally-induced neuronal plasticity that mediates heat**

tolerance BioRxiv Preprint. DOI: https://doi.org/10.1101/2024.08.06.606929.

Amr M, Volpe FM (2012) **Seasonal influences on admissions for mood**

disorders and schizophrenia in a teaching psychiatric hospital in Egypt

J Affect Disord **137**:56-60.

Augustine V, Gokce SK, Lee S, Wang B, J. Davidson T, Reimann F, Gribble F,

Deisseroth K, Lois C, Oka Y (2018) **Hierarchical neural architecture**

underlying thirst regulation *Nature* **555**:204-209.

Barriopedro D, Fischer EM, Luterbacher J, Trigo RM, Garcia-Herrera R (2011)

The hot summer of 2010: redrawing the temperature record map of

Europe *Science* **332**:220-224.

Basu R, Gavin L, Pearson D, Ebisu K, Malig B (2018) **Examining the**

association between apparent temperature and mental health related

emergency room visits in California *Am J Epidemiol* **187**:726-735.

Beas BS, Wright BJ, Skirzewski M, Leng Y, Hyun JH, Koita O, Ringelberg N,

Kwon HB, Buonanno A, Penzo MA (2018) **The locus coeruleus drives**

disinhibition in the midline thalamus via a dopaminergic mechanism *Nat*

Neurosci **21**:963-973.

Beecher ME, Eggett D, Erekson D, Rees LB, Bingham J, Klundt J, Bailey RJ,

Ripplinger C, Kirchhoefer J, Gibson R, Griner D, Cox JC, Boardman RD (2016)

Sunshine on my shoulders: weather, pollution, and emotional distress *J*

Affect Disord **205**:234-238.

Bhatnagar S, Dallman M (1998) **Neuroanatomical basis for facilitation of**

hypothalamic–pituitary–adrenal responses to a novel stressor after

chronic stress *Neuroscience* **84**:1025-1039.

Bridges MT, Tulapurkar ME, Shah NG, Singh IS, Hasday JD (2012)

Tolerance for chronic heat exposure is greater in female than male mice

Int J Hyperthermia **28**:747-755.

Cassuto Y (1968) **Metabolic adaptations to chronic heat exposure in the**

golden hamster *Am J Physiol* **214**:1147-1151.

Cruz B, Vozella V, Borgonetti V, Bullard R, Bianchi PC, Kirson D, Bertotto LB,

Bajo M, Vlkolinsky R, Messing RO, Zorrilla EP, Roberto M (2024)

Chemogenetic inhibition of central amygdala CRF-expressing neurons

decreases alcohol intake but not trauma-related behaviors in a rat model

of post-traumatic stress and alcohol use disorder *Mol Psychiatry* Online

ahead of print 21 March 2024; doi: 10.1038/s41380-024-02514-8.

Cure M (1989) **Plasma corticosterone response in continuous versus**

discontinuous chronic heat exposure in rat *Physiol Behav* **45**:1117-1122.

Curtis GR, Oakes K, Barson JR (2021) E**xpression and Distribution of**

Neuropeptide-Expressing Cells Throughout the Rodent Paraventricular

Nucleus of the Thalamus *Frontiers in Behavioral Neuroscience* **14**:634163.

Deuis JR, Dvorakova LS, Vetter I (2017) **Methods Used to Evaluate Pain**

Behaviors in Rodents *Front Mol Neurosci* **10**:284.

Dominiak M, Swiecicki L, Rybakowski J (2015) **Psychiatric hospitalizations**

for affective disorders in Warsaw, Portland: Effects of season and

intensity of sunlight *Psychiatry Res* **229**:287-294.

Do-Monte FH, Quiñones-Laracuente K, Quirk GJ (2015) **A temporal shift in**

the circuits mediating retrieval of fear memory *Nature* **519**:460-463.

Fausnacht DW, Kroscher KA, McMillan RP, Martello L, Baumgard LH, Sellsby

JT, Hulver MW, and Rhoads RP (2021) **Heat Stress Reduces Metabolic**

Rate While Increasing Respiratory Exchange Ratio in Growing Pigs

Animals **11**:215

García-Herrera R, Díaz J, Trigo RM, Fischer EM (2010) **A review of the**

European summer heat wave of 2003 *Crit Rev Environ Sci Technol* **40**:267-

306.

Ghasemi M, Navidhamidi M, Rezaei F, Azizikia A, Mehranfard N (2022)

Anxiety and hippocampal neuronal activity: Relationship and potential

mechanisms *Cog Affect Behav Neurosci* **22**:431-449.

Grover LM, Kim E, Cooke JD, Holmes WR (2009) **LTP in hippocampal area**

CA1 is induced by burst stimulation over a broad frequency range

centered around delta *Learn Mem* **16**:69-81.

Hansen A, Bi P, Nitscheke M, Ryan P, Pisaniello D, Tucker G (2008) **The**

effect of heat waves on mental health in a temperate Australian city

Environ Health Perspect **116**:1369-1375.

Heilbronner U, Kampen MV, Flüggeg G (2004) **The Alpha-2B Adrenoceptor**

in the Paraventricular Thalamic Nucleus is Persistently Upregulated by

Chronic Psychosocial Stress *Cell Mol Neurobiol* **24**:815-831.

Herman JP (2013) **Neural control of chronic stress adaptation** *Front*

Behav Neurosci **7**: 61.

Heydendael W, Sharma K, Iyer V, Luz S, Piel D, Beck S, Bhatnagar S (2011)

Orexins/hypocretins act in the posterior paraventricular thalamic

nucleus during repeated stress to regulate facilitation to novel stress

Endocrinology **152**:4738-4752.

Hwong, Udaka Y, Kauer JA (2008) **High-Frequency Afferent Stimulation**

Induces Long-Term Potentiation of Field Potentials in the Ventral

Tegmental Area *Neuropsychopharmacology* **33**:1704-1712.

Jin Y, Hu YX, Han DP, Wang M (2011) **Chronic heat stress weakened the**

innate immunity and increased the virulence of highly pathogenic avian

influenza virus H5N1 in mice *J Biomed Biotechnol* **2011**:367846.

Jurik A, Auffenberg E, Klein S, Deussing JM, Schmid RM, Wotjak CT,

Thoeringer CK (2015) **Roles of prefrontal cortex and paraventricular**

thalamus in affective and mechanical components of visceral nociception. *Pain* **156**:2479-2491.

Kawakami Y, Sielski R, and Kawakami T (2018) **Mouse Body Temperature Measurement Using Infrared Thermometer During Passive Systemic Anaphylaxis and Food Allergy Evaluation** *J Vis Exp* **139**:58391.

Kenny GP, Yardley J, Brown C, Sigal RJ, Jay O (2010) **Heat stress in older individuals and patients with common chronic diseases** *CMAJ* **182**:1053-1060.

Knowland D, Lilascharoen V, Pacia CP, Shin S, Wang EHJ, Lim BK (2017) **Distinct Ventral Pallidal Neural Populations Mediate Separate Symptoms of Depression** *Cell* **170**:284-297.

Lee HJ, Kim L, Joe SH, Suh KY (2002) **Effects of season and climate on the first manic episode of bipolar affective disorder in Korea** *Psychiatry Res* **113**:151-159.

Li YH, Li S, Wei CG, Wang HY, Sui N, Kirouac GJ (2010) **Orexins in the paraventricular nucleus of the thalamus mediate anxiety-like responses in rats** *Psychopharmacology (Berl)* **212**:251-265.

Malmquist A, Hjerpe M, Glaas E, Karlsson H, Lassi T (2002) **Elderly People's Perceptions of Heat Stress and Adaptation to Heat: An Interview Study** *Int J Environ Res Public Health* **19**: 3775.

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M (2018) **DeepLabCut: markerless pose estimation of user-defined body parts with deep learning** *Nat Neuroscience* **21**:1281-11289.

McMorris T, Swain J, Smith M, Corbett J, Delves S, Sale C, Harris RC, Potter J (2006) **Heat stress, plasma concentrations of adrenaline, noradrenaline, 5-hydroxytryptamine and cortisol, mood state and cognitive performance**. *Int J Psychophysiol* **61**:204-215.

Mitchell D, Heaviside C, Vardoulakis S, Huntingford C, Masato G, Guillod PB, Frumhoff P, Bowery A, Wallom D and Allen M (2016) **Attributing human mortality during extreme heat waves to anthropogenic climate change**. *Environ Res* Lett **11**:74006.

Mora C, Dousset B, Caldwell IR, Powell FE, Geronimo RC, Bielecki CR, Counsell CWW, Dietrich BS, Johnston ET, Louis LV, Lucas MP, McKenzie MM, Shea AG, Tseng Han, Giambelluca TW, Leon LR, Hawkins E & Trauernicht C (2017) **Global risk of deadly heat** *Nat Clim Change* **7**:501-506.

Morrison SF, Nakamura K (2011) **Central neural pathways for thermoregulation** *Front Biosci* **16**:74-104.

Mulder RT, Cosgriff JP, Smith AM, Joyce PR (1990) **Seasonality of mania in New Zealand** *Aust N Z J Psychiatry* **24**:187-190.

Murray KO, Clanton TL, Horowitz M (2022) **Epigenetic responses to heat: From adaptation to maladaptation** *Exp Physiol* **107**:1144-1158.

Oppermann E, Kjellstrom T, Lemke B, Otto M, Lee JKW (2021) **Establishing intensifying chronic exposure to extreme heat as a slow onset event with implications for health, wellbeing, productivity, society and economy** *Curr Opin Environ Sustain* **50**:225-235.

Pantoni MM, Herrera GM, Van Alstyne KR, Anagnostaras SG (2020) **Quantifying the Acoustic Startle Response in Mice Using Standard Digital Video** *Front Behav Neurosci* **14**:83.

Penzo MA, Gao Claire (2021) **The paraventricular nucleus of the thalamus: an integrative node underlying homeostatic behavior** *Trends Neurosci* **44**:538-549.

Penzo MA, Robert V, Tucciarone J, De Bundel D, Wang MH, Van Aelst L, Darvas M, Parada LF, Palmiter R, He M, Huang ZJ, Li B (2015) **The paraventricular thalamus controls a central amygdala fear circuit** *Nature* **519**:455-459.

Pereira TD, Tabris N, Matsliah A, Turner D, Li JY, Ravindranath S, Papadoyannis ES, Normand E, Deutsch D, Wang Z, McKenzie-Smith GC, Mitelut CC, Castro MD, D'Uva J, Kislin M, Sanes DH, Kocher SD, Wang SSH, Falkner AL, Shaervitz JW, Murthy M (2022) **SLEAP: A deep learning system for multi-animal pose tracking** *Nat Method* **19**:486-495.

Périard JD, Travers GJS, Racinais S, Sawka MN (2016) **Cardiovascular adaptations supporting human exercise-heat acclimation** *Autonomic Neuroscience* **196**:52-62

Perica MI, Luna B (2023) **Impact of stress on excitatory and inhibitory markers of adolescent cognitive critical period plasticity** *Neurosci Biobehav Rev* **153**:105378.

Podstawski R, Boryslawski K, Pomianowski A, Krystkiewicz W, Żurek P (2021)

Endocrine Effects of Repeated Hot Thermal Stress and Cold Water

Immersion in Young Adult Men *Am J Mens Health* **15**:15579883211008339.

Podstawski R, Boryslawski K, Pomianowski A, Krystkiewicz W, Boraczyński T,

Mosler D, Wasik J, Jaszczur-Nowicki J (2021) **The Effects of Repeated**

Thermal Stress on the Physiological Parameters of Young Physically

Active Men Who Regularly Use the Sauna: A Multifactorial Assessment

Int J Environ Res Public Health **18**:11503

Privitera M, Ferrari KD, von Ziegler LM, Sturman O, Duss SN, Floriou-Servou

A, Germain PL, Vermeiren Y, Wyss MT, P De Deyn P, Weber B, Bohacek J

(2020) **A complete pupillometry toolbox for real-time monitoring of locus**

coeruleus activity in rodents *Nat Protoc* **15**:2301-2320.

Ren SC, Wang YL, Yue FG, Cheng XF, Dang RZ, Qiao QC, Sun XQ, Li X,

Jiang Q, Yao JW, Qin H, Wang GZ, Liao X, Gao D, Xia JX, Zhang J, Hu B,

Yan JA, Wang YJ, Xu M, Han YY, Tang XD, Chan XW, He C, Hu ZA (2018)

The paraventricular thalamus is a critical thalamic area for wakefulness

Science **362**:429-434.

Robine JM, Cheung SLK, Le Roy S, Van Oyen H, Griffiths C, Michel JP,

Herrmann FR (2008) **Death toll exceeded 70 000 in Europe during the**

summer of 2003 *C R Biol* **331**:171-178.

Rousset B, Cure M, Jordan D, Kervran A, Bornet H, Mornex R (1984)

Metabolic alterations induced by chronic heat exposure in the rat: the

involvement of thyroid function *Pflugers Arch* **401**:64-70.

Sampath V, Shalakhti O, Veidis E, Efobi JAI, Shamji MH, Agache I, Skevaki C,

Renz H, Nadeau KC (2023) **Acute and chronic impacts of heat stress on**

planetary health *Allergy* **78**:2109-2120.

Shao J, Gao DS, Liu YH, Chen SP, Liu NA, Zhang L, Zhou XY, Xiao Q, Wang

LP, Hu HL, Wang F (2022) **Cav3.1-driven bursting firing in ventromedial**

hypothalamic neurons exerts dual control of anxiety-like behavior and

energy expenditure *Mol Psychiatry* **27**:2901-2913.

Sidor MM, McClung C (2014) **Timing matters: using optogenetics to**

chronically manipulate neural circuitry and rhythms *Front Behav Neurosci*

8:41.

Spencer SJ, Fox JC, Day TA (2004) **Thalamic paraventricular nucleus**

lesions facilitate central amygdala neuronal responses to acute

psychological stress *Brain Res* **997**:234-237.

Tan CL, Cooke EK, Leib DE, Lin YC, Daly GE, Zimmerman CA, Knight ZA

(2016) **Warm-sensitive neurons that control body temperature** *Cell*

167:47-59.

Tollefson J (2023) **Earth's hottest month: these charts show what**

happened in July and what comes next *Nature* **620**:703-704.

Trang PM, Rocklöv J, Giang KB, Van Minh H, Tinh LT, Nilsson M (2015)

Weather variations and hospital admissions for depressive disorders: a

case study in Hanoi *Ann Psychiatry Ment Health* **5**:1095.

Venkatachalam K, Montell C. TRP channels (2007) **Annu Rev Biochem**

76:387-417.

Vertes RP, Linley SB, Hoover WB (2015) **Limbic circuitry of the midline**

thalamus *Neurosci Biobehav Rev* **54**:89-107.

Wang L, Liu FD, Luo Y, Zhu LQ, Li GH (2015) **Effect of acute heat stress on**

adrenocorticotropic hormone, cortisol, interleukin-2, interleukin-12 and

apoptosis gene expression in rats *Biomed Rep* **3**:425-429.

Watabe-Uchida M, Zhu L, Ogawa SK, Vamanrao A, Uchida N (2012) **Whole-**

Brain Mapping of Direct Inputs to Midbrain Dopamine Neurons *Neuron*

74:858-873.

Whitney DK, Sharma V, Kueneman K (1999) **Seasonality of manic-**

depressive illness in Canada *J Affect Disord* **55**:99-105.

Xia G, Han Y, Meng F, He Y, Srisai D, Farias M, Dang MH, Palmiter RD, Xu Y,

Wu Q (2021) **Reciprocal control of obesity and anxiety-depressive**

disorder via a GABA and serotonin neural circuit *Mol Psychiatry* **26**:2837-

2853.

Yu S, Qualls-Creekmore E, Rezai-Zadeh K, Jiang YY, Berthoud HR, Morrison

CD, Derbenev AV, Zsombok A, Münzberg H (2016) **Glutamatergic preoptic**

area neurons that express leptin receptors drive temperature-dependent

body weight homeostasis *J Neurosci* **36**:5034-5046.

Zhang GW, Shen L, Tao C, Jung AH, Peng B, Li Z, Zhang LI, Whit Tao HZ

(2021) **Medial preoptic area antagonistically mediates stress-induced**

anxiety and parental behavior *Nat Neurosci* **24**:516-528.

Zhang QH, Hao JW, Li GL, Ji XJ, Zhou M, Yao YM (2020) **Long-lasting**

neurobehavioral alterations in burn-injured mice resembling post-

traumatic stress disorder in humans *Exp Neurol* **323**:113084.

Zhang Y, Kerman IA, Laque A, Nguyen P, Faouzi M, Louis GW, Jones JC,

Rhodes C, Münzberg H (2011) **Leptin-receptor-expressing neurons in the**

dorsomedial hypothalamus and median preoptic area regulate

sympathetic brown adipose tissue circuits *J Neurosci* **31**:1873-1884.

Zhao ZD, Yang WZ, Gao CC, Fu X, Zhang W, Zhou Q, Chen WP, Ni XY, Lin

JK, Yang J, Xu XH, Shen WL (2016) **A hypothalamus circuit that controls**

body temperature *Proc Natl Acad Sci USA* **114**:2042-2047.

Zhu YT, Wang RB, Wang YH (2016) **A Comparative Study of the Impact of**

Theta-Burst and High-Frequency Stimulation on Memory Performance

Front Hum Neurosci **10**:19.

1140

Figure legends

Figure 1. Chronic heat exposure produces negative emotional valence

and hyperarousal states but not depression-like behaviors. (A)

Experimental schematics. Mice (n=10 in each group) were divided into Control

and Heat groups and conducted with chronic exposure to room temperature

and heat conditions, respectively, followed by behavioral tests. **(B, C)** The

heatmap of representative tracking trace examples in elevated plus maze

(EPM) test and the time spent in the open arms of EPM (Mann-Whitney

unpaired two-tailed U test; U=0, ***p<0.001). **(D, E)** The heatmap of

representative tracking trace examples in three-chamber test (TCT) and the

interaction time with an unfamiliar male mouse relative to the inanimate object

in TCT (Mann-Whitney unpaired two-tailed U test; U=13, ***p=0.0039). **(F, G)**

The heatmap of representative tracking trace examples in female encounter

test (FET) and the time surrounding the unfamiliar female mouse compared to

an unfamiliar male mouse in FET (Mann-Whitney unpaired two-tailed U test;

U=10, **p=0.0015). **(H, I)** The 1st-time attack latency (Mann-Whitney unpaired

two-tailed U test; U=21, *p=0.0288) and the attack durations (Mann-Whitney

unpaired two-tailed U test; U=23, *p=0.0433) in the aggression test. **(J, K)**

Visualized and representative acoustic startle response example (left panel)

and corresponding labeled body parts' skeletons (shown as purple dots) (right

panel) in acoustic startle response (ASR) test from Control and Chronic Heat

group.**(L, M)** The startle amplitude (Mann-Whitney unpaired two-tailed U test;

U=6, ***p=0.0003) and the pre-pulse / pulse ratio (Mann-Whitney unpaired

two-tailed U test; U=26, p=0.0753) in ASR test. **(N-Q)** The immobile time in

the forced swim test (FST) (Mann-Whitney unpaired two-tailed U test; U=35,

p=0.2799) and in the tail suspension test (TST) (Mann-Whitney unpaired two-

tailed U test; U=26, p=0.0753; The percentage of sucrose intake in the

sucrose preference test (SPT) (Mann-Whitney unpaired two-tailed U test;

U=49.5, p=0.4887); The time spent in the center of the open field test (OFT)

(Mann-Whitney unpaired two-tailed U test; U=35.5, p=0.2888). *p<0.05,

**p<0.01, ***p<0.001, N.S.: not significant.

Figure 2. Involvement of the hypothalamic preoptic area to posterior paraventricular thalamus projections. (A) The strategy of virus injection followed by heat exposure-induced c-Fos staining (n=3 mice). **(B)** The representative microphotograph showed the expression of rabies virus in the pPVT regions. **(C)** Rabies virus retro-labeled POA neurons. Upper: magnification: x4, scale bar: 400 μm; Lower: magnification: x10, scale bar: 40 μm; The top right corner of the lower picture: magnification: x60, scale bar: 5 μm. **(D)** The representative microphotograph and quantification showed that most of pPVT rabies virus retro-labeled POA neurons co-stained with heat exposure-induced c-Fos. Upper: magnification: x10, scale bar: 40 μm; Lower and left: magnification: x20, scale bar: 20 μm. **(E)** The strategy of virus injection (n=5 mice) and patch-clamp recording was performed on POA expressing ChR2-mCherry neurons and pPVT neurons using potassium and cesium, low chloride internal solutions, respectively. Representative traces showed that POA expressing ChR2-mCherry neurons exhibited robust firing in response to optical stimulation at different frequencies (n=15 neurons from 5 mice). **(F, G)** Representative traces showed that blue light stimulation evoked either excitatory postsynaptic current (EPSC) which could be blocked by cyanquixaline (CNQX, 10 μM) or inhibitory postsynaptic current (IPSC) which could be blocked by picrotoxin (PTX, 100 nM). **(H)** Pie chart showed the projection types recorded on pPVT neurons (n=63 neurons from 10 mice). **(I)** The representative recorded pPVT neuron was visualized by biocytin staining and was found being surrounded by POA expressing ChR2-mCherry terminals. Magnification: x60, scale bar: 20 μm. **(J, K)** The representative trace showed that the application of tetrodotoxin (TTX) eliminated the oPSC

1198 held at -70 mV while the addition of 4-amino-pyridine (4-AP, 1mM) recovered

1199 it and the quantification (n=24 neurons from 10 mice).

1200

Figure 3. Activity changes of POA recipient pPVT neurons throughout

chronic heat exposure. (A) Experimental schematics. Mice (n=5) were

stereotaxically injected with Cre-dependent GCaMP into the pPVT, followed

by the implantation of optical fiber and chronic calcium recording. **(B)**

Representative microphotographs showed the virus expression in the POA

and pPVT regions, scale bar: all 40 μm. **(C)** From the top to the bottom: the

representative calcium events (on the left panel) and the calcium events from

different mice (on the right panel) on different days. Each vertical stripe in the

heatmap represents one calcium event for each mouse. Statistical analysis of

the frequency and amplitude of POA recipient pPVT neurons' calcium events

compared for **(D)** Pre and During heat exposure on day 1 (Paired, parametric,

two tailed t-test; Frequency ($t=6.278$, $df=4$, $P=0.0033$); Amplitude ($t=4.344$,

$df=4$, $*p=0.0122$)); **(E)** Pre heat on day 1 and After heat on day 2 (Paired,

parametric, two tailed t-test; Frequency ($t=0.2787$, $df=4$, $p=0.7943$); Amplitude

($t=1.726$, $df=4$, $p=0.1595$)); **(F)** Pre heat on day 1 and During heat exposure

on day 21 (Paired, parametric, two tailed t-test; Frequency ($t=6.124$, $df=4$,

$**p=0.0036$); Amplitude ($t=4.704$, $df=4$, $**p=0.0093$)); **(G)** Pre heat on day 1

and after Chronic heat on day 22 (Paired, parametric, two tailed t-test;

Frequency ($t=6.216$, $df=4$, $**p=0.0034$); Amplitude ($t=1.36$, $df=4$, $p=0.2454$));

(H) During heat exposure on day 1 and day 21 (Paired, parametric, two tailed

t-test; Frequency (*t=1.242, df=4, p=0.2821*); Amplitude (*t=0.9424, df=4,*

p=0.3993)). * $p<0.05$, ** $p<0.01$, N.S.: not significant.

1223

Figure 4. POA recipient pPVT neurons are sufficient and necessary for

chronic heat exposure-induced negative emotional valence and

hyperarousal states in mice. (A) Experimental schematics. Mice (n=10 in

each group) were stereotaxically injected with either AAV1-hSyn-EGFP or

AAV9-CaMKII-ChR2-mCherry into the POA, followed by the implantation of

optical fiber into pPVT, chronic optogenetic activation, and behavioral tests.

(B, C) The heatmap of representative tracking trace examples in EPM test

and the time spent in the open arms (Mann-Whitney unpaired two-tailed U

test; U=2, ***$p<0.001$). **(D, E)** The heatmap of representative tracking trace

examples in TCT and the interaction time with an unfamiliar male mouse

relative to the inanimate object (Mann-Whitney unpaired two-tailed U test,

U=8; ***$p=0.0007$). **(F, G)** The heatmap of representative tracking trace

examples in FET and the time surrounding the unfamiliar female mouse

compared to an unfamiliar male mouse (Mann-Whitney unpaired two-tailed U

test; U=13, **$p=0.0039$). **(H, I)** The 1st-time attack latency (Mann-Whitney

unpaired two-tailed U test; U=8, $p=0.007$) and the attack durations (Mann-

Whitney unpaired two-tailed U test; U=19, *$p=0.0185$) in the aggression test.

(J, K) Visualized acoustic startle response examples (upper panel) and

corresponding labeled body parts' skeletons (lower panel) from hSyn-EGFP

group and ChR2-mCherry group. **(L, M)** The startle amplitude (Mann-Whitney

unpaired two-tailed U test; U=17, *$p=0.0115$) and the pre-pulse / pulse ratio

(Mann-Whitney unpaired two-tailed U test; U=48, p=0.9118) in ASR test. **(N)**

Experimental schematics. Mice (n≥6 in each group) were stereotaxically

injected with AAV1-hSyn-Cre-EGFP in the POA and either AAV9-Ef1a-Dio-

eNpHR3.0-EGFP or AAV8-hSyn-Dio-EGFP into the pPVT, followed by the

implantation of optical fiber and behavioral tests. **(O)** Representative

microphotographs showed the expression of AAV1-hSyn-Cre-EGFP and

AAV9-hSyn-Dio-eNpHR3.0 within the POA and pPVT, respectively.

Magnification: x4, scale bar: 100 μm (upper panel) and 250 μm (lower panel).

(P, Q) The time (One-way repeated measures ANOVA with Tukey post-hoc

test; $F_{(2, 27)}$=83.03, ***p<0.001; Control vs. Chronic heat, ***p<0.001;

Chronic heat vs. Chronic heat + inhibition, ***p<0.001) and the entry numbers

(One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 20.92)}$=27.96, ***p<0.001; Control vs. Chronic heat, ***p<0.001; Chronic heat

20.92)=27.96, ***p<0.001; Control vs. Chronic heat, ***p<0.001; Chronic heat

vs. Chronic heat + inhibition, ***p<0.001) into the open arms of EPM. **(R)** The

time spent with an unfamiliar male mouse compared to an inanimate object in

TCT (One-way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 27)}$=5.381, *p=0.0108; Control vs. Chronic heat, *p=0.0292; Chronic heat vs.

27)=5.381, *p=0.0108; Control vs. Chronic heat, *p=0.0292; Chronic heat vs.

Chronic heat + Inhibition, *p=0.0175). **(S)** The ratio of time with an unfamiliar

female mouse compared to an unfamiliar male mouse in FET (One-way

repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 27)}$=10.94,

***p=0.0003; Control vs. Chronic Heat, ***p=0.0002; Chronic Heat vs. Chronic

Heat + Inhibition, *p=0.0335). **(T, U)** The 1[st] attack latency (One-way repeated

measures ANOVA with Tukey post-hoc test; $F_{(2, 17)}$=7.426, *p=0.0048;

Control vs. Chronic Heat, *p=0.0157; Chronic Heat vs. Chronic Heat +

Inhibition, *p=0.0063) and the attack durations (One-way repeated measures

ANOVA with Tukey post-hoc test; F(2, 17)=13.38, ***p=0.0003; Control vs.

Chronic Heat, ***p=0.0003; Chronic Heat vs. Chronic Heat + Inhibition,

p=0.0051) in the aggression test. **(V, W) The startle amplitude (One-way

repeated measures ANOVA with Tukey post-hoc test; F(2, 21)=21.03,

****p<0.0001; Control vs. Chronic Heat, ***p<0.001; Chronic Heat vs. Chronic

Heat + Inhibition, ***p<0.001) and the pre-pulse / pulse ratio (One-way

repeated measures ANOVA with Tukey post-hoc test; F(2, 21)=3.802,

*p=0.039; Control vs. Chronic Heat, p=0.2616; Chronic Heat vs. Chronic Heat

+ Inhibition, p=0.2246) in ASR test. *p<0.05, **p<0.01, ***p<0.001, N.S.: not

significant.

1280

Figure 5. Chronically activated POA recipient pPVT neurons exhibited

exaggerated response to stressful situations. (A) Experimental

schematics. Mice (n=5) were stereotaxically injected with AAV1-hSyn-Cre-

EGFP in the POA and AAV9-hSyn-Flex-jGCaMP8F-WPRE in the pPVT,

followed by the implantation of optical fiber and calcium recording. **(B)** The

number of entries into the open arms of EPM (Paired, parametric, two tailed t-

test; t=6.901, df=4, **p=0.0023). **(C)** The number of instances where the mice

paused at the center area followed by back to closed arms of EPM (Paired,

parametric, two tailed t-test; t=9, df=4, ***p=0.0008). **(D)** Pie chart showed the

changes of the percentage of pause and then running back to closed arms. **(E)**

The changes of running speed in EPM (Mann-Whitney unpaired two-tailed U

test; U=139, **p=0.0028). **(F-H)** Heatmaps showed the calcium activities of

POA recipient pPVT neurons when mice performed the pause-and-run-back-to-closed-arms behavior for pre (n=13 trials from 5 mice) and post conditions (n=47 trials from 5 mice), Z-Score calcium signals, and statistical comparison (Paired, parametric, two tailed t-test; t=4.387, df=4, *p=0.0118). **(I)** The time spent in open arms of EPM (Paired, parametric, two tailed t-test; t=3.649, df=4, *p=0.0218). **(J)** The total number of the behavioral event: back from open to closed arms (Paired, parametric, two tailed t-test; t=6.901, df=4, **p=0.0023).

(K) Pie chart showed the changes of the number of running episodes from open to closed arms for pre and post conditions. **(L)** The changes of running speed in EPM (Mann-Whitney unpaired two-tailed U test; U=23, ***p=0.0005).

(M-O) Heatmaps showed the calcium activities of POA recipient pPVT neurons when mice performed fast running from open to closed arms for pre (n=14 trials from 5 mice) and post conditions (n=13 trials from 5 mice), Z-Score calcium signals, and statistical comparison (Paired, parametric, two tailed t-test; t=3.087, df=4, *p=0.0367). **(P)** The total distances of mice travelled in the chamber previously subjected to chronic heat (Paired, parametric, two tailed t-test; t=6.876, df=4, **p=0.0023). **(Q)** The changes of motion speed for pre and post conditions from one representative mouse. **(R)** The number of instances with motion speed exceeding 400 mm/s (Paired, parametric, two tailed t-test; t=5.100, df=4, ***p=0.007). **(S)** The number of fast running episodes for pre and post conditions. **(T-V)** Heatmaps showed the calcium activities of POA recipient pPVT neurons when the mice performed fast running in the previous chronic heat-exposed chamber (n=90 trials from 5 mice) compared to pre-heat condition (n=38 trials from 5 mice),

1317 Z-Score calcium signals, and statistical comparison (Paired, parametric, two

1318 tailed t-test; t=5.456, df=4, **p=0.0055). *$p<0.05$, **$p<0.01$, ***$p<0.001$.

1319

Figure 6. Increased pre- and post-synaptic excitability of pPVT neurons

but saturated circuitry neuroplasticity capacity following chronic heat

exposure. (A) Experimental schematics. *In vitro* brain slice recording was

performed on mice from three groups (n=16 neurons from 3 mice in the Room

temperature group, n=20 neurons from 4 mice in the Acute heat group, and

n=20 neurons from 4 mice in the Chronic heat group). **(B)** Representative

traces of miniature postsynaptic currents from three groups. Duration: 12 s.

Scale bar: 10 pA. **(C, D)** The changes of mIPSCs' amplitude (One-way

repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 54)}=8.226$,

***$p=0.0008$, Room temperature vs. Acute heat, ***$p=0.0005$; Room

temperature vs. Chronic heat, *$p=0.0268$) and frequency (One-way repeated

measures ANOVA with Tukey post-hoc test; $F_{(2, 53)}=1.346$, $p=0.269$, Room

temperature vs. Acute heat, $p=0.4327$; Room temperature vs. Chronic heat,

$p=0.9787$) of pPVT neurons. **(E, F)** The changes of mEPSCs' amplitude (One-

way repeated measures ANOVA with Tukey post-hoc test; $F_{(2, 52.98)}=0.1995$,

$p=0.8198$, Room temperature vs. Acute heat, $p=0.8433$; Room temperature

vs. Chronic heat, $p=0.8433$) and frequency (One-way repeated measures

ANOVA with Tukey post-hoc test; $F_{(2, 50.1)}=2.981$, $p=0.0598$, Room

temperature vs. Acute heat, $p=0.2931$; Room temperature vs. Chronic heat,

*$p=0.0473$) of pPVT neurons. **(G)** The representative traces of action potential

of pPVT neurons upon 100 pA current injection. **(H)** The changes of

excitability of pPVT neurons when different currents were injected to the patched pPVT neurons (n=11 neurons from 3 mice in each groups; Mann-Whitney unpaired two-tailed U test; 30 pA: U=22.5, **p=0.0091; 40 pA: U=31, *p=0.048; 50 pA: U=29.5, *p=0.0412; 60 pA: U=25, *p=0.0167; 70 pA: U=29.5, *p=0.0396). **(I)** The changes of rheobase of action potential (Mann-Whitney unpaired two-tailed U test; U=28.5, *p=0.0345). **(J)** Experimental schematics. Mice (n=5 mice in each group) stereotaxically injected with AAV9-CaMKII-ChR2-mCherry into the POA were then divided into the Room temperature and Chronic heat groups. Sagittal slices of mice were prepared for LTP induction and recording. **(K)** The representative traces showed pPVT neurons from mice exposed to room temperature and chronic heat exhibited different amplitude of oEPSCs after blue light-mediated high frequency stimulation. **(L)** The LTP induction and recording of POA to pPVT pathway from slices of mice exposed to room temperature and chronic heat conditions after blue light stimulation at 30 Hz (n=14 neurons from control group and 18 neurons from chronic heat group). **(M)** Statistical comparison of the amplitude of oEPSCs at different time points (Two-way repeated measures ANOVA with Sidak post-hoc test; Interaction: $F_{(3, 120)}$=0.3486, p=0.7902; Optical stimulation main effect: $F_{(3, 120)}$=76.2, ***p<0.001; Time points effect: $F_{(3, 120)}$=0.8215, p=0.4844; 1st min: Room temperature vs. Chronic heat: ***p<0.001; 10th min: Room temperature vs. Chronic heat: p=0.0001; 20th min: Room temperature vs. Chronic heat: ***p<0.001; 30th min: Room temperature vs. Chronic heat: **p=0.0024). *p<0.05, **p<0.01, ***p<0.001, N.S.: not significant.

Figure 7. A working model of neural circuit mechanisms underlying POA

recipient pPVT neurons-mediated chronic heat exposure-induced

negative emotional valence and hyperarousal states. Different from acute

heat exposure, chronic heat exposure-induced enhancement in excitatory

inputs to pPVT and saturated neuroplasticity contributed to the increased

membrane excitability underlie the heightened activities of POA recipient

pPVT neurons, rendering mice become more susceptible to stressful

situations manifested as negative emotional valence and hyperarousal states.

A Rabies virus

pPVT

Heat Exposure (38±2℃) → POA c-Fos staining

B Rabies virus in pPVT

Bregma -1.67 mm

3 V

pPVT

400 um

C Bregma 0.25 mm

400 um

POA

5 um

40 um

D Bregma 0.25 mm

40 um

Rabies virus / c-Fos

20 um

Overlap (%)

POA

E AAV9-hSyn-ChR2-mCherry

Potassium, Low Cl⁻

POA terminal

Cesium / Potassium low Cl⁻

pPVT

POA

5Hz

10Hz

20Hz

30Hz

F ACSF   +CNQX   +CNQX & PTX

30 pA

0.05 s

+10 mV

G ACSF   +PTX   +PTX & CNQX

-70 mV

30 pA

0.05 s

H EPSCs   IPSCs

12.7% (8/63)

52.4% (33/63)

34.9% (22/63)

EPSCs & IPSCs

I Biocytin / ChR2-mCherry

20 um

J oPSCs
+TTX
+4-AP

40 pA

200 ms

K Normalized oPSCs (%)

Control   TTX   TTX+4-AP