Novel replication-competent reporter-expressing Rift Valley Fever Viruses for

molecular studies.

Aitor Nogales[a,*],[#], Celia Alonso[a,*], Sandra Moreno[a], Gema Lorenzo[a], Belén

Borrego[a], Luis Martinez-Sobrido[b], Alejandro Brun[a,#].

[a]Centro de Investigación en Sanidad Animal (CISA), Instituto Nacional de

Investigación y Tecnología Agraria y Alimentaria, Consejo Superior de

Investigaciones Científicas (INIA-CSIC), Valdeolmos, 28130 Madrid, Spain

[b]Texas Biomedical Research Institute, San Antonio, TX, United States.

*Contributed equally.

[#]Corresponding authors.

Running Head: Rift Valley fever viruses expressing reporter genes

[#]Address correspondence to Alejandro Brun, brun@inia.csic.es.

ABSTRACT

Rift Valley fever virus (RVFV) is a mosquito-borne zoonotic disease that

causes severe disease in both domestic and wild ungulates and humans, making it a

significant threat to livestock and public health. The RVFV genome consists of three

single-stranded, negative-sense RNA segments differing in size: Small (S), Medium

(M) and Large (L). Segment S encodes the virus nucleoprotein N and the virulence-

associated factor non-structural (NSs) protein in opposite orientations, separated by

an intergenic region (IGR). To overcome the current need of using secondary

techniques to detect the presence of RVFV in infected cells, we used T7-driven polymerase plasmid-based reverse genetics to generate replication-competent recombinant (r)RVFV expressing Nanoluciferase (Nluc) or Venus fluorescent proteins. These reporter genes were used as valid surrogates to track the presence of RVFV in mammalian and insect cells. Notably, we explored the genome plasticity of RVFV and compared four different strategies by modifying the viral segment S in order to introduce the reporter gene foreign sequences. The reporter-expressing rRVFV were stable and able to replicate in cultured mammalian and insect cells, although to a lesser extent than the recombinant wild-type (WT) counterpart. Moreover, rRVFV expressing reporter genes were validated to identify neutralizing antibodies or compounds with antiviral activity. *In vivo*, all mice infected with the reporter-expressing rRVFV displayed an attenuated phenotype, although at different levels. These rRVFV expressing reporter genes provide a novel approach to better understand the biology and pathogenesis of RVFV, and represent an excellent biotechnological tool for developing new therapeutics against RVFV infections.

IMPORTANCE

Rift Valley fever virus (RVFV) is a mosquito-borne virus and zoonotic agent threat that can be deadly to domestic or wild ungulates, and humans. In this work, we used reverse genetics approaches to explore the genome plasticity of RVFV by generating a set of recombinant (r)RVFV that express fluorescent or luminescent proteins to track viral infection. All the generated reporter-expressing rRVFV were able to propagate in mammalian or insect cells, and in a mouse model of infection. Our studies may contribute to advances in research on RVFV and other bunyaviruses and pave the way for the development of novel vaccines and the

identification of new antivirals for the prophylactic and therapeutic treatment,
respectively, of RFVF infections.

KEY WORDS: Rift valley fever virus (RVFV), segment S, reporter gene, NanoLuc
luciferase, Venus fluorescent protein, neutralizing antibody, antiviral, mouse model.

INTRODUCTION

Rift Valley fever (RVF) is a re-emerging zoonotic disease endemic to Africa,
the Middle East, and the Indian Ocean with recurrent outbreaks in domestic or wild
ruminants resulting in spillover into humans (1-3). Rift Valley fever virus (RVFV) is a
phlebovirus belonging to the family *Phenuiviridae* that is transmitted by *Aedes* and
Culex mosquito species, but may also be transmitted through direct contact with
body fluids of infected animals (2). Human infections in endemic regions are usually
asymptomatic or can cause mild illness. However, in some cases, infected persons
develop more severe symptoms that range from ocular lesions to encephalitis and
haemorrhagic fever or death (1). Moreover, RVF disease leads to high abortion rates
in herds and neonatal mortality, making it a significant threat to livestock of affected
regions (4). Due to its virulence, economic impact and lack of efficient vaccines or
antivirals, RVFV is considered a potential biothreat and is listed as a select agent by
both the U.S. Health and Human Services (HHS) and the U.S. Department of
Agriculture (USDA). Moreover, RVFV is included, together with other pathogens, in
the Blueprint list by the World Health Organization (WHO) which are prioritized for
research and development (R&D) of countermeasures (5). Therefore, research to
better understand RVFV pathogenesis and to develop effective prophylactic

vaccines and therapeutic strategies is urgently needed for the control of this

important human viral pathogen.

RVFV virions are icosahedral and contain a lipid envelope enclosing a

nucleocapsid, and the viral genome is formed by three single-stranded RNA

segments with negative and ambisense polarity and different sizes: the large (L),

medium (M), and small (S) (6, 7). The 3′ and 5′ ends of the coding regions contain

the regulatory untranslated regions (UTRs). The L segment encodes the RNA-

dependent RNA polymerase (RdRp) (8, 9). The M segment encodes a polyprotein,

which, upon protease processing, gives the two membrane glycoproteins (Gn and

Gc) that form heterodimers in the viral envelope (10-13), and a nonstructural protein

(NSm) involved in modulation of apoptosis (14). The mRNA of the M segment may

also translate an additional 78 kDa protein via the NSm and Gn open reading frames

(ORFs) (15). The S segment encodes the virus nucleoprotein (N) and the virulence-

associated factor non-structural (NSs) in opposite orientations, separated by an

intergenic region (IGR) (16-18). Together with the viral RNA, the N and RdRp

proteins constitute the viral replication complex. While, the genomic L and M RNAs

serve as templates for the synthesis of their cognate antigenomic RNAs and

mRNAs, the segment S uses an ambisense coding strategy for the expression of N

and NSs proteins in opposite orientation. The genomic S segment serves as a

template for the synthesis of N mRNA and the antigenomic S segment serves as the

template for the synthesis of NSs mRNA.

Reverse genetics methods have provided researchers with a powerful

experimental approach for generating recombinant RVFV (rRVFV), which have

allowed researchers to study important aspects of the biology and replication of

RVFV, identify host factors, study markers of pathogenesis, and generate live-

attenuated vaccines (19-21). The generation of reporter-expressing rRVFV has been described previously and engineered to contain four-segments: an authentic L segment, an S segment that encodes N and enhanced green fluorescent protein (eGFP), and two M-type segments that encode either NSmGn or Gc (22). This rRVFV-4s eGFP was able to reach high titters in mammalian BSR cells, but not in insect C6/36 cells (22). Moreover, the virus was highly attenuated in mice (22). In addition, the generation of rRVFV lacking NSs (ΔNSs) expressing either eGFP or firefly luciferase (FLuc) in place of NSs have also been described (23). The growth curves of recombinant wild-type (WT) and ΔNSs-eGFP or ΔNSs-Fluc were similar in Vero cells, but the growth of the reporter viruses in mice was highly attenuated compared to their WT counterpart (23). Moreover, while the eGFP sequence was well tolerated in the viral genome, FLuc sequence was unstable when expressed by the rRVFV (23).

In this work we used a new experimental strategy to generate replication-competent, reporter-expressing rRVFV in which heterologous genes are inserted into the viral genome without the need for segment duplication or gene deletion. Notably, we used four different approaches that also provided information related to the genome plasticity of RVFV. While numerous reporter genes with different features have been described, fluorescent and bioluminescent proteins are becoming the preferred choice for researchers due to their high sensitivity and the continuous improvement of the technologies associated with their detection (24-28). Therefore, we have generated a set of rRVFV expressing fluorescent (Venus) or luciferase (Nluc) reporter genes. All viruses were characterized *in vitro* and *in vivo* and proven to be a valid method for studying viral replication and propagation or for the identification of antivirals and neutralizing antibodies (NAbs).

RESULTS

Generation of replication-competent rRVFV expressing reporter genes

To generate replication-competent, reporter-expressing rRVFV (strain 56/74), the sequence of Nluc or Venus and the porcine teschovirus 1 (PTV-1) 2A autoproteolytic cleavage site were cloned upstream or downstream of the ORF of the N or NSs proteins in the segment S (**Figure 1**). We chose Nluc due to its small size (~20 kDa), stability, and brightness (24, 29). The fluorescent protein Venus was also selected because of the intensity of the signal and previous experience with other RNA viruses, where the insertion of Venus and Nluc in the genome was well tolerated (30-33). Importantly, the UTRs and the IGR of the S segment were not modified. We then used T7-pol plasmid-based reverse genetics approaches to generate a set of rRVFV expressing Nluc or Venus at different genome positions. Notably, 4 Nluc-expressing rRVFV, covering the 4 strategies used, were recovered, namely Nluc2ANSs, NSs2ANluc, N2ANluc, and Nluc2AN. However, only 3 Venus-expressing rRVFV were rescued: Venus2ANSs, NSs2AVenus, and Venus2AN (**Figure 1**). We were not able to successful rescue the N2AVenus even after multiple attempts. Moreover, a rRVFV WT (rWT) was also generated for comparison.

Growth properties and plaque phenotype of reporter-expressing rRVFV in mammalian cells

We next evaluated the fitness of Nluc-expressing (**Figures 2A and B**) or Venus-expressing (**Figures 2C and D**) rRVFVs in mammalian cells by assessing their growth properties and compare them to those of rWT. To this end, confluent monolayers of Vero E6 cells were infected with the different viruses at the same

multiplicity of infection (MOI; 0.01), and the presence of virus in culture supernatants

was quantified at different times post-infection (p.i).

Replication of Nluc-expressing rRVFVs was delayed and did not reach titters

similar to those of the rWT (**Figure 2A**). Moreover, higher replication levels were

observed when Nluc was inserted next to the UTRs (Nluc2AN and Nluc2ANSs),

instead of next to the IGR (N2ANluc and NSs2ANluc) and when the reporter gene

was fused to the NSs ORF (Nluc2ANSs and NSs2ANluc) instead of the N ORF

(Nluc2AN and N2ANluc) (**Figure 2A**). In addition, Nluc activity in culture

supernatants was evaluated at the same time points (**Figure 2B**). Nluc activity

increased in a time-dependent manner, peaking at 72 h p.i. (the last time point

evaluated in our study), most likely because the cytopathic effect (CPE) caused by

viral infection resulted in the release and accumulation of Nluc in the culture

supernatants. Interestingly, viruses expressing Nluc in the same ORF that the viral N

gene (Nluc2AN and N2ANluc) displayed higher levels of Nluc activity (**Figure 2B**).

This is most likely because of the higher levels of N expression during viral infection

as compared to NSs (34).

Similarly, the viral growth of Venus-expressing rRVFVs was delayed as

compared with that of rWT, and the replication of Venus2AN was more affected than

the replication of Venus2ANSs (**Figures 2C and D**). Venus expression was detected

as early as 24 h p.i. and steadily increased, in a time-dependent manner, until 72 h

p.i., when visualization was reduced due to a virus-induced CPE. Importantly, all

reporter-expressing rRVFVs were still able to reach high viral titters in culture

supernatants ($\sim 10^{5/6}$ PFU/ml) but in all cases were reduced as compared to rWT

($\sim 10^{7}$ PFU/ml) (**Figure 2**). Our data suggest that the insertion of Nluc in the viral

genome is better tolerated than the insertion of Venus. In addition, our results also

provide some clues regarding the regions of RVFV segment S that can be more

suitable for the expression of foreign sequences. Therefore, results indicate that

reporter-expressing rRVFVs may be used to track and quantify the dynamics of viral

infection in vitro by using either fluorescent or luciferase reporter genes.

We also evaluated the plaque phenotype of the rRVFVs in Vero E6 and

compared it to the rWT virus (**Figure 3**). For that, Vero E6 cells were infected and at

72 h p.i., immunostaining using specific antibodies against Nluc or GFP (for Venus

detection), and crystal violet staining. As expected, we detected Nluc or Venus-

positive viral plaques only in cells infected with Nluc- or Venus-expressing rRVFVs,

respectively. In addition, all the Nluc- or Venus-positive plaques colocalized with the

plaques detected by crystal violet staining, confirming that all the rRVFV-infected

plaques expressed the Nluc or Venus reporter gene. Moreover, plaque sizes were in

agreement with viral replication kinetics (**Figure 2**), since the sizes of the viral foci

produced by reporter-expressing rRVFVs were smaller than those of the produced

by the rWT virus (**Figure 3**), most likely due to the effect of the reporter gene on viral

fitness. Overall, these data and our previous studies with other reporter-expressing

replication-competent viruses (30, 32, 35, 36) demonstrate that Nluc and Venus can

be successfully expressing from the viral genome using the PTV-1 2A site, and

reporter gene expression can be used as valid surrogates to monitor viral infection

without the need for secondary approaches to detect the presence of the virus in

infected cells, or the presence of virus in culture supernatants.

In vitro **characterization of reporter-expressing rRVFVs**

To assess the expression of Nluc and Venus, confluent monolayers of Vero

E6 cells were mock-infected or infected with rWT and Nluc-expressing rRVFVs

(**Figure 4A**) or Venus-expressing rRVFVs (**Figure 4B**). Then, the cells were

analyzed by indirect immunofluorescence microscopy using antibodies against

RVFV, Nluc (**Figure 4A**), or GFP (for Venus detection) (**Figure 4B**). As expected,

only rRVFV-infected cells were stained with the anti-RVFV polyclonal antibody (pAb).

Accordingly, the expression of Nluc and Venus was detected only in cells infected

with Nluc- (**Figure 4A**) or Venus-expressing (**Figure 4B**) rRVFVs, respectively.

Interestingly, in the absence of cytoplasmatic Nluc detection, the presence of

characteristic intranuclear NSs filaments was readily detected in both Nluc2ANSs

and NSs2ANluc viruses although with a fluorescence intensity signal much stronger

in the latter. In contrast, the Nluc signal coloc.alizes along the RVFV signal in the

cytoplasmic compartment in cells infected with either N2ANluc or Nluc2AN viruses.

The presence of intranuclear NSs accumulation was barely detectable in both

NSs2AVenus and Venus2ANSs viruses in which both RVFV and Venus signals

colocalize in the cytoplasmic compartment as of Venus2AN virus.

Next, protein expression in Vero E6 cells was evaluated by Western blot at 24

and 48 h p.i. (**Figure 5**). Total cell extracts from either mock-, rWT-, or reporter-

expressing rRVFV-infected Vero E6 cells were tested using antibodies specific for

RVFV, N or NSs, Nluc (**Figure 5A**) or Venus (**Figure 5B**) reporters, and for the

loading control, actin. Western blot analysis revealed specific bands corresponding

to the expected molecular sizes of the viral Gn (54 kDa) and Gc (56 kDa), N and

NSs proteins. The N protein expressed by N2ANluc virus displays a slightly higher

molecular mass as a consequence of the proteolytic mechanism that leaves the

linker and part of the 2A peptide sequence at the C-term of the fusion partner.

Interestingly, this shift is also observed for NSs (NSs2ANluc) although only at 24 h

p.i. In addition, we observed specific bands for Nluc or the fusion protein of Nluc with

NS or N (**Figure 5A**) and for Venus or the fusion protein Venus with NS or N (**Figure**

5A), indicating that the proteolytic efficiency of the 2A peptide sequence used was

not 100%, as previously described (37, 38). Moreover, protein expression was

greater at 48 h p.i. In the case of cell extracts infected with Venus-expressing

rRVFVs, we also observed bands of lower molecular size than expected, which

could indicate Venus degradation.

Intracellular expression of Nluc and Venus in infected cells

To determine whether differences in Nluc activity in cell culture supernatants

were due to differences in the levels of Nluc secretion *versus* its accumulation in

infected cells, confluent monolayers of Vero E6 cells were infected (MOI of 0.01) with

Nluc-expressing viruses, and the presence of virus in culture supernatants was

evaluated at 48 h p.i. (**Figure 6A**). In addition, Nluc activity in the supernatant or in

cell extracts was quantified (**Figure 6B**). There was a correlation between Nluc

activity in the supernatants and that in the cell extracts, suggesting that differences in

Nluc activity were not a consequence of a greater secretion into the supernatant or

cell accumulation of Nluc; indeed, these differences were most likely due to

differences in the Nluc position in the viral genome. However, the observed

intranuclear detection of Nluc aggregates (**Figure 4**), resembling NSs filamentous

structures, might correlate with lower availability of soluble Nluc in cells infected with

the NSs2ANluc expressing virus.

Next, Vero E6 cells infected with Venus-expressing rRVFVs were evaluated

by flow cytometry to quantify the percentage of cells expressing Venus, N protein, or

Venus and N protein at 24 h. p.i, using the anti N monoclonal antibody (mAb)

2B1conjugated to Alexa 674 (**Figure 6C**). Notably, the percentage of Venus+ cells

was greater than the percentage of N+ cells. Moreover, 100% of the Venus+ cells

were also N+. In contrast, and as expected, the rWT infection control was detected

only after N staining. Our data indicate that the expression of Venus during viral

infection could be a more sensitive method to detect the presence of the virus in

infected cells than antigen staining, even one of the higher proteins expressed during

viral infection (e.g. N), at least, at early times post infection.

***In vitro* stability of the reporter-expressing rRVFV**

The genetic stability of reporter-expressing viruses is important for their use

since can limit some of their applications (30, 32). Therefore, we evaluated the *in*

vitro stability of all the reporter-expressing rRVFVs generated (**Figure 7**). To that

end, Nluc- or Venus-expressing rRVFVs were passaged in duplicate three

consecutive times in Vero E6 cells, and the percentage of reporter-expressing

viruses was determined by plaque assay. In the case of Venus-expressing viruses,

plaques were observed directly under a fluorescence microscope. For the

expression of Nluc, plaques were amplified in 96-well plates, and then, Nluc activity

in the cell culture supernatant was evaluated at 48 h.p.i. rRVFVs that express Nluc

or Venus were 100% stable at passage two (virus working stock) and passage 3

(first amplification passage) in all cases. Interestingly, the rRVFVs expressing the

reporter genes close to the IGR (NSs2ANluc, N2ANluc, and NSs2AVenus) were

more stable than the viruses encoding Nluc or Venus next to the UTRs (Nluc2ANSs,

Nluc2AN, and Venus2ANSs), where a gradual loss of Nluc or Venus expression was

observed from passages 5 or 4, respectively (**Figure 7**). Venus2AN was the

exception since it expressed stable fluorescent levels up to 5 passages.

Use of rRVFV expressing Nluc for the identification of antivirals

Identification of compounds that inhibit RVFV normally requires the use of secondary approaches to evaluate viral infection (39-42). The use of reporter-expressing viruses compatible with high-throughput screening for the identification of antivirals may be an interesting alternative to traditional assays. To demonstrate that Nluc-expressing rRVFVs are a valid surrogate for identifying antivirals, we examined the ability of ribavirin to inhibit RVFV infection using the Nluc-expressing rRVFVs NSs2ANluc, Nluc2ANSs, and Nluc2AN (**Figures 8A and B**). We did not include the virus N2ANluc in the assay, because this rRVFV was the most affected in viral replication (**Figure 2**). At high concentrations of ribavirin (starting concentration of 1000 μM), all Nluc-expressing rRVFV were inhibited in a dose-dependent manner, as determined by Nluc activity in cell culture supernatants (**Figure 8A**). Importantly, the IC_{50} values were similar for all the viruses tested (**Figure 8B**) and consistent with those described in the literature (39-43).

Use of rRVFV expressing Nluc for the identification of NAbs

NAbs are a key immunological outcome for the induction of protective immunity after vaccination to prevent RVFV infections. Similar to the previously described antiviral test, viral neutralization assays typically involve the use of secondary methods to measure viral infection. Because the presence of Nluc in the culture supernatants is a valid surrogate of viral replication, an Nluc-based microneutralization assay was developed using the Nluc-expressing rRVFVs. Confluent monolayers of Vero E6 cells were infected with Nluc-expressing rRVFV, which had previously been incubated with sheep sera obtained before or at 14 days post-vaccination with an experimental RVFV vaccine (20) (**Figures 8C and D**).

Then, at 48 h p.i. Nluc activity in the culture supernatants was quantified (**Figure 8C**).

As expected, all viruses were neutralized, in a dose-dependent matter, by sera

collected at 14 days post-vaccination. Using sigmoidal dose-response curves, we

determined the 50% neutralization titter (NT_{50}) obtained by the quantification of Nluc

(**Figure 8D**). Importantly, the NT_{50} was similar for all the tested viruses. These data

indicate that our Nluc-based-microneutralization assays may be a suitable method

for the rapid identification and quantification of antivirals and NAbs.

Growth properties of rRVFV in C6/36 mosquito cells

RVFV is an arbovirus, with mosquitoes playing an important role in the natural

transmission cycle. Therefore, the growth properties of the reporter-expressing

rRVFVs were also investigated in the *Aedes albopictus* clone   mosquito larvae

derived cell line (**Figure 9**). For that, C6/36 cells were infected (MOI; 0.01) with the

different viruses, and the presence of virus in culture supernatants was quantified at

3, 5 and 7 days p.i. As expected, we observed differences in the levels of viral

replication among the rRVFVs. Nluc-expressing rRVFVs replicated to a lesser extent

than the rWT. In addition, as previously observed in Vero E6 cells (**Figure 2**), higher

replication levels were observed when Nluc was cloned in the ORF of the NSs gene

(Nluc2ANSs vs Nluc2AN or NSs2ANluc vs N2ANluc) (**Figure 9A**). Furthermore, the

presence of Nluc in cell culture supernatants was measured at the same time points.

Nluc activity increased in a time-dependent manner, peaking at 7 days p.i. (the last

time point evaluated). Similar to Vero E6 cells, viruses expressing Nluc from the

ORF of the viral N gene displayed also high levels of Nluc activity, most likely

because of the higher levels of N expression during viral infection than NSs (34)

(**Figure 9B**). In contrast to what was detected in Vero E6, the NSs2ANluc virus also

had high Nluc activity level. The viral replication of Venus-expressing rRVFVs was also delayed as compared with that of the rWT (**Figure 9C and D**). Venus expression was detected as early as 3 days p.i. (first time point evaluated) and steadily increased in a time-dependent manner, until 7 days p.i. (last time point evaluated). Therefore, the data indicate that Nluc- and Venus-expressing rRVFVs may also be used to track the dynamics of viral infection in mosquito cells.

Pathogenicity and replication of Nluc-expressing rRVFV in mice

Next, we evaluated the pathogenicity of Nluc-expressing rRVFV in BALB/C mouse model of infection (**Figure 10**). To that end, groups of mice (n = 5) were inoculated intraperitoneally with 10^3, 10^4 or 10^5 PFU of each rRVFV and body weight loss (**Figure 10A**) and survival (**Figure 10B**) were monitored for 18 days. Compared with rWT-infected mice, Nluc-expressing rRVFVs were attenuated. With a lower dose of 10^3 PFU, all mice inoculated with rWT lost weight and all of them succumbed to viral infection. However, only rRVFV NSs2ANluc displayed virulence level comparable to the rWT, and although moderate body weight loss occurred between days 7 and 9, four of the 5 mice died. When animals were infected with 10^4 PFU, only two of five mice inoculated with rWT died, and none of the mice infected with the Nluc-expressing rRVFVs showed signs of morbidity and all of them survived viral infection. Interestingly, with the higher dose of 10^5 PFU, three animals inoculated with rWT and 2 inoculated with rRVFV Nluc2ANSs succumbed to the infection, although without significant body weight loss. The absence of a correlation between the dose and virulence was somehow expected, since this phenomenon has been shown before for RVFV, including in some of our studies (44, 45).

Given that RVFV is isolated from blood and causes viremia, it was speculated

that the presence of Nluc could be detected in blood samples from infected mice, as

occurs with Nluc-expressing bluetongue virus, which is also isolated from blood

samples of infected mice (32). To prove this, Nluc activity was measured in plasma

samples from infected animals and days 3 and 5 p.i. However, Nluc activity was not

detected in any of the plasma samples (data not shown), even presence of the virus

was detected at the same times post-infection by qRT-PCR (**Figure 10C**).

Replication levels were higher at 3 days p.i in all cases and a clear correlation

between mortality and leves of viral RNA in blood was not observed (**Figure 10C**).

To study seroconversion in the animals, serum samples were collected at 18 d p.i,

and the presence of antibodies against RVFV N was analyzed by ELISA (**Figure**

10D). The presence of anti-N antibodies is typically used as a sign of infection and

replication and as a valid serological diagnostic method (20, 46-49). We observed

seroconversion in all the mice that survived infection (**Figure 10D**), suggesting that

Nluc-expressing rRVFVs were able to replicate, although they are mostly attenuated

in vivo. Altogether, these data indicate that the our expression strategy of Nluc from

the segment S contributes to in vivo attenuation depending on the insertion position,

at least in the mouse model of infection used.

DISCUSSION

The establishment of reverse genetics systems for RVFV and other

bunyaviruses has allowed researchers to investigate virus replication, mechanisms

of viral pathogenesis, and the generation of live-attenuated vaccines (7, 17, 19, 47,

50, 51). Reporter-expressing, replication-competent rRVFV represents a powerful

tool for basic and/or translational studies (9, 19, 22, 23). Previously, reporter-

expressing rRVFVs were generated using a four segment strategy or by substituting

the viral NSs sequence with the reporter gene (22, 23). In this work, we have used

four new different strategies to insert a reporter gene in the viral segment S, avoiding

duplication of viral segments or removal of non-structural viral genes; and, therefore,

representing a more biological approach than previous strategies. In this new

approach, foreign sequences were expressed from the same viral N or NSs ORFs

using the PTV-1 2A cleavage site, an approach that has been successfully

implemented for the development of other replication-competent, reporter-expressing

viruses with high success (32, 37, 38). Similar to these previous studies, we have

been able to successfully rescue using this novel 2A approach RVFV expressing

fluorescent and luciferase reporter genes. Based on the different properties of

fluorescent and luciferases proteins (25-28, 30) it is important to carefully considered

the advantages and disadvantages of these reporter genes before performing

experiments. For example, *in vitro*, luciferases may be more appropriate for

quantitative determinations (27, 29, 52). However, fluorescent proteins are a better

choice to observe localization in cells (25, 26). Thus, in this manuscript we used the

same innovative approach to generate rRVFVs expressing luciferase (Nluc) or

fluorescent (Venus) reporter proteins. However, other reporter luciferase or

fluorescent proteins can be inserted into the genome of RVFV using the same

experimental strategy to rescue similar viruses. Moreover, based on the restuls

obtained in this study, we have been able to have a better understanding on the

plasticity of the RVFV genome and segment S to accommodate extra foreign

sequences and how that could results in levels of expression of foreign proteins and

attenuation of recombinant viruses expressing these reporter genes from the locus of

the vial N or NSs ORFs.

Subsequently, the expression of the Nluc and Venus was confirmed, and the

virological phenotypes of Nluc- and Venus-expressing viruses were characterized.

As expected, the infection of mammalian Vero E6 or mosquito C6/36 cells with

reporter-expressing rRVFV was visualized in real time without the need for

secondary methods. Moreover, the expression of Venus and Nluc revealed similar

kinetics that correlated with the levels of viral replication, further demonstrating the

feasibility of using Nluc or Venus as valid surrogates of viral replication and infectivity

without the need of secondary approaches to identify the presence of the virus in

infected cells or cell culture supernatnats, respectively. Compared to rWT, replication

of the reporter-expressing rRVFVs was affected, although, in general, all viruses

growth with high viral titers. Importantly, all the rRVFVs were stable during multiple

passages in cell culture, which is desirable for studies using reporter-expressing

viruses (30, 32, 35). However, we identified interesting differences regarding the

strategy used to insert the reporter genes. Greater expression was observed when

reporter genes were expressed from the locus of the viral N protein. On the other

hand, rRVFV encoding Nluc or Venus from the locus of the NSs protein replicated to

a greater extent but had reduced levels of reporter gene expression. Notably, if

reporter genes were inserted after the UTR sequence, we observed higher viral titers

than thos of recombinant viruses expressing the reporter gene before the UTR

sequence. However, the mechanisms responsible for these differences remain to be

investigated and will be the subject of our follow up studies aimed to assess foreign

gene expression from the RVFV S segment based on their location in the viral

genome.

Notably, we developed and validated a Nluc-based microneutralization assay

that allow us to rapidly identify antiviral compounds and NAbs against RVFV. Our

novel reporter-based mincroneutralization assay represent an excellent option to

interrogate large libraries of compounds or antibodies to identify those with antiviral

and neturalizing activity against RVFV, including their feasibility to implement in HTS

approaches to rapidly and sensitivity screen for novel antivirals or, in the case of

NAbs, the efficacy of novel vaccine candidates. In addition, the virulence of Nluc-

expressing rRVFVs was assessed in BALB/C mice. In this mouse model of infection,

reporter-expressing viruses were attenuated as compared to rWT, and mortality was

only observed in mice inoculated with Nluc2ANSs or NSs2ANluc viruses.

Importantly, all reporter rRVFVs were able to replicate in this mouse model.

However, the reporter rRVFVs could display a more virulent phenotype in other

mouse strains that are more susceptible to RVFV infection (53). It has been shown

that BALB/C mice are slightly more resistant than other mouse strains, where

different disease phenotypes (hepatic, neurological and mixed) can be clearly

distinguished (54, 55). In addition, differences depending on the sex and age of the

animals have been reported, with males being more susceptible than females (this

work) (44, 56). Interestingly, all mice produced anti-N antibodies, suggesting that the

viruses were able to infect and replicate (20, 46-49), although causing limited or no

pathogenicity in terms of body weight loss or mortality.

NSs is localized both in the cytoplasm and nucleus of infected cells and it is

considered the major virulence factor of RVFV. NSs form nuclear filamentous

structures through homo-oligomerization (57) and inhibit host innate immune

responses through different strategies, including inhibition of the type I interferon

(IFN) system (58), and induction of host cellular transcription shut-off (59). It has

been suggested that NSs filament formation is dispensable for efficient replication in

vitro and appears to be important but insufficient for virulence in vivo (60). In two of

our strategies used to generate reporter-expressing rRVFVs, we introduced the

foreign sequence adjacent to the NSs sequence. This could affect some of the

antiviral functions of NSs, including the formation of filamentous structures, and thus

explain the attenuated phenotype in vivo. However, filament like structures

resembling those of NSs were clearly observed in the virus expressing NSs2ANluc

and to a lesser extent in the Nluc2ANSs virus. Interestingly, the detection of

filaments could be accomplished by anti-Nluc antibody, indicating that unprocessed

NSs-Nluc or Nluc-NSs fused proteins are transported to the nucleus of the infected

cells. How this fact correlates with the observed virulence levels in vivo of both

rRVFVs remains to be elucidated.

The formation of the RVFV N-RNA complex protects the vRNA and replication

intermediates from nucleases and is key for viral replication-transcription, and

efficient packaging of the viral genome (61, 62). It has been observed in

crystallographic structures that the N protein adopts two major conformations, open

and closed (61, 63-65). It cannot be discarded that by inserting the reporter gene

next to the viral N sequence, the proper function of the N protein during viral infection

could be affected, due to the intrinsic proteolytic mechanism of the PTV-2A peptide

in which the cleavage site lies between the Gly-Pro bond at the end of the peptide,

therefore 21 extra aminoacids are added to the end of the viral nucleoprotein. This

could explain the observed attenuation in vivo and reduced viral fitness in vitro

compared with those of the rWT or with rRVFVs expressing the reporter gene on the

NSs side.

In addition, although the UTRs and IGR of the viral segment S were not

directly modified in our strategy, the insertion of foreign sequences could introduce

some changes in the RNA structure or unknown cis regulatory sequences of the viral

genomic and/or antigenomic sequences. In fact, it was observed, that reporter-expressing rRVFVs containing the reporter gene next to the 5´ or 3´ UTRs displayed better fitness than viruses in which the heterologous genes were inserted next to the IGR (17, 66-68). Notably, our strategy could also be useful for developing and testing novel live RVFV vaccines or vaccine vectors with different degrees of attenuation.

MATERIALS AND METHODS

Cell lines and viruses

Vero E6 cells (ATCC/CRL/1586) were routinely maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 5% fetal bovine serum (FBS), 1% 100x nonessential amino-acids, 100 U/mL penicillin, 100 µg/mL streptomycin, and 2 mM L glutamine. The cells were incubated at 37°C in the presence of 5% CO_2. BSR-T7 cells (32, 69) stably expressing the bacteriophage T7 RNA polymerase were maintained in the same manner as above except for the addition of 1 mg/mL of Geneticin every 2-3 passages. C6/36 *Aedes albopictus* cells (ATCC CRL–1660) were grown in Eagle's Minimum Essential Medium (MEM) supplemented with 10% fetal calf serum (FCS), L-glutamine (2 mM), gentamicin (50 µg/ml), and MEM vitamin Solution (Sigma) at 28°C.

The sequence of the WT South African RVFV strain 56/74 (parental virus), which was used for the synthesis of the reverse genetic system, was previously described (70) . GenBank: OM744400.1 (segment L), OM744401.1 (segment M) and OM744402.1 (segment S). Virus stocks were propagated in Vero E6 cells and titrated by plaque assay as previously described (20, 46, 47). All the experiments were conducted in the biological security facility of the Animal Health Research Center (CISA, INIA-CSIC).

Plasmids

All pBlueScript II SK (+) reverse genetics (RG) plasmids were synthetized *de novo* by Biomatik and designed to contain the viral cDNAs flanked upstream by a T7 RNA polymerase promoter sequence and downstream by the hepatitis delta virus ribozyme (HDVR) and T7 RNA polymerase terminator sequences (32). To generate replication-competent reporter-expressing RVFVs and study viral genome plasticity, we engineered a set of recombinant modified segments S using four different strategies for the cloning of reporter genes. Importantly, the 5′- and 3′- UTRs and the IGR were not modified. A common heterologous sequence was designed containing a linker (GSG), the porcine teschovirus-1 (PTV-1) 2A autoproteolytic cleavage site (ATNFSLLKQAGDVEENPGP), and a multicloning site (AgeI/BglII/NheI) for the insertion of foreign sequences (30, 32, 36). Then, the 4 master plasmids containing the heterologous sequence before or after the 5′- and 3′- UTRs or the IGR were used for the cloning of the Venus or Nluc reporter genes. Reporter genes were amplified by PCR using specific primers containing AgeI or NheI sites. Then, the PCR products and the 4 modified segment S plasmids were digested, and the reporter genes were cloned using standard molecular biology techniques. Plasmid constructs were confirmed by sequencing (Macrogen). Master plasmids containing the modified segment S

-pBS-2A-N: The reporter gene without a stop codon was cloned between the 3′UTR of the genomic vRNA and the N ORF. This plasmid was used to generate Venus2AN and Nluc2AN viruses.

-pBS-N-2A: The reporter gene with a stop codon was cloned between the N ORF and the IGR. This plasmid was used to generate N2AVenus and N2ANluc viruses.

-pBS-NSs-2A: The reporter gene with a stop codon was cloned between the NSs

ORF and the IGR. This plasmid was used to generate NSs2AVenus and NSs2ANluc

viruses.

-pBS-2A-NSs: The reporter gene without a stop codon was cloned between the

5´UTR of the genomic vRNA and the NSs ORF. This plasmid was used to generate

Venus2ANSs and Nluc2ANSs viruses.

Rescue of recombinant viruses

To rescue rRVFV, monolayers of BSR-T7 cells (T25 flasks, 80% of

confluence, duplicates) were cotransfected with 1 μg of each of the reverse genetics

plasmids encoding the RVFV cDNAs using Lipofectamine 3000 (Invitrogen) in Opti-

MEM I reduced serum medium (Invitrogen) according to the manufacturer's

instructions. Twenty four hours after transfection, the transfection mixture was

replaced with 5 mL of growth medium. After 72-96 h, the cell culture supernatants

were collected, clarified, and used to infect fresh monolayers of Vero E6 cells. Once

the CPE was observed (48-96 h), the medium was harvested and clarified. Clonal

rRVFV were selected by plaque assays, and virus stocks were propagated in Vero

E6 cells.

Virus growth kinetics

To evaluate virus multicycle growth kinetics, confluent Vero E6 cells (12-well

plate format, 5×10^5 cells/well, triplicates) or C6/36 cells (T25 flasks, 4×10^6

cells/well, triplicates) were infected at an MOI of 0.01. After 1 h of virus adsorption at

37°C (Vero E6) or 28ºC (C6/36), the inoculum was discarded, the cells were

washed, and 1.5 mL (Vero E6) or 5 mL (C6/36) of growth medium was added. At the

indicated times post infection (24, 48, and 72 h for Vero E6; 3, 5, and 7 days for

C6/36), 150 µL of the supernatant was removed and stored at -80°C. Virus titers

were determined in Vero E6 cells by plaque assay (plaque-forming units, PFU/mL),

as previously described (20, 46, 47, 70). The mean value and standard deviation

(SD) were calculated using GraphPad Prism version 8.0.1 (GraphPad Software Inc.).

In addition, cells infected with Venus-expressing rRVFV were imaged using a Zeiss

Axio fluorescence microscope (Zeiss), and the presence of Nluc in the culture

supernatants was quantified using Nano-Glo luciferase substrate (Promega)

following the manufacturer's specifications.

Plaque assay and immunostaining

Confluent monolayers of Vero E6 cells (6-well plate format, 10^6 cells/well)

were infected with the indicated viruses for 1 h at room temperature, overlaid with

agar, and incubated at 37°C. At 3 days p.i., the cells were fixed with 10%

paraformaldehyde (PFA), and the overlays were removed. The cells were then

permeabilized (0.01% Triton X-100 in PBS) for 15 min at room temperature, and

prepared for immunostaining as previously described (30). Primary monoclonal or

polyclonal antibodies (mAbs and pAbs, respectively) antibodies against Nluc (mouse

mAb; R&D Systems) for Nluc-expressing viruses, GFP (goat pAb, Rockland) for

Venus-expressing viruses, and anti-N (mAb, 2B1) (51) were used. The Vectastain

ABC kit and DAB HRP substrate kit; (Vector) were used for the immunostaining

following the manufacturers' specifications. The plates were stained with crystal

violet after the immunostaining.

Protein gel electrophoresis and Western blot analysis

Cell extracts from either mock- or virus-infected (MOI, 0.1) Vero E6 cells were

lysed at 24 or 48 h p.i. in radioimmunoprecipitation assay (RIPA) buffer, and proteins

were separated by denaturing electrophoresis. The membranes were blocked for 1 h

with 5% dried skim milk in TBS containing 0.1% Tween 20 (T-TBS) and incubated

overnight at 4°C with specific primary mAbs and pAbs: Nluc (mouse mAb; R&D

Systems) for Nluc-expressing viruses, GFP (goat pAb, Rockland) for Venus-

expressing viruses, NSs (pool of mAb; 5C3A1B2, 2A10B1B and 5F12C1B3) kindly

provided by M. Groschup (50), and anti-RVFV (rabbit pAb generated in the

laboratory) (20). A mAb against actin (Sigma) was used as an internal loading

control. Bound primary antibodies were detected with horseradish peroxidase (HRP)-

conjugated secondary antibodies against the different species (mouse, rabbit or

goat). Proteins were detected by chemiluminescence (Thermo Fisher Scientific)

following the manufacturer's recommendations and photographed using a Bio-Rad

digital ImageStation.

Indirect immunofluorescence assays

Vero E6 cells were mock infected or infected (MOI, 0.1) with the indicated

rRVFV. At 24 h p.i., the cells were fixed with 4% paraformaldehyde (PFA) and

permeabilized with 0.5% Triton X-100 in PBS for 15 min at room temperature and

blocked with PBS/5% FBS. After blocking, specific primary mAbs or pAbs for Nluc

(mouse mAb; R&D Systems) for Nluc-expressing viruses, GFP (goat pAb, Rockland)

for Venus-expressing viruses, and anti-RVFV (rabbit pAb generated in the

laboratory) (20) were diluted in blocking solution and used to separately detect

protein expression using Alexa Fluor 594-conjugated secondary antibodies

(Invitrogen). Images were taken with a Zeiss Axio fluorescence microscope (Zeiss).

Flow cytometry assays

Vero E6 cells were mock infected or infected (MOI, 0.01) with the indicated rRVFV WT or Venus-expressing viruses. At 24 h p.i., the cells were harvested from the culture, centrifuged for 2 min at 2,000 x g and incubated for 30 min at 4ºC with a mAb against the N (2B1) conjugated to Alexa 700 (Invitrogen). The cells were washed and fixed with fixation buffer (BioLegend) for 20 minutes at room temperature in the dark. The cells were analyzed by flow cytometry by gating on green (Venus) and red fluorescence (mAb anti-N 2B1 Alexa 647) (51) using a CyFlow Cube 8 cytometer (Sysmex). The data were analyzed using FlowJo v10 software.

Stability of reporter-expressing viruses in cultured cells

After generation of virus working stocks (P2), confluent monolayers of Vero E6 cells (24-well plate format, duplicates) were infected (MOI, 0.01) with the reporter expressing viruses and incubated until a 70% CPE was observed. Tissue culture supernatants were then harvested and 1/10 of the collected medium was used for the infection of fresh Vero E6 cells for a total of 3 passages. During each viral passage, cultured supernatants were collected and plaque assays were carried out as described above. Venus-expressing plaques were visualized directly using a Zeiss Axio fluorescence microscope (Zeiss). To determine the stability of Nluc-expressing viruses, plaques were used to infect fresh Vero E6 cells (96-well plate format, 2.5×10^4 cells/well), and Nluc activity in the tissue culture supernatants was determined at 48 h.p.i., as indicated above. Approximately 30 plaques per passage were evaluated for each virus.

Reporter-based microneutralization assays

Nluc microneutralization assays were performed as previously described (30-32). Pre- and post-challenge Sheep sera from a previous experiment carried out in the laboratory were serially diluted 2-fold (starting dilution, 1/20). Then, 200 PFU of virus was added to the antibody dilutions, and the mixture was incubated for 1 h at 37ºC. Vero E6 cells (96-well plate format, 5×10^4 cells/well, triplicates) were then infected with the serum-virus mixture for 1 h at 37ºC. Then, the serum-virus mixture was removed and 150 µl of fresh growth medium was added and incubated for 48 h at 37°C. The Nluc activity in the culture supernatants was quantified using Nano-Glo luciferase substrate (Promega) and a microplate reader as described previously. The luciferase values of virus-infected cells in the absence of serum were used to calculate 100% viral infection. Cells in the absence of viral infection were used to calculate the luminescence background. Triplicate wells were used to calculate the average and SD of neutralization. The 50% neutralization titter (NT_{50}) was determined by use of a sigmoidal dose-response curve (GraphPad Prism, v8.0.1 software).

Antiviral assays

Confluent monolayers of Vero E6 cells (96-well plate format, 5×10^4 cells/well, triplicates) were infected with 200 PFU of the Nluc-reporter rRVFVs. After 1 h of infection, the fresh medium was supplemented with 2-fold serial dilutions (starting concentration, 1000 µM) of ribavirin (TCI), and the cells were incubated for 48 h at 37°C. Then, Nluc expression was quantified as indicated above. Triplicate wells were used to calculate the mean and SD of inhibition. The 50% inhibitory concentration

(IC_{50}) was determined by use of a sigmoidal dose-response curve (GraphPad Prism,

v8.0.1 software).

Mouse experiments

The animals were housed under pathogen-free conditions at the biosafety

level 3 (BSL3) animal facility of the Animal Health Research Center (CISA-

INIA/CSIC), Madrid (Spain). Animal experimental protocols were approved by the

Ethical Review Committee at the CISA-INIA/CSIC and Comunidad de Madrid (Permit

number: PROEX079.6/22), in strict accordance with EU guidelines 2010/63/UE

about the protection of animals used for experimentation and the Spanish Animal

Welfare Act 32/2007. For pathogenicity studies, cohorts of 8-to-12-week-old BALB/C

female mice (N = 5/group) were intraperitoneally inoculated with the indicated

viruses at the indicated dose. Then, the animals were monitored daily for 18 days for

clinical symptoms (such as malaise, respiratory distress, and lack of movement),

body weight loss relative to the initial weight, and mortality. At 72 or 96 h after

infection, blood samples were taken by submandibular puncture and tested for viral

RNA by RT-qPCR (51, 71) to monitor viremia. To assess the presence of Nluc in

blood samples, plasma from 50 µL of the same whole blood sample was obtained,

and the Nluc activity was measured using a Nano-Glo Luciferase Assay Kit

(Promega) and a microplate reader (32). Serum samples to be used in antibody

assays were collected at 18 d.p.i. (at the end of the experiment) by submandibular

puncture.

ELISA

Antibodies against N were detected by an in-house ELISA (46). Briefly, for the

detection of antibodies against N, sera collected at 18 d.p.i were tested (Dilution

1/100) in ELISA plates adsorbed with 50 ng/well of purified recombinant N protein

produced from *Escherichia coli* and diluted in carbonate buffer (pH 9.6). The wells

were blocked with 5% BSA in PBS and 0.05% Tween 20 (PBS-T). Then, the bound

antibodies were detected with conjugated goat anti-mouse-IgG-HRP, and the bound

conjugates were detected using TMB for 10 min, followed by the addition of one

volume of stopping solution (3N H_2SO_4). Optical densities were measured at 450 nm

(OD450).

Statistical analysis

Microsoft Excel (Microsoft Corporation) and GraphPad Prism version 10.1.2

(GraphPad Software, San Diego, CA, USA) software were used to analyze the data.

Microsoft Excel was used to perform some of the calculations and to visualize the

raw data. GraphPad Prism software was used to perform statistical analysis of all the

data. Growth kinetics were analyzed using two-way ANOVA the Dunnett's multiple

comparisons test. A one-way ANOVA with Tukey's method was used to evaluate

viral titters comparing between groups. To evaluate the Nluc activity between the

different groups, two-way ANOVA by Tukey's method was used. Statistical

differences between survival curves were analyzed using log-rank test. Viremia

levels were analyzed using the Dunnett's method. A P value lower than 0.05 was

considered significant in all cases.

ACKNOWLEDGEMENTS

This study was partially funded by the Ramon y Cajal Incorporation grant

(RYC-2017) from the Spanish Ministry of Science, Innovation, and Universities and

the European Union´s Horizon Europe Research and Innovation Programme (grant

agreement 101046133), and by grant PID2021-122567OB-I00 funded by MCIN/ AEI

/10.13039/501100011033/ and by ERDF "A way of making Europe".

FIGURES

Figure 1. Schematic representation of the RVFV segment S. RVFV segment S

and its viral products are indicated by blue (N) or gray (NSs) boxes. The black and

striped red boxes denote untranslated regions and intergenic regions, respectively.

The sequences of the reporter and 2A are indicated in green and red boxes,

respectively.

Figure 2. Multicycle growth kinetics. Vero E6 cells (12-well plates, triplicates)

were infected (MOI of 0.01) with rRVFV WT (rWT) or with recombinant viruses

expressing Nluc (**A and B**) or Venus (**C and D**). At the indicated h.p.i. (24, 48 and

72), viral titers in culture supernatants were evaluated by plaque assay (PFU/ml),

and the same rWT representation was used in A and D (A and D). Comparison of

the replication curves of viruses expressing Nluc or Venus versus rWT at different

post-infection times revealed statistically significant differences (Dunnett's method). *

$P < 0.05$; ** $P < 0.0021$; *** $P < 0.0002$; **** $P < 0.0001$. The data represent the

means ± SDs of triplicate samples. At the same time, post-infection Nluc activity was

evaluated (RLU: Relative Light Units) (**B**), and Venus expression was analyzed using

a fluorescence microscope (**C**). Representative images are shown. Scale bars,

100 µm.

Figure 3. Plaque assays. Vero E6 cells (10^6 cells/well, 6-well plate format) were infected with rRVFV WT (rWT) or with recombinant viruses expressing Venus (top) or Nluc (bottom) and incubated at 37ºC for 3 days. Plaques were evaluated by immunostaining with the mAb anti-N (2B1), the goat pAb anti-GFP (top) or the mAb anti-Nluc (bottom). Immunostaining of rWT was performed with the mAb anti-N (2B1) or a combination of the goat pAb anti-GFP and the mAb anti-Nluc.

Figure 4. Analysis of protein expression by IFA. Vero E6 cells were infected with the indicated virus at an MOI of 0.1 or mock infected. At 24 h p.i., the cells were fixed and permeabilized. Protein expression was examined by IFA using a rabbit pAb against RVFV or mAb for Nluc (**A**) or a goat pAb against GFP (**B**). Scale bars, 100 µm.

Figure 5. Analysis of protein expression by WB. Vero E6 cells were infected with Nluc- (**A**) or Venus-expressing (**B**) rRVFV at an MOI of 0.01 or mock infected. At 24 or 48 h p.i, cell extracts were prepared in RIPA buffer. Protein expression was examined by Western blotting using specific antibodies for viral proteins: pAb against RVFV for N and Gn, a pool of mAbs against NSs (after stripping of pAb). In addition, the expression of the reporter genes Nluc (**A**) and Venus (**B**) was evaluated. Actin was used as a loading control. The numbers on the left indicate the molecular size of the protein markers (in kilodaltons).

Figure 6. Nluc activity in tissue culture supernatant *versus* cell extracts. Vero E6 cells (24-well plates, triplicates) were infected (MOI of 0.01) with recombinant

viruses expressing Nluc. At 48 h p.i., viral titers in culture supernatants were

evaluated by plaque assay (PFU/ml). The differences between the groups were

calculated using one-way ANOVA with the Tukey method. * $P < 0.05$; *** $P < 0.0002$;

**** $P < 0.0001$ (**A**). Nluc activity in tissue culture supernatants (TCSs) and cell

extracts was measured. RLU: Relative Light Units. The differences between the

groups were calculated using two-way ANOVA with the Tukey method. * $P < 0.05$;

*** $P < 0.0002$; **** $P < 0.0001$. (**B**). The data represent the means ± SDs of triplicate

samples. (**C**) Flow cytometry analysis. Vero E6 cells were infected (MOI of 0.01) with

recombinant viruses expressing Venus. At 24 h p.i., the cells were harvested from

the culture and stained with a mAb against N conjugated with Alexa 700. The cells

were analyzed by flow cytometry by gating on green (Venus) and red fluorescence

(mAb anti-N 2B1) and the percentage of N+, Venus+ or Venus+/N+ is indicated.

Figure 7. Genetic stability of reporter-expressing RVFVs in vitro. (**A**) After the

generation of virus working stocks (P2), the recombinant viruses were passaged up

to 3 times in Vero E6 cells (P3 to P5). Stability is expressed as the percentage of

plaques that expressed Nluc (**B**) or Venus (**C**) reporter genes in each passage.

Figure 8. A luciferase-based microneutralization assay for assessing RVFV

antivirals and neutralizing antibodies. (**A and B**) **Antiviral.** Vero E6 cells (96-well

plates, 2×10^4 cells/well, triplicates) were infected with 200 PFU of Nluc-expressing

rRVFV and incubated with 2-fold serial dilutions (starting concentration, 1000 µM) of

ribavirin. Inhibition of viral replication was evaluated by quantifying Nluc reporter

expression at 48 h p.i. The differences between the groups were calculated using

two-way ANOVA with the tukey method. * $P < 0.05$; *** $P < 0.0002$; **** $P < 0.0001$.

(**A**). The percent inhibition (IC_{50}) was calculated using sigmoidal dose-response curves (**B**). As an internal control, infected cells were not treated with antivirals. The percentage of viral inhibition was normalized to that with infection in the absence of antivirals. The data are presented as the means ± SDs of the results determined in triplicate. (**C and D**) **Neutralizing antibodies.** Two hundred PFU of Nluc-expressing rRVFV were pre-incubated with 2-fold serial dilutions (starting dilution, 1:20) of sheep sera collected before or after vaccination for 1 h. Subsequently, Vero E6 cells (96-well plates, 2×10^4 cells/well, triplicates) were infected with the sera-virus mixture. (**A**) Virus neutralization was determined by quantitating Nluc reporter expression at 48 h p.i. (**B**) The percent neutralization (NT_{50}) was calculated using sigmoidal dose-response curves. Infected cells in the absence of serum and mock-infected cells were used as internal controls. The data are presented as the means ± SDs of the results determined in triplicate.

Figure 9. Multicycle growth kinetics in insect cells. C6/36 mosquito cells (T25 flasks, triplicates) were infected (MOI of 0.01) with rRVFV WT (rWT) or with recombinant viruses expressing Nluc (**A and B**) or Venus (**C and D**). At the indicated days p.i. (3, 5, and 7), viral titers in culture supernatants were evaluated by plaque assay (PFU/ml), and the same rWT representation was used in A and C. Dunnett's method was used to statistically compare viruses expressing Nluc or Venus versus rWT. * $P < 0.05$; ** $P < 0.0021$; *** $P < 0.0002$; **** $P < 0.0001$ (**A and C**). The data represent the means ± SDs of triplicate samples. At the same time, p.i. Nluc activity (RLU: Relative Light Units) was measured in (**B**), and Venus expression was evaluated using a fluorescence microscope (**D**). Representative images are shown. Scale bars, 100 µm.

Figure 10. In vivo virulence of Nluc-expressing rRVFV. Groups of 8-to-12-week-

old BALB/C female mice (N= 5/group) were infected with 10^3, 10^4 or 10^5 PFU/mouse

of the indicated viruses: rWT or the Nluc-expressing viruses. Weight loss (**A**) and

survival (**B**) were evaluated daily for 18 days. The statistical differences between the

curves of recombinant viruses expressing NLuc versus rWT were calculated using

the log-rank test, with a P-value < 0.05. (**C**) Viremia determined by RT-qPCR in

EDTA-treated blood samples collected at day 3 and 5 p.i. Cutoff: Cq ≥ 37 (dotted

grey line). Points represent individual Cq value for each mouse, and lines of the

corresponding colour represent the mean Cq value of each group. Dunnett's method

was used to statistically compare viruses expressing Nluc versus rWT. * P < 0.05; **

P < 0.0021; *** P < 0.0002; **** P < 0.0001. (**D**) Antibody responses in surviving

mice were evaluated by ELISA (using purified N protein) on day 18 p.i., to confirm

seroconversion. Mouse sera were diluted 1:100.

REFERENCES

1.   Mroz C, Gwida M, El-Ashker M, El-Diasty M, El-Beskawy M, Ziegler U, et al. Seroprevalence of Rift Valley fever virus in livestock during inter-epidemic period in Egypt, 2014/15. BMC Vet Res. 2017;13(1):87.

2.   Samy AM, Peterson AT, Hall M. Phylogeography of Rift Valley Fever Virus in Africa and the Arabian Peninsula. PLoS Negl Trop Dis. 2017;11(1):e0005226.

3.   Sang R, Arum S, Chepkorir E, Mosomtai G, Tigoi C, Sigei F, et al. Distribution and abundance of key vectors of Rift Valley fever and other arboviruses in two ecologically distinct counties in Kenya. PLoS Negl Trop Dis. 2017;11(2):e0005341.

4.   Fawzy M, Helmy YA. The One Health Approach is Necessary for the Control of Rift Valley Fever Infections in Egypt: A Comprehensive Review. Viruses. 2019;11(2).

5.   Jonkmans N, D'Acremont V, Flahault A. Scoping future outbreaks: a scoping review on the outbreak prediction of the WHO Blueprint list of priority diseases. BMJ Glob Health. 2021;6(9).

6.   Grobbelaar AA, Weyer J, Leman PA, Kemp A, Paweska JT, Swanepoel R. Molecular epidemiology of Rift Valley fever virus. Emerg Infect Dis. 2011;17(12):2270-6.

7.   Terasaki K, Murakami S, Lokugamage KG, Makino S. Mechanism of tripartite RNA genome packaging in Rift Valley fever virus. Proc Natl Acad Sci U S A. 2011;108(2):804-9.

8.   Muller R, Argentini C, Bouloy M, Prehaud C, Bishop DH. Completion of the genome sequence of Rift Valley fever phlebovirus indicates that the L RNA is negative sense and codes for a putative transcriptase-replicase [corrected]. Nucleic Acids Res. 1991;19(19):5433.

9.   Muller R, Poch O, Delarue M, Bishop DH, Bouloy M. Rift Valley fever virus L segment:
correction of the sequence and possible functional role of newly identified regions conserved in
RNA-dependent polymerases. J Gen Virol. 1994;75 (Pt 6):1345-52.
10.   Huiskonen JT, Overby AK, Weber F, Grunewald K. Electron cryo-microscopy and single-
particle averaging of Rift Valley fever virus: evidence for GN-GC glycoprotein heterodimers. J Virol.
2009;83(8):3762-9.
11.   Rusu M, Bonneau R, Holbrook MR, Watowich SJ, Birmanns S, Wriggers W, et al. An assembly
model of rift valley Fever virus. Front Microbiol. 2012;3:254.
12.   Sherman MB, Freiberg AN, Holbrook MR, Watowich SJ. Single-particle cryo-electron
microscopy of Rift Valley fever virus. Virology. 2009;387(1):11-5.
13.   Collett MS, Purchio AF, Keegan K, Frazier S, Hays W, Anderson DK, et al. Complete nucleotide
sequence of the M RNA segment of Rift Valley fever virus. Virology. 1985;144(1):228-45.
14.   Won S, Ikegami T, Peters CJ, Makino S. NSm protein of Rift Valley fever virus suppresses
virus-induced apoptosis. J Virol. 2007;81(24):13335-45.
15.   Kreher F, Tamietti C, Gommet C, Guillemot L, Ermonval M, Failloux AB, et al. The Rift Valley
fever accessory proteins NSm and P78/NSm-GN are distinct determinants of virus propagation in
vertebrate and invertebrate hosts. Emerg Microbes Infect. 2014;3(10):e71.
16.   Giorgi C, Accardi L, Nicoletti L, Gro MC, Takehara K, Hilditch C, et al. Sequences and coding
strategies of the S RNAs of Toscana and Rift Valley fever viruses compared to those of Punta Toro,
Sicilian Sandfly fever, and Uukuniemi viruses. Virology. 1991;180(2):738-53.
17.   Ikegami T, Won S, Peters CJ, Makino S. Characterization of Rift Valley fever virus
transcriptional terminations. J Virol. 2007;81(16):8421-38.
18.   Lara E, Billecocq A, Leger P, Bouloy M. Characterization of wild-type and alternate
transcription termination signals in the Rift Valley fever virus genome. J Virol. 2011;85(23):12134-45.
19.   Tercero B, Makino S. Reverse Genetics System for Rift Valley Fever Virus. Methods Mol Biol.
2024;2733:101-13.
20.   Moreno S, Lorenzo G, Lopez-Valinas A, de la Losa N, Alonso C, Charro E, et al. Safety and
Efficacy upon Infection in Sheep with Rift Valley Fever Virus ZH548-rA2, a Triple Mutant Rescued
Virus. Viruses. 2024;16(1).
21.   Calvo-Pinilla E, Marin-Lopez A, Moreno S, Lorenzo G, Utrilla-Trigo S, Jimenez-Cabello L, et al.
A protective bivalent vaccine against Rift Valley fever and bluetongue. NPJ Vaccines. 2020;5(1):70.
22.   Wichgers Schreur PJ, Oreshkova N, Moormann RJ, Kortekaas J. Creation of Rift Valley fever
viruses with four-segmented genomes reveals flexibility in bunyavirus genome packaging. J Virol.
2014;88(18):10883-93.
23.   Billecocq A, Gauliard N, Le May N, Elliott RM, Flick R, Bouloy M. RNA polymerase I-mediated
expression of viral RNA for the rescue of infectious virulent and avirulent Rift Valley fever viruses.
Virology. 2008;378(2):377-84.
24.   Hall MP, Unch J, Binkowski BF, Valley MP, Butler BL, Wood MG, et al. Engineered luciferase
reporter from a deep sea shrimp utilizing a novel imidazopyrazinone substrate. ACS Chem Biol.
2012;7(11):1848-57.
25.   Shaner NC, Patterson GH, Davidson MW. Advances in fluorescent protein technology. J Cell
Sci. 2007;120(Pt 24):4247-60.
26.   Shaner NC, Steinbach PA, Tsien RY. A guide to choosing fluorescent proteins. Nat Methods.
2005;2(12):905-9.
27.   Welsh DK, Noguchi T. Cellular bioluminescence imaging. Cold Spring Harb Protoc.
2012;2012(8).
28.   Zhao H, Doyle TC, Coquoz O, Kalish F, Rice BW, Contag CH. Emission spectra of
bioluminescent reporters and interaction with mammalian tissue determine the sensitivity of
detection in vivo. J Biomed Opt. 2005;10(4):41210.
29.   Stacer AC, Nyati S, Moudgil P, Iyengar R, Luker KE, Rehemtulla A, et al. NanoLuc reporter for
dual luciferase imaging in living animals. Mol Imaging. 2013;12(7):1-13.

30.   Nogales A, Avila-Perez G, Rangel-Moreno J, Chiem K, DeDiego ML, Martinez-Sobrido L. A
Novel Fluorescent and Bioluminescent Bireporter Influenza A Virus To Evaluate Viral Infections. J
Virol. 2019;93(10).

31.   Chiem K, Nogales A, Lorenzo M, Morales Vasquez D, Xiang Y, Gupta YK, et al. Identification of
In Vitro Inhibitors of Monkeypox Replication. Microbiol Spectr. 2023;11(4):e0474522.

32.   Utrilla-Trigo S, Jimenez-Cabello L, Marin-Lopez A, Illescas-Amo M, Andres G, Calvo-Pinilla E,
et al. Engineering recombinant replication-competent bluetongue viruses expressing reporter genes
for in vitro and non-invasive in vivo studies. Microbiol Spectr. 2024;12(3):e0249323.

33.   Chiem K, Park JG, Morales Vasquez D, Plemper RK, Torrelles JB, Kobie JJ, et al. Monitoring
SARS-CoV-2 Infection Using a Double Reporter-Expressing Virus. Microbiol Spectr.
2022;10(5):e0237922.

34.   Brennan B, Welch SR, Elliott RM. The consequences of reconfiguring the ambisense S
genome segment of Rift Valley fever virus on viral replication in mammalian and mosquito cells and
for genome packaging. PLoS Pathog. 2014;10(2):e1003922.

35.   Breen M, Nogales A, Baker SF, Martinez-Sobrido L. Replication-Competent Influenza A
Viruses Expressing Reporter Genes. Viruses. 2016;8(7).

36.   DiPiazza A, Nogales A, Poulton N, Wilson PC, Martinez-Sobrido L, Sant AJ. Pandemic 2009
H1N1 Influenza Venus reporter virus reveals broad diversity of MHC class II-positive antigen-bearing
cells following infection in vivo. Sci Rep. 2017;7(1):10857.

37.   Kim JH, Lee SR, Li LH, Park HJ, Park JH, Lee KY, et al. High cleavage efficiency of a 2A peptide
derived from porcine teschovirus-1 in human cell lines, zebrafish and mice. PLoS One.
2011;6(4):e18556.

38.   Liu Z, Chen O, Wall JBJ, Zheng M, Zhou Y, Wang L, et al. Systematic comparison of 2A
peptides for cloning multi-genes in a polycistronic vector. Sci Rep. 2017;7(1):2193.

39.   Atkins C, Freiberg AN. Recent advances in the development of antiviral therapeutics for Rift
Valley fever virus infection. Future Virol. 2017;12(11):651-65.

40.   Kende M, Lupton HW, Rill WL, Levy HB, Canonico PG. Enhanced therapeutic efficacy of
poly(ICLC) and ribavirin combinations against Rift Valley fever virus infection in mice. Antimicrob
Agents Chemother. 1987;31(7):986-90.

41.   Lang Y, Li Y, Jasperson D, Henningson J, Lee J, Ma J, et al. Identification and evaluation of
antivirals for Rift Valley fever virus. Vet Microbiol. 2019;230:110-6.

42.   Scharton D, Bailey KW, Vest Z, Westover JB, Kumaki Y, Van Wettere A, et al. Favipiravir (T-
705) protects against peracute Rift Valley fever virus infection and reduces delayed-onset neurologic
disease observed with ribavirin treatment. Antiviral Res. 2014;104:84-92.

43.   Geerling E, Murphy V, Mai MC, Stone ET, Casals AG, Hassert M, et al. Metal coordinating
inhibitors of Rift Valley fever virus replication. PLoS One. 2022;17(9):e0274266.

44.   Graham VA, Easterbrook L, Kennedy E, Rayner E, Findlay-Wilson S, Flett L, et al. Pathogenesis
of Rift Valley Fever Virus in a BALB/c Mouse Model Is Affected by Virus Culture Conditions and Sex of
the Animals. Viruses. 2023;15(12).

45.   Alkan CJ-C, E.; Ikegami, T. . Distinct Pathological Changes in Preweaning Mice Infected with
Live-Attenuated Rift Valley Fever Virus Strains. . Viruses. 2024;16(7), 999.

46.   Borrego B, Brun A. A Hyper-Attenuated Variant of Rift Valley Fever Virus Generated by a
Mutagenic Drug (Favipiravir) Unveils Potential Virulence Markers. Front Microbiol. 2020;11:621463.

47.   Borrego B, Moreno S, Lopez-Valinas A, de la Losa N, Weber F, Nunez JI, et al. Identification of
Single Amino Acid Changes in the Rift Valley Fever Virus Polymerase Core Domain Contributing to
Virus Attenuation In Vivo. Front Cell Infect Microbiol. 2022;12:875539.

48.   Crespo Leon F, Gutierrez Diez F, Rodriguez Ferri F, Leon Vizcaino L, Cuello Gijon F, Gimeno EJ,
et al. The translation into Spanish of the OIE Manual of diagnostic tests and vaccines for terrestrial
animals (mammals, birds and bees): problems, solutions and conclusions. Rev Sci Tech.
2005;24(3):1095-104.

49.   Cullinane AA, Garvey M. A review of diagnostic tests recommended by the World
Organisation for Animal Health. Rev Sci Tech Oie. 2021;40(1):75-89.
50.   Borrego B, Moreno S, de la Losa N, Weber F, Brun A. The Change P82L in the Rift Valley
Fever Virus NSs Protein Confers Attenuation in Mice. Viruses. 2021;13(4).
51.   Moreno S, Calvo-Pinilla E, Devignot S, Weber F, Ortego J, Brun A. Recombinant Rift Valley
fever viruses encoding bluetongue virus (BTV) antigens: Immunity and efficacy studies upon a BTV-4
challenge. PLoS Negl Trop Dis. 2020;14(12):e0008942.
52.   Tran V, Moser LA, Poole DS, Mehle A. Highly sensitive real-time in vivo imaging of an
influenza reporter virus reveals dynamics of replication and spread. J Virol. 2013;87(24):13321-9.
53.   Cartwright HN, Barbeau DJ, McElroy AK. Rift Valley Fever Virus Is Lethal in Different Inbred
Mouse Strains Independent of Sex. Front Microbiol. 2020;11:1962.
54.   Cartwright HN, Barbeau DJ, Doyle JD, Klein E, Heise MT, Ferris MT, et al. Genetic diversity of
collaborative cross mice enables identification of novel rift valley fever virus encephalitis model.
PLoS Pathog. 2022;18(7):e1010649.
55.   Lathan R, Simon-Chazottes D, Jouvion G, Godon O, Malissen M, Flamand M, et al. Innate
Immune Basis for Rift Valley Fever Susceptibility in Mouse Models. Sci Rep. 2017;7(1):7096.
56.   Tokuda S, Do Valle TZ, Batista L, Simon-Chazottes D, Guillemot L, Bouloy M, et al. The genetic
basis for susceptibility to Rift Valley fever disease in MBT/Pas mice. Genes Immun. 2015;16(3):206-
12.
57.   Kohl A, di Bartolo V, Bouloy M. The Rift Valley fever virus nonstructural protein NSs is
phosphorylated at serine residues located in casein kinase II consensus motifs in the carboxy-
terminus. Virology. 1999;263(2):517-25.
58.   Le May N, Mansuroglu Z, Leger P, Josse T, Blot G, Billecocq A, et al. A SAP30 complex inhibits
IFN-beta expression in Rift Valley fever virus infected cells. PLoS Pathog. 2008;4(1):e13.
59.   Billecocq A, Spiegel M, Vialat P, Kohl A, Weber F, Bouloy M, et al. NSs protein of Rift Valley
fever virus blocks interferon production by inhibiting host gene transcription. J Virol.
2004;78(18):9798-806.
60.   Li S, Zhu X, Guan Z, Huang W, Zhang Y, Kortekaas J, et al. NSs Filament Formation Is
Important but Not Sufficient for RVFV Virulence In Vivo. Viruses. 2019;11(9).
61.   Hayashi M, Schultz EP, Lanchy JM, Lodmell JS. Time-Resolved Analysis of N-RNA Interactions
during RVFV Infection Shows Qualitative and Quantitative Shifts in RNA Encapsidation and
Packaging. Viruses. 2021;13(12).
62.   Overby AK, Popov VL, Pettersson RF, Neve EP. The cytoplasmic tails of Uukuniemi Virus
(Bunyaviridae) G(N) and G(C) glycoproteins are important for intracellular targeting and the budding
of virus-like particles. J Virol. 2007;81(20):11381-91.
63.   Ferron F, Li Z, Danek EI, Luo D, Wong Y, Coutard B, et al. The hexamer structure of Rift Valley
fever virus nucleoprotein suggests a mechanism for its assembly into ribonucleoprotein complexes.
PLoS Pathog. 2011;7(5):e1002030.
64.   Raymond DD, Piper ME, Gerrard SR, Skiniotis G, Smith JL. Phleboviruses encapsidate their
genomes by sequestering RNA bases. Proc Natl Acad Sci U S A. 2012;109(47):19208-13.
65.   Raymond DD, Piper ME, Gerrard SR, Smith JL. Structure of the Rift Valley fever virus
nucleocapsid protein reveals another architecture for RNA encapsidation. Proc Natl Acad Sci U S A.
2010;107(26):11769-74.
66.   Murakami S, Terasaki K, Narayanan K, Makino S. Roles of the coding and noncoding regions
of rift valley Fever virus RNA genome segments in viral RNA packaging. J Virol. 2012;86(7):4034-9.
67.   Piper ME, Sorenson DR, Gerrard SR. Efficient cellular release of Rift Valley fever virus
requires genomic RNA. PLoS One. 2011;6(3):e18070.
68.   Tercero B, Terasaki K, Narayanan K, Makino S. Mechanistic insight into the efficient
packaging of antigenomic S RNA into Rift Valley fever virus particles. Front Cell Infect Microbiol.
2023;13:1132757.

69.   Buchholz UJ, Finke S, Conzelmann KK. Generation of bovine respiratory syncytial virus (BRSV)
from cDNA: BRSV NS2 is not essential for virus replication in tissue culture, and the human RSV
leader region acts as a functional BRSV genome promoter. J Virol. 1999;73(1):251-9.
70.   Borrego B, de Avila AI, Domingo E, Brun A. Lethal Mutagenesis of Rift Valley Fever Virus
Induced by Favipiravir. Antimicrob Agents Chemother. 2019;63(8).
71.   Mwaengo D, Lorenzo G, Iglesias J, Warigia M, Sang R, Bishop RP, et al. Detection and
identification of Rift Valley fever virus in mosquito vectors by quantitative real-time PCR. Virus Res.
2012;169(1):137-43.

Figure 1

A)

WT

N SSs

1,7 kb

N NSs

N2AReporter

N 2A Reporter SSN

N 2A Reporter NSs

Reporter2AN

Reporter 2A N SSN

Reporter 2A N NSs

NSs2AReporter

N Reporter 2A NSs

N Reporter 2A NSs

Reporter2ANSs

N NSs 2A Reporter

N NSs 2A Reporter

B) Recombinant viruses generated in this study

Nluc2ANSs
NSs2ANluc
N2ANluc
Nluc2AN
Venus2ANSs
NSs2AVenus
Venus2AN

Figure 2

Figure 3

Figure 4

A)

Rows (top to bottom): DAPI, Anti-RVFV, Anti-Nluc, Merge

Columns (left to right): Mock, rWT, Nluc2ANSs, NSs2ANluc, Nluc2AN, N2ANluc

B)

Rows (top to bottom): DAPI, Anti-RVFV, Anti-GFP, Merge

Columns (left to right): Mock, rWT, Venus2ANSs, NSs2AVenus, Venus2AN

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10