Falsifying computational models of endothelial cell network formation through quantitative comparison with *in vitro* models

Tessa M. Vergroesen[1], Vincent Vermeulen[1], Roeland M. H. Merks[1,2]

[1]Institute of Biology, Leiden University, Leiden, The Netherlands

[2]Mathematical Institute, Leiden University, Leiden, The Netherlands

Abstract

During angiogenesis, endothelial cells expand the vasculature by migrating from existing blood

vessels, proliferating and collectively organizing into new capillaries. *In vitro* and *in vivo*

experimentation is instrumental for identifying the molecular players and cell behaviour that regulate

angiogenesis. Alongside experimental work, computational and mathematical models of endothelial

cell network formation have helped to analyse if the current molecular and cellular understanding of

endothelial cell behaviour is sufficient to explain the formation of endothelial cell networks. As input,

the models take (a subset of) the current knowledge or hypotheses of single cell behaviour and capture

it into a dynamical, mathematical description. As output, they predict the multicellular behaviour

following from the actions of many individual cells, *i.e.*, formation of a vascular-like network.

Paradoxically, computational modelling based on different assumptions, *i.e.*, completely different,

sometimes non-intersecting sets of observed single cell behaviour, can reproduce the same

angiogenesis-like multicellular behaviour, making it practically impossible to decide which, if any, of

these models is correct. Here we present dynamical analyses of time-lapses of *in vitro* endothelial cell

network formation experiments and compare these with dynamic analyses of three mathematical

models: (1) the cell elongation model; (2) the contact-inhibited chemotaxis model; and (3) the

mechanical cell-cell communication model. We extract a variety of dynamical characteristics of

endothelial cell network formation using a custom time-lapse video analysis pipeline in ImageJ. We

compare the dynamical network characteristics of the *in vitro* experiments to those of the cellular

networks produced by the computational models. We test the response of the *in silico* dynamical cell

network characteristics to changes in cell density and make related changes in the *in vitro* experiments.

Of the three computational models that we have considered, the cell elongation model best captures

the remodelling phase of *in vitro* endothelial cell network formation. Furthermore, in the *in vitro*

model, the final size and number of lacunae in the network are independent of the initial cell density.

This observation is also reproduced in the cell elongation model, but not in the other two models that

we have considered. Altogether, we present an approach to model validation based on comparisons of

time-resolved data and variations of model conditions.

Keywords: angiogenesis, image analysis, model falsification, endothelial cells, cellular Potts model

Author Summary

Understanding how blood vessels grow and organise into well-structured networks is crucial for many

clinical applications, from wound healing to cancer treatment. The growth of blood vessels, known as

angiogenesis, involves endothelial cells migrating, proliferating, and forming new vascular structures.

Angiogenesis is studied through biological experiments, but mathematical models are also used to test

whether our understanding of cell behaviour is sufficient to explain cell network formation. Models

assuming chemotaxis, elongated cell shape, contact inhibition or mechanical cell-cell communication

often produce similar results despite the large differences in the underlying assumptions, making it

difficult to determine which model is most accurate. In this study, we analysed time-lapse videos of

endothelial cell network formation and compared these with three computational models: (1) the cell

elongation model, (2) the contact-inhibited chemotaxis model, and (3) the mechanical cell-cell

communication model. By examining the dynamics of network formation and testing how they change

with cell density, we found that the cell elongation model best captures key aspects of the real-life cell

network remodelling process. This approach highlights the importance of time-resolved data in

evaluating computational models and provides a framework for refining our understanding of

angiogenesis.

Introduction

During embryogenesis, endothelial precursor cells, angioblasts, migrate and differentiate to form the first blood vessels. This process of blood vessel formation by *de novo* endothelial cell (EC) production, is called vasculogenesis [1]. After vasculogenesis, the primary vascular network is expanded by angiogenesis. Angiogenesis, the formation of new blood vessels from existing ones, is crucial for physiological and pathological mechanisms such as embryonic development, wound healing and tumour growth. Pro-angiogenic factors, including vascular endothelial growth factor (VEGF)-A, stimulate sprouting in ECs, which make up the inner lining of blood vessels [2–4]. The ECs proliferate and migrate into the extracellular environment and form a vascular network that supplies the surrounding tissue with nutrients and oxygen and removes waste products. A key regulatory factor determining the final structure of the new blood vessel network is the composition of the extracellular matrix (ECM), the scaffold of proteins that surrounds the blood vessels and provides structural support to the newly formed multicellular network [5]. Specific integrin binding sites on ECM proteins like fibronectin and laminin allow cells to exert stresses on the ECM and sense the mechanical properties of their environment [6]. Cells also secrete enzymes which digest the ECM, releasing ECM-retained growth factors, including VEGF-A, that stimulate EC migration [7]. Additionally, cell-cell interactions, like Notch signalling and vascular endothelium cadherin (VE-cadherin) adhesions, regulate individual cell behaviour within the network and separate ECs into leading tip- and proliferating stalk cells [4]. In response to VEGF-A, cells upregulate Delta-like 4 (DLL4), a transmembrane Notch ligand that activates NOTCH1 in neighbouring cells [8]. NOTCH1 activation subsequently downregulates VEGF receptor (VEGFR)-2 expression, reducing the neighbouring cell's susceptibility to VEGF-A, resulting in less angiogenic potential of the cells [8]. Similarly, VE-cadherin binding inhibits VEGFR-2 signalling in neighbouring cells [9]. The combination of extracellular, intercellular and intracellular regulation makes angiogenesis a complex and multiscale process. An understanding of how each of these regulatory mechanisms contribute to the final network formation is necessary for biomedical applications where angiogenesis needs to be stimulated, steered or reduced, like tissue engineering [10] or vascular renormalization [11].

One of the more commonly used *in vitro* models to test the effect of environmental changes on EC

network formation is the tube formation assay (TFA). In this assay ECs are placed on top of a

basement membrane-like substrate, where the ECs, under the right conditions, spontaneously organise

into networks of interconnected tubes [12]. With this assay the influence of various chemical and

mechanical stimuli on EC networks has been investigated by comparing the branch density of the final

EC network between conditions [13]. In the analysis of TFA outcomes it is often assumed that a

higher density of branches indicates a higher angiogenic potential of the cells (*i.e.*, more sprouting).

However, the temporal development of the network over time may contain key information on the

dynamic processes regulating angiogenesis. Merks *et al*. (2006), therefore, measured the density of

branch points in developing vascular networks *in vitro* at regular intervals and observed that the

network density dropped slowly over time [14]. Parsa *et al*. (2011), analysed time-lapse videos of *in*

vitro EC network formation in more detail, and described the development of EC networks as a

carefully orchestrated sequence of five subevents: (1) rearrangement and aggregation, (2) spreading,

(3) elongation and formation of cell–cell contacts, (4) plexus stabilization, and (5) plexus

reorganization [15]. The first two stages describe the attachment of the cells to the substrate directly

after cell seeding. The rate and the extent of cell spreading on the substrate have been linked to

substrate stiffness and composition [6], and are assumed to be influenced by the traction forces cells

are able to exert on the substrate [16]. Later stages in the tube formation assay rely on a combination

of cell-cell and cell-ECM interactions: Rüdiger *et al*. (2020) observed a collapse of EC networks on

soft, laminin matrices, confirming the importance of the ECM as a scaffold to provide support for the

network to stabilise [5].

Computational modelling is a helpful tool for proposing hypotheses and testing whether (a

combination of) proposed developmental mechanisms are sufficient to explain biological observations

[17]. Different computational models of angiogenesis all consider angiogenesis as the assembly of

ECs into a vascular-like network structure [18–20]. For example, in Manoussaki *et al*. (1996), the *in*

vitro matrix remodelling behaviour of ECs on a compliant substrate is described with a continuum

model, in which ECs are described as local densities using a system of differential equations [21].

Similarly, in Serini *et al*. (2003), ECs are modelled to migrate upwards a chemoattractant gradient

with a continuum model [22]. In Palachanis *et al*. (2015), ECs are considered as identical 2D ellipses,

using a single-particle model [23]. Köhn-Luque *et al*., (2013) used a hybrid continuum and cell-based

model to model the effect of VEGF retention in the ECM on EC network formation [7]. In cell-based

models of angiogenesis, ECs are described as discrete areas of variable shapes, where multicellular

behaviour arises from the response of single cells to inputs from their microenvironment [24,25].

Similarly, Vega *et al*. (2020) used a hybrid continuum and cell-based model to model the interaction

between Notch- and VEGF signalling and cell-ECM adhesion [26].

Interestingly, both continuum models, particle-based models, as well as cell-based models are

successful in mimicking EC network formation, and different hypotheses, or model inputs, can in

some cases lead to a qualitatively similar output. In this work we focus on three cell-based models as

previously developed in our group [14,27,28]. The "cell elongation model" assumes that ECs are

attracted towards one another through autocrine/paracrine signalling by secreting a chemoattractant

[14]. In this model it was observed that cells self-organise into vascular-like structures if the cells have

an elongated shape. Observations by Parsa *et al*. (2011) support the necessity for ECs to elongate to

form networks, as cells elongate before they start forming a network and continue to elongate as the

network develops [15]. The "contact inhibition model" also assumes that ECs are attracted to one

another through a secreted chemoattractant, but here it was assumed that VE-cadherin binding

mediates contact inhibition of chemotaxis, following observations by Dejana *et al*. (2004) that VE-

cadherin-VEGFR2 complex forming limits cell proliferation [29] and that in the yolk sac of VE-

cadherin double-knockout mice and in *in vitro* mouse allantois explants ECs failed to form vascular

networks and aggregated into isolated islands [28,30]. In disagreement with VEGF guided cell-cell

attraction-based models, Rüdiger *et al*. (2020) observed persistent network formation in the absence of

a VEGF-A gradient and they argue that the *in vitro* network formation is driven by mechanical

communication [5]. The importance of cell-ECM interactions is further supported by observations by

Stéphanou *et al*. (2007), where they observed faster and more lacunae in network formation on

fibrinogen gels of intermediate rigidity [31]. Finally, the "mechanical cell-cell communication model"

assumes that ECs are attracted to one another solely through mechanical interaction with the ECM

[27]. In this cell-ECM interaction model ECs induce strain in the ECM through contractile forces and

migrate up the strain gradient. With these many models that successfully reproduce multicellular

network formation based on different observed EC behaviour, we need to critically examine these

models to see which assumptions, if any, are correct.

Thus, for a more complete understanding of the integration of different mechanisms driving EC

network formation, we must quantitatively compare these computational models to *in vitro*

experiments by systematically changing model parameters and *in vitro* experimental setup, and

iteratively refining the models based on the outcomes of the comparison. As an initial step, we will

critically examine three models originating from our own research group: (1) The cell elongation

model [14]; (2) the contact inhibition model [28]; and (3) the mechanical cell-cell communication

model [27]. We select model parameters from the original work [14,27,28] and from published global

sensitivity analyses [32], then we fit the models to our own *in vitro* observations, and finally we

present additional parameter examinations. For the comparison, dynamical analyses of *in vitro*

experiments are required. Here we present new dynamic time-lapse videos of tube formation assays

and a new analysis pipeline based on ImageJ, and demonstrate its use for model selection for network

formation. Commercial and open-source image analysis tools are available for quantitative analyses of

EC networks [33–36]. However, these tools are designed for single frame end-point analyses and

therefore less suitable for high-throughput dynamical analyses of large quantities of data. Our novel

image analysis pipeline is suitable to dynamically analyse and compare *in vitro* and simulated

networks. The pipeline allows us to reliably extract network features over time, such as the number of

branches and their length and the average size and number of lacunae. In this study we use this

pipeline to compare the output of three cellular Potts models of 2D EC network formation [14,27,28]

to *in vitro* experiments over time at different cell densities and find that the distance at which cells are

able to communicate with each other determines the features of the final, stabilised network, but the

speed and manner at which the networks stabilize depends on other characteristics of the cell

communication, like the speed at which the cell attracting signal spreads.

Results

Quantification of *in vitro* endothelial cell network dynamics

To understand and compare the dynamic behaviour during *in vitro* and *in silico* EC network formation, we developed an image analysis pipeline to extract network characteristics from 2D EC network formation time-lapses and applied it to *in vitro* and *in silico* videos. To capture *in vitro* EC network formation, ECs were seeded on a Matrigel matrix and imaged in phase contrast for 24 hours with an interval of 15 minutes (Fig 1A). The ECs in the time-lapse images were segmented from the background using a combination of Gaussian denoising and local variance extraction, followed by Huang thresholding (S1 Fig A). Network features were extracted from the segmented time-lapses for each timestep using FIJI plugins Analyze Skeleton [37] and Analyze Particles (Fig 1B). For the *in silico* networks, we selected three cell-based computational models of 2D EC network formation [14,27,28] and extracted the network characteristics directly from the output images using the same image analysis pipeline as the segmented *in vitro* time-lapses.

Fig 1. Image analysis pipeline for *in vitro* phase contrast time-lapses. A) Graphic summary of *in*

vitro experiment. B) Networks were segmented from the background and labelled in FIJI. Insert is 350

x 350 µm². C) ECs were seeded on Matrigel, and network formation is captured in phase contrast for

24 hours with intervals of 15 minutes. Insert is 500 x 500 µm². D-G) Quantitative analysis of (D-F)

lacunae and (G-H) branches under standard conditions. Shaded area represents a standard deviation of

eight wells on one slide.

In the *in vitro* culture, under standard conditions, the ECs started forming cell-cell connections within

the first hour after seeding on a Matrigel matrix and thereafter self-organised into fully connected

networks within six hours (Fig 1C). We were able to divide the development of the network

characteristics over time in three distinct parts, which we describe following Parsa *et al*. (2011) [15],

as: (1) The first hour resembled stages 1 and 2, "rearrangement and aggregation", followed by

"spreading" (Fig 1C). In these stages the ECs were randomly dispersed on top of the Matrigel and

started to form attachments to the ECM. (2) The next 6 hours matched Parsa's stage 3, "elongation and

formation of cell-cell contact". During this stage new connections formed between ECs, which is

reflected in an increase in the number of lacunae from 0.03 ± 0.09 to 5.16 ± 1.05 per mm² (Fig 1D).

(3) Hereafter, the number of lacunae decreased linearly in time ($R^2 = 0.82$, p-value < 0.001),

resembling Parsa's stage 4 and 5: "plexus stabilization" and "reorganization". The average lacuna area

displayed a linear increase due to a higher occurrence of lacuna merging and smaller lacunae closing

compared to the formation of new lacunae (Fig 1E) ($R^2 = 0.89$, p-value < 0.001). The number of

branches within the network continuously decreased and their average length increased (Figs 1G and

H). During the first three stages branches predominantly increased in length due to merging of branch

sprouts, but during stages 4 and 5 we observed an increase in merging of existing branches.

In silico models can be adjusted to resemble *in vitro* networks

To compare the dynamics of *in silico* networks to the *in vitro* networks we selected model parameters

that mimic the *in vitro* experiment. To represent the inner well of the *in vitro* experiments (Fig 1A),

ECs were simulated within a circular field with a diameter of 1900 lattice sites for the two chemotaxis

models (1 lattice site = 2 x 2 µm) and 760 lattice sites for the mechanical model (1 lattice site = 5 x 5

µm) (Fig 2A). To determine the target area and length of the simulated cells, we measured cell areas

and diameters in fluorescently labelled ECs within network time-lapses (S2 Fig A). We observed an

average cell area of 1052 ± 596 µm² after three hours (S2 Fig B), with a cell length of 90 ± 21 µm

for elongated cells (S2 Fig C). Based on these measurements we set the target area for the simulated

cells to 250 lattice sites (chemotaxis models) and 40 lattice sites (mechanical model), which

corresponds to a physical area of 1000 µm². In the cell elongation model, we set the target length to 45

lattice sites, which corresponds to 90 µm.

Fig 2. Quantitative comparisons between two chemotaxis-based models and a mechanical model

of 2D EC network formation to *in vitro* time-lapse results. A) The model parameters are adjusted to

resemble the *in vitro* situation and network characteristics are extracted using the same image analysis

pipeline as the *in vitro* networks. B) Simulated ECs were placed on a circular grid and run for 2880

Monte Carlo steps (MCS). In the chemotaxis models ECs (red) excrete a chemoattractant (grey) that

diffuses through the medium (white). In the mechanical model ECs (white) exert strain on their

environment (green/blue heatmap) (Field area is 500 x 500 μm²). C) A comparison of the number of

lacunae per mm² for chemoattractant diffusion lengths ranging from 10 μm to 100 μm for the cell

elongation model and the contact inhibition model after 2880 MCS. In magenta the *in vitro* number of

lacunae per mm² after 24 hours. Inserts show the cell elongation model for a diffusion length of 10, 50

and 100 μm. D-G) Quantitative analysis of (D, E) lacunae and (F, G) branches *in vitro* (magenta) and

in the cell elongation model (blue), the contact inhibition model (golden) and the mechanical model

(green). H) The number of sprouts, branches connected to a single node *in vitro*, in the cell elongation

model, the contact inhibition model and the mechanical model. ($D = 5.0 \cdot 10^{-13} \mathrm{m}^2\mathrm{s}^{-1}; \epsilon = 1.02 \cdot$

$10^{-4}\mathrm{s}^{-1}; \alpha = 1 \cdot 10^{-3}\mathrm{s}^{-1}$). Shaded area represents a standard deviation of eight simulations or eight

in vitro time-lapses.

In the cell elongation model, ECs elongated towards a target length, secreted a chemoattractant and

migrated upwards the chemoattractant gradient [14]. In the contact inhibition model ECs secreted a

chemoattractant and migrated upwards the chemoattractant gradient, but locally inhibited chemotaxis

in their neighbours (Fig 2A) [28]. To find the parameters that describe the chemoattractant in the two

chemotaxis models for which the models most resemble the *in vitro* networks (*i.e.*, the diffusion

coefficient, the decay rate and the secretion rate) we investigated the effect of the diffusion length of

the chemoattractant on the network formation. Diffusion length (l_D) depends on the diffusion

coefficient (D) and the decay rate (ϵ) of the chemoattractant according to $l_D = \sqrt{\frac{D}{\epsilon}}$. Therefore, we

altered the diffusion length from $l_D = 10$ μm to $l_D = 100$ μm with steps of 10 μm by varying the

decay rate of the chemoattractant, while keeping the diffusion coefficient constant at $D = 5 \cdot$

$10^{-13}\mathrm{m}^2\mathrm{s}^{-1}$. (Fig 2C). For a full list of the selected parameters, see S6 Table. *In vitro*, the average

number of lacunae per mm² after 24 hours was 3.30 ± 0.83, with an average lacuna area of 0.18 ± 0.04

mm² (Fig 1D and E). When we increased the diffusion length in the two chemotaxis model, we

observed that the average number of lacunae decreased (Fig 2C), in agreement with previous work

[14,28,32].

Based on the outcome of the comparison of the number of lacunae (Fig 2C) and branches (S2 Fig D)

per mm², we selected a decay rate of $\epsilon = 1.02 \cdot 10^{-4} s^{-1}$, which corresponds to a diffusion length of

$l_D = 70$ µm. This resulted in an average number of lacunae per mm² after 2880 MCS of 3.27 ± 0.47,

with an average area of 0.15 ± 0.02 mm² for the cell elongation model and 4.88 ± 0.35 lacunae per

mm², with an average area of 0.10 ± 0.01 mm² for the contact inhibition model (Figs 2D and E). With

the selected diffusion length of $l_D = 70$ µm, we observed some similarities between the dynamics of

the model outcomes and the *in vitro* networks: During the initial relaxation time (100 MCS) the ECs in

the models reached their target size and shape, which mimics stages 2 and 3 of the 5 stages described

in [15]: cell spreading and elongation. Next, we looked at the formation of cell-cell connections as

ECs started excreting a chemoattractant. The increase in cell-cell connections is reflected by a

decrease in number of branches (Fig 2F). The transition towards stage 4, stabilization of the network,

was more gradual in the contact inhibition model than in the *in vitro* networks and in the cell

elongation model (Fig 2E). The number of lacunae decreased faster in the cell elongation model than

in the contact inhibition model (Fig 2E) and for the first 1680 MCS we observed fewer and larger

lacunae in the contact inhibition model than in the cell elongation model. The number of branches was

initially larger in the cell elongation model than in the contact inhibition model. In agreement with the

in vitro networks, the branch number decreased quasi-exponentially in the cell elongation model,

whereas in the contact inhibition model the number of branches decreased more linearly (Fig 2F).

Both chemotaxis-based models have longer branches than the *in vitro* networks (Fig 2G). This

difference in number of branches and their length could partially be explained by a higher density of

sprouts detected *in vitro* than in the models (Fig 2H). Sprouts are branches that are connected to a

single node and the Analyze Skeleton plugin splits an existing branch into two where a sprout appears,

resulting in two shorter branches.

In the mechanical model, the ECs are assumed to stiffen the ECM through the exertion of traction

forces on their environment and move upwards the stiffness gradient (Fig 2B) [27]. We selected a

Young's modulus of 12 kPa, with a Poisson's ratio of 0.45, based on the original work. For a full list

of the selected parameters, see S7 Table. For the selected parameters, cells formed a dense network of

branches with small lacunae. The composition of this network was similar to an early-stage EC

network. This suggests that in the present form of the mechanical model, the mechanical model

successfully reproduces the network formation phases (phases 1-3), but it did not describe the

remodelling phase well.

Another noticeable difference between the three cellular Potts models and the *in vitro* EC networks

was the composition and thickness of the branches (Figs 1B and 2B, S3 Fig). *In vitro* cells are able to

move on top of each other to form a semi-3D structure, consisting of thin branches and thick nodes.

This is reflected in a decrease in the cell-covered area over time (S2 Fig E). In the strictly 2D *in silico*

models, however, cells are unable to move on top of each other and the cell-covered area remains

constant (S2 Fig E). Therefore, the models form increasingly thick branches and larger nodes as the

cell density increases (S3 Fig).

Increasing cell density results in an initial increase in number of branches and lacunae *in vitro*

To investigate whether the models respond similarly to a change in the initial conditions as *in vitro*

networks, we next analysed network formation at different cell densities. Previous *in vitro* studies on

primary ECs observed an inability for the cells to form tubes under or over a threshold density, similar

to a percolation threshold [5,22]. Consistent with these previous results, we observed that HMEC-1

cells retained the ability to form cell-cell connections and short branches at 1000 ECs per well (80

cells/mm²) (Fig 3A). However, they were not able to form a network spanning the full well. At a cell

density of more than 2000 ECs per well (160 cells mm²), we observed the development of a fully

connected network spanning the entire well (Fig 3A). With an increase in cells density a higher

number of lacunae and branches formed within the first hours (Figs 3B and D). However, for over

4000 cells per well (320 cells mm²), the networks remodelled to the same number of lacunae and

branches and average lacuna and branch sizes (Figs 3B-D).

Fig 3. A comparison of the response of *in vitro* networks and model outcomes to higher EC seeding density. A) Segmented *in vitro* networks with an inner well cell count of 1000 (top) and 4000 (bottom). B-E) Quantitative analysis of (B, C) branches and (D, E) lacunae for different cell densities. Scale bar = 500 µm. Shaded area represents standard deviation of 5 or 6 wells from two separate experiments.

The cell elongation model resembles the *in vitro* networks in its response to changes in cell density

Whereas a higher cell density caused an initial increase in the number of branches per mm² for all three *in silico* models, this difference diminished as the network remodelled in the two chemotaxis-based models at a diffusion length of 70 µm (Figs 4B and E), similar to what we observed *in vitro* (Fig 3B). The mechanical model had a much higher number of branches per mm², and the difference between branches per mm² for different cell densities converged much slower than in the chemotaxis-based models and the *in vitro* time-lapses (Figs 3B and 4A-I). Out of the three models, the later dynamics of the cell elongation model most resembled the *in vitro* network remodelling, which appears to be independent to a change in cell density, even when the model is subjected to a range of diffusion lengths (Figs 4A-C and S4 Fig).

Fig 4. A comparison of the response of model outcomes to a higher EC density. Quantitative

analysis of the number of branches for different cell densities with different diffusion lengths (l_D) in

the (A-C) cell elongation model and (D-F) the contact inhibition model, and (G-I) different stiffness

sensitivity (β) in the mechanical model. ($D = 5.0 \cdot 10^{-13} \mathrm{m}^2 \mathrm{s}^{-1}; \epsilon = 1.02 \cdot 10^{-4} \mathrm{s}^{-1}; \alpha = 1 \cdot$

$10^{-3} \mathrm{s}^{-1}$). Shaded area represents the standard deviation of six simulations.

Discussion

In biomedical applications, like tissue engineering [10] or tumour vascular normalization [11],

angiogenesis is stimulated, redirected or reduced to properly vascularize the tissue of interest.

Computational models can help understand how different environmental factors and cellular cues

affect various aspects of EC migration and network formation, and help identify effective regulatory

targets. Many computational models of EC network formation, based on assumptions like cell-cell

attraction [28], cell shape effects [14,23], cell-ECM interaction [21,27], or a combination of the former

[38], have been successful at simulating the formation of EC networks that qualitatively resemble

those observed *in vitro*. To obtain more insight into the extent to which the mechanisms described by

these computational models are a sufficient explanation for EC network formation *in vitro* we selected

three previously published, cell-based models of 2D EC network formation and compared these with

time-lapse data of EC network formation *in vitro*: (1) the cell elongation model [14]; (2) the contact-

inhibited chemotaxis model [28]; and (3) the mechanical cell-cell communication model [27]. We

examined their dynamical behaviour using a range of quantifications and compared it to time-lapses

data of *in vitro* EC network formation.

To quantify dynamical features of both *in vitro* EC network formation and outcomes of cellular Potts

models of 2D EC network formation, we developed a custom image analysis pipeline (Figs 1B, 2A

and S1 Fig A). To compare the models to our *in vitro* time-lapses we selected parameters from the

original work [14,27,28] and from published sensitivity analyses [39], and fitted parameters to our

own *in vitro* observations (S2 Figs A-C). We used the outcome of our initial dynamical analysis of the

in vitro and simulated EC networks to find a parameter space in which the network features of the

chemotaxis-based model outcomes resembled those of the *in vitro* EC networks (Figs 2C and S2 Fig

D). For the mechanical cell-cell communication model we were not able to find such a parameter

regime. The mechanical model was very sensitive and often lost its ability to form networks after

small changes in the explored parameters. Its high computational cost also made it difficult to perform

large scale parameter sweeps.

We then asked how the *in vitro* model responded to systematic changes in the culture conditions, with

the aim of mimicking the same conditions *in silico*. Among the conditions tested (S5 Fig) only cell

seeding density affected network formation *in silico* and *in vitro* to a sufficient degree to allow for

assessment of the correctness of the models (Figs 3 and 4). Interestingly, both chemotaxis-based

models showed similar dynamic behaviour as the *in vitro* networks for the selected parameter regime

(Figs 3 and 4 and S4 Fig). They showed an initial cell-cell contact forming phase, followed by a

remodelling phase, similar to Parsa's stages 3 to 5 of *in vitro* network formation: "elongation and

formation of cell-cell contacts", "plexus stabilization", and "plexus reorganization". In the mechanical

model, however, the ECs quickly form connections, mimicking early *in vitro* network formation, but

there is little network remodelling in the model outcomes.

We found that, among the models tested, the dynamic behaviour of the lacunae in the cell elongation

model matched the dynamics of the *in vitro* networks most closely (Figs 2D and E). In the cell

elongation model [14] the ECs actively elongate before they migrate upwards a chemoattractant

gradient. Because of their shape, the elongated ECs can form long-range connections to other ECs and

can thus form branches and close lacunae more rapidly than rounded ECs (Figs 2E and F). Due to the

slower movement along the short axis for elongated cells, branches are not very mobile and merge

slowly as the network stabilizes [40]. This results in a gradual decrease in lacunae per mm², which

mimics the remodelling phase of *in vitro* network formation quite well (Fig 2E). In the contact

inhibition model [28] however, network remodelling proceeds more slowly than in the elongated EC

model (Figs 2D and E). Also, in the contact inhibition model, for a short diffusion length ($l_D = 20$

µm), the number of branches per mm² remains dependent on the cell density (Fig 4D), which is not

observed *in vitro* or in the cell elongation model. The cell elongation model also predicts that networks

can form at lower cell densities than the contact inhibition model (Figs 4 and S4 Fig). Thus, the cell

elongation seems more robust in reproducing the remodelling stage of *in vitro* EC network formation

than the contact inhibition model.

There were also clear discrepancies between the cell elongation model and *in vitro* network formation.

For example, Parsa *et al*. (2011) observed that *in vitro* ECs aggregate and spread first, before

elongating [15]. Thus, *in vitro*, the earliest stages of network formation seem not driven by elongation,

as assumed by the cell elongation model. Also, the cell elongation model was unable to explain the

observation that network formation depends on substrate stiffness [5,41,42], as well as substrate

density (S5 Fig F) [6] and substrate thickness [43]. Such observations may be better explained by

models based on mechanical cell-cell communication [21,27]. Rüdiger *et al*. (2020) argue that

mechanical forces are the main driver behind *in vitro* EC network formation [5]. In this study they

continuously perfused the ECs during *in vitro* network formation to wash away any possible

chemoattractant gradient, and network formation still occurs. To further argue for mechanical

regulation, they show that ECs compress the substrate up to a distance of 30 µm, with measurable

ECM displacement up to 100 µm. These distances fall within the chemical signalling range of what we

observed in the two chemotaxis models similar to the *in vitro* networks (Figs 2C and S2 Fig D). In the

mechanical model we observed strains up to 40 µm from the cell boundary in the investigated

parameter regime (S2 Fig F). However, this was not enough to form networks similar to those

observed *in vitro* (Fig 2). Finding parameter regimes for the mechanical model in which it

quantitatively reproduces the *in vitro* dynamics might make it possible to explore the effect of ECM

straining on EC network formation in more detail. To find new parameter regimes in the mechanical

model we could combine our presented image-analysis pipeline with topological data analysis [44]

and novel high-throughput parameter inference methods such as FitMultiCell [45], which combines

the iteratively changing of parameters, the running of simulations and the assessment of the model

outcomes in one tool, to streamline the model fitting process. Also, a number of improvements are

possible in the mechanical model, which so far focused on synthetic, fully elastic substrates such as

polyacrylamide gels, which cannot be deformed or remodelled. In cell cultures using natural ECM

materials such as Matrigel, the ECs tend to deform and remodel the substrate over time [5,46], altering

the substrate's material properties and its response to stress. The absence of ECM remodelling in the

mechanical model could explain why the EC networks in the mechanical model stabilizes relative

quickly, whereas *in vitro* and in the two chemotaxis-based models the EC networks continuously

evolve as the simulated chemoattractant gradient gradually stabilizes over time. The effect of matrix

remodelling, in the form of strain-stiffening, on EC network formation dynamics could be investigated

in vitro with synthetic gels with tuneable stiffness [47,48]. With these gels the travel distance of

mechanical perturbations on the system can be altered systematically, to measure its effect on EC

network formation [16]. With the use of synthetic gels, it is also possible to wash away any

chemoattractant from the system, without the possible interference of ECM retention, which might

have influenced the network formation in Rüdiger *et al.* (2020). *In silico* models have considered

matrix remodelling by using continuum equations of matrix advection due to pulling by the ECs

[21,49]. A similar approach was integrated in a recent hybrid CPM-PDE model of network formation

[38]. In these models, the number of junctions was reported to increase over time, in contrast to our *in*

vitro observations (Fig 1C-H) and those of others [15,50], where the network remodels into a less

dense network within 24 hours. In our ongoing work, we are integrating detailed models of the

structural remodelling of fibrous ECMs into models of angiogenesis [51,52], which in the future may

give more insight into the potential role of matrix remodelling.

In the two chemotaxis-based models we assumed that each MCS represents 30 seconds in real time

and each pixel is 2 μm. This was chosen such that the mean speed of the cells became 5 μm/h^{-1},

which, it was previously argued, complies with *in vivo* observations [14,53]. This allows us to

translate the physical processes like chemoattractant diffusion and decay rate to physical units

(m^2s^{-1}; s^{-1}). We fixed the diffusion coefficient at $D = 5 \cdot 10^{-13}$ m^2s^{-1}. The main driver behind

angiogenesis is generally assumed to be VEGF-A [54]. The diffusion coefficient of VEGF-A is in the

order of 10^{-11} m^2s^{-1} in collagen-I, 10^{-10} m^2s^{-1} in Matrigel and $10^{-9}m^2s^{-1}$ in water [55–57],

which is 100 to 10000-fold higher than what we implemented in the model. However, ECM and EC

membrane receptor retention of VEGF-A could affect VEGF-A diffusion, resulting in a smaller

effective diffusion coefficient and decay rate [7]. Köhn-Luque *et al*., (2013) show an aggregation of

fluorescently labelled VEGF in the substrate surrounding HUVECs on Matrigel within 5-10 minutes

after VEGF addition [7]. Additionally, it is yet unclear whether VEGF-A diffusion is the main driver

behind angiogenic network formation *in vitro*. Especially since Rüdiger *et al*. (2020) observed EC

network formation under constant perfusion to avoid the emergence of a chemical gradient [5]. With a

diffusion coefficient of $D = 5 \cdot 10^{-13}$ m^2s^{-1} we observe simulated networks most like *in vitro* at a

diffusion length of $l_D = 70$ μm (Fig 2C). With a diffusion coefficient of $D = 10^{-10}$ m^2s^{-1}, the decay

rate would need to be $\epsilon = 2 \cdot 10^{-2}s^{-1}$ for a diffusion length of $l_D = 70$ μm, since $l_D = \sqrt{\frac{D}{\epsilon}}$, which is

much faster than the 90 minute half-life of VEGF-A measured *in vitro* [58]. *In vivo*, VEGF-A has been

shown to have a half-life of 33.7 minutes [59].

There are some steps in this image analysis pipeline that require optimization. For example, we

implemented a watershed step in the pipeline to correct for breaks in EC branches where thin

stretched-out ECs are misinterpreted as background during segmentation. This watershed step can

result in an overrepresentation of connections between branches, but overall, we observe an accurate

and consistent result (S1 Fig B). To circumvent the need for segmentation corrections and other

suboptimal parameter settings in the segmentation a neural network could be trained to perform the

segmentation steps necessary for the network quantification [60]. This network can be trained to

recognize thin stretches of ECs where conventional image analysis methods, which rely mostly on

contrast, might mistake them for background. However, the training of a neural network requires a lot

of correctly annotated training data and undertrained networks will give suboptimal results. More

interestingly, a neural network could be trained to recognize network features that are more difficult to

extract from segmented images automatically, like cell division events. Another interesting addition to

the image analysis pipeline would be the inclusion of topological data analysis (TDA) [44]. The time-

lapses of the segmented branches and lacunae could be regarded as multidimensional graphs suitable

for further analysis using TDA. TDA could also be expanded to work on 3D time-lapses to analyse 3D

in vitro EC sprouting assays [61,62], 3D vessels-on-a-chip sprouting assays [63] or even *in vivo*

systems, like the developing zebrafish embryo [64]. 3D *in vitro* systems have the benefit of both

having a high degree of control over the experimental environment, as allowing for 3D vascular

growth and EC migration. ECs *in vivo* migrate both in 2D over the basal membrane, as in 3D during

sprouting into the ECM [65]. 2D models of EC network formation can be regarded as a projection of a

3D system and adapted to study different migration types, but recent improvements in computational

power have allowed the increased usage of more complex 3D models of angiogenesis [66,67]. 3D

models, both *in vitro*, *in vivo* and *in silico* allow for a greater understanding of the complex

interactions ECs have with the fibrous ECM during sprouting [68].

Conventional tube formation assays to assess the sprouting ability of ECs are usually only examined

for a single characteristic (*e.g.* the number of branches) at a single time point [33]. However, we

observed that network features converged over time (Figs 3B-E). This indicates that an endpoint

observation might not suffice to examine the influence of a compound or knockout on collected cell

behaviour before a stable network has been formed. We also observed variation between different

samples of the same experimental conditions. This can be explained by variations in fitness of

different passage numbers of ECs [13], handling of the ECs prior to the experiment, and small

fluctuations in cell seeding density. Cell density naturally affects the distance between cells right after

seeding, and can therefore affect the ability of cells to sense their neighbours at the start of the

experiment [5,13,15]. Cells also influence the experimental environment by digesting nutrients and

excreting signalling molecules [69], but also by actively remodelling the ECM [5,70]. Therefore, it is

important that the exact number of seeded ECs is also considered when comparing samples within 6

hours after seeding or at low cell densities.

In this work we have shown the importance of dynamic analyses of EC network formation by showing

how EC network features converge over time. We hope to have paved the way for a more systematic

comparison of computational models to biological experiments, and we have indicated areas of

improvement for existing cellular Potts models of 2D EC network formation. In the future we plan to

expand our analysis of EC networks through more sophisticated data-analysis techniques such as

topological data analysis [44] and Sobol' analysis [32] for falsifying and refining our understanding of

EC network formation using *in vitro* and *in silico* modelling in hopes that such techniques can be

generalized to the dynamics of other models of biological morphogenesis.

Methods

Cell culture

Immortalised human microvascular endothelial cells (HMEC-1) were maintained in T75 flasks

(Sarstedt Inc) in MCDB-131 medium (Gibco) supplemented with 10 % fetal calf serum (FCS), 10

ng/mL epidermal growth factor (EGF) (Sigma-Aldrich), 1 µg/mL hydrocortisone (HC) (Thermo

Scientific Chemicals), 1 % GlutaMAX (Gibco) and 1 % penicillin/ streptomycin (Gibco). Cells were

cultured in a humidified incubator at 37 °C and 5 % CO2 and split twice per week. Passages used were

between 9 and 17.

Tube formation assay

Cells at 70-80% confluency were plated on growth factor reduced (GFR) Matrigel (Corning) on
Angiogenesis μ-slides (Ibidi) at a density of 10,000 cells per well unless stated otherwise. For
fluorescent plasma membrane labelling cells were incubated with PKH67, PKH26 or CellVue Claret
(Sigma-Aldrich) in Diluent C as described in [15].

Imaging

All images were captured at a magnification of 10X using a Nikon Ti inverted phase contrast
microscope equipped with a DS-Ri2 camera or a Zeiss Axio Observer inverted phase contrast
microscope equipped with an AxioCam 705 mono camera with a motorised stage for 24 hours with
15-minute intervals. Fluorescent images were acquired using the Zeiss Colibri multicolour LED light
source. Full well images were created by stitching multiple fields together with the NIS-elements and
ZEN Blue software.

Network characterization

Time-lapse images of the full wells were processed and quantified using a custom Fiji/ImageJ [71]
pipeline (S1 Fig A). For the segmentation of cells from background the following steps were
performed for each slice in the time-lapse images: (1) High-frequency features were amplified by first
convoluting the phase-contrast image with a Gaussian filter with sigma value of 10 µm, then
subtracting the resulting low-frequency image from the original image and finally adding the resulting
high-frequency image back to the original image. (2) Hereafter, the local variance in the sharpened
images was computed with a radius of 8.75 µm. The local variances were convoluted with a Gaussian
filter with a sigma value of 3.5 µm. (3) From the resulting image a threshold was computed with
automatic Huang thresholding to create a segmented mask of the network. (4) The mask was then
corrected for fused lacunae by producing a distance map of the lacunae larger than 2000 µm² and
watershed segmenting the distance map with a threshold set to 60. Hereafter the smaller lacunae were
added back to the watershed-segmented network for the final mask. The watershed step was omitted
for the first hour of the time-lapse, since no lacunae have formed yet. The branches and nodes within

the binary masks were analysed with the Skeletonize 2D/3D and Analyze Skeleton (2D/3D) [37]

plugins applied on a ten times scaled down image to avoid a very noisy skeleton. Lacunae between

4000 µm² and 11.3 mm² with a circularity between 0.2 and 1.0 were analysed with the Analyze

Particles plugin. The resulting output includes the number, size, position and shape of the lacunae, and

the number, length, position and connectivity of the branches. All results were exported to CSV files

for further analysis in RStudio [72]. This analysis pipeline was automated to process large quantities

of time-lapse images sequentially without user input to maximize reproducibility and minimize

personal bias in the data analysis. For the validation of the pipeline, we had six independent

participants manually encircle three lacunae and compared the average lacuna area of their selections

to automatically segmented images. The automatic segmentation underestimated the lacunae as

selected by the participants by 4.2 ± 11.7 % of the average area (n = 15). The difference between two

different participants encircling the same lacunae was 4.5 ± 2 % of average area (n = 3) (S1 Fig B).

Computational model

The three computational models of EC network formation [14,27,28] are all based on hybrid cellular

Potts models (CPMs). Hybrid CPMs couple a dynamical description of cell behaviour based on the

CPM with a detailed model of the cellular micro-environment, specifically the concentration of

extracellular signalling molecules or strains and stresses in the ECM. The cellular Potts model is a

cell-based model that focuses on modelling cell-cell interactions and predicting collective cell

behaviour [73]. Cells are represented as collections of lattice sites $\vec{x}$ with an identical index σ_x on a

grid, where each index labels a single cell. Each lattice site represents an area of 2 x 2 µm² in the

chemotaxis models and 5 x 5 µm² in the mechanical model. To mimic the circular *in vitro* dishes, we

labelled all lattice sites outside of a circle with a diameter of 3800 µm sites as boundary sites $\sigma_x = -1$

in the cell elongation and contact inhibition models and as a fixed node in the mechanical model.

Apart from the shapes of the simulation domains, models were used as described previously

[14,27,28]. Briefly, during a Monte Carlo step (MCS), a time-step in the simulation representing 30

seconds, cells change their shape and move through the lattice by attempting to copy its index to a

neighbouring lattice site. There are N copy attempts per MCS, with N the number of lattice sites. If the

copy attempt results in a lower effective energy of the system, the copy attempt is accepted. With an

effective energy H,

$H = \sum_{\vec{x},\vec{x}'} J_{\sigma_{\vec{x}},\sigma_{\vec{x}'}} \left(1 - \delta_{\sigma_{\vec{x}},\sigma_{\vec{x}'}}\right) + \lambda_A \sum_{\sigma}(a_\sigma - A_\sigma)^2 + \lambda_L \sum_{\sigma}(l_\sigma - L_\sigma)^2,$

where $J_{\sigma_{\vec{x}},\sigma_{\vec{x}'}}$ represents the bond energy between a lattice site $\vec{x}$ and its eight second-order neighbours

$\vec{x}'$, and λ_A and λ_L represent energy penalties for deviation from a designated target area (A_σ) or target

length (L_σ), which we set at 50 and 5.0 respectively. If the copy attempt does not result in a lower

effective energy, the copy is accepted with Boltzmann probability:

$P(\Delta H) = e^{-\Delta H/T}$

where T is the "cellular temperature" of the system, a measure of cell motility, which we set at 50, and

ΔH the change in the effective energy function.

In the cell elongation [14] and contact inhibition [28] models cells secrete a chemoattractant c, which

diffuses and degrades in the ECM,

$\frac{\partial c(\vec{x},t)}{\partial t} = \alpha(1 - \delta_{\sigma_{\vec{x}},0}) - \varepsilon\delta_{\sigma_{\vec{x}},0}c(\vec{x},t) + D\nabla^2 c(\vec{x},t),$

where $\delta_{\sigma_{\vec{x}},0} = 1$ outside the cells, α is the secretion rate and ε the degradation rate of the

chemoattractant and D the diffusion constant of the chemoattractant. An additional term is added to the

energy function, which promotes copies upwards the chemoattractant gradient:

$\Delta H_{chemotaxis} = -\mu(c(\vec{x}') - c(\vec{x})),$

Where μ is the strength of the chemotactic response, which we set at 1000. For details see [14,28].

In the mechanical cell-cell communication model [27], cells were able to deform the ECM through

cell-shape dependent traction forces. The deformations of the ECM are calculated using the finite

element method. An additional term is added to the energy function, such that cells respond to the

ECM stiffness:

$\Delta H_{durotaxis} = -g(\vec{x}, \vec{x}')\lambda_{durotaxis}\left(h\big(E(\epsilon_1)\big)(\vec{v}_1 \cdot \vec{v}_m)^2 + h\big(E(\epsilon_2)\big)(\vec{v}_2 \cdot \vec{v}_m)^2,$

where $g(\vec{x}, \vec{x}') = 1$ for cell extensions and $g(\vec{x}, \vec{x}') = -1$ for cell retractions, $\lambda_{\mathrm{durotaxis}}$ is a parameter

which we set at 24, $h(E)$ mimics the influence of the ECM stiffness E as a function of ϵ_1 and ϵ_2, the

principal strains, on the tension in the cell-ECM connection, $\vec{v}_m$ gives the copy direction, and v_1 and

v_2 are the strain orientations.

The preference for higher stiffness E is implemented as a sigmoid function:

$$h(E) = \frac{\alpha}{1 + \exp\left(-\beta(E - E_{\mathrm{tr}})\right)},$$

where α sets the strength of the durotactic response, E_{tr} determines the stiffness where half this

strength is reached and β represents the stiffness sensitivity which determines the steepness of the

curve. For details see [27].

Statistical analysis

564   P-values between groups were calculated using an ANOVA test followed by a student t-test. Dataset

565   significance was defined as $p \leq 0.05$ (*); $p < 0.01$ (**); $p < 0.001$ (***); $p > 0.05$ (ns).

Acknowledgments

This publication is part of the 'Mathematics-based strategies for repairing tumour blood vessels'

project with project number 865.17.004 of the Vici research program which is financed by the Dutch

Research Council (https://www.nwo.nl/). The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript. This work was performed using the

compute resources from the Academic Leiden Interdisciplinary Cluster Environment (ALICE)

provided by Leiden University.

References

1.   Risau W. Mechanisms of angiogenesis. Nature. 1997;386: 671–674. doi:10.1038/386671a0

2.   De Palma M, Biziato D, Petrova TV. Microenvironmental regulation of tumour angiogenesis.
Nat Rev Cancer. 2017;17: 457–474. doi:10.1038/nrc.2017.51

3.   Jiang X, Wang J, Deng X, Xiong F, Zhang S, Gong Z, et al. The role of microenvironment in
tumor angiogenesis. J Exp Clin Cancer Res. 2020;39: 204. doi:10.1186/s13046-020-01709-5

4.   Potente M, Gerhardt H, Carmeliet P. Basic and Therapeutic Aspects of Angiogenesis. Cell.
2011;146: 873–887. doi:10.1016/j.cell.2011.08.039

5.   Rüdiger D, Kick K, Goychuk A, Vollmar AM, Frey E, Zahler S. Cell-Based Strain Remodeling
of a Nonfibrous Matrix as an Organizing Principle for Vasculogenesis. Cell Rep. 2020;32:
108015. doi:10.1016/j.celrep.2020.108015

6.   Reinhart-King CA, Dembo M, Hammer DA. Endothelial Cell Traction Forces on RGD-
Derivatized Polyacrylamide Substrata. Langmuir. 2003;19: 1573–1579. doi:10.1021/la026142j

7.   Köhn-Luque A, Back W de, Yamaguchi Y, Yoshimura K, Herrero MA, Miura T. Dynamics of
VEGF matrix-retention in vascular network patterning. Phys Biol. 2013;10: 066007.
doi:10.1088/1478-3975/10/6/066007

8.   Hellström M, Phng L-K, Hofmann JJ, Wallgard E, Coultas L, Lindblom P, et al. Dll4 signalling
through Notch1 regulates formation of tip cells during angiogenesis. Nature. 2007;445: 776–780.
doi:10.1038/nature05571

9.   Lampugnani MG, Zanetti A, Corada M, Takahashi T, Balconi G, Breviario F, et al. Contact
inhibition of VEGF-induced proliferation requires vascular endothelial cadherin, β-catenin, and
the phosphatase DEP-1/CD148. J Cell Biol. 2003;161: 793–804. doi:10.1083/jcb.200209019

10. Mastrullo V, Cathery W, Velliou E, Madeddu P, Campagnolo P. Angiogenesis in Tissue
Engineering: As Nature Intended? Front Bioeng Biotechnol. 2020;8: 188.
doi:10.3389/fbioe.2020.00188

11. Jain RK. Antiangiogenesis strategies revisited: from starving tumors to alleviating hypoxia.
Cancer Cell. 2014;26: 605–622. doi:10.1016/j.ccell.2014.10.006

12. Nowak-Sliwinska P, Alitalo K, Allen E, Anisimov A, Aplin AC, Auerbach R, et al. Consensus
guidelines for the use and interpretation of angiogenesis assays. Angiogenesis. 2018;21: 425–
532. doi:10.1007/s10456-018-9613-x

13. DeCicco-Skinner KL, Henry GH, Cataisson C, Tabib T, Gwilliam JC, Watson NJ, et al.
Endothelial Cell Tube Formation Assay for the In Vitro Study of Angiogenesis. J Vis Exp JoVE.
2014; 51312. doi:10.3791/51312

14. Merks RMH, Brodsky SV, Goligorksy MS, Newman SA, Glazier JA. Cell elongation is key to
in silico replication of in vitro vasculogenesis and subsequent remodeling. Dev Biol. 2006;289:
44–54. doi:10.1016/j.ydbio.2005.10.003

15. Parsa H, Upadhyay R, Sia SK. Uncovering the behaviors of individual cells within a
multicellular microvascular community. Proc Natl Acad Sci U S A. 2011;108: 5133–5138.
doi:10.1073/pnas.1007508108

16. Califano JP, Reinhart-King CA. Substrate Stiffness and Cell Area Predict Cellular Traction
Stresses in Single Cells and Cells in Contact. Cell Mol Bioeng. 2010;3: 68–75.
doi:10.1007/s12195-010-0102-6

17. Sharpe J. Computer modeling in developmental biology: growing today, essential tomorrow.
Development. 2017;144: 4214–4225. doi:10.1242/dev.151274

18. Zhang Y, Wang H, Oliveira RHM, Zhao C, Popel AS. Systems biology of angiogenesis
signaling: Computational models and omics. WIREs Mech Dis. 2022;14: e1550.
doi:10.1002/wsbm.1550

19.   Subramanian A, Zakeri P, Mousa M, Alnaqbi H, Alshamsi FY, Bettoni L, et al. Angiogenesis
621   goes computational – The future way forward to discover new angiogenic targets? Comput
622   Struct Biotechnol J. 2022;20: 5235–5255. doi:10.1016/j.csbj.2022.09.019

623   20.   Miura T. Theoretical Models of Vascular Pattern Formation. Fukuoka Igaku Zasshi Hukuoka
624   Acta Medica. 2016;107: 161–168.

625   21.   Manoussaki D, Lubkin SR, Vernon RB, Murray JD. A mechanical model for the formation of
626   vascular networks in vitro. Acta Biotheor. 1996;44: 271–282. doi:10.1007/BF00046533

627   22.   Serini G, Ambrosi D, Giraudo E, Gamba A, Preziosi L, Bussolino F. Modeling the early stages
628   of vascular network assembly. EMBO J. 2003;22: 1771–1779. doi:10.1093/emboj/cdg176

629   23.   Palachanis D, Szabó A, Merks RMH. Particle-based simulation of ellipse-shaped particle
630   aggregation as a model for vascular network formation. Comput Part Mech. 2015;2: 371–379.
631   doi:10.1007/s40571-015-0064-5

632   24.   Metzcar J, Wang Y, Heiland R, Macklin P. A Review of Cell-Based Computational Modeling in
633   Cancer Biology. JCO Clin Cancer Inform. 2019;3: 1–13. doi:10.1200/CCI.18.00069

634   25.   Merks RMH, Glazier JA. A cell-centered approach to developmental biology. Phys Stat Mech Its
635   Appl. 2005;352: 113–130. doi:10.1016/j.physa.2004.12.028

636   26.   Vega R, Carretero M, Travasso RDM, Bonilla LL. Notch signaling and taxis mechanisms
637   regulate early stage angiogenesis: A mathematical and computational model. PLOS Comput
638   Biol. 2020;16: e1006919. doi:10.1371/journal.pcbi.1006919

639   27.   Oers RFM van, Rens EG, LaValley DJ, Reinhart-King CA, Merks RMH. Mechanical Cell-
640   Matrix Feedback Explains Pairwise and Collective Endothelial Cell Behavior In Vitro. PLOS
641   Comput Biol. 2014;10: e1003774. doi:10.1371/journal.pcbi.1003774

642   28.   Merks RMH, Perryn ED, Shirinifard A, Glazier JA. Contact-Inhibited Chemotaxis in De Novo
643   and Sprouting Blood-Vessel Growth. PLOS Comput Biol. 2008;4: e1000163.
644   doi:10.1371/journal.pcbi.1000163

645   29.   Dejana E. Endothelial cell–cell junctions: happy together. Nat Rev Mol Cell Biol. 2004;5: 261–
646   270. doi:10.1038/nrm1357

647   30.   Gory-Fauré S, Prandini M-H, Pointu H, Roullot V, Pignot-Paintrand I, Vernet M, et al. Role of
648   vascular endothelial-cadherin in vascular morphogenesis. Development. 1999;126: 2093–2102.
649   doi:10.1242/dev.126.10.2093

650   31.   Stéphanou A, Meskaoui G, Vailhé B, Tracqui P. The rigidity in fibrin gels as a contributing
651   factor to the dynamics of *in vitro* vascular cord formation. Microvasc Res. 2007;73: 182–190.
652   doi:10.1016/j.mvr.2006.12.002

653   32.   Boas SEM, Navarro Jimenez MI, Merks RMH, Blom JG. A global sensitivity analysis approach
654   for morphogenesis models. BMC Syst Biol. 2015;9: 85. doi:10.1186/s12918-015-0222-7

655   33.   Carpentier G, Berndt S, Ferratge S, Rasband W, Cuendet M, Uzan G, et al. Angiogenesis
656   Analyzer for ImageJ — A comparative morphometric analysis of "Endothelial Tube Formation
657   Assay" and "Fibrin Bead Assay." Sci Rep. 2020;10: 11568. doi:10.1038/s41598-020-67289-8

658   34.   Live Cell Analysis Instruments. In: Sartorius [Internet]. [cited 6 May 2023]. Available:
659   https://www.sartorius.com/en/products/live-cell-imaging-analysis/live-cell-analysis-instruments

35.   Tube Formation Image Analysis | WimTube | Wimasis. [cited 6 May 2023]. Available:
https://www.wimasis.com/en/WimTube

36.   Tube Formation FastTrack AI Image Analysis Software. In: ibidi [Internet]. [cited 6 May 2023].
Available: https://ibidi.com/software-and-image-analysis/189-tube-formation-fasttrack-ai-image-
analysis.html

37.   Arganda-Carreras I, Fernández-González R, Muñoz-Barrutia A, Ortiz-De-Solorzano C. 3D
reconstruction of histological sections: Application to mammary gland tissue. Microsc Res Tech.
2010;73: 1019–1029. doi:10.1002/jemt.20829

38.   Chiang H, Chung C-A. Simulation of Soluble and Bound VEGF-stimulated in vitro Capillary-
like Network Formation on Deformed Substrate. PLOS Comput Biol. 2024;20: e1012281.
doi:10.1371/journal.pcbi.1012281

39.   Boas SEM, Merks RMH. Tip cell overtaking occurs as a side effect of sprouting in
computational models of angiogenesis. BMC Syst Biol. 2015;9: 86. doi:10.1186/s12918-015-
0230-7

40.   Palm MM, Merks RMH. Vascular networks due to dynamically arrested crystalline ordering of
elongated cells. Phys Rev E. 2013;87: 012725. doi:10.1103/PhysRevE.87.012725

41.   Califano JP, Reinhart-King CA. The effects of substrate elasticity on endothelial cell network
formation and traction force generation. 2009 Annual International Conference of the IEEE
Engineering in Medicine and Biology Society. 2009. pp. 3343–3345.
doi:10.1109/IEMBS.2009.5333194

42.   Noerr PS, Zamora Alvarado JE, Golnaraghi F, McCloskey KE, Gopinathan A, Dasbiswas K.
Optimal mechanical interactions direct multicellular network formation on elastic substrates.
Proc Natl Acad Sci. 2023;120: e2301555120. doi:10.1073/pnas.2301555120

43.   Vernon RB, Angello JC, Iruela-Arispe L, Lane TF, Sage EH. Reorganization of Basement
Membrane Matrices by Cellular Traction Promotes the Formation of Cellular Networks in vitro.
Lab Invest. 1992;66: 536–547.

44.   Nardini JT, Stolz BJ, Flores KB, Harrington HA, Byrne HM. Topological data analysis
distinguishes parameter regimes in the Anderson-Chaplain model of angiogenesis. PLOS
Comput Biol. 2021;17: e1009094. doi:10.1371/journal.pcbi.1009094

45.   Alamoudi E, Schälte Y, Müller R, Starruß J, Bundgaard N, Graw F, et al. FitMultiCell:
simulating and parameterizing computational models of multi-scale and multi-cellular processes.
Bioinformatics. 2023;39: btad674. doi:10.1093/bioinformatics/btad674

46.   Davidson CD, Wang WY, Zaimi I, Jayco DKP, Baker BM. Cell force-mediated matrix
reorganization underlies multicellular network assembly. Sci Rep. 2019;9: 12.
doi:10.1038/s41598-018-37044-1

47.   Wang WY, Kent RN, Huang SA, Jarman EH, Shikanov EH, Davidson CD, et al. Direct
comparison of angiogenesis in natural and synthetic biomaterials reveals that matrix porosity
regulates endothelial cell invasion speed and sprout diameter. Acta Biomater. 2021;135: 260–
273. doi:10.1016/j.actbio.2021.08.038

48.   Dietrich M, Roy HL, Brückner DB, Engelke H, Zantl R, Rädler JO, et al. Guiding 3D cell
migration in deformed synthetic hydrogel microstructures. Soft Matter. 2018;14: 2816–2826.
doi:10.1039/C8SM00018B

49. Tranqui L, Tracqui P. Mechanical signalling and angiogenesis. The integration of cell–extracellular matrix couplings. Comptes Rendus Académie Sci - Ser III - Sci Vie. 2000;323: 31–47. doi:10.1016/S0764-4469(00)00110-4

50. Huuskes BM, DeBuque RJ, Kerr PG, Samuel CS, Ricardo SD. The Use of Live Cell Imaging and Automated Image Analysis to Assist With Determining Optimal Parameters for Angiogenic Assay in vitro. Front Cell Dev Biol. 2019;7. doi:10.3389/fcell.2019.00045

51. Tsingos E, Bakker BH, Keijzer KAE, Hupkes HJ, Merks RMH. Hybrid cellular Potts and bead-spring modeling of cells in fibrous extracellular matrix. Biophys J. 2023;122: 2609–2622. doi:10.1016/j.bpj.2023.05.013

52. Keijzer KAE, Tsingos E, Merks RMH. How cells align to structured collagen fibrils: a hybrid cellular Potts and molecular dynamics model with dynamic mechanosensitive focal adhesions. Front Cell Dev Biol. 2025;12. doi:10.3389/fcell.2024.1462277

53. Rupp PA, Czirók A, Little CD. αvβ3 integrin-dependent endothelial cell dynamics in vivo. Development. 2004;131: 2887–2897. doi:10.1242/dev.01160

54. Carmeliet P. VEGF as a Key Mediator of Angiogenesis in Cancer. Oncology. 2005;69: 4–10. doi:10.1159/000088478

55. Jeong GS, Han S, Shin Y, Kwon GH, Kamm RD, Lee S-H, et al. Sprouting Angiogenesis under a Chemical Gradient Regulated by Interactions with an Endothelial Monolayer in a Microfluidic Platform. Anal Chem. 2011;83: 8454–8459. doi:10.1021/ac202170e

56. Miura T, Tanaka R. *In vitro* Vasculogenesis Models Revisited - Measurement of VEGF Diffusion in Matrigel. Math Model Nat Phenom. 2009;4: 118–130. doi:10.1051/mmnp/20094404

57. Shin Y, Han S, Jeon JS, Yamamoto K, Zervantonakis IK, Sudo R, et al. Microfluidic assay for simultaneous culture of multiple cell types on surfaces or within hydrogels. Nat Protoc. 2012;7: 1247–1259. doi:10.1038/nprot.2012.051

58. Kleinheinz J, Jung S, Wermker K, Fischer C, Joos U. Release kinetics of VEGF165 from a collagen matrix and structural matrix changes in a circulation model. Head Face Med. 2010;6: 17. doi:10.1186/1746-160X-6-17

59. Eppler SM, Combs DL, Henry TD, Lopez JJ, Ellis SG, Yi J-H, et al. A target-mediated model to describe the pharmacokinetics and hemodynamic effects of recombinant human vascular endothelial growth factor in humans. Clin Pharmacol Ther. 2002;72: 20–32. doi:10.1067/mcp.2002.126179

60. Maška M, Ulman V, Delgado-Rodriguez P, Gómez-de-Mariscal E, Nečasová T, Guerrero Peña FA, et al. The Cell Tracking Challenge: 10 years of objective benchmarking. Nat Methods. 2023;20: 1010–1020. doi:10.1038/s41592-023-01879-y

61. Beter M, Abdollahzadeh A, Pulkkinen HH, Huang H, Orsenigo F, Magnusson PU, et al. SproutAngio: an open-source bioimage informatics tool for quantitative analysis of sprouting angiogenesis and lumen space. Sci Rep. 2023;13: 7279. doi:10.1038/s41598-023-33090-6

62. Kempers L, van der Bijl I, van Stalborch A-MD, Ponsioen B, Margadant C. Fast *in vitro* protocol for the visualization and quantitative high-throughput analysis of sprouting angiogenesis by confocal microscopy. STAR Protoc. 2021;2: 100690. doi:10.1016/j.xpro.2021.100690

744   63.   van Duinen V, Zhu D, Ramakers C, van Zonneveld AJ, Vulto P, Hankemeier T. Perfused 3D
745   angiogenic sprouting in a high-throughput in vitro platform. Angiogenesis. 2019;22: 157–165.
746   doi:10.1007/s10456-018-9647-0

747   64.   Schuermann A, Helker CSM, Herzog W. Angiogenesis in zebrafish. Semin Cell Dev Biol.
748   2014;31: 106–114. doi:10.1016/j.semcdb.2014.04.037

749   65.   Michaelis UR. Mechanisms of endothelial cell migration. Cell Mol Life Sci CMLS. 2014;71:
750   4131–4148. doi:10.1007/s00018-014-1678-0

751   66.   Swat MH, Thomas GL, Belmonte JM, Shirinifard A, Hmeljak D, Glazier JA. Multi-Scale
752   Modeling of Tissues Using CompuCell3D. Methods Cell Biol. 2012;110: 325–366.
753   doi:10.1016/B978-0-12-388403-9.00013-8

754   67.   Jafari Nivlouei S, Soltani M, Shirani E, Salimpour MR, Travasso R, Carvalho J. A multiscale
755   cell-based model of tumor growth for chemotherapy assessment and tumor-targeted therapy
756   through a 3D computational approach. Cell Prolif. 2022;55: e13187. doi:10.1111/cpr.13187

757   68.   Rauff A, LaBelle SA, Strobel HA, Hoying JB, Weiss JA. Imaging the Dynamic Interaction
758   Between Sprouting Microvessels and the Extracellular Matrix. Front Physiol. 2019;10: 1011.
759   doi:10.3389/fphys.2019.01011

760   69.   Eelen G, de Zeeuw P, Treps L, Harjes U, Wong BW, Carmeliet P. Endothelial Cell Metabolism.
761   Physiol Rev. 2018;98: 3–58. doi:10.1152/physrev.00001.2017

762   70.   Wang X, Khalil RA. Matrix Metalloproteinases, Vascular Remodeling, and Vascular Disease.
763   Adv Pharmacol San Diego Calif. 2018;81: 241–330. doi:10.1016/bs.apha.2017.08.002

764   71.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
765   source platform for biological-image analysis. Nat Methods. 2012;9: 676–682.
766   doi:10.1038/nmeth.2019

767   72.   RStudio Team. RStudio: Integrated Development Environment for R. Boston, MA: RStudio,
768   PBC; 2022. Available: http://www.rstudio.com/

769   73.   Glazier JA, Graner F. Simulation of the differential adhesion driven rearrangement of biological
770   cells. Phys Rev E. 1993;47: 2128–2154. doi:10.1103/PhysRevE.47.2128

 # Supporting information

S1 Fig. The image analysis pipeline uses automatic segmentation to avoid bias in time-lapse

analysis A) A detailed scheme of the image analysis pipeline showing intermediate images. B) A

comparison between lacunae area measured by independent peers and automatically segmented

lacunae.

S2 Fig. **Additional images and graphs** A) ECs were labelled using membrane linker dyes and

segmented from the background. B) The average area of fluorescently labelled ECs over 24 hours. (n

> 600 per timestep) C) The average cell length of elongated cells over time (Area between 500 and

1500 μm²; roundness below 0.25) (n > 6 per timestep). D) A comparison of the number of branches

per mm² for chemoattractant diffusion lengths ranging from 10 μm to 100 μm for the cell elongation

model and the contact inhibition model after 2880 MCS. In magenta the *in vitro* number of branches

per mm² after 24 hours. E) Cell covered area was measured as a percentage of the total well area.

Shaded areas represent the standard deviation. F) ECs (white) in the mechanical model exert strain on

their environment indicated by blue green heatmap. G) Histogram shows the relative frequency of

lacuna areas in the computational models of endothelial cell network formation after 24 hours (2880

MCS) $(D = 5.0 \cdot 10^{-13} \mathrm{m}^2 \mathrm{s}^{-1}; \epsilon = 1.02 \cdot 10^{-4} \mathrm{s}^{-1}; \alpha = 1 \cdot 10^{-3} \mathrm{s}^{-1}; n = 8)$.

S3 Fig. Overview images of simulated networks for different cell densities after 2880 MCS. A)

Cell elongation model. B) Contact inhibition model. C) Mechanical model.

S4 Fig. Overview of dynamical analysis of simulated networks for different cell densities and diffusion length or stiffness sensitivity. A-D) Cell elongation model. E-H) Contact inhibition model. I-L) Mechanical model. Overview images are 4000 cells after 2880 MCS.

S5 Fig. Overview of branch length analysis of *in vitro* endothelial cell networks for different

experimental conditions. A) HMEC-1 cells were seeded at different densities. B) 10 mM l-lactate

was added to the MCDB-131 medium prior to the start of the tube formation assay. C) HMEC-1 cells

were seeded on either growth factor reduced Matrigel or regular Matrigel. HCl was added to the

medium prior to the start of the tube formation assay to reduce its pH to 6.0. D) Fetal calf serum was

added to the medium in different percentages. E) HMEC-1 cells were seeded on either growth factor

reduced Matrigel or regular Matrigel. VEGF-A was added to the medium. F) HMEC-1 cells were

seeded on either regular Matrigel diluted in DMEM high-glucose.

807 **S6 Table. Chemotaxis model parameters**

Parameter	Value	Unit
Pixel size	2e-6	m
Field diameter	1900	pixels
CPM temperature	50	-
Cell target area	250	pixels
Initial cell diameter	12	pixels
Strength of area constraint	50	-
Strength of length constraint	5	-
Connectivity dissipation	5000	$N\,m^{-1}$
Cell-medium contact cost	20	-
Cell-cell contact cost	40	-
Chemotaxis strength	1000	$N\,m^{-1}$
Relaxation time	100	MCS
Border energy	100	$N\,m^{-1}$
Order of neighbourhood	2	-
Diffusion coefficient	5e-13	$m^2\,s^{-1}$
Secretion rate	1e-3	s^{-1}
PDE time increment	2	s
PDE iterations per MCS	15	-

809 **S7 Table. Mechanical model parameters**

Parameter	Value	Unit
Pixel size	0.000005	m
Field diameter	760	pixels
CPM temperature	1	-
Cell target area	40	pixels
Strength of area constraint	10000	-
Cell force	1	$N\,m^{-1}$
Cell-medium contact cost	500000	m^{-1}
Cell-cell contact cost	1000000	m^{-1}
Young's modulus	12000	Pa
Poisson's ratio	0.45	-
Accuracy of solver	0.00001	-
Strength of durotaxis	24	-
Threshold for stiffness preference	15000	Pa
Strain stiffening parameter	0.1	-