1 **Pleiotropy increases parallel selection signatures during adaptation from**

2 **standing genetic variation**

4 Wei-Yun Lai[1,2], Sheng-Kai Hsu[1,2], Andreas Futschik[3] and Christian Schlötterer[1*]

[1]Institut für Populationsgenetik, Vetmeduni Vienna, Vienna, Austria.

[2]Vienna Graduate School of Population Genetics, Vetmeduni Vienna, Vienna, Austria.

[3]Department of Applied Statistics, Johannes Kepler University Linz, Linz, Austria.

[*]Correspondence: christian.schloetterer@vetmeduni.ac.at; Tel.: +43-1-25077-4300.

Abstract

The phenomenon of parallel evolution, whereby similar genomic and phenotypic changes

occur across replicated pairs of population or species, is widely studied. Nevertheless, the

determining factors of parallel evolution remain poorly understood. Theoretical studies have

proposed that pleiotropy, the influence of a single gene on multiple traits, is an important

factor. In order to gain a deeper insight into the role of pleiotropy for parallel evolution from

standing genetic variation, we characterized the interplay between parallelism, polymorphism

and pleiotropy. The present study examined the parallel gene expression evolution in 10

replicated populations of *Drosophila simulans*, which were adapted from standing variation

to the same new temperature regime. The data demonstrate that parallel evolution of gene

expression from standing genetic variation is positively correlated with the strength of

pleiotropic effects. The ancestral variation in gene expression is, however, negatively

correlated with parallelism. Given that pleiotropy is also negatively correlated with gene

expression variation, we conducted a causal analysis to distinguish cause and correlation and

evaluate the role of pleiotropy. The causal analysis indicated that both direct (causative) and

indirect (correlational) effects of pleiotropy contribute to parallel evolution. The indirect

effect is mediated by historic selective constraint in response to pleiotropy. This results in

parallel selection responses due to the reduced standing variation of pleiotropic genes. The

direct effect of pleiotropy is likely to reflect a genetic correlation among adaptive traits,

which in turn gives rise to synergistic effects and higher parallelism.

Background

Distinguishing selection from stochastic changes continues to be a major challenge in evolutionary research. The evidence for selection mostly comes from parallel evolution, where similar genes, traits or functions are identified in replicate populations exposed to similar environments [1-4]. Given that non-parallel evolution does not preclude the possibility of selection [5, 6], there has been a sustained interest in elucidating the factors that influence the extent of genetic parallelism (i.e., how similar is the response of the same gene to selection in multiple populations from the same/similar ecological niche(s)) [7].

It has been proposed that the degree of pleiotropy, the number of traits affected by a single gene, may be a potential factor associated with parallelism [7-10]. The majority of hypotheses regarding the influence of pleiotropy on parallel selection responses are (implicitly) related to the ideas of Fisher [11], later modified by Orr [12], focusing on the cost of pleiotropy. This concept posits that any new mutation in a pleiotropic gene results in a shift in trait values away from their optimum across multiple traits. New mutations in highly pleiotropic genes are more likely to have negative fitness effects than mutations in less pleiotropic genes as more traits are displaced from their optima. This "cost of pleiotropy" model [12] is evidenced by the lack of sequence divergence observed in genes with a higher level of pleiotropy [13-16]. Consequently, it can be postulated that less pleiotropic genes are more likely to contribute to adaptation over long evolutionary timescales, given that they have a higher probability of accumulating beneficial genetic variants. This results in a more parallel selection response in different populations/species [17].

Fisher's model of pleiotropy is simplistic in its assumption of uncorrelated traits and uncorrelated selection. It has been proposed that the influence of pleiotropy on the selection response strongly depends on the genetic correlation of the affected traits and the underlying fitness function which connects phenotypic values with fitness [18-20]. If the fitness effects of the affected traits are positively correlated (synergistic pleiotropy), this may result in stronger net selection [21] and consequently in more parallel selection signatures [22]. This implies that the correlation of fitness effects determines whether pleiotropy increases or decreases parallel selection signatures.

The impact of pleiotropy on parallel evolution has typically been considered in the context of long-term adaptation via de novo mutations. However, many studies of parallel evolution

have focused on short-term adaptation [e. g. 23, 24-30], where standing genetic variation

plays a more significant role for selection responses than de novo mutations. Nevertheless, it

remains unclear whether a similar impact of pleiotropy on evolutionary parallelism is to be

expected for standing genetic variation.

Indeed, the empirical evidence regarding the role of pleiotropy for intra-specific adaptive

responses is somewhat inconclusive. A gene expression study in European grayling

populations adapted to different temperature regimes provided compelling evidence for the

constraining effect of pleiotropy [31], which supports the cost of complexity concept. A

comparison of gene expression changes in natural *Drosophila* populations along a

temperature cline and populations evolved in the laboratory to different temperature regimes

demonstrated the importance of correlated fitness effects [21]. A direct test of the influence

of pleiotropy on parallel evolution was conducted on 19 independent contrasts of stickleback

populations [9]. The authors discovered that pleiotropic genes were more likely to exhibit

parallel adaptation signatures [9], thereby supporting the importance of synergistic

pleiotropy.

Previous studies on pleiotropy and parallelism did not account for polymorphism (standing

genetic variation), which plays a central role for selection responses during short-term

adaptation. Although natural variation provides the basis for adaptive responses, it is not

well-documented how ancestral variation (polymorphism) determines the level of genetic

parallelism during short-term adaptation in replicate populations. Moreover, polymorphism

of a gene is also affected by its pleiotropic level while the expectations regarding the

influence of pleiotropy on the extent of polymorphism varies. The cost of complexity predicts

that pleiotropic genes with uncorrelated fitness effects will be subject to stronger purifying

selection, which will result in reduced polymorphism. In the case of multiple selected traits

with aligned fitness effect, synergistic pleiotropy may facilitate the occurrence of selective

sweeps and thereby reducing variation in pleiotropic genes. Alternatively, it may be possible

that once pleiotropic alleles have been established in a population, their multivariate fitness

effects lead to a balanced polymorphism, which in turn maintains higher levels of variation.

Empirical studies have observed a negative correlation between the level of pleiotropy and

sequence/expression polymorphism [32, 33]. This suggests that pleiotropy may preferentially

reduce variation, either by purifying selection, synergistic effects, or a combination of both.

In this study we employed gene expression analysis to investigate the interplay between
polymorphism and pleiotropy on the degree of parallelism of adaptive responses. We
assessed the selective response by measuring gene expression differences between
populations from different environments, a widely employed approach [e. g. 34, 35-40].
Specifically, we compared gene expression in the ancestral *D. simulans* population with that
of flies that had evolved for 100 generations in a novel hot temperature regime. To infer
parallel evolution, the gene expression change was measured in 10 replicate populations. By
focusing on the evolutionary response, we circumvent the challenge of connecting the
phenotype to an unknown fitness landscape. Furthermore, we take advantage of well-
established pleiotropy estimates [41-44] to evaluate the influence of pleiotropy on parallel
gene expression evolution. The natural variation in gene expression was determined by
examining the differences in expression levels among individual flies from the ancestral *D.*
simulans population.

We observed significant interplays between pleiotropy, ancestral variation, and parallel
selection responses among genes. The parallelism of gene expression evolution is positively
associated with pleiotropy but negatively associated with the ancestral variation in gene
expression. Given that pleiotropy was also negatively associated with ancestral variation, it is
crucial to differentiate between statistical causality and correlation. Causal analysis revealed
that pleiotropy affects parallel evolution both directly and indirectly via ancestral variation. It
is likely that the indirect effects operate via the reduction of ancestral variation in pleiotropic
genes. The direct effects of pleiotropy on parallelism are probably best explained by
synergistic pleiotropy which results in a stronger selection response.

Methods

<u>Estimating gene pleiotropy</u>

We approximated the pleiotropy of each gene with two alternative estimators, network

connectivity and tissue specificity (τ).

τ indicates how specific the expression of a gene is across different tissues. τ was estimated

for each gene using the adult male expression profiles on flyatlas2 [45] as:

$\tau = \frac{\sum_i [1 - \frac{gene\ expression_i}{gene\ expression_{max}}]}{N-1}$, where N is the number of tissues examined and i indicates each of

them [46]. If a gene is only expressed in one tissue, τ will equal to 1 while it equals 0 when a

gene is expressed at the same level across all tissues. The relationship between τ and

pleiotropy is based on the idea that the genes expressed in many tissues are more likely to

affect multiple traits than genes expressed in fewer tissues [41, 44, 47, 48]. Hence, we used 1-

τ to indicate the pleiotropic effect of a gene.

Network connectivity has also been used as proxy for pleiotropy, as more connected genes

are likely to be more pleiotropic [16, 42, 43, 49]. We used published information about the

transcriptional regulatory network of *Drosophila*, which was inferred from several sources,

including genome-wide chromatin immuno-precipitation, conserved transcription factor

binding motifs, gene expression profiles across different development stages, and chromatin

modification profiles among several cell types [50]. The connectivity for each gene was

estimated as the sum of adjacencies between the focal gene and other genes in the network.

The significant positive correlation (rho=0.54) between 1-τ and network connectivity

suggests that both estimates capture a similar, but not identical, signature of the pleiotropic

properties among the genes (Supplementary figure 1).

We note that approximating pleiotropic effects in *D. simulans* by using data-rich estimates

from the close relative *Drosophila melanogaster* could potentially compromise the

interpretability of our results. However, comparative analyses of tissue-specific gene

expression have shown an extremely high consistency across species in animals [51] and

plants [52, 53]. Hence, these data suggest that at least for tissue specificity, one of the

pleiotropy measures used, estimates from one species can be safely used for close relatives.

<u>Experimental evolution and common garden experiment</u>

The procedures of the evolution and common garden experiments were described in [5, 35,

54]. Briefly, 202 isofemale lines collected from Florida, USA were used to constitute 10

outbred *Drosophila simulans* populations, which have been exposed for more than 100

generations to a laboratory experiment at 28/18 °C with 12hr light/12hr dark photoperiod.

The census population size of each replicate population is 1000 to 1250 adult flies.

Each founder isofemale line was maintained at a small population size (typically less than 50

individuals) at 18 °C on standard laboratory food. Adaptation to the laboratory environment

with the residual heterogeneity or de novo mutation is unlikely due to the small effective

population size during the maintenance of each line [5]. This is supported by the observation

that 50 generations of maintenance as isofemale lines does not result in significant genetic

differentiation from the ancestral population [55]. Furthermore, mutations that occur in the

laboratory are mostly recessive [56], and the effects are likely to be masked because after two

generations of random mating during common garden maintenance, most individuals will be

heterozygous for isofemale line-specific variants.

The collection of samples for RNA-Seq was preceded by two generations of common garden

experiment (CGE). As described previously [5, 35, 54], at generation 103 of the evolution

experiment, five biological replicates of reconstituted ancestral populations were generated

by pooling five mated females from 184 founder isofemale lines [57] and reared together

with 10 independently evolved populations for two generations with controlled egg density

(400 eggs/bottle) at the same temperature regime as in the evolution experiment. Three

biological replicates were included for each independently evolved population (each

biological replicate was cultivated separately for the two generations of common garden).

From each biological replicate, a sample of 50 five-day-old adult males were collected

(Figure 1a).

<u>Quantifying the evolutionary parallelism in gene expression across ten replicate populations</u>

The raw RNA-Seq count table of five replicates of the reconstituted ancestral population and

10 independently evolved populations, each with three biological samples, was taken from

(Jaksic et al., 2020). The raw read counts of each gene were normalized with the TMM

method implemented in edgeR [58]. Only genes with at least 0.1 normalized counts per

million base (CPM) across all samples were considered for further analysis. Because we are

interested in the evolutionary response in each independently evolved population, we

contrasted the three biological samples from each evolved population to the five biological
samples from the ancestral population. The differential expression (DE) analysis was done
separately in each of the ten evolved populations. For DE analysis, we utilized negative
binomial generalized linear modeling implemented in edgeR to fit the expression to the
model $Y = E + \varepsilon$, in which Y stands for gene expression, E is the effect of evolution and ε is
the random error. Likelihood ratio tests were performed to test the effect of evolution. P-
values was adjusted using Benjamini-Hochberg false discovery rate (FDR) correction [59].
After we identified DE genes in each evolved population, we constructed a replicate-
frequency spectrum (RFS; [5]). We considered genes with significant changes in the same
direction in at least three evolved populations as putatively adaptive genes.

To quantify the degree of parallel gene expression evolution, we calculated the expression
changes in response to evolution (log$_2$FC; fold change in expression between ancestral and
evolved sample; see below) of each putatively adaptive gene (n=1,775) in each evolved
sample.

$$log_2FC = \log\left(\frac{y_{evo}}{\overline{y_{anc}}}\right),$$

210   (1)
211 where y_{evo} is the expression value (CPM) per gene per evolved sample, and $\overline{y_{anc}}$ is the
212 overall mean expression value (CPM) per gene across five ancestral samples.
213 We calculated the ratio of between-evolution population variation (MS$_{pop}$) and residual (MS$_e$)
214 among the expression changes (log$_2$FC), which is denoted as F.

$$F = \frac{MS_{pop}}{MS_e}$$

215   (2)

$$MS_{pop} = \frac{\sum_{i=1}^{10}(\bar{X}_i - \bar{X})^2}{10 - 1}$$

216   (3)

$$MS_e = \frac{\sum_{i=1}^{10}\sum_{j=1}^{3}(X_{ij} - \bar{X}_i)^2}{30 - 10}$$

217   (4)
Where X_{ij} represent the log$_2$FC value for the j[th] biological sample in the i[th] evolution
population. i=1, 2, …,10, j=1, 2, 3.
As F reflects the heterogeneity of the evolutionary response for a gene across ten
evolutionary populations, we used the reciprocal of F (i.e., 1/F) to quantify the degree of
parallelism in gene expression evolution.

<u>Estimating gene expression variation in an outbred ancestral population</u>

The variation in expression of each gene within an outbred ancestral population was

estimated from 20 adult males sampled from an outbred ancestral population [60]. We

distinguished biological variation and measurement error of each gene using the statistical

method implemented in edgeR [58], where the biological variance across individual

(biological coefficient of variation; BCV; defined in [58]) of each gene can be estimated (tag-

wised dispersion). BCV^2 was used to represent the expression variance relative to the mean

of each gene within the population. To illustrate that the gene expression variation is not

explained by random measurement error, we took the data from two ancestral population

replicates, which are considered to be genetically identical and calculated BCV^2 of expression

levels across individuals for each population replicates separately. We found that both mean

and variance in expression for each gene was similar in two ancestral population replicates

(mean: rho>0.99; variance: rho=0.8, p-value<2.2e-16), suggesting that sequencing noise does

not mask the biological variability.

<u>Causal analysis</u>

To infer the causal relationship between three factors: pleiotropy (Pl), ancestral variation (A)

and parallelism (Pa), we applied causal analysis. The causal analysis is built upon a

previously published statistical framework [61]. We consider five possible causal

relationships between pleiotropic effects, ancestral variation and parallelism (Figure 5). Our

goal is to determine which of the five models is best supported by the data. The selection is

based on Bayesian information criteria (BIC) which are calculated from maximum likelihood

of the respective models. The likelihood of each model is represented as:

$$\mathcal{L}(\text{Model I}) = p(Pa|A)p(A|Pl)p(Pl)$$

247 $$(5)$$

$$\mathcal{L}(\text{Model II}) = p(Pa|Pl)p(A|Pl)p(Pl)$$

248 $$(6)$$

$$\mathcal{L}(\text{Model III}) = p(Pa|A, Pl)p(A|Pl)p(Pl)$$

249 $$(7)$$

$$\mathcal{L}(\text{Model IV}) = p(Pa|A, Pl)p(A)p(Pl)$$

250 $$(8)$$

$$\mathcal{L}(\text{Model V}) = p(Pa)p(A)p(Pl)$$

251 $$(9)$$
252

p(Pl) is the probability distribution of the pleiotropic effect. To utilize normal probability

density function in the derivation of the likelihoods for each joint probability distributions,

we transformed the random variables Pa and A to be normally distributed (natural log

transformation on 1/F and BCV^2). The exact forms of these likelihoods are given in

supplementary information.

For each model, we maximized the likelihood and estimated the parameters using standard

maximum likelihood methods. Based on the maximum likelihood, we computed the BICs for

each model; the model with the smallest BIC is the one best support by the data.

BIC is calculated as:

$$\text{BIC} = -2\ln(\mathcal{L}_i) + k_i ln(n)$$

$$(8)$$

Where $\mathcal{L}_i$ is the likelihood for model I-V, k_i is the corresponding number of free parameters

and n is total number of putatively selected genes. The analysis was performed independently

on two measures of pleiotropy (tissue specificity and network connectivity).

Estimating the sizes of direct and indirect pleiotropic effects on parallelism

We used path analysis to estimate the effect sizes of direct and indirect pleiotropic effects on

parallelism [62]. We standardized all three transformed variables (Parallelism (Pa),

Pleiotropy (Pl), Ancestral variation (A); see supplementary information) by subtracting the

mean and dividing by the standard deviation for the following analysis.

First, we fit the regression model for all three variables across all putatively adaptive genes:

$$y_{Pa} = \beta_1 x_A + \beta_2 x_{Pl} + \varepsilon$$

276　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　(9)

277　where y_{Pa} stands for parallelism, x_A is ancestral variation, x_{Pl} is pleiotropy measure and ε is

278　random error. β_1 and β_2 are the regression coefficients corresponding to x_A and x_{Pl},

279　respectively.

281　Second, we fit another regression model for ancestral variation and pleiotropy measure across

282　all putatively adaptive genes:

$$y_A = \beta_3 x_{Pl} + \varepsilon$$

283　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　(10)

284　where y_A stands for ancestral variation, x_{Pl} is pleiotropy measure and ε is random error. β_3 is

285　the regression coefficient corresponding to x_{pl}.

287　The sizes of the direct effect of pleiotropy on parallelism will be β_2. The size of the indirect

288　effect of pleiotropy on parallelism via ancestral variation will then be the product of $\beta_1 \times \beta_3$.

290　The analysis was performed independently on two different measures of pleiotropy.

291
Computer simulations

In our interpretation of the indirect effect of pleiotropy on parallelism, we assumed that

ancestral variation in gene expression is negatively correlated with its parallel evolution.

Because this has not been previously demonstrated, we used computer simulations to

illustrate how the level of standing genetic variation impacts the parallelism of adaptive

responses after a shift in trait optimum. For simplicity, we considered four genes equally

contributing to a fitness-associated trait. The phenotype (expression level) of each gene is

controlled by a different number of genetic loci (5, 15, 30 and 50) with equal effect. The

underlying assumption is that these four genes exhibit different levels of pleiotropy and thus

experience varying strength of purifying selection, retaining different levels of genetic

variation. Using mimicrEE2 [63], we evolved 10 replicate populations (N=300) derived from

the same set of 189 natural *Drosophila* haplotypes [5] in same selection regime with a shift in

trait optimum of one standard deviation relative to the ancestral phenotypic distribution. After

100 generations, we determined the evolutionary parallelism of the four genes using the same

approach as for the empirical data. Each simulation was repeated 100 times and one out of

four genes was randomly picked from each simulation run to generate Figure 4.

Results

The parallelism of expression evolution differs among genes

We investigated the degree of parallelism in gene expression evolution, using an RNA-Seq dataset from an experimental evolution study in which 10 replicated populations with the same genetic variation were exposed to the same environment for more than 100 generations [29, 54, 60] (Figure 1a). While previous studies have focused on the shared parallel response across replicated populations [29, 54, 60], we re-analyzed the RNA-Seq data to quantify the degree of evolutionary parallelism across different genes. We identified 1,775 putatively selected genes with a significant change in gene expression in the same direction in response to the novel environment in at least three populations. This conditioning on multiple replicate populations with a significant change in the same direction reduces false positives due to drift [9, 64]. Relaxing the threshold of responding populations provides qualitatively similar results.

Barghi et al. (2019) introduced the replicate frequency spectrum to describe the parallel genomic selection response in replicate populations. The replicate frequency spectrum of DE genes shows considerable heterogeneity among populations, with many genes displaying significant gene expression changes only in a few populations (Figure 1b). The replicate frequency spectrum is strongly influenced by the significance threshold used and provides an incomplete quantitative measure of parallelism. To obtain a more quantitative measure of parallelism, we evaluated the between-population variation in the evolutionary response (log_2FC) of each putatively adaptive gene (n=1,775) while controlling for measurement error (F). The parallelism of evolution for each gene is then quantified as 1/F (Figure 1b; see Materials and Methods). A higher value of 1/F indicates a more parallel evolutionary response among replicated populations (Supplementary figure 2). Importantly, the measure of parallelism varies between genes (Figure 1c) and is positively correlated with the number of populations in which a significant change in gene expression is observed for each gene (Supplementary figure 3).

Pleiotropy is positively associated with the degree of parallelism in gene expression evolution

Pleiotropy is one of the factors that influence the degree of parallel evolution [7-10]. Therefore, we tested whether and how the pleiotropic effect of a gene influences the degree of parallelism of its expression evolution in a new environment. We correlated two different

measures of pleiotropy (1-τ (tissue specificity) and network connectivity; see materials and

methods) of each putatively adaptive gene (n=1,775) with the parallelism of its evolutionary

responses (1/F). We observed a significant positive correlation between the parallelism and

the degree of pleiotropy, (rho=0.26, p-value<2.2e-16 for 1-τ and rho=0.11, p-value<5.6e-07

for network connectivity) (Figure 2 and Supplementary figure 4a). This suggests that

pleiotropy may enhance parallel changes in gene expression during evolution.

Pleiotropy constrains the ancestral (natural) variation in gene expression

The negative effect of pleiotropy on genetic variation over longer time scales, mostly

between species, is well documented [13-16], but it not yet clear whether the same effect is

seen for adaptation from standing genetic variation. We reconstituted the ancestral population

from 189 isofemale lines [60] and estimated the expression variance (BCV^2; see materials

and methods) of each putatively adaptive gene (n=1,775) in two replicate sets of 20

individuals. Since the variance in gene expression is positively correlated with genetic

variation in cis-regulatory regions [65] and the heritability of gene expression is high [66], we

assumed that the observed variation in gene expression has a genetic basis, rather than being

noise (see discussion). We observed a highly significant negative correlation between the

level of pleiotropy and the magnitude of ancestral variation in gene expression (Figure 3a and

Supplementary figure 4b; rho=-0.34, p-value<2.2e-16 for 1-τ and rho=-0.37, p-value<2.2e-16

for network connectivity). To exclude that this correlation was driven by our filtering for

significant expression changes in at least three populations, we repeated these analyses with

all genes (n=9,882) and also observed a significant correlation (rho=-0.45, p-value<2.2e-16

for 1-τ and rho=-0.38, p-value<2.2e-16 for network connectivity, data not shown). Hence, we

suggest that the negative relationship between pleiotropy and polymorphism/diversity [32, 33]

(i.e. reduced variants, lower frequencies and/or lower allelic effects) can be extended to gene

expression.

Evolutionary parallelism is negatively associated with ancestral variation

For a better understanding of the factors determining parallel evolution, we examined the

influence of ancestral variation in gene expression on the parallel evolution of gene

expression. We observed a significant negative correlation between ancestral variation and

parallelism of the evolutionary response (rho=-0.22, p-value<2.2e-16; Figure 3b). We suggest

that this pattern could be explained by genetic redundancy [5, 67], which describes the

phenomenon that a polygenic trait can shift its phenotypic value through different combinations of contributing loci: When the diversity of the contributing loci is reduced (either through fewer loci, lower ancestral frequencies or smaller variation in effect size), fewer combinations of the contributing loci are available for adaptation, resulting in more parallel adaptive responses. To confirm this, we performed computer simulations of replicated polygenic adaptation and found that lower ancestral variation (i.e. fewer contributing loci) leads to more parallel trait evolution (Figure 4). Hence, that natural variation in the ancestral population could be one of factor determining the parallel adaptive response in replicate populations. Although we focused on the phenotypic response to match gene expression, it is interesting to note that a similar result was obtained for parallelism at the genetic level with different levels of standing genetic variation [68]. If pleiotropy reduces variation, the positive correlation between pleiotropy and parallelism may be an indirect (correlative) effect, rather than a direct consequence of pleiotropy.

Testing causality

Our previous analyses indicated that parallelism is affected by pleiotropy and ancestral variation. Because pleiotropy is negatively correlated with ancestral variation for both pleiotropy measures (Figure 3a and supplementary figure 4b), it is not clear whether pleiotropy affects parallelism directly or indirectly through the negative correlation with ancestral variation. We used causal analysis, which based on a Bayesian statistical framework [61], to disentangle the causal relationship between the three different parameters. Since the directionality of pleiotropic effects on ancestral variation has been previously demonstrated, and ancestral variation is unlikely to influence the pleiotropy of a gene, we considered five possible causal relationships between pleiotropic effects, ancestral variation, and parallelism (Figure 5). In the first model, parallelism is determined by ancestral variation, which is shaped by pleiotropic, but pleiotropy has no direct effect on parallelism. In model II, the pleiotropic effect affects ancestral variation and parallelism independently, but ancestral variation has no causal effect on parallelism. In model III, pleiotropy affects both ancestral variation and parallelism, and ancestral variation also determines parallelism. In model IV, parallelism is determined by ancestral variation and pleiotropy, but pleiotropic effects are assumed to have no effect on the ancestral variation. This serves as a null model to confirm the indirect effect of pleiotropy on parallelism via ancestral variation. We also add a global null model with no correlation between these three factors (model V, Figure 5). The analysis

was performed independently for two measures of pleiotropy (tissue specificity and network

connectivity).

For tissue specificity, our data were better explained by model III (i.e., model III has the

lowest BIC, Table 1), suggesting that both direct and indirect effects of pleiotropy determine

the degree of parallelism of adaptive gene expression evolution. On the other hand, when

pleiotropy was estimated by network connectivity, model I had a slightly lower BIC than

model III (Table 1). Thus, our causal analysis does not provide evidence for a direct effect of

network connectivity. We further quantified the strength of the direct and indirect effects of

pleiotropy on parallelism using a path analysis (see materials and methods). The path analysis

confirmed the significant direct effect of tissue specificity, but network connectivity has no

statistically significant direct effect on the parallelism of gene expression evolution (Table 2

and Table S1). Although the relevance of direct effects of pleiotropy on parallel gene

expression evolution differs for the two pleiotropy estimates, the similarity of the indirect

effect sizes for both pleiotropy measures is striking (Table 2). Taken together, these results

suggest that pleiotropy may enhance the parallel evolutionary response of gene expression

directly and indirectly through its influence on ancestral variation.

Discussion

This study examined the influence of pleiotropy on the parallelism of the adaptive response from standing genetic variation. By examining the role of natural variation, which is important for parallelism of short-term evolution, we extended the scope of previous studies. Because differences in genetic background and selection strength among replicated populations cannot be controlled in natural populations and may affect the degree of parallelism, we relied on experimental evolution for our study. We took advantage of ten replicated populations that have evolved from the same genetic background in the same hot environment for more than 100 generations [5].

The adaptive response during adaptation to the new hot temperature regime was measured by changes in gene expression. This approach has the advantage that pleiotropic effects and their associated fitness function, which are typically not known, are already included in the adaptive response. Furthermore, we were able to rely on two well-established measures of pleiotropy to study the interplay with parallelism and natural variation.

A key finding of our study is that that adaptation from standing genetic variation results in complex interplays between pleiotropy, ancestral variation, and parallelism. Of particular interest is the observed negative correlation between pleiotropy and ancestral variation in gene expression, which confirms previous observations [32, 33]. This suggests that the primary effect of pleiotropy is not to maintain variation. Rather, stronger purifying selection removes pleiotropic variants from the population, resulting in a reduced standing genetic variation. The reduced variation in more pleiotropic genes could reflect historical hard sweeps or purifying selection. A similar pattern is seen over longer evolutionary timescales: less sequence divergence in pleiotropic genes [13-16].

An important difference between adaptation from de novo mutations and standing genetic variation concerns the implications for parallel evolution. While pleiotropic genes are less likely to contribute to parallel evolution from de novo mutations [7], our results show that adaptation from standing genetic variation favors parallel selection responses of pleiotropic genes (Figure 2, Supplementary Figure 4a). It is important to note that many previous studies focused on sequence variation, whereas we used expression variation. While it is possible that these two types of variation behave differently, we do not believe that this is the case as

sequence polymorphism and expression variation are correlated [69]. Furthermore, a recent

study focusing on genomic changes [9] reached similar conclusions.

A very important assumption in our interpretation is that differences in gene expression

variation in the ancestral population reflect regulatory sequence polymorphism rather than

noise. The high correlation in the variance of gene expression between two replicate groups

of individuals indicates that our measures do not reflect technical noise but have a biological

basis. However, stochastic fluctuations in gene expression have also been observed among

cells in single-celled and multicellular organisms and the variance in gene expression differs

among genes [70-72]. In this case, the differences in gene expression variance are biological

in nature, but not associated with sequence variation. We do not believe that variation in the

control of gene expression levels among genes can explain our pattern because *Drosophila* is

a multicellular organism and we used whole body RNA-Seq data for our analysis. Therefore,

even if expression is controlled to different extents among genes, by averaging across

multiple cells, this does not affect our results. The only scenario we can imagine in which

gene expression noise could propagate across the many cells of a multicellular organism is

that during early development, differences in gene expression of a key regulator could affect

downstream gene expression throughout the entire organism. However, we do not think this

is a likely scenario, especially in the light of previous results showing that genetic variation in

cis-regulatory regions correlates with expression variation [65]. Furthermore, whole-body

gene expression in *Drosophila* is highly heritable, suggesting that gene expression is not

stochastic [66]. Nevertheless, we would like to point out that our observations are valid

irrespective of whether the correlation between parallelism and expression variation is the

result of genetic polymorphism or genetic control of expression noise.

Based on the significant correlation results, the question arose as to whether the positive

correlation between pleiotropy and parallelism, is causal (direct effect) or merely

correlational (indirect effect). Regardless of the measure of pleiotropy, our causal analysis

showed a consistent indirect effect of pleiotropy on parallelism (Table 1, 2). The estimated

indirect effects were very similar for both measures of pleiotropy (Table 2). This indirect

effect of pleiotropy could be explained by the reduced variation in the expression of more

pleiotropic genes in the ancestral population leading to a more parallel selection response of

gene expression. For the direct effect, however, the results differed between the two measures

of pleiotropy. Only tissue specificity had a significant direct effect, which was even larger

than the indirect effect (Table 2). No significant direct effect was found for network connectivitiy. The discrepancy between the two measures of pleiotropy is particularly interesting given their significant correlation (Supplementary Figure 1). This suggests that both measures capture aspects of pleiotropy that differ in their biological implications. The direct effects of pleiotropy, most likely arise from the positively correlated fitness effects of non-focal traits (synergistic pleiotropy). The synergistic pleiotropic effects result in a stronger fitness advantage, which translates into a more parallel selection response - as previously shown [22]. Given that temperature is one of the most important abiotic factors, it is expected that positive genetic correlations between temperature-adaptive traits have been selected over long evolutionary timescales, probably prior to the speciation of *D. simulans*. Thus, it is conceivable that positive pleiotropy for temperature adaptation related traits is common and thus determined the positive contribution of pleiotropy to the parallel evolution observed here.

In a concluding remark, we would like to emphasize that although pleiotropy and ancestral variation were found to be significantly associated with the degree of parallelism of gene expression evolution, it is likely that additional factors also contribute. For example, diminishing return epistasis, population size (drift), and the contribution of gene expression to the selected traits may also affect the parallelism of gene evolution. Further investigation of other potential factors will continue to advance our understanding of parallelism and the predictability of evolution.

References

1.   Colosimo PF, Hosemann KE, Balabhadra S, Villarreal G, Jr., Dickson M, Grimwood J, et al. Widespread parallel evolution in sticklebacks by repeated fixation of Ectodysplasin alleles. Science. 2005;307(5717):1928-33. Epub 2005/03/26. doi: 10.1126/science.1107239. PubMed PMID: 15790847.

2.   Westram AM, Faria R, Johannesson K, Butlin R. Using replicate hybrid zones to understand the genomic basis of adaptive divergence. Molecular ecology. 2021;30(15):3797-814. Epub 2021/02/28. doi: 10.1111/mec.15861. PubMed PMID: 33638231.

3.   van der Zee MJ, Whiting JR, Paris JR, Bassar RD, Travis J, Weigel D, et al. Rapid genomic convergent evolution in experimental populations of Trinidadian guppies (Poecilia reticulata). Evol Lett. 2022;6(2):149-61. Epub 2022/04/08. doi: 10.1002/evl3.272. PubMed PMID: 35386829; PubMed Central PMCID: PMCPMC8966473.

4.   Zong SB, Li YL, Liu JX. Genomic Architecture of Rapid Parallel Adaptation to Fresh Water in a Wild Fish. Molecular biology and evolution. 2021;38(4):1317-29. Epub 2020/11/05. doi: 10.1093/molbev/msaa290. PubMed PMID: 33146383; PubMed Central PMCID: PMCPMC8480189.

5.   Barghi N, Tobler R, Nolte V, Jaksic AM, Mallard F, Otte KA, et al. Genetic redundancy fuels polygenic adaptation in *Drosophila*. PLoS biology. 2019;17(2):e3000128. Epub 2019/02/05. doi: 10.1371/journal.pbio.3000128. PubMed PMID: 30716062.

6.   Barghi N, Hermisson J, Schlötterer C. Polygenic adaptation: a unifying framework to understand positive selection. Nature reviews Genetics. 2020;21(12):769-81. Epub 2020/07/01. doi: 10.1038/s41576-020-0250-z. PubMed PMID: 32601318.

7.   Bolnick DI, Barrett RDH, Oke KB, Rennison DJ, Stuart YE. (Non) Parallel Evolution. Annu Rev Ecol Syst. 2018;49:303-30.

8.   Conte GL, Arnegard ME, Best J, Chan YF, Jones FC, Kingsley DM, et al. Extent of QTL Reuse During Repeated Phenotypic Divergence of Sympatric Threespine Stickleback. Genetics. 2015;201(3):1189-200. Epub 2015/09/19. doi: 10.1534/genetics.115.182550. PubMed PMID: 26384359; PubMed Central PMCID: PMCPMC4649644.

9.   Rennison DJ, Peichel CL. Pleiotropy facilitates parallel adaptation in sticklebacks. Molecular ecology. 2022;31(5):1476-86. Epub 2022/01/09. doi: 10.1111/mec.16335. PubMed PMID: 34997980; PubMed Central PMCID: PMCPMC9306781.

10.   Stern DL. The genetic causes of convergent evolution. Nature reviews Genetics. 2013;14(11):751-64. Epub 2013/10/10. doi: 10.1038/nrg3483. PubMed PMID: 24105273.

11.   Fisher RA. The genetical theory of natural selection. Oxford: Clarendon Press; 1930.

12.   Orr HA. Adaptation and the cost of complexity. Evolution; international journal of organic evolution. 2000;54(1):13-20. Epub 2000/08/11. PubMed PMID: 10937178.

13.   Fraser HB, Hirsh AE, Steinmetz LM, Scharfe C, Feldman MW. Evolutionary rate in the protein interaction network. Science. 2002;296(5568):750-2. Epub 2002/04/27. doi: 10.1126/science.1068696. PubMed PMID: 11976460.

14.   Hahn MW, Kern AD. Comparative genomics of centrality and essentiality in three eukaryotic protein-interaction networks. Molecular biology and evolution.

2005;22(4):803-6. Epub 2004/12/24. doi: 10.1093/molbev/msi072. PubMed PMID:
15616139.

15.   Masalia RR, Bewick AJ, Burke JM. Connectivity in gene coexpression networks
negatively correlates with rates of molecular evolution in flowering plants. PloS one.
2017;12(7):e0182289. Epub 2017/08/02. doi: 10.1371/journal.pone.0182289. PubMed
PMID: 28759647; PubMed Central PMCID: PMCPMC5536297.

16.   Josephs EB, Wright SI, Stinchcombe JR, Schoen DJ. The Relationship between
Selection, Network Connectivity, and Regulatory Variation within a Population of
Capsella grandiflora. Genome biology and evolution. 2017;9(4):1099-109. doi:
10.1093/gbe/evx068. PubMed PMID: WOS:000406755800024.

17.   Hansen TF. Is modularity necessary for evolvability? Remarks on the
relationship between pleiotropy and evolvability. Biosystems. 2003;69(2-3):83-94. doi:
Pii S0303-2647(02)00132-6
Doi 10.1016/S0303-2647(02)00132-6. PubMed PMID: WOS:000182602700002.

18.   Blows MW. A tale of two matrices: multivariate approaches in evolutionary
biology. Journal of evolutionary biology. 2007;20(1):1-8. Epub 2007/01/11. doi:
10.1111/j.1420-9101.2006.01164.x. PubMed PMID: 17209986.

19.   Lande R, Arnold SJ. The Measurement of Selection on Correlated Characters.
Evolution; international journal of organic evolution. 1983;37(6):1210-26. Epub
1983/11/01. doi: 10.1111/j.1558-5646.1983.tb00236.x. PubMed PMID: 28556011.

20.   Lande R. Quantitative Genetic Analysis of Multivariate Evolution, Applied to
Brain:Body Size Allometry. Evolution; international journal of organic evolution.
1979;33(1Part2):402-16. Epub 1979/03/01. doi: 10.1111/j.1558-
5646.1979.tb04694.x. PubMed PMID: 28568194.

21.   Thorholludottir DAV, Nolte V, Schlötterer C. Temperature driven gene expression
evolution in natural and laboratory populations highlights the crucial role of correlated
fitness effects for polygenic adaptation. Evolution; international journal of organic
evolution. 2023. Epub 2023/07/17. doi: 10.1093/evolut/qpad132. PubMed PMID:
37455661.

22.   MacPherson A, Nuismer SL. The probability of parallel genetic evolution from
standing genetic variation. Journal of evolutionary biology. 2017;30(2):326-37. Epub
2016/11/02. doi: 10.1111/jeb.13006. PubMed PMID: 27801996.

23.   Rivas-Sanchez DF, Melo-Florez L, Aragon A, Pardo-Diaz C, Salazar C, Montgomery
SH, et al. Parallel evolution of behavior, physiology, and life history associated with
altitudinal shifts in forest type in Heliconius butterflies. Evolution; international journal
of organic evolution. 2023;77(6):1458-67. Epub 2023/04/19. doi:
10.1093/evolut/qpad062. PubMed PMID: 37075171.

24.   Huang Y, Lack JB, Hoppel GT, Pool JE. Parallel and population-specific gene
regulatory evolution in cold-adapted fly populations. Genetics. 2021;218(3). Epub
2021/05/15. doi: 10.1093/genetics/iyab077. PubMed PMID: 33989401; PubMed
Central PMCID: PMCPMC8864734.

25.   Heckley AM, Pearce AE, Gotanda KM, Hendry AP, Oke KB. Compiling forty years
of guppy research to investigate the factors contributing to (non)parallel evolution.
Journal of evolutionary biology. 2022;35(11):1414-31. Epub 2022/09/14. doi:
10.1111/jeb.14086. PubMed PMID: 36098479.

26.   Fischer EK, Song Y, Hughes KA, Zhou W, Hoke KL. Nonparallel transcriptional
divergence during parallel adaptation. Molecular ecology. 2021;30(6):1516-30. Epub
2021/02/02. doi: 10.1111/mec.15823. PubMed PMID: 33522041.

27.   Elmer KR, Fan S, Kusche H, Spreitzer ML, Kautt AF, Franchini P, et al. Parallel
evolution of Nicaraguan crater lake cichlid fishes via non-parallel routes. Nature
communications. 2014;5:5168. Epub 2014/10/28. doi: 10.1038/ncomms6168. PubMed
PMID: 25346277.
28.   Butlin RK, Saura M, Charrier G, Jackson B, Andre C, Caballero A, et al. Parallel
evolution of local adaptation and reproductive isolation in the face of gene flow.
Evolution; international journal of organic evolution. 2014;68(4):935-49. Epub
2013/12/05. doi: 10.1111/evo.12329. PubMed PMID: 24299519; PubMed Central
PMCID: PMCPMC4261988.
29.   Hsu SK, Belmouaden C, Nolte V, Schlötterer C. Parallel gene expression evolution
in natural and laboratory evolved populations. Molecular ecology. 2021;30(4):884-94.
Epub 2020/09/27. doi: 10.1111/mec.15649. PubMed PMID: 32979867; PubMed
Central PMCID: PMCPMC7891358.
30.   Otte KA, Nolte V, Mallard F, Schlötterer C. The genetic architecture of
temperature adaptation is shaped by population ancestry and not by selection regime.
Genome biology. 2021;22(1). doi: ARTN 211
10.1186/s13059-021-02425-9. PubMed PMID: WOS:000673395800001.
31.   Papakostas S, Vollestad LA, Bruneaux M, Aykanat T, Vanoverbeke J, Ning M, et al.
Gene pleiotropy constrains gene expression changes in fish adapted to different thermal
conditions. Nature communications. 2014;5:4071. Epub 2014/06/04. doi:
10.1038/ncomms5071. PubMed PMID: 24892934; PubMed Central PMCID:
PMCPMC4059932.
32.   Lemos B, Meiklejohn CD, Hartl DL. Regulatory evolution across the protein
interaction network. Nature genetics. 2004;36(10):1059-60. Epub 2004/09/21. doi:
10.1038/ng1427. PubMed PMID: 15378060.
33.   Mahler N, Wang J, Terebieniec BK, Ingvarsson PK, Street NR, Hvidsten TR. Gene
co-expression network connectivity is an important determinant of selective constraint.
PLoS genetics. 2017;13(4). doi: ARTN e1006402
10.1371/journal.pgen.1006402. PubMed PMID: WOS:000402549200001.
34.   Nourmohammad A, Rambeau J, Held T, Kovacova V, Berg J, Lassig M. Adaptive
Evolution of Gene Expression in Drosophila. Cell Rep. 2017;20(6):1385-95. doi:
10.1016/j.celrep.2017.07.033. PubMed PMID: 28793262.
35.   Hsu SK, Jaksic AM, Nolte V, Lirakis M, Kofler R, Barghi N, et al. Rapid sex-specific
adaptation to high temperature in *Drosophila*. Elife. 2020;9. Epub 2020/02/23. doi:
10.7554/eLife.53237. PubMed PMID: 32083552; PubMed Central PMCID:
PMCPMC7034977.
36.   Yu X, Wei P, Zhao S, Chen Z, Li X, Zhang W, et al. Population transcriptomics
uncover the relative roles of positive selection and differential expression in
Batrachium bungei adaptation to the Qinghai-Tibetan plateau. Plant Cell Rep.
2023;42(5):879-93. Epub 2023/03/28. doi: 10.1007/s00299-023-03005-w. PubMed
PMID: 36973418.
37.   Velotta JP, Wegrzyn JL, Ginzburg S, Kang L, Czesny S, O'Neill RJ, et al.
Transcriptomic imprints of adaptation to fresh water: parallel evolution of
osmoregulatory gene expression in the Alewife. Molecular ecology. 2017;26(3):831-48.
Epub 2016/12/25. doi: 10.1111/mec.13983. PubMed PMID: 28012221.
38.   Tang Q, Gu Y, Zhou X, Jin L, Guan J, Liu R, et al. Comparative transcriptomics of 5
high-altitude vertebrates and their low-altitude relatives. Gigascience. 2017;6(12):1-9.
Epub 2017/11/18. doi: 10.1093/gigascience/gix105. PubMed PMID: 29149296;
PubMed Central PMCID: PMCPMC5729692.

39.   O'Neil ST, Dzurisin JD, Williams CM, Lobo NF, Higgins JK, Deines JM, et al. Gene
expression in closely related species mirrors local adaptation: consequences for
responses to a warming world. Molecular ecology. 2014;23(11):2686-98. Epub
2014/04/29. doi: 10.1111/mec.12773. PubMed PMID: 24766086.
40.   Margres MJ, Wray KP, Hassinger ATB, Ward MJ, McGivern JJ, Moriarty Lemmon E,
et al. Quantity, Not Quality: Rapid Adaptation in a Polygenic Trait Proceeded Exclusively
through Expression Differentiation. Molecular biology and evolution.
2017;34(12):3099-110. Epub 2017/09/30. doi: 10.1093/molbev/msx231. PubMed
PMID: 28962003.
41.   McShea DW. Functional complexity in organisms: Parts as proxies. Biol Philos.
2000;15(5):641-68. doi: Doi 10.1023/A:1006695908715. PubMed PMID:
WOS:000165088500002.
42.   Proulx SR, Promislow DE, Phillips PC. Network thinking in ecology and evolution.
Trends in ecology & evolution. 2005;20(6):345-53. Epub 2006/05/17. doi:
10.1016/j.tree.2005.04.004. PubMed PMID: 16701391.
43.   He X, Zhang J. Toward a molecular understanding of pleiotropy. Genetics.
2006;173(4):1885-91. Epub 2006/05/17. doi: 10.1534/genetics.106.060269. PubMed
PMID: 16702416; PubMed Central PMCID: PMCPMC1569710.
44.   Mank JE, Hultin-Rosenberg L, Zwahlen M, Ellegren H. Pleiotropic constraint
hampers the resolution of sexual antagonism in vertebrate gene expression. Am Nat.
2008;171(1):35-43. Epub 2008/01/04. doi: 10.1086/523954. PubMed PMID:
18171149.
45.   Leader DP, Krause SA, Pandit A, Davies SA, Dow JAT. FlyAtlas 2: a new version of
the Drosophila melanogaster expression atlas with RNA-Seq, miRNA-Seq and sex-
specific data. Nucleic acids research. 2018;46(D1):D809-D15. Epub 2017/10/27. doi:
10.1093/nar/gkx976. PubMed PMID: 29069479; PubMed Central PMCID:
PMCPMC5753349.
46.   Dean R, Mank JE. Tissue Specificity and Sex-Specific Regulatory Variation Permit
the Evolution of Sex-Biased Gene Expression. Am Nat. 2016;188(3):E74-84. Epub
2016/08/09. doi: 10.1086/687526. PubMed PMID: 27501094.
47.   Allen SL, Bonduriansky R, Chenoweth SF. Genetic constraints on
microevolutionary divergence of sex-biased gene expression. Philos Trans R Soc Lond B
Biol Sci. 2018;373(1757). Epub 2018/08/29. doi: 10.1098/rstb.2017.0427. PubMed
PMID: 30150225; PubMed Central PMCID: PMCPMC6125734.
48.   Watanabe K, Stringer S, Frei O, Mirkov MU, de Leeuw C, Polderman TJC, et al. A
global overview of pleiotropy and genetic architecture in complex traits (vol 51,1339,
2019). Nature genetics. 2020;52(3):353-. doi: 10.1038/s41588-019-0571-z. PubMed
PMID: WOS:000511537400002.
49.   Hamala T, Gorton AJ, Moeller DA, Tiffin P. Pleiotropy facilitates local adaptation
to distant optima in common ragweed (Ambrosia artemisiifolia). PLoS genetics.
2020;16(3):e1008707. Epub 2020/03/27. doi: 10.1371/journal.pgen.1008707. PubMed
PMID: 32210431; PubMed Central PMCID: PMCPMC7135370.
50.   Marbach D, Roy S, Ay F, Meyer PE, Candeias R, Kahveci T, et al. Predictive
regulatory models in Drosophila melanogaster by integrative inference of
transcriptional networks. Genome research. 2012;22(7):1334-49. Epub 2012/03/30.
doi: 10.1101/gr.127191.111. PubMed PMID: 22456606; PubMed Central PMCID:
PMCPMC3396374.
51.   Mao YX, Li Y, Yang Z, Xu N, Zhang S, Wang X, et al. Comparative transcriptome
analysis between rhesus macaques (Macaca mulatta) and crab-eating macaques (M.

fasicularis). Zool Res. 2024;45(2):299-310. Epub 2024/03/15. doi:
10.24272/j.issn.2095-8137.2023.322. PubMed PMID: 38485500; PubMed Central
PMCID: PMCPMC11017088.
52. Davidson RM, Gowda M, Moghe G, Lin H, Vaillancourt B, Shiu SH, et al.
Comparative transcriptomics of three Poaceae species reveals patterns of gene
expression evolution. The Plant journal : for cell and molecular biology.
2012;71(3):492-502. Epub 2012/03/27. doi: 10.1111/j.1365-313X.2012.05005.x.
PubMed PMID: 22443345.
53. Julca I, Ferrari C, Flores-Tornero M, Proost S, Lindner AC, Hackenberg D, et al.
Comparative transcriptomic analysis reveals conserved programmes underpinning
organogenesis and reproduction in land plants. Nat Plants. 2021;7(8):1143-59. Epub
2021/07/14. doi: 10.1038/s41477-021-00958-2. PubMed PMID: 34253868.
54. Jaksic AM, Karner J, Nolte V, Hsu SK, Barghi N, Mallard F, et al. Neuronal function
and dopamine signaling evolve at high temperature in Drosophila. Molecular biology
and evolution. 2020. Epub 2020/05/14. doi: 10.1093/molbev/msaa116. PubMed PMID:
32402077.
55. Nouhaud P, Tobler R, Nolte V, Schlötterer C. Ancestral population reconstitution
from isofemale lines as a tool for experimental evolution. Ecol Evol. 2016;6(20):7169-
75. doi: 10.1002/ece3.2402. PubMed PMID: 27895897; PubMed Central PMCID:
PMCPMC5114691.
56. Charlesworth B, Charlesworth D. Elements of Evolutionary Genetics. Greenwood
Village, Colorado: Roberts and Company Publishers; 2010.
57. Nouhaud P, Mallard F, Poupardin R, Barghi N, Schlötterer C. High-throughput
fecundity measurements in *Drosophila*. Sci Rep. 2018;8:4469.
58. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics.
2010;26(1):139-40. doi: 10.1093/bioinformatics/btp616. PubMed PMID: 19910308;
PubMed Central PMCID: PMC2796818.
59. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and
powerful approach to multiple testing. J R Stat Soc Ser B. 1995;57:289-300.
60. Lai WY, Schlötterer C. Evolution of phenotypic variance in response to a novel
hot environment. Molecular ecology. 2022;31(3):934-45. Epub 2021/11/15. doi:
10.1111/mec.16274. PubMed PMID: 34775658.
61. Schadt EE, Lamb J, Yang X, Zhu J, Edwards S, GuhaThakurta D, et al. An
integrative genomics approach to infer causal associations between gene expression
and disease. Nature genetics. 2005;37(7):710-7. doi: 10.1038/ng1589. PubMed PMID:
WOS:000230196400017.
62. Wright S. Correlation and causation Part I. Method of path coefficients. J Agric
Res. 1920;20:0557-85. PubMed PMID: WOS:000188347400034.
63. Vlachos C, Kofler R. MimicrEE2: Genome-wide forward simulations of Evolve and
Resequencing studies. PLoS computational biology. 2018;14(8):e1006413. Epub
2018/08/17. doi: 10.1371/journal.pcbi.1006413. PubMed PMID: 30114186; PubMed
Central PMCID: PMCPMC6112681.
64. Bohutinska M, Vlcek J, Yair S, Laenen B, Konecna V, Fracassetti M, et al. Genomic
basis of parallel adaptation varies with divergence in *Arabidopsis* and its relatives.
Proceedings of the National Academy of Sciences of the United States of America.
2021;118(21). Epub 2021/05/19. doi: 10.1073/pnas.2022713118. PubMed PMID:
34001609; PubMed Central PMCID: PMCPMC8166048.

65.   Hämälä T, Guiltinan MJ, Marden JH, Maximova SN, dePamphilis CW, Tiffin P. Gene
Expression Modularity Reveals Footprints of Polygenic Adaptation in *Theobroma cacao*.
Molecular biology and evolution. 2020;37(1):110-23. Epub 2019/09/11. doi:
10.1093/molbev/msz206. PubMed PMID: 31501906.
66.   Ayroles JF, Carbone MA, Stone EA, Jordan KW, Lyman RF, Magwire MM, et al.
Systems genetics of complex traits in *Drosophila melanogaster*. Nature genetics.
2009;41(3):299-307. doi: 10.1038/ng.332. PubMed PMID: WOS:000263640200010.
67.   Laruson AJ, Yeaman S, Lotterhos KE. The Importance of Genetic Redundancy in
Evolution. Trends in ecology & evolution. 2020. Epub 2020/05/23. doi:
10.1016/j.tree.2020.04.009. PubMed PMID: 32439075.
68.   Thompson KA, Osmond MM, Schluter D. Parallel genetic evolution and speciation
from standing variation. Evol Lett. 2019;3(2):129-41. Epub 2019/07/11. doi:
10.1002/evl3.106. PubMed PMID: 31289688; PubMed Central PMCID:
PMCPMC6591551.
69.   Lawniczak MK, Holloway AK, Begun DJ, Jones CD. Genomic analysis of the
relationship between gene expression variation and DNA polymorphism in Drosophila
simulans. Genome biology. 2008;9(8):R125. Epub 2008/08/14. doi: 10.1186/gb-2008-
9-8-r125. PubMed PMID: 18700012; PubMed Central PMCID: PMCPMC2575515.
70.   Swain PS, Elowitz MB, Siggia ED. Intrinsic and extrinsic contributions to
stochasticity in gene expression. Proceedings of the National Academy of Sciences of the
United States of America. 2002;99(20):12795-800. Epub 2002/09/19. doi:
10.1073/pnas.162041399. PubMed PMID: 12237400; PubMed Central PMCID:
PMCPMC130539.
71.   Elowitz MB, Levine AJ, Siggia ED, Swain PS. Stochastic gene expression in a single
cell. Science. 2002;297(5584):1183-6. Epub 2002/08/17. doi:
10.1126/science.1070919. PubMed PMID: 12183631.
72.   Blake WJ, M KA, Cantor CR, Collins JJ. Noise in eukaryotic gene expression.
Nature. 2003;422(6932):633-7. Epub 2003/04/11. doi: 10.1038/nature01546. PubMed
PMID: 12687005.
788
789

Acknowledgments

We thank Viola Nolte for preparing all RNA-Seq libraries and supervising the maintenance of the evolution experiment. We thank Dagný Ásta Rúnarsdóttir for the suggestions on the early version of the manuscript and all member of the Institut für Populationsgenetik for discussion. Ana Marija Jakšić, Neda Barghi, François Mallard and Kathrin Otte performed the common garden experiment. Illumina sequencing was done at the VBCF NGS Unit (www.vbcf.ac.at). This research was funded in whole or in part by the Austrian Science Fund (FWF) [W1225, P32935] and the European Research Council (ERC) [ArchAdapt]. For open access PURPOSES, the author has applied a CC BY public copyright license to any author-accepted manuscript version arising from this submission..

Author contribution
W.Y.L, S.K.H. and C.S. conceived the study. A.F. contributed statistical advice, in particular for the causal analysis. W.Y.L. performed the data analysis. S.K.H. performed the simulations. W.Y.L. and C.S. wrote the manuscript. W.Y.L., S.K.H., A.F. and C.S. revised and edited the final manuscript.

Competing interests
The authors declare no competing interests.

Data accessibility statement
All sequencing data are available in European Nucleotide Archive (ENA) under the accession number PRJEB35504, PRJEB35506 and PRJEB37011. Other raw data and scripts are available on github (https://github.com/cloudweather34/pleiotropy).

814
Figure 1. Overview of the experimental procedures (a) and the variation in parallel gene

expression evolution across genes (b,c). **a.** Experimental Evolution: ten replicated

populations seeded from one common founder population have been evolving for >100

generations in a hot laboratory environment at 18 and 28°C. Common Garden Experiment: at

the 103[rd] generation of the evolution experiment, the ten evolved populations (each with three

biological replicates) were maintained together with the reconstituted ancestral population

(with five biological replicates) for two generations in the same environment as in the

evolution experiment. After two generations in common garden, 50 males from each

biological replicates were pooled and subjected to RNA-Seq. **b**. Replicate frequency

spectrum. Number of populations (x-axis) in which a given gene experienced a significant

change in gene expression. The y-axis indicates the number of genes in each category. Most

of the genes experienced a significant change in gene expression in few evolved populations

while much fewer genes were significant in all 10 populations. This pattern suggests that the

parallelism in gene expression evolution differs across genes. **c.** The distribution of gene

expression evolution parallelism ($\log(1/F)$) across genes. Larger $\log(1/F)$ values indicate

more parallel evolution of a gene. The exhibit variation suggests that genes varied in their

parallelism of its expression evolution.

Figure 2. Association between the magnitude of evolution parallelism and the strength of pleiotropy. The distribution of evolution parallelism (log(1/F)) of different genes is shown in boxplots binned by their strength of pleiotropy (1-). Overall, the strength of pleiotropy was positively correlated with evolution parallelism (=0.26, p-value<2.2e-16).

838
839

Figure 3. Association between ancestral variation in gene expression with the strength of pleiotropy (1-τ) (a) and with the evolution parallelism (b). a. The distribution of gene expression variation in the ancestral population (log (BCV^2)) of different genes is shown in boxplots binned by their strength of pleiotropy (1-τ). Overall, the strength of pleiotropy was negatively correlated with ancestral variation in gene expression (ρ=-0.34, p-value<2.2e-16). **b.** The distribution of evolutionary parallelism (log(1/F)) of different genes is shown in boxplots binned by their strength of ancestral variation. The strength of parallelism is negatively correlated with the ancestral variation in gene expression (ρ=-0.22, p-value<2.2e-16).

Figure 4. Computer simulations illustrate the negative effect of ancestral variation on parallel evolution. Computer simulations assume a fitness related trait determined by the expression of four genes. The expression of these four genes is determined by a different number of genetic loci, reflecting the influence of pleiotropy on ancestral genetic variation. A shift in trait optimum was used to illustrate the connection between ancestral variation in expression (x-axis) and parallelism of expression change across ten evolved populations (y-axis). The negative correlation between them (rho = -0.26, p-value < 1.77e-07) suggests that the gene with less ancestral variation are resulting in more parallel responses.

862
Figure 5. Schematic figure illustrating the causal models evaluated. Five possible
relationships between pleiotropic effects, ancestral variation, and evolutionary parallelism. $\mathcal{L}$
denotes the likelihood of each model given the data. P is the probability or conditional
probability of the measurements for each gene; *Pa* is the parallelism level and *A* is the level
of ancestral variation in gene expression. *Pl* is the pleiotropic effects. See materials and
methods for a more detailed description.
869

870 **Table 1. BIC value for Model I-V. The model with the smallest BIC is the one best**
871 **support by the data.**
872

Model	BIC value (Network connectivity)	BIC value (Tissue specificity)
I	9258.395	8958.125
II	9316.852	8972.626
III	9264.769	8939.342
IV	9490.017	9162.090
V	9550.309	9247.540

873
874

Table 2. The size of direct and indirect pleiotropic effects on the evolutionary parallelism.

	Direct effect	Indirect effect
1-τ	0.127	0.055 (-0.181, -0.332) *
Network connectivity	0.026	0.068 (-0.210, -0.337) *

* Coefficient estimates (β_2, β_3) whose product provides the indirect effect.