Genomic data reveal a north-south split and introgression history of blood fluke (*Schistosoma haematobium*) populations from across Africa

Roy N Platt II[1], Egie E. Enabulele[1], Ehizogie Adeyemi[2], Marian O Agbugui[3], Oluwaremilekun G Ajakaye[4], Ebube C Amaechi[5], Chika E Ejikeugwu[6], Christopher Igbeneghu[7], Victor S Njom[8], Precious Dlamini[9], Grace A. Arya[1], Robbie Diaz[1], Muriel Rabone[10], Fiona Allan[10], Bonnie Webster[10], Aidan Emery[10], David Rollinson[10, 11], Timothy J.C. Anderson[1]

[1] Texas Biomedical Research Institute, San Antonio TX, United States

[2] Department of Pathology, University of Benin Teaching Hospital, Edo State, Nigeria

[3] Department of Biological Sciences, Edo State University, Uzairue, Nigeria

[4] Department of Animal and Environmental Biology, Adekunle Ajasin University, Nigeria

[5] Department of Zoology, University of Ilorin, Kwara State, Nigeria

[6] Enugu State University of Science and Technology, Nigeria

[7] Department of Medical Laboratory Science, Ladoke Akintola University of Technology, Nigeria

[8] Department of Applied Biology and Biotechnology, Enugu State University of Science and Technology, Nigeria

[9] Central Public Health Offices, Manzini, Swaziland

[10] Science Department, Natural History Museum, London, United Kingdom

[11] Global Schistosomiasis Alliance, London, United Kingdom

Keywords: schistosomes, parasite, *Schistosoma bovis*, hybridization, introgression desert, admixture

Abstract: The human parasitic fluke, *Schistosoma haematobium* hybridizes with the livestock parasite *S. bovis* in the laboratory, but the extent of hybridization in nature is unclear. We analyzed 34.6 million single nucleotide variants in 162 samples from 18 African countries, revealing a sharp

genetic discontinuity between northern and southern *S. haematobium*. We found no evidence for

recent hybridization. Instead the data reveal admixture events that occurred 257-879 generations

ago in northern *S. haematobium* populations. Fifteen introgressed *S. bovis* genes are

approaching fixation in northern *S. haematobium* with four genes potentially driving adaptation.

We identified 19 regions that were resistant to introgression; these were enriched on the sex

chromosomes. These results (i) demonstrate strong barriers to gene flow between these species,

(ii) indicate that hybridization may be less common than currently envisaged, but (iii) reveal

profound genomic consequences of interspecific hybridization between schistosomes of medical

and veterinary importance.

Funding - This work was funded by NIH NIAID (R01 AI166049).

Introduction

Hybridization and the transfer of alleles via introgression is an important source of genetic variation between species (1). This process allows for allelic variants, which have already been preselected in a donor species, to be introduced into the genome of a recipient species in a single generation. By comparison, it may take multiple generations for random mutation and selection to deliver comparable levels of genetic variation in the absence of introgression (2). As a result, introgressive hybridization, can result in rapid evolution of new genetic traits in hybridizing species. Hybridization between human and animal parasites can lead to the emergence of parasites with novel traits such as increased pathogenicity (3), expanded host range (4), altered transmission dynamics (5) and drug resistance (6). Understanding the frequency and impact of such hybridization events is critical for devising effective disease intervention strategies.

Members of the blood fluke genus *Schistosoma* parasitize a range of mammal species and cause substantial morbidity and economic loss (7). One pair of species, *S. haematobium*, a human parasite, and *S. bovis*, an ungulate parasite common in domestic livestock, are genetically divergent (3-5% and 18% divergent in the nuclear and mitochondrial genomes respectively), but can hybridize and produce viable offspring when given the opportunity (8). Given the close proximity between humans and their livestock and the regular use of the same water bodies, the potential for hybridization between these species is a particular concern and a significant effort has been mounted to identify, monitor and map *S. haematobium* and *S. bovis* hybrids (9). Multiple studies have reported mitochondrial and/or ribosomal DNA from *S. bovis* in *S. haematobium* populations (for examples see 8, 10, 11, 12). The high frequency of individuals with discordant mitochondrial and nuclear markers has been used to argue that hybridization is common and that the zoonotic threat of *S. bovis* should be considered in human schistosomiasis control programs (13).

However, several more recent multi-marker and genomic studies using single nucleotide variants (SNVs), microsatellites markers and whole genome sequence assemblies have suggested that hybridization between *S. haematobium* and *S. bovis* may not occur as frequently as previously postulated. Exome (14), whole genome (15), and microsatellite data (16), and others (17-20) failed to identify evidence of contemporary hybridization in field-collected parasites. Instead, these studies indicate that *S. bovis* and *S. haematobium* are genetically distinct and do not hybridize frequently but evidence for historical hybridization is clearly evident within genomes of *S.*

haematobium. As a result, some *S. bovis* genes have introgressed into the *S. haematobium*
population and reached high frequency; evidence of a potential, adaptive introgression event (14,
15, 17, 19).

In this study, we build upon previous work and try to address knowledge gaps by analyzing a
comprehensive dataset of 34.6 million genome-wide SNVs from *S. bovis* (n=21) and *S.*
haematobium (n=141) samples collected from 18 countries across the African continent. Many of
these samples presented discordant mitochondrial and ribosomal DNA profiles and were
categorized as *S. haematobium-bovis* hybrids. This expanded dataset, and recent availability of
a high-quality *S. haematobium* genome assembly (21), allows for a more detailed examination of
the genetic relationships between these two species and the potential consequences of
hybridization on their evolution and epidemiology. Our results: (i) reveal a strong discontinuity
between northern and southern *S. haematobium* populations; (ii) define similar genomic
introgression profiles in *S. haematobium* sampled from locations 3,002 Km apart; (iii) fine-map
introgressed genome regions and identify putative genes driving adaptive introgression; (iv)
identify two distinct lineages of *S. bovis*-like mitochondrial DNA in northern *S. haematobium*,
consistent with rare, ancient introgression and (v) identify "introgression deserts" on the ZW
chromosomes consistent with the sex chromosomes maintaining species integrity. These results
enhance our understanding of *Schistosoma spp.* epidemiology, with important implications for
control efforts.

Results

DNA Sequencing and Genotyping – We examined 219 *Schistosoma* samples from 24 countries
(Figure 1A). Just over 80% (n=176) of the samples were collected as part of this study with the
remaining 43 samples made available through open access resources. Median genome coverage
per sample was 29.7x. After filtering, the final dataset contained 35,817,757 total SNVs, 7,206,957
common (minor allele frequency; MAF>0.05%) SNVs, and 446,162 common unlinked SNVs
genotyped from 162 samples (141 *S. haematobium* and 21 *S. bovis*; Figure 1B). NCBI Short Read
Archive (SRA) accessions and sample metadata are available in Supplemental Table 1.

Population structure and ancestry - We examined relationships among samples with a PCA of
355,715 unlinked, common, autosomal SNVs (Figure 1C). Each of our samples fell into one of 3
K-means clusters along PC1 and PC2. The three clusters corresponded with (a) *S. haematobium*

individuals from northern Africa, (b) southern Africa, and (c) all *S. bovis* samples. The northern

population includes samples collected in Cameroon, Cote d' Ivoire, Egypt Gambia, Guinea

Bissau, Liberia, Mali, Niger, Nigeria, Senegal and Sudan. The southern population includes

samples collected from Angola, Eswatini, Kenya, Madagascar, Namibia, Tanzania, Uganda,

Zambia, and Zanzibar. In general, the equator was used to delineate the northern and southern

populations. No samples were placed intermediate between the *S. haematobium* and *S. bovis*

clusters: we found no evidence for F1 *S. haematobium-bovis* hybrids among these samples. The

weighted, Weir and Cockerham F_{ST} (22, 23) between the *S. bovis* and *S. haematobium* samples

was high ($F_{ST} \geq 0.74\text{-}0.79$). We observed strong subdivision between northern and SE *S.*

haematobium populations ($F_{ST} = 0.16$; Figure 1C) with multiple peaks (Figure 2D). There were

275,657 sites showing fixed differences ($F_{ST} = 1$) between *S. bovis* and *S. haematobium*

(Supplemental Table 2).

We used Admixture v1.3.0 (24) to quantify ancestry among the samples (Figure 1D, Supplemental

Figure 1). We found that when $k = 2$, *S. bovis* and south African *S. haematobium* individuals were

exclusively assigned different ancestry components. By contrast *S. haematobium* samples

collected from north Africa was a composite of the two population components including 0.5-

26.2% (median = 4.2%) of the *S. bovis* population component. Nigeria was an outlier (Kruskal-

Wallis H test statistic = 7915, *P*-value = 0.0049; Supplemental Figure 2), showing significantly

higher levels of introgression together with extremely high variance in introgression levels (2.8 –

26.2%): further analyses of Nigerian samples will be of considerable interest.

<u>Reference biases</u> - We tested the data for read-mapping reference biases, which could occur if

non-*haematobium* species map poorly to the *S. haematobium* reference assembly

(GCF_000699445.3). Mapping rates were 77.7%, 82.8% and 76.3% for *S. bovis*, northern *S.*

haematobium and southern *S. haematobium* populations (Figure 1A). A *t*-Test failed to identify

differential mapping rates between *S. haematobium* and *S. bovis* (*p* = 0.397), suggesting that

reference bias does not significantly contribute to the results observed.

<u>Phylogenetics</u> - The species tree generated using SVDquartets and nuclear SNVs revealed a

well-resolved topology. We examined 2,500,000 random quartets, representing 8.82% of all

possible distinct quartets. Of these, 18.5% (n = 463,571) were incompatible with the final tree

(Figure 3). *S. haematobium* and *S. bovis* were resolved into two clades and *S. haematobium*

population tips generally fell into clades reflecting geographic relationships.

On average, we were able to assemble 15,558.4 bp of the mitochondrial genome for each sample 4,757 of which were phylogenetically informative sites. The mitochondrial phylogeny (Figure 4) reveals two major mitochondrial haplotypes, one containing *S. haematobium* individuals from across Africa and another clade containing all of the *S. bovis* and 38 north African *S. haematobium*. The presence of *S. haematobium* samples within a larger *S. bovis* clade is consistent with *S. bovis* mitochondrial introgression into *S. haematobium* that has been frequently reported in field samples (ex. 25). Within the *S. bovis* clade, all *S. haematobium* samples with the introgressed *S. bovis* mitochondria fell into two monophyletic groups, clades "A" and "B". mtDNA haplotypes from these two clades were from samples widely distributed in northern Africa. For example, the same clade "A" haplotype was found in samples from Egypt, Niger and Cote d' Ivoire (>3,300 km apart). The clade "B" haplotype was found in Niger, Nigeria and Cote d' Ivoire, a linear distance of 1,171 km. Bootstrap support for each of these major clades was strong (100%). Phylogenetic trees in Newick format are available in the supplementary materials.

Hybridization and Introgression - We used four methods to identify signatures of hybridization and introgression between *S. haematobium* and *S. bovis*. These methods include the f_3, *D*-statistic, local ancestry assignment (RFmix; 26) and phylogenetic discordance (TWISST; 27). First, the (NW: SE, Sb) f_3 =-0.128 (SE= $0.8e^{-3}$, z-score=-156.4) was significantly negative indicating the north African *S. haematobium* population contains *S. bovis* ancestry. Next, we used the *D*-statistic to test for introgression between *S. haematobium* and *S. bovis* while accounting for lineage sorting (Figure 2E). We averaged *D* in 10Kb blocks. *D* was significantly positive (*D*=0.46, σ_M=0.007, n=30,278) indicating introgression between north African *S. haematobium* and *S. bovis*.

We examined the landscape of introgression across the genome using local ancestry with RFmix. We used 38 southern African *S. haematobium* lacking *S. bovis* introgression and 13 *S. bovis* samples to serve as reference panels for "pure" *S. haematobium* and *S. bovis*. RFmix results showed that ancestry across the genome was not uniformly distributed in the north African population (Figure 2A). Within the north African population, *S. bovis* ancestry blocks ranged in frequency from 0-100% at any particular locus. Each north African *S. haematobium* sample contained 4.1-22.0% *S. bovis* ancestry (median 7.0%). By comparison the median *S. bovis* ancestry was 0.02% and 100% in the southern *S. haematobium* and *S. bovis* control samples, respectively.

We used TWISST v67b9a66 (27) as an independent method for identifying local introgression.
TWISST, measures shifting gene tree frequencies across the genome. Trees were generated
from 37,200 non-overlapping, 10kb, sliding windows. On average (mean) each window contained
657.3 SNVs. We examined the three possible topologies between northern *S. haematobium*,
southern *S. haematobium* and *S. bovis* using *S. margrebowiei* (GCA_944470205.2; 28) as an
outgroup (Figure 2B). The expected species tree, with a monophyletic clade of *S. haematobium*,
sister to *S. bovis* was the most common with a mean weight of 0.876 across the genome. The
discordant topology uniting northern *S. haematobium* and *S. bovis* was the second most abundant
topology (weight = 0.085) compared to the topology with southern *S. haematobium* and *S. bovis*
(weight = 0.039).
181
We examined the genome for regions that are devoid of introgressed *S. bovis* in the northern *S.*
haematobium population. There were 918 genomic regions lacking *S. bovis* alleles with a median
size of 35.8Kb (Figure 2F). With log transformation and robust Z-scores we identified 19 genome
regions that were significant outliers in terms of length ranging from 1.13-6 Mb (median 1.67 Mb).
Thirteen of the 19 genome regions were on the ZW scaffold and accounted for 32% (28.6 Mb) of
its total length.
188

189   <u>Introgression profiles in different countries</u> - We examined the pattern of introgression in individual
190   countries of north Africa (Supplemental Figure 3) as determined by RFMix. The overall patterns
191   of introgression across the genome were consistent between north African populations. Pairwise
192   comparisons of introgressed allele frequencies between countries were positively correlated (*r* =
193   0.59-0.8; Supplemental Figure 4) despite distances spanning up to 3,000 km.
194

<u>Impact of introgression on nucleotide diversity and genetic differentiation of *S. haematobium*</u> - We
masked introgressed alleles within individual genomes, and recalculated π, F_{ST} and a PCA
(Figure 5). Prior to masking, mean nucleotide diversity (π), was 2.3-fold greater in the northern
(π = 2.991 x 10^{-3}) vs southern (π = 1.278 x 10^{-3}) *S. haematobium*, and π was 3.3-fold greater in
S. bovis (π = 8.329 x 10^{-3}) than the entirety of *S. haematobium* (π = 2.523 x 10^{-3}). After masking,
northern African *S. haematobium* nucleotide diversity was reduced to nearly identical levels seen
in the south population: π_{NW} = 2.991 x 10^{-3} to π_{NW} = 1.07 x 10^{-3}. By comparison, removing
introgressed alleles had no impact on F_{ST} (F_{ST} = 0.154) between northern and southern *S.*
haematobium. Additionally, the structure of the PCA was retained, demonstrating that the
differentiation between NW and SE Africa is not driven by introgression.

Both PCA (Fig 1 and 5) analyses and Admixture (k=3) plots (Supplemental Figure 1) recovered
one *S. bovis* and two *S. haematobium* population components. The distribution of these
components, shows a clear distinction between countries in the north and south, with a boundary
that extends from Cameroon to Somalia and into Madagascar (Figure 5A). This boundary
generally coincides with the distribution of *S. bovis* in north and central Africa (29) with a few
exceptions (30, 31). The division of the *S. haematobium* into northern and southern populations
was consistent among analyses with one exception. Madagascar was an intermediate population
in Admixture analysis (k=3). In the PCA, samples from Madagascar area assigned to the southern
cluster, but they form a distinct subgroup that is intermediate between the remaining southern
and northern samples.

<u>Dating Introgression</u> - We used the size of introgressed haplotype blocks to estimate the number
of generations since hybridization for each *S. haematobium* sample in north Africa (Supplemental
Figure 5). This gave estimated hybridization dates of ~257-879 generations ago (Median – 426
generations, 95% confidence intervals = 281.6-764 generations). *S. haematobium* generation
time varies from 3-4 (32) months in lab populations, but is estimated to be 6-12 months in wild
populations (33). These generation times imply that admixture between *S. haematobium* and *S.*
bovis occurred ~106 years ago assuming four generations per year (high transmission) or 426
years assuming one generation per year (low and / or seasonal transmission). Dating estimates
varied between countries: median estimates are lowest in Egypt (286.9) and highest in Nigeria
(565) despite their relatively close proximity. A one-way ANOVA indicated significant differences
in the number of generations since hybridization between countries (p-value = $1.3e^{-10}$;
Supplemental Figure 5).

<u>Selection and adaptive introgression</u> - We examined *S. haematobium* and *S. bovis* populations
for signatures of selection using normalized, xpEHH (Figure 2C). We found 996 statistically
significant xpEHH values after multiple test correction. We combined values within 1Mb to identify
15 genome regions with signatures of positive selection in the northern population and five in the
southern population (Supplemental Table 3). The median normalized xpEHH in each of these
regions was >|6| and the windows ranged in size from 3 bp to 709,928 bp (mean 139,942 bp).

We combined selection and introgression analyses to identify genome regions showing evidence
for adaptive introgression of *S. bovis* alleles into the northern *S. haematobium* population. These

regions contained outlier values for selection (xpEHH), elevated Patterson's D (D ≥ 0) indicative

of introgression, high levels of S. *bovis* ancestry (>95%) and significant differentiation from

southern *S. haematobium* (F_{ST} ≥ 95[TH] percentile). Two genome regions met these criteria;

chromosome four (NC_067199.1: 28,476,500-28,813,500) and chromosome five

(NC_067200.1:9,773,000-10,447,000). These genome regions span 1.01 Mb and 15 genes; eight

on chr four and seven on chr five (Table 1). Of the 74,955 SNVs in these regions, 989 are

nonsense or missense mutations. We found 37 missense SNVs where the *S. bovis* allele is at or

near fixation in the northern population (Supplemental Table 4). All of these variants are on chr

four and fall within four genes; leishmanolysin-like peptidase, a Rho GTPase-activating protein

35, Jumonji domain-containing protein six (JMJD6_4), and Jumonji domain-containing protein six

(JMJD6_3).

Discussion

Our analysis of >38 million SNVs provides compelling evidence that *S. haematobium* and *S. bovis*

are genetically well differentiated. This conclusion is supported by multiple lines of evidence: high

F_{ST} values (F_{ST} ≥ 0.74-0.79; Figure 1C; Figure 2D), distinction by PCA (Figure 1C), strong

differentiation by ancestry components in Admixture analyses (Figure 1D) and well supported

monophyletic clades in the nuclear species tree (Figure 3). The agreement between these

approaches suggests that strong barriers to gene flow exist between these two species.

Our analysis revealed that northern African *S. haematobium* are genetically differentiated (F_{ST} =

0.16) from the southern population. The boundary between these populations appears to extend

from Cameroon, Gabon, the Central African Republic, South Sudan, and Ethiopia (Figure 5A).

When introgressed *S. bovis* alleles were removed from genomic data, F_{ST} between these

populations remained unchanged (F_{ST} = 0.154). Hence, genetic differences between the northern

and southern populations do not appear to be driven by introgressed *S. bovis* alleles.

Our results indicate barriers to gene flow exist between northern and southern *S. haematobium*.

The southern *S. haematobium* clade is monophyletic within a larger clade of northern African *S.*

haematobium (Figure 3). This indicates that *S. haematobium* originated in one of the northern

African countries and is consistent with previous work that identified the Arabian Peninsula/Asia

as a potential ancestral source population (34). It is possible that the two populations are defined

by the distribution of their intermediate hosts. Regional differences in parasite compatibility with

their intermediate snail hosts can occur within limited geographical areas (35). *S. haematobium*

from North Africa and the Middle East are transmitted by the *Bulinus truncatus/tropicus* species
complex and parasites from the Afro-tropical region are transmitted by snails of the *Bulinus*
africanus group (36). The presence of such a barrier has important implications for our
understanding of the ecological and epidemiological factors that shape the distribution and
dynamics of these two parasite populations. Further investigation at the population boundaries
may provide new insights into biological differences and incompatibilities between northern and
southern *S. haematobium* populations.

Four aspects of our results support an ancient introgression hypothesis. First, each of the north
African *S. haematobium* samples contain low levels of *S. bovis* ancestry with the exception of the
sole Cameroonian sample. Percentages of *S. bovis* ancestry per individual are similar across
multiple analyses: introgressed haplotype blocks from RFMix account for 4.1-22% of individual
genomes in the northern *S. haematobium* population, while the population component associated
with *S. bovis* in Admixture ranges from 5-26.2% at *K*=2.

Second, the landscape of introgressed alleles across the genome is consistent across north
African samples (Supplemental Figure 3) and positively correlated (Supplemental Figure 4)
despite being separated by ≤3,000 Km. For this profile to be conserved across such a broad
distance suggests (A) it occurred in an ancestor of the north African *S. haematobium* or (B)
introgressed alleles provided a selective advantage that spread throughout the north African
population. Our data support the later with the nuclear phylogeny (Figure 3) showing that the
northern population is paraphyletic; the southern population is a monophyletic clade within this
group and it lacks introgressed *S. bovis* alleles. We also observe that some introgressed alleles
have reached high frequency in the north African population and show signs of selection (Figure
2). Finally, our data suggest that there is a barrier to gene flow/migration between northern and
southern *S. haematobium* populations, restricting dispersal of introgressed alleles to the southern
population.

Third, mitochondrial DNA provides insights into a minimal number of hybridization events. 58% of
northern *S. haematobium* samples contain introgressed S*. bovis* mitochondria (Figure 4). If the
introgressed *S. bovis* mitochondria were the result of contemporary hybridization, we would
expect sister relationships between *S. bovis* and *S. haematobium* at the terminal branches of the
tree. However, we find that introgressed *S. haematobium* individuals with introgressed *S. bovis*
mitochondrial genomes form two monophyletic clades. Clade "A" contains samples from Egypt,

Niger, and Cote d' Ivoire, and Clade "B" contains samples from Niger, Nigeria, and Cote d' Ivoire;

each clade spanning >1,000 Km. The most parsimonious interpretation of the phylogeny is that

the introgressed *S. bovis* mitochondria share two distinct origins and imply at least two admixture

events resulting from mating between a *S. bovis* female and *S. haematobium* male that occurred

in the distant past. We note that laboratory crosses between *S. haematobium* are often

asymmetric, and may only produce offspring when male *S. haematobium* are mated with female

S. bovis (37) or produce more male offspring (32). As females are the heterogametic sex, F1

females are expected to show reduced fitness (Haldane's rule; 38). This may contribute to the

limited number of *S. bovis* mtDNA lineages observed in *S. haematobium populations*.

Fourth, introgressed *S. bovis* nuclear loci are heavily fragmented within the *S. haematobium*

genomes indicating multiple generations since introgression. Our estimates of time since

introgression span 257-879 generations ago (95% confidence interval). Introgressed loci were

measured in tens of kilobases (median = 76.3 Kb) and the largest blocks extended into the

megabases (max = 4.05 Mb). This contrasts with early generation hybrids which would have

introgressed block lengths spanning, or nearly spanning entire chromosomes (39). One Nigerian

sample contained ~25% introgressed DNA, consistent with expectations for a F2 backcross.

However, the maximum introgressed fragment size in a Nigerian sample was only 2.83 Mb and

median block size in these samples ranged from 47.1-97.6 Kb indicating multiple recombination

events. We found that the time since introgression was significantly different between multiple

countries (Figure 6). For example, neighboring countries Niger (453 generations) and Nigeria

(565 generations) were not significantly different, but introgression in Cote d' Ivoire (385

generations) appears to have occurred more recently. The variation in the estimates of

generations since introgression are consistent with several regional introgression events.

Alternatively, variation in age estimates between countries could reflect extrinsic factors like

seasonality or intervention strategies that could lengthen or reduce generation times within sub-

populations. If this were the cause, it is possible that the number of generations that have lapsed

since an introgression event may vary between countries.

S. bovis shows 3.3-fold higher diversity than *S. haematobium*, while genetic diversity (π) is 2.3-

fold greater in the north *S. haematobium* population than in the south African *S. haematobium*

population. When the introgressed *S. bovis* alleles are removed from the analyses, this difference

in genetic diversity between the north and south *S. haematobium* populations is reduced to just

1.05-fold and π is not significantly different (Figure 5B). By contrast, F_{ST} values between northern

and southern *S. haematobium* are consistent whether introgressed alleles are considered (F_{ST} =
0.16) or not (F_{ST} = 0.154) and the relationship among samples in the PCAs is nearly identical
when introgressed alleles are included or excluded. These results indicate (i) that the elevated π
in northern African *S. haematobium* results from *S. bovis* introgression and (ii) that northern and
southern *S. haematobium* populations are well differentiated even after removing introgressed *S.*
bovis alleles.

Given that introgressed *S. bovis* alleles have persisted in the northern *S. haematobium*
population, we examined the data for signals of adaptive introgression. We found two introgressed
genome regions with signals of positive selection in the northern population on chr four
(NC_067199.1:28,348,440-28,877,530) and chr five (NC_067200.1:9,712,340-10,514,400).
Despite the convergence of signals to these regions, we were not able to identify variants driving
selection in these regions. We found 37 missense SNVs where the *S. bovis* allele was nearly
fixed in the northern population, but none withstood multiple test correction for directional
selection. These variants occur in four genes (WormBaseParaSite v18.0; 40), a Rho GTPase-
activating protein 35 (n_{SNVs} = 30; MS3_00007803) a Leishmanolysin-like peptidase (n_{SNVs} = 4;
MS3_00007802), and two members of the Jumonji domain-containing protein 6 family; JMJD6_4
(n_{SNVs} = 1; MS3_00010935) and JMJD6_3 (n_{SNVs} = 2; MS3_00010934). The same
Leishmanolysin-like peptidase (Table 1) has been identified as a candidate for adaptive
introgression from *S. bovis* into *S. haematobium* in two previous studies (14, 15). Genes in same
invadolysin gene family are known to modulate the snail host immune system in Schistosoma
mansoni (41, 42) and this particular gene has been associated with cell migration and invasion in
other parasitic taxa (43).

We also observed genomic regions on three chromosomes of the *S. haematobium* samples
where *S. bovis* introgression is rare or absent (Figure 2F). These introgression deserts may
contain hybrid incompatibility loci that result in reduced fitness of early generation hybrids and
present barriers to further introgression. Thirteen of the 19 regions occur on the sex
chromosomes. This is consistent with expectations given that a reduced recombination and
smaller effective populations size of the sex chromosomes expose deleterious, introgressed
alleles to strong selection pressures (44) and could lead to female sterility as predicted by
Haldane's rule.

Understanding hybridization and introgression between *S. haematobium* and *S. bovis* is important

for disease control. If hybridization between these species is infrequent, then there may be

minimal benefit in linking strategies that manage both human (*S. haematobium*) and livestock (*S.*

bovis) *Schistosoma* species. Consistent with this, our results from these samples suggest that

hybridization between these species is rare, and gene flow is insufficient to break down strong

reproductive barriers between these species. However, adaptive introgression has introduced *S.*

bovis alleles into *S. haematobium* populations. This is a clear example of alleles being transferred

between livestock and human parasites through introgression. Some *S. bovis* alleles have

reached high frequency and are likely selectively advantageous. Future work should aim to

understand how the introgressed *S. bovis* variants contribute to the fitness of *S. haematobium*

individuals. The strong differentiation between northern *S. haematobium* populations, carrying

introgressed *S. bovis* alleles and southern *S. haematobium* populations, with no introgression, is

of particular interest. Additionally, future work should examine differences between northern and

southern *S. haematobium* populations, and test whether they are reproductively isolated.

Several limitations to our study and its conclusions should be noted. First, our results indicate that

hybridization between *S. haematobium* and *S. bovis* is rare and ancient when measured on the

continental scale. While early generation hybrids between *S. haematobium* and *S. bovis* may be

found with further sampling, it is clear that strong barriers to gene flow maintain species integrity.

Second, recombination rates have not been quantified in *S. haematobium* so our estimates of

age of admixture are based on recombination rates measured in *Schistosoma mansoni* (45). To

improve the accuracy of these estimates, direct measures of recombination rates from *S.*

haematobium genetic crosses are needed. Third, our results pertain to *S. haematobium* and *S.*

bovis. Extant hybridization between other schistosome species (*S. haematobium*/*S. guineensis*

and *S. bovis*/*S. curassoni*) have been documented in field collected samples with genomic data

(46, 47). Our results suggest that *Schistosoma* species pairs may form a spectrum in hybridization

frequency and compatibility. Future work to understand the factors that impact hybridization and

present barriers to gene flow between schistosomes species pairs will be of great interest, and

can provide a more nuanced understanding of hybridization and potential implications for

schistosome control.

Online Methods

Sample collection: description, ethics, and identification – We used samples or data from three sources. i) The first dataset was generated from samples provided by the Schistosomiasis Collection At the Natural History Museum (48) which is housed at the Natural History Museum (London). SCAN samples consisted of individual miracidia and cercariae preserved on Whatman FTA cards (49). We analyzed 114 *S. haematobium* and *S. bovis* samples from 123 individual hosts (snails or humans) and 12 Africa countries. ii) In addition to the SCAN samples, we collected nine adult *Schistosome* worms, presumed to be *S. bovis*, from the intestines of routinely slaughtered cattle from meat vendors at three abattoirs located in Auchi, Benin City, and Enugu in Nigeria. In the laboratory, the mesenteric vessels of each purchased intestines were visually inspected for schistosome parasites. Adult schistosomes were recovered using forceps and washed in saline solution. Adult pairs were separated into males and females before being stored in 96% ethanol for subsequent DNA isolation analyses. iii) Finally, for the third source of data we used whole genome sequence data from NCBI (14, 15, 17, 21, 28, 50).

Samples provided by the SCAN repository were originally collected in accordance with protocols approved by local, state, and national authorities, including the Ministry of Health. The Imperial College Research Ethics Committee (ICREC) at Imperial College London, in conjunction with ongoing Schistosomiasis Control Initiative (SCI) activities, provided additional ethical guidance for samples collected through the CONTRAST program. Ethical clearance and study protocols for Nigerian samples were approved by the National Health Research Ethics Committee of Nigeria (NHREC) (protocol number: NHREC/01/01/2007– 30/10/2020 and approval number: NHREC/01/01/2007– 29/03/2021) and the Institutional Review Board (IRB) of University of Texas Health, San Antonio Texas, United States of America (protocol number: HSC20180612H). Informed consent was obtained from all participants, with processes tailored to ensure understanding and voluntary participation. All data were anonymized to protect participant privacy, and schistosomiasis-positive individuals were treated with a single dose of praziquantel (40 mg/kg). For livestock parasite collection, approval was secured from local veterinarians. No animals were euthanized for research purposes; *Schistosoma* samples were collected during routine activities at abattoirs. Further details on collection methods, ethical approvals, and data availability for public samples can be found in their respective publications documented in Supplemental Table 1.

Provisional species identifications were assigned to cercariae and miracidia based on sampled host. For example, miracidia hatched from eggs collected from human urine samples were

assumed to be *S. haematobium* while miracidia hatched from eggs in cattle feces were assumed
to be *S. bovis*. Cercariae collected from snails were identified by Sanger sequencing the
mitochondrial cox1 region and the ribosomal internal transcribed spacer (ITS) rDNA region as
previously described (20). Downstream genetic analysis with whole genome SNVs was used to
confirm and reassign species identifications where necessary.

Library prep and sequencing – DNA from single parasites stored on FTA cards was subjected to
whole-genome amplification (WGA) using methods previously described in (49). DNA was
extracted from single male adult *S. bovis* worms using the DNeasy® Blood and Tissue kit before
subsequent WGA. We quantified amount of schistosome DNA in each WGA sample by real time
quantitative PCR (qPCR) reactions using the single copy gene α-tubulin 1 gene markers primers
(*S. haematobium*: forward [GGT GGT ACT GGT TCT GGT TT], reverse [AAA GCA CAA TCC
GAA TGT TCT AA]; *S. bovis*: forward [ATG GCC TCG TTA TCA ACC AT], reverse [TGG CCT
CGT TAT CAA CCA TA] following previously described protocol in (49). DNA sequencing libraries
were generated from 500 ng of DNA per sample using the KAPA Hyperplus kit protocol with the
following modifications: i) enzymatic fragmentation at 37°C for 10 minutes, ii) adapter ligation at
20°C for an hour, and iii) 4 cycles of library PCR amplification. After qPCR quantification of each
library with KAPA Library Quantification Kits, samples with similar concentrations were combined
into pools for sequencing at 4nM, while samples with disparate concentrations were equalized in
10 mM Tris-HCl pH 8.5 before pooling. Libraries were sequenced with 150 bp paired-end reads
on two Illumina NovaSeq flowcell. All resulting reads were deposited in the NCBI Short Read
Archive under BioProject PRJNA636746 and are documented in Supplemental Table 1.

Computing environment - Analyses were conducted on the Texas Biomedical Research Institute's
high-performance computing cluster, with worker nodes containing 96 cores and 1 TB of memory.
Computational environments were managed using Conda v22.9.0. Environmental recipe files,
Jupyter notebooks, and other code can is archived on GitHub
(github.com/nealplatt/sch_hae_scan v0.1z) and at DOI:10.5281/zenodo.13124718.

Read filtering and Mapping - Raw reads were quality trimmed with trimmomatic v0.39 (51) using
the following parameters: LEADING:10, TRAILING:10, SLIDINGWINDOW:4:15, MINLEN:36,
ILLUMINACLIP:2:30:10:1:true. This command removed low quality bases at the beginning and
ends of the reads, removed portions of the read where quality dropped below a minimum
threshold, trimmed adapter sequences and discarded reads <36 nts. We then mapped the

trimmed reads to the *S. haematobium* reference genome, GCF_000699445.3 (21) with BBMap

v38.18 (52). On average the *S. haematobium* and *S. bovis* (GCA_944470425.1) genome

assemblies are ~97% similar across their genomes (17) which should minimally affect reference

biases when mapping short reads. However, to avoid reference biases we used the 'vslow' and

'minid=0.8' options with BBMap and discarded ambiguously mapping reads ('ambig=toss').

Genotyping, phasing, and filtering - Mapped reads were sorted with SAMtools v1.13 (53) and

checked for duplicates with GATK v4.2.0.0's (54) mark_duplicates. Then single nucleotide

variants (SNVs) were genotyped with HaplotypeCaller and GenotypeGVCFs. To make the

dataset more manageable, we genotyped each chromosome separately using the -L

option. Next, we removed all indels and hard filtered SNVs based on QualByDepth ("QD < 2.0"),

RMSMappingQuality (MQ < 30.0), FisherStrand (FS > 60.0), StrandOddsRation (SOR > 3.0),

MappingQualityRankSumTest (MQRankSum < -12.5), and ReadPosRankSumTest

(ReadPosRankSum < -8.0) with GATK's VariantFiltration. We removed multi-allelic sites, and

sites with genotype quality (GQ) <20 or read depth (DP) <8 with VCFtools v0.1.16 (55).   After

these filters were applied we removed genomic sites that were genotyped in ≤50% of individuals

and then any individuals that were genotyped at ≤50% of sites.

SNVs on each chromosome were phased using Beagle v 5.2_21Apr21.304 (56) in windows of

20cm and a 10cm overlap. Assuming a uniform recombination rate similar to *S. mansoni* across

the genome, these values are comparable to a 6.5 Mb window and a 3.25 Mb step size (45). We

used a burn in of 20 iterations and 60 iterations for the phasing run. All phased chromosome

VCFs were combined into a single file using vcfcombine from vcflib v1.0.3 (57) before an

additional round of post-phase filtering.

In some cases, multiple miracidia were analyzed from a single host potentially adding highly

related samples to our dataset and skewing the downstream results. To remove these, we

examined kinship coefficients in our samples using the autosomal chromosomes and the "–

unrelated" function in king v2.2.7 (58). This parameter identifies second-degree relatives within

the dataset that can be removed prior to downstream analyses. Next, we generated a set of SNVs

that were common (minor allele frequency; MAF > 0.05) and unlinked. Unlinked loci were filtered

with Plink v1.90b6.21 (59) by removing linked SNVs with a pairwise r^2 > 0.2. This filter was applied

in 25 Kb sliding windows with a 5kb steps. Finally, we used SnpEff v5.1 (60), to identify the impact

of these SNVs on the amino acid sequence in coding regions. To do this we imported the *S.*

haematobium reference genome (GCF_000699445.3) along with the associated GenBank

annotations to create a custom database.

<u>Principal Component Analyses</u> - We used a series of tools to explore population structure in our

data sets. We used common (minor allele frequency; MAF>0.05), unlinked, autosomal SNVs and

Plink v1.90b6.21 (59) to generate a principal component analysis (PCA) to examine relationships

among the samples. We used a *K*-means clustering algorithm to assign each sample to between

1 and 10 populations with the Kmeans() function in sklearn.cluster v1.2.0 (61). We then used the

Elbow method (62) to examine distortion in the model and determine the optimal number of

clusters in the data. Once we identified the optimal number of clusters, we assigned each sample

within a cluster, and those designations were used to differentiate the *S. haematobium*

populations using analyses as below. These cluster assignments were also used validate the

assumed species identify of each sample.

<u>Admixture</u> - We examined the ancestry of each sample with Admixture v1.3.0 (24) and the same

unlinked, autosomal SNV dataset from the PCA analyses. However, we further thinned the SNV

data with VCFtools v0.1.16 (55) ensuring that no two SNVs were within 10kb of each other. This

step minimizes any potential effects of linkage on the results. We ran Admixture v1.3.0 (24),

allowing for 2 to 20 possible population components, and used the cross-validation error to

determine the optimal range (63). Additionally, we randomly selected individuals with ≥99.999%

S. bovis or *S. haematobium* ancestry in the k=2 analysis to serve as reference samples for each

species in downstream analyses.

<u>Nucleotide diversity (π) and Fixation index (F_{ST})</u> - We used scikit-allel v1.3.5 (64) to calculate

nucleotide diversity (π) and the fixation index (F_{ST}) in sliding windows of 10 kb using autosomal,

common   (MAF   >0.05)   SNVs   allel.windowed_diversity()   and

allel.windowed_weir_cockerham_fst() functions. The weighted, Weir-Cockerham F_{ST} (22, 23)

was measured between species (*S. haematobium* vs. *S. bovis*) and between the *K*-means

populations. Next, we used the reference panel, described above, to identify ancestry informative

sites between the *S. haematobium* and *S. bovis* samples. We used scikit-allel v1.3.5's (64)

allel.weir_cockerham_fst() to calculate F_{ST} at individual sites. Only sites where $F_{ST} = 1$ where

retained.

<u>Biogeography</u> – *S. haematobium* was split into two groups based on the *K*-means clustering
analysis of the PCA results. At k=2 Admixture differentiated *S. haematobium* and *S. bovis*
samples, but at k=3 Admixture broadly confirmed the presence of two different *S. haematobium*
populations. We used the admixture proportion (Q) from k=3, to visualize how the populations
were distributed across Africa. The presence of this ancestry component was extrapolated into
unsampled geographic regions using the OrdinaryKriging() function implemented in pykrige
v1.7.0 with a linear variogram model (65). Geographic distances between samples were
calculated with the haversine() v2.8.0 function (https://pypi.org/project/haversine/).
550
<u>Genome-wide tests for introgression</u> - We used a series of tests to explore the presence of
introgression between *S. bovis* and the *S. haematobium* populations. First, we used
average_patterson_f3() from scikit-allel v1.3.5 (64) to calculate a normalized f_3 (66) averaged
across blocks of 500 SNVs. Next, we tested for gene flow using the *D*-statistic, also known as the
ABBA BABA test (67). We used *S. margrebowiei* (GCA_944470205.2; 28) as the outgroup (O),
S. bovis as the donor population (P3), and the *S. haematobium K*-means populations as the
recipients (P1 and P2). We measured *D* across the genome in 500 SNV blocks with
moving_patterson_d() in scikit-allel v1.3.5 (64). Introgressed loci were defined when $D>0+2\sigma$.
559
<u>Local Ancestry Assignment</u> - For local ancestry assignment, we used RFMix v2.03-r0 (26) and
TWISST v67b9a66 (27). RFMix v2.03-r0 (26) uses a random forest approach to assign local
ancestry to genomic segments by comparing samples to reference panels. For this, we used the
reference samples selected from the Admixture analyses. We generated a genetic map using a
uniform recombination rate estimated from *S. mansoni* crosses (1 centimorgan = 287,000 bp;
45). The remainder of the parameters were set to the default.
566
TWISST v67b9a66 (27) uses gene trees sampled from across the genome to identify potentially
introgressed loci. It does this by iteratively sampling subtrees from the gene tree and calculating
relative support for each of the possible species trees. We generated gene trees from loci
containing 500, phased, common (MAF >0.05) SNVs with RAxML-NG v1.1 (68). For each locus
we searched for the 10 best trees and then bootstrapped the best tree for 100 replicates using
the GTR+ASC_LEWIS substitution model and *S. margrebowiei* as an outgroup. Nodes supported
in ≤10 bootstrap replicates were collapsed with Newick Utilities v1.6 (69). The collapsed trees
were used as input for TWISST v67b9a66 (27). Samples were assigned to their *K*-means
population.

Selection - We compared selection in the *S. haematobium* intra populations using cross-population extended haplotype homozygosity (xpEHH; 70). Unphased xpEHH was measured with selscan v2.0.0 (71). The resulting unphased xpEHH values were normalized with norm v1.3.0 and the '--xpehh flag'. Bonferroni corrected p-values were assigned to each site. Sites with a corrected p-value < 0.01 were considered to be experiencing putative directional selection between the two *S. haematobium* populations.

Identifying putative adaptive introgression –We searched the genome for regions with F_{ST}, Patterson's D, local ancestry, and xpEHH values indicative of adaptive introgression. To do this we examined how these values were distributed across the genome in sliding windows of 337 Kb and 3,370 bp step size; values equivalent to 1% and 0.01% of the autosomal genome. Specifically, we were looking for regions of the genome that are among the most highly differentiated between the two schistosome populations (F_{ST} >= 95th percentile), with statistically significant signals of introgression (Patterson's D > 0) and directional selection (xpEHH p-value < 0.01), and the *S. bovis* alleles are at high frequency in the northern or southern *S. haematobium* populations (>95%). Windows that met these criteria were then merged together if they were within 10Kb of each other to identify loci containing signals of adaptive introgression.

Autosomal Species Tree - To better understand the relationships among the samples, used SVDquartets (72) as implemented in PAUP* v4.0.a.build166 (73) to generate a species tree. We examined 2.5m random quartets along with 100 standard bootstrap replicates. Nodes in the gene trees supported by <10% of bootstrap replicates were collapsed Newick Utilities v1.6 (69).

Dating introgression - Recombination acts to continuously break down introgressed haplotypes. As a result, the size of introgressed haplotype blocks is directly related to the number or generations since hybridization (74). This can be roughly estimated with the formula G=1/LP where G is generations, L is the average length of introgression haplotypes in Morgans, and P is the proportion of the genome from the major parent (75). We identified introgressed blocks and their lengths (L) for each individual with RFMix v2.03-r0 (26) and P was estimated using Admixture (represented as *q*). A one-way ANOVA was used to identify differences in age estimates between populations (countries).

<u>Introgression Deserts</u> – Some regions of the genome may be resistant to introgression. This could
present as large regions lacking introgressed alleles. We used the RFMix results to identify
regions of the genome where *S. bovis* ancestry was 0% in the north African *S. haematobium*
populations. We log-transformed the length of each region and assigned robust Z-scores. Putative
introgression deserts were regions with robust Z-scores > 3.

<u>Mitochondrial genome assembly and phylogeny</u> - We used GetOrganelle v1.7.7.0 (76) to *de novo*
assemble mitochondrial genomes. Specifically, we used the animal_mt model and 10 rounds of
assembly with *k*-mer sizes of 21, 45, 65, 85, and 105. The mitochondrial contigs were then
scaffolded with RagTag v2.1.0 (77, 78) and RAxML-NG v1.1 (68) was used to generate a
maximum likelihood tree of the mitochondrial genomes. We used a GTR+G substitution model to
and 100 starting trees. Nodal support was assessed with 1,000 bootstrap replicates.

Acknowledgements

We thank Sandy Smith and John Heaner for providing computational support at Texas Biomedical
Research Institute's High-Performance Computing Cluster. We would like to thank Frederic
Chevalier, Winka Le Clec'h, and Kathrin Bailey for providing feedback during the course of this
project. This research was funded by the National Institute of Allergy and Infectious Diseases
(NIAD R01 AI166049-01). Schistosomiasis Collections at the Natural History Museum (SCAN)
was supported by Wellcome Trust grant 1045958/Z/13/Z. The field work referenced in this study
relied on various organizations and funding bodies across multiple countries. In Angola, funding
was provided by the Calouste Gulbenkian Foundation in collaboration with the Centro de
Investigação em Saúde de Angol, the Programa Nacional de Doenças Tropicais Negligenciadas
and the Ministério da Saúde de Angola. In Côte d'Ivoire and Niger, the research was funded by
SCORE through the Bill & Melinda Gates Foundation, RISEAL-Niger, Université Félix Houphouët-
Boigny, and the Centre Suisse de Recherches Scientifiques en Côte d'Ivoire. The Eswatini
National Control Program, Environmental Health Unit, and the School Health Program contributed
significantly to the work in Eswatini. The CONTRAST project in Kenya, Uganda, and Zambia was
funded by the European Commission, the Kenya Medical Research Institute, National Museums
of Kenya, Uganda Ministry of Health, and the University of Zambia. In Liberia, the UK Department
for International Development provided funding via SCI (now Unlimit Health) and the Ministry of
Health of Liberia. Madagascar's efforts were funded through SCAN, Unlimit Health, and the
Madagascar Ministry of Public Health. Work in Namibia was funded by the END Fund and the
Namibia Ministry of Health and Social Services. In Zanzibar, the ZEST project was funded by

SCORE, the Ministry of Health (Zanzibar), and the Public Health Laboratory—Ivo de Carneri,

Pemba.

Individual collectors are listed in the Supplemental Table 1.

Author Contributions

Roy N Platt II: Conceptualization, Formal Analysis, Writing – Original Draft, Writing - Review &

Editing. Elisha Enabulele: Conceptualization, Formal Analysis, Investigation, Resources, Writing

– Original Draft, Writing - Review & Editing. Ehizogie Adeyemi: Resources. Marian O Agbugui:

Resources. Oluwaremilekun G Ajakaye: Resources. Ebube C Amaechi: Resources. Chika E

Ejikeugwu: Resources. Christopher Igbeneghu: Resources. Victor S Njom: Resources. Precious

Dlamini: Resources. Grace A. Arya: Investigation. Robbie Diaz: Investigation. Muriel Rabone:

Resources, Writing - Review & Editing. Fiona Allan: Resources, Writing - Review & Editing.

Bonnie Webster: Resources, Writing - Review & Editing. Aidan Emery: Resources, Writing -

Review & Editing. David Rollinson: Resources, Writing - Review & Editing. Timothy J.C.

Anderson: Conceptualization, Formal Analysis, Writing – Original Draft, Writing - Review &

Editing, Supervision.

Competing interests

The authors declare no competing interests.

Corresponding authors

Correspondence to Roy N. Platt II or Timothy J. C. Anderson.

References

1.　　Taylor SA, Larson EL. Insights from genomes into the evolutionary importance and prevalence of
hybridization in nature. Nature ecology & evolution. 2019;3(2):170-7. Epub 2019/01/31. doi:
10.1038/s41559-018-0777-y. PubMed PMID: 30697003.
2.　　Aguillon SM, Dodge TO, Preising GA, Schumer M. Introgression. Current biology : CB.
2022;32(16):R865-r8. Epub 2022/08/24. doi: 10.1016/j.cub.2022.07.004. PubMed PMID: 35998591;
PMCID: PMC10581619.
3.　　Mixão V, Gabaldón T. Hybridization and emergence of virulence in opportunistic human yeast
pathogens. Yeast (Chichester, England). 2018;35(1):5-20. Epub 2017/07/07. doi: 10.1002/yea.3242.
PubMed PMID: 28681409; PMCID: PMC5813172.
4.　　Depotter JR, Seidl MF, Wood TA, Thomma BP. Interspecific hybridization impacts host range and
pathogenicity of filamentous microbes. Current opinion in microbiology. 2016;32:7-13. Epub
2016/04/27. doi: 10.1016/j.mib.2016.04.005. PubMed PMID: 27116367.

5.   Ravazi A, Oliveira J, Madeira FF, Nunes GM, Reis YVD, Oliveira ABB, Azevedo LMS, Galvão C,
Azeredo-Oliveira MTV, Rosa JAD, Alevi KCC. Climate and Environmental Changes and Their Potential
Effects on the Dynamics of Chagas Disease: Hybridization in Rhodniini (Hemiptera, Triatominae). Insects.
2023;14(4). Epub 2023/04/27. doi: 10.3390/insects14040378. PubMed PMID: 37103193; PMCID:
PMC10143345.
6.   King KC, Stelkens RB, Webster JP, Smith DF, Brockhurst MA. Hybridization in Parasites:
Consequences for Adaptive Evolution, Pathogenesis, and Public Health in a Changing World. PLoS
pathogens. 2015;11(9):e1005098. Epub 2015/09/04. doi: 10.1371/journal.ppat.1005098. PubMed PMID:
26336070; PMCID: PMC4559376.
7.   Colley DG, Bustinduy AL, Secor WE, King CH. Human schistosomiasis. Lancet (London, England).
2014;383(9936):2253-64. Epub 2014/04/05. doi: 10.1016/s0140-6736(13)61949-2. PubMed PMID:
24698483; PMCID: PMC4672382.
8.   Webster BL, Diaw OT, Seye MM, Webster JP, Rollinson D. Introgressive hybridization of
Schistosoma haematobium group species in Senegal: species barrier break down between ruminant and
human schistosomes. PLoS Negl Trop Dis. 2013;7(4):e2110. Epub 2013/04/18. doi:
10.1371/journal.pntd.0002110. PubMed PMID: 23593513; PMCID: PMC3617179.
9.   Webster JP, Neves MI, Webster BL, Pennance T, Rabone M, Gouvras AN, Allan F, Walker M,
Rollinson D. Parasite Population Genetic Contributions to the Schistosomiasis Consortium for
Operational Research and Evaluation within Sub-Saharan Africa. The American journal of tropical
medicine and hygiene. 2020;103(1_Suppl):80-91. Epub 2020/05/14. doi: 10.4269/ajtmh.19-0827.
PubMed PMID: 32400355; PMCID: PMC7351308 role in the study design, data collection and analysis,
decision to publish, or preparation of the manuscript.
10.   Léger E, Borlase A, Fall CB, Diouf ND, Diop SD, Yasenev L, Catalano S, Thiam CT, Ndiaye A, Emery
A, Morrell A, Rabone M, Ndao M, Faye B, Rollinson D, Rudge JW, Sène M, Webster JP. Prevalence and
distribution of schistosomiasis in human, livestock, and snail populations in northern Senegal: a One
Health epidemiological study of a multi-host system. The Lancet Planetary health. 2020;4(8):e330-e42.
Epub 2020/08/18. doi: 10.1016/s2542-5196(20)30129-7. PubMed PMID: 32800151; PMCID:
PMC7443702.
11.   Huyse T, Webster BL, Geldof S, Stothard JR, Diaw OT, Polman K, Rollinson D. Bidirectional
introgressive hybridization between a cattle and human schistosome species. PLoS pathogens.
2009;5(9):e1000571. Epub 2009/09/05. doi: 10.1371/journal.ppat.1000571. PubMed PMID: 19730700;
PMCID: PMC2731855.
12.   Sene-Wade M, Marchand B, Rollinson D, Webster BL. Urogenital schistosomiasis and
hybridization between Schistosoma haematobium and Schistosoma bovis in adults living in Richard-Toll,
Senegal. Parasitology. 2018;145(13):1723-6. Epub 2018/09/07. doi: 10.1017/s0031182018001415.
PubMed PMID: 30185248.
13.   Díaz AV, Walker M, Webster JP. Reaching the World Health Organization elimination targets for
schistosomiasis: the importance of a One Health perspective. Philosophical transactions of the Royal
Society of London Series B, Biological sciences. 2023;378(1887):20220274. Epub 2023/08/21. doi:
10.1098/rstb.2022.0274. PubMed PMID: 37598697; PMCID: PMC10440173.
14.   Platt RN, McDew-White M, Le Clec'h W, Chevalier FD, Allan F, Emery AM, Garba A, Hamidou AA,
Ame SM, Webster JP, Rollinson D, Webster BL, Anderson TJC. Ancient Hybridization and Adaptive
Introgression of an Invadolysin Gene in Schistosome Parasites. Molecular biology and evolution.
2019;36(10):2127-42. Epub 2019/06/30. doi: 10.1093/molbev/msz154. PubMed PMID: 31251352;
PMCID: PMC6759076.
15.   Rey O, Toulza E, Chaparro C, Allienne JF, Kincaid-Smith J, Mathieu-Begné E, Allan F, Rollinson D,
Webster BL, Boissier J. Diverging patterns of introgression from Schistosoma bovis across S.

haematobium African lineages. PLoS pathogens. 2021;17(2):e1009313. Epub 2021/02/06. doi:
10.1371/journal.ppat.1009313. PubMed PMID: 33544762; PMCID: PMC7891765.
16.   Djuikwo-Teukeng FF, Kouam Simo A, Allienne JF, Rey O, Njayou Ngapagna A, Tchuem-Tchuente
LA, Boissier J. Population genetic structure of Schistosoma bovis in Cameroon. Parasites & vectors.
2019;12(1):56. Epub 2019/01/27. doi: 10.1186/s13071-019-3307-0. PubMed PMID: 30678712; PMCID:
PMC6346511.
17.   Oey H, Zakrzewski M, Gravermann K, Young ND, Korhonen PK, Gobert GN, Nawaratna S, Hasan
S, Martínez DM, You H, Lavin M, Jones MK, Ragan MA, Stoye J, Oleaga A, Emery AM, Webster BL,
Rollinson D, Gasser RB, McManus DP, Krause L. Whole-genome sequence of the bovine blood fluke
Schistosoma bovis supports interspecific hybridization with S. haematobium. PLoS pathogens.
2019;15(1):e1007513. Epub 2019/01/24. doi: 10.1371/journal.ppat.1007513. PubMed PMID: 30673782;
PMCID: PMC6361461.
18.   Boon NAM, Mbow M, Paredis L, Moris P, Sy I, Maes T, Webster BL, Sacko M, Volckaert FAM,
Polman K, Huyse T. No barrier breakdown between human and cattle schistosome species in the
Senegal River Basin in the face of hybridisation. International journal for parasitology. 2019;49(13-
14):1039-48. Epub 2019/11/18. doi: 10.1016/j.ijpara.2019.08.004. PubMed PMID: 31734338.
19.   Kincaid-Smith J, Tracey A, de Carvalho Augusto R, Bulla I, Holroyd N, Rognon A, Rey O, Chaparro
C, Oleaga A, Mas-Coma S, Allienne JF, Grunau C, Berriman M, Boissier J, Toulza E. Morphological and
genomic characterisation of the Schistosoma hybrid infecting humans in Europe reveals admixture
between Schistosoma haematobium and Schistosoma bovis. PLoS Negl Trop Dis. 2021;15(12):e0010062.
Epub 2021/12/24. doi: 10.1371/journal.pntd.0010062. PubMed PMID: 34941866; PMCID: PMC8741037.
20.   Pennance T, Allan F, Emery A, Rabone M, Cable J, Garba AD, Hamidou AA, Webster JP, Rollinson
D, Webster BL. Interactions between Schistosoma haematobium group species and their Bulinus spp.
intermediate hosts along the Niger River Valley. Parasites & vectors. 2020;13(1):268. Epub 2020/05/26.
doi: 10.1186/s13071-020-04136-9. PubMed PMID: 32448268; PMCID: PMC7247258.
21.   Stroehlein AJ, Korhonen PK, Lee VV, Ralph SA, Mentink-Kane M, You H, McManus DP, Tchuenté
LT, Stothard JR, Kaur P, Dudchenko O, Aiden EL, Yang B, Yang H, Emery AM, Webster BL, Brindley PJ,
Rollinson D, Chang BCH, Gasser RB, Young ND. Chromosome-level genome of Schistosoma
haematobium underpins genome-wide explorations of molecular variation. PLoS pathogens.
2022;18(2):e1010288. Epub 2022/02/16. doi: 10.1371/journal.ppat.1010288. PubMed PMID: 35167626;
PMCID: PMC8846543.
22.   Weir BS, Cockerham CC. Estimating F-statistics for the analysis of population structure.
Evolution; international journal of organic evolution. 1984;38(6):1358-70. Epub 1984/11/01. doi:
10.1111/j.1558-5646.1984.tb05657.x. PubMed PMID: 28563791.
23.   Bhatia G, Patterson N, Sankararaman S, Price AL. Estimating and interpreting FST: the impact of
rare variants. Genome Res. 2013;23(9):1514-21. Epub 2013/07/19. doi: 10.1101/gr.154831.113.
PubMed PMID: 23861382; PMCID: PMC3759727.
24.   Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated
individuals. Genome Res. 2009;19(9):1655-64. Epub 2009/08/04. doi: 10.1101/gr.094052.109. PubMed
PMID: 19648217; PMCID: PMC2752134.
25.   Angora EK, Allienne JF, Rey O, Menan H, Touré AO, Coulibaly JT, Raso G, Yavo W, N'Goran EK,
Utzinger J, Balmer O, Boissier J. High prevalence of Schistosoma haematobium × Schistosoma bovis
hybrids in schoolchildren in Côte d'Ivoire. Parasitology. 2020;147(3):287-94. Epub 2019/11/16. doi:
10.1017/s0031182019001549. PubMed PMID: 31727202; PMCID: PMC10317610.
26.   Maples BK, Gravel S, Kenny EE, Bustamante CD. RFMix: a discriminative modeling approach for
rapid and robust local-ancestry inference. American journal of human genetics. 2013;93(2):278-88. Epub
2013/08/06. doi: 10.1016/j.ajhg.2013.06.020. PubMed PMID: 23910464; PMCID: PMC3738819.

27.   Martin SH, Van Belleghem SM. Exploring Evolutionary Relationships Across the Genome Using
Topology Weighting. Genetics. 2017;206(1):429-38. Epub 2017/03/28. doi:
10.1534/genetics.116.194720. PubMed PMID: 28341652; PMCID: PMC5419486.
28.   Comparative genomics of the major parasitic worms. Nature genetics. 2019;51(1):163-74. Epub
2018/11/07. doi: 10.1038/s41588-018-0262-1. PubMed PMID: 30397333; PMCID: PMC6349046.
29.   Moné H, Mouahid G, Morand S. The distribution of Schistosoma bovis Sonsino, 1876 in relation
to intermediate host mollusc-parasite relationships. Advances in parasitology. 1999;44:99-138. Epub
1999/11/24. doi: 10.1016/s0065-308x(08)60231-6. PubMed PMID: 10563396.
30.   Pennance T, Ame SM, Amour AK, Suleiman KR, Allan F, Rollinson D, Webster BL. Occurrence of
Schistosoma bovis on Pemba Island, Zanzibar: implications for urogenital schistosomiasis transmission
monitoring. Parasitology. 2018;145(13):1727-31. Epub 2018/08/09. doi: 10.1017/s0031182018001154.
PubMed PMID: 30086805; PMCID: PMC7116046.
31.   Stothard JR, Lockyer AE, Kabatereine NB, Tukahebwa EM, Kazibwe F, Rollinson D, Fenwick A.
Schistosoma bovis in western Uganda. Journal of helminthology. 2004;78(3):281-4. Epub 2004/10/08.
doi: 10.1079/joh2004239. PubMed PMID: 15469635.
32.   Polack B, Mathieu-Bégné E, Vallée I, Rognon A, Fontaine JJ, Toulza E, Thomas M, Boissier J.
Experimental Infections Reveal Acquired Zoonotic Capacity of Human Schistosomiasis Trough
Hybridization. The Journal of infectious diseases. 2024;229(6):1904-8. Epub 2024/04/26. doi:
10.1093/infdis/jiae152. PubMed PMID: 38669235.
33.   King CH, Muchiri EM, Ouma JH. Evidence against rapid emergence of praziquantel resistance in
Schistosoma haematobium, Kenya. Emerging infectious diseases. 2000;6(6):585-94. Epub 2000/11/15.
doi: 10.3201/eid0606.000606. PubMed PMID: 11076716; PMCID: PMC2640915.
34.   Webster BL, Emery AM, Webster JP, Gouvras A, Garba A, Diaw O, Seye MM, Tchuente LA,
Simoonga C, Mwanga J, Lange C, Kariuki C, Mohammed KA, Stothard JR, Rollinson D. Genetic diversity
within Schistosoma haematobium: DNA barcoding reveals two distinct groups. PLoS Negl Trop Dis.
2012;6(10):e1882. Epub 2012/11/13. doi: 10.1371/journal.pntd.0001882. PubMed PMID: 23145200;
PMCID: PMC3493392.
35.   Rollinson D, Stothard JR, Southgate VR. Interactions between intermediate snail hosts of the
genus Bulinus and schistosomes of the Schistosoma haematobium group. Parasitology. 2001;123
Suppl:S245-60. Epub 2002/01/05. doi: 10.1017/s0031182001008046. PubMed PMID: 11769287.
36.   Brown DS. Freshwater snails of Africa and their medical importance: CRC press; 1994.
37.   Taylor MG. Hybridisation experiments on five species of African schistosomes. Journal of
helminthology. 1970;44(3):253-314. Epub 1970/01/01. doi: 10.1017/s0022149x00021969. PubMed
PMID: 5505355.
38.   Haldane JBJJog. Sex ratio and unisexual sterility in hybrid animals1922;12:101-9.
39.   Harris K, Nielsen R. The Genetic Cost of Neanderthal Introgression. Genetics. 2016;203(2):881-
91. Epub 2016/04/03. doi: 10.1534/genetics.116.186890. PubMed PMID: 27038113; PMCID:
PMC4896200.
40.   Harris TW, Arnaboldi V, Cain S, Chan J, Chen WJ, Cho J, Davis P, Gao S, Grove CA, Kishore R, Lee
RYN, Muller HM, Nakamura C, Nuin P, Paulini M, Raciti D, Rodgers FH, Russell M, Schindelman G, Auken
KV, Wang Q, Williams G, Wright AJ, Yook K, Howe KL, Schedl T, Stein L, Sternberg PW. WormBase: a
modern Model Organism Information Resource. Nucleic Acids Res. 2020;48(D1):D762-d7. Epub
2019/10/24. doi: 10.1093/nar/gkz920. PubMed PMID: 31642470; PMCID: PMC7145598.
41.   Hambrook JR, Kaboré AL, Pila EA, Hanington PC. A metalloprotease produced by larval
Schistosoma mansoni facilitates infection establishment and maintenance in the snail host by interfering
with immune cell function. PLoS pathogens. 2018;14(10):e1007393. Epub 2018/10/30. doi:
10.1371/journal.ppat.1007393. PubMed PMID: 30372490; PMCID: PMC6224180.

42.   Hambrook JR, Hanington PC. A cercarial invadolysin interferes with the host immune response
and facilitates infection establishment of Schistosoma mansoni. PLoS pathogens. 2023;19(2):e1010884.
Epub 2023/02/03. doi: 10.1371/journal.ppat.1010884. PubMed PMID: 36730464; PMCID: PMC9928134.
43.   McHugh B, Krause SA, Yu B, Deans AM, Heasman S, McLaughlin P, Heck MM. Invadolysin: a
novel, conserved metalloprotease links mitotic structural rearrangements with cell migration. The
Journal of cell biology. 2004;167(4):673-86. Epub 2004/11/24. doi: 10.1083/jcb.200405155. PubMed
PMID: 15557119; PMCID: PMC2172566.
44.   Fraïsse C, Sachdeva H. The rates of introgression and barriers to genetic exchange between
hybridizing species: sex chromosomes vs autosomes. Genetics. 2021;217(2). Epub 2021/03/17. doi:
10.1093/genetics/iyaa025. PubMed PMID: 33724409; PMCID: PMC8045713.
45.   Criscione CD, Valentim CL, Hirai H, LoVerde PT, Anderson TJ. Genomic linkage map of the human
blood fluke Schistosoma mansoni. Genome Biol. 2009;10(6):R71. Epub 2009/07/02. doi: 10.1186/gb-
2009-10-6-r71. PubMed PMID: 19566921; PMCID: PMC2718505.
46.   Berger DJ, Léger E, Sankaranarayanan G, Sène M, Diouf ND, Rabone M, Emery A, Allan F, Cotton
JA, Berriman M, Webster JP. Genomic evidence of contemporary hybridization between Schistosoma
species. PLoS pathogens. 2022;18(8):e1010706. Epub 2022/08/09. doi: 10.1371/journal.ppat.1010706.
PubMed PMID: 35939508; PMCID: PMC9387932.
47.   Landeryou T, Rabone M, Allan F, Maddren R, Rollinson D, Webster BL, Tchuem-Tchuenté LA,
Anderson RM, Emery AM. Genome-wide insights into adaptive hybridisation across the Schistosoma
haematobium group in West and Central Africa. PLoS Negl Trop Dis. 2022;16(1):e0010088. Epub
2022/02/01. doi: 10.1371/journal.pntd.0010088. PubMed PMID: 35100291; PMCID: PMC8803156.
48.   Emery AM, Allan FE, Rabone ME, Rollinson D. Schistosomiasis collection at NHM (SCAN).
Parasites & vectors. 2012;5:185. Epub 2012/09/05. doi: 10.1186/1756-3305-5-185. PubMed PMID:
22943137; PMCID: PMC3453491.
49.   Le Clec'h W, Chevalier FD, McDew-White M, Allan F, Webster BL, Gouvras AN, Kinunghi S,
Tchuem Tchuenté LA, Garba A, Mohammed KA, Ame SM, Webster JP, Rollinson D, Emery AM, Anderson
TJC. Whole genome amplification and exome sequencing of archived schistosome miracidia.
Parasitology. 2018;145(13):1739-47. Epub 2018/05/29. doi: 10.1017/s0031182018000811. PubMed
PMID: 29806576; PMCID: PMC6193844.
50.   Young ND, Jex AR, Li B, Liu S, Yang L, Xiong Z, Li Y, Cantacessi C, Hall RS, Xu X, Chen F, Wu X,
Zerlotini A, Oliveira G, Hofmann A, Zhang G, Fang X, Kang Y, Campbell BE, Loukas A, Ranganathan S,
Rollinson D, Rinaldi G, Brindley PJ, Yang H, Wang J, Wang J, Gasser RB. Whole-genome sequence of
Schistosoma haematobium. Nature genetics. 2012;44(2):221-5. Epub 2012/01/17. doi: 10.1038/ng.1065.
PubMed PMID: 22246508.
51.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114-20. Epub 2014/04/04. doi: 10.1093/bioinformatics/btu170. PubMed
PMID: 24695404; PMCID: PMC4103590.
52.   Bushnell B. BBMap: a fast, accurate, splice-aware aligner. Lawrence Berkeley National
Lab.(LBNL), Berkeley, CA (United States), 2014.
53.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. The
Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. Epub 2009/06/10.
doi: 10.1093/bioinformatics/btp352. PubMed PMID: 19505943; PMCID: PMC2723002.
54.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D,
Gabriel S, Daly M, DePristo MA. The Genome Analysis Toolkit: a MapReduce framework for analyzing
next-generation DNA sequencing data. Genome Res. 2010;20(9):1297-303. Epub 2010/07/21. doi:
10.1101/gr.107524.110. PubMed PMID: 20644199; PMCID: PMC2928508.
55.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,
Marth GT, Sherry ST, McVean G, Durbin R. The variant call format and VCFtools. Bioinformatics.

2011;27(15):2156-8. Epub 2011/06/10. doi: 10.1093/bioinformatics/btr330. PubMed PMID: 21653522;
PMCID: PMC3137218.

56.   Browning BL, Tian X, Zhou Y, Browning SR. Fast two-stage phasing of large-scale sequence data.
American journal of human genetics. 2021;108(10):1880-90. Epub 2021/09/04. doi:
10.1016/j.ajhg.2021.08.005. PubMed PMID: 34478634; PMCID: PMC8551421.

57.   Garrison E, Kronenberg ZN, Dawson ET, Pedersen BS, Prins P. A spectrum of free software tools
for processing the VCF variant call format: vcflib, bio-vcf, cyvcf2, hts-nim and slivar. PLoS Comput Biol.
2022;18(5):e1009123. Epub 2022/06/01. doi: 10.1371/journal.pcbi.1009123. PubMed PMID: 35639788;
PMCID: PMC9286226.

58.   Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen W-MJB. Robust relationship
inference in genome-wide association studies2010;26(22):2867-73.

59.   Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker
PI, Daly MJ, Sham PC. PLINK: a tool set for whole-genome association and population-based linkage
analyses. American journal of human genetics. 2007;81(3):559-75. Epub 2007/08/19. doi:
10.1086/519795. PubMed PMID: 17701901; PMCID: PMC1950838.

60.   Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. A program
for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the
genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly. 2012;6(2):80-92. Epub 2012/06/26.
doi: 10.4161/fly.19695. PubMed PMID: 22728672; PMCID: PMC3679285.

61.   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer P,
Weiss R, Dubourg VJtJomLr. Scikit-learn: Machine learning in Python2011;12:2825-30.

62.   Kodinariya TM, Makwana PRJIJ. Review on determining number of Cluster in K-Means
Clustering2013;1(6):90-5.

63.   Evanno G, Regnaut S, Goudet JJMe. Detecting the number of clusters of individuals using the
software STRUCTURE: a simulation study2005;14(8):2611-20.

64.   Miles A, R M, Ralph P, Kelleher J, Pisupati R, Rae S, Millar T. scikit-allel: v1.3.5. 2022.

65.   Murphy BS. PyKrige: Development of a Kriging Toolkit for Python. 2014.

66.   Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, Genschoreck T, Webster T, Reich
D. Ancient admixture in human history. Genetics. 2012;192(3):1065-93. Epub 2012/09/11. doi:
10.1534/genetics.112.145037. PubMed PMID: 22960212; PMCID: PMC3522152.

67.   Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai W, Fritz
MH, Hansen NF, Durand EY, Malaspinas AS, Jensen JD, Marques-Bonet T, Alkan C, Prüfer K, Meyer M,
Burbano HA, Good JM, Schultz R, Aximu-Petri A, Butthof A, Höber B, Höffner B, Siegemund M,
Weihmann A, Nusbaum C, Lander ES, Russ C, Novod N, Affourtit J, Egholm M, Verna C, Rudan P,
Brajkovic D, Kucan Ž, Gušic I, Doronichev VB, Golovanova LV, Lalueza-Fox C, de la Rasilla M, Fortea J,
Rosas A, Schmitz RW, Johnson PLF, Eichler EE, Falush D, Birney E, Mullikin JC, Slatkin M, Nielsen R, Kelso
J, Lachmann M, Reich D, Pääbo S. A draft sequence of the Neandertal genome. Science (New York, NY).
2010;328(5979):710-22. Epub 2010/05/08. doi: 10.1126/science.1188021. PubMed PMID: 20448178;
PMCID: PMC5100745.

68.   Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: a fast, scalable and user-
friendly tool for maximum likelihood phylogenetic inference. Bioinformatics. 2019;35(21):4453-5. Epub
2019/05/10. doi: 10.1093/bioinformatics/btz305. PubMed PMID: 31070718; PMCID: PMC6821337.

69.   Junier T, Zdobnov EM. The Newick utilities: high-throughput phylogenetic tree processing in the
UNIX shell. Bioinformatics. 2010;26(13):1669-70. Epub 2010/05/18. doi: 10.1093/bioinformatics/btq243.
PubMed PMID: 20472542; PMCID: PMC2887050.

70.   Sabeti PC, Varilly P, Fry B, Lohmueller J, Hostetter E, Cotsapas C, Xie X, Byrne EH, McCarroll SA,
Gaudet R, Schaffner SF, Lander ES, Frazer KA, Ballinger DG, Cox DR, Hinds DA, Stuve LL, Gibbs RA,
Belmont JW, Boudreau A, Hardenbol P, Leal SM, Pasternak S, Wheeler DA, Willis TD, Yu F, Yang H, Zeng

C, Gao Y, Hu H, Hu W, Li C, Lin W, Liu S, Pan H, Tang X, Wang J, Wang W, Yu J, Zhang B, Zhang Q, Zhao H,
Zhao H, Zhou J, Gabriel SB, Barry R, Blumenstiel B, Camargo A, Defelice M, Faggart M, Goyette M, Gupta
S, Moore J, Nguyen H, Onofrio RC, Parkin M, Roy J, Stahl E, Winchester E, Ziaugra L, Altshuler D, Shen Y,
Yao Z, Huang W, Chu X, He Y, Jin L, Liu Y, Shen Y, Sun W, Wang H, Wang Y, Wang Y, Xiong X, Xu L, Waye
MM, Tsui SK, Xue H, Wong JT, Galver LM, Fan JB, Gunderson K, Murray SS, Oliphant AR, Chee MS,
Montpetit A, Chagnon F, Ferretti V, Leboeuf M, Olivier JF, Phillips MS, Roumy S, Sallée C, Verner A,
Hudson TJ, Kwok PY, Cai D, Koboldt DC, Miller RD, Pawlikowska L, Taillon-Miller P, Xiao M, Tsui LC, Mak
W, Song YQ, Tam PK, Nakamura Y, Kawaguchi T, Kitamoto T, Morizono T, Nagashima A, Ohnishi Y, Sekine
A, Tanaka T, Tsunoda T, Deloukas P, Bird CP, Delgado M, Dermitzakis ET, Gwilliam R, Hunt S, Morrison J,
Powell D, Stranger BE, Whittaker P, Bentley DR, Daly MJ, de Bakker PI, Barrett J, Chretien YR, Maller J,
McCarroll S, Patterson N, Pe'er I, Price A, Purcell S, Richter DJ, Sabeti P, Saxena R, Schaffner SF, Sham PC,
Varilly P, Altshuler D, Stein LD, Krishnan L, Smith AV, Tello-Ruiz MK, Thorisson GA, Chakravarti A, Chen
PE, Cutler DJ, Kashuk CS, Lin S, Abecasis GR, Guan W, Li Y, Munro HM, Qin ZS, Thomas DJ, McVean G,
Auton A, Bottolo L, Cardin N, Eyheramendy S, Freeman C, Marchini J, Myers S, Spencer C, Stephens M,
Donnelly P, Cardon LR, Clarke G, Evans DM, Morris AP, Weir BS, Tsunoda T, Johnson TA, Mullikin JC,
Sherry ST, Feolo M, Skol A, Zhang H, Zeng C, Zhao H, Matsuda I, Fukushima Y, Macer DR, Suda E, Rotimi
CN, Adebamowo CA, Ajayi I, Aniagwu T, Marshall PA, Nkwodimmah C, Royal CD, Leppert MF, Dixon M,
Peiffer A, Qiu R, Kent A, Kato K, Niikawa N, Adewole IF, Knoppers BM, Foster MW, Clayton EW, Watkin J,
Gibbs RA, Belmont JW, Muzny D, Nazareth L, Sodergren E, Weinstock GM, Wheeler DA, Yakub I, Gabriel
SB, Onofrio RC, Richter DJ, Ziaugra L, Birren BW, Daly MJ, Altshuler D, Wilson RK, Fulton LL, Rogers J,
Burton J, Carter NP, Clee CM, Griffiths M, Jones MC, McLay K, Plumb RW, Ross MT, Sims SK, Willey DL,
Chen Z, Han H, Kang L, Godbout M, Wallenburg JC, L'Archevêque P, Bellemare G, Saeki K, Wang H, An D,
Fu H, Li Q, Wang Z, Wang R, Holden AL, Brooks LD, McEwen JE, Guyer MS, Wang VO, Peterson JL, Shi M,
Spiegel J, Sung LM, Zacharia LF, Collins FS, Kennedy K, Jamieson R, Stewart J. Genome-wide detection
and characterization of positive selection in human populations. Nature. 2007;449(7164):913-8. Epub
2007/10/19. doi: 10.1038/nature06250. PubMed PMID: 17943131; PMCID: PMC2687721.
71.   Szpiech Z. selscan 2.0: scanning for sweeps in unphased data. bioRxiv; 2021.
72.   Chifman J, Kubatko L. Identifiability of the unrooted species tree topology under the coalescent
model with time-reversible substitution processes, site-specific rate variation, and invariable sites.
Journal of theoretical biology. 2015;374:35-47. Epub 2015/03/21. doi: 10.1016/j.jtbi.2015.03.006.
PubMed PMID: 25791286.
73.   Swofford DLJhpcfe. PAUP: phylogenetic analysis using parsimony (and other methods), version
4.0 beta2002.
74.   Harrison RG, Larson EL. Hybridization, introgression, and the nature of species boundaries. The
Journal of heredity. 2014;105 Suppl 1:795-809. Epub 2014/08/26. doi: 10.1093/jhered/esu033. PubMed
PMID: 25149255.
75.   Schumer M, Cui R, Powell DL, Rosenthal GG, Andolfatto P. Ancient hybridization and genomic
stabilization in a swordtail fish. Molecular ecology. 2016;25(11):2661-79. Epub 2016/03/05. doi:
10.1111/mec.13602. PubMed PMID: 26937625.
76.   Jin JJ, Yu WB, Yang JB, Song Y, dePamphilis CW, Yi TS, Li DZ. GetOrganelle: a fast and versatile
toolkit for accurate de novo assembly of organelle genomes. Genome Biol. 2020;21(1):241. Epub
2020/09/12. doi: 10.1186/s13059-020-02154-5. PubMed PMID: 32912315; PMCID: PMC7488116.
77.   Alonge M, Lebeigle L, Kirsche M, Jenike K, Ou S, Aganezov S, Wang X, Lippman ZB, Schatz MC,
Soyk S. Automated assembly scaffolding using RagTag elevates a new tomato system for high-
throughput genome editing. Genome Biol. 2022;23(1):258. Epub 2022/12/16. doi: 10.1186/s13059-022-
02823-7. PubMed PMID: 36522651; PMCID: PMC9753292.

78.   Alonge M, Soyk S, Ramakrishnan S, Wang X, Goodwin S, Sedlazeck FJ, Lippman ZB, Schatz MC.
RaGOO: fast and accurate reference-guided scaffolding of draft genomes. Genome Biol. 2019;20(1):224-.
doi: 10.1186/s13059-019-1829-6. PubMed PMID: 31661016.

967
968
Figure 1. – Sampling localities, sample summary and the population structure of *Schistosoma*
haematobium* and *S. bovis. (A) Collection locations for samples used in this study. Where exact
coordinates for samples were not readily available we used the country capital as the collection locality.
Two populations of *S. haematobium* were identified; northern and southern. The southern population in
red-yellow and the northern population in blue to green. (B) A principal component analysis of 355,715
unlinked, common (MAF>0.05), autosomal variants. The three clusters correspond to *S. bovis,* and the
northern and southern *S. haematobium* populations. Weighted, Weir-Cockerham F_{ST} values between
these populations are shown.
977

978
Figure 2. – Local measurements of differentiation, introgression and selection across the genome.
(A) The weighted Weir-Cockerham fixation index (FST) between northern and southern Africa, S.
haematobium populations was measured across the genome in 10Kb windows. These results indicated
multiple, highly differentiated regions between the two populations. (B) Patterson's *D* statistic was

measured to determine if high FST regions were the result of *S. bovis* alleles present in northern
populations. *D* measured across the genome was significantly positive indicating the presence of gene
flow between S. bovis and north African *S. haematobium* populations. (C) The frequency of *S. bovis*
ancestry across the genome in the northern *S. haematobium* population was estimated using RFmix. While
the percentage of *S. bovis* alleles in the population are low overall, the *S. bovis* alleles are at or near fixation
at loci on Chr4 and Chr5. (D) Gene tree topology weightings across the genome depicting the possible
relationships between the northern and south *S. haematobium* populations and *S. bovis* using TWISST.
Each locus across the genome is shown as stacked bar plots. While both tools use different methods to
depict the relationships between these taxa they recover similar results. (E) Differential selection between
S. haematobium populations was measured across the genome with extended haplotype homozygosity
(xpEHH). Positive values indicate positive selection in the northern population and negative values indicate
positive selection in the southern population. Significant xpEHH values (p<0.05) after multiple test
correction are highlighted in red. (F) Multiple regions of the north African *S. haematobium* genome lacked
introgressed *S. bovis* alleles. Regions that are longer than expected by chance are shown (Z-score$_{Length}$ >
3) in red. These regions, also known as introgression deserts, where not randomly distributed across the
genome, with 75% of them occurring on the sex chromosome. Results for FST and Patterson's D are
shown after Gaussian smoothing (sigma=3). Pink vertical lines indicate putative regions of adaptive
introgression.

1001

1002

Figure 3 – Species tree of *S. haematobium* and *S. bovis* populations – SVDquartets species tree generated from autosomal SNVs. All nodes were supported by >95% of bootstrap replicates. Phylogenetic relationships between the species can be used to differentiate extant vs ancestral hybridization (inset). The tree shows that both *S. haematobium* and *S. bovis* are monophyletic. Biogeographic partitioning within the tree indicates that *S. haematobium* originated in northern Africa and expanded into southern Africa.

1009
Figure 4. Mitochondrial tree of *S. haematobium* and *S. bovis* - A gene tree was recovered from
mitochondrial genome assemblies from each sample. Bootstrap support at select nodes is shown.
Phylogenetic relationships between the species can be used to differentiate extant vs ancestral
hybridization (inset). Two well supported clades of *S. haematobium* contain an introgressed *S. bovis*
mitotype, designated as "A" and "B". Both the "A" and "B" clades contain samples from north Africa. All
south African samples are found within a single clade of the remaining *S. haematobium* samples

1016

Figure 5 content (panels A, B, C):

Panel A: Map of Africa with Admixture Q1 color scale (1.0 to 0.0). Axis labels: 10°N, 0°, 10°S, 20°S, 30°S; 10°W, 0°, 10°E, 20°E, 30°E, 40°E, 50°E.

Panel B: Box plot of π (y-axis: 0.000, 0.005, 0.010, 0.015, 0.020) for categories: *S. bovis*, *S. haematobium*, *S. haem.* (North), *S. haem.* (South), *S. haem.* (North) masked.

Panel C: PCA plot, PC2 (24.4) vs PC1 (78.0). Labels: Sh (South), S. bovis, Sh (North), $F_{ST}=0.79$, $F_{ST}=0.16$, $F_{ST}=0.78$.

Species
S. haematobium ○
S. bovis ⬡

Countries
Cameroon, Cote d' Ivoire, Egypt, Gambia, Guinea Bissau, Liberia, Mali, Niger, Nigeria, Senegal, Sudan, Angola, Eswanti, Ethiopia, Kenya, Madagascar, Namibia, Tanzania, Uganda, Zambia, Zanzbiar

1017

Figure 5 Biogeography of *S. haematobium* is not determined by introgressed *S. bovis* alleles. *S. haematobium* samples were split into two populations by PCA, Admixture and phylogenetic analyses. (A) We used Kriging interpolation to examine the distribution of these populations across Africa using the population component that differentiates the *S. haematobium* populations from one another (B) Nucleotide diversity (π) was calculated in 10kb sliding windows after masking introgressed *S. bovis* alleles present in the northern *S. haematobium population*. π is higher in the northern African *S. haematobium* compared to the southern population. When introgressed *S. bovis* alleles are masked, π is similar for both the southern and northern populations. (C) After masking introgressed *S. bovis* alleles the PCA is similar to Figure 1C. The similarity between the two PCAs show that the genetic differentiation between the northern and southern *S. haematobium* populations is not driven by introgressed *S. bovis* alleles.

1028

1029

Supplemental Figure 1. Whole genome ancestry assignment with Admixture. We examined multiple
different population(k) sizes with Admixture. (A) At k=3, *S. haematobium* and *S. bovis* were separated, and
two general populations were identified within the *S. haematobium* samples corresponding to a northern
and southern population. (B) An optimal number of populations (k=5) (Evanno *et al.* 2005) shows clear
distinctions between the two *S. haematobium* populations.   Finally, (C) we generated a reference panel of
samples that maximized the *S. haematobium* and *S. bovis* population components from the k=3 results to
run a supervised admixture analysis and assign samples as either *S. haematobium* or *S. bovis*. This
analysis shows almost all of the northern *S. haematobium* samples contained low levels of *S. bovis* ancestry
at k=2, but this percentage various as more population components are added. Country Codes are as
follows: "AGO": Angola, "CMR": Cameroon, "CIV": Cote d' Ivoire, "EGY": Egypt, "SWZ": Eswanti, "ETH":
Ethiopia, "GMB": Gambia, "GNB": Guinea Bissau, "KEN": Kenya, "LBR": Liberia, "MDG": Madagascar,
"MLI": Mali, "NAM": Namibia, "NER": Niger, "NGA": Nigeria, "SEN": Senegal, "SDN": Sudan, "TZA":
Tanzania, "UGA": Uganda, "ZMB": Zambia, "ZAN": Zanzibar.

1043

1044

1045

Supplemental Figure 2. Comparison of non-*S. haematobium* ancestry calculated from Admixture (k=2) in each north African sample – Ancestry of each sample was assigned to up to two different population components with Admixture. These two components were maximized in samples from southern Africa or S. bovis samples. By comparison, north African samples were a composite of these two populaitions with low, but varying levels fo the *S. bovis* compponent found in each individual. The population component corresponding to *S. bovis* ancestry was significantly higher in Nigerian that in other north African countries (Kruskal-Wallis H test statistic = 7.915, P-value = 0.0049).

1053

1054

1055 **Supplemental Figure 3.** *S. bovis* **allele frequency across the genome within** *S. haematobium*
1056 **samples from Northern African countries -** The frequency of *S. bovis* ancestry across the genome is
1057 shown for each of the northwest African countries. In general, the distribution of *S. bovis* alleles is similar
1058 for each population. This consistency is an indicator of an ancient introgression event. The dotted line
1059 indicates 95% allele frequency.

1060

1061

1062 **Supplemental Figure 4. Pairwise comparison of introgressed *S. bovis* allele frequencies within**
1063 **northern *S. haematobium* samples by country –** Introgressed *S. bovis allele* frequency is positivley
1064 correlated between countries. Pearson's correlation coefficient (*R*) is >0.586 in all comparisons. The
1065 correlation of introgressed allele freuqencies between populations up to 3,338 Km apart is expected under
1066 an ancient introgression scenario.

1067

1068

Supplemental Figure 5 – Estimated number of generations since admixture with *S. bovis.* We estimated the number of generations since admixture for each sample in the northern *S. haematobium* population by examining the length of introgressed *S. bovis* loci with in the genomes. Individual estimates for each sample for each country as grey points. Results from a one-way ANOVA indicated that age estimates varied significantly between countries. Countries with a single individual, two haplotypes, are shown in blue and were not included in the ANOVA analyses. A "*" indicates p-values < 0.05. Differences in ages may indicate multiple introgression events.

1076

Table1. Genes containing outlier loci.

Chrom	Location	Gene_ID	Gene Name
Chr4	NC_067199.1:28466881-28497752	MS3_00007802	Leishmanolysin-like peptidase
Chr4	NC_067199.1:28497929-28529268	MS3_00007803	MS3_00007803
Chr4	NC_067199.1:28531133-28546611	MS3_00007804	RAD50
Chr4	NC_067199.1:28562068-28562732	MS3_00000457	JMJD6_1
Chr4	NC_067199.1:28571061-28634110	MS3_00010935	JMJD6_4
Chr4	NC_067199.1:28662329-28782419	MS3_00010934	JMJD6_3
Chr4	NC_067199.1:28742276-28747614	MS3_00007805	TY3BI_12
Chr4	NC_067199.1:28785057-28816546	MS3_00010936	JMJD6_5
Chr5	NC_067200.1:9933321-9974793	MS3_00011123	MS3_00011123
Chr5	NC_067200.1:9989479-10043873	MS3_00011124	AK2_3
Chr5	NC_067200.1:10117557-10118821	MS3_00011125	MS3_00011125
Chr5	NC_067200.1:10187316-10209172	MS3_00011126	TSC2
Chr5	NC_067200.1:10219534-10229257	MS3_00009120	MDP1_1
Chr5	NC_067200.1:10245731-10246840	MS3_00000691	MS3_00000691
Chr5	NC_067200.1:10414911-10585361	MS3_00011127	MS3_00011127