The *Cryptococcus neoformans* STRIPAK complex controls genome stability, sexual development, and virulence

Patricia P. Peterson[1], Jin-Tae Choi[2], Ci Fu[3], Leah E. Cowen[3], Sheng Sun[1], Yong-Sun Bahn[2], and Joseph Heitman[1*]

1 Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, North Carolina, USA.

2 Department of Biotechnology, College of Life Science and Biotechnology, Yonsei University, Seoul, Republic of Korea.

3 Department of Molecular Genetics, University of Toronto, Toronto, Ontario, Canada.

* Corresponding author
Email: heitm001@duke.edu (JH)

Short title: Pathobiological roles of the *C. neoformans* STRIPAK complex

Abstract

The eukaryotic serine/threonine protein phosphatase PP2A is a heterotrimeric enzyme composed of a scaffold A subunit, a regulatory B subunit, and a catalytic C subunit. Of the four known B subunits, the B''' subunit (known as striatin) interacts with the multi-protein striatin-interacting phosphatase and kinase (STRIPAK) complex. Orthologs of STRIPAK components were identified in *C. neoformans*, namely PP2AA/Tpd3, PP2AC/Pph22, PP2AB'''/Far8, STRIP/Far11, SLMAP/Far9, and Mob3. Structural modeling, protein domain analysis, and detected protein-protein interactions suggest *C. neoformans* STRIPAK is assembled similarly to the human and fungal orthologs. Here, STRIPAK components Pph22, Far8, and Mob3 were functionally characterized. Whole-genome sequencing revealed that mutations in STRIPAK complex subunits lead to increased segmental and chromosomal aneuploidy, suggesting STRIPAK functions in maintaining genome stability. We demonstrate that *PPH22* is a haploinsufficient gene: heterozygous *PPH22/pph22Δ* mutant diploid strains exhibit defects in hyphal growth and sporulation and have a significant fitness disadvantage when grown in competition against a wild-type diploid. Deletion mutants *pph22Δ, far8Δ*, and *mob3Δ* exhibit defects in mating and sexual differentiation, including impaired hyphae, basidia, and basidiospore production. Loss of either *PPH22* or *FAR8* leads to growth defects at 30ºC, severely reduced growth at elevated temperature, abnormal cell morphology, and impaired virulence. The *pph22Δ* and *far8Δ* mutants are also unable to grow in the presence of the calcineurin inhibitors cyclosporine A or FK506, and thus these mutations are synthetically lethal with loss of calcineurin activity. Conversely, *mob3Δ* mutants display increased thermotolerance, capsule production, and melanization, and are hypervirulent in a murine infection model. Taken together, these findings reveal that the *C. neoformans* STRIPAK complex plays an important role in genome stability, vegetative growth, sexual development, and virulence in this prominent human fungal pathogen.

Introduction

Eukaryotic organisms utilize dynamic signaling networks to respond and adapt to changes in their internal and external environments. Sensing of environmental stimuli triggers a cascade of downstream events, eliciting a coordinated cellular response tightly orchestrated by interconnected signal transduction pathways. The function and activity of signaling components within these pathways are modulated through post-translational modifications, including protein phosphorylation, which is governed by the balanced actions of kinases and phosphatases. Phosphorylation and dephosphorylation of target proteins by kinases and phosphatases are coordinated by finely-tuned mechanisms to ensure signaling pathways are turned on and off as needed, maintaining cellular homeostasis. Among eukaryotic phosphatases, protein phosphatase 2A (PP2A) plays a pivotal role in governing numerous cellular processes, including cell cycle progression, proliferation, apoptosis, metabolism, and stress responses [1].

PP2A is a heterotrimeric enzyme composed of a scaffold A subunit, a regulatory B subunit, and a catalytic C subunit. Among the four known B subunits, the B''' subunit (known as striatin) interacts with the striatin-interacting phosphatase and kinase (STRIPAK) complex [2-5]. First identified in mammals, striatin proteins act as scaffolds to assemble the other STRIPAK subunits, forming a large, multifunctional signaling complex. In addition to the PP2A holoenzyme, STRIPAK is comprised of a striatin-interacting protein (STRIP), monopolar spindle-one-binder related protein (Mob3/Phocein), sarcolemmal membrane-associated protein (SLMAP), small coiled-coil protein (SIKE), cerebral cavernous malformation protein (CCM3), and a germinal center kinase (GCKIII) [3]. Orthologs of the mammalian STRIPAK complex have been identified in many fungi, including *Sordaria macrospora, Neurospora crassa, Saccharomyces cerevisiae, Schizosaccharomyces pombe, Aspergillus nidulans, Fusarium graminearum,* and several other ascomycete species [6-22].

In mammals and fungi, the STRIPAK complex connects signal transduction pathways to regulate numerous aspects of cell growth and development, such as TORC2 signaling and actin cytoskeleton remodeling in *S. cerevisiae,* and the Hippo pathway and regulation of tissue growth in humans [14, 23, 24]. In fungi, STRIPAK plays a critical role in morphogenesis and sexual development, including control of cell fusion, hyphal elongation, fruiting body formation, fertility, nuclear division, and sporulation [25, 26]. The *S. cerevisiae* STRIPAK counterpart, the Far complex, is implicated in pheromone-induced cell cycle arrest during mating, acts as an antagonist to TORC2 at the endoplasmic reticulum (ER), and inhibits mitophagy at the mitochondrial membrane [14, 27-30]. In *N. crassa*, homologs of STRIP and Striatin act on two MAP kinase pathways, downstream of the cell wall integrity (CWI) and pheromone response pathways, modulating fungal self-signaling and developmental morphogenesis [8]. The STRIPAK complex also governs virulence in plant fungal pathogens including *Magnaporthe oryzae* and several *Fusarium* species [12, 13, 31-33].

Although the STRIPAK complex has been characterized in pathogenic and non-pathogenic ascomycetes, it has not yet been elucidated in a basidiomycete or notable human fungal pathogen. The basidiomycetous yeast and opportunistic pathogen *Cryptococcus neoformans* is a clinically relevant and genetically tractable model for studying the molecular mechanisms underlying fungal pathogenesis in humans. *C. neoformans* infections occur following inhalation of spores or desiccated yeast cells from the environment and are more prevalent in immunocompromised hosts. Systemic cryptococcosis can lead to lethal meningoencephalitis, accounting for approximately 20% of HIV/AIDS-related deaths annually, thus representing a significant burden of global fungal diseases [34-36]. *C. neoformans* possesses virulence traits necessary to cause disease, including melanization, extracellular polysaccharide capsule production, resistance to oxidative stress, and thermotolerance [37-39].

Understanding the regulatory pathways underlying *C. neoformans* virulence is critical for

developing targeted therapies against cryptococcosis.

Due to the key roles of STRIPAK in controlling cell growth, developmental processes,

and pathogenicity in other fungi, this study aimed to functionally characterize the cellular

functions of the *C. neoformans* STRIPAK complex. We identified protein orthologs for six

STRIPAK complex subunits, three of which, Pph22 (PP2A catalytic subunit)*,* Far8 (regulatory

B‴ subunit)*,* and Mob3, were characterized. Genome sequencing of *pph22Δ*, *far8Δ*, and *mob3Δ*

deletion mutants revealed frequent segmental and whole chromosomal aneuploidy, suggesting

a role in maintaining genome stability. We demonstrate that *PPH22* is not an essential gene,

though the loss of one functional copy leads to haploinsufficiency in a heterozygous diploid

strain, and the *pph22Δ* mutation incurs a significant fitness cost in a haploid. Similar to findings

in other fungi, *C. neoformans* STRIPAK is important for mating, hyphal formation, and

sporulation. Both Pph22 and Far8 are crucial for high-temperature growth, stress response, and

virulence. Surprisingly, *mob3Δ* mutants exhibited increased thermotolerance, melanization, and

capsule production, resulting in hypervirulence in a murine model of *C. neoformans* infection.

Taken together, these studies shed light on the organization and function of the STRIPAK

complex in *C. neoformans* and provide a foundation for future identification of its cellular targets

to understand the mechanisms mediating its various roles.

Results

Identification of *Cryptococcus neoformans* STRIPAK complex components

To identify the *C. neoformans* STRIPAK complex, BLAST searches (blastp) were performed with known *Homo sapiens* and *S. cerevisiae* STRIPAK components and the *C. neoformans* H99 genome (Figure 1A). Significant alignments were found for the protein phosphatase 2A catalytic C subunit (CNAG_02177/Pph22), the scaffold A subunit (CNAG_07914/Tpd3), the regulatory B‴ subunit (CNAG_00073/Far8), the striatin-interacting protein (CNAG_00008/Far11), the tail-anchor domain protein (CNAG_04838/Far9), and the striatin-associated protein (CNAG_04629/Mob3). No ortholog of the coiled-coil domain protein (Far3/7 in *S. cerevisiae,* SIKE in *H. sapiens,* Sci1 in *S. macrospora,* SipB in *A. nidulans,* and Csc4 in *S. pombe*) was identified, possibly due to its small size and low sequence similarity [16]. BLAST analysis of the STRIPAK-associated kinase with the sequences of the three mammalian kinases Stk24, Stk25, and Stk26 [3], which are part of the GCKIII family of kinases, produced three significant alignments in *C. neoformans* (CNAG_03290, CNAG_00405, and CNAG_05274). Reciprocal BLAST analyses confirmed an orthologous relationship between the proteins. With the exception of *PPH22* [40]*,* none of the *C. neoformans* STRIPAK complex components were annotated on FungiDB (https://fungidb.org) and thus are referred to here based on *S. cerevisiae* or *H. sapiens* nomenclature.

Multi-sequence alignments revealed the *C. neoformans* STRIPAK components share 23%-86% identity and 42%-93% similarity with the *S. cerevisiae* and *H. sapiens* orthologs, with greater homology to the human than the yeast proteins (Figure 1B). Mob3 from *C. neoformans* only aligned with the human protein, as *S. cerevisiae* lacks Mob3. The STRIPAK components also exhibit highly conserved structures and domain architecture [26] (Figure 1B). For example, CnTpd3 contains 15 tandem HEAT (huntingtin-elongation-A subunit-TOR) repeats, which mediate interactions with the B and C subunits to form the PP2A holoenzyme [1]. Eukaryotic

striatin proteins share a coiled-coil domain and a WD40 repeat domain, forming a β-propeller.

Human striatin interacts with the PP2A A and C subunits via the coiled-coil domain [41, 42]. The

tail-anchor domain protein CnFar9 contains a conserved FHA (forkhead-associated) domain

and a small, hydrophobic transmembrane domain at its C-terminus. In fungi and mammals,

mutations in the tail-anchor domain of Far9/SLMAP lead to changes in their subcellular

localization, and the membrane association of Far9 with ER and mitochondria is important for its

functions [14, 29, 43].

Due to the evolutionarily conserved nature of the *C. neoformans* STRIPAK complex

protein sequences and domains (Figure 1B), it was predicted that the complex would also have

a similar three-dimensional structure and organization. Recently, the core of the human

STRIPAK complex was resolved at high resolution by cryo-EM [42]. The *C. neoformans* proteins

were aligned with the model of the human STRIPAK core with the program ChimeraX [44]

(EMD-22650, PDB-7k36) (Figure 1C). The predicted and known protein structures are

remarkably similar. The *C. neoformans* STRIPAK complex was predicted with AlphaFold2

multimer [45] (Figure 1D). This structure prediction included Far9, which has not been identified

in the cryo-EM model or crystal structures of human STRIPAK [41, 42]. The predicted

CnSTRIPAK complex model showed a linear arrangement of the proteins Mob3-Far11-Pph22-

Tpd3, similar to the human complex. The Far8 WD40 domain lies in close proximity to Mob3,

and the coiled-coil domain is at the base of the complex. The middle of the Far9 protein

contains a large coiled-coil domain (approximately 135 amino acids) that interacts closely with

the coiled-coil of Far8, stabilizing the core complex. Taken together, these results suggest that

the key components of the *C. neoformans* STRIPAK complex are significantly conserved in their

domain architecture, three-dimensional structure, and assembly. Our data also suggest the

CnSTRIPAK complex is more similar to human STRIPAK than to the *S. cerevisiae* counterpart.

Yeast two-hybrid analysis of subunit interactions of the STRIPAK complex

We next sought to detect physical interactions between STRIPAK components predicted to be in close proximity. To this end, yeast two-hybrid analysis of each CnSTRIPAK subunit was performed (Figure 2A). Plasmids encoding fusion proteins of the Gal4 DNA-binding domain and CnSTRIPAK subunits Pph22, Tpd3, Far8, Far11, Far9, and Mob3 were generated and transformed into the yeast two-hybrid reporter strain Y2HGold. Similarly, plasmid constructs encoding the Gal4 transcriptional activation domain fused to individual STRIPAK subunits were transformed into reporter strain Y187. A 6-by-6 crossing between Y2HGold and Y187 strains carrying plasmids encoding GBD and GAD fusion proteins was conducted and assayed for Gal4-dependent expression of the *ADE2* and *HIS3* reporter genes (Figure S1)*.* For Pph22 and Tpd3, because the GBD fusions alone were able to activate reporter gene expression, only the GAD fusions of these two proteins were analyzed to determine their interactions with GBD-fused STRIPAK components. Positive interactions between Mob3 and Far8 were detected in both configurations, suggesting specific and robust binding between these two proteins (Figure 2A). Positive protein-protein interactions were also detected for Far9-Tpd3, Far11-Pph22, and Far11-Tpd3 (Figure 2A). This yeast two-hybrid analysis provides further support for the predicted CnSTRIPAK model (Figures 1D and 2B).

Mutations in genes encoding the STRIPAK complex components lead to genome instability

To analyze the function of the *C. neoformans* STRIPAK complex, deletion mutations were generated for genes encoding STRIPAK complex subunits. We first selected the genes encoding the PP2A trimeric enzyme, Pph22, Tpd3, and Far8, for targeted deletion mutagenesis. While we were able to successfully generate *far8Δ* deletion strains, we failed to obtain *pph22Δ*

and *tpd3Δ* mutant strains in H99α or KN99**a** haploid wild-type strains, after multiple attempts of

both biolistic transformation and CRISPR-Cas9 gene deletion approaches, suggesting that

PPH22 and *TPD3* might be essential for cell viability. In *S. cerevisiae,* the deletion of *TPD3*

causes growth defects [46], and deletions of *PPH21* and *PPH22* are synthetically lethal [47]. In

C. neoformans, a genome-wide functional analysis of phosphatases suggested that *PPH22* is a

putative essential gene [40]. Conversely, *FAR8* and other striatin subunit homologs have not

been characterized as essential in other species.

We then took a different approach to test the essentiality of *PPH22* and *TPD3,* as well as

the other STRIPAK component encoding genes *FAR9*, *FAR11*, and *MOB3*. Specifically, we

sought to delete one of the two alleles for each gene in a wild-type diploid strain CnLC6683,

which is a fusion product between the congenic strain pair KN99**a** and KN99α, and then isolate

haploid deletion mutants by sporulating and dissecting meiotic progeny. Using CRISPR-Cas9

directed mutagenesis, we successfully obtained *PPH22/pph22Δ* and *MOB3/mob3Δ*

heterozygous mutants in strain CnLC6683 background. We were unable to isolate heterozygous

TPD3/tpd3Δ, *FAR9/far9Δ*, or *FAR11/far11Δ* mutant strains for this study; therefore, we

concentrated our analyses on *PPH22/pph22Δ* and *MOB3/mob3Δ*. Four *PPH22/pph22Δ* strains

were generated from two independent transformations, and three *MOB3/mob3Δ* strains from

two independent transformations. PCR genotyping and Illumina whole-genome sequencing

confirmed heterozygosity at the *PPH22* and *MOB3* loci. Analysis of read depth revealed multiple

segmental and chromosomal aneuploidies throughout the genome (Figure 3A), including

increased coverage for entire chromosomes, and both increased and decreased coverage in

segments of chromosomes. Whole chromosome 13 trisomy was observed in *PPH22/pph22Δ-3,*

4 and *MOB3/mob3Δ-1, 2, 3* strains, while partial trisomy of this chromosome was seen in

PPH22/pph22Δ-1. Partial trisomy for additional chromosomes was observed in other cases:

chr.4 in *PPH22/pph22Δ-1*; chr. 3 in *PPH22/pph22Δ-2*; chr. 6 in *PPH22/pph22Δ-3* and

MOB3/mob3Δ-1; and chr.9 and chr. 10 in *MOB3/mob3Δ*-1. Segmental monosomy was also

observed in chr. 2 and chr. 4 in *PPH22/pph22Δ*-4; and chr. 10 in *MOB3/mob3Δ*-1, 3.

PPH22/pph22Δ-1 also exhibited increased coverage for chr. 1 and chr. 6, although to a lesser

extent.

We next assessed whether the genetic changes present in the heterozygous mutant

diploid populations would be inherited and maintained in meiotic progeny. *PPH22/pph22Δ* and

MOB3/mob3Δ cells were incubated on MS medium at room temperature for at least three weeks

to induce self-filamentation. Basidiospores were dissected, and haploid *pph22Δ* and *mob3Δ*

deletion mutant progeny were validated through PCR genotyping (Figure S2). Haploid *pph22Δ*

deletion mutants were obtained from the dissection of spores from *PPH22/pph22Δ*-2, *3*, and *4*,

and *mob3Δ* mutants were similarly obtained from *MOB3/mob3Δ*-2. Whole-genome sequencing

supported the PCR analysis, showing no reads mapping to the *PPH22* or *MOB3* genes.

Therefore, *PPH22* and *MOB3* are not essential in *C. neoformans*. Single colonies from *pph22Δ*

and *mob3Δ* deletion mutants were passaged at least three times prior to whole-genome

sequencing to obtain a pure cell population and reduce the likelihood of sequencing a

population of cells with mixed genotypes. Interestingly, whole-genome sequencing analysis

showed that whole chromosome aneuploidy was present in some but not all haploid mutant

progeny, and that the genomic profile of the progeny often did not mirror the genomic profile of

the parent (Figures 3A and 3B). Haploid *pph22Δ* progeny from *PPH22/pph22Δ*-2 exhibited

increased coverage for chr. 3 and chr. 8, while the parental diploid strain was euploid for those

chromosomes. One other haploid strain showed aneuploidy that was not present in its diploid

parent: P3 dissected from *MOB3/mob3Δ*-2 was disomic for chr. 7 (Figures 3A and 3B). The

most common aneuploidy present in the heterozygous diploid progenitor strains was associated

with chr. 13 (Figure 3A), which was found in P2 from *PPH22/pph22Δ*-4 and P1-P3 from

MOB3/mob3Δ-2. No ploidy changes were detected in any of the haploid and diploid control

strains from each experiment. These results suggest that the *PPH22* and *MOB3* genes play

roles in maintaining genome stability.

Genetic analysis of *FAR8* revealed a related role in genome stability. The *far8Δ* mutants

were successfully isolated via biolistic transformation in the congenic H99α and YL99**a** haploid

strains. Deletion of *FAR8* was confirmed by PCR genotyping and Illumina whole-genome

sequencing (Figure S3). Read-depth analyses indicated that the four independent *far8Δ* mutants

in H99α had rampant aneuploidy throughout the genome, suggesting that they were diploid

instead of haploid (Figure 3C). Similar results were obtained from whole-genome sequencing of

YL99**a** *far8Δ* strains (Figure S4). Among seven independently obtained *far8Δ* mutant strains, all

were aneuploid for at least six out of 14 chromosomes, and the only chromosome that was

never found to be duplicated was chr. 3. The parental H99α and YL99**a** isolates were haploid,

confirming the aneuploidy occurred after deletion of *FAR8.* Complementation of the *far8Δ*

mutation in the H99α background strain *far8Δ-1* yielded a euploid population, suggesting

reintroduction of *FAR8* prevents formation of aneuploidy.

Fluorescence-activated cell sorting (FACS) analysis was conducted to determine the

DNA content in the STRIPAK mutants (Figures 3D and S4). A wild-type haploid shows a major

1C peak and a minor 2C peak, while a wild-type diploid has a major 2C peak and a minor 4C

peak. The *pph22Δ* mutants displayed either one single merged, widened peak, or additional

widened peaks at 4C or larger, indicating that there is a heterogeneous population of cells with

different DNA content with modest to severe aneuploidy. Another possibility is that *pph22Δ*

mutants form clusters or have defects in cytokinesis, as described later in Figure 4C. FACS

analysis of *far8Δ* mutants showed two major peaks at 2C and 4C, suggesting that these strains

are largely diploid. FACS data from the *far8Δ::FAR8* complemented strain had a similar peak

profile to the 1N control strain H99α, indicating that it is haploid. The *mob3Δ* mutants were also

analyzed by FACS and exhibited only two peaks at 1C and 2C, indicating they are haploid

(Figure S4).

Taken together, whole-genome sequencing and FACS analysis showed that *pph22Δ*,

far8Δ, and *mob3Δ* mutations cause genome instability, resulting in aneuploidy caused by whole

chromosome and segmental duplication/deletion. Our data also suggest that the deletion of

FAR8 might have led to genome endoreplication, which was followed by rampant chromosomal

losses resulting in aneuploidy, suggesting a role in cell cycle control. These results demonstrate

that the *C. neoformans* STRIPAK complex is important for maintaining genome stability.

PPH22 is a haploinsufficient gene in diploid *C. neoformans*

During self-filamentation, we observed that *PPH22/pph22Δ* strains showed delayed

production of hyphae, basidia, and basidiospores compared to the wild-type control. To analyze

this further, equal amounts of WT/WT (CnLC6683) and *PPH22/pph22Δ* diploid cells were plated

on MS media and incubated at room temperature. After 1 to 2 weeks, the wild-type diploid strain

produced abundant hyphae and basidia, and basidia with spore chains were also visible.

However, no hyphae were seen in the *PPH22/pph22Δ* strains at that time. After 3 to 4 weeks of

incubation, the patch of *PPH22/pph22Δ-1* had robust filamentation (Figure 4A), and hyphae,

basidia, and spore chains were observed at high magnification. However, patches of

PPH22/pph22Δ-2, 3, 4 cells exhibited only minimal hyphae, basidia, and spores. Hyphae,

basidia, and basidiospores of the self-filamenting diploid strains were observed by scanning

electron microscopy (SEM) (Figure 4B). The basidia in *PPH22/pph22Δ* strains exhibited

abnormal and fewer spore formation, with spores produced in irregular clusters instead of

chains. The morphology of the spores was also atypical, with significant variation in shape and

size. There were also instances of ectopic hyphal growth, as seen in *PPH22/pph22Δ-4*, where

basidia appeared to produce a new hyphal filament instead of spores (Figure 4B).

The defects in sexual development observed in *PPH22/pph22Δ* strains led us to

hypothesize that *PPH22* might be haploinsufficient. When WT/WT and *PPH22/pph22Δ* strains

were grown to mid-logarithmic phase in YPD liquid media at 30°C and cell morphology was

observed using differential interference contrast (DIC) microscopy, *PPH22/pph22Δ* cells were

typically larger and formed abnormal clusters compared to wild-type diploid cells (Figure 4C).

Some *PPH22/pph22Δ* cells were elongated and showed incomplete budding. The

PPH22/pph22Δ-1-4 strains also exhibited severe growth defects compared to the WT/WT strain

in liquid culture (Figure 4D). In competition assays to compare the fitness of the WT/WT and

PPH22/pph22Δ strains, the *pph22Δ* deletion allele in the heterozygous diploid strain conferred a

significant competitive disadvantage compared to the wild type (Figure 4E). To determine if this

disadvantage was due to reduced viability of *PPH22/pph22Δ* strains, *PPH22/pph22Δ* and

WT/WT cells that had been grown in YPD overnight cultures for the competition assays were

serially diluted and plated onto YPD medium at 30°C (Figure S5). There was no observable

difference in growth between the wild-type and heterozygous mutant diploid strains, indicating

cells were similarly viable at the time of competition. The significant reduction in the

PPH22/pph22Δ population at the end of the competition assay likely results from its inherent

growth defects, leading to reduced fitness compared to the wild type. Thus, the loss of one

PPH22 allele in a diploid leads to defects in sexual development, cell morphology, and

competitive growth, indicating *PPH22* is a haploinsufficient gene. This may be attributable to

loss of one copy of *PPH22*, aneuploidy arising due to reduction in *PPH22* level, or both.

Haploid *pph22Δ* mutants exhibit severe growth defects and frequently accumulate suppressor mutations

To further characterize the functions of *PPH22*, haploid *pph22Δ* mutants were obtained for phenotypic analysis. Self-filamenting *PPH22/pph22Δ-1-4* strains produced sufficient basidiospores for dissection after 4 to 6 weeks of incubation on MS plates. *pph22Δ* mutant colonies were much smaller than wild-type colonies on the dissection plate (Figure 5A). The severe growth defects of *pph22Δ* mutants likely explain why *PPH22* was initially thought to be essential. *pph22Δ* mutants formed tan colonies on YPD plates incubated at 30°C. After prolonged incubation, faster-growing white colonies appeared, possibly as a result of suppressor mutations (referred to as *pph22Δ suppressor* or *pph22Δ sup* mutants) (Figure 5B). While *pph22Δ* strains grew poorly compared to wild type at 24°C and 30°C, produced tan colonies, and could not grow at 33°C, the *pph22Δ sup* strains exhibited near wild-type growth at 24°C, 30°C, and 33°C, and produced whiter colonies (Figure 5C).

Ten *pph22Δ suppressor* strains were subjected to whole-genome sequencing to identify the causative suppressor mutation(s). Variant calling failed to identify any significant single-nucleotide polymorphisms (SNPs) or insertion/deletion mutations. However, all 10 *pph22Δ suppressor* mutant strains exhibited segmental aneuploidy on chr. 6 in a shared overlapping region of ~200 kbp (Figure 5D) and were euploid for the remainder of their genomes. Read depth in this region ranged from 2X to 11X coverage compared to the mean coverage of the remainder of the chromosome. The *PPH22* gene lies in this region, but as expected, no reads were found mapping to the *PPH22* locus. All 10 isolates shared an approximately 15 kbp region adjacent to the centromere, encompassing 7 genes (Figures 5D and 5E). One notable candidate for the *pph22Δ suppressor* gene is *PPG1,* which encodes a serine/threonine type 2A-like protein phosphatase involved in the cell wall integrity pathway [48] that shares 56% identity and 73% protein sequence similarity with Pph22. Studies from *S. cerevisiae* have shown that

313   Ppg1, and not Pph22, interacts with Far11 to regulate assembly of the Far complex [49, 50],

314   which may have pleiotropic roles beyond those of the STRIPAK complex. Our data suggest that

315   an increased dosage of *PPG1* may suppress the mutant phenotypes of *pph22Δ*.

STRIPAK is important for mating and sexual development in *C. neoformans*

To investigate roles for the *C. neoformans* STRIPAK complex in sexual development, we

analyzed the effects of mutations in STRIPAK components on mating and the sexual cycle using

PPH22/pph22Δ and *MOB3/mob3Δ* heterozygous diploid strains compared to WT/WT (Figure

6A). As observed previously, *PPH22/pph22Δ* exhibited severe self-filamentation defects.

Similarly, *MOB3/mob3Δ* displayed defects in sexual development, showing a smaller area of

self-filamentation and shortened hyphae compared to the wild type. Although *MOB3/mob3Δ* was

able to produce basidia and basidiospores, this occurred only near the edge of the cell patch,

indicating that *MOB3* is also important for sexual development in a self-filamenting diploid.

We next examined the role of the STRIPAK complex during bisexual mating. Mutant

cells (*pph22Δ*, *pph22Δ suppressor, mob3Δ*, and *far8Δ*) were mixed with wild-type cells of the

opposite mating type, spotted onto MS media, and incubated at room temperature (Figure 6B).

The H99α x KN99**a** cross served as a wild-type control. Initial crosses involving *pph22Δ* and

far8Δ strains did not show any signs of mating, even after 8 weeks. This issue could be due to

the growth defects of these strains on MS media, leading to them being outcompeted by the

wild-type partner. Additional crosses were conducted utilizing two strategies to promote mating:

(1) mixing mutant and wild-type cells in a 10:1 ratio, and (2) growing mutant cells alone for two

to three days on MS media before plating wild-type cells on top. These strategies enabled

mating in *pph22Δ* x WT and *far8Δ* x WT crosses. *pph22Δ*, *pph22Δ suppressor,* and *far8Δ*

crosses with WT could form large branches of hyphae extending from the mating patch, similar

to the wild-type control. The *pph22Δ* x WT cross could produce hyphae, but mostly bald basidia

heads with almost no spore formation. This phenotype was partially rescued in the *pph22Δ*

suppressor mutants, which produced basidia with basidiospores after mating, albeit to a lesser

extent than the wild-type control.

Mating of *far8Δ* x WT strains is an unusual case, as the *far8Δ* mutant contains a largely

diploid genome. This results in a diploid-by-haploid cross followed by triploid meiosis. Similar to

pph22Δ x WT crosses, we observed mostly bald basidia heads with no spores in *far8Δ* x WT

crosses, though some basidia with spore chains were found at low frequency. Crossing WT with

the *far8Δ*::*FAR8* complemented strain successfully restored mating and sporulation efficiency to

the wild-type level. The *mob3Δ* and wild-type cells formed stunted hyphae and a smaller area of

hyphal growth, resembling the self-filamentation of the *MOB3/mob3Δ* heterozygous diploid

strain, but produced abundant basidia and basidiospore chains. We did not observe mating

during bilateral crosses between STRIPAK component deletion mutants (Figure S6). This

suggests that there are additive interactions between the defects caused by each individual

deletion of either *PPH22*, *FAR8*, or *MOB3*. In summary, the STRIPAK complex plays a critical

role in various aspects of sexual development in *C. neoformans.*

Phenotypes of *pph22Δ* and *far8Δ* mutants in response to nutrients, temperature, and

stress

Next, the role of STRIPAK in vegetative growth and stress response was analyzed.

pph22Δ and *far8Δ* mutant strains, along with *pph22Δ suppressors*, *far8Δ*::*FAR8,* and isogenic

wild-type control strains were serially diluted and spotted onto media under different nutrient,

temperature, and stress conditions (Figures 7A and 7B). *pph22Δ* mutants exhibited severe

growth defects on YPD, which was exacerbated by supplementation with 1 M sorbitol (Figure

7A). This finding could explain why we were unable to generate a haploid *pph22Δ* deletion

mutant via biolistic transformation, during which sorbitol medium was used to provide osmotic

support for cells to recover from puncture by DNA-coated gold particles [51].

The *pph22Δ* mutants were then tested on YPD with rapamycin, which inhibits TORC1

and mimics nutrient starvation (Figure 7A). Interestingly, *pph22Δ* grew slightly better than the

wild type, while the *pph22Δ #1 sup1* and *pph22Δ #2 sup2* strains exhibited robust growth on this

medium. This suggests that *pph22Δ* leads to rapamycin tolerance, and that the suppressor

mutations of *pph22Δ*, potentially via overexpression of the phosphatase *PPG1*, can confer

further resistance to rapamycin. On YPD medium at 37°C, neither *pph22Δ* nor *pph22Δ sup*

strains grew, indicating *PPH22* is required for high-temperature growth, and the suppressor

mutation does not rescue the defect. The *pph22Δ* and *pph22Δ sup* cells also failed to grow in

the presence of the immunosuppressive drugs FK506 or cyclosporine A at 30°C, both of which

inhibit the activity of the PP2B phosphatase, calcineurin, illustrating synthetic lethality due to the

loss of two phosphatases.

A similar role in nutrient and stress response was investigated for *FAR8.* In serial dilution

assays, *far8Δ* cells grew more slowly on YPD at 30°C and formed smaller colonies compared to

the wild type (Figure 7B). This growth defect was partially rescued by the supplementation of

sorbitol. Contrary to our observations of *pph22Δ* cells grown in the presence of rapamycin, *far8Δ*

strains were severely growth impaired on this medium compared to wild type, suggesting that

these strains have different responses to TORC1 inhibition. *far8Δ-1* and *-2* strains exhibited

severe growth defects on YPD at 37°C, while *far8Δ-3* and *-4* strains (constructed in the YL99**a**

background) exhibited almost no growth. Similar to *pph22Δ* mutants, *far8Δ* strains did not grow

in the presence of FK506 or cyclosporine A at 30°C, suggesting a synthetic lethal interaction

between the deletion of *FAR8* and the inhibition of calcineurin by FK506 or cyclosporine A.

These findings suggest that PP2A is important for cell growth under various stress

conditions and that *PPH22* in particular may have a role in response to osmotic stress or in

maintaining cell wall integrity. The mutants were tested for sensitivity to cell wall-disrupting

agents (calcofluor white and caffeine) (Figure 7C). *pph22Δ* mutants were unable to grow in the

presence of calcofluor white, which was partially rescued in the *pph22Δ #1 sup1* and *pph22Δ #2*

sup2 strains. In the presence of calcofluor white, *far8Δ* mutants exhibited a subtle growth defect

compared to WT, similar to YPD alone. On media supplemented with caffeine, both wild-type

and *pph22Δ* cells grew slightly poorer and formed smaller colonies than on YPD, while the

pph22Δ sup strains exhibited strong caffeine sensitivity. *far8Δ* cells showed even higher caffeine

sensitivity and exhibited almost no growth. Therefore, *pph22Δ* and *far8Δ* mutants display

opposite phenotypes in response to calcofluor white and caffeine. DIC microscopy images of

pph22Δ and *far8Δ* cells grown in nutrient-limiting synthetic complete (SC) medium also revealed

changes in cell morphology (Figure 7D). Specifically, while cells of the wild-type strain were

homogenously round and typical in size, both *pph22Δ* and *far8Δ* mutants produced

heterogenous cell populations, which were mixtures of normal-looking cells with elongated cells

(*pph22Δ*) or considerably enlarged cells (*far8Δ*), suggesting both *PPH22* and *FAR8* are

important for faithful mitotic cytokinesis.

The above growth analyses of the mutant strains indicated severe growth impairment

when exposed to conditions affecting cell wall integrity. We hypothesized that *pph22Δ* and *far8Δ*

mutants might exhibit hypersensitivity to the clinically relevant antifungal drug fluconazole, which

targets the ergosterol biosynthesis pathway and weakens the fungal cell membrane. We

investigated this with Etest strips (Figure 7E). As predicted, the *far8Δ* strain showed a larger

zone of inhibition and increased susceptibility to fluconazole compared to the wild-type control.

Surprisingly, *pph22Δ* mutants were completely resistant to fluconazole at the full range of

concentrations on the Etest strip, suggesting an MIC (minimum inhibitory concentration) value

greater than 256 μg/mL. A *pph22Δ sup* mutant was also highly resistant to fluconazole. Similar

results were obtained for independent *pph22Δ*, *pph22Δ sup,* and *far8Δ* mutant strains (Figure

S7).

PPH22 and FAR8 are crucial for melaninization, capsule production, and virulence in *C.*

neoformans

The role of PP2A-STRIPAK components, *PPH22* and *FAR8*, was next addressed in the

production of two important virulence factors in *C. neoformans*: melanin and the polysaccharide

capsule. On Niger seed agar, *pph22Δ* mutants showed severe growth impairment, complicating

the assessment of melaninization (Figure 8A). The *pph22Δ sup* strains also exhibited growth

defects but produced some melanin pigment. The *far8Δ* strains failed to produce melanin on this

media and remained white, similar to the *lac1Δ* mutant control. Copper availability is known to

regulate melaninization [52], prompting us to test whether the melanin defect in *far8Δ* strains is

copper-dependent. Melanin production of the mutants was assessed on L-DOPA medium,

supplemented with either copper sulfate or the copper chelator, bathocuproine disulfonate

(BCS) (Figure 8B). For controls, we used strains H99α (produces melanin), H99α *cbi1Δ ctr4Δ*

(produces melanin only when copper is present), and H99α *lac1Δ* (melanin production

defective). Under the copper-deficient condition (BCS), *pph22Δ* produced minimal melanin,

which was partially restored in the suppressor mutant, resembling observations on Niger seed

medium. The *far8Δ* strains did not produce melanin in the presence of BCS. When copper

sulfate was added, the *pph22Δ* strains did not show increased melaninization, but the *pph22Δ*

#1 sup1 strain did show a copper-dependent increase in pigment. The *far8Δ* strains were only

able to produce melanin when copper was added to the media, suggesting that their inability to

make melanin is partially linked to defects in maintaining copper homeostasis.

Capsule production was assessed in WT, *pph22Δ, pph22Δ sup, far8Δ*, and *far8Δ::FAR8*

strains by growing cells in RPMI medium at 30°C, and then analyzing capsule size with India ink

(Figure 8C). The *pph22Δ* mutants grew very slowly in this nutrient-poor medium, produced no

visible capsule, and formed elongated cells. The *pph22Δ sup* strains produced small capsules

with normal cell morphology (Figure 8C, top panel). The *far8Δ* strains produced capsules

relatively similar in size to the wild type, but the cells were significantly larger, appearing as titan

cells, which are enlarged cells greater than 15 μm in diameter [53, 54]. Cells larger than 10 μm

accounted for ~30% of the total cell population. This experiment was repeated to include a

WT/WT diploid control, ruling out the possibility that the large cell size was due to the diploid

nature of the *far8Δ* strains. Cells were grown in RPMI media for two days and their cell

diameters were quantified (Figure 8D). Both *far8Δ-1* and *far8Δ-2* strains produced significantly

larger cells than the WT haploid and diploid control cells, with diameters ranging from around 5

to 23 μm. Subsequent FACS analysis showed that *far8Δ* cells from RPMI were still primarily

diploid (peaks at 2C and 4C) and did not become polyploid, which is typical of titan cells (Figure

S8). Therefore, we can conclude that deletion of *FAR8* leads to an abundant production of

diploid titan cells in nutrient-limiting conditions.

The virulence of *pph22Δ* and *far8Δ* mutants was tested in a murine infection model. As

expected, given their temperature-sensitive growth defects, *pph22Δ, pph22Δ sup,* and *far8Δ*

mutants could not cause disease upon intranasal instillation in mice before the experiment was

terminated after 45 days (Figure 8E). Four mice from each cohort were randomly selected and

sacrificed on day 14 for fungal burden analysis in the brain and lungs (Figure 8F). The *pph22Δ*

and *far8Δ* mutants could not persist in the lungs or disseminate to the central nervous system.

However, cells from the *pph22Δ #1 sup1* strain were able to persist in the lungs and

disseminate to the brain in one out of four infected mice, although at a significantly reduced

level. These results indicate that *PPH22* and *FAR8* are required for the virulence of *C.*

neoformans.

Deletion of STRIPAK complex subunit *MOB3* leads to hypervirulence

Phenotypic analyses demonstrated that the *C. neoformans* STRIPAK complex is crucial

for sexual development and multiple aspects of vegetative growth. The *pph22Δ* and *far8Δ*

mutants exhibited dramatic phenotypes under different stress conditions. It was hypothesized

that *mob3Δ* mutant strains might present similar phenotypes. However, *mob3Δ* mutants did not

exhibit any growth defects compared to the wild-type strain on YPD media at 30°C or 37°C, on

YNB, or in the presence of rapamycin, CFW, caffeine, FK506, or CsA (Figures 9A and S9). At a

higher temperature of 39°C, the *mob3Δ* mutants grew significantly better than the wild type.

Additionally, *mob3Δ* mutant strains also grew slightly better than the wild type at 37°C in the

presence of 5% CO_2.

Next, we investigated virulence factor production in the *mob3Δ* mutants. When

examining the capsule, we noted an increase in capsule size in *mob3Δ* compared to wild type

(Figure 9B). Quantification of capsule thickness from wild-type and *mob3Δ* cells revealed that

mob3Δ produced significantly larger capsules, with some cells having capsules reaching nearly

4.5 μm (Figure 9C). For melanin production, *mob3Δ* strains produced significantly more melanin

than the wild-type strain, which was visible after 24 and 48 hours of incubation on L-DOPA

medium (Figure 9D). Collectively, these results indicate that the deletion of *MOB3* leads to

increased heat and CO_2 tolerance, and enhanced production of capsule and melanin,

suggesting possible alterations in virulence potential.

The virulence of *mob3Δ* mutants was tested in a murine inhalation model of *C.*

neoformans infection. Four-week-old mice were infected intranasally with *mob3Δ #16* strain or

an isogenic KN99α wild-type strain. Animals infected with the *mob3Δ #16* strain exhibited

significantly decreased survival, with several succumbing to death prior to 7 days post-infection

(Figure 9E). The fungal burden of lung and brain tissues was assessed at 14 days post-infection

and the *mob3Δ* mutation led to increased proliferation in the lungs and dissemination into the

brain (Figure 9F).

To confirm that the changes in virulence were indeed due to the *mob3Δ* mutation, the

infection experiment was repeated with two other independent mutant strains, *mob3Δ #1* and

mob3Δ #8. For this experiment, five- to six-week-old mice were used to rule out any influence of

smaller mouse size on disease progression. Fungal burden assays were performed at day 7

post-infection instead of day 14 to better understand how rapidly *mob3Δ* cells disseminate from

the lungs of infected animals to the brain. The survival curves indicated that animals infected

with *mob3Δ #1* and *mob3Δ #8* succumbed to death significantly faster than those infected with

the wild type (Figure 9G). Fungal burden was also measured at higher levels for *mob3Δ* mutants

in the lungs and brain tissue than the control. Notably, CFUs were obtained from the brains of all

four mice in both *mob3Δ #1* and *mob3Δ #8* groups, whereas no CFUs were found in the brains

of animals infected with wild-type KN99α at 7 days post-infection. These findings demonstrate

that the *mob3Δ* mutation significantly enhances the virulence of *C. neoformans.*

Discussion

The STRIPAK signaling complex is highly conserved across eukaryotic species and

regulates numerous developmental processes. In this study, we characterized the organization

of the *C. neoformans* STRIPAK complex and determined its role in genome stability,

development, and virulence. The *C. neoformans* STRIPAK complex is composed of the PP2A

subunits Tpd3, Far8, and Pph22, along with associated proteins Far9, Far11, and Mob3.

Sequence analysis, structural modeling, and protein interaction analysis revealed that *C.*

neoformans STRIPAK closely resembles the human ortholog. AlphaFold multimer structure

prediction showed an interaction between the coiled-coil domains of Far8 and Far9. In humans,

SLMAP (Far9) and the SIKE protein are predicted to form a regulatory STRIPAK subcomplex,

which was not detected in the cryo-EM structure of the STRIPAK core [42]. The variable

localization of fungal Far9-STRIPAK to the ER, nuclear envelope, or mitochondria is associated

with its role in connecting signal transduction pathways [55]. Similarly, the organization and

localization of *C. neoformans* STRIPAK may depend on the membrane association of Far9.

STRIPAK's role in cell cycle control is conserved among eukaryotes, and in humans,

STRIPAK dysfunction is correlated with genome instability and DNA damage [56]. Here, we

demonstrate a specific role for STRIPAK complex subunits Pph22, Far8, and Mob3 in

maintaining genome stability. In a diploid, deletion of one copy of the *PPH22* or *MOB3* genes

leads to both segmental and whole chromosome aneuploidy. The aneuploid chromosomes vary

between mutant strains, based upon differences in the read coverage maps of *PPH22/pph22Δ,*

MOB3/mob3Δ, *pph22Δ*, and *mob3Δ* isolates, with chromosome 13 being most frequently

duplicated. The *far8Δ* mutants were generated in haploid backgrounds but underwent near

whole-genome endoreplication to become primarily diploid, as supported by FACS analysis. Our

data reinforce the conserved role of *C. neoformans* STRIPAK in maintaining genome stability.

The aneuploidy observed in *pph22Δ*, *far8Δ*, and *mob3Δ* mutants complicates phenotypic

analysis; however, analysis of multiple independent mutants with differing chromosomal

aneuploidy strengthens the hypothesis that the phenotypes described may be attributable to

direct effects of *pph22Δ*, *far8Δ*, and *mob3Δ* mutations, rather than indirect effects of aneuploidy.

STRIPAK's involvement in sexual development regulation is well-documented in fungi,

and we uncovered a similar role in *C. neoformans.* Deletion mutations in *PPH22*, *MOB3*, and

FAR8 led to defects in sexual development when mated with the wild type, affecting hyphal

initiation and elongation, basidia formation, and sporulation. Deletion of *PPH22* and *FAR8*

orthologs in the filamentous fungi *S. macrospora* and *N. crassa* affects cell fusion, fruiting body

formation, and septation, leading to obstruction of the sexual life cycle [9, 11, 57]. Filamentous

ascomycetes also contain homologs of Mob3, which is absent in the ascomycetous yeasts like

S. cerevisiae or *S. pombe*, suggesting a more specialized function in the sexual life cycle. In

Cryptococcus, a basidiomycetous yeast, Mob3 appears to play a similar role in sexual

development. Further study of the STRIPAK components in *C. neoformans* will address the

mechanisms underlying STRIPAK regulation of sexual development.

Mutations in the PP2A catalytic subunit lead to severe growth defects or lethality in other

fungal species [25] and *PPH22* has been assumed to be essential in *C. neoformans* [40]. We

concluded that *PPH22* is not essential but *pph22Δ* does lead to highly reduced fitness during

vegetative growth and under various stress conditions, likely due to severely diminished PP2A

activity. However, interestingly, the growth defects observed in *pph22Δ* mutants were not only

abolished by the addition of rapamycin, both *pph22Δ* and *pph22Δ sup* strains exhibited greater

rapamycin resistance than the wild type. These findings demonstrate that inhibition of TORC1

effectively rescues growth defects due to *pph22Δ*, suggesting PP2A and TORC1 counteract

each other's activities. Another notable phenotype of *pph22Δ* mutants is their inability to grow

on YPD with 1 M sorbitol. It is possible that the presence of sorbitol in YPD medium causes

hyperosmotic stress, and *pph22Δ* cells are unable to activate an adequate stress response to

maintain osmotic homeostasis. The High-Osmolarity Glycerol (HOG) pathway, well-

characterized in *S. cerevisiae* and *Cryptococcus,* is essential for cells to adapt to fluctuating

osmotic conditions [58, 59]. *PPH22* may modulate the HOG pathway or coordinate a response

with other stress signaling pathways to maintain cell growth during hyperosmotic stress. Our

data highlight the importance of PP2A in regulating numerous growth processes in *C.*

neoformans.

Another critical protein phosphatase in *C. neoformans* is protein phosphatase 2B,

calcineurin, which is required for growth at high temperatures and virulence [60], and is the

target of the immunosuppressive drugs FK506 and cyclosporine A (CsA) [61]. We found that

deletion of PP2A catalytic and regulatory subunits, *PPH22* and *FAR8,* also led to an inability to

grow at elevated temperatures or in the presence of FK506 or CsA at 30°C, suggesting a

synthetic lethal relationship with calcineurin inhibition. In fungal pathogens, calcineurin also

plays key roles in host temperature tolerance, sexual development, morphological transitions,

cell wall integrity, drug tolerance, and ER stress response [62]. It is possible that PP2A and

calcineurin share some overlapping functions by dephosphorylating common targets, with one

phosphatase partially compensating for the other's loss. However, the fact that calcineurin is

required for growth at high temperatures and that FK506 or CsA only inhibit cell growth at 37°C

suggests that PP2A has distinct functions that are independent of calcineurin.

For *mob3Δ* mutants, apart from their mating defects, we did not observe any other

significant fitness costs due to the deletion mutation, as seen in *pph22Δ* and *far8Δ* strains.

Strikingly, both our *in vitro* and *in vivo* analyses demonstrate that *mob3Δ* mutants are

hypervirulent. Deletion of *MOB3* leads to increased thermotolerance, CO_2 tolerance, melanin

production, and capsule production, all of which are important for survival in the host

environment. These combined effects likely caused decreased survival of mice, elevated

proliferation in the lungs, and increased brain dissemination in murine infection models. A role

for *MOB3* in virulence or pathogenicity has not been described in studies on the STRIPAK

complex in pathogenic ascomycetes, suggesting it may uniquely negatively regulate virulence in

Cryptococcus. Despite increased virulence and pathogenicity due to *mob3Δ* deletion, there are

likely other fitness trade-offs beyond the host environment in nature that could have limited the

occurrence or persistence of mutations in this key gene during *Cryptococcus* evolution. Further

study is needed to explore host-pathogen interactions involving *mob3Δ* mutants.

While *C. neoformans* is primarily a haploid yeast species, relatively stable diploid

isolates have been described and used in genetic studies to elucidate gene functions [63-66].

One limitation of previous studies analyzing heterozygous diploid gene deletion mutants by

sporulation and dissection is the nature of the parental diploid isolates. In most cases, these

diploids have been selected by fusing complementary auxotrophic parental isolates, resulting in

a prototrophic diploid, but with auxotrophic mutations (such as *ade2* and *ura5*) segregating in

the haploid F1 progeny. Additionally, in some cases, such as the *C. neoformans* diploid strain

AI187, one of the parental isolates resulted from UV irradiation of strain H99α and contains

numerous nucleotide variants throughout the genome, attributable to UV mutagenesis [67, 68].

Thus, analyzing gene function based on the segregation of F1 haploid progeny from a

heterozygous deletion mutant in the AI187 background has limitations due to the segregation

of *ade2*, *ura5*, and additional heterozygous genetic factors. In this study, we capitalized on the

generation of a new diploid strain (CnLC6683), resulting from a spontaneous fusion of two

congenic strains, KN99α and KN99**a**, which has no auxotrophic mutations or mutations

introduced by mutagenesis. This approach provides a robust platform for analyzing essential

genes, as well as genes that are critical for cell growth and challenging to delete in haploid

genetic backgrounds.

The data in this study demonstrate that *C. neoformans* STRIPAK is vital for genome

stability, sexual development, and virulence. Functional characterizations also reveal both

shared and unique roles for STRIPAK components Pph22*, Far8,* and Mob3 in regulating cell

growth, stress responses, and virulence processes. These results are consistent with its

established function as a signaling hub in other organisms. However, the regulators and

downstream effectors of the STRIPAK complex remain unclear. Investigations into the

mechanisms through which STRIPAK can modulate various cellular processes may broaden our

understanding of its comprehensive functions across diverse eukaryotes. Further studies of *C.*

neoformans STRIPAK, including transcriptional and phosphoproteomic analyses, as well as

analyses of the subcellular localization of its components, will provide insights into how

STRIPAK regulates key signaling pathways in this important human fungal pathogen, as well as

in other fungal species and beyond.

Materials and methods

Ethics statement

All animal experiments in this manuscript were approved by the Duke University Institutional Animal Care and Use Committee (IACUC) (protocol #A098-22-05). Animal care and experiments were conducted according to IACUC ethical guidelines.

Strains, media, and growth conditions

C. neoformans strains used in this study are listed in Table S1. Strains were stored as 20% glycerol stocks at -80°C. Fresh cultures were revived and maintained on YPD (1% yeast extract, 2% Bacto Peptone, 2% dextrose) agar medium at 30°C. *Cryptococcus* transformants were selected on YPD medium supplemented with 100 µg/mL nourseothricin (NAT) or 200 µg/mL neomycin (G418). Strains were grown in YPD, synthetic complete (SC) (0.67% yeast nitrogen base with ammonium sulfate and without amino acids, 0.2% amino acid drop-out mix, 2% dextrose), or RPMI 1640 (Sigma-Aldrich R1383, 2% dextrose) liquid medium at either 30°C or 37°C, as indicated. For plate growth assays, strains were cultivated on YPD, Murashige and Skoog (MS) (Sigma-Aldrich M5519), YNB (0.67% yeast nitrogen base with ammonium sulfate and without amino acids, 2% dextrose), Niger seed (7% Niger seed, 0.1% dextrose), or L-3,4-dihydroxyphenylalanine (L-DOPA) (7.6 mM L-asparagine monohydrate, 5.6 mM glucose, 22 mM KH_2PO_4, 1 mM $MgSO_4.7H_2O$, 0.5 mM L-DOPA, 0.3 µM thiamine-HCl, 20 nM biotin, pH 5.6). All plate media were prepared with 2% Bacto agar. To induce copper sufficiency or deficiency, L-DOPA plates were supplemented with 10 µM $CuSO_4$ or 10 µM of the copper chelator bathocuproine disulfonate (BCS). To analyze cell wall-associated phenotypes, sorbitol (1 M), caffeine (0.5 mg/mL), and calcofluor white (3 mg/mL) were added to YPD medium. To analyze cell growth in response to immunosuppressive agents, rapamycin (100 ng/mL), FK506 (1 µg/mL), and cyclosporine A (100 µg/mL) were added to YPD medium. Fluconazole Etest was

633   performed using 0.016-256 µg/mL MIC test strips (Liofilchem 921470). For capsule analysis,

634   strains were incubated for 2 days in RPMI media at 30°C or 37°C, followed by negative staining

635   with India ink. For serial dilution assays, fresh cells were diluted to a starting OD_{600} of 0.1,

636   serially diluted 20-fold, and spotted onto plates for the indicated media and temperature

637   conditions. Plates were incubated for 2 to 7 days and photographed daily.

Generation of marker-free *Cryptococcus neoformans* diploid strain CnLC6683

Wild-type KN99α and KN99**a** cells were mixed in equal numbers and incubated on V8

medium for 24 hours at room temperature in the dark to allow cells to fuse. The cells were then

harvested and plated on YPD to isolate single colonies, with 100-300 colonies per plate.

Colonies from YPD were replica-plated onto filament agar medium (1X YNB without amino acids

or ammonium sulfate, 0.5% glucose, 4% agar) and incubated at room temperature in the dark.

Self-filamentous colonies were isolated, and single colonies were purified on filament agar.

Clones with a stable self-filamentation phenotype on filament agar were recovered on YPD and

maintained as yeast colonies. Colony PCR was performed to confirm the presence of both **a**

and α mating types, and flow cytometry and whole genome sequencing were used to confirm

the ploidy of the resulting diploid, marker-free strain CnLC6683 (Figures 3A and 3D).

Construction of mutant strains

Deletion mutant strains were generated in the *C. neoformans* H99α, YL99**a,** or

KN99**a**/KN99α backgrounds. To generate the deletion alleles, the nourseothricin

acetyltransferase gene expression cassette (NAT) or the neomycin resistance gene expression

cassette (NEO) were amplified from plasmids pAI3 and pJAF12, respectively. Approximately 1

kbp homologous 5′ and 3′ regions of the targeted genes were amplified from H99 genomic

DNA. These homologous arms were assembled with the drug resistance marker with overlap

PCR, as previously described [69, 70]. The *PPH22/pph22Δ* and *MOB3/mob3Δ* heterozygous

mutant diploid strains were generated via CRISPR-Cas9-directed mutagenesis. A codon-

optimized version of *CAS9* for *C. neoformans* was PCR-amplified from plasmid pBHM2403 with

universal primers M13F and M13R [71]. The desired target sequences for the sgRNA constructs

were designed using the Eukaryotic Pathogen CRISPR guide RNA/DNA Design Tool

(EuPaGDT) with default parameters [72]. The selected 20-nt guide sequence was added to the

primers. The *Cryptococcus* U6 promoter, 20-nt guide sequence, scaffold, and 6T terminator

were assembled using single-joint PCR with plasmid pBHM2329 as a template [71]. Gene

deletion, sgRNA, and Cas9 expression constructs were introduced into *Cryptococcus* cells via

electroporation, as previously described [73]. To generate *far8Δ* mutants, we employed

homologous recombination in the *C. neoformans* H99α and YL99**a** backgrounds, using gene

disruption cassettes containing *NAT* or *NEO* markers. Mutants were constructed according to

previously described methods [51, 74]. For the *far8Δ::FAR8-NEO* complemented strain, full-

length *FAR8* was amplified from H99α genomic DNA and cloned into a pNEO plasmid via the

Gibson assembly method. After confirming integration of the *FAR8* gene into the plasmid

through sequencing analysis, the plasmid was linearized with enzymatic digestion (BglII) and

targeted reintegration of *FAR8-NEO* at the native locus was performed via biolistic

transformation. Stable transformants from the YPD+NAT or YPD+NEO selection plates were

screened by diagnostic PCR to confirm cassette integration at the endogenous locus. Positive

transformants from diagnostic PCR were further confirmed via Illumina whole genome

sequencing, which demonstrated the absence of reads mapping to the open reading frame in

the deletion mutants, reads mapping to both the open reading frame and the deletion allele in

the heterozygous diploid mutants, and integration of the *FAR8-NEO* construct at the

endogenous locus in the *far8Δ::FAR8-NEO* complemented strain. Primers used in this study are

shown in Table S2.

Yeast two-hybrid assay

DNA segments encoding Pph22, Tpd3, Far8, Far9, Far11, and Mob3 were amplified by PCR from *C. neoformans* H99 cDNA and cloned into the BamHI and EcoRI restriction sites of pGADT7, a Gal4 transcriptional activation domain vector, and pGBKT7, a Gal4 DNA-binding domain vector (Takara Bio). Yeast two-hybrid strains Y187 (Clontech Laboratories) and Y2HGold (Takara Bio) were transformed with plasmids expressing GAD and GBD constructs, respectively, via the high-efficiency method [75]. The transformed strains were crossed to generate diploid cells co-expressing GAD and GBD fusion constructs. For the analysis of *ADE2* and *HIS3* reporter gene expression under the control of the Gal4-dependent promoter, transformants were grown on synthetic dextrose (SD) dropout medium (0.67% yeast nitrogen base, 2% dextrose) minus leucine, tryptophan, or histidine. On media with histidine (to select for Gal4-dependent expression of reporter gene *ADE2)* Ade+ strains were scored as white and Ade$^-$ strains as red or pink. Amino acids and uracil were added at standard concentrations to support auxotrophic growth requirements.

Flow cytometry analysis

Fluorescence-activated cell sorting (FACS) analysis to determine *Cryptococcus* ploidy was performed as previously described [76], with some modifications. Wild-type and mutant strains were grown on YPD medium at 30°C overnight, harvested, and washed with PBS. The cells were then fixed in 70% ethanol for 16 hours at 4°C. Fixed cells were pelleted and washed with 1 mL NS buffer (10 mM Tris-HCl, 0.25 M sucrose, 1 mM EDTA, 1 mM $MgCl_2$, 0.1 mM $ZnCl_2$, 0.4 mM phenylmethylsulfonyl fluoride, and 7 mM β-mercaptoethanol). After centrifugation, the cells were treated with RNase (0.5 mg/mL) and stained with propidium iodide (10 μg/mL) in a 200 μL suspension of NS buffer for 2 hours in the dark. Then, 50 μL of the stained cells were diluted into 2 mL of 50 mM Tris-HCl, pH=8.0, and submitted to the Duke Cancer Institute Flow Cytometry Shared Resource for analysis. Fluorescence was measured using a BD FACSCanto

707   flow cytometer and analyzed with BD FACSDiva software. Approximately 15,000 events were

708   analyzed for each sample.

Scanning electron microscopy (SEM) and microscopic quantification

For sample preparation for SEM from self-filamenting diploid strains, an agar slice of the

plated cells was fixed in a solution of 4% formaldehyde and 4% glutaraldehyde for 16 hours at

4°C. The fixed cells were then gradually dehydrated in a graded ethanol series (30%, 50%,

70%, and 95%), with a one-hour incubation at 4°C for each concentration. This was followed by

three washes with 100% ethanol, each for 1 hour at room temperature. The samples were

further dehydrated using a Ladd CPD3 Critical Point Dryer and coated with a layer of gold using

a Denton Desk V Sputter Coater (Denton Vacuum, USA). Hyphae, basidia, and basidiospores

were observed with a scanning electron microscope with an EDS detector (Apreo S,

ThermoFisher, USA).

Brightfield and differential interference contrast (DIC) microscopy images were

visualized with an AxioScop 2 fluorescence microscope and captured with an AxioCam MRm

digital camera (Zeiss, Germany). Consistent exposure times were used for all images analyzed.

Cell body sizes were measured using the measurement tool in ImageJ/Fiji. The thickness of

polysaccharide capsules were calculated using the Quantitative Capture Analysis program,

which uses the exclusion zone generated by India ink to differentiate the capsule from the cell

body of individual cells [77]. Statistical differences in cell body diameter between groups were

determined by one-way ANOVA with Dunnett's multiple comparisons test, and the statistical

difference in capsule thickness was determined using an unpaired *t*-test.

Self-filamentation and mating analysis

To monitor self-filamentation efficiency, the wild-type diploid strain CnLC6683

(KN99**a**/KN99α) and the indicated heterozygous mutant diploid strains were grown overnight in

YPD liquid media. The cultures were diluted to an OD$_{600}$ of 1.0, and 4 µL was spotted onto MS

plates. The plates were incubated at room temperature in the dark and monitored for signs of

filamentation and sporulation for at least 4 weeks. For mating analyses, strains were grown

overnight in YPD liquid media. *MAT**a*** cells were mixed with *MAT*α cells in equal amounts and

spotted onto MS plates. For crosses involving *pph22*Δ and *far8*Δ strains, mutant and wild-type

cells were spotted in a 10:1 ratio to compensate for the growth defects of *pph22*Δ and *far8*Δ on

MS media. First, *pph22*Δ and *far8*Δ mutant cells grown overnight in YPD liquid media were

spotted onto MS plates and allowed to pre-grow for two days without the presence of the wild-

type partner. After this pre-incubation, wild-type cells of the opposite mating type were spotted

on top of the mutant cells. A H99α x KN99**a** cross served as a wild-type control on each mating

plate. Mating efficiencies between groups were compared based on data obtained from at least

4 biological replicates.

Whole-genome sequencing, ploidy, and SNP analysis

Genomic DNA for whole-genome sequencing was extracted from saturated 4 mL YPD

cultures with the MasterPure Yeast DNA Purification Kit (LGC Biosearch Technologies,

MPY80200). The precipitated DNA was dissolved in 35 µL of 1x TE buffer (100 mM Tris-HCl, 10

mM EDTA, pH 8.0), and the concentration was estimated using Qubit. Illumina sequencing was

performed at the Duke Sequencing and Genomic Technologies core facility

(https://genome.duke.edu) with Novaseq 6000, providing 250 bp paired-end reads. The Illumina

sequences were mapped to the H99α genome assembly using Geneious software. The

resulting BAM files were converted to TDF format, and read coverage was visualized in IGV to

estimate ploidy for each chromosome. To assess significant changes in read coverage for each

chromosome within a sample, regions of the genome were colored based upon a *Z*-score >1.96

or <-1.96 and a *P*-value <0.05. *Z*-scores between -1.96 and 1.96 with a *P*-value >0.05 were

considered statistically insignificant. For SNP calling, the Illumina sequences were mapped to

the H99α genome assembly using the Geneious default mapper in five iterations. Variant calling

was performed using mapped read files, with parameters set to a 0.9 variant frequency and a

minimum of 100x coverage per variant. Illumina sequences from H99α and KN99**a**/KN99α

(CnLC6683) served as controls for SNP calling analysis.

Competition assay

KN99**a**/KN99α and *PPH22/pph22Δ* diploid strains were cultured overnight at 30°C in

liquid YPD or YPD+NAT, respectively. Cells were adjusted to equal densities using OD_{600}

measurements and mixed in equal numbers in a 4 mL YPD co-culture. The cell density at the

onset of competition was confirmed by plating the mixed cell dilution on YPD+NAT to select for

PPH22/pph22Δ mutants and on YPD to determine the total cell count. This plating process was

repeated at 24 and 48 hours to calculate the cell density of each strain in the co-culture. The

data presented are based on four biological replicates, each with three technical replicates.

Murine infection model

C. neoformans inoculum was prepared by culturing cells in 5 mL YPD on a tissue culture

roller drum at 30°C for approximately 16 hours. Cells were collected by centrifugation, washed

twice with sterile phosphate-buffered saline (PBS), and the cell density was determined with a

hemocytometer. The final cell concentration was adjusted to 4×10^6/mL in PBS. Four- to five-

week-old A/J mice (Jackson Laboratory, USA) were utilized for the murine intranasal infection

model (n=14 for each group, 7 male and 7 female). Mice were anesthetized with isoflurane and

infected by intranasal instillation of 25 μL inoculum (10^5 cells). Mice survival was monitored daily,

and euthanasia was performed via CO_2 exposure upon reaching humane endpoints, including

greater than 20% weight loss, reduced grooming and mobility, or a hunched appearance. For

fungal burden analysis, four mice (2 male, 2 female) from each group were randomly selected

and euthanized via CO_2 exposure at 7 or 14 days post-infection. The brain and lungs were

dissected and homogenized in 1 mL sterile PBS using bead-beating. Organ homogenates were

plated onto YPD agar containing antibiotics (100 µg/mL ampicillin, 30 µg/mL chloramphenicol) to

isolate fungal colonies. Survival data were plotted using Kaplan-Meier curves and statistically

analyzed through log-rank (Mantel-Cox) test. Statistical analyses of fungal burdens were

performed using either Mann-Whitney U test or one-way ANOVA with Dunnett's multiple

comparisons test. Data plotting and analysis of mouse survival and fungal burden was

performed with GraphPad Prism v 10.2.3.

Acknowledgements

PPP is supported by NIH/NIAID T32 grant AI052080-20 as a Tri-I MMPTP fellow. This

work is also supported by NIH/NIAID R01 grants AI039115-27, AI050113-20, and AI172451-02

awarded to JH. LEC is supported by the Canadian Institutes of Health Research (CIHR)

Foundation grant (FDN-154288) and is a Canada Research Chair (Tier 1) in Microbial

Genomics & Infectious Disease. JH and LEC are co-directors of the CIFAR Fungal Kingdom:

Threats & Opportunities program. This work is also supported by National Research Foundation

funded by the Korean government (MSIT) (2021R1A2B5B03086596, 2021M3A9I4021434, and

2018R1A5A1025077 to YSB). We thank Dr. Vikas Yadav, Dr. Maria Isabel Navarro-Mendoza,

and Dr. Zhengchang Liu for their critical reading of this manuscript. We also thank laboratory

manager Anna Floyd Averette for constant support, Dr. Corinna Probst for providing the *cbi1Δ*

ctr4Δ strain, and Dr. Erica Washington for guidance on protein structure 3D modeling. We are

grateful to Dr. Ulrich Kück for his advice and expertise. We commend the Duke Sequencing and

Genomic Technologies Core Facility and the core's director Dr. Devi Swain Lenz for their

assistance.

References

1.   Janssens V, Goris J. Protein phosphatase 2A: a highly regulated family of serine/threonine phosphatases implicated in cell growth and signalling. Biochem J. 2001;353(Pt 3):417-39. doi: 10.1042/0264-6021:3530417. PubMed PMID: 11171037.

2.   Gordon J, Hwang J, Carrier KJ, Jones CA, Kern QL, Moreno CS, et al. Protein phosphatase 2A (PP2A) binds within the oligomerization domain of striatin and regulates the phosphorylation and activation of the mammalian Ste20-Like kinase Mst3. BMC biochemistry. 2011;12:54. doi: 10.1186/1471-2091-12-54. PubMed PMID: 21985334.

3.   Goudreault M, D'Ambrosio LM, Kean MJ, Mullin MJ, Larsen BG, Sanchez A, et al. A PP2A phosphatase high density interaction network identifies a novel striatin-interacting phosphatase and kinase complex linked to the cerebral cavernous malformation 3 (CCM3) protein. Mol Cell Proteomics. 2009;8(1):157-71. doi: 10.1074/mcp.M800266-MCP200. PubMed PMID: 18782753.

4.   Kean MJ, Ceccarelli DF, Goudreault M, Sanches M, Tate S, Larsen B, et al. Structure-Function Analysis of Core STRIPAK Proteins: A Signaling Complex Implicated In Golgi Polarization. J Biol Chem. 2011;286(28):25065-75. doi: 10.1074/jbc.M110.214486. PubMed PMID: 21561862.

5.   Ribeiro PS, Josue F, Wepf A, Wehr MC, Rinner O, Kelly G, et al. Combined functional genomic and proteomic approaches identify a PP2A complex as a negative regulator of Hippo signaling. Mol Cell. 2010;39(4):521-34. doi: 10.1016/j.molcel.2010.08.002. PubMed PMID: 20797625.

6.   Bernhards Y, Poggeler S. The phocein homologue SmMOB3 is essential for vegetative cell fusion and sexual development in the filamentous ascomycete *Sordaria macrospora*. Curr Genet. 2011;57(2):133-49. doi: 10.1007/s00294-010-0333-z. PubMed PMID: 21229248.

7.   Elramli N, Karahoda B, Sarikaya-Bayram O, Frawley D, Ulas M, Oakley CE, et al. Assembly of a heptameric STRIPAK complex is required for coordination of light-dependent multicellular fungal development with secondary metabolism in *Aspergillus nidulans*. PLoS Genet. 2019;15(3):e1008053. doi: 10.1371/journal.pgen.1008053. PubMed PMID: 30883543.

8.   Dettmann A, Heilig Y, Ludwig S, Schmitt K, Illgen J, Fleissner A, et al. HAM-2 and HAM-3 are central for the assembly of the *Neurospora* STRIPAK complex at the nuclear envelope and regulate nuclear accumulation of the MAP kinase MAK-1 in a MAK-2-dependent manner. Mol Microbiol. 2013;90(4):796-812. doi: 10.1111/mmi.12399. PubMed PMID: 24028079.

9.   Bloemendal S, Bernhards Y, Bartho K, Dettmann A, Voigt O, Teichert I, et al. A homologue of the human STRIPAK complex controls sexual development in fungi. Mol Microbiol. 2012;84(2):310-23. doi: 10.1111/j.1365-2958.2012.08024.x. PubMed PMID: 22375702.

10.   Radchenko D, Teichert I, Poggeler S, Kuck U. A Hippo Pathway-Related GCK Controls Both Sexual and Vegetative Developmental Processes in the Fungus *Sordaria macrospora*. Genetics. 2018;210(1):137-53. doi: 10.1534/genetics.118.301261. PubMed PMID: 30012560.

11.   Beier A, Teichert I, Krisp C, Wolters DA, Kuck U. Catalytic Subunit 1 of Protein Phosphatase 2A Is a Subunit of the STRIPAK Complex and Governs Fungal Sexual Development. mBio. 2016;7(3). doi: 10.1128/mBio.00870-16. PubMed PMID: 27329756.

12.   Zhang H, Mukherjee M, Kim JE, Yu W, Shim WB. Fsr1, a striatin homologue, forms an endomembrane-associated complex that regulates virulence in the maize pathogen *Fusarium verticillioides*. Mol Plant Pathol. 2018;19(4):812-26. doi: 10.1111/mpp.12562. PubMed PMID: 28467007.

13.   Islam KT, Bond JP, Fakhoury AM. FvSTR1, a striatin orthologue in *Fusarium virguliforme*, is required for asexual development and virulence. Appl Microbiol Biotechnol. 2017;101(16):6431-45. doi: 10.1007/s00253-017-8387-1. PubMed PMID: 28643182.

14.   Pracheil T, Liu Z. Tiered assembly of the yeast Far3-7-8-9-10-11 complex at the endoplasmic reticulum. J Biol Chem. 2013;288(23):16986-97. doi: 10.1074/jbc.M113.451674. PubMed PMID: 23625923.

15.   Nordzieke S, Zobel T, Franzel B, Wolters DA, Kuck U, Teichert I. A fungal sarcolemmal membrane-associated protein (SLMAP) homolog plays a fundamental role in development and localizes to the nuclear envelope, endoplasmic reticulum, and mitochondria. Eukaryot Cell. 2015;14(4):345-58. doi: 10.1128/EC.00241-14. PubMed PMID: 25527523.

16. E JR, Nordzieke S, Valerius O, Braus GH, Poggeler S. A novel STRIPAK complex component
mediates hyphal fusion and fruiting-body development in filamentous fungi. Mol Microbiol.
2018;110(4):513-32. doi: 10.1111/mmi.14106. PubMed PMID: 30107058.

17. Frey S, Reschka EJ, Poggeler S. Germinal Center Kinases SmKIN3 and SmKIN24 Are
Associated with the *Sordaria macrospora* Striatin-Interacting Phosphatase and Kinase (STRIPAK)
Complex. PLoS One. 2015;10(9):e0139163. doi: 10.1371/journal.pone.0139163. PubMed PMID:
26418262.

18. Wang CL, Shim WB, Shaw BD. Aspergillus nidulans striatin (StrA) mediates sexual development
and localizes to the endoplasmic reticulum. Fungal Genet Biol. 2010;47(10):789-99. doi:
10.1016/j.fgb.2010.06.007. PubMed PMID: 20601045.

19. Green KA, Becker Y, Fitzsimons HL, Scott B. An *Epichloe festucae* homologue of MOB3, a
component of the STRIPAK complex, is required for the establishment of a mutualistic symbiotic
interaction with *Lolium perenne*. Mol Plant Pathol. 2016;17(9):1480-92. doi: 10.1111/mpp.12443.
PubMed PMID: 27277141.

20. Gautier V, Tong LCH, Nguyen TS, Debuchy R, Silar P. PaPro1 and IDC4, Two Genes Controlling
Stationary Phase, Sexual Development and Cell Degeneration in *Podospora anserina*. J Fungi
(Basel). 2018;4(3). doi: 10.3390/jof4030085. PubMed PMID: 29997371.

21. Wang CL, Shim WB, Shaw BD. The *Colletotrichum graminicola* striatin orthologue Str1 is
necessary for anastomosis and is a virulence factor. Mol Plant Pathol. 2016;17(6):931-42. doi:
10.1111/mpp.12339. PubMed PMID: 26576029.

22. Frost A, Elgort MG, Brandman O, Ives C, Collins SR, Miller-Vedam L, et al. Functional
repurposing revealed by comparing *S. pombe* and *S. cerevisiae* genetic interactions. Cell.
2012;149(6):1339-52. doi: 10.1016/j.cell.2012.04.028. PubMed PMID: 22682253.

23. Couzens AL, Knight JD, Kean MJ, Teo G, Weiss A, Dunham WH, et al. Protein interaction
network of the mammalian Hippo pathway reveals mechanisms of kinase-phosphatase
interactions. Sci Signal. 2013;6(302):rs15. doi: 10.1126/scisignal.2004712. PubMed PMID:
24255178.

24. Bae SJ, Ni L, Osinski A, Tomchick DR, Brautigam CA, Luo X. SAV1 promotes Hippo kinase
activation through antagonizing the PP2A phosphatase STRIPAK. Elife. 2017;6. doi:
10.7554/eLife.30278. PubMed PMID: 29063833.

25. Kuck U, Stein V. STRIPAK, a Key Regulator of Fungal Development, Operates as a
Multifunctional Signaling Hub. J Fungi (Basel). 2021;7(6). doi: 10.3390/jof7060443. PubMed
PMID: 34206073.

26. Kuck U, Radchenko D, Teichert I. STRIPAK, a highly conserved signaling complex, controls
multiple eukaryotic cellular and developmental processes and is linked with human diseases. Biol
Chem. 2019. doi: 10.1515/hsz-2019-0173. PubMed PMID: 31042639.

27. Pracheil T, Thornton J, Liu Z. TORC2 signaling is antagonized by protein phosphatase 2A and the
Far complex in *Saccharomyces cerevisiae*. Genetics. 2012;190(4):1325-39. doi:
10.1534/genetics.111.138305. PubMed PMID: 22298706.

28. Kemp HA, Sprague GF, Jr. Far3 and five interacting proteins prevent premature recovery from
pheromone arrest in the budding yeast *Saccharomyces cerevisiae*. Mol Cell Biol.
2003;23(5):1750-63. doi: 10.1128/MCB.23.5.1750-1763.2003. PubMed PMID: 12588993.

29. Innokentev A, Furukawa K, Fukuda T, Saigusa T, Inoue K, Yamashita SI, et al. Association and
dissociation between the mitochondrial Far complex and Atg32 regulate mitophagy. Elife. 2020;9.
doi: 10.7554/eLife.63694. PubMed PMID: 33317697.

30. Furukawa K, Innokentev A, Kanki T. Mitophagy regulation mediated by the Far complex in yeast.
Autophagy. 2021;17(4):1042-3. doi: 10.1080/15548627.2021.1885184. PubMed PMID:
33530805.

31. Chen A, Liu N, Xu C, Wu S, Liu C, Qi H, et al. The STRIPAK complex orchestrates cell wall
integrity signalling to govern the fungal development and virulence of *Fusarium graminearum*. Mol
Plant Pathol. 2023;24(9):1139-53. doi: 10.1111/mpp.13359. PubMed PMID: 37278525.

32. Yamamura Y, Shim WB. The coiled-coil protein-binding motif in *Fusarium verticillioides* Fsr1 is
essential for maize stalk rot virulence. Microbiology (Reading). 2008;154(Pt 6):1637-45. doi:
10.1099/mic.0.2008/016782-0. PubMed PMID: 18524918.

33.   Du Y, Shi Y, Yang J, Chen X, Xue M, Zhou W, et al. A serine/threonine-protein phosphatase PP2A
909   catalytic subunit is essential for asexual development and plant infection in *Magnaporthe oryzae*.
910   Curr Genet. 2013;59(1-2):33-41. doi: 10.1007/s00294-012-0385-3. PubMed PMID: 23269362.

911   34.   Rajasingham R, Govender NP, Jordan A, Loyse A, Shroufi A, Denning DW, et al. The global
912   burden of HIV-associated cryptococcal infection in adults in 2020: a modelling analysis. Lancet
913   Infect Dis. 2022;22(12):1748-55. doi: 10.1016/S1473-3099(22)00499-6. PubMed PMID:
914   36049486.

915   35.   Bongomin F, Gago S, Oladele RO, Denning DW. Global and Multi-National Prevalence of Fungal
916   Diseases-Estimate Precision. J Fungi (Basel). 2017;3(4). doi: 10.3390/jof3040057. PubMed
917   PMID: 29371573.

918   36.   Zhao Y, Lin J, Fan Y, Lin X. Life Cycle of *Cryptococcus neoformans*. Annu Rev Microbiol.
919   2019;73:17-42. doi: 10.1146/annurev-micro-020518-120210. PubMed PMID: 31082304.

920   37.   Casadevall A, Coelho C, Cordero RJB, Dragotakes Q, Jung E, Vij R, et al. The capsule of
921   *Cryptococcus neoformans*. Virulence. 2019;10(1):822-31. doi: 10.1080/21505594.2018.1431087.
922   PubMed PMID: 29436899.

923   38.   Garcia-Solache MA, Casadevall A. Global warming will bring new fungal diseases for mammals.
924   mBio. 2010;1(1). doi: 10.1128/mBio.00061-10. PubMed PMID: 20689745.

925   39.   Lee D, Jang EH, Lee M, Kim SW, Lee Y, Lee KT, et al. Unraveling Melanin Biosynthesis and
926   Signaling Networks in *Cryptococcus neoformans*. mBio. 2019;10(5). doi: 10.1128/mBio.02267-19.
927   PubMed PMID: 31575776.

928   40.   Jin JH, Lee KT, Hong J, Lee D, Jang EH, Kim JY, et al. Genome-wide functional analysis of
929   phosphatases in the pathogenic fungus *Cryptococcus neoformans*. Nat Commun.
930   2020;11(1):4212. doi: 10.1038/s41467-020-18028-0. PubMed PMID: 32839469.

931   41.   Chen C, Shi Z, Zhang W, Chen M, He F, Zhang Z, et al. Striatins contain a noncanonical coiled
932   coil that binds protein phosphatase 2A A subunit to form a 2:2 heterotetrameric core of striatin-
933   interacting phosphatase and kinase (STRIPAK) complex. J Biol Chem. 2014;289(14):9651-61.
934   doi: 10.1074/jbc.M113.529297. PubMed PMID: 24550388.

935   42.   Jeong BC, Bae SJ, Ni L, Zhang X, Bai XC, Luo X. Cryo-EM structure of the Hippo signaling
936   integrator human STRIPAK. Nature structural & molecular biology. 2021;28(3):290-9. doi:
937   10.1038/s41594-021-00564-y. PubMed PMID: 33633399.

938   43.   Byers JT, Guzzo RM, Salih M, Tuana BS. Hydrophobic profiles of the tail anchors in SLMAP
939   dictate subcellular targeting. BMC Cell Biol. 2009;10:48. doi: 10.1186/1471-2121-10-48. PubMed
940   PMID: 19538755.

941   44.   Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, et al. UCSF ChimeraX:
942   Meeting modern challenges in visualization and analysis. Protein Sci. 2018;27(1):14-25. doi:
943   10.1002/pro.3235. PubMed PMID: 28710774.

944   45.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate protein
945   structure prediction with AlphaFold. Nature. 2021;596(7873):583-9. doi: 10.1038/s41586-021-
946   03819-2. PubMed PMID: 34265844.

947   46.   van Zyl W, Huang W, Sneddon AA, Stark M, Camier S, Werner M, et al. Inactivation of the protein
948   phosphatase 2A regulatory subunit A results in morphological and transcriptional defects in
949   *Saccharomyces cerevisiae*. Mol Cell Biol. 1992;12(11):4946-59. PubMed PMID: 1328868.

950   47.   Sneddon AA, Cohen PT, Stark MJ. *Saccharomyces cerevisiae* protein phosphatase 2A performs
951   an essential cellular function and is encoded by two genes. Embo J. 1990;9(13):4339-46.
952   PubMed PMID: 2176150.

953   48.   Gerik KJ, Donlin MJ, Soto CE, Banks AM, Banks IR, Maligie MA, et al. Cell wall integrity is
954   dependent on the PKC1 signal transduction pathway in *Cryptococcus neoformans*. Mol Microbiol.
955   2005;58(2):393-408. doi: 10.1111/j.1365-2958.2005.04843.x. PubMed PMID: 16194228.

956   49.   Furukawa K, Fukuda T, Yamashita SI, Saigusa T, Kurihara Y, Yoshida Y, et al. The PP2A-like
957   Protein Phosphatase Ppg1 and the Far Complex Cooperatively Counteract CK2-Mediated
958   Phosphorylation of Atg32 to Inhibit Mitophagy. Cell reports. 2018;23(12):3579-90. doi:
959   10.1016/j.celrep.2018.05.064. PubMed PMID: 29925000.

960   50.   Niphadkar S, Karinje L, Laxman S. The PP2A-like phosphatase Ppg1 mediates assembly of the
961   Far complex to balance gluconeogenic outputs and enables adaptation to glucose depletion.
962   PLoS Genet. 2024;20(3):e1011202. doi: 10.1371/journal.pgen.1011202. PubMed PMID:
963   38452140.

51. Toffaletti DL, Rude TH, Johnston SA, Durack DT, Perfect JR. Gene transfer in *Cryptococcus*
neoformans by use of biolistic delivery of DNA. J Bacteriol. 1993;175(5):1405-11. doi:
10.1128/jb.175.5.1405-1411.1993. PubMed PMID: 8444802.

52. Walton FJ, Idnurm A, Heitman J. Novel gene functions required for melanization of the human
pathogen *Cryptococcus neoformans*. Mol Microbiol. 2005;57(5):1381-96. doi: 10.1111/j.1365-
2958.2005.04779.x. PubMed PMID: 16102007.

53. Okagaki LH, Strain AK, Nielsen JN, Charlier C, Baltes NJ, Chretien F, et al. Cryptococcal cell
morphology affects host cell interactions and pathogenicity. PLoS Pathog. 2010;6(6):e1000953.
doi: 10.1371/journal.ppat.1000953. PubMed PMID: 20585559.

54. Zaragoza O, Garcia-Rodas R, Nosanchuk JD, Cuenca-Estrella M, Rodriguez-Tudela JL,
Casadevall A. Fungal cell gigantism during mammalian infection. PLoS Pathog.
2010;6(6):e1000945. doi: 10.1371/journal.ppat.1000945. PubMed PMID: 20585557.

55. Kuck U, Stein V. STRIPAK, a Key Regulator of Fungal Development, Operates as a
Multifunctional Signaling Hub. J Fungi (Basel). 2021;7(6):443. doi: 10.3390/jof7060443. PubMed
PMID: 34206073.

56. Hwang J, Pallas DC. STRIPAK complexes: structure, biological function, and involvement in
human diseases. Int J Biochem Cell Biol. 2014;47:118-48. doi: 10.1016/j.biocel.2013.11.021.
PubMed PMID: 24333164.

57. Fu C, Iyer P, Herkal A, Abdullah J, Stout A, Free SJ. Identification and characterization of genes
required for cell-to-cell fusion in *Neurospora crassa*. Eukaryot Cell. 2011;10(8):1100-9. doi:
10.1128/EC.05003-11. PubMed PMID: 21666072.

58. Hohmann S. Osmotic stress signaling and osmoadaptation in yeasts. Microbiol Mol Biol Rev.
2002;66(2):300-72. doi: 10.1128/MMBR.66.2.300-372.2002. PubMed PMID: 12040128.

59. Bahn YS, Jung KW. Stress signaling pathways for the pathogenicity of *Cryptococcus*. Eukaryot
Cell. 2013;12(12):1564-77. doi: 10.1128/EC.00218-13. PubMed PMID: 24078305.

60. Odom A, Muir S, Lim E, Toffaletti DL, Perfect J, Heitman J. Calcineurin is required for virulence of
Cryptococcus neoformans. EMBO J. 1997;16(10):2576-89. doi: 10.1093/emboj/16.10.2576.
PubMed PMID: 9184205.

61. Hemenway CS, Heitman J. Calcineurin. Structure, function, and inhibition. Cell Biochem Biophys.
1999;30(1):115-51. doi: 10.1007/BF02737887. PubMed PMID: 10099825.

62. Yadav V, Heitman J. Calcineurin: The Achilles' heel of fungal pathogens. PLoS Pathog.
2023;19(7):e1011445. doi: 10.1371/journal.ppat.1011445. PubMed PMID: 37410706.

63. Sia RA, Lengeler KB, Heitman J. Diploid strains of the pathogenic basidiomycete *Cryptococcus*
neoformans are thermally dimorphic. Fungal Genet Biol. 2000;29(3):153-63. doi:
10.1006/fgbi.2000.1192. PubMed PMID: 10882532.

64. Davidson RC, Blankenship JR, Kraus PR, de Jesus Berrios M, Hull CM, D'Souza C, et al. A PCR-
based strategy to generate integrative targeting alleles with large regions of homology.
Microbiology (Reading). 2002;148(Pt 8):2607-15. doi: 10.1099/00221287-148-8-2607. PubMed
PMID: 12177355.

65. Hull CM, Davidson RC, Heitman J. Cell identity and sexual development in *Cryptococcus*
neoformans are controlled by the mating-type-specific homeodomain protein Sxi1alpha. Genes
Dev. 2002;16(23):3046-60. doi: 10.1101/gad.1041402. PubMed PMID: 12464634.

66. Idnurm A. A tetrad analysis of the basidiomycete fungus *Cryptococcus neoformans*. Genetics.
2010;185(1):153-63. doi: 10.1534/genetics.109.113027. PubMed PMID: 20157004.

67. Perfect JR, Toffaletti DL, Rude TH. The gene encoding phosphoribosylaminoimidazole
carboxylase (*ADE2*) is essential for growth of *Cryptococcus neoformans* in cerebrospinal fluid.
Infect Immun. 1993;61(10):4446-51. doi: 10.1128/iai.61.10.4446-4451.1993. PubMed PMID:
8406836.

68. Sudarshan S, Davidson RC, Heitman J, Alspaugh JA. Molecular analysis of the *Cryptococcus*
neoformans ADE2 gene, a selectable marker for transformation and gene disruption. Fungal
Genet Biol. 1999;27(1):36-48. doi: 10.1006/fgbi.1999.1126. PubMed PMID: 10413613.

69. Lin X, Chacko N, Wang L, Pavuluri Y. Generation of stable mutants and targeted gene deletion
strains in *Cryptococcus neoformans* through electroporation. Med Mycol. 2015;53(3):225-34. doi:
10.1093/mmy/myu083. PubMed PMID: 25541555.

70.   Davidson RC, Cruz MC, Sia RA, Allen B, Alspaugh JA, Heitman J. Gene disruption by biolistic
1019   transformation in serotype D strains of *Cryptococcus neoformans*. Fungal Genet Biol.
1020   2000;29(1):38-48. doi: 10.1006/fgbi.1999.1180. PubMed PMID: 10779398.
1021   71.   Huang MY, Joshi MB, Boucher MJ, Lee S, Loza LC, Gaylord EA, et al. Short homology-directed
1022   repair using optimized Cas9 in the pathogen *Cryptococcus neoformans* enables rapid gene
1023   deletion and tagging. Genetics. 2022;220(1). doi: 10.1093/genetics/iyab180. PubMed PMID:
1024   34791226.
1025   72.   Peng D, Tarleton R. EuPaGDT: a web tool tailored to design CRISPR guide RNAs for eukaryotic
1026   pathogens. Microb Genom. 2015;1(4):e000033. doi: 10.1099/mgen.0.000033. PubMed PMID:
1027   28348817.
1028   73.   Fan Y, Lin X. Multiple Applications of a Transient CRISPR-Cas9 Coupled with Electroporation
1029   (TRACE) System in the *Cryptococcus neoformans* Species Complex. Genetics.
1030   2018;208(4):1357-72. doi: 10.1534/genetics.117.300656. PubMed PMID: 29444806.
1031   74.   Jung KW, Lee KT, So YS, Bahn YS. Genetic Manipulation of *Cryptococcus neoformans*. Curr
1032   Protoc Microbiol. 2018;50(1):e59. doi: 10.1002/cpmc.59. PubMed PMID: 30016567.
1033   75.   Gietz D, St Jean A, Woods RA, Schiestl RH. Improved method for high efficiency transformation
1034   of intact yeast cells. Nucleic Acids Res. 1992;20(6):1425. PubMed PMID: 1561104.
1035   76.   Tanaka R, Taguchi H, Takeo K, Miyaji M, Nishimura K. Determination of ploidy in *Cryptococcus*
1036   *neoformans* by flow cytometry. J Med Vet Mycol. 1996;34(5):299-301. PubMed PMID: 8912162.
1037   77.   Dragotakes Q, Casadevall A. Automated Measurement of Cryptococcal Species Polysaccharide
1038   Capsule and Cell Body. J Vis Exp. 2018;(131). doi: 10.3791/56957. PubMed PMID: 29364243.

1039

 Figure legends

 Figure 1. Conservation analysis of the *C. neoformans* STRIPAK complex. (A) Table listing

 STRIPAK complex subunits used in BLAST analysis to identify protein orthologs in *C.*

 neoformans. (B) Sequence alignment of STRIPAK protein orthologs in *C. neoformans, S.*

 cerevisiae, and *H. sapiens*. Conserved protein domains are labeled in the schematic diagram:

 HEAT (Huntingtin, elongation factor 3, protein phosphatase 2A, TOR1), FHA (forkhead-

 associated domain), TM (transmembrane domain), and Mob1 (monopolar spindle-one-binder

 protein). (C) The *Cryptococcus* STRIPAK complex superimposed on the electron density map

 and ribbon model of the human STRIPAK complex using the program ChimeraX. The human

 complex is shown in grey, and the corresponding *C. neoformans* proteins are labeled in color.

 (D) ColabFold structure prediction of CnSTRIPAK, with dotted lines representing pseudobonds.

 Figure 2. Yeast two-hybrid analysis of STRIPAK complex subunit interactions. (A) *S. cerevisiae*

 cells carrying plasmids encoding the Gal4 DNA-binding domain (GBD) or the Gal4

 transcriptional activation domain (GAD), fused to individual STRIPAK subunits, were crossed.

 The resulting diploid cells were grown on synthetic dextrose medium with or without histidine.

 Interaction between bait and prey proteins activates Gal4, driving the expression of *ADE2* and

 HIS3, making cells less red (+histidine plates) and allowing growth on medium lacking histidine

 (-histidine plates). Results are representative of two independent experiments. (B) Illustration of

 positive protein-protein interactions detected in the yeast two-hybrid assay.

 Figure 3. Genome instability in STRIPAK complex mutants. (A) Read depth analyses from

 whole-genome sequencing of the WT/WT (CnLC6683) parental diploid strain and heterozygous

 mutant diploid strains. The WT/WT genome is euploid, while *PPH22/pph22Δ* and *MOB3/mob3Δ*

strains exhibit multiple instances of segmental and whole chromosome aneuploidy, with dark

blue and orange highlighting regions/chromosomes with increased and decreased read depth,

respectively. (B) Chromosome maps of *pph22Δ* and *mob3Δ* strains show aneuploidy in some,

but not all, haploid progeny. Progeny can also acquire segmental or chromosomal aneuploidies

absent in the parental strain. "P" stands for meiotic progeny from the indicated parent. (C)

Coverage of *far8Δ* mutants generated in the H99 background reveals duplication of most

chromosomes, suggesting a diploid state. Complementation of the deletion allele in the

far8Δ::*FAR8* strain restores euploidy or results in haploidy. Regions of read coverage in the

genome maps in (A-C) are shaded based on $Z > 1.96$, $P < 0.05$ (dark blue); or $Z < -1.96$, $P < 0.05$

(orange). (D) FACS analysis of *pph22Δ* and *far8Δ* strains. Cells were stained with propidium

iodide to measure DNA content via flow cytometry. Peaks represent relative DNA content during

G_1/S and G_2/mitotic phases. H99 and WT/WT (CnLC6683) (KN99**a**/KN99α) served as 1N and

2N controls, respectively. The number sign (#) after a strain name indicates F1 meiotic progeny

dissected from a heterozygous mutant diploid parental strain. Graphs are representative of two

biological replicates.

1079

Figure 4. Deletion of *PPH22* in a diploid background leads to haploinsufficiency. (A) Self-

filamentation of WT/WT (CnLC6683) and *PPH22/pph22Δ* diploid strains on MS medium. Strains

were grown on MS and incubated at room temperature (24°C) in the dark for 3 to 4 weeks

before images were taken. Each spot represents an independent experiment. A representative

image from one out of four independent experiments for each strain is shown. (B) Scanning

electron microscopy (SEM) analysis of basidia and basidiospores from WT/WT and

PPH22/pph22Δ strains. Samples were prepared following incubation on MS media for four

weeks [shown in (A)]. The black arrowhead in image *PPH22/pph22Δ*-4 shows a hyphal filament

being produced from a basidia head. Scale bars represent 5 µm. (C) Differential interference

contrast (DIC) microscopy images of WT/WT and *PPH22/pph22Δ* cells grown to mid-logarithmic

phase in YPD liquid media at 30°C. *PPH22/pph22Δ* cells showed elongated cells, increased cell

size, incomplete budding, and abnormal cell clusters. Cultures were grown in biological

replicates for analysis. Scale bars represent 10 μm. (D) Growth rates of WT/WT and

PPH22/pph22Δ strains grown in YPD liquid culture at 30°C. Cell density was quantified as

cells/mL, and logarithmic growth was modeled using nonlinear regression. (E) Competition

assay of WT/WT versus *PPH22/pph22Δ*. The growth of strains in the coculture is expressed as

a percentage of the total number of cells at the indicated time point. The data correspond to the

mean of four biological replicates ± standard deviation.

1098

Figure 5. *pph22Δ* mutants frequently accumulate suppressor mutations. (A) Representative

image showing the growth of spores dissected from the *PPH22/pph22Δ-4* strain on YPD

medium. Wild-type spores are marked with a square, and *pph22Δ* deletion mutant spores with a

triangle. The image was taken after five days of incubation at 30°C. (B) *pph22Δ* strains exhibit

very slow growth and form tan-colored colonies. After prolonged incubation, large white colonies

appear (indicated by black arrowheads), resulting from suppressor mutations. The image shows

a YPD plate incubated at 30°C for one week. (C) Wild-type (KN99), isogenic *pph22Δ,* and

pph22Δ suppressor (*pph22Δ sup*) strains were serially diluted and spotted onto YPD medium at

24°C, 30°C, and 33°C. Images were taken after 5 days of incubation. (D) *pph22Δ sup* strains

show shared aneuploidy on chromosome 6. The diagram displays read coverage from whole-

genome sequencing, with chromosomal coordinates at the bottom. The region of increased

coverage shared by all 10 *pph22Δ sup* strains is indicated by a red box, which spans 7 genes.

(E) Table listing the 7 genes with a complete ORF included in the ~15 kb boxed region from (D).

The read depth in this region is at least 3 times higher than the average depth of the rest of

chromosome 6. The gene ID numbers, from top to bottom in the table, are CNAG_02240,

CNAG_02239, CNAG_02238, CNAG_02237, CNAG_02236. CNAG_02235, and CNAG_02234.

1115

Figure 6. Deletion mutations in *PPH22, FAR8,* and *MOB3* lead to defects in sexual

differentiation. (A) *PPH22/pph22Δ* and *MOB3/mob3Δ* strains exhibit defects in self-filamentation

compared to WT/WT (CnLC6683). Images were taken after four weeks of incubation on MS

plates at room temperature. Scale bars represent 200 µm. (B) Mating efficiency of STRIPAK

complex mutants. *pph22Δ*, *pph22Δ sup*, *far8Δ*, *mob3Δ*, and *far8Δ::FAR8* cells were co-cultured

with isogenic wild-type cells of the opposite mating type (H99α or KN99**a**) on MS plates. An

H99α x KN99**a** cross served as a control. Black arrows indicate basidia that did not produce

spores in KN99**a** x α *pph22Δ* and KN99**a** x α *far8Δ* crosses. Scale bars represent 100 µm and

50 µm in the middle and bottom panels, respectively.

1125

Figure 7. Phenotypic analyses of *pph22Δ* and *far8Δ* mutants. (A) WT (KN99**a**) and isogenic

pph22Δ and *pph22Δ sup* strains were serially diluted and plated on YPD, YPD+1 M sorbitol,

YPD+100 ng/mL rapamycin, YPD+1 µg/mL FK506, and YPD+100 µg/mL cyclosporine A (CsA)

at 30°C, as well as YPD at 37°C. Images were taken between 2 and 5 days of incubation. This

set is from the same experiment as Figure 5C. (B) WT (H99α) and isogenic *far8Δ-1, far8Δ-2,*

and *far8Δ::FAR8*, along with WT (YL99**a**) and isogenic *far8Δ-3* and *far8Δ-4* strains, were serially

diluted and plated on the indicated media and temperature conditions. (C) WT, *pph22Δ*, *pph22Δ*

sup, far8Δ, and *far8Δ::FAR8* cells were serially diluted and spotted onto YPD supplemented with

3 mg/mL calcofluor white (CFW) or 0.5 mg/mL caffeine. Plates were incubated at 30°C and

images were taken after 3 days. (D) DIC microscopy images depicting cell morphology of WT

(H99), *pph22Δ,* and *far8Δ* strains grown to mid-logarithmic phase in synthetic complete (SC)

media at 30°C. Scale bars represent 10 µm. (E) Fluconazole Etest to analyze drug susceptibility

in WT (H99), *far8Δ*, *pph22Δ,* and *pph22Δ sup* strains. Cells were grown in an overnight culture

in YPD to saturation, and then spread onto YPD plates before adding FLC Etest strips. Plates

were incubated at 30°C and images were taken after 48 hours. MIC values are shown below the

plate images. Drug sensitivities are representative of two biological replicates.

1142

Figure 8. Role of *PPH22* and *FAR8* in production of virulence factors and pathogenicity. (A) WT

(KN99**a**, H99α, or YL99**a**) and isogenic *pph22Δ, pph22Δ sup, far8Δ,* and *far8Δ::FAR8* strains

were serially diluted and plated onto Niger seed agar medium to induce melanin production. The

lac1Δ mutant was included as a negative control. Plates were incubated at 30°C for 7 days in

the top panel and 4 days in the bottom panel. (B) Analysis of copper-dependent melanin

production in *pph22Δ* and *far8Δ* mutants. The indicated strains were grown in a YPD overnight

culture to saturation and spotted onto L-DOPA plates supplemented with either 10 µM of a

copper chelator, bathocuproine disulfonate (BCS)*,* or 10 µM copper(II) sulfate (Cu$_2$SO$_4$). The

cbi1Δ ctr4Δ double mutant, which can only produce melanin in the copper-supplemented

condition, served as a positive control. Plates were incubated at 30°C for 2 days. (C) Analysis of

capsule formation by India ink staining of WT (KN99**a** or H99α), *pph22Δ, pph22Δ sup, far8Δ,*

and *far8Δ::FAR8* cells. Strains were grown for 3 days at 30°C in RPMI media to induce capsule

formation. Cells were harvested, resuspended in PBS, and stained with India ink. The scale bar

for the top panel of DIC images is 10 µm, and the scale bar for the bottom panel of images is 5

µm. (D) Cell size analysis of WT (H99α), WT/WT (CnLC6683) (KN99**a**/KN99α), *far8Δ,* and

far8Δ::FAR8 complemented strains. Cells analyzed were from the same experiment as (C).

Images were analyzed with ImageJ/Fiji. Data are presented as a scatter dot plot with the

indicated mean cell diameter. Statistical significance was calculated using one-way ANOVA with

Dunnett's multiple comparisons test (****, *P* <0.0001; ns, not significant) (WT, n=190; WT/WT,

n=104; *far8Δ-1*, n=151; *far8Δ-2,* n=162; *far8Δ::FAR8,* n=140). (E, F) Virulence of WT (KN99**α**),

pph22Δ, *pph22Δ sup,* *far8Δ,* and *far8Δ::FAR8* cells in a murine model of *C. neoformans*

infection via the intranasal inhalation route. Equal numbers of male and female A/J mice (n=14

per group) were inoculated with 10^5 cells and monitored for 45 days post-infection. (E) Survival

rates of mice infected with the indicated strains. Ten mice per strain were analyzed. Dashed line

indicates the time of fungal burden analysis (****, *P* <0.0001). (F) Brain and lungs were

harvested from four randomly selected mice per group at 14 days post-infection to quantify CFU

per gram of organ tissue (One-way ANOVA; ****, *P* <0.0001; ***, *P* <0.001; **, *P* <0.01; ns, not

significant).

1171

Figure 9. *mob3Δ* mutants are hypervirulent. (A) Thermotolerance of WT (KN99**a**) and isogenic

mob3Δ strains. Cells were serially diluted, spotted onto YPD at 30°C, 37°C, 39°C, and 37°C

with 5% CO_2, and incubated for two days. (B) WT (KN99**a**) and *mob3Δ* cells were grown in

RPMI at 37°C for three days, harvested, washed with PBS, and stained with India ink to analyze

capsule formation. (C) Capsule thickness measurements were made by subtracting the cell

body diameter from the capsule diameter and dividing by 2. Data are presented in a scatter dot

plot with the mean capsule thickness indicated for each strain. Statistical analysis was

performed using the Mann-Whitney U test (****, *P* <0.0001) (WT, n=214; *mob3Δ*, n=222). (D)

Melanization of WT (KN99**a**) and *mob3Δ* strains on L-DOPA medium at 30°C after 24 and 48

hours of incubation. (E-H) Equal numbers of male and female A/J mice were infected

intranasally with 10^5 cells of the indicated WT and isogenic *mob3Δ* mutant strains and analyzed

for survival rate (n=10) and fungal burden (n=4). (E) Survival analysis of WT (KN99α) and

mob3Δ #16 infected mice. Mice were 4 weeks old at the time of infection. The dashed line

indicates the day at which fungal burden analysis was performed (***, *P* <0.001). (F) CFUs per

gram of lung and brain tissue recovered from organs harvested at 14 days post-infection (Mann-

Whitney U test; *, *P*=0.014). (G) Survival rates of 5-week-old mice infected with WT (KN99α),

mob3Δ #1, and *mob3Δ #8* strains. The dashed line represents the day post-infection of fungal

burden analysis (***, *P* <0.001). (H) CFUs per gram of lung and brain tissue recovered from

organs harvested at 7 days post-infection (One-way ANOVA; **, *P* <0.01; *, *P* <0.05).

Figure 1.

A

STRIPAK Subunit	Gene Name
PP2A catalytic subunit	*PPH21/22 (Sc)*
PP2A scaffold subunit	*TPD3 (Sc)*
PP2A B''' subunit	*FAR8 (Sc)* *STRIATIN (Hs)*
Striatin-interacting protein	*FAR11 (Sc)* *STRIP (Hs)*
Tail-anchor domain protein	*FAR9/10 (Sc)* *SLMAP (Hs)*
Striatin-associated protein	*MOB3 (Hs)*
Coiled-coil domain protein	*FAR3/7 (Sc)* *SIKE (Hs)*
GCKIII family kinase	*STK24/25/26 (Hs)*

(Sc) *Saccharomyces cerevisiae*
(Hs) *Homo sapiens*

Figure 2.

Figure 3.

Figure 4.

Figure 5.

A
PPH22/pph22Δ-4 progeny

□ WT △ *pph22Δ*

B

C

	YPD 24°C	YPD 30°C	YPD 33°C
WT			
pph22Δ #1			
pph22Δ #1 sup1			
WT			
pph22Δ #2			
pph22Δ #2 sup2			

D

Chromosome 6

pph22Δ

pph22Δ
pph22Δ #1 sup1
pph22Δ #2 sup2
pph22Δ #5 sup1
pph22Δ #8 sup1
pph22Δ #9 sup1
pph22Δ #9 sup2
pph22Δ #10 sup1
pph22Δ #10 sup3
pph22Δ #11 sup1
pph22Δ #12 sup1

0   724,048   939,752   1,422,463

E

Gene annotation	Gene name
cytoplasmic protein	*VID27*
26S protease regulatory subunit 4	*RPT2*
ubiquitin-conjugating enzyme E2 G2	*UBC7*
cytoskeleton assembly control protein	*SLA2*
type 2a-like phosphatase	***PPG1***
hypothetical protein	CNAG_02235
large subunit ribosomal protein	*RPL6b*

Figure 6.

Figure 7.

Figure 8.

Figure 9.