Engineering synthetic agonists for targeted activation of Notch signaling

David H. Perez[1], Daniel Antfolk[1], Xiomar E. Bustos[1], Elliot Medina[1], Shiun Chang[1], Ahmed A.

Ramadan[1], Paulo C. Rodriguez[1], David Gonzalez-Perez[1], Daniel Abate-Daga[1], Vincent C.

Luca[1*]

[1] Department of Immunology, Moffitt Cancer Center, Tampa, FL 33602, USA

*Corresponding author, email: vince.luca@moffitt.org

10 ABSTRACT

Notch signaling regulates cell fate decisions and has context-dependent tumorigenic or tumor

suppressor functions. Although there are several classes of Notch inhibitors, the mechanical force

requirement for Notch receptor activation has hindered attempts to generate soluble agonists. To

address this problem, we engineered synthetic Notch agonist (SNAG) proteins by tethering

affinity-matured Notch ligands to antibodies or cytokines that internalize their targets. This

bispecific format enables SNAGs to "pull" on mechanosensitive Notch receptors, triggering their

activation in the presence of a desired biomarker. We successfully developed SNAGs targeting

six independent surface markers, including the tumor antigens PDL1, CD19, and HER2, and the

immunostimulatory receptor CD40. SNAGs targeting CD40 increase expansion of central

memory γδ T cells from peripheral blood, highlighting their potential to improve the phenotype

and yield of low-abundance T cell subsets. These insights have broad implications for the

pharmacological activation of mechanoreceptors and will expand our ability to modulate Notch

signaling in biotechnology.

25 INTRODUCTION

The Notch pathway is a cell-to-cell communication system that regulates embryonic development,

tissue homeostasis, and immune cell differentiation. The signaling of mechanosensitive Notch

receptors is tightly regulated[1] and aberrant Notch activity causes several human diseases[2]. For

example, loss-of-function mutations in Notch components are linked to the development of aortic

valve disease (Notch1), Alagille syndrome (Notch2, Jagged1), CADASIL (Notch3), spondylocostal

dysostosis (DLL3)[3–6]. In cancer, Notch functions as a tumor suppressor or oncogene depending

on the cell type, and both loss-of-function and hyperactivating mutations influence tumorigenesis

and disease progression[7]. Notch is also pleiotropic with respect to its guidance of cell fate

decisions, in that Notch activation stimulates either proliferation or differentiation in different stem

cell populations[8]. These diverse functions suggest that Notch agonists and antagonists may each

be viable therapeutics in certain biomedical contexts[9].

The role of Notch in T cell biology has led to the development of several Notch-based strategies

for enhancing cancer immunotherapy. Notch signaling is important for several natural stages of T

cell maturation[10], and ex vivo Notch activation is required for the differentiation of T cells from

hematopoietic stem cells (HSCs)[11]. This latter function may potentially be used to generate

allogeneic T cells for "off-the-shelf" adoptive T cell or chimeric antigen receptor (CAR) T cell

therapies. More recently, Notch activation has been shown to enhance the antitumor function of

fully mature, activated T cells[12]. Genetic overexpression of an activated form of Notch[13], as well

as culturing T cells in the presence of Notch1-specific antibodies[14] or ligand-expressing cells[15],

were each associated with improved tumor clearance in various animal models of cancer. Detailed

analysis of the T cells used in these studies revealed that these phenotypes were due to the

Notch-stimulated induction of exhaustion-resistant or stem-like phenotypes.

At the molecular level, Notch receptors are massive (~290kD) transmembrane proteins that are

activated by a distinctive, mechanical force-driven mechanism[16–18,1]. Notch signaling is initiated

when a Delta-like (DLL) or Jagged (JAG) ligand forms a *trans*-interaction with a Notch receptor

on the surface of an adjacent cell[1,19–21]. Endocytosis of the ligand then generates a "pulling" force

that propagates to the negative regulatory region (NRR) of Notch [16,22]. This pulling destabilizes

the NRR, which exposes internal cleavage sites for processing by the intramembrane proteases

ADAM10 (S2 cleavage) and γ-secretase (S3 cleavage)[23,24]. Following these proteolytic events,

the Notch intracellular domain (NICD) translocates to the nucleus to function as a transcriptional

co-activator[25].

Although Notch inhibitors are widely available, the requirement for mechanical force in Notch

activation has precluded the development of soluble agonists[9,18]. Specifically, these agents are

challenging to engineer because they must somehow "pull" on the Notch receptor despite lacking

a method of force generation. Several strategies have been developed to activate Notch receptors

in vitro through mimicry of the physiological activation process. Notch signaling may be induced

through co-culture of Notch-expressing cells and ligand-expressing cells, by culturing Notch-

expressing cells on plates coated with ligands or antibodies, or by administration of ligand-coated

microbeads[14,26,27]. By contrast, only a single antibody targeting Notch3, A13, has been reported

to function as a soluble agonist[28]. Binding of this antibody promotes the unfolding of metastable

Notch3 NRR domains, which in turn exposes the S2 site for proteolytic cleavage [29]. Unfortunately,

this NRR unfolding approach has been ineffective for receptor subtypes with stable NRRs (e.g.,

Notch1/2/4), and the lack of soluble agonists remains a significant void in our biochemical toolkit

for manipulating the Notch pathway.

In this study, we engineered bispecific proteins that stimulate activation of Notch signaling in

desired cellular contexts. We successfully developed SNAGs that enhance the signaling of weakly

activating JAG1 ligands, as well as those that selectively activate Notch in the presence of the

tumor biomarkers PDL1, CD19, and HER2. Furthermore, we show that SNAGs targeting the

immunostimulatory receptor CD40 improve the expansion of γδ T cells from mixed cell

populations. The modularity and versatility of this SNAG platform provide a blueprint for the

development of a diverse repertoire of Notch-based biologics.

RESULTS

Soluble DLL4 ligand multimers do not activate Notch signaling. As an initial attempt to

generate Notch agonists, we investigated whether soluble oligomers of an affinity-matured DLL4

ligand (Delta^{MAX}) activate Notch signaling[30]. Delta^{MAX} contains ten mutations that increase its

affinity for human Notch receptors by 500- to 1000-fold, making it a more potent activator than

DLL4 in co-culture and plate-bound formats[30]. We hypothesized that this increased affinity,

coupled with receptor crosslinking through multimerization, could introduce tension in the absence

of an endocytic pulling force. To test this hypothesis, we incubated Notch1-Gal4 mCitrine reporter

cells[31] with soluble and immobilized Delta^{MAX} multimers (Fig. 1a-c). Delta^{MAX} dimers were

generated through the C-terminal addition of a dimeric human IgG1 Fc domain (Fig. 1b), and

tetramers were generated by pre-mixing a 4:1 molar ratio of biotinylated Delta^{MAX} with streptavidin

(SA, Fig. 1c). We found that neither the monomers nor the multimers induced reporter activity. By

contrast, the plated Delta^{MAX} ligands potently stimulated Notch1 activation (Fig. 1a-c). This

indicates that the receptor crosslinking by Delta^{MAX}-Fc dimers and Delta^{MAX}-SA tetramers is

insufficient for signaling activation.

Design of synthetic Notch agonists. To develop soluble Notch agonists, we engineered

bispecific proteins that recapitulate the endocytosis-linked activation mechanism of DLL and JAG

ligands (Fig. 1d). SNAGs were created by fusing Delta^{MAX} to the N-terminus of biomarker-targeting

antibody fragments via a flexible (GS)$_5$ linker, or by fusing Delta^{MAX} and antibody fragments to the

N- and C-termini of a dimeric IgG1 Fc domain (Fig. 1e). These design concepts are intended to

form a "molecular bridge" between Notch-expressing cells and cells that express a given surface

protein. Conceptually, SNAGs should then activate Notch if the enforced interactions induce

endocytic or tensile force capable of unfolding the NRR.

SNAGs rescue the signaling of a signaling-deficient DLL4 mutant. To demonstrate proof-of-

concept, we tested whether SNAGs could rescue the activity of a signaling-deficient DLL4 mutant.

Loss-of-function DLL4 cells were generated by expressing a "headless" DLL4 truncation where

the Notch-binding C2 and DSL domains[20,32] were replaced with a BC2 epitope tag (BC2-DLL4[HL])

(Fig. 2a, Fig. S1a)[33]. BC2-SNAGs were then generated by fusing Delta[MAX] to a BC2-specific

nanobody (Figs. 2a-b, Fig. S2a). We found that BC2-DLL4[HL] cells alone did not activate signaling

in a Notch1-Gal4 mCitrine reporter assay, whereas the addition of 1 nM to 100 nM concentrations

of SNAGs stimulated a dose-dependent increase in reporter activity (Fig. 2c, Fig. S3a-c).

Monomeric BC2-SNAGs containing the $(GS)_5$ linker (BC2-SNAG) stimulated a ~6-fold increase

in Notch1 signaling, whereas dimeric BC2-SNAG Fc fusion proteins (BC2-SNAG[Fc]) were more

effective and induced a ~10-fold increase (Fig. 2c). Importantly, administration of the monomeric

or dimeric BC2-SNAGs alone did not substantially increase Notch1 reporter activity, indicating

that a mixture of target-expressing and non-expressing cells is required for SNAG-mediated

activation (Fig. 2c).

SNAGs bolster the activity of weakly-signaling JAG1 ligands. DLL or JAG ligands

preferentially signal through certain Notch receptor subtypes, and JAG1 is a particularly weak

activator of Notch1[34]. Therefore, we tested whether a JAG1-targeting SNAG (JAG1-SNAG[Fc])

could potentiate JAG1-Notch1 signaling. In the JAG1-SNAG[Fc] construct, Delta[MAX] and an scFv

derived from the JAG1-targeting antibody B70[35] were fused to the N- and C-termini of an IgG1 Fc

domain as described above (Fig. S2b). We then performed a signaling assay to measure the

activation of Notch1 reporter cells by JAG1-overexpressing HEK293 cells (JAG1-293 cells) in the

presence or absence of the SNAG. We found that addition of the JAG1-SNAG[Fc] increased Notch1

reporter activity by ~4-fold compared to JAG1-293 cells alone, and that JAG1-293 cells did not

stimulate a significant increase in reporter activity (Fig. 2d, Fig. S1b). We also evaluated the

JAG1-SNAG[Fc] with a JAG1 H268Q "Nodder" mutant that causes Alagille syndrome-like symptoms

in mice by decreasing JAG1-Notch1 binding (Fig. S1c). Addition of the JAG1-SNAG[Fc] to

cocultures of Notch1 and JAG1[H268Q] cells increased JAG1 signaling by up to 7-fold compared to

JAG1[H268Q] cells alone (Fig. 2e). These data indicate that SNAGs can function as "signaling

enhancers" by potentiating the activity of endogenous or mutated ligands.

SNAGs targeting tumor antigens activate Notch in mixed cell populations. We next tested whether SNAGs targeting the tumor antigens PD-L1, CD19, or HER2 (Fig. 3a) can stimulate Notch activation. There is mounting evidence that Notch signaling enhances the function of activated T cells[13–15], and SNAGs localized to the tumor microenvironment have the potential to stimulate localized activation of tumor-associated lymphocytes. For these SNAGs, the targeting arms were derived from antibody-drug conjugates (ADCs) that were pre-selected for their ability to induce target internalization. We hypothesized that SNAGs incorporating ADC antibodies could thus mimic the physiological endocytosis mechanism of DLL or JAG ligands.

We generated monomeric and dimeric PDL1-SNAGs by fusing Delta^{MAX} to a single-chain variable fragment (scFv) derived from the ADC antibody Atezolizumab[36,37]. In the monomeric PDL1-SNAG, Delta^{MAX} and the scFv were connecting using a $(GS)_5$ linker, and in the dimeric PDL1-SNAG (PDL1-SNAG^{Fc}), Delta^{MAX} and the scFv were fused to the N- and C-termini of an IgG1 Fc domain (Fig. 3a, Fig. S2a). Unexpectedly, addition of the monomeric PDL1-SNAG to a 1:1 mixture of Notch1 reporter cells and PDL1-expressing MDA-MB-231 cells did not activate Notch1 (Fig. 3b, Fig. S1d). However, the dimeric PDL1-SNAG^{Fc} protein stimulated a ~7-fold increase in Notch1 signaling in the coculture, suggesting that multimerization or avidity-enhancement may be required for SNAGs to effectively target biomarkers other than Notch ligands (Fig. 3b). Neither the PDL1-SNAG nor the PDL1-SNAG^{Fc} substantially increased Notch1 reporter activity in the absence of MDA-MB-231 cells. Because of the increased efficacy of the dimeric SNAGs (Fig. 2c, Fig. 3b), we designed subsequent CD19- and HER2-SNAGs using only the Fc-fusion format (Fig. 3a).

SNAGs do not activate signaling on cells expressing both Notch1 and PD-L1. Given the ubiquitous expression of Notch1 in mammalian cells, it is conceivable that SNAGs could activate signaling when Notch1 and the target protein are both present on the cell surface. To test this possibility, we cultured MDA-MB-231 cells in the presence of soluble Delta^{MAX}-Fc, PDL1-SNAG^{Fc}, or immobilized Delta^{MAX}-Fc and monitored the levels activated Notch1 by Western Blot (Fig. 3c). We found that the plated Delta^{MAX}-Fc protein stimulated high levels of Notch1 activation, whereas the PDL1-SNAG^{Fc} did not induce signaling over the background levels observed for soluble Delta^{MAX}-Fc alone (Fig. 3c). The inability of SNAGs to activate Notch1 in MDA-MB-231 cells suggests that the present design does not enable sufficient intercellular crosslinking in cultures of cells expressing both Notch1 and the biomarker.

Development of SNAGs targeting CD19 and HER2. To generate a SNAG targeting the B-lymphocyte antigen CD19, we fused an scFv derived from the CD19-targeting ADC loncastuximab[38] to the C-terminus of Delta^MAX-Fc (CD19-SNAG^Fc, Fig. S2a). The CD19-SNAG was then added to Notch1 reporter cells or to co-cultures of Notch1 reporter cells and CD19-overexpressing 3T3 fibroblast cells (Fig. S1e). We found that the CD19-SNAG^Fc protein stimulated up to a 6-fold increase in reporter activity in the co-culture compared to untreated Notch1 cells (Fig. 3d). To generate a SNAG targeting the breast cancer antigen HER2 (HER2-SNAG^Fc), we replaced the CD19-targeting arm with an scFv derived from the HER2-targeting ADC trastuzumab[39] (Fig. S2c). Addition of the HER2-SNAG^Fc to a mixed culture of Notch1 reporter cells and HER2-expressing SK-BR-3 breast cancer cells induced a 6-fold increase in reporter activity (Fig. 3e, Fig. S1f) at the highest concentration tested (100 nM), which is similar to the level of activation we observed for the PDL1-SNAG^Fc and the CD19-SNAG^Fc constructs (Fig. 3b, 3d). In the absence of biomarker-expressing cells, neither the CD19-SNAG^Fc nor the HER2-SNAG^Fc stimulated a significant increase in signaling compared to Delta^MAX-Fc alone (Fig. 3d-e). Our collective development of PD-L1, CD19, and HER2 SNAGs demonstrates that SNAGs can facilitate signaling by engaging cell surface proteins beyond endogenous ligands.

Endocytosis is required for SNAG-mediated Notch activation. Ligand endocytosis is important for Notch activation[40], and this process is regulated by ubiquitination of DLL or JAG ICDs by the E3 ligase Mindbomb1[41–43]. To determine whether endocytosis occurs with a SNAG targeting a surface protein other than a natural Notch ligand, we performed an immunofluorescent endocytosis assay utilizing CD19-SNAG^Fc in CD19-expressing cells. CD19-SNAG^Fc coupled with a fluorescent secondary antibody (Alexafluor 647-labeled anti-Fc) bound strongly to the surface of the CD19-expressing cells when the mixture was incubated on ice (Fig 4a, Fig. S4a-c), and the contours of the cells were identified by staining for filamentous actin (Fig. S4a). Incubating the cells at 37 °C after attaching CD19-SNAG^Fc-647 to cells allowed for cellular functions, including endocytosis, to resume. Visualizing the cells after a 15 min incubation at 37 °C showed that the majority of CD19-SNAG^Fc is internalized (Fig. 4b, Fig. S4e-d).

To test whether endocytosis is necessary for SNAG function, we co-administered SNAGs with the dynamin-dependent endocytosis inhibitor Dynasore. We found that Dynasore completely ablated the activity of CD19-SNAG^Fc in co-cultures of Notch1- and CD19-expressing cells, indicating that endocytosis is required for SNAG-mediated activation utilizing CD19 as a biomarker (Fig. 4c). We

further found that BC2-SNAGs targeting BC2-DLL4[HL] were unable to activate Notch1 in co-
cultures of Notch1 and BC2-DLL4[HL] cells in the presence of Dynasore, confirming that
endocytosis is also required for SNAG-mediated rescue of DLL4 signaling (Fig 4d-e.)
Interestingly, we found that immobilized SNAGs were also unable to activate Notch1 in the
presence of Dynasore, suggesting that endocytosis in the Notch-receptor cell is essential for
Notch activation by plated ligands (Fig. 4f). These studies demonstrate that Notch activation by
plated ligands, SNAGs targeting a DLL4 loss-of-function mutant, and SNAGs targeting tumor
antigens each depend on endocytosis. However, it is currently unclear whether endocytosis of
the receptor, ligand, or both, is essential for SNAG function.

SNAGs targeting CD40L increase the expansion of γδ T cells. We next generated a SNAG
targeting CD40 (CD40-SNAG), an immunostimulatory receptor that undergoes endocytosis upon
binding to CD40 ligand (CD40L)[44]. We hypothesized that a CD40-SNAG could be used to activate
Notch in T cells in the presence of CD40[+] B cells, as these cell types are known to colocalize in
germinal centers, peripheral blood, the tumor microenvironment, and other biological contexts[45].
To generate a CD40-SNAG, we fused Delta[MAX] and the ECD of CD40L to the N- and C-termini of
a trimeric leucine zipper[46] (Fig. 5a, Fig. S2d), respectively. This trimeric scaffold was selected
instead of an Fc domain for the CD40-SNAG because CD40L is naturally a homotrimer. We found
that addition of the CD40-SNAG robustly activated Notch in mixed cultures of Notch1 reporter
cells and CD40-expressing OCI-Ly3 cells (Fig. S1g), but only weakly in Notch1 reporter cells
alone (Fig. 5b). Notably, OCI-Ly3 cells used in this assay are non-adherent, indicating that SNAGs
may facilitate Notch activation between adherent cells and those grown in suspension.

There has been a growing interest in γδ T cell-based cancer immunotherapies, including the
adoptive transfer of unmodified γδ T cells and the development of γδ T CAR T cells[47–49]. It was
previously shown that Notch1 and Notch2 are important for γδ T cell antitumor function[50], and that
Notch stimulation generally induces memory phenotypes in mature and activated T cells[15,14].
Therefore, we sought to determine whether the CD40-SNAG could be used to improve the
expansion and phenotype of γδ T cells isolated from peripheral blood mononuclear cells
(PBMCs). To determine the ratio of CD40-expressing and non-expressing cells, we analyzed
PBMCs by flow cytometry. We found that a representative PBMC culture contained 18.6% CD19+
CD40[+] B cells and 71.9% CD3[+] CD40[-] T cells (Fig. 5c), suggesting that the T cell fraction would
be amenable to stimulation with the CD40-SNAG.

To test the effect of the CD40-SNAG on γδ T cells, we administered increasing amounts of CD40-

SNAG during an established γδ T cell expansion protocol[51]. Addition of the CD40-SNAG

increased both the relative amount and total amount of γδ T cells recovered at all concentrations

tested (Fig. 6a-c). We found that the highest CD40-SNAG concentration (500 ng/mL) was

associated with >4-fold increase in the total number of γδ T cells compared to untreated PBMCs

or those treated with Delta[MAX]-Fc or trimerized CD40L alone. Treatment with 500 ng/mL CD40-

SNAG also induced a ~4-fold increase in the number of central memory cells and a significant

reduction in the number of effector memory γδ T cells (Fig. 6d). This bias towards the central

memory subset is a desirable outcome given that central memory T cells have increased survival

and antitumor function compared with effector memory T cells and effector T cells[52].

DISCUSSION

The development of soluble agonists has been an enduring challenge in the Notch field[9,53]. The

SNAG platform described here offers a solution to this problem and provides a framework for the

development of a diverse array of Notch activating biologics. Such agents have a wide range of

potential translational applications, particularly in cancers where Notch functions as a tumor

suppressor[7], T cell manufacturing[11,27], T cell immunotherapy[13–15], wound healing[54], and other

areas of regenerative medicine. These first-generation SNAGs were engineered using an Fc-

fusion format used in clinically viable protein drugs, which may also help to accelerate *in vivo*

translation.

In their present form, SNAGs facilitate potent activation of Notch signaling in mixed populations

of cells. However, we anticipate that the design may be tuned to further optimize SNAG function.

For example, it may be preferable to engineer SNAGs such that the target binding arm has a

higher binding affinity than the Notch-binding arm to improve specificity and tissue distribution.

Such strategies have been successfully employed both for bispecific inhibitory antibodies[55] and T

cell engagers[56]. Additionally, higher-order oligomers beyond the dimeric and trimeric SNAG

scaffolds tested here may lead to increased signaling potency. Future studies will focus on

optimizing affinity and multimerization to maximize signaling while maintaining favorable

biochemical properties.

One surprising observation was that PDL1-SNAGs did not activate signaling on cells expressing

both PDL1 and Notch1. We speculate that these SNAGs engage the two targets in *cis* on the

surface of a single cell, as opposed to bridging PDL1 and Notch1 proteins between cells, and that

cis interactions do not introduce sufficient tension to unfold the NRR. This may be attributed to
the restricted diffusion of SNAGs in the 2-dimensional environment of the membrane, which can
promote preferential *cis* interactions by increasing the local concentration. Previous studies have
shown that *cis* inhibition of Notch signaling occurs when ligands and receptors are expressed on
the same cell[31,57], and it appears that SNAGs are similarly unable to activate Notch in this context.
Regardless, the ability of SNAGs to mediate unidirectional signaling enables highly selective
targeting, which could minimize the risks of potential toxicity from global Notch agonism.

There has been a longstanding interest in using Notch stimulation to manufacture "off-the-shelf"
CD4[+] or CD8[+] T cells for immunotherapy[58]. By contrast, the field of γδ T cell immunotherapy
emerged more recently and the role of Notch in their biology is less thoroughly understood[47].
These cells represent only 1-10% of the T cell population in healthy donors, and protocols for the
manufacture of pure populations of functional γδ T cells are continuously being optimized[59,60]. We
showed that the addition of a CD40-SNAG can greatly increase both the total number of γδ T
cells and the fraction of cells with desirable central memory phenotypes. This expansion was not
observed upon the addition of trimerized CD40L, suggesting that CD40 signaling alone is not able
to recapitulate the effects of the CD40-SNAG. In future studies, it will be interesting to probe
whether Notch stimulation in γδ T cells is associated with enhanced antitumor function akin to
what is observed for conventional activated T cells[14], or whether there are synergistic effects
between Notch activation in T cells and CD40 activation in B cells.

Although SNAGs are effective in mixed cell populations, the development of "unconditional"
agonists that do not rely on a secondary target remains a challenging problem. Thus far, it appears
that the metastable NRR of Notch3 is uniquely susceptible to antibody-mediated
destabilization[29,61]. The engineering of agonists targeting other Notch receptors with more stable
NRRs may require alternative solutions. The successful activation of Notch1 with ligands
immobilized on beads[27] or DNA origami structures[62] suggests that oligomerization may be an
effective strategy, but these methods are not currently viable for in vivo applications. Despite these
limitations, the development of SNAGs represents a key first step towards the widespread
development of Notch activating molecules for basic and translational research.

AUTHOR CONTRIBUTIONS
V.C.L. and D.H.P. wrote the manuscript. V.C.L., D.H.P, D.A., and D.A.D. designed the
experiments. D.H.P. cloned the SNAG constructs, purified the proteins, and performed the

signaling assays. E.M. and D.G.P. generated the Delta^{MAX} constructs. D.A. performed the Notch

activation assays in MDA-MB-231 cells and immunofluorescent endocytosis assays. X.E.B.

performed T cell expansion assays. P.C.R. and S.C. assisted with immunology experiments.

V.C.L. supervised the project and edited the manuscript.

ACKNOWLEDGEMENTS.

This project was supported by NIH R35GM133482 (V.C.L. and D.A.), the Sigrid Juselius

Foundation (D.A.) and NIH R35 Diversity Supplement R35GM133482-03S2 (E.M.). V.C.L. is a

Rita Allen Scholar. Shared resources were provided by the Moffitt Cancer Center Support Grant

NIH P30CA076292.

COMPETING INTERESTS

V.C.L. is a consultant on unrelated projects for Cellestia Biotech, Remunix, and Curie.Bio. The

remaining authors have no competing interests. V.C.L. and D.H.P. have filed provisional patents

(serial numbers 63/548,615 and 63/663,744) based on the described technology.

 FIGURES AND FIGURE LEGENDS

Figure. 1. Design concept for synthetic Notch agonists. (a) Flow cytometry histogram overlay of Notch1 reporter cells stimulated by soluble or plated (non-specifically adsorbed) Delta^{MAX}. The cartoon depicts the site-specifically biotinylated Delta^{MAX}(N-EGF5) construct. (b) Histogram overlay of Notch1 reporter cells stimulated with soluble or plated Delta^{MAX}-Fc protein. (c) Histogram overlay of Notch1 reporter cells stimulated with plated or soluble Delta^{MAX}-SA tetramers. (d) Cartoon schematic depicting the ECDs of Notch1 and DLL4 interacting during canonical Notch activation. The NRR and ligand-binding domains (LBD) of Notch1 (EGF domains 8-12) are shaded. (e) Schematic of a generalized SNAG construct alongside a cartoon depicting SNAG-mediated Notch activation.

Figure. 2. SNAGs rescue the signaling of binding-deficient DLL4 and JAG1 mutants. (a)
Cartoon schematic depicting a SNAG binding to Notch1 and a loss-of-function DLL4 mutant. The
"headless" loss-of-function DLL4 protein (DLL4^{HL}) was generated by replacing the Notch-binding
C2-DSL region with a BC2 peptide epitope recognized by the anti-BC2 nanobody. (b) Cartoon
schematic depicting the multivalent binding of a dimeric Fc-tagged SNAG (BC2-SNAG^{Fc}) to
Notch1 and "headless" DLL4. (c-e) Fluorescent reporter assays were used to evaluate SNAG-
mediated activation of Notch1 in cocultures with HEK293T cells expressing DLL4^{HL} (c), JAG1 (d),
or JAG1^{H268Q} (e). Increasing concentrations (1 nM, 10 nM, or 100 nM) of the indicated SNAGs
were added to Notch1-Gal4 Citrine reporter cells alone, or to a 1:1 mixture of Notch1 reporter
cells and HEK293 cells expressing DLL4^{HL}, and fluorescence was measured by flow cytometry. A
representative experiment from three biological replicates is shown. Mean fluorescence intensity
(MFI) was normalized to the mean MFI of Notch1 reporter cells alone. Error bars represent the
standard deviation of three technical replicates with the *P* value by Student's *t* test shown above
each comparison.

Figure. 3. SNAGs targeting tumor antigens activate Notch in mixed cell populations. (a) Cartoon schematics of PDL1, CD19, and HER2 SNAGs. (b) PDL1-SNAG-mediated activation of Notch1 was evaluated in a fluorescent reporter assay. Increasing concentrations (1 nM, 10 nM, or 100 nM) of Delta^{MAX}-Fc, PDL1-SNAG, or PDL1-SNAG^{Fc} were added to Notch1-Gal4 mCitrine reporter cells alone, or a 1:1 mixture of Notch1 reporter cells and MDA-MB-231 cells. (c) Activation of Notch1 in MDA-MB-231 cells, which express both PDL1 and Notch1, was assessed by Western Blot using an antibody against the activated NICD. (d, e). Notch1 activation by CD19-SNAGs and HER2-SNAGs, was evaluated using a fluorescent reporter assay. Increasing concentrations (1 nM, 10 nM, or 100 nM) of Delta^{MAX}-Fc or each SNAG^{Fc} were added to Notch1-Gal4 mCitrine reporter cells alone, or a 1:1 mixture of Notch1 reporter cells and CD19-overexpressing 3T3 cells (d) or HER2-expressing SK-BR-3 cells (e). For b, d, and e, a representative experiment from three biological replicates is shown. Mean fluorescence intensity (MFI) was normalized to the mean MFI of Notch1 reporter cells alone. Error bars represent the standard deviation of three technical replicates with the *P* value by Student's *t* test shown above each comparison.

373
374
Figure 4. SNAG-mediated Notch activation requires endocytosis. (a) Representative
immunofluorescence images of fluorescently labeled CD19-SNAGs (magenta) used to stain the
surface of CD19 expressing 3T3 cells that were kept on ice. CD19-SNAGs visualized by Alexa
Fluor anti-Fc 647. To visualize the contours of the cells, the actin cytoskeleton was stained using
phalloidin-488 (green). Nuclei counterstained by Hoechst 33342 (blue). Scale, 20 mm. (b)
Representative immunofluorescence images of fluorescently labeled CD19-SNAGs used to stain
the surface of CD19 expressing 3T3 cells, followed by washing away unbound SNAGs and
subjecting the cells to a 15 min incubation in a 37 ºC incubator to resume cellular processes
including endocytosis. After 15 min the cells were fixed and stained in parallel with the no
endocytosis samples. Scale, 20 mm. (d-e) Flow cytometry histogram overlays depicting Notch1
reporter activity induced by soluble CD19-SNAG^{Fc}, BC2-SNAG, or BC2-SNAG^{Fc} in the presence
or absence of Dynasore. (c) Notch1 reporter cells were co-cultured with CD19-overexpressing
3T3 cells. In (d) and (e), Notch1 reporter cells were co-cultured with HEK293 cells expressing
DLL4^{HL}. (f) Flow cytometry histogram overlay depicting Notch1 reporter activity induced by
immobilized BC2-SNAG in the presence or absence of Dynasore. A representative histogram is
shown for each experimental condition from one of three biological replicates.
391
392

393
394
395
Figure 5. Trimeric CD40-SNAGs activate Notch1 signaling. (a) Cartoon schematic depicting a
SNAG binding to Notch1 and CD40 receptor on adjacent cell. The CD40-SNAG was generated
by using a GCN4 leucine zipper to trigger dimerization of both the Notch1- and CD40-targeting
domains. (b) A fluorescent reporter assay was used to evaluate SNAG-mediated activation of
Notch1. Increasing concentrations (1 nM, 10 nM, or 100 nM) of Delta^{MAX}-Fc or CD40-SNAG were
added to Notch1-Gal4 mCitrine reporter cells alone or a 1:2 mixture of Notch1 reporter cells and
OCI-Ly3 cells that have high expression of the CD40 receptor and fluorescence was measured
by flow cytometry. A representative experiment from three biological replicates is shown. Mean
fluorescence intensity (MFI) was normalized to the mean MFI of Notch1 reporter cells alone. Error
bars represent the standard deviation of three technical replicates with the *P* value by Student's *t*
test shown above each comparison. (c) Flow cytometry plots showing the percentage of B cells
and T cells on a healthy donor sample, gated on lymphoid, single, live cells. The percentage of
CD40+ cells in the B cell subpopulation (CD19+) are shown in the top panel, and the T cell
subpopulation (CD3+) on shown in the bottom panel.

Figure. 6. CD40-SNAGs improve the expansion and phenotype of γδ T cells. (a) Bar chart
showing the percentage of live cells in culture with different concentrations of CD40-SNAG or
controls; gated on lymphoid, single cells. (b) Percentage of γδ TCR expression in T cells cultured
with CD40-SNAG vs controls. The histogram to the right shows data from a representative
biological replicate; gated on lymphoid, single, viable, CD3+ cells. (c) Bar graph showing the total
number of cells in culture by each condition (left) and the relative number of γδ TCR+ cells (right).
(d) Bar charts representing the distribution of different differentiation phenotypes in γδ TCR+ T
cells cultured with CD40-SNAG or controls. A representative flow cytometry zebra plot from one
healthy donor is shown to the right. Cells were analyzed and counted on day seven, in the bar
charts each symbol correspond to an independent donor (biological replicate), p values over the
bars were calculated using paired t-tests comparing to the control.

MATERIALS AND METHODS

Protein expression and purification

All SNAG sequences were cloned into a pAcGP67A vector for insect cell production containing an N-terminal gp67 signal peptide and C-terminal 8xHis-tag. Monomeric SNAGs were generated by fusing a truncated version of the Delta^{MAX} protein spanning from the N-terminus to EGF5 (N-EGF5) fused to a biomarker-targeting scFv or nanobody using a flexible (GS)$_5$ linker. Dimeric SNAG^{Fc} constructs were generated by fusing Delta^{MAX} (N-EGF5) and the biomarker targeting module to the N- and C-termini of a human IgG1 Fc domain, respectively. All SNAG^{Fc} constructs contained short GSG-linkers between the Fc sequence and Delta^{MAX} or the targeting module. Published sequences of atezolizumab, trastuzumab, and loncastuximab[63] were converted into a scFv format prior to being incorporated into SNAGs, and the sequence of the BC2-specific nanobody[33] was obtained from the Protein Data Bank (PDB ID 5VIN). Each scFv was generated by fusing the C-terminus of the variable heavy (V$_H$) domain to the N-terminus of the variable light (V$_L$) domain with a (GGGGS)$_3$ linker. Biotinylated Delta^{MAX}(N-EGF5) protein was generated through enzymatic modification of a C-terminal biotin acceptor peptide (BirA tag) as previously described[30]. The "headless" loss-of-function DLL4^{HL} mutant was generated by replacing the C2 and DSL domains of human DLL4 with the BC2-peptide sequence, which was connected to the N-terminus of EGF1 by a short GSG-linker. The DLL4^{HL} construct was cloned into a pLenti-IRES-Puro vector for mammalian expression.

All SNAG constructs in this study were expressed for by infecting *Trichoplusia ni* insect cell cultures (Expression Systems) at a density of 2×10^6 cells ml^{-1} with recombinant Baculovirus. Culture supernatants were harvested after 48h, and proteins were purified by nickel and size-exclusion chromatography. Biotinylated proteins were site-specifically modified using BirA ligase and excess biotin was removed by purifying the proteins on a size-exclusion column. Protein purity was assessed by SDS–PAGE using TGX 12% Precast gels (Bio-Rad). All proteins were flash-frozen in liquid nitrogen and stored at −80 °C following purification.

Cell culture and generation of cell lines

Mammalian cells were cultured at 37 °C, with a humidified atmosphere of 5% CO2, washed with Dulbecco's PBS (DPBS, Corning), and detached with trypsin–EDTA 0.25% (Gibco) for subculturing or cell-based assays. Notch reporter cell lines CHO-K1 N1-Gal4 were a gift from Dr. Michael Elowitz (California Institute of Technology)[31]. Briefly, transfections of HEK293T cells were carried out with packaging vectors VSV-G and d8.9 in the presence of polyethyleneimine at a ratio

of 4:1 (DNA:polyethyleneimine). HER2[+] SK-BR-3 cells, human CD19-overexpressing 3T3 cells,

PD-L1[+] MDA-MB-231, and CD40[+] OCI-Ly3 cells were gifts from Drs. Brian Czerniecki, Fred

Locke, Eric Lau, and John Cleveland, respectively (Moffit Cancer Center). HEK293T, SK-BR-3,

3T3 mouse fibroblast, and MDA-MB-231 cells were cultured in high-glucose DMEM (Cytiva)

supplemented with 10% FBS (peak serum) and 2% penicillin/streptomycin (Gibco). Puromycin

5 µg ml^{-1} was added to HEK293T cell cultures to maintain homogeneous populations of receptor-

expressing cells. CHO-K1 N1-Gal4 cells were cultured in minimum essential medium Eagle-alpha

modification (α-MEM, Cytiva) supplemented with 10% FBS (peak serum), 2%

penicillin/streptomycin (Gibco), 400 µg ml−1 of zeocin (Alfa aesar) and 600 µg ml−1 of geneticin

(Gibco). Expression of receptors on the cell surface was confirmed by flow cytometry (BD Accuri

C6 plus) staining the cell lines with anti-hDLL4 PE, anti-JAG1 APC, anti-hPDL1 FITC, anti-hHER2

(anti-IgG FITC), anti-hCD19 FITC, and anti-CD40 PE in DMEM supplemented with 10% FBS for

1 h at 4 °C.

Notch activation with Delta[MAX] multimers

On day one, biotinylated Delta[MAX], Delta[MAX] tetramers formed with streptavidin, or Delta[MAX]-Fc

were reconstituted in DPBS and adsorbed to tissue culture 96-well plates (Costar) for 1 h at 37 °C.

The wells were then washed three times with 200 µl of DPBS to remove unbound proteins. Next,

CHO-K1 N1-Gal4 cells were detached with trypsin–EDTA 0.25% (Gibco) and manually counted.

Appropriate dilutions were prepared in α-MEM media to ensure 30,000 CHO-K1 N1-Gal4 cells

per well in a volume of 50 mL. Cells were transferred to the ligand-coated plates and cultured for

24 h at 37 °C in 5% CO2. On day two, CHO-K1 N1-Gal4 cells were washed with 200 µl DPBS,

detached with 30 µL of trypsin–EDTA 0.25%, and quenched with 170 mL of α-MEM media. Finally,

cells were resuspended, and the H2B-mCitrine signal was measured by flow cytometry (BD Accuri

C6 plus). CHO-K1 N1-Gal4 cells alone were used as the control. The measurements represent

the mean fluorescent intensity as fold-change of Notch activation ± s.d. of three technical

replicates. Notch activation was normalized to wells containing CHO-K1 N1-Gal4 cells alone.

Notch activation with SNAGs in coculture of cells expressing the target tumor biomarker

On day one, cells expressing the target receptor of the SNAG (signal-sending cells) were

detached with trypsin–EDTA, counted manually, and dilutions prepared such that 50 mL of DMEM

containing 15,000 signal-sender cells were added to wells of a tissue culture 96-well plate. The

next day, CHO-K1 N1-Gal4 reporter cells (signal-receiver cells) were detached with trypsin–

EDTA, and 50 µl of α-MEM media containing 30,000 cells were added to the tissue culture 96-

well plate containing the signal-sending cells after combining with the indicated Delta^{MAX} or SNAG

protein. For the CD40-SNAG experiments, the difference was that an equal amount of signal-

sending and signal-receiving cells were added same day. Wells without signal-sending cells were

used to determine background activation of Notch by Delta^{MAX} and SNAGs. When testing

inhibition of endocytosis, 80 µM of the Dynamin inhibitor I (Dynasore, Sigma) was added to the

mixture of Notch reporter cells with protein and added to the tissue culture 96-well plate containing

the signal-sending cells. Notch activation was measured as previously described.

Western blot detection of Notch1 activation by the PDL1-SNAG^{Fc} in MDA-MB-231 cells.

Delta^{MAX} (100 nM protein in 600 mL of DPBS) was non-specifically adsorbed to a single well of a

12-well plate for 1 hour at 37 ºC as a positive control for Notch1 activation. The positive control

well and three additional wells were then seeded with 200 x 10^3 cells with MDA-MB-231 cells.

The plate was centrifuged at 400 x g for 4 min to ensure cells were retained at the bottom of each

well, and then the media of all wells was discarded. In the first uncoated well, 600 mL of DMEM

was added as a negative control. The second well was filled with 600 mL of media containing 100

nM of Delta^{MAX}-Fc to monitor Notch1 activation by soluble ligand. The third was filled with 600 mL

of media containing 100 nM PDL1-SNAG^{Fc}. The following day, the media was aspirated from all

four wells, and the samples were resuspended in 60 mL of Laemli sample buffer with 5% b-

mercaptoethanol to lyse cells, followed by boiling at 100 °C for 4 min. Lastly, the samples were

analyzed by western blotting using equal protein amounts of cell lysates separated by SDS–PAGE

(12% Mini-PROTEAN TGX Precast Protein Gels, Bio-Rad) and transferred to PVDF membranes

using an iBlot2 Gel Transfer Device (Thermo Fisher Scientific). The membranes were blocked in

3% BSA + 0.1% TBS-Tween. Primary antibodies were anti-Notch1 (D1E11 rabbit mAb, Cell

Signaling Technology, 1:1,000), anti-cleaved Notch1 (Val1744 rabbit mAb, Cell Signaling

Technology, 1:1,000), and b-actin (rabbit polyclonal Ab, Cell Signaling Technology, 1:1,000).

Secondary antibody anti-Rabbit IgG conjugated to HRP (Goat polyclonal Ab, Vector Laboratories,

1:8,000) was used for detection of proteins using SuperSignal West Pico PLUS

Chemiluminescent Substrate (Thermo Fisher Scientific). Images were acquired using a Chemidoc

Imaging System and analyzed with Image-Lab v.6 software (Bio-Rad).

Immunofluorescent cell staining. For endocytosis assays, cells were grown on glass-like

polymer bottoms in 24 well black frame plates (Cellvis). For visualization of CD19-SNAG^{Fc} protein

binding, 500 nM protein was preincubated with anti-Fc 647 (Alexa Fluor) at 1:200 dilution for 1h

on rotation in +4°C. The CD19-SNAG^{Fc}-647 solution was added to cells on ice that were further

kept in +4°C for 1 h. For endocytosis, the incubation was followed by washing away non-bound

CD19-SNAG^Fc-647 with PBS, and 37°C DMEM added to the cells followed by a 15 min incubation

in a 37°C incubator. After incubation of CD19-SNAG^Fc-647 with or without endocytosis, the cells

were fixed in 3% paraformaldehyde and permeabilized with 0.15% Triton X-100 in PBS for 10 min

at RT. Nonspecific binding was blocked by incubation in 3% BSA in PBS with 0.05% Triton X-100

and 0.1M glycine for 60 min at RT. Cells were further stained for filamentous actin with Alexa 488

conjugated to phalloidin (Invitrogen) for 45 min to visualize contours of the individual cells.

Hoechst 33342 (Invitrogen) was used to counterstain nuclei. Images were acquired using a

Keyence BZ-X710 microscope using a Nikon Plan Apo 20x objective. The far-red channel

(magenta) was processed with the de-haze function in the BZ-X710LE analyzer software. A

minimum of 100 cells were imaged for each condition.

γδ expansion from PBMCs with CD40-SNAG

Peripheral blood mononuclear cells (PBMCs) were isolated from healthy-donor buffy coats using

an established density gradient protocol[51]. γδ expansion was performed using zoledronic acid in

addition or not of CD40-SNAG. Briefly, ten million cells per condition were resuspended in RPMI

[supplemented with 5%FBS, antibiotic, IL-2 (100 IU/ml) and zoledronic acid (4μM)] at 1x106

cells/ml. The cells were spiked with CD40-SNAG at three different concentrations (50, 100 or

500ng/ml), Delta^MAX-Fc (100ng/ml), or CD40L trimer (100ng/ml), and cultured in 24-well plates

(2ml/well) for three days at 37C (5% CO2). On day three, the full media was replaced by new

media with zoledronic acid and CD40-SNAG, Delta^MAX-Fc, or CD40L trimer as on day one. Two

days later, cells were lifted from the plates, centrifuged, and resuspended in new RMPI media

[5%FBS, antibiotic, IL-2 (100 IU/ml)] without additional molecules. At day seven, T cells were

counted, and their phenotype was assessed by surface staining and flow cytometry (using

markers for human anti-CD3 – BV711, anti-Vd2 TCR – FITC, anti-CD45RA – BV421, anti-CD27

– BV605, and aqua live/dead fluorescent dye).

REFERENCES

1. Sprinzak, D. & Blacklow, S. C. Biophysics of Notch Signaling. *Annu Rev Biophys* **50**, 157–189

(2021).

2. Zhou, B. *et al.* Notch signaling pathway: architecture, disease, and therapeutics. *Sig*

Transduct Target Ther **7**, 1–33 (2022).

3. Garg, V. *et al.* Mutations in NOTCH1 cause aortic valve disease. *Nature* **437**, 270–274 (2005).

4. Li, L. *et al.* Alagille syndrome is caused by mutations in human Jagged1, which encodes a

ligand for Notch1. *Nat. Genet.* **16**, 243–251 (1997).

5. Joutel, A. *et al.* Notch3 mutations in CADASIL, a hereditary adult-onset condition causing

stroke and dementia. *Nature* **383**, 707–710 (1996).

6. Bulman, M. P. *et al.* Mutations in the human delta homologue, DLL3, cause axial skeletal

defects in spondylocostal dysostosis. *Nat Genet* **24**, 438–441 (2000).

7. Radtke, F. & Raj, K. The role of Notch in tumorigenesis: oncogene or tumour suppressor?

Nat Rev Cancer **3**, 756–767 (2003).

8. Hori, K., Sen, A. & Artavanis-Tsakonas, S. Notch signaling at a glance. *Journal of Cell Science*

126, 2135–2140 (2013).

9. Andersson, E. R. & Lendahl, U. Therapeutic modulation of Notch signalling--are we there

yet? *Nat Rev Drug Discov* **13**, 357–378 (2014).

10. Brandstadter, J. D. & Maillard, I. Notch signalling in T cell homeostasis and differentiation.

Open Biol **9**, 190187 (2019).

11. Schmitt, T. M. & Zúñiga-Pflücker, J. C. Induction of T Cell Development from Hematopoietic

Progenitor Cells by Delta-like-1 In Vitro. *Immunity* **17**, 749–756 (2002).

12. Kelliher, M. A. & Roderick, J. E. NOTCH Signaling in T-Cell-Mediated Anti-Tumor Immunity

and T-Cell-Based Immunotherapies. *Front Immunol* **9**, 1718 (2018).

13. Sierra, R. A. *et al.* Rescue of notch-1 signaling in antigen-specific CD8+ T cells overcomes

tumor-induced T-cell suppression and enhances immunotherapy in cancer. *Cancer Immunol*

Res **2**, 800–811 (2014).

14. Wilkens, A. B. *et al.* NOTCH1 signaling during CD4+ T-cell activation alters transcription

factor networks and enhances antigen responsiveness. *Blood* **140**, 2261–2275 (2022).

15. Kondo, T. *et al.* Notch-mediated conversion of activated T cells into stem cell memory-like T

cells for adoptive immunotherapy. *Nat Commun* **8**, 15338 (2017).

16. Meloty-Kapella, L., Shergill, B., Kuon, J., Botvinick, E. & Weinmaster, G. Notch ligand

endocytosis generates mechanical pulling force dependent on dynamin, epsins, and actin.

Dev. Cell **22**, 1299–1312 (2012).

17. Wang, X. & Ha, T. Defining Single Molecular Forces Required to Activate Integrin and Notch

Signaling. *Science* **340**, 991–994 (2013).

18. Gordon, W. R. *et al.* Mechanical Allostery: Evidence for a Force Requirement in the

Proteolytic Activation of Notch. *Dev. Cell* **33**, 729–736 (2015).

19. Rebay, I. *et al.* Specific EGF repeats of Notch mediate interactions with Delta and Serrate:

implications for Notch as a multifunctional receptor. *Cell* **67**, 687–699 (1991).

20. Luca, V. C. *et al.* Structural biology. Structural basis for Notch1 engagement of Delta-like 4.

Science **347**, 847–853 (2015).

21. Luca, V. C. *et al.* Notch-Jagged complex structure implicates a catch bond in tuning ligand

sensitivity. *Science* eaaf9739 (2017) doi:10.1126/science.aaf9739.

22. Gordon, W. R. *et al.* Structural basis for autoinhibition of Notch. *Nat. Struct. Mol. Biol.* **14**,

295–300 (2007).

23. De Strooper, B. *et al.* A presenilin-1-dependent gamma-secretase-like protease mediates

release of Notch intracellular domain. *Nature* **398**, 518–522 (1999).

24. Brou, C. *et al.* A novel proteolytic cleavage involved in Notch signaling: the role of the

disintegrin-metalloprotease TACE. *Mol. Cell* **5**, 207–216 (2000).

25. Schroeter, E. H., Kisslinger, J. A. & Kopan, R. Notch-1 signalling requires ligand-induced

proteolytic release of intracellular domain. *Nature* **393**, 382–386 (1998).

26. Varnum-Finney, B. *et al.* Immobilization of Notch ligand, Delta-1, is required for induction of

notch signaling. *J. Cell. Sci.* **113 Pt 23**, 4313–4318 (2000).

27. Trotman-Grant, A. C. *et al.* DL4-μbeads induce T cell lineage differentiation from stem cells

in a stromal cell-free system. *Nat Commun* **12**, 5023 (2021).

28. Li, K. *et al.* Modulation of Notch Signaling by Antibodies Specific for the Extracellular

Negative Regulatory Region of NOTCH3. *J. Biol. Chem.* **283**, 8046–8054 (2008).

29. Tiyanont, K., Wales, T. E., Siebel, C. W., Engen, J. R. & Blacklow, S. C. Insights into Notch3

Activation and Inhibition Mediated by Antibodies Directed Against its Negative Regulatory

Region. *J Mol Biol* **425**, 3192–3204 (2013).

30. Gonzalez-Perez, D. *et al.* Affinity-matured DLL4 ligands as broad-spectrum modulators of

Notch signaling. *Nat Chem Biol* 1–9 (2022) doi:10.1038/s41589-022-01113-4.

31. Sprinzak, D. *et al.* Cis-interactions between Notch and Delta generate mutually exclusive

signalling states. *Nature* **465**, 86–90 (2010).

32. Cordle, J. *et al.* A conserved face of the Jagged/Serrate DSL domain is involved in Notch

trans-activation and cis-inhibition. *Nat Struct Mol Biol* **15**, 849–857 (2008).

33. Braun, M. B. *et al.* Peptides in headlock – a novel high-affinity and versatile peptide-binding

nanobody for proteomics and microscopy. *Sci Rep* **6**, (2016).

34. Andersson, E. R. *et al.* Mouse Model of Alagille Syndrome and Mechanisms of Jagged1

Missense Mutations. *Gastroenterology* **154**, 1080–1095 (2018).

35. Lafkas, D. *et al.* Therapeutic antibodies reveal Notch control of transdifferentiation in the

adult lung. *Nature* **528**, 127–131 (2015).

36. Powles, T. *et al.* MPDL3280A (anti-PD-L1) treatment leads to clinical activity in metastatic

bladder cancer. *Nature* **515**, 558–562 (2014).

37. Xiao, D. *et al.* Development of bifunctional anti-PD-L1 antibody MMAE conjugate with

cytotoxicity and immunostimulation. *Bioorganic Chemistry* **116**, 105366 (2021).

38. Zammarchi, F. *et al.* ADCT-402, a PBD dimer-containing antibody drug conjugate targeting

CD19-expressing malignancies. *Blood* **131**, 1094–1105 (2018).

39. Lewis Phillips, G. D. *et al.* Targeting HER2-positive breast cancer with trastuzumab-DM1, an

antibody-cytotoxic drug conjugate. *Cancer Res* **68**, 9280–9290 (2008).

40. Parks, A. L., Klueg, K. M., Stout, J. R. & Muskavitch, M. A. Ligand endocytosis drives receptor

dissociation and activation in the Notch pathway. *Development* **127**, 1373–1385 (2000).

41. Koo, B.-K. *et al.* An obligatory role of mind bomb-1 in notch signaling of mammalian

development. *PLoS One* **2**, e1221 (2007).

42. McMillan, B. J. *et al.* A tail of two sites: a bipartite mechanism for recognition of notch

ligands by mind bomb E3 ligases. *Mol Cell* **57**, 912–924 (2015).

43. Cao, R. *et al.* Structural requirements for activity of Mind bomb1 in Notch signaling.

Structure (2024) doi:10.1016/j.str.2024.07.011.

44. Yellin, M. J. *et al.* CD40 molecules induce down-modulation and endocytosis of T cell surface

T cell-B cell activating molecule/CD40-L. Potential role in regulating helper effector function.

J Immunol **152**, 598–608 (1994).

45. Mitchison, N. A. T-cell–B-cell cooperation. *Nat Rev Immunol* **4**, 308–312 (2004).

46. Naito, M. *et al.* CD40L-Tri, a novel formulation of recombinant human CD40L that effectively

activates B cells. *Cancer Immunol Immunother* **62**, 347–357 (2013).

47. Mensurado, S., Blanco-Domínguez, R. & Silva-Santos, B. The emerging roles of γδ T cells in

cancer immunotherapy. *Nat Rev Clin Oncol* **20**, 178–191 (2023).

48. Ganapathy, T., Radhakrishnan, R., Sakshi, S. & Martin, S. CAR γδ T cells for cancer

immunotherapy. Is the field more yellow than green? *Cancer Immunol Immunother* **72**, 277–

286 (2023).

49. Frieling, J. S. *et al.* γδ-Enriched CAR-T cell therapy for bone metastatic castrate-resistant

prostate cancer. *Sci Adv* **9**, eadf0108.

50. Gogoi, D., Dar, A. A. & Chiplunkar, S. V. Involvement of Notch in activation and effector

functions of γδ T cells. *J Immunol* **192**, 2054–2062 (2014).

51. Peters, C., Kouakanou, L., Oberg, H.-H., Wesch, D. & Kabelitz, D. *In vitro* expansion of Vγ9Vδ2

T cells for immunotherapy. in *Methods in Enzymology* (eds. Galluzzi, L. & Rudqvist, N.-P.) vol.

631 223–237 (Academic Press, 2020).

52. Liu, Q., Sun, Z. & Chen, L. Memory T cells: strategies for optimizing tumor immunotherapy.

Protein Cell **11**, 549–564 (2020).

53. Medina, E., Perez, D. H., Antfolk, D. & Luca, V. C. New tricks for an old pathway: emerging

Notch-based biotechnologies and therapeutics. *Trends Pharmacol Sci* S0165-

6147(23)00213–4 (2023) doi:10.1016/j.tips.2023.09.011.

54. Bonnici, L., Suleiman, S., Schembri-Wismayer, P. & Cassar, A. Targeting Signalling Pathways in

Chronic Wound Healing. *International Journal of Molecular Sciences* **25**, 50 (2024).

55. Mazor, Y. *et al.* Enhanced tumor-targeting selectivity by modulating bispecific antibody

binding affinity and format valence. *Sci Rep* **7**, 40098 (2017).

56. Haber, L. *et al.* Generation of T-cell-redirecting bispecific antibodies with differentiated

profiles of cytokine release and biodistribution by CD3 affinity tuning. *Sci Rep* **11**, 14397

(2021).

57. del Álamo, D., Rouault, H. & Schweisguth, F. Mechanism and Significance of cis-Inhibition in

Notch Signalling. *Current Biology* **21**, R40–R47 (2011).

58. Boyd, N. *et al.* 'Off-the-Shelf' Immunotherapy: Manufacture of CD8+ T Cells Derived from

Hematopoietic Stem Cells. *Cells* **10**, 2631 (2021).

59. Tan, W. K., Tay, J. C., Zeng, J., Zheng, M. & Wang, S. Expansion of Gamma Delta T Cells - A

Short Review on Bisphosphonate and K562-Based Methods. *Journal of Immunological*

Sciences **2**, (2018).

60. Verkerk, T. *et al.* Isolation and expansion of pure and functional γδ T cells. *Front. Immunol.*

15, (2024).

61. Xu, X. *et al.* Insights into Autoregulation of Notch3 from Structural and Functional Studies of

Its Negative Regulatory Region. *Structure* **23**, 1227–1235 (2015).

62. Smyrlaki, I. *et al.* Soluble and multivalent Jag1 DNA origami nanopatterns activate Notch

without pulling force. *Nat Commun* **15**, 465 (2024).

63. Abanades, B. *et al.* The Patent and Literature Antibody Database (PLAbDab): an evolving

reference set of functionally diverse, literature-annotated antibody sequences and

structures. *Nucleic Acids Research* **52**, D545–D551 (2024).

698