1
2
3
4   **Title:** Phylogenomic prediction of interaction networks in the presence of gene duplication
5

6   **Authors:**
7   Evan S Forsythe[1,2,3,*], Tony C Gatts[4], Linnea E Lane[2], Chris deRoux[4], Monica Berggren[4],
8   Elizabeth A Rehmann[3], Emily N Zak[2], Trinity Bartel[2], Luna L'Argent[3], and Daniel B Sloan[4]
9

Affiliations:
[1]Oregon State University, Department of Integrative
[2]Oregon State University-Cascades, Biology Program
[3]Oregon State University-Cascades, Biochemistry and Molecular Biology Program
[4]Colorado State University, Department of Biology
[*]Correspondence
16
17

Abstract:
Assigning gene function from genome sequences is a rate-limiting step in molecular biology
research. A protein's position within an interaction network can potentially provide insights into
its molecular mechanisms. Phylogenetic analysis of evolutionary rate covariation (ERC) in protein
sequence has been shown to be effective for large-scale prediction of functional relationships and
interactions. However, gene duplication, gene loss, and other sources of phylogenetic
incongruence are barriers for analyzing ERC on a genome-wide basis. Here, we developed
ERCnet, a bioinformatic program designed to overcome these challenges, facilitating efficient all-
vs-all ERC analyses for large protein sequence datasets. We simulated proteome datasets and
found that *ERCnet* achieves combined false positive and negative error rates well below 10% and
that our novel 'branch-by-branch' length measurements outperforms 'root-to-tip' approaches in
most cases, offering a valuable new strategy for performing ERC. We also compiled a sample set
of 35 angiosperm genomes to test the performance of *ERCnet* on empirical data, including its
sensitivity to user-defined analysis parameters such as input dataset size and branch-length
measurement strategy. We investigated the overlap between *ERCnet* runs with different species
samples to understand how species number and composition affect predicted interactions and to
identify the protein sets that consistently exhibit ERC across angiosperms. Our systematic
exploration of the performance of *ERCnet* provides a roadmap for design of future ERC analyses
to predict functional interactions in a wide array of genomic datasets. *ERCnet* code is freely
available at *https://github.com/EvanForsythe/ERCnet*.
38
39

40
Introduction:

Proteins that interact with each other often exhibit correlated rates of sequence evolution across a phylogeny, which have been attributed to shared selective pressures or reciprocal coevolution (Clark et al. 2012). The statistical signature of these interactions is known as evolutionary rate covariation (ERC). ERC analyses have been applied in a wide variety of organisms (Steenwyk et al.; Goh et al. 2000; Ramani and Marcotte 2003; Sato et al. 2005; Clark and Aquadro 2010; De Juan et al. 2013; Wolfe and Clark 2015; Yan et al. 2019; Forsythe et al. 2021; Rei Liao et al. 2022; Tao et al. 2024), and bioinformatic tools have been developed to use ERC to identify novel interaction partners (Wolfe and Clark 2015; Steenwyk et al. 2021) and correlated molecular and phenotypic evolution (Asar et al. 2023; Duchêne et al. 2024). Moreover, ERC analyses are beginning to be applied in a genome-wide manner to yield large sets of predicted protein-protein interactions (i.e, interactome networks) (Steenwyk et al. 2021; Steenwyk et al. 2022). A network view of genetic interactions has proven valuable in revealing the interconnected nature of biological systems (Mao et al. 2009; Rao and Dixon 2019; Wright et al. 2024), and ERC has the potential to be a powerful and efficient method to infer such interaction networks.

As we scale ERC analyses to more taxonomic groups and to larger numbers of genes, phylogenetic incongruence arising from gene duplication/loss, reticulate evolution, and incomplete lineage sorting becomes increasingly problematic (Degnan and Rosenberg 2013; Hahn and Nakhleh 2015). These processes can all lead to trees inferred from different loci (i.e. gene trees) having different branching patterns (i.e., topologies) from each other. Discordant gene trees make it difficult to directly compare evolutionary rates across gene trees because it is unclear which branches correspond to each other. Prior studies have overcome this challenge through combinations of (1) focusing on groups with low levels of phylogenetic incongruence and gene duplication/loss, (2) filtering gene families to retain only gene families with one-to-one orthology, (3) constraining gene tree topologies to the accepted species tree, and (4) measuring branch lengths in a root-to-tip manner that does not depend on topology. These strategies have worked well when the underlying level of phylogenetic incongruence is minimal; however, they may be insufficient in taxonomic groups such as plants that tend to experience more complex genome evolution with extensive gene and whole-genome duplication (Wendel 2015; Panchy et al. 2016; Forsythe et al. 2020). Thus, additional methods are needed to perform ERC analyses in such taxa. A common approach to performing analyses in the face of phylogenetic incongruence is to incorporate gene-tree/species-tree (GT/ST) reconciliation (Vernot et al. 2008; Stolzer et al. 2012; Wu et al. 2014; Comte et al. 2020). GT/ST reconciliation regards phylogenetic incongruence not as noise, but as a signal with which to infer evolutionary events such as gene duplication. Importantly, GT/ST reconciliation also provides a framework for interpreting multiple gene trees in the shared context of a prevailing species tree, which can provide a valuable basis for making direct comparisons between discordant gene trees.

We previously developed a GT/ST reconciliation-based approach to performing genome-wide ERC in plants (Forsythe et al. 2021) that operated at the level of a 'one-vs-all' comparison strategy, in which a single locus was tested against a genome-wide set of genes. The biological findings from this approach suggested that large groups of functionally interacting proteins exhibit correlated evolution. However, the limited lens of one-vs-all comparisons only allowed for indirect tests of this hypothesis. A network view of the interactome is needed to assess whether large sets of proteins with correlated rates of evolution share functional relationships. Such a network view

requires ERC to be applied in an 'all-vs-all' manner that tests all pairwise combinations of proteins.
This added dimensionality introduces and exacerbates scaling challenges to ERC analyses.

Here, we present *ERCnet*, a program for performing all-vs-all ERC analyses in the presence of
phylogenetic incongruence. We describe our phylogenomic workflow, which automates and
parallelizes the computationally intensive analytical steps, and our novel approach to performing
branch-by-branch ERC analysis even in trees with gene duplicates. We compile simulated and
empirical test datasets to test *ERCnet* performance in response to user-defined analysis parameters
and experimental design choices, thus providing an evidence-based roadmap for future ERC
analyses.

Results:

Overcoming gene duplication/loss/incongruence with branch length reconciliation

ERCnet (*https://github.com/EvanForsythe/ERCnet*) is a computational pipeline designed to
predict functional interactions among proteins based on phylogenetic signatures of correlated
evolution and overcome key scaling challenges associated running all-vs-all analyses on a
genome-wide level (Fig. 1). *ERCnet* uses the output of the gene family clustering program
Orthofinder (Emms and Kelly 2015, 2019)*,* thus making it easy to plug *ERCnet* into existing
genomic workflows. Our pipeline employs phylogenetic bootstrap resampling and GT/ST
reconciliation strategies to account for phylogenetic incongruence and uncertainty, thus
accommodating gene families that have undergone duplication and loss. *ERCnet* automates and
parallelizes the computational steps needed to achieve genome-wide network analyses of ERC and
consists of four major analytical steps, (1) phylogenomic analyses, (2) GT/ST reconciliation, (3)
ERC analyses, and (4) network analyses (See Methods; Fig 1A).

In addition to automating standard phylogenetic analyses, *ERCnet* provides a novel procedure for
comparing rates in a branch-by-branch (BXB) manner even when comparing gene trees with
paralogs, missing taxa, and/or topological incongruence. Some previous implementations of ERC
have employed a root-to-tip (R2T) methodology for measuring branch lengths. The R2T approach
simplifies the task of parsing gene trees (especially when incongruence is present); however, it
results in statistical non-independence among datapoints used in correlation analyses because
different species can share internal branches within a tree. This statistical shortcoming has been
previously recognized (Yan et al. 2019; Forsythe et al. 2021; Smith et al. 2024) but was largely
viewed as a 'necessary evil' due to lack of a method for traversing discordant gene trees to extract
relevant branches for direct comparisons.

Our novel method, branch-length reconciliation (BLR), provides a strategy for avoiding the
statistical shortcomings of the R2T method. BLR extracts individual internal and external branch
length measurements in a species-tree-aware manner, which allows for direct comparisons across
gene trees. The major barrier to performing BXB-based ERC between incongruent phylogenetic
trees has been that it is difficult to systematically define which branches (if any) are shared between
two gene trees with potentially different taxon-composition and topology. The premise of GT/ST
reconciliation methods is that all gene trees evolve within the context of an overarching species
tree. This means that the species tree is the common denominator that connects all gene trees in a
phylogenomic dataset. BLR uses the species tree as the common comparison point by tallying
branch length information from individual gene trees in context of the species tree. BLR

accomplishes this task by incorporating the labeled coalescent tree reconciliation structure
employed in *DLCpar* (Wu et al. 2014), which provides information that maps gene tree nodes to
species tree nodes. Using this information, BLR iteratively extracts the branch(es) on each gene
tree that correspond to each species tree branch. During this process BLR also tracks whether a
duplication event occurred in the gene tree and, if so, takes the average branch length from the
resulting paralogs. When a gene tree lacks information for a given species tree branch, BLR stores
an "NA" for that branch. The output of BLR is a table of branch length measurements for each
gene tree that are standardized by the species tree branches, allowing comparison across gene trees.
This approach provides the ability to extract branch-specific evolutionary rate information from
gene trees with incongruent topologies and histories of gene duplication and loss, enabling ERC
analyses to include larger gene sets and be applied to a wider range of taxa.

The all-by-all ERC analyses implemented in *ERCnet* yield large tables of ERC results, including
p-values calculated from Pearson and/or Spearman correlation analyses. These ERC results are
filtered according to user-defined cutoffs to retain the pairwise combinations displaying significant
p-values and R^2 values, which we refer to as "ERC hits". The full set of significant ERC hits form
the edges in an interactome network, which can be further analyzed to identify emergent properties,
such as clustered modules of cofunctional proteins. Below, we systematically explore the
performance of *ERCnet* on simulated and empirical genomic datasets to identify and optimize the
parameters that influence computational prediction of interaction networks.

Accuracy of ERCnet on simulated datasets

We simulated protein sequences for 21 species under a model in which 100 of the 1000 gene
families underwent co-accelerated rates of protein evolution (Fig. 2A and B). We used our
simulated dataset to assess false positive and negative rates of ERC hits under different *p* and R^2
value significance cutoffs and with alternative methodologies for measuring branch lengths (Fig.
2C-F). We found that our BXB method is more conservative than the R2T method and that the
Spearman correlation method is more conservative than the Pearson method. The rates obtained
from dual filtering (Pearson and Spearman) were very similar to the rates obtained from the
Spearman filter alone, meaning Spearman correlation filter acts as the bottleneck in nearly all
cases. The combination of BXB and Spearman leads to an especially conservative approach,
yielding fewer than 10 total ERC hits and false positive and negative rates approaching 0 and 1
respectively. We summed the false positive and negative rates to identify the overall parameter
combination with the lowest combined error rate and found [*BXB; Pearson; P<0.05; R^2>0.5*] to
show the best performance, with a false positive rate of 6.5×10^{-4} (0.065%) and false negative rate
of 0. By this metric, the top five error-minimizing combinations all made use of the BXB method
and Pearson correlation coefficient, indicating these methods are favorable for minimizing false
positive and negative error. The best performing R2T run [*R2T; Pearson; P<0.00001; R^2>0.2*]
yielded a false positive rate of 0.032 (3.2%) and a false negative rate of 0.

Next, we simulated a separate proteome dataset under a model in which rate acceleration takes
place in an uncorrelated manner, meaning any ERC hit obtained from this dataset would be
considered a 'spurious' ERC hit (Fig. S1). Consistent with the results above, BXB performed in
a far more conservative manner compared to R2T on the non-accelerated dataset. R2T showed
more than 60,000 spurious hits, whereas the highest number of spurious hits in a BXB run was
19,395. Taken together, these results show that *ERCnet* achieves very promising levels of accuracy
on simulated data and that our BXB method will be a valuable tool in minimizing error in ERC
analyses.

Proteome coverage and statistical power for ERCnet analyses of sample datasets

ERC has been applied in a variety of organisms with considerable success. However, a systematic
evidence-based exploration of the power, accuracy, and efficiency of the ERC method on real
genomic datasets has never been performed. A consequential decision in ERC analyses is the
number of species to include. Currently, researchers have little guidance in making taxon-sampling
choices for ERC analyses, resulting in somewhat arbitrary choices that can likely have a substantial
impact on ERC performance. To address this limitation, we compiled a set of 35 plant proteomes
(i.e., protein sequences from all annotated genes in the genome), spanning angiosperm diversity
(Table S1). We randomly subsampled sets of these proteomes to create test datasets of varying
sizes. The resulting datasets contained $n=10$, $n=15$, $n=20$, and $n=25$ in-group taxa (see Methods).
Five random replicates were performed for each n value, resulting in a total of 20 datasets, each of
which was subjected to full *ERCnet* analyses (Fig. 3).

The first step of *ERCnet* is to filter *Orthofinder* data to identify the gene families ("orthogroups")
that are suitable for phylogenomic analyses by applying filtering cutoffs based on minimum
number of species represented and maximum number of gene copies per species. *ERCnet* provides
a parameter scan option for users to make an informed choice on filtering settings (see Methods).
To understand how filtering parameters affect dataset size, we applied a filtering formula across
all datasets (see Methods) and found that datasets with a larger number of species yield a smaller
number of genes retained after filtering (Fig. 4A). While datasets with a larger number of species
tend to have more total gene families, these gene families tend to be filtered out at a
disproportionally high rate based on both the minimum species representation filter and the
maximum number of gene copies per species filter (Table S2), leading to a net loss of proteome
coverage at larger dataset sizes.

The gene families that pass initial filtering are next run through the phylogenomic pipeline and
ultimately subjected to all-vs-all ERC correlation analyses. Each pairwise ERC comparison
between two gene trees constitutes a correlation analysis. The number of points for each correlation
corresponds to the number of branches shared between the two gene trees being compared, which
is variable because each gene may have experienced different degrees of gene duplication and loss.
The number of overlapping branches is also dependent on the method used to measure branch
lengths (*i.e.* R2T vs BXB). We found that datasets with larger number of species yield a higher
average number of overlapping branches (Fig. 4B and C). This result is expected because larger
datasets yield larger gene trees, which inherently have more branches. Comparing R2T results
(Fig. 4B) to BXB results (Fig. 4C), we see that BXB tends to yield a higher number of overlapping
branches, which stems from including internal branches under BXB, while the R2T methodology
is limited to the number of species in the in-group. For example, a gene tree with four ingroup
species will only have four R2T measurements but will have six branches when internal branches
are included. In general, the number of overlapping branches will directly impact the sample size
of correlation analyses, suggesting that BXB has the potential to add statistical power to ERC
analyses. Thus, there is a clear tradeoff between the proportion of the genome that can be probed
with *ERCnet* (Fig. 4A) and the potential statistical power of ERC analyses (Fig. 4B and C).

Size and functional clustering of networks across datasets:

We used our simulation results (Fig. 2 and S1) to guide our choice of significance filtering criteria
to use on our empirical datasets. We chose [*Pearson; P<0.0001; $R^2>0.4$*] to obtain a reasonable
balance between false positive and negative error across both R2T and BXB-based networks. We

applied this filter to obtain an interactome network for each *ERCnet* run (Fig. 5, S2, and S3). In
general, we found that R2T analyses returned a much higher number of ERC hits than BXB
analyses, resulting in larger networks in terms of both nodes and edges (Fig. 5A vs B). The large
number of hits observed for R2T analyses could partially result from pseudoreplication due to
statistical non-independence of internal branch lengths (described above). Another general pattern
is that networks show more variability between replicates for lower *n*-values. This is likely at least
partially driven by randomness playing a larger role at smaller sample sizes in our taxon sampling
from a fixed-size pool (Fig. 3), highlighting that selection of individual species has a large impact
on ERC-based networks at small *n*-values. Interestingly, we do not detect clear trends between *n*
and the size of networks. For R2T-based networks, there is some evidence that small *n*-values
yield networks with fewer edges (Fig. 5A), consistent with idea that R2T-based methods may not
achieve a sufficient number of overlapping branches to yield significant ERC hits (see Fig. 4B).
BXB ERC analyses, on the other hand, display the highest number of ERC hits for intermediate *n*
values (Fig 5B). This result suggests that there could be a 'sweet-spot' that balances the number
of genes retained after filtering (see Fig. 4A) and the number of overlapping branches (see Fig.
4C) to achieve coverage and statistical power for BXB-based ERC analyses. Taken together, the
differences in trends for R2T versus BXB-based analyses indicate that the optimal dataset size
depends on the branch-length measuring method.

The number of ERC hits yielded by *ERCnet* analyses defines the total number of edges in the
interaction network output with each run of *ERCnet*. The size of the network is an important
component of *ERCnet* performance; however, it is also important to understand contribution of
biological signal versus noise in empirical ERC networks. We sought a summary statistic that can
serve as a proxy for biological signal in ERC networks. To this end, we incorporated subcellular
localization annotations from *Arabidopsis thaliana* and used the graph-theory assortativity
coefficient to quantify the degree to which co-localized proteins are clustered in interaction
networks (Fig. 5C-D). For BXB analyses, we found that all but one of the *ERCnet* runs yielded
networks with significantly positive assortativity coefficient, providing another promising
indication that *ERCnet* successfully detects 'true' biological signal at a level that outweighs
background noise. BXB shows significantly positive assortativity across all replicates (Fig 5D).
There is not a clear trend between *n* and clustering, but, similar to network size (Fig 5A-B), there
is much more variability between replicates at *n*=10. R2T shows a slight positive relationship
between *n* and assortativity. This includes a clear outlier replicate [n=10; replicate=4] that shows
negative assortativity coefficient, suggesting that noise and error outweigh biological signal under
this combination of species and the R2T branch length method. These results highlight that internal
biological validation will play an important role in interpreting ERC-based interaction networks.

Overlap in interactions predicted across multiple datasets:

In addition to understanding how different datasets and parameters affect the number of ERC hits
produced by *ERCnet*, it is also important to understand the consistency with which individual
protein-protein interactions are predicted across different datasets. Therefore, we compared the
overlap of specific *ERCnet* hits (i.e. pairs of proteins) across different taxon sampling datasets
(Fig. 6). We found that there were no ERC hits that persisted across all 20 of the R2T and BXB
ERCnet runs. More than half of the ERC hits runs were non-overlapping 'singletons' (i.e. found
in a single *ERCnet* run); however, there were hits with a degree overlap was as high as 19-way
overlap and 15-way overlap among R2T and BXB *ERCnet* runs, respectively. We might expect a
background level of overlap, even if ERC hits were driven entirely by noise, so we generated

randomized replicates in which the identities of the interaction partners were randomly swapped
within each *ERCnet* run (Fig. 6, grey bars). We found that the observed level of overlap exceeds a
randomized background, with far more ERC hits occupying rightward tail of the distribution (high
degree of overlap) in our observed data. This result shows that ERC signal rises above background
noise and persists across *ERCnet* runs that use different input data. However, the general
orientation of distributions toward low-degree overlap highlights the importance of species
sampling on *ERCnet* results.

Moreover, our sampling approach provides a unique opportunity to identify pairs of proteins that
show a consensus signature of correlated evolution across *ERCnet* runs. We compiled the ERC
hits from the three highest levels of overlap for R2T and BXB runs (Table 1). This shortlist of
'consensus' ERC hits contains 17 R2T consensus hits and eight BXB consensus hits. Using the
Arabidopsis thaliana homolog from each gene family, we obtained gene names, subcellular
localization information, and descriptions and from *The Arabidopsis Information Resource*
(TAIR) (Lamesch et al. 2012). We found that 8/17 and 5/8 ERC hits show evidence of subcellular
co-localization for R2T and BXB runs, respectively. 9/17 and 2/8 ERC hits show at least some
evidence of co-functionality based on gene descriptions for R2T and BXB runs, respectively.
Taking co-localization and co-functionality information together, we find that 10/17 and 6/8 ERC
hits show evidence of plausible interaction by at least one of these annotation metrics for R2T and
BXB runs, respectively.

A dominant pattern from both R2T and BXB consensus ERC hits is that chloroplast-localized
proteins appear to be especially prevalent. 12/13 ERC hits that exhibit co-localization share
localization in the chloroplast, suggesting that there are fundamental dynamics to chloroplast
biology that drives consistent patterns of ERC across a variety of angiosperm datasets. Among the
R2T consensus hits, members of the plastid caseinolytic protease (Clp) complex are especially
prevalent; we observe two ERC hits among members of the Clp complex *sensu stricto* as well as
three ERC hits between a Clp protein and a plastid ribosomal protein. These results are consistent
with prior results, in which R2T and related methods revealed strong ERC among Clp and
ribosomal proteins (Forsythe et al. 2021; Gatts et al. 2024). However, it is notable that previous
studies specifically sampled species known *a priori* to exhibit accelerated plastid genome
evolution, whereas the species sampled here chosen without regard to plastid evolutionary rates.
Therefore, the results shown here provide the new insight that ERC signatures spanning the plastid
proteostasis apparatus are prevalent in a more general and widespread sense in angiosperms.
Similar to R2T consensus hits, BXB consensus hits appear to be enriched for organelle-localized
proteins. However, they otherwise lack a clear pattern of functional enrichment, instead touching
on several areas of plant cellular biology such as cellular signaling, the cell cycle, chloroplast and
mitochondrial organization, and retrograde signaling.

Parallel processing and computational runtime

Scaling ERC analyses to whole proteomes involves computationally intensive statistical steps that
are best suited for high-performance computing (HPC). *ERCnet* provided built-in and
customizable parallelization at the most computationally intensive steps, allowing users to tailor
ERCnet to perform optimally in a variety of computing environments. We ran *ERCnet* on our test
datasets on a server, utilizing 48 parallel threads (Fig 7). Through our early testing of *ERCnet*, we
found that the pairwise ERC analysis steps (implemented with ERC_analyses.py) suffered a
performance deficit when run on more than 24 parallel threads, perhaps due to a parallel computing
phenomenon known as 'thrashing', in which increased resource allocation comes at an overhead

cost for the operating system to schedule, resulting in reduced efficiency. Therefore, we run this step using only 24 threads. Total runtimes for full *ERCnet* analyses ranged from ~12-43 hours. We find that all-by-all ERC step constitutes a larger portion of the total runtime step at *n*=10, whereas the phylogenomic analyses step rules the runtime at larger *n*-values (Fig. 7A). Somewhat counterintuitively, *n*=10 runs exhibited some of the longest total runtimes. These patterns are likely a product of the complex relationship between proteome coverage after filters, size of individual genes trees, and the fact that processing in the phylogenomics step scale linearly with number of genes in the dataset, while all-by-all ERC analyses is proportional to the square of the number of genes. We tested the performance of the phylogenomics steps in a highly parallel versus modestly parallel architecture and found that 48-thread parallelization decreased runtime 4.2-fold on average compared to four threads (Fig. 7B). While the maximum number of species included in our test datasets (25 ingroup species) is modest in comparison to some implementations in non-plant taxa (Steenwyk et al. 2022; Tao et al. 2024), the built-in parallelization features of *ERCnet* show potential to help enable high-throughput analyses in reasonable timeframes with modest computational resources.

Discussion:

Assessing significance filters and error rates with simulations of protein evolution

Our simulations revealed *ERCnet* can achieve low rates of false positive and negative error. Using the most conservative filters, *ERCnet* achieved false positive rates below 10% in most cases for R2T and below 1% in most cases for BXB, both of which are an encouraging indication that *ERCnet* will be a useful and reliable predictive tool. However, it should be noted that our simulations were designed for parameter exploration and proof-of-principle and likely do not capture important sources of noise that impact real datasets. Developing highly realistic simulations and incorporating gold-standard plant interaction data into ERC-based networks represents an important area of future research for the field.

Our simulations revealed that our BXB method is a relatively conservative approach. This is desirable in applications in which researchers seek high-confidence interactions; however, our results demonstrate that BXB combined with the Spearman correlation method produces false negative rates approaching 100%, meaning BXB should not be combined with the Spearman method. Taken together, our results outline a useful roadmap to help users select *ERCnet* options to tailor the stringency of significance thresholds to their experimental priorities.

Optimized design for ERC analyses

The choice of how many taxa to include is typically made in the early stages of planning an ERC study, yet the field lacks clear guidance in this area. A wide range of taxon sample sizes have been used in past studies, ranging from 18 budding yeast species (Clark et al. 2012) to 472 insect species (Tao et al. 2024). To our knowledge, past choices are mainly driven by the availability of publicly available genome/proteome sequences and are likely aimed at including the maximum possible number of taxa under the rationale that this is favorable because it increases statistical power. The datasets tested here represent only a modest range in size and extend to smaller numbers of species than have typically been used in ERC, yet we were able to detect patterns across this range that draw the 'more is more' rationale into question. Most notably, we demonstrate a tradeoff between the proportion of the proteome that passes filters as suitable to study via ERC and the potential

statistical power of those ERC analyses gained in large datasets (Fig. 4). Our gene family filtering criteria are more lenient than the criteria typically applied in ERC studies, but even so, our largest datasets (25 ingroup species), restricted the number of gene families passing filters to <7,000 proteins (Fig. 4A), which contain only ~30% of Arabidopsis proteins. This suggests that there is at least some benefit to limiting sampling sets to a modest number of species. Consistent with this idea, we did not observe a clear impact of dataset size on the functional clustering of ERC networks, with datasets as small as n=10 showing significant clustering (although with a larger degree of variance across replicate sample datasets) (Fig. 5). Indeed, branch-length measurement strategy (R2T vs BXB) may have a largest influence on *ERCnet* performance, with notable differences in error rate on simulated data (Figs. 2 and S1), network size (Fig. 5), and network overlap (Fig. 6 and Table 1). The cases in which BXB showed a substantial (Figs. 2 and S1) or incremental (Fig 5 C-D) improvement over R2T may be owed to the additional internal branch length datapoints made available in the BXB method or by the ability to reduce the statistical non-independence/pseudo-replication problem inherent to the R2T method (see Results). In any case, our results suggest that combining a moderately sized dataset with the BXB branch length method could present an optimal balance of proteome coverage and statistical power.

It should be noted that the test set of species included here were selected based on model organism status and phylogenetic representation, but not based on organismal biology or ecology features. This is a major difference from some prior studies (Williams et al. 2019; Forsythe et al. 2021) that have incorporated *a priori* knowledge of important transitions in evolutionary rates into taxon sampling decisions. The use of ERC to explore functional modules that coevolve during specific evolutionary transitions (Gatts et al. 2024) or to explore how rate variation interacts with complex categorical (Redlich et al. 2023) or quantitative trait evolution (Kowalczyk et al. 2019) is an exciting new direction (Hu et al. 2019; Treaster et al. 2023; Yan et al. 2023). In such cases, it may make sense to prioritize sampling species that represent important evolutionary transitions or trait values. Although we have not explored this dimension of taxon sampling here, it is possible that *a priori* information about phenotypic and evolutionary rate variation could alter the dynamics of how dataset size influences ERC analyses.

Dataset dependence and consensus signatures of ERC across angiosperm datasets

There are countless biological and technical factors that could lead to a given ERC hit not being detected in a given sampling set. The highly dimensional nature of all-by-all ERC analyses mean that differences in genome quality and organismal biology can become amplified, leading to vastly inconsistent ERC-based networks. These factors could explain why the ERC hits produced by different runs of *ERCnet* show only a modest degree of overlap (Fig 6). While the level of overlap we observed clearly exceeds a random background, the observation that most ERC hits were found in only a single run raises concerns about the reproducibility of ERC results. On the other hand, the lack of overlapping hits may suggest that ERC signatures reflect the specific biology of the taxa included in an *ERCnet* analysis. This demonstrates that applying ERC in unique sets of taxa has the potential to identify novel interactions. Our results also highlight that ERC does not return all protein-protein interactions in the cell. Instead, ERC is equipped to return the interactions that have been influenced by variable selection pressures across the chosen taxa.

Given the dataset dependence of many ERC hits, it could be especially informative to identify the shortlist of ERC hits that are detected across multiple datasets and *ERCnet* runs because the identity of the proteins involved in these interactions could reveal fundamental drivers of plant

molecular evolution. We find that interactions among chloroplast-localized proteins constitute a large portion of our consensus ERC hits (Table 1), including interactions among functional categories that have been documented in previous plastid-nuclear ERC work. However, the prevalence of chloroplast-related proteins in this study is especially novel and striking because the taxa used here were not selected with chloroplast function in mind. Despite this (and perhaps because of this), our consensus ERC hits revealed several new interactions that had not been identified in prior plastid-nuclear focused ERC analyses. These hits include proteins with little functional information on TAIR, meaning understanding these interactions could lead to novel insights into chloroplast biology. Beyond the novel predictions of gene function provided by our consensus ERC hits, our results raise the question: why do chloroplast-localized proteins exhibit ERC to such an exceptional degree? The answer to this question remains elusive and is outside of the scope of this study, but perhaps the consensus approach that is made possible by *ERCnet* will lay the groundwork for addressing this type of question.

In general, our consensus ERC results resemble prior attempts to identify the 'strongest' set of ERC hits (Forsythe et al. 2021; Gatts et al. 2024) in that our shortlist is a 'mixed bag' that includes some extremely intuitive predicted interactions but also pairs of proteins that seemingly lack a clear functional connection, at least based on summary information from TAIR. For example, we predict an interaction between *SHOT1-binding ATPase 4* and *SHOT1-binding ATPase 2*, which share identical TAIR descriptions, supporting co-functionality. On the other hand, our ERC hit connecting *Fibrillin10* (located in the nucleus and chloroplast; involved in structural activity) with *Ribosomal Protein UL15M* (located in the mitochondrion) does not inherently suggest co-functionality at first glance; however, this ERC hit could represent a functional pathways that spans compartments, such as proteostasis or retrograde signaling, which ERC may be adept at detecting (Forsythe et al. 2021; Gatts et al. 2024; Little et al. 2024). In any case, understanding which of these 'surprising' ERC hits constitute novel functional insights versus noise remains an ongoing challenge to interpretating ERC results; however, our approach of identifying ERC hits spanning randomly sampled datasets could present a valuable means of pinpointing the groups of proteins whose interaction and coevolution are indicative of the most widespread drivers of intermolecular co-evolution in plants.

Materials and methods

Proteome simulations

We used the python module, *pyvolve* (v. 0.9.0), to simulate protein evolution of 1000 gene families and combined the resulting protein sequences into proteomes for 21 species (Spielman and Wilke 2015). To simulate each gene family, first we generated a randomly branching bifurcating tree with each branch length randomly sampled from a distribution between 0.01 and 0.05. Trees generated in this way were used as guide trees to simulate protein sequences using *pyvovle*. Protein sequence lengths were randomly set to between 200 and 1000 amino acids. All proteins with simulated with the "jtt" model of evolution. We designated tree 1-100 as our 'co-acceleration' trees, which were used to simulate protein sequences that underwent accelerated evolution along specific branches of the tree. We simulated this acceleration by randomly selecting five branches on the tree and multiplying the branch length by ten. Protein families 1-100 were simulated using

this accelerated tree as guide tree. Protein families 101-1000 were simulated with the original un-accelerated guide tree. After all protein families were simulated, the protein family sequences were deconstructed and moved into 21 proteome files, which were input into the standard *ERCnet* workflow (Fig. 2). For this dataset, a true-positive ERC hit was defined as a significant ERC hit in which both interaction partners were from protein families 1-100. The false positive rate was calculated as the number of significant ERC hits in which at least one interaction partner was from protein family 101-1000 divided by the total number of comparisons of that type. The false negative rate was defined was 1 minus the number of true-positives divided by the total number comparisons between pairs of proteins families from 1-100.

As a separate simulation-based test, we simulated a separate set of proteomes with a critical difference being that the five accelerated branches were randomly chosen separately for each of the 100 accelerated gene trees. While this dataset still includes 100 accelerated protein families, each is accelerated in a different set of branches, meaning they are not 'co-accelerated', thus serving as a negative control for *ERCnet* detection of correlated evolution (Fig. S1). In this dataset, any significant ERC hit was considered a 'spurious' ERC hit.

All simulations used in this study can be reproduced with the *Simulations.py* script. Calculations of false positive/negative rates can be reproduced with the *Sim_error_track.py* script. Both scripts are available on the *ERCnet* GitHub page.

Obtaining and processing randomized plant proteome datasets

We created a pool of 35 plant proteome sequence dataset (Table S1), which included a mixture of crop species and model organisms, chosen for their general high-quality genomic resources. Sequences were obtained from the *Phytozome* plant comparative genomics portal (release 13) (Goodstein et al. 2012). We downloaded the 'primary transcripts only' version fasta files for each species to avoid splice isoforms being mistaken for paralogs. Fasta files were reformatted to standardize sequence ID formatting (e.g. "Atha__AT1G06950" for an *Arabidopsis thaliana* protein sequence), using *SeqKit* Unix tools (Shen et al. 2016).

From our pool of 35 plant genomes, we randomly selected species for datasets of size $n=10$, $n=15$, $n=20$, and $n=25$, where n refers to the number of ingroup species included. We performed five replicates for each n value, each with its own randomly selected set of taxa. All datasets included *Amborella trichopoda* as an outgroup and *Arabidopsis thaliana* as a common focal species for cross-dataset comparisons. Three replicates of randomly selected taxa were performed for each n value, resulting in a total of 20 datasets, each of which was subjected to full *ERCnet* analyses.

ERCnet analyses require clustered gene families as input, which we obtained by running *Orthofinder* (v2.2.5) (Emms and Kelly 2015, 2019). We ran *Orthofinder* separately on each test dataset using the following command.

```
orthofinder -f <path/to/dir/containing/proteomes/> -y -X -M msa -t <number of
threads available on computing system>
```

Orthofinder species tree outputs were manually inspected to verify that *Amborella trichopoda* was the outgroup. When this was not the case, we re-ran *Orthofinder* with a constrained species tree consistent with established species relationships (Jansen et al. 2007). *Orthofinder* "hierarchical orthogroups' (HOGs) for each dataset were used as input for each *ERCnet* run.

Setting up computing environment and installing ERCnet dependencies

ERCnet has been tested extensively on multiple MacOS and Linux-based computers. For this study, we ran all *ERCnet* analyses on a node on a HPC Linux server equipped with dual AMD EPYC 7713 processors, each featuring 64 physical cores with a base clock speed of 2.0 GHz, for a combined total of 128 physical cores. The system includes 1 TB (16x64 GB) of PC4-25600 3200 MHz DDR4 ECC RDIMM RAM, two 240 GB Micron 5400 PRO Series SATA SSDs for the operating system, and a 256 TB storage capacity. We created two *Conda* virtual environments to accomplish *ERCnet* steps that require *Python 3* and *Python 2*. Our environments utilized *Python* 3.9.7 and *Python* 2.7.18, respectively. The *Conda* environments, including *ERCnet* dependencies, can be recreated using the .yml files provided with the *ERCnet* package on *GitHub*. All *ERCnet* runs presented in this study were performed using *ERCnet* (v1.2.0) (see *ERCnet* documentation for release information).

Phylogenomic analyses

The first steps of *ERCnet* are implemented in the *Phylongenomics.py* script, which filters gene families by user defined parameters (see -r and -p flag below). Next, multiple sequence alignments are generated using the L-INS-I algorithm in *MAFFT* (v7.515) (Katoh and Standley 2013). Next, *TAPER* (Zhang et al. 2021) is (optionally) used to identify outlier sequences and filter out noise from heterogeneity. *GBLOCKS (v0.91b)* (Castresana 2000) is then used to trim poorly aligned regions from alignments. Alignments that are shorter than 100 amino acids (set with the '-l 100' flag) following trimming are discarded. Below is an example command used to call Phylogenomics.py, including descriptions of all required arguments. For detailed descriptions of optional arguments, see the *ERCnet* GitHub page.

```
./Phylogenomics.py -j <Job Name> -o <OrthoFinder Results Files> -m <Threads> -
s -p 4 -r 11 -T <Julia Environment Path> -n 1
```

The -j flag indicates the desired name for the output folder for all *ERCnet* outputs. This 'jobname' is used to help *ERCnet* find needed input/output files throughout the *ERCnet* workflow. The -o flag indicates the path to *Orthofinder* results used in the analyses. The -m flag allows for parallel multithreading to speed up analysis time by using *Joblib* (see below). The -s flag allows the user to provide a custom species mapping file to help *ERCnet* assign gene names to species. The -p and -r flags are used during an initial filtering step to remove gene families with too many paralogs or too little species representation, which are unlikely to be useful in *ERCnet* analyses (e.g. gene families consisting of only 100 paralogs from one species). The -p flag indicates the maximum number of paralogs per species allowed in each gene family. For example, "-p 3" would cause *ERCnet* to exclude a gene family that contains four or more paralogs for any species. Note that *Orthofinder's* HOGs (Emms and Kelly 2019) split deep paralogs into separate gene families, so the paralogs detected during this filtering step represent only relatively recent gene duplicates. We set -p to 3 for all *ERCnet* runs used in this study. The -r flag controls the minimum number of species represented in each gene family. For example, "-r 6" indicates that at least six unique ingroup species must be present in a gene family in order for *ERCnet* to retain the gene family. For the randomized datasets of varying size used in this study, we used set -r to the ceiling of (n/2)+1. For example, for n=20 dataset, r was set to 11. The -T flag is (optionally) used to tell *ERCnet* where to find the *Julia* installation used to run *TAPER*. The -n flag allows the user to specify the species tree node from which to obtain *Orthofinder* HOG subtrees (Emms and Kelly 2019).

To achieve parallelization for the steps that represent potential computational bottlenecks (Fig. 1),
we use *Joblib*, an open-source *Python* library that uses *C* and *loky* to circumvent *Python's* Global
Interpreter Lock and spawn parallel worker processes.

Gene-tree/species-tree reconciliation

Once gene trees are generated for all retained gene families, *ERCnet* performs a GT/ST
reconciliation step via the *GTST_reconciliation.py* script. GTST_reconciliation.py employs
DLCpar (v2.0.1 Kellis) to infer gene duplication, coalescence, and loss, which allows *ERCnet* to
reconcile conflicting gene trees and species and perform BLR (See Results and Fig 1B). While
ERCnet generally runs inside a *Python 3* environment, *DLCpar* requires *Python 2*. We use *Conda*
environments to easily toggle between *Python 2* and *Python 3*. Once a *Python* 2 environment is
activated, *GTST_reconciliation.py* can be called with the following command:

```
./GTST_reconciliation.py -j <Job Name>
```

536 The -j flag requires the same job name from the previous step to continue the analysis.

537 *ERC analyses with branch length reconciliation*

538 After GT/ST reconciliation, *ERCnet* performs ERC analyses in an all-vs-all fashion by comparing
539 the branch lengths for every possible pairwise combination of gene trees. To account for lineage-
540 specific differences in rate of evolution, each branch length on each gene tree is normalized by the
541 genome-wide average for the corresponding branch before correlation analyses are performed.
542 Below is an example script used to call ERC_analysis.py.

543 ```
./ERC_analyses.py -j <Job Name> -m <Threads> -s <Focal Species> -b <Branch
544 Reconciliation Method>
```

The -j flag continues to use the same job name from previous steps. The -m flag determines how
many threads used for parallelizing the analysis. The -s flag indicates which species should be the
focal species for the analysis. The -b flag controls which branch length correlation method (R2T
or BXB) is used. For this study, we ran *ERCnet* with both methods in order to compare results.

*Network analyses*

The final step of *ERCnet* filters ERC results to retain ERC hits according to user-defined cutoffs
and filtering criteria. *Network_analysis.py* includes options to filter results by R-squared values
and p-values obtained from Pearson and/or Spearman correlation analyses. Users choose to filter
by raw p-value or p-values that are multiple-test corrected with the Benjamini-Hochberg false
discovery rate (FDR) method (Benjamini and Hochberg 1995). The script also includes options
for building network graphs to visualize interaction networks. Nodes indicate gene families and
edges between nodes indicate significant ERC signal. Many options for network visualization,
clustering, and export are available (see *ERCnet* GitHub page). Additionally, *Network_analysis.py*
provides the option to evaluate clustering of user-defined functional gene annotations by
calculating the assortativity coefficient (Newman 2003) using the *igraph* package (Csardi and
Nepusz 2006). *Network_analyses.py* was run with the following command:

```
./Network_analyses.py -j <Job Name> -m <Branch method used>  -y <Clustering
562  algorithm> -s <Focal Species> -f <ERC results filename>
```

563 The -j flag continues the same job name from previous steps. The -m flag informs the script which
564 branch length method was used from the previous step. The -f flag indicates which ERC results
565 file should be analyzed in the event that prior step was run multiple times. The -y flag allows the
566 user to specify which clustering method they would like for generating network plots. The -s flag
567 specifics which focal species the user is providing. The -f flag provides the script with the ERC
568 results tsv file the user wishes to use for this portion of the analysis. Multiple ERC results files can
569 be generated by *ERC_analysis.py*.

570 *Analysis of ERC hit overlap*

571 To ask if ERC hits from multiple *ERCnet* runs overlap more than would be expected by chance,
572 we performed a permutation test in which we randomized the ERC results table. The ERC results
573 table includes a "Gene A" and "Gene B" corresponding to the two gene families being compared
574 for ERC signature in a given pairwise combination. First, to gauge overall overlap between
575 different *ERCnet* runs, we simply counted the number of *ERCnet* runs in which a given Gene A
576 and Gene B combination are present among the ERC hits. To achieve a 'null distribution', we
577 randomized the Gene B column of the tables and again calculated ERC hit overlap across *ERCnet*
578 runs. We replicated this procedure ten times to get an average degree of overlap for *ERCnet* runs
579 with different parameters. To analyze consensus ERC hits, we extracted the ERC hits present in
580 the top three levels of overlap for R2T and BXB *ERCnet* runs (Table 1) and obtained functional
581 annotation data for the *Arabidopsis thaliana* paralog in each gene family from TAIR (Lamesch et
582 al. 2012).

584 **Works Cited**
585 **Asar Y, Sauquet H, and Ho SYW**. Evaluating the Accuracy of Methods for Detecting Correlated
586   Rates of Molecular and Morphological Evolution. Syst Biol. 2023:**72**(6):1337–1356.
587   https://doi.org/10.1093/sysbio/syad055
588 **Benjamini Y and Hochberg Y**. Controlling the False Discovery Rate : A Practical and Powerful
589   Approach to Multiple Testing. Royal Statistical Society. 1995:**57**(1):289–300.
590 **Castresana J**. Selection of conserved blocks from multiple alignments for their use in
591   phylogenetic analysis. Mol Biol Evol. 2000:**17**(4):540–552.
592   https://doi.org/10.1093/oxfordjournals.molbev.a026334
593 **Clark NL, Alani E, and Aquadro CF**. Evolutionary rate covariation reveals shared functionality
594   and coexpression of genes. Genome Res. 2012:**22**(607):714–720.
595   https://doi.org/10.1101/gr.132647.111.Guerois
596 **Clark NL and Aquadro CF**. A novel method to detect proteins evolving at correlated rates:
597   Identifying new functional relationships between coevolving proteins. Mol Biol Evol.
598   2010:**27**(5):1152–1161. https://doi.org/10.1093/molbev/msp324
599 **Comte N, Morel B, Hasic D, Gueguen L, Boussau B, Daubin V, Penel S, Scornavacca C, Gouy M,**
600   **Stamatakis A, et al.** Treerecs: An integrated phylogenetic tool, from sequences to
601   reconciliations. Bioinformatics. 2020:**36**(18):4822–4824.
602   https://doi.org/10.1093/bioinformatics/btaa615
603 **Csardi G and Nepusz T**. The igraph software package for complex network research. InterJournal
604   Complex Systems. 2006:**Complex Sy**(1695):1695.

Degnan JH and Rosenberg NA. Discordance of species trees with their most likely gene trees: A unifying principle. Mol Biol Evol. 2013:**30**(12):2709–2713. https://doi.org/10.1093/molbev/mst160

Duchêne DA, Duchêne S, Stiller J, Heller R, and Ho SYW. ClockstaRX: Testing Molecular Clock Hypotheses with Genomic Data. Genome Biol Evol. 2024:**16**(4). https://doi.org/10.1093/gbe/evae064

Emms DM and Kelly S. OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. Genome Biol. 2015:**16**(1):157. https://doi.org/10.1186/s13059-015-0721-2

Emms DM and Kelly S. OrthoFinder: Phylogenetic orthology inference for comparative genomics. Genome Biol. 2019:**20**(1):1–14. https://doi.org/10.1186/s13059-019-1832-y

Forsythe ES, Nelson ADL, and Beilstein MA. Biased gene retention in the face of introgression obscures species relationships. Genome Biol Evol. 2020:**12**(9):1646–1663. https://doi.org/10.1093/GBE/EVAA149

Forsythe ES, Williams AM, and Sloan DB. Genome-wide signatures of plastid-nuclear coevolution point to repeated perturbations of plastid proteostasis systems across angiosperms. Plant Cell. 2021:1–18. https://doi.org/10.1093/plcell/koab021

Gatts TC, Rehmann EA, Sloan DB, and Forsythe ES. Correlated evolutionary rates reveal novel components and cross-compartment connectivity in plant proteostasis systems. 2024. https://doi.org/10.1101/2024.08.22.609246

Goh CS, Bogan AA, Joachimiak M, Walther D, and Cohen FE. Co-evolution of proteins with their interaction partners. J Mol Biol. 2000:**299**(2):283–293. https://doi.org/10.1006/jmbi.2000.3732

Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U, Putnam N, et al. Phytozome: A comparative platform for green plant genomics. Nucleic Acids Res. 2012:**40**(D1):1178–1186. https://doi.org/10.1093/nar/gkr944

Hahn MW and Nakhleh L. Irrational exuberance for resolved species trees. The Society for the Study of Evolution. 2015:**70**(1):7–17. https://doi.org/10.1111/evo.12832

Hu Z, Sackton TB, Edwards S V., and Liu JS. Bayesian Detection of Convergent Rate Changes of Conserved Noncoding Elements on Phylogenetic Trees. Mol Biol Evol. 2019:**36**(5):1086–1100. https://doi.org/10.1093/molbev/msz049

Jacob L. Steenwyk, Megan A. Phillips, Feng Yang, Swapneeta S. Date, Todd R. Graham, Judith Berman, Chris Todd Hittinger, and Antonis Rokas. An orthologous gene coevolution network provides insight into eukaryotic cellular and genomic structure and function. Sci Adv. 2022.

Jansen RK, Cai Z, Raubeson LA, Daniell H, Depamphilis CW, Leebens-Mack J, Müller KF, Guisinger-Bellian M, Haberle RC, Hansen AK, et al. Analysis of 81 genes from 64 plastid genomes resolves relationships in angiosperms and identifies genome-scale evolutionary patterns. Proc Natl Acad Sci U S A. 2007:**104**(49):19369–19374. https://doi.org/10.1073/pnas.0709121104

De Juan D, Pazos F, and Valencia A. Emerging methods in protein co-evolution. Nat Rev Genet. 2013:**14**(4):249–261. https://doi.org/10.1038/nrg3414

647 **Katoh K and Standley DM**. MAFFT multiple sequence alignment software version 7:
648 Improvements in performance and usability. Mol Biol Evol. 2013:**30**(4):772–780.
649 https://doi.org/10.1093/molbev/mst010
650 **Kowalczyk A, Meyer WK, Partha R, Mao W, Clark NL, and Chikina M**. RERconverge: An R
651 package for associating evolutionary rates with convergent traits. Bioinformatics.
652 2019:**35**(22):4815–4817. https://doi.org/10.1093/bioinformatics/btz468
653 **Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, Muller R, Dreher K,
654 Alexander DL, Garcia-Hernandez M, et al.** The Arabidopsis Information Resource (TAIR):
655 Improved gene annotation and new tools. Nucleic Acids Res. 2012:**40**(D1):1202–1210.
656 https://doi.org/10.1093/nar/gkr1090
657 **Little J, Chikina M, and Clark NL**. Evolutionary rate covariation is a reliable predictor of co-
658 functional interactions but not necessarily physical interactions. 2024.
659 https://doi.org/10.7554/eLife
660 **Mao L, Van Hemert JL, Dash S, and Dickerson JA**. Arabidopsis gene co-expression network and
661 its functional modules. BMC Bioinformatics. 2009:**10**:1–24. https://doi.org/10.1186/1471-
662 2105-10-346
663 **Newman MEJ**. Mixing patterns in networks. Phys Rev E. 2003:**67**(2):026126.
664 https://doi.org/10.1103/PhysRevE.67.026126
665 **Panchy N, Lehti-Shiu M, and Shiu SH**. Evolution of gene duplication in plants. Plant Physiol.
666 2016:**171**(4):2294–2316. https://doi.org/10.1104/pp.16.00523
667 **Priedigkeit N, Wolfe N, and Clark NL**. Evolutionary Signatures amongst Disease Genes Permit
668 Novel Methods for Gene Prioritization and Construction of Informative Gene-Based
669 Networks. PLoS Genet. 2015:**11**(2):1–17. https://doi.org/10.1371/journal.pgen.1004967
670 **Ramani AK and Marcotte EM**. Exploiting the co-evolution of interacting proteins to discover
671 interaction specificity. J Mol Biol. 2003:**327**(1):273–284. https://doi.org/10.1016/S0022-
672 2836(03)00114-1
673 **Rao X and Dixon RA**. Co-expression networks for plant biology: Why and how. Acta Biochim
674 Biophys Sin (Shanghai). 2019:**51**(10):981–988. https://doi.org/10.1093/abbs/gmz080
675 **Redlich R, Kowalczyk A, Tene M, Sestili HH, Foley K, Saputra E, Clark N, Chikina M, Meyer WK,
676 and Pfenning A**. RERconverge Expansion: Using Relative Evolutionary Rates to Study
677 Complex Categorical Trait Evolution. Biorxiv. 2023.
678 https://doi.org/10.1101/2023.12.06.570425
679 **Rei Liao J-Y, Friso G, Forsythe ES, Michel EJS, Williams AM, Boguraev SS, Ponnala L, Sloan DB,
680 and van Wijk KJ**. Proteomics, phylogenetics, and co-expression analyses indicate novel
681 interactions in the plastid CLP chaperone-protease system. Journal of Biological Chemistry.
682 2022:101609. https://doi.org/10.1016/j.jbc.2022.101609
683 **Sato T, Yamanishi Y, Kanehisa M, and Toh H**. The inference of protein-protein interactions by co-
684 evolutionary analysis is improved by excluding the information about the phylogenetic
685 relationships. Bioinformatics. 2005:**21**(17):3482–3489.
686 https://doi.org/10.1093/bioinformatics/bti564
687 **Shen W, Le S, Li Y, and Hu F**. SeqKit: A cross-platform and ultrafast toolkit for FASTA/Q file
688 manipulation. PLoS One. 2016:**11**(10). https://doi.org/10.1371/journal.pone.0163962

689   **Smith CH, Mejia-Trujillo R, and Havird JC**. Mitonuclear compatibility is maintained despite
690   relaxed selection on male mitochondrial DNA in bivalves with doubly uniparental
691   inheritance. Evolution (N Y). 2024. https://doi.org/10.1093/evolut/qpae108

692   **Spielman SJ and Wilke CO**. Pyvolve: A flexible python module for simulating sequences along
693   phylogenies. PLoS One. 2015:**10**(9). https://doi.org/10.1371/journal.pone.0139047

694   **Steenwyk JL, Buida TJ, Labella AL, Li Y, Shen XX, and Rokas A**. PhyKIT: A broadly applicable UNIX
695   shell toolkit for processing and analyzing phylogenomic data. Bioinformatics.
696   2021:**37**(16):2325–2331. https://doi.org/10.1093/bioinformatics/btab096

697   **Steenwyk JL, Phillips MA, Yang F, Date SS, Graham TR, Berman J, Hittinger CT, and Rokas A**. An
698   orthologous gene coevolution network provides insight into eukaryotic cellular and
699   genomic structure and function. https://doi.org/10.1101/2021.07.09.451830

700   **Stolzer M, Lai H, Xu M, Sathaye D, Vernot B, and Durand D**. Inferring duplications, losses,
701   transfers and incomplete lineage sorting with nonbinary species trees. Bioinformatics.
702   2012:**28**(18):i409–i415. https://doi.org/10.1093/bioinformatics/bts386

703   **Tao M, Chen J, Cui C, Xu Y, Xu J, Shi Z, Yun J, Zhang J, Ou G-Z, Liu C, et al.** Identification of a
704   longevity gene through evolutionary rate covariation of insect mito-nuclear genomes. Nat
705   Aging. 2024. https://doi.org/10.1038/s43587-024-00641-z

706   **Treaster S, Deelen J, Daane JM, Murabito J, Karasik D, and Harris MP**. Convergent genomics of
707   longevity in rockfishes highlights the genetics of human life span variation.

708   **Vernot B, Stolzer M, Goldman A, and Durand D**. Reconciliation with non-binary species trees. J
709   Comput Biol. 2008:**15**(8):981–1006. https://doi.org/10.1089/cmb.2008.0092

710   **Wendel JF**. The wondrous cycles of polyploidy in plants. Am J Bot. 2015:**102**(11):1753–1756.
711   https://doi.org/10.3732/ajb.1500320

712   **Williams AM, Friso G, Wijk KJ Van, and Sloan DB**. Extreme variation in rates of evolution in the
713   plastid Clp protease complex. The plan journal. 2019:1–17.
714   https://doi.org/10.1111/tpj.14208

715   **Wolfe NW and Clark NL**. ERC analysis: Web-based inference of gene function via evolutionary
716   rate covariation. Bioinformatics. 2015:**31**(23):3835–3837.
717   https://doi.org/10.1093/bioinformatics/btv454

718   **Wright SN, Colton S, Schaffer L V, Pillich RT, Churas C, Pratt D, and Ideker T**. State of the
719   Interactomes: an evaluation of molecular networks for generating biological insights.
720   bioRxiv. 2024. https://doi.org/10.1101/2024.04.26.587073

721   **Wu Y, Rasmussen MD, Bansal MS, and Kellis M**. Most parsimonious reconciliation in the
722   presence of gene duplication , loss , and deep coalescence using labeled coalescent trees.
723   2014:475–486. https://doi.org/10.1101/gr.161968.113.Freely

724   **Yan H, Hu Z, Thomas GWC, Edwards S V., Sackton TB, and Liu JS**. PhyloAcc-GT: A Bayesian
725   Method for Inferring Patterns of Substitution Rate Shifts on Targeted Lineages Accounting
726   for Gene Tree Discordance. Mol Biol Evol. 2023:**40**(9).
727   https://doi.org/10.1093/molbev/msad195

728   **Yan Z, Ye G, and Werren JH**. Evolutionary Rate Correlation between Mitochondrial-Encoded and
729   Mitochondria-Associated Nuclear-Encoded Proteins in Insects. Mol Biol Evol.
730   2019:**36**(5):1022–1036. https://doi.org/10.1093/molbev/msz036

731   **Zhang C, Zhao Y, Braun EL, and Mirarab S**. TAPER: Pinpointing errors in multiple sequence
732   alignments despite varying rates of evolution. Methods Ecol Evol. 2021:**12**(11):2145–2158.
733   https://doi.org/10.1111/2041-210X.13696
734
735
736

737 **Table 1: Consensus ERC hits from angiosperm taxon-sampling datasets.** ERC hits found in
738 the top three categories of overlap for BXB or R2T *ERCnet* runs. (**Meth/Overlap.**) The branch
739 length method and a number of overlapping *ERCnet* runs the hit was found in parentheses.
740 (**AGI[s]**) Presence of Multiple Arabidopsis Gene Identifiers (AGIs) indicates that multiple
741 Arabidopsis paralogs were present in the gene family. In cases of multiple paralogs, we used to
742 first one (underlined) to obtain TAIR information. (**Name and Description**) Gene name and
743 description from TAIR. (**Loc.**) The subcellular localization as indicated by 'Cellular Component'
744 Gene Ontology terms. (**Loc. evid.**) Indicates whether the "Loc." columns show at least some
745 overlap between Gene A and Gene B *(Yes/No/NA)*. (**Desc. evid.**) Qualitative assessment of whether
746 the information in 'Description' columns points to cofunctionality/interaction between Gene A and
747 Gene B *(None, Weak, Strong, Unknown)*. 'Unknown' indicates lack of sufficient information in at
748 least one of the two genes.

Meth./ Overlap.	Gene A				Gene B				Loc. evid.	Desc. evid.
	AGI(s)	Name	Description	Loc.	AGI(s)	Name	Description	Loc.		
R2T (19)	AT1G09130	CLPR3	ATP-dependent caseinolytic (Clp) protease/crotonase family protein	chloro., mito.	AT2G27460	ATSEC23D	Sec23 homolog, forms a distinct clade with SEC23A. Mutants have defects in pollen exine patterning, tapetal development and pollen intine formation	cytoplasm, endoplasmic reticulum exit site	No	None
R2T (18)	AT1G09130	CLPR3	ATP-dependent caseinolytic (Clp) protease/crotonase family protein	chloro., mito.	AT3G44890	RIBOSOMAL PROTEIN BL9C	Plastid ribosomal protein CL9. The mRNA is cell-to-cell mobile	chloro., cytos.	Yes	Strong
R2T (18)	AT1G29340	PLANT U-BOX 17	Encodes a protein containing a UND, a U-box, and an ARM domain. This protein has E3 ubiquitin ligase activity. It is required for cell death and full resistance specified by Arabidopsis RPM1 and RPS4 resistance proteins against Pseudomonas syringae pv tomato. The mRNA is cell-to-cell mobile	chloro.	AT1G68020	TREHALOSE -6-PHOSPHATASE SYNTHASE S6	Encodes an enzyme putatively involved in trehalose biosynthesis. The protein has a trehalose synthase (TPS)-like domain and a trehalose phosphatase (TPP)-like domain. It can complement a yeast mutant lacking both of these activities suggesting that this is a bifunctional enzyme	chloro.	Yes	None
R2T (18)	AT1G14000	VH1-INTERACTING KINASE	Encodes a protein with similarity to members of the C1 subgroup of MAP kinase kinase kinases. Interacts physically with the receptor kinase BRL2/VH1 and appears to be involved in auxin and brassinosteriod signaling. The mRNA is cell-to-cell mobile.	extracell., nuc., vacuole	AT2G27460	ATSEC23D	Sec23 homolog, forms a distinct clade with SEC23A. Mutants have defects in pollen exine patterning, tapetal development and pollen intine formation	cytopl, ER exit site	No	Weak
R2T (18)	AT1G06440	NA	Ubiquitin carboxyl-terminal hydrolase family protein	chloro.	AT5G63100	NA	S-adenosyl-L-methionine-dependent methyltransferases superfamily protein	chloro., mito.	Yes	Weak
R2T (18)	AT3G49730 AT5G65820	NA	Tetratricopeptide repeat (TPR)-like superfamily protein	mito.	AT4G14590	DEFECTIVE IN SNRNA PROCESSING 2, EMBRYO DEFECTIVE 2739	embryo defective 2739	nuc.	No	None
R2T (18)	AT1G11750	CLPP6	One of several nuclear-encoded ClpPs (caseinolytic protease). Contains a highly conserved catalytic triad of Ser-type proteases (Ser-His-Asp)	chloro.	AT3G44890	RIBOSOMAL PROTEIN BL9C	Plastid ribosomal protein CL9. The mRNA is cell-to-cell mobile.	chloro., cytos.	Yes	Strong
R2T (18)	AT1G09130	CLPR3	ATP-dependent caseinolytic (Clp) protease/crotonase family protein	chloro., mito.	AT1G49970	CLPR1	Encodes a ClpP-related sequence. Though similar to ClpP proteins, this does not contains the highly conserved catalytic triad of Ser-type proteases (Ser-His-Asp)	chloro., nuc.,	Yes	Strong
R2T (17)	AT1G49970	CLPR1	Encodes a ClpP-related sequence. Though similar to ClpP proteins, this does not contains the highly conserved catalytic triad of Ser-type proteases (Ser-His-Asp)	chloro., nuc.	AT5G09995	REGULATOR OF FATTY ACID SYNTHESIS 1	Regulates de novo fatty acid synthesis by modulating hetACCase distribution.	chloro.	Yes	Weak
R2T (17)	AT1G09130	CLPR3	ATP-dependent caseinolytic (Clp) protease/crotonase family protein	chloro., mito.	AT1G11750	CLPP6	One of several nuclear-encoded ClpPs (caseinolytic protease). Contains a highly conserved catalytic triad of Ser-type proteases (Ser-His-Asp)	chloro.	Yes	Strong
R2T (17)	AT1G29340	PLANT U-BOX 17	Encodes a protein containing a UND, a U-box, and an ARM domain. This protein has E3 ubiquitin ligase activity. It is required for cell death and full resistance specified by Arabidopsis RPM1 and RPS4 resistance proteins against Pseudomonas syringae pv tomato. The mRNA is cell-to-cell mobile	chloro.	AT2G27460	ATSEC23D	Sec23 homolog, forms a distinct clade with SEC23A. Mutants have defects in pollen exine patterning, tapetal development and pollen intine formation	cytop, ER exit site	No	None
R2T (17)	AT4G14590	DEFECTIVE IN SNRNA PROCESSING 2	embryo defective 2739	Nuc.	AT5G50280	ECD2, EMB1006	Responsible for chloroplast gene expression and group II intron splicing of several genes. Associated with the expression of ribosomal genes and accumulation of chloroplast	chloro.	No	Strong

							ribosomes. Critically important for early chloroplast development in cotyledon			
R2T (17)	AT2G27460	ATSEC23D	Sec23 homolog , forms a distinct clade with SEC23A.Mutants have defects in pollen exine patterning, tapetal development and pollen intine formation.	cytop., ER exit site	AT3G49730 AT5G65820	NA	Tetratricopeptide repeat (TPR)-like superfamily protein;(source:Araport11)	mito.	No	None
R2T (17)	AT1G14000	VH1-INTERACTING KINASE	Encodes a protein with similarity to members of the C1 subgroup of MAP kinase kinase kinases. Interacts physically with the receptor kinase BRL2/VH1 and appears to be involved in auxin and brassinosteriod signaling. The mRNA is cell-to-cell mobile	extracell., nuc., vacuole	AT4G29060	EMBRYO DEFECTIVE 2726	Involved in chloroplast biogenesis and early embryo development. May function as an EF-Ts to regulate plastid translation	chloro., plasma memb.	No	None
R2T (17)	AT1G33340	PICALM8	ENTH/ANTH/VHS superfamily protein	clathrin-coated vesicle	AT1G68020	TREHALOSE-6-PHOSPHATASE SYNTHASE S6	Encodes an enzyme putatively involved in trehalose biosynthesis. The protein has a trehalose synthase (TPS)-like domain and a trehalose phosphatase (TPP)-like domain. It can complement a yeast mutant lacking both of these activities suggesting that this is a bifunctional enzyme	Chloro.	No	None
R2T (17)	AT1G09130	CLPR3	ATP-dependent caseinolytic (Clp) protease/crotonase family protein	chloro., mito.	AT4G17560 AT5G47190	RIBOSOMAL PROTEIN BL19CZ	Ribosomal protein L19 family protein	chloro., cytos.	Yes	Strong
R2T (17)	AT1G09130	CLPR3	ATP-dependent caseinolytic (Clp) protease/crotonase family protein	chloro., mito.	AT3G58800	NA	secretion-regulating guanine nucleotide exchange factor	nuc.	No	None
BXB (15)	AT1G29850	PROGRAMMED CELL DEATH PROTEIN 5	Encodes a protein that by its interaction with HAM acetyltransferases plays an important role during DNA damage responses induced by UV-B radiation and participates in programmed cell death programs	cytos., nuc.	AT5G60340	ADENYLATE KINASE 6	Encodes a nuclear adenylate kinase that interacts with a putative homolog of Rps14, AtRPS14-1 and affects the elongation of cells in the stem	nuc.	Yes	None
BXB (14)	AT4G36580 AT2G18330	SHOT1 BINDING ATPASE 4	Homologue of animal ATPase Family AAA Domain-Containing Protein 3 (ATAD3), which is involved in mitochondrial nucleoid organization; interacts with SHOT1	plasma memb.	AT5G16930 AT3G03060	SHOT1 BINDING ATPASE 2	Homologue of animal ATPase Family AAA Domain-Containing Protein 3 (ATAD3), which is involved in mitochondrial nucleoid organization; interacts with SHOT1	chloro., mito.	No	Strong
BXB (14)	AT1G51110	FIBRILLIN10	PAP/fibrillin (ECM1) localized to chloroplasts; involved in structural activity	chloro., nuc.	AT5G27560	DUF1995 domain protein	NA	chloro.	Yes	Un-known
BXB (14)	AT3G09150	GENOMES UNCOUPLED 3	Required for biosynthesis of the tetrapyrrole phytochrome chromophore phytochromobilin. Encodes phytochromobilin synthase, a ferredoxin-dependent biliverdin reductase. It is necessary for coupling the expression of some nuclear genes to the functional state of the chloroplast	chloro.	AT5G27560	DUF1995 domain protein	NA	chloro.	Yes	Un-known
BXB (13)	AT1G51110	FIBRILLIN10	PAP/fibrillin (ECM1) localized to chloroplasts; involved in structural activity	chloro., nuc.	AT5G64670	RIBOSOMAL PROTEIN UL15M	Ribosomal protein L18e/L15 superfamily protein	mito.	No	None
BXB (13)	AT1G05385	LOW PSII ACCUMULATION 19	Encodes a Psb27 homolog involved in photosystem II biogenesis	chloro., cyto.	AT1G51110	FIBRILLIN 10	PAP/fibrillin (ECM1) localized to chloroplasts; involved in structural activity	chloro., nuc.	Yes	Weak
BXB (13)	AT1G51110	FIBRILLIN10	PAP/fibrillin (ECM1) localized to chloroplasts; involved in structural activity	chloro., nuc.	AT4G27700	NA	Rhodanese/Cell cycle control phosphatase superfamily protein	chloro., cytoskel.	Yes	None
BXB (13)	AT1G10790	NA	hydroxyproline-rich glycoprotein family protein	NA	AT5G25640	NA	Rhomboid-related intramembrane serine protease family protein	NA	NA	Un-known

752 **Fig. 1: *ERCnet* workflow and algorithm development.** (A) Four-step analytical workflow used
753 to analyze *Orthofinder* results (input) and generate an ERC-based interaction network and
754 accompanying summary statistics (output). Stars indicate steps that employ parallel computing.
755 (B) The major analytical steps of the novel 'branch length reconciliation' procedure used to
756 calculate branch lengths on a branch-by-branch basis, using species tree (ST) and gene tree (GT)
757 information. Rounded arrows indicate iterative processes.
758

759
760 **Fig. 2: Simulated protein sequences to assess *ERCnet* error rates.** (A) The random phylogenetic
761 tree used to simulate the background rates of protein evolution. (B) Tree with the same topology
762 but in which five randomly selected branch lengths (bold branches) were multiplied by 10 in order
763 to simulate co-acceleration for 100 of the 1000 protein families. (C-F) False positive/negative error
764 rates of *ERCnet* runs using different branch length methods (R2T vs BXB) as well as different
765 correlation calculation methods (Pearson vs Spearman) and p-value and R^2 cutoffs as significance
766 thresholds. Legend shows the branch length method and correlation methods. "Pearson and
767 Spearman" means that ERC hits were only deemed significant if passing filters according to both
768 methods. (C-D) False positive (C) and negative (D) rates assessed across several p-value cutoff
769 values. R^2 cutoff was held constant at ≥ 0.20. (E-F) False positive (E) and negative (F) rates
770 assessed across several R^2 cutoff values. P-value cutoff was held constant at ≤ 0.05.
771

Fig. 3: Angiosperm taxon-sampling datasets used to assess *ERCnet* performance. (Left) Phylogenetic tree of the full pool of species included for random sub-sampling. (Right) Presence-absence plot indicating the species included in random datasets of each size. Five replicates were performed for each dataset size. *Amborella trichopoda* was included as an outgroup for all replicates and *Arabidopsis thaliana* was included as a common ingroup representative for all replicates.

Fig. 4: Proteome coverage and overlapping branches after *ERCnet* filtering. (A) The number of proteins retained for phylogenomic analysis after several phases of quality-control filtering that occur during the first steps of *ERCnet*. These number represents the number of proteins that are tested for interaction during 'all-vs-all' ERC analyses at later steps of *ERCnet*. (B-C) The number of overlapping branches (i.e. points on correlation plots) among pairs of proteins during all-vs-all ERC analyses. For the root-to-tip method (B), 'branches' refers to paths from root of tree to each tip. For the branch-by-branch method (C), 'branches' refers to the common branches determined by our 'branch length reconciliation' method.

793
794 **Fig. 5: Composition and functional clustering of *ERCnet* interaction networks.** (A-B) The
795 number of nodes (points) and edges (x's) in networks obtained using the root-to-tip (A) and branch-
796 by-branch method (B). Note the log scale for panel A. (C-D) The assortativity coefficient estimated
797 from. Filled points indicate the assortativity coefficient is significantly greater than the randomized
798 null distribution (z-score from randomized permutation test). Significant positive assortativity
799 indicates clustering of traits across a network. The trait measured here was the predicted targeting
800 (plastid, mitochondrial, other) of the proteins in the network.

814
815 **Fig. 6: Overlap in *ERCnet* hits between runs.** (A-B) Bar plots describing the number of hits
816 plotted by the number of *ERCnet* replicate runs they appeared in. Blue bars show observed values,
817 and gray bars show averages from ten randomized replicates. Lines indicate standard error from
818 randomized replicates.

820
821 **Fig. 7: Performance of *ERCnet* using parallel processing** (A) Runtime of the analytical steps of
822 *ERCnet* parallelized with 48 threads. The reconciliation and network analyses steps omitted
823 because they are very fast relative to the phylogenomics steps and pairwise ERC steps and
824 contribute negligibly to the overall runtime. (B) Runtime of the *Phylogenomics* steps of *ERCnet*
825 on either four threads or 48 threads. The highly parallelized 48-thread runs were approximately 4-
826 fold faster than four-thread runs.
827

829 **Table S1: Proteome data used for random sample datasets.**

Species	File name	Common name	Version	Clade
Arabidopsis thaliana	Atha	Thale cress	Araport11	Eudicot
Amborella trichopoda	Atri	NA	v1.0	Outgroup
Ananas comosus	Acom	Pineapple	v3	Monocots
Asparagus officinalis	Aoff	Asparagus	V1.1	Monocots
Brachypodium distachyon	Bdis	Brachypodium	v3.1	Monocots
Dioscorea alata	Dala	Purple yam	v2.1	Monocots
Hordeum vulgare	Hvul	Barley	r1	Monocots
Musa acuminata	Macu	Banana	v1	Monocots
Miscanthus sinensis	Msin	Chinese silver grass	v7.1	Monocots
Oryza sativa	Osat	Rice	v7.0	Monocots
Oropetium thomaeum	Otho	Dessication tolerant grass	v1.0	Monocots
Panicum hallii	Phal	Hall's panic grass	v2.0	Monocots
Panicum virgatum	Pvir	Switchgrass	v5.1	Monocots
Sorghum bicolor	Sbic	Sorghum	v3.1.1	Monocots
Spirodela polyrhiza	Spol	Duckweed	v2	Monocots
Setaria viridis	Svir	Wild foxtail millet	v2.1	Monocots
Triticum aestivum	Taes	Triticum	v2.2	Monocots
Zostera marina	Zmar	Eelgrass	v2.2	Monocots
Zea mays	Zmay	Maize	RefGen_v4	Monocots
Arabidopsis lyrata	Alyr	Sand cress	v2.1	Eudicot
Beta vulgaris	Bvul	Sugar beet	EL10_1.0	Eudicot
Carica papaya	Cpap	Papaya	ASGPBv0.4	Eudicot
Capsella rubella	Crub	Pink sheapard's purse	v1.1	Eudicot
Eutrema salsugineum	Esal	Saltwater cress	v1.0	Eudicot
Populus trichocarpa	Ptri	California poplar	v4.1	Eudicot
Vitis vinifera	Vvin	Common grape vine	v2.1	Eudicot
Fragaria vesca	Fves	Strawberry	v4.0.a2	Eudicot
Eucalyptus grandis	Egra	Ecalyptus	v2.0	Eudicot
Helianthus annuus	Hann	Sunflower	r1.2	Eudicot
Mimulus guttatus	Mgut	Monkeyflower	v2.0	Eudicot
Prunus persica	Pper	Peach	v2.1	Eudicot
Solanum lycopersicum	Slyc	Tomato	TAG2.4	Eudicot
Spinacia oleracea	Sole	Spinach	Spov3	Eudicot
Vigna unguiculata	Vung	Black-eyed pea	v1.2	Eudicot
Lactuca sativa	Lsat	Lettuce	V8	Eudicot

831 **Table S2: Gene families filtered from sample datasets.**

Spec-ies in data-set (n)	Rep-licate	Repres-ent-ation filter cutoff (r)	Para-log filter cutoff (p)	Number of gene families					Proportion of gene families			
				Re-tained	Dis-qualified by r	Dis-qualified by p	Dis-qualified by p and r	Ortho-finder output	Re-tained	Dis-qualified by r	Dis-qualified by p	Dis-qualified by p and r
10	1	6	4	9106	20191	2886	4472	37337	0.24	0.54	0.08	0.12
10	2	6	4	9842	18711	2249	4491	36026	0.27	0.52	0.06	0.12
10	3	6	4	8837	20403	2855	4251	36942	0.24	0.55	0.08	0.12
10	4	6	4	10982	20112	3298	4132	39108	0.28	0.51	0.08	0.11
10	5	6	4	10262	10161	1616	2645	25458	0.40	0.40	0.06	0.10
15	1	9	4	8565	24280	3238	4740	41522	0.21	0.58	0.08	0.11
15	2	9	4	7399	28322	4502	5516	46259	0.16	0.61	0.10	0.12
15	3	9	4	9391	22459	2592	4935	40124	0.23	0.56	0.06	0.12
15	4	9	4	8719	23148	3089	5190	40858	0.21	0.57	0.08	0.13
15	5	9	4	9154	23719	2586	4889	41088	0.22	0.58	0.06	0.12
20	1	11	4	8446	30865	3639	5982	49696	0.17	0.62	0.07	0.12
20	2	11	4	9422	21912	2487	4954	39622	0.24	0.55	0.06	0.13
20	3	11	4	8524	26066	3369	5379	44091	0.19	0.59	0.08	0.12
20	4	11	4	7984	30249	3686	6317	48894	0.16	0.62	0.08	0.13
20	5	11	4	7413	29484	4678	6257	48514	0.15	0.61	0.10	0.13
25	1	14	4	6571	32019	4823	7513	51627	0.13	0.62	0.09	0.15
25	2	14	4	6316	34568	5324	8036	54869	0.12	0.63	0.10	0.15
25	3	14	4	6802	33350	5114	7119	53090	0.13	0.63	0.10	0.13
25	4	14	4	6439	35191	5002	7704	54956	0.12	0.64	0.09	0.14
25	5	14	4	6622	31182	4888	7115	50521	0.13	0.62	0.10	0.14

832
833

834
835
836

837
838 **Fig. S1: Simulations of spurious ERC hits.** Simulations performed under an uncorrelated
839 acceleration model. (A) Unaccelerated tree used as the background species tree. For each protein
840 family simulation, five branches randomly chosen branches were accelerated, leading to
841 uncorrelated patterns of acceleration between different protein families. (B-C) Number of
842 'spurious' ERC hits in networks obtained across different P-value (B) and R^2 value significance
843 filtering cutoffs.

844
845

847
848 **Fig. S2: Network graphs for R2T-based networks.** Network graphs from empirical replicate
849 datasets obtained using the root-to-tip method. Number of ingroup species and replicate number
850 are indicated by "spec" and "rep", respectively.
851
852

854
855 **Fig. S3: Network graphs for BXB-based networks.** Network graphs from empirical replicate
856 datasets obtained using the branch-by-branch method. Number of ingroup species and replicate
857 number are indicated by "spec" and "rep", respectively.
858