TITLE PAGE

PTEN neddylation aggravates CDK4/6 inhibitor resistance in breast cancer

Fan Liu [1,2]†, Weixiao Liu [1,2]†, Yawen Tan [3]†, Yaxin Shang [1], Sihui Ling [1], Xiaokun Jiang [4], Zhen Zhang [4], Shiyao Sun [3], Ping Xie [1,2]*

1 Department of Cell Biology, Laboratory for Clinical Medicine, Capital Medical University, Beijing 100069, China.

2 Beijing lnstitute of Hepatology, Beijing Youan Hospital, Capital Medical University, Beijing 100069, China.

3 Department of Breast and Thyroid Surgery, The Second People's Hospital of Shenzhen, Shenzhen, Guangdong 518035, China.

4 Department of Neurosurgery, Shandong Provincial Hospital affiliated to Shandong First Medical University, Jinan, Shandong 250021, China.

*Correspondence: Ping Xie (xiep@ccmu.edu.cn).

† These authors contributed equally to this work: Fan Liu, Weixiao Liu, Yawen Tan.

COMPETING INTERESTS

The authors declare that there are no competing financial, commercial, or professional interest that could influence the performance, interpretation, or reporting of the research presented in this manuscript. All authors have read and approved this declaration of interests, confirming that it accurately reflects their current situation.

PTEN neddylation aggravates CDK4/6 inhibitor resistance in breast cancer

Fan Liu [1,2]†, Weixiao Liu [1,2]†, Yawen Tan [3]†, Yaxin Shang [1], Sihui Ling [1], Xiaokun Jiang [4], Zhen Zhang [4], Shiyao Sun [3], Ping Xie [1,2]*

1 Department of Cell Biology, Laboratory for Clinical Medicine, Capital Medical University, Beijing 100069, China.

2 Beijing lnstitute of Hepatology, Beijing Youan Hospital, Capital Medical University, Beijing 100069, China.

3 Department of Breast and Thyroid Surgery, The Second People's Hospital of Shenzhen, Shenzhen, Guangdong 518035, China.

4 Department of Neurosurgery, Shandong Provincial Hospital affiliated to Shandong First Medical University, Jinan, Shandong 250021, China.

*Correspondence: Ping Xie (xiep@ccmu.edu.cn).

† These authors contributed equally to this work: Fan Liu, Weixiao Liu, Yawen Tan.

ABSTRACT

The gradual emergence of a novel therapeutic approach lies in the restoration of tumor suppressive machinery. PTEN is a crucial negative regulator of the PI3K/Akt signaling pathway. Protein neddylation modification contributes to PTEN inactivation and fuels breast cancer progression. Here, we highlight that an elevated level of PTEN neddylation is markedly associated with resistance to palbociclib, a CDK4/6 inhibitor used in luminal subtype breast cancer patients. Mechanistically, PTEN neddylation activates the PI3K/Akt signaling pathway, and more notably, upregulates the activity of the AP-1 transcription factor. Our data showed that PTEN neddylation stabilizes JUND, a transcription factor involved in the AP-1 complex, by

disrupting its interaction with the E3 ubiquitin ligase ITCH. Consequently, activated JUND leads to the release of cytokines and chemokines, which in turn may drive an inflammatory tumor microenvironment, potentially contributing to drug resistance. Then, we identified Echinacoside as a potent inhibitor of PTEN neddylation both *in vivo* and *in vitro* by disrupting its interaction with XIAP, the E3 ligase responsible for PTEN neddylation. Combination of Echinacoside effectively overcome resistance to palbociclib in breast cancer treatment. These findings highlight targeting PTEN neddylation as a promising strategy for restoring tumor suppressor activity and overcoming resistance.

INTRODUCTION

Breast cancer is a heterogeneous disease with multiple intrinsic tumor subtypes. The luminal breast cancer subtype, characterized by the expression of hormone receptor-positive (HR+) and human epidermal growth factor receptor 2-negative (HER2-) accounts for 70 % of breast cancer patients [1, 2]. Cyclin D-dependent Kinases 4 and 6 inhibitors (CDK4/6i) have been approved by the FDA for the treatment of luminal breast cancer [3-5]. Although patients with advanced disease have benefited from CDK4/6 inhibitors, drug resistance still remains a significant clinical challenge [6-8]. Many factors contribute to CDK4/6i resistance, including CDK4/6 and Cyclin D1 overexpression, p16 amplification, loss of p53, loss of retinoblastoma (Rb) and over-activation of CDK2. Notably, resistance to CDK4/6 inhibitors is also prominently characterized by over-activated PI3K/Akt signaling [9-13]. However, although PI3K inhibitors could prevent resistance to CDK4/6 inhibitors, they failed to re-sensitize cells once resistance had been acquired [14, 15]. PTEN (phosphatase and tension homologue deleted from chromosome 10) is a well-known negative regulator of the PI3K signaling pathway and governing fundamental cellular processes [16-19]. Recently, PTEN loss is considered as a bona fide mechanism of resistance to CDK4/6 inhibitors. Loss of PTEN increases PI3K/Akt activity and increases activity of CDK4/6 [20-22].

Therefore, targeting the restoration of PTEN tumor suppressor activity is believed to be a more effective approach in tumor treatment [23, 24].

Nedd8, a ubiquitin-like protein covalently conjugated to substrates similar to ubiquitin, mediates protein post-translational modification called neddylation [25]. Neddylation involves the covalent attachment of Nedd8 to a substrate through an ATP-dependent cascade involving the Nedd8-activating enzyme (E1, UBA3/APPBP1), Nedd8-conjugating enzyme (E2, UBE2M and UBE2F), and Nedd8 ligases (E3). The substrates of neddylation include both cullin proteins and non-cullin proteins [26, 27]. Over-active neddylation modification has been indicated to be closely correlated with cancer development. The first-in-class neddylation E1 inhibitor, MLN4924 (pevonedistat), has entered in several phases of clinical trials for tumor therapy [28]. Additionally, several inhibitors targeting E2s have shown well anti-tumor activity in mice models [29]. Up to now, none of them is granted by FDA for the treatment of cancer. It's still necessary to discover more potent and selective neddylation inhibitors for clinical trials. Our previous study found that high levels of PTEN neddylation, which lead to PTEN inactivation, enhance the activity of the PI3K/Akt pathway, promote PTEN nuclear import, and accelerate breast tumor proliferation and invasion [30, 31]. Here, we found that PTEN neddylation is elevated in luminal-type breast cancer patients with resistance to palbociclib by promoting JUND protein stability and AP-1 transcriptional activation. Furthermore, we proposed that Echinacoside (ECH), a natural phenylethanoid glycoside in *Echinacea angustifolia* DC, acts as a potent PTEN neddylation inhibitor. Combined treatment of ECH therapeutically blunted palbociclib resistance and enhanced drug sensitivity. Therefore, we provide an innovative therapeutic strategy to limit cancer drug resistance through PTEN reactivation by blocking neddylation modification.

RESULTS

Resistance to CDK4/6 inhibitor upregulates PTEN neddylation in breast cancer

We previously found that elevated PTEN neddylation in breast cancer correlates with poor clinical outcomes [30], and we confirmed those results here (Fig. 1A, B). We observed that PTEN neddylation was upregulated in all breast cancer subtypes, particularly in luminal-type cancers, while PTEN was downregulated in all subtypes without distinction (Fig. 1C). Detailed pathological information was showed in Supplementary Table 1. Palbociclib, one kind of CDK4/6 inhibitor, has been widely used for the treatment of advanced breast cancer patients. Using the antibody that specifically recognized PTEN neddylation, our results showed that PTEN neddylation was markedly upregulated in palbociclib-resistant breast cancer patients with recurrence *in situ* and liver metastasis, while PTEN remained lowly expressed in drug-resistant patients (Fig. 1D). Pathological data of the patients were provided (Supplementary Fig. 1A). Next, we monitored MMTV-*PyMT* (C57BL/6J background) transgenic mice, which spontaneously develop luminal breast tumors, as a model for late-stage luminal breast cancer [32-34]. Following spontaneous tumor formation, breast tumors exhibited a significant reduction in volume and weight after about 20 days of palbociclib treatment, suggesting that this period corresponded to the drug-sensitive phase of palbociclib. Notably, as the duration of continuous treatment was extended, the tumor volumes and weights ceased to decline further, indicating the emergence of drug resistance. This drug resistance could persist throughout the 50-day treatment period (Fig. 1E-H). Moreover, compared with the low-dose group (75 mg/kg) that developed drug resistance at 20 days, the high-dose group (100 mg/kg) showed that the tumor volume and weight no longer continued to decrease after 25 days, indicating the onset of the drug resistance. Palbociclib-resistant tumors showed no significant difference in size or weight compared to controls, and were marked larger than in the sensitive group at 35-day (Fig. 1I-L).

Rb is a tumor suppressor that is phosphorylated during the transition from G1 to S phase of the cell cycle by CDK4/6. When CDK4/6 inhibitors are ineffective, phosphorylation of the Rb protein would become uncontrolled [35]. Therefore, the increased phosphorylation of the Rb protein in

the mice also indicated the occurrence of drug resistance (Fig. 1M, N input group; Supplementary Fig. 1B). More importantly, high levels of PTEN neddylation were detected in palbociclib-resistant breast tumor tissues of MMTV-*PyMT* mice, while the expression of PTEN remained no obviously changed (Fig. 1M, N and Supplementary Fig. 1B). Additionally, we found that the IC50 of breast tumor cells isolated from palbociclib-resistant MMTV-*PyMT* mice was significantly higher compared to the palbociclib-sensitive group (Fig. 1O), indicating the development of drug resistance. Taken together, our data revealed that PTEN neddylation exhibited elevated levels in palbociclib-resistant breast cancer.

Increased XIAP leads to enhanced PTEN neddylation in palbociclib-resistant breast cancer

To explore the underlying mechanisms of elevated PTEN neddylation in palbociclib-resistant breast cancer, we generated acquired palbociclib-resistant breast cancer stable cell lines (MCF-7-PR). Dose-response curves results showed that palbociclib-resistant cells were more resistant to palbociclib than the parental cells (Fig. 2A). Compared with the parental cells, the palbociclib-resistant cells exhibited accelerated G1/S phase and cell proliferation ability after treatment of palbociclib (Fig. 2B, C). Consistently, PTEN neddylation was upregulated in acquired palbociclib-resistant breast cancer cells (Fig. 2D). Moreover, the ubiquitination of PTEN was unchanged in palbociclib-resistant cells (Supplementary Fig. 1C). Compared to other drug-resistant breast cancer cell lines, such as MCF-7 sublines resistant to doxorubicin (MCF-7/ADR), cisplatin (MCF-7/DDP), and paclitaxel (MCF-7/PTX), PTEN neddylation was specifically increased in palbociclib-resistant breast cancer cells (Fig. 2E). Short-term drug treatment could induce adaptive resistance in cells [13]. We noticed that the phosphorylation of Rb, initially reduced by palbociclib treatment, recovered after prolonged drug exposure, which indicated adaptive resistance was occurred (Fig. 2F). Under the adaptive resistance, PTEN neddylation was increased and could be reduced by MLN4924, which was the inhibitor of neddylation E1 (Fig. 2F, G left panels). For ER negative cells, such as glioma cells T98G, show reduced palbociclib

sensitivity [36]. As expected, no obvious changes in PTEN neddylation or p-Rb were detected

when the T98G cells were treated with palbociclib (Fig. 2G right panels).

Then, we intended to explore the mechanism of high PTEN neddylation in palbociclib-resistant

breast cancer. Notably, our data revealed that XIAP, the E3 ligase for PTEN neddylation [30],

was upregulated in palbociclib-resistant breast cancer cells, while no significant changes were

observed in Nedd8 E1-UBA3, Nedd8 E2-Ubc12, and Nedd8 (Fig. 2H). Consistent with previous

studies showing compensatory CDK2 activation upon resistance elevates XIAP mRNA levels [37,

38], we observed increased XIAP and CDK2 mRNA in both drug-resistant cells and mouse

models (Fig. 2I, 2J). The increased XIAP enhanced the interaction with PTEN in both early

adaptative palbociclib-resistant and acquired palbociclib-resistant breast cancer cells (Fig. 2K-M).

The above data suggested that high PTEN neddylation in palbociclib-resistant breast cancer cells

was likely due to XIAP upregulation and increased XIAP-PTEN interaction.

PTEN neddylation enhances the protein stability of JUND

In previous studies, we have demonstrated that Nedd8 is covalently attached to lysine residues at

positions 197 and 402 of PTEN [30]. To examine the role of PTEN neddylation on CDK4/6i

resistance, we performed RNA-seq in palbociclib-resistant breast cancer cells (Fig. 3A) and

PTEN neddylation deficient cells (2KR, K197 and K402 sites mutated to R) (Fig. 3B). Other than

the PI3K/Akt signaling pathway, which has been believed to be highly associated with CDK4/6i

resistance and PTEN neddylation, we noticed that MAPK, as well as JAK/STAT signaling

pathways were also enriched (Fig. 3A, B). GSEA analysis further indicated that MAPK signaling

pathway obviously was also enriched in both palbociclib-resistant breast cancer cells (Fig. 3C)

and PTEN neddylation deficient cells (Fig. 3D). Similar data were shown in the related dataset

(GSE222367) (Supplementary Fig. 2A). AP-1 (Activator Protein-1) transcription factors are

heterodimers or homodimers composed of JUN and FOS proteins. AP-1 is known to be activated

by MAPK signaling pathways through the phosphorylation of JUN and FOS proteins, and in turn,

AP-1 also regulates the expression of genes involved in MAPK signaling pathways [39]. The reporter gene assays revealed that AP-1 transcription factor activity was significantly increased after palbociclib resistance developed (Fig. 3E). Further heat map data showed that palbociclib resistance led to the upregulation of AP-1 related transcription factors, including *FOSB*, *FOSL*, *c-FOS*, *c-JUN*, *JUND*, and *JUNB* (Fig. 3F and Supplementary Fig. 2B). Conversely, PTEN neddylation deficiency decreased the expression of those genes (Fig. 3G). Then, we detected the protein levels of the AP-1 transcriptional factors and noticed that there was a markable increase of JUND in palbociclib-resistant cells (Fig. 3H). Meanwhile, the activated form of JUND, phosphorylated JUND, was also upregulated when cells were resistant to palbociclib (Fig. 3I). Similar results were also found in early adaptive palbociclib-resistant MCF-7 cells and palbociclib-resistant breast cancer tissues in MMTV-*PyMT* mice, but not in ER-negative T98G cells (Fig. 3J, K). Next, analysis of nuclear and cytoplasmic fractions revealed that, in palbociclib-resistant breast cancer cells, JUND and phosphorylated JUND were significantly elevated, primarily in the nucleus (Fig. 3L). Finally, the results showed that JUND was also markedly upregulated in palbociclib-resistant breast cancer patients (Fig. 3M). These results strongly suggest that the MAPK/AP-1 signaling pathway is a key mechanism of resistance to palbociclib, and PTEN neddylation upregulates JUND protein expression while regulating AP-1 activity.

PTEN neddylation decreases JUND ubiquitination

Next, we intended to explore the mechanism by which PTEN neddylation promotes JUND protein expression, IP analysis indicated that the ubiquitination of JUND was obviously decreased in palbociclib-resistant breast cancer cells (Fig. 4A, B). We generated a fusion protein (PTEN-Nedd8) by linking Nedd8 to the C-terminus of PTEN to mimic neddylated PTEN. Neddylation would promote PTEN nuclear translocation. Therefore, a nuclear localization sequence (NLS) was fused to the N-terminus of PTEN to generate NLS-PTEN for comparison [30, 40, 41]. Then we observed an increased expression level of JUND and phosphorylated JUND in

sh*PTEN*+PTEN-Nedd8 MCF-7 cells, but the opposite effect was found in sh*PTEN*+PTEN-NLS MCF-7 cells (Fig. 4C). Compared with NLS-PTEN, JUND showed a higher affinity for neddylated PTEN (Fig. 4D). Moreover, enhanced interaction between neddylated PTEN and JUND reduced JUND ubiquitination (Fig. 4E). Next, we confirmed that JUND expression was significantly increased in tumor tissues compared with matched adjacent normal tissues, particularly in luminal-type breast cancer (Supplementary Fig. 2C, D). By comparison, the levels of JUND and neddylated PTEN were positively correlated in luminal-type breast cancer (Supplementary Fig. 2E, F). Detailed pathological information was in Supplementary Table 2.

To date, no specific ubiquitin ligase for JUND has been reported. Through UbiBrowser 2.0 [42, 43], ITCH (also known as AIP4) was predicted to be the E3 of JUND with the highest affinity among the interactive protein candidates (Supplementary Fig. 2G). As expected, overexpression of ITCH reduced the levels of JUND protein and its phosphorylation, while *ITCH* knockdown showed opposing effects; however, this effect diminished following the onset of palbociclib resistance (Fig. 4F, G). Overexpression the wild-type of ITCH but not the ligase-deficient ITCH C830A mutant [44] promoted ubiquitination of JUND (Fig. 4H and Supplementary Fig. 2H). Similarly, knockdown of *ITCH* reduced JUND ubiquitination (Fig. 4I). Considering that neddylated PTEN in the nuclear intended to regulate the protein-protein interactions, we first used molecular docking with AlphaFold 2.0 to evaluate whether neddylation of PTEN affects the interaction between ITCH and JUND. The results indicated that neddylated PTEN could occupy the binding region between ITCH and JUND, whereas PTEN without Nedd8 could not (Fig. 4J). We then evaluated the interaction between ITCH and JUND and found that JUND was unable to bind to ITCH in sh*PTEN*+PTEN-Nedd8 MCF-7 cells with PTEN-Nedd8 overexpression (Fig. 4K-M). Furthermore, JUND failed to interact with ITCH in the palbociclib-resistant breast cancer cells compared to the palbociclib-sensitive cells (Fig. 4N). In addition, we noticed that PTEN neddylation increased the mRNA or protein levels of JUND, but not ITCH (Supplementary Fig.

2I). It is known that the proto-oncogene *JUN* is positively autoregulated by its product, JUN/AP-1

[45]. We proposed that PTEN neddylation enhanced JUND protein levels, which established

positive feedback loop on the transcriptional level of *JUND* gene. Collectively, the results

indicate that during drug resistance, elevated PTEN neddylation interferes with the interaction

between JUND and its ubiquitin ligase, reducing JUND ubiquitination and increasing its protein

stability.

Identification of Echinacoside as a PTEN neddylation inhibitor

The above data indicate that PTEN neddylation is a key factor contributing to resistance against

palbociclib treatment. Next, we aimed to explore the possibility of disrupting the interaction

between PTEN and XIAP to reduce PTEN neddylation in tumor cells, and thus overcoming

resistance to palbociclib. Consistent with previous study [30, 46], XIAP was co-

immunoprecipitated with PTEN *in vivo* (Supplementary Fig. 3A). The truncations containing the

BIR2 domain of XIAP and the C2 domain of PTEN interacted with each other (Fig. 5A and

Supplementary Fig. 3B-D). Then, molecular docking and *in vitro* pull-down assay indicated that

Y154 and N234 sites at XIAP, as well as S207 and R335 sites at PTEN were key binding sites on

the interaction surface (Fig. 5B, C). On the basis of the binding determinants of the XIAP-PTEN

interface, structure-based virtual screening was performed to find potential XIAP-PTEN PPI

(protein and protein interaction) inhibitors (Fig. 5D). Ultimately, we selected top five compounds,

guided by the top-ranking binding models and docking scores from the Enamine libraries

containing protein-protein interacting compounds and the Bioactive Compound Library Plus at

Med Chem Express LLC (Supplementary Fig. 3E). No PPI inhibitors affected the protein and

mRNA levels of PTEN (Supplementary Fig. 3F, G). All the five compounds inhibited PTEN

neddylation, with Echinacoside (ECH) and Z219118374 showing the most significant inhibition

effect (Supplementary Fig. 3H). We noticed that Echinacoside (ECH), a natural phenylethanoid

glycoside, had the highest molecular docking score and formed hydrogen bonds with the critical

PTEN residues S207 and R335 that interacted with XIAP (Fig. 5E, F). ECH is one of the main phenylethanol glycosides isolated and purified from *cistanche tubulosa*, which has been used as traditional Chinese herbal medicine with anti-senility and antifatigue effects [47, 48]. Subsequently, we employed surface plasmon resonance and microscale thermophoresis assay to evaluate the interaction between ECH and PTEN. As illustrated in Fig. 5G and 5H, ECH showed a high-affinity binding with PTEN. In addition, TSA assay showed that the inflection temperature (Ti) of GST-PTEN increased after co-incubation with ECH (Supplementary Fig. 3I). Next, we monitored ECH binding with PTEN using the cellular thermal shift assay. The obvious shift in the melting curve indicated increased PTEN protein stability and high-affinity binding upon addition of ECH (Fig. 5I). Furthermore, strengthened XIAP-PTEN interaction was found under the drug resistance (Fig. 5J, K). However, ECH effectively disrupted the interaction between XIAP and PTEN in both palbociclib-sensitive and palbociclib-resistant cells *in vivo.* (Fig. 5J, K and Supplementary Fig. 3J). The pull-down assay revealed a clear dose-dependent decrease in the interaction between XIAP and the PTEN C2 domain following ECH treatment. (Fig. 5L). Moreover, ECH did not affect the dimerization of PTEN (Supplementary Fig. 3K, L).

To determine whether ECH could disrupt PTEN neddylation, we performed *in vivo* and *in vitro* neddylation assays. The results revealed that ECH reduced PTEN neddylation, but not ubiquitination (Fig. 5M). *In vitro* neddylation assay indicated that ECH directly interfered XIAP mediated PTEN neddylation (Fig. 5N). As expected, elevated JUND levels, JUND phosphorylation, and Akt phosphorylation, which increased after palbociclib resistance, were diminished following ECH treatment (Fig. 5O). By inhibiting PTEN neddylation, ECH reduced JUND mRNA levels, protein expression, and phosphorylation in a concentration-dependent manner, while exhibiting no effect on ITCH (Supplementary Fig. 3M, N). Additionally, there was a significant increase in JUND ubiquitination and decrease in AP-1 transcriptional activity following ECH treatment (Fig. 5P, Q). Taken together, we identified Echinacoside as a potent

268   PTEN neddylation inhibitor that eliminates PTEN neddylation by inhibiting the interaction

269   between XIAP and PTEN.

ECH depends on PTEN neddylation to suppress tumorigenesis

Then, we intended to explore the role of ECH in tumorigenesis. ECH exhibited a significant

tumor-suppressive effect, manifested in a dose-dependent manner, on cellular proliferation,

colony formation, and cell migration across various tumor cell lines (Supplementary Fig. 4A-D).

We measured drug toxicity of ECH in CD-1 mice and found no significant differences in body

weight (Supplementary Fig. 5A, B), blood cell counts or composition (Supplementary Fig. 5C), or

serum ALT levels (Supplementary Fig. 5D), indicating no obvious liver injury. Additionally,

ECH treatment showed no signs of toxicity or organ damage based on weight and histopathology

of vital organs (Supplementary Fig. 5E-G). Thus, we speculated that ECH might have substantial

anti-tumor efficacy with minimal adverse effects.

To be note, deletion of *PTEN* significantly weakened the inhibitory effects of ECH on cell

proliferation (Fig. 6A, B) and tumor invasion (Fig. 6C). Dose-response curves results showed that

PTEN neddylation deficient cells were more sensitive to palbociclib than the sh*PTEN*+PTEN-WT

cells (Fig. 6D). Moreover, palbociclib inhibited cell proliferation (Fig. 6E, F) and increased the

cell death (Fig. 6G) in PTEN neddylation deficient cells. However, ECH was unable to inhibit

tumor cell proliferation when PTEN neddylation was deficient (Fig. 6H, I). Additionally, in

PTEN neddylation-deficient cells, ECH no longer reduced the protein levels of JUND and p-

JUND (Fig. 6J), nor did it decrease AP-1 activity (Fig. 6K). Furthermore, CUT&Tag analysis

performed on PTEN neddylation-deficient cells indicated an enrichment of the PI3K/Akt

signaling pathway and the MAPK/AP-1 signaling pathway (Fig. 6L, M). The data demonstrated

that in the absence of PTEN neddylation, JUND exhibited reduced enrichment at the promoters of

downstream genes associated with inflammation and immune responses, including CSF2, IL-17,

IL-6, MMP13, and IL-1β (Fig. 6N). Due to the significant changes in these genes, an

immunoassay array was performed. As shown in Fig. 6O and 6P, the results demonstrated a reduction in multiple cytokines and chemokines, including G-CSF, GM-CSF, IFN-γ, IL-1RA, IL-1β, IL-8, CCL2/4, IL-17, and VEGF, following ECH treatment. However, no alterations were observed in the PTEN neddylation deficient cells subsequent to ECH exposure. Similar results of G-CSF, IL-1β, IL-17 were also approved by ELISA assay (Fig. 6Q). Therefore, the above evidence suggested that ECH inhibited oncogenic effects, at least to some extent, by interfering with PTEN neddylation.

Combination ECH enhances the sensitivity of palbociclib in breast cancer

We next aimed to explore whether ECH could overcome palbociclib resistance in breast cancer. As expected, the tumor weight and volume were both markedly declined in the palbociclib resistance MMTV-*PyMT* mice following the treatment of ECH (Fig. 7A-C). Similar results were also obtained from the primary breast cancer cells isolated from MMTV-*PyMT* mice (Supplementary Fig. 6A, B), and the tumor organoids established from them (Supplementary Fig. 6C-E). In immunoassay arrays, there was an obvious reduction of GM-CSF and G-CSF following ECH treatment (Fig. 6O, P). G-CSF is well known to function as a mobilizing agent for neutrophils, while GM-CSF influences the development and functionality of both neutrophils and macrophages [49-51]. Immunohistochemistry and flow cytometry showed a significant reduction in the population of active neutrophils labeled by Ly6G$^+$, CD11b$^+$, and CD62L$^-$ in palbociclib-resistant MMTV-*PyMT* mice after ECH treatment (Fig. 7D, E). We confirmed a decrease in JUND, phosphorylation of Akt, PTEN neddylation, and XIAP levels, with no notable changes in neddylation of E1-UBA3, E2-Ubc12, and Nedd8 in palbociclib-resistant breast cancer tissues from MMTV-*PyMT* mice after ECH treatment (Supplementary Fig. 6F-I). The combination of ECH with palbociclib showed significantly decreased cell proliferation compared with monotherapy (Fig. 7F). In the ER-negative T98G cells, which was resistance to palbociclib, the co-administration of ECH increased sensitivity to palbociclib, thereby suppressing tumor cell

proliferation (Supplementary Fig. 7A, B). We finally intended to confirm the efficacy of combination therapy using tumor organoid models derived from the human luminal breast cancer patient. Both ECH and palbociclib monotherapies demonstrated tumor growth suppression effects, but the combination treatment exhibited even better anti-tumor growth efficacy compared with the monotherapies (Fig. 7G, H). Immunofluorescence staining showed a significant reduction in PTEN neddylation following ECH treatment in the E-cadherin positive organoids (Fig. 7I). Taken together, these findings collectively illustrate that ECH serves as an effective sensitizing agent for palbociclib in breast cancer, with the combined therapy showing an enhanced anti-tumor activity.

DISCUSSION

For luminal breast cancer patients, the standard treatment consists of CDK4/6 inhibitors combined with endocrine therapy. However, since resistance to CDK4/6 inhibitors is inevitable, there is considerable focus on developing therapeutic strategies to overcome this resistance. Here, we demonstrate that a post-translational modification of PTEN, namely neddylation, contributes to CDK4/6 inhibitor resistance in breast cancer. We firstly prepared antibodies that specifically recognize PTEN neddylation modification. Then, increased PTEN neddylation was observed in breast cancer patients resistant to palbociclib, as well as in both palbociclib-resistant breast cancer cells and spontaneous breast cancer MMTV-*PyMT* mice. Our previous data defined neddylation as a critical modification of PTEN, which inhibits PTEN tumor suppressor function and activates the PI3K/Akt signaling pathway, revealing a previously unidentified tumor-promoting role of neddylated PTEN [30]. Palbociclib resistance upregulated PTEN neddylation *via* the E3 ligase XIAP, consistent with previous studies suggesting that compensatory CDK2 activation upon resistance increases XIAP mRNA levels [37, 38]. This dynamic resulted in enhanced binding to PTEN, thereby contributing to the augmented PTEN neddylation and hyperactivated PI3K/Akt signaling pathway observed following palbociclib resistance.

PIK3CA mutations occur in approximately 40 % of luminal breast cancers, and activation of the PI3K/Akt signaling pathway is prominent as cancers become resistant to CDK4/6 inhibitors [15]. While PI3K inhibitors hold promise in mitigating the onset of CDK4/6 inhibitor resistance, their efficacy in rescuing cells from established resistance falls short, as they are unable to reverse the acquired insensitivity [52-54]. Apart from the upregulation of PI3K/Akt signaling pathway, our study showed that the increase in PTEN neddylation after palbociclib resistance led to the over-activation of the MAPK/AP-1 signaling pathway. Mechanistically, we identified that neddylated PTEN interacted with JUND. JUND is a component protein of the activating protein-1 (AP-1) transcription factor and is considered as a classical proto-oncogene. Dysfunctional AP-1 transcriptional activity is associated with tumor growth, invasion, migration and drug resistance [39, 55, 56]. So far, it is known that CDK4/6 inhibition reprograms the breast cancer enhancer landscape by stimulating AP-1 transcriptional activity [57]. However, the role of AP-1 in regulating resistance to CDK4/6 inhibitors remains unclear. Here, we reported that PTEN neddylation led to an obviously enhancement in JUND protein stability in palbociclib resistance breast cancer cells and mouse models. The expression of JUND in palbociclib-resistant patients showed a positive correlation with the PTEN neddylation modification level and a negative correlation with PTEN expression. Up to day, no ubiquitin ligase has been clearly reported to catalyze the ubiquitination mediated degradation of JUND. We identified that ITCH as a polyubiquitination E3 ligase for JUND. However, under the palbociclib-resistant conditions, ITCH no longer promoted JUND ubiquitination and reduced its protein stability. Further structural docking and immunoprecipitation assays showed that neddylated PTEN interfered with the interaction between JUND and ITCH, reducing the ubiquitination of JUND catalyzed by ITCH, thereby preventing the degradation of JUND. Therefore, the increased modification of PTEN neddylation after palbociclib resistance enhanced the protein stability and phosphorylation level of JUND, thereby activating the AP-1 transcription factor activity. Thus, further data

showed that the expression levels of JUND downstream target genes such as CSF2, IL-1β, IL-6, IL-17, and MMP13 were increased. These results propose a novel role for PTEN neddylation in upregulating AP-1 transcription factor activity and activating the MAPK/AP-1 signaling pathway during resistance to palbociclib treatment.

Neddylation is indispensable in various cellular processes and homeostasis, but excessive activation of this modification has been observed in numerous tumors. Small molecules targeting neddylation activating enzymes (E1), such as MLN4924 (trade name pevonedistat) and TAS4464, have shown promise in clinical trials for cancer treatment [28, 58]. DI-404, DI-591, NAcM-HIT, NAcM-OPT, and WS-383 are competitive neddylation conjugating enzymes (E2) inhibitors that target UBE2M-DCN1 interaction [59-61]. HA-9104 is a targeted inhibitor of another E2-UBE2F [62]. Given the limited number of neddylation E1 and E2 enzymes, inhibiting neddylation broadly may pose inevitable cytotoxicity and drug safety concerns. Therefore, for tumor treatment, more specific and targeted drugs are needed to address neddylation modifications. No small molecule drugs specifically targeting neddylation substrates have been reported, but we believe that would offer greater specificity, reduced drug side effects, and potentially more potent anti-tumor effects. XIAP has been identified as a specific ligase for PTEN neddylation both *in vivo* and *in vitro* [30]. Next, we employed a specific small molecule inhibitor, Echinacoside (ECH), to interfere with the dynamic interplay between XIAP and PTEN, ultimately suppressing the neddylation of PTEN both *in vivo* and *in vitro*. ECH, a key efficacious constituent found in *Echinacea*, is also abundantly present in various other natural plant sources, including the roots of *Scrophulariae*, *Rehmanniae*, and *Cistanches Herba*, among others [45, 46]. It has been reported that ECH inhibits tumor metastasis *via* suppressing the PI3K/AKT signaling pathway, but the mechanism remains unclear [63-65]. Our research demonstrates that ECH exhibits excellent anti-tumor efficacy through its effect on PTEN neddylation. For the first time, our study demonstrates that ECH is a selective inhibitor targeting neddylation substrates.

Although the resistance mechanisms associated with CDK4/6 inhibitors in breast cancer have been documented, but not fully elucidated. For instance, the concurrent use of PI3K inhibitors like alpelisib or buparlisib with CDK4/6 inhibitors is discouraged due to safety concerns [14, 15]. Since PTEN is a key suppressor in the PI3K/Akt pathway, restoring its tumor-suppressive function could more effectively inhibit this pathway [23]. More importantly, our data reported that ECH obviously elevates the sensitivity of palbociclib, and ECH in combination with palbociclib would more effectively reduce the tumor growth. In the tumor microenvironment, the role of neutrophils is complex and multifaceted, highly correlated with the development of drug resistance [66, 67]. Our data showed that the number of activated neutrophils increases after palbociclib resistance, accompanied by an increase in various pro-inflammatory factors and growth factors, such as G-CSF, GM-CSF, IL-1β, VEGF. Treatment with ECH was observed to decrease activated neutrophils following palbociclib resistance. Future study needs to explore the role of activated neutrophils in ROS production and the immunosuppressive microenvironment, and discuss the function of other different terms of neutrophils. Ultimately, our findings unlock a promising therapeutic strategy by focusing on neddylation to effectively restore PTEN tumor suppressor activity, thereby enhancing the efficacy of palbociclib. It highlights a deeper understanding of CDK4/6i resistance mechanisms and the development of advanced combination therapies in optimizing breast cancer treatment outcomes.

METHODS AND MATERIALS

Patients and mice

The clinical samples were approved by the department of breast and thyroid surgery at the Second People's Hospital of Shenzhen and Hunan Cancer Hospital. Informed consent was obtained from all subjects or their relatives. The MMTV-*PyMT* mice (*PyMT*/WT mice, female, background C57BL/6J (RRID: IMSR_JAX: 000664)) were purchased from Cyagen Biosciences. The MMTV-*PyMT* mice (*PyMT*/WT mice, female, background FVB/NJGpt (RRID: IMSR_GPT: T004993))

were kindly gifted by the Second People's Hospital of Shenzhen. CD-1 (ICR, RRID: IMSR_CRL: 022) mice were purchased from Beijing Vital River Laboratory Animal Technology Co., Ltd. Animals were assigned to experimental groups using simple randomization.

Cell culture and reagents

All mammalian cells were cultured following ATCC recommended protocols and in medium supplemented with 10 % fetal bovine serum (FBS, VivaCell BIOSCIENCES C04001-500) and 1 % Penicillin-Streptomycin (Gene-Protein Link P08X14). MCF-7 (RRID: CVCL_0031), HEK293T (RRID: CVCL_0063), HCT116 (RRID: CVCL_0291), A549 (RRID: CVCL_0023), MDA-MB-231 (RRID: CVCL_0062), MCF-7 sh*PTEN*, MCF-7 sh*PTEN*+PTEN-Nedd8, MCF-7 sh*PTEN*+NLS-PTEN, MCF-7 sh*PTEN*+PTEN-WT and MCF-7 sh*PTEN*+PTEN-2KR cells were cultured in Dulbecco's modified Eagle's medium (DMEM) with high glucose (Thermo Fisher Scientific). T-47D (RRID: CVCL_0553) and T98G (RRID: CVCL_0556) cells were cultured in RPMI-1640 medium (Thermo Fisher Scientific C11875500BT). MCF-7 sublines resistant to doxorubicin (MCF-7/ADR) were purchased from Sunncell Biosciences and were cultured in RPMI-1640 medium supplemented with 500 ng/ml Doxorubicin hydrochloride. MCF-7 sublines resistant to cisplatin (MCF-7/DDP) were purchased from Shanghai Mei Xuan Biological Technology Co., Ltd and were cultured in RPMI-1640 medium supplemented with 500 ng/ml Cisplatin. MCF-7 sublines resistant to paclitaxel (MCF-7/PTX) were purchased from BLUEFBIO (Shanghai, China) and were cultured in Dulbecco's modified Eagle's medium (DMEM) with high glucose supplemented with 200 ng/ml Paclitaxel. All cell lines were authenticated by STR profiling.

Palbociclib (HY-50767), Pevonedistat (HY-70062), Echinacoside (HY-N0020), Maltotetraose (HY-N2464), MG132 (HY-13259), Cisplatin (HY-17394), Paclitaxel (HY-B0015), Doxorubicin hydrochloride (HY-15142), Z21911837, Z45457765 and Z1549417413 were purchased from *Med Chem Express*. *ITCH* and negative control small interfering RNA (siRNA) were acquired from

142 GenePharma (Suzhou GenePharma Co.,Ltd). The sequences of siRNAs are shown in

143 Supplementary Table 4.

Establishment of palbociclib-resistant cells

To generate the palbociclib-resistant cells, the cells were cultured in media supplemented with

increasing concentrations of palbociclib for over 6 months. The concentration of palbociclib were

began at 100 nmol/L and were maintained at a final dose of 6 μmol/L, and the palbociclib-

resistant cells were named as MCF-7-PR cells (MCF-7 palbociclib-resistant cells). For all

experiments, the resistant cells were refreshed with palbociclib-free media for 48 h before

treatment.

Antibodies

The following antibodies were used: Phospho-Rb (Ser807/811) (Cat# 8516, RRID:

AB_11178658), Rb (Cat# 9309, RRID: AB_823629), Cyclin D1 (Cat# 2922, RRID:

AB_2228523), PTEN (Cat# 9188, RRID: AB_2253290), Ubiquitin (Cat# 20326, RRID:

AB_3064918), Akt (Cat# 9272, RRID: AB_329827), Phospho-Akt (Ser473) (Cat# 9271, RRID:

AB_329825), Phospho-Akt (Thr308) (Cat# 13038, RRID: AB_2629447), Phospho-JUND

(Ser100) (Cat# 9164, RRID: AB_330892), CUL1 (Cat# 4995, RRID: AB_2261133), XIAP (Cat#

2042, RRID: AB_2214870), Nedd8 (Cat# 2754, RRID: AB_659972), c-JUN (Cat# 9165, RRID:

AB_2130165), JUNB (Cat# 3753, RRID: AB_2130002), c-FOS (Cat# 2250, RRID: AB_2247211)

and FOSB (Cat# 2251, RRID: AB_2106903) antibodies were purchased from Cell Signaling

Technology. Ly6G (Cat# sc-53515, RRID: AB_783639), CD3 (Cat# sc-18843, RRID:

AB_627010), CD4 (Cat# sc-53042, RRID: AB_629062), CD8 (Cat# sc-53063, RRID:

AB_629231), CD11b (Cat# sc-1186, RRID: AB_626884), NEDP1 (Cat# sc-271498, RRID:

AB_10649498) and Cyclin E (Cat# sc-377100, RRID: AB_2923122) antibodies were purchased

from Santa Cruz Biotechnology. F4/80 (Cat# 28463-1-AP, RRID: AB_2881149), GAPDH (Cat#

60004-1-Ig, RRID: AB_2107436), UBA3 (Cat# 68067-1-Ig, RRID: AB_2918808), Ubc12 (Cat#

CL488-67482, RRID: AB_3084351), ITCH (Cat#20920-1-AP, RRID: AB_2878765) and GST

(Cat# HRP-66001, RRID: AB_2883833) antibodies were purchased from Proteintech. Flag (Cat#

M185-3L, RRID: AB_11123930), Myc (Cat# M047-3, RRID: AB_591112) and HA (Cat# M132-

3, RRID: AB_10207271) antibodies were purchased from MBL. JUND (Cat# A5433, RRID:

AB_2863501) and FOSL1 (Cat# A5372, RRID: AB_2766182) antibodies were purchased from

ABclonal. Goat anti-Rabbit IgG (H□+□L) Secondary Antibody (Cat# 65-6120, RRID:

AB_2533967) and Goat anti-Mouse IgG (H□+□L) Secondary Antibody (Cat# 31430, RRID:

AB_228307) were purchased from Thermo Fisher Scientific. PTEN-Nedd8 (K402) antibody was

purchased from PTM BioLab (Hang Zhou) Co., Inc.

Establishment of palbociclib-resistant mice

477 The palbociclib-sensitive MMTV-*PyMT* mice were obtained by continuing palbociclib (75 mg/kg

478 or 100 mg/kg) gavage for 20 days. Palbociclib-resistant MMTV-*PyMT* mice were obtained by

479 continuing palbociclib (75 mg/kg or 100 mg/kg) gavage for 35 or 50 days.

Immunoprecipitation and immunoblotting

For immunoprecipitation assays, cells were lysed in EBC lysis buffer (0.5 % NP-40, 50 mM Tris,

pH 7.6, 120 mM NaCl, 1 mM EDTA, 1 mM Na_3VO_4, 50 mM NaF and 1 mM β-mercaptoethanol)

supplemented with protease inhibitor cocktail (Roche, 11836170001). Immunoprecipitations were

performed using the indicated primary antibody and Dynabeads™ protein G beads (10004D) at

4 °C. The immunoprecipitants were washed at least three times in EBC lysis buffer before being

resolved by SDS-PAGE and immunoblotted with indicated antibodies.

***In vivo* modification assays**

488 Cells were solubilized in modified lysis buffer (50 mM Tris, pH 7.4, 150 mM NaCl, 10 %

489 glycerol, 1 mM EDTA, 1 mM EGTA, 1 % SDS, 1 mM Na_3VO_4, 1 mM DTT and 10 mM NaF)

490 supplemented with a protease inhibitor cocktail. The cell lysate was incubated at 60 °C for 10 min.

491 The lysate was then diluted 10 times with modified lysis buffer without SDS and incubated with

the indicated antibody for 3 h at 4 °C before adding Dynabeads™ protein G beads (10004D).

Then the lysate was rotated gently for 8 h at 4 °C. The immunoprecipitants were washed at least

three times in wash buffer (50 mM Tris, pH 7.4, 150 mM NaCl, 10 % glycerol, 1 mM EDTA, 1

mM EGTA, 0.1 % SDS, 1 mM DTT and 10 mM NaF). Proteins were analyzed by western

blotting with indicated antibodies.

In vitro modification assays

In vitro neddylation was performed using recombinant purified enzymes. GST-PTEN and GST-

Ubc12 were expressed in Escherichia coli BL21 (DE3). 0.5 μg of GST-PTEN and 0.5 μg of GST-

Ubc12 were incubated with 2 μg of Nedd8, 10 ng of E1 (APPBP1 UBA3) and 0.5 μg of E3 (His-

XIAP) in a total reaction volume of 30 μL Nedd8 conjugation Rxn Buffer Kit (Boston Biochem,

SK-20). Nedd8 (UL-812), Nedd8 E1 (APPBP1/UBA3) (E-313), His-XIAP were purchased from

Boston Biochem. Samples were incubated at 30 □ for 1 h, and reactions were terminated with 2 ×

SDS-PAGE loading buffer (20 mM Tris-HCl, pH 6.8, 100 mM DTT, 2 % SDS, 20 % glycerol and

0.016 % Bromophenol blue) before western blotting. If needed, the neddylation reaction sample

was first purified with GST-tag Purification Resin.

Dual Luciferase assay

MCF-7 cells were transfected using Lipofectamine 3000 (Invitrogen) according to manufacturer's

protocol. After 48 hours, cells were lysed in 100 μL of a passive lysis buffer (Promega).

Luciferase activity was measured with the Dual Luciferase Assay System (Promega) according to

the manufacturer's protocol.

GST pull down

The PTEN or PTEN C2 domain sequence was inserted into the pGEX-4T-3 vector (Amersham).

To detect the direct binding, bacteria-expressed GST-tagged proteins were immobilized on

BeyoGold™ GST-tag Purification Resin (P2250) and then incubated with Myc-tagged proteins

for 8 h at 4 □ under rotation. Beads were washed with GST-binding buffer (100 mM NaCl, 50

mM NaF, 2 mM EDTA, 1 % NP-40 and protease inhibitor mixture) three times and proteins were eluted, followed by immunoblotting.

Molecular docking and virtual screening

The molecular docking of PTEN (PDB ID5R) and XIAP (PDB 1C9Q) was performed using the reliable online docking server ZDOCK (http: //zdock.umassmed.edu/). The interaction between PTEN and XIAP was visualized using PYMOL version 1.7.0.3. All ten docking complexes were used for our analysis. The cryo-EM structure of human PTEN (PDB ID5R) was used to perform structure-based virtual screening via a workflow application of Glide in Maestro (Schrödinger Maestro 11.4). The Protein-Protein Interaction Library and the MCE Bioactive Compound Library Plus were selected as screening databases. The LigPrep module of Maestro was used to generate the three-dimensional (3D) conformations of the compounds. At the very beginning, the compounds were screened using the high-throughput virtual screening module and the top-ranked 20 % candidates were selected on the basis of the Glide G-score. Then these candidates were redocked using the standard precision and extra precision mode in turn. Finally, 200 compounds with shape rationality and structural diversity were picked for assay.

Surface plasmon resonance binding assays (SPR)

Surface plasmon resonance experiments were performed on a BIACORE 3000 biosensor system (GE Healthcare) according to the manufacturer's instructions at 25 °C. PTEN protein was immobilized onto a CM5 chip (GE Healthcare), which was activated using a 1: 1 mixture of 1-ethyl-3-(3-dimethyllaminopropyl) carbodiimide (EDC) and N-hydroxysuccinimide (NHS) at a flow rate of 10□ μL/min. ECH was diluted in PBS, and then injected over the PTEN surface at a rate of 30 μL/min flow rate. The duration of protein binding time was set to 180 s, after which the running buffer was injected at the same rate for 300 s. The binding kinetics were processed and calculated by BIAevaluation software.

Microscale thermophoresis assays (MST)

Microscale thermophoresis assays detected interactions between PTEN and ECH. The concentration of PTEN protein was held constantly at 150 nM, whereas the concentrations of ECH were gradient-diluted (0.0153 μM-500 μM). After brief incubation, the samples were loaded into MST-standard glass capillaries. The measurements were performed on a MST machine (NanoTemper Technologies, Munich, Germany).

Cellular thermal shift assay (CETSA)

HEK293T cells (1×10^7) which transfected Myc-PTEN were resuspended in PBS containing 1 % protease inhibitors. The cells were then lysed by liquid nitrogen through three cycles of freezing and thawing. The resulting cell lysates were centrifuged at 12 000 g for 15 min at 4 °C to collect the supernatants. Subsequently, the samples were divided into 7 equal parts and heated at temperatures of 45 °C, 48 °C, 51 °C, 58 °C, 65 °C, 69 °C, and 72 °C for 3 min each. After heating, the samples were centrifuged at 12 000 g for 15 min at 4 °C, and the resulting supernatants were collected and boiled with loading buffer. The levels of the target proteins were determined using western blotting.

Fluorescence microscopy

After fixation with 4 % paraformaldehyde and permeabilization in 0.2 % Triton X-100 (PBS), cells were incubated with the indicated antibodies for 12 h at 4 °C, followed by incubation with goat anti rabbit IgG H&L Alexa Fluor 488 or 568 antibody for 1 h at 37 °C. The nuclei were stained with DAPI, and images were visualized with a Leica confocal microscopy.

Tumor organoid culture

The breast cancer tissues were minced and digested with 5 mL of TrypLE (Gibco 12605-010) digestion solution at 37 °C for 1 hour. The digestion was stopped by adding 5 mL of culture medium, followed by centrifugation at 300 g for 5 minutes. The cell pellet was washed with 10 mL of PBS and then filtered through a 70 µm cell strainer. The filtered suspension was centrifuged again at 300 g for 5 minutes. Pre-chilled pipette tips were used to resuspend the

bottom cells in 50 µL of pre-chilled organoid complete medium and keep them on ice. Then, 250

µL of Matrigel was added to the cell suspension. A pre-warmed 96-well plate was removed, and

30 µL of each drop of the cell suspension was aspirated into the well of the plate using a pre-

chilled pipette tip. The plate was quickly inverted and placed in a 37 °C, 5 % CO☐ incubator to

allow the matrix gel to solidify for 15 minutes. After solidification, an appropriate volume of

organoid complete medium was added to each well.

CUT&Tag sequencing

CUT&Tag assay was performed using NovoNGS CUT&Tag 3.0 High-Sensitivity Kit (for

Illumina, Novoprotein scientific Inc., Cat#N259-YH01-01A). Briefly, $1\square\times 10^5$ cells were

harvested freshly. Then 37 % formaldehyde was gently added to cells and these samples were

incubated at room temperature for 2 min. Cross-linking was stopped by addition of 1.25 M

glycine and then the samples were washed with the wash buffer. The cells were enriched by ConA

Beads and resuspended by 50☐µL primary antibody buffer of anti-JUND antibody, and incubated

overnight at 4 °C. Then discarded the primary antibody buffer and added 100☐µL anti-rabbit IgG

antibody buffer for 1☐h at a dilution of 1: 100. Then the beads were washed for 3 times using

antibody buffer and then incubated with protein A/G-Tn5 transposome for 1 h and washed for 3

times by ChiTaq buffer. Cells were resuspended in 50☐µL Tagmentation buffer and incubated at

37 °C for 1 h. The incubation was stopped by adding 10 µL 10 % SDS at 55 °C for 10 min. The

DNA fragments were extracted by Tagment DNA extract beads and amplified using 5 ×

AmpliMix. Then the DNA was re-extracted by DNA clean beads for sequencing.

Chromatin immunoprecipitation (ChIP) and qChIP

ChIP assays were performed using the EZ-Magna ChIP A/G Chro matin Immunoprecipitation Kit

(17-10086, Merck Millipore) according to the manufacturer's recommendations. In brief, the

harvested MCF-7 cells were cross-linked with 1 % formaldehyde at 37 °C for 20 min, lysed, and

sonicated to generate 200 to 1000 bp DNA fragments. Antibodies against JUND and rabbit IgG

592 were used for immunoprecipitation, and the precipitated DNA fragments were subjected to qPCR

593 amplification. The primers used were listed in Supplementary Table 3.

Flow cytometry analysis and FACS sorting

595 For detection of apoptosis, cells were co-stained with Annexin-V-phycoerythrin and amino

596 actinomycin D (7-AAD) according to the manufacturer's instructions. Stained cells were sorted

597 with fluorescence-activated cell sorting (FACS). The isolated cell suspension was washed with

598 PBS, then stained with specific mAbs at indicated concentration in PBS solution containing 1 %

599 FBS. Antibody staining was performed at 4 °C for 30 min and the stained cells were analyzed on

500 an LSRFortessa cell analyzer (BD Biosciences). FACS sorting routinely yielded cell purity levels

501 of over 90 %. Data were analyzed using FlowJo software (RRID: SCR_008520).

Cohort and immunohistochemistry

The breast tissue microarrays were purchased from Shanghai Biochip Company (HBreD090Bc03)

and Shanghai Zhuoli Biotech Company (ZL-Brcsur1801). ZL-Brcsur1801 was immunostained

with PTEN and Nedd8-PTEN (K402) antibodies (Fig. 1A-C). Detailed pathological information

was in Supplementary Table 1. HBreD090Bc03 was immunostained with Nedd8-PTEN (K402)

and JUND antibodies (Supplementary Fig. 2C-F), which are available in Supplementary Table 2.

In brief, tissues were fixed with 4 % PFA and embedded in paraffin. The samples were then

attached to slides, washed and incubated with 3 % H_2O_2 for 10 min. Afterwards, these sections

were blocked and then incubated with a primary antibody at 4 □°C overnight. After washing with

PBS, the sections were incubated with secondary antibody for 1 h at room temperature. Finally,

the sections were washed and incubated with DAB substrate for visualization. After washing and

dehydration, the sections were sealed with coverslips. All staining was assessed by a quantitative

imaging method; the percentage of immunostaining and the staining intensity were recorded. An

H-score was calculated using the following formula: H-score□=□Σ (PI□×□I)□=□(percentage of

cells of weak intensity□×□1)□+□(percentage of cells of moderate

intensity $\square\times\square$2)$\square+\square$(percentage of cells of strong intensity $\square\times\square$3). PI indicates the percentage of positive cells versus all cells.

Multiplex immunohistochemical

Pheno 7-plex IHC kit (Pheno, PVB2002) was used according to manufacturer's protocol. Slides were deparaffinised in xylene and rehydrated with ethanol and double distilled water. Antigen was retrieved by citrate buffer and heated to boil in microwave for approximately 20 $\square$min, sections were then blocked with bovine serum albumin at room temperature for 10 $\square$min. Primary antibodies were incubated at 37 °C for 30$\square$ min, secondary antibody conjugated with horseradish peroxidase was incubated at room temperature for 10 $\square$min. Tyramide signal amplification was performed with 1: 100 in antibody dilution buffer diluted fluorophores and incubated at room temperature for 10$\square$ min. Nuclei were stained with DAPI after primary antibodies were stained.

ELISA assay

The quantitation of IL-1β, IL-17A and G-CSF levels in MCF-7 cells were performed using ELISA assay kit (Solarbio, SEKM-0002, SEKH-0026, and SEKH-0058) according to the manufacturer's instructions.

Immunoassay arrays

Cytokine concentrations of supernatants of indicated cells were measured using the CBA (Cytometric Bead Array, RayBiotech FAH-STRM-1) following the manufacturer' s instruction. Read samples on a flow cytometer and analyze using FCAP Array software (BD Biosciences).

Statistical analysis

All data were representative of no fewer than three independent experiments. All data were presented as means ± S.D. Data were analyzed using SPSS 19.0 (IBM Corp., Armonk, NY, USA, RRID: SCR_002865) and GraphPad Prism 8 (GraphPad Software, Inc., La Jolla, CA, RRID: SCR_002798). The differences between groups were assessed by Student's *t*-test. The correlations were analyzed using the pearson correlation test. A value of *P* < 0.05 was considered to be

statistically significant.

DATA AVAILABILITY

RNA-seq data have been deposited at SRA (PRJNA1143082 and PRJNA1144658) and are publicly available as of the date of publication. All data are available in the main text or the supplementary materials.

REFERENCES

1. Nolan E, Lindeman GJ, Visvader JE. Deciphering breast cancer: from biology to the clinic. Cell. 2023;186:1708-1728.

2. Harbeck N, Gnant M. Breast cancer. Lancet. 2017;389:1134-1150.

3. Giuliano M, Schettini F, Rognoni C, Milani M, Jerusalem G, Bachelot T, et al. Endocrine treatment versus chemotherapy in postmenopausal women with hormone receptor-positive, HER2-negative, metastatic breast cancer: a systematic review and network meta-analysis. Lancet Oncol. 2019;20:1360-1369.

4. Goel S, Bergholz JS, Zhao JJ. Targeting CDK4 and CDK6 in cancer. Nat Rev Cancer. 2022;22:356-372.

5. Morrison L, Loibl S, Turner NC. The CDK4/6 inhibitor revolution - a game-changing era for breast cancer treatment. Nat Rev Clin Oncol. 2024;21:89-105.

6. O'Sullivan CC, Clarke R, Goetz MP, Robertson J. Cyclin-Dependent Kinase 4/6 Inhibitors for Treatment of Hormone Receptor-Positive, ERBB2-Negative Breast Cancer: A Review. JAMA Oncol. 2023;9:1273-1282.

7. Portman N, Alexandrou S, Carson E, Wang S, Lim E, Caldon CE. Overcoming CDK4/6 inhibitor resistance in ER-positive breast cancer. Endocr Relat Cancer. 2019;26:R15-R30.

8. Lloyd MR, Spring LM, Bardia A, Wander SA. Mechanisms of Resistance to CDK4/6 Blockade in Advanced Hormone Receptor-positive, HER2-negative Breast Cancer and Emerging Therapeutic Opportunities. Clin Cancer Res. 2022;28:821-830.

9. Cai Z, Wang J, Li Y, Shi Q, Jin L, Li S, et al. Overexpressed Cyclin D1 and CDK4 proteins are responsible for the resistance to CDK4/6 inhibitor in breast cancer that can be reversed by PI3K/mTOR inhibitors. Sci China Life Sci. 2023;66:94-109.

10. Palafox M, Monserrat L, Bellet M, Villacampa G, Gonzalez-Perez A, Oliveira M, et al. High p16 expression and heterozygous RB1 loss are biomarkers for CDK4/6 inhibitor resistance in ER(+) breast cancer. Nat Commun. 2022;13:5258.

11. Kudo R, Safonov A, Jones C, Moiso E, Dry JR, Shao H, et al. Long-term breast cancer response to CDK4/6 inhibition defined by TP53-mediated geroconversion. Cancer Cell. 2024;42:1919-1935.

12. Zhou Y, Jin X, Ma J, Ding D, Huang Z, Sheng H, et al. HDAC5 Loss Impairs RB Repression of Pro-Oncogenic Genes and Confers CDK4/6 Inhibitor Resistance in Cancer. Cancer Res. 2021;81:1486-1499.

13. Herrera-Abreu MT, Palafox M, Asghar U, Rivas MA, Cutts RJ, Garcia-Murillas I, et al. Early Adaptation and Acquired Resistance to CDK4/6 Inhibition in Estrogen Receptor-Positive Breast Cancer. Cancer Res. 2016;76:2301-2313.

14. Wander SA, Cohen O, Gong X, Johnson GN, Buendia-Buendia JE, Lloyd MR, et al. The Genomic Landscape of Intrinsic and Acquired Resistance to Cyclin-Dependent Kinase 4/6 Inhibitors in Patients with Hormone Receptor-Positive Metastatic Breast Cancer. Cancer Discov. 2020;10:1174-1193.

15. Clark AS, Makhlin I, DeMichele A. Setting the Pick: Can PI3K Inhibitors Circumvent CDK4/6 Inhibitor Resistance? Clin Cancer Res. 2021;27:371-373.

16. Di Cristofano A, Pandolfi PP. The multiple roles of PTEN in tumor suppression. Cell. 2000;100:387-390.

17. Song MS, Salmena L, Pandolfi PP. The functions and regulation of the PTEN tumour suppressor. Nat Rev Mol Cell Biol. 2012;13:283-296.

18. Lee YR, Chen M, Pandolfi PP. The functions and regulation of the PTEN tumour suppressor: new modes and prospects. Nat Rev Mol Cell Biol. 2018;19:547-562.

19. Papa A, Pandolfi PP. The PTEN-PI3K Axis in Cancer. Biomolecules. 2019;9:153.

20. Costa C, Wang Y, Ly A, Hosono Y, Murchie E, Walmsley CS, et al. PTEN Loss Mediates Clinical Cross-Resistance to CDK4/6 and PI3Kα Inhibitors in Breast Cancer. Cancer Discov. 2020;10:72-85.

21. Dosil MA, Mirantes C, Eritja N, Felip I, Navaridas R, Gatius S, et al. Palbociclib has antitumour effects on Pten-deficient endometrial neoplasias. J Pathol. 2017;242:152-164.

22. Lee JS, Yost SE, Li SM, Cui Y, Frankel PH, Yuan YC, et al. Genomic Markers of CDK4/6 Inhibitor Resistance in Hormone Receptor Positive Metastatic Breast Cancer. Cancers. 2022;14:3159.

23. Lee YR, Chen M, Lee JD, Zhang J, Lin SY, Fu TM et al. Reactivation of PTEN tumor suppressor for cancer treatment through inhibition of a MYC-WWP1 inhibitory pathway. Science. 2019;364:eaau0159.

24. Lin YX, Wang Y, Ding J, Jiang A, Wang J, Yu M, et al. Reactivation of the tumor suppressor PTEN by mRNA nanoparticles enhances antitumor immunity in preclinical models. Sci Transl Med. 2021;13:eaba9772.

25. Kamitani T, Kito K, Nguyen HP, Yeh ET. Characterization of NEDD8, a developmentally down-regulated ubiquitin-like protein. J Biol Chem. 1997;272:28557-62.

26. Enchev RI, Schulman BA, Peter M. Protein neddylation: beyond cullin-RING ligases. Nat

Rev Mol Cell Biol. 2015;16:30-44.

27. Zhang S, Yu Q, Li Z, Zhao Y, Sun Y. Protein neddylation and its role in health and diseases.

Signal Transduct Target Ther. 2024;9:85.

28. Soucy TA, Smith PG, Milhollen MA, Berger AJ, Gavin JM, Adhikari S, et al. An inhibitor of

NEDD8-activating enzyme as a new approach to treat cancer. Nature. 2009;458:732-736.

29. Fu DJ, Wang T. Targeting NEDD8-activating enzyme for cancer therapy: developments,

clinical trials, challenges and future research directions. J Hematol Oncol. 2023;16:87.

30. Xie P, Peng ZQ, Chen YJ, Li HC, Du MG, Tan YW, et al. Neddylation of PTEN regulates its

nuclear import and promotes tumor development. Cell Res. 2021;31:291-311.

31. Du MG, Peng ZQ, Gai WB, Liu F, Liu W, Chen YJ, et al. The Absence of PTEN in Breast

Cancer Is a Driver of MLN4924 Resistance. Front Cell Dev Biol. 2021;9:667435.

32. Chakrabarti R, Wei Y, Hang J, Hang X, Andres Blanco M, Choudhury A, et al. ΔNp63

promotes stem cell activity in mammary gland development and basal-like breast cancer by

enhancing Fzd7 expression and Wnt signalling. Nat Cell Biol. 2014;16:1-13.

33. Wan L, Lu X, Yuan S, Wei Y, Guo F, Shen M, et al. MTDH-SND1 interaction is crucial for

expansion and activity of tumor-initiating cells in diverse oncogene- and carcinogen-induced

mammary tumors. Cancer cell. 2014;26:92-105.

34. Guerriero JL, Sotayo A, Ponichtera HE, Castrillon JA, Pourzia AL, Schad S, et al. Class IIa

HDAC inhibition reduces breast tumours and metastases through anti-tumour macrophages.

Nature. 2017;543:428-432.

35. Knudsen ES, Knudsen KE. Tailoring to RB: Tumour suppressor status and therapeutic

response. Nat Rev Cancer. 2008;8:714-724.

36. Guiley KZ, Stevenson JW, Lou K, Barkovich KJ, Kumarasamy V, Wijeratne TU, et al. p27

allosterically activates cyclin-dependent kinase 4 and antagonizes palbociclib inhibition. Science.

2019;366:6471.

37. Dong F, Guo W, Zhang L, Wu S, Teraishi F, Davis JJ, et al. Downregulation of XIAP and

induction of apoptosis by the synthetic cyclin-dependent kinase inhibitor GW8510 in non-small

cell lung cancer cells. Cancer Biol Ther. 2006;5:165-170.

38. Chen Q, Chen K, Guo G, Li F, Chen C, Wang S, et al. A critical role of CDKN3 in Bcr-Abl-

mediated tumorigenesis. PloS One. 2014;9:e111611.

39. Eferl R, Wagner EF. AP-1: a double-edged sword in tumorigenesis. Nat Rev Cancer.

2003;3:859-868.

40. Liu F, Wagner S, Campbell RB, Nickerson JA, Schiffer CA, Ross AH. PTEN enters the

nucleus by diffusion. J Cell Biochem. 2005;96:221-234.

41. Planchon SM, Waite KA, Eng C. The nuclear affairs of PTEN. J Cell Sci. 2008;121:249-253.

42. Li Y, Xie P, Lu L, Wang J, Diao L, Liu Z, et al. An integrated bioinformatics platform for

investigating the human E3 ubiquitin ligase-substrate interaction network. Nat Commun.

2017;8:347.

43. Wang X, Li Y, He M, Kong X, Jiang P, Liu X, et al. UbiBrowser 2.0: a comprehensive

resource for proteome-wide known and predicted ubiquitin ligase/deubiquitinase-substrate

interactions in eukaryotic species. Nucleic Acids Res. 2022;50:D719-D728.

44. Paul J, Singh AK, Kathania M, Elviche TL, Zeng M, Basrur V, et al. IL-17-driven intestinal

fibrosis is inhibited by Itch-mediated ubiquitination of HIC-5. Mucosal Immunol. 2018;11:427-

436.

45. Angel P, Hattori K, Smeal T, Karin M. The jun proto-oncogene is positively autoregulated by its product, Jun/AP-1. Cell. 1988;55:875-85.

46. Van Themsche C, Leblanc V, Parent S, Asselin E. X-linked inhibitor of apoptosis protein (XIAP) regulates PTEN ubiquitination, content, and compartmentalization. J Biol Chem. 2009;284:20462-20466.

47. Facino RM, Carini M, Aldini G, Saibene L, Pietta P, Mauri P. Echinacoside and caffeoyl conjugates protect collagen from free radical-induced degradation: a potential use of Echinacea extracts in the prevention of skin photodamage. Planta Med. 1995;61:510-514.

48. Wang W, Jiang S, Zhao Y, Zhu G. Echinacoside: A promising active natural products and pharmacological agents. Pharmacol Res. 2023;197:106951.

49. Chen BD, Clark CR, Chou TH. Granulocyte/macrophage colony-stimulating factor stimulates monocyte and tissue macrophage proliferation and enhances their responsiveness to macrophage colony-stimulating factor. Blood. 1988;71:997-1002.

50. Spiekermann K, Roesler J, Emmendoerffer A, Elsner J, Welte K. Functional features of neutrophils induced by G-CSF and GM-CSF treatment: differential effects and clinical implications. Leukemia. 1997;11:466-478.

51. Kolaczkowska E, Kubes P. Neutrophil recruitment and function in health and inflammation. Nat Rev Immunol. 2013;13:159-175.

52. Burgoyne AM, Vann KR, Joshi S, Morales GA, Vega FM, Singh A, et al. A triple action CDK4/6-PI3K-BET inhibitor with augmented cancer cell cytotoxicity. Cell Discov. 2020;6:49.

53. Agostinetto E, Debien V, Marta GN, Lambertini M, Piccart-Gebhart M, de Azambuja E. CDK4/6 and PI3K inhibitors: A new promise for patients with HER2-positive breast cancer. Eur J Clin Invest. 2021;51:e13535.

54. Lee CL, Cremona M, Farrelly A, Workman JA, Kennedy S, Aslam R, et al. Preclinical

evaluation of the CDK4/6 inhibitor Palbociclib in combination with a PI3K or MEK inhibitor in

colorectal cancer. Cancer Biol Ther. 2023;24:2223388.

55. Shaulian E, Karin M. AP-1 as a regulator of cell life and death. Nat Cell Biol. 2002;4:E131-

E136.

56. Ruiz EJ, Lan L, Diefenbacher ME, Riising EM, Da Costa C, Chakraborty A, et al. JunD, not

c-Jun, is the AP-1 transcription factor required for Ras-induced lung cancer. JCI Insight

2021;6:e124985.

57. Watt AC, Cejas P, DeCristo MJ, Metzger-Filho O, Lam EYN, Qiu X, et al. CDK4/6 inhibition

reprograms the breast cancer enhancer landscape by stimulating AP-1 transcriptional activity. Nat

cancer. 2021;2:34-48.

58. Yamamoto N, Shimizu T, Yonemori K, Kitano S, Kondo S, Iwasa S, et al. A first-in-human,

phase 1 study of the NEDD8 activating enzyme E1 inhibitor TAS4464 in patients with advanced

solid tumors. Invest New Drugs. 2021;39:1036-1046.

59. Zhou H, Lu J, Liu L, Bernard D, Yang CY, Fernandez-Salas E, et al. A potent small-molecule

inhibitor of the DCN1-UBC12 interaction that selectively blocks cullin 3 neddylation. Nat

Commun. 2017;8:1150.

60. Zhou H, Zhou W, Zhou B, Liu L, Chern TR, Chinnaswamy K, et al. High-Affinity

Peptidomimetic Inhibitors of the DCN1-UBC12 Protein-Protein Interaction. J Med Chem.

2018;61:1934-1950.

61. Zheng YC, Guo YJ, Wang B, Wang C, Mamun MAA, Gao Y, et al. Targeting neddylation

E2s: a novel therapeutic strategy in cancer. J Hematol Oncol. 2021;14:57.

62. Xu T, Ma Q, Li Y, Yu Q, Pan P, Zheng Y, et al. A small molecule inhibitor of the UBE2F-

CRL5 axis induces apoptosis and radiosensitization in lung cancer. Signal Transduct Target Ther.

2022;7:354.

63. Liu J, Tang N, Liu N, Lei P, Wang F. Echinacoside inhibits the proliferation, migration,

invasion and angiogenesis of ovarian cancer cells through PI3K/AKT pathway. J Mol Histol.

2022;53:493-502.

64. Liang H, Yin G, Shi G, Liu Z, Liu X, Li J. Echinacoside regulates PI3K/AKT/HIF-1α/VEGF

cross signaling axis in proliferation and apoptosis of breast cancer. Anal Biochem.

2024;684:115360.

65. Wei J, Zheng Z, Hou X, Jia F, Yuan Y, Yuan F, et al. Echinacoside inhibits colorectal cancer

metastasis via modulating the gut microbiota and suppressing the PI3K/AKT signaling pathway. J

Ethnopharmacol. 2024;318:116866.

66. Giese MA, Hind LE, Huttenlocher A. Neutrophil plasticity in the tumor microenvironment.

Blood. 2019;133:2159-2167.

67. Teh JLF, Aplin AE. Arrested Developments: CDK4/6 Inhibitor Resistance and Alterations in

the Tumor Immune Microenvironment. Clin Cancer Res. 2019;25:921-927.

AUTHOR CONTRIBUTIONS

Conceptualization: PX; Methodology: FL, WXL, YXS, SHL, XKJ; Formal Analysis: FL;

Investigation: PX, FL; Resources: SYS, YWT, ZZ; Writing-Original Draft: PX; Funding

Acquisition: PX.

FUNDING

This work was supported by the National Outstanding Youth Science Fund Project of National

Natural Science Foundation of China (No. 82122052), National Natural Science Foundation of

China (NSFC) (No. 32471305), Outstanding Young Talents Program in Chinese Institutes for Medical Research (No. CX23YQA03).

COMPETING INTERESTS

The authors declare that there are no competing financial, commercial, or professional interest that could influence the performance, interpretation, or reporting of the research presented in this manuscript. All authors have read and approved this declaration of interests, confirming that it accurately reflects their current situation.

ETHICS APPROVAL

All animals were handled in strict accordance to the "Guide for the Care and Use of Laboratory Animals" and the "Principles for the Utilization and Care of Vertebrate Animals", and all animal work was approved by Capital Medical University Ethics Committee. The clinical samples were approved by the department of breast and thyroid surgery at the Second People's Hospital of Shenzhen and Hunan Cancer Hospital. Informed consent was obtained from all subjects or their relatives.

ACKNOWLEDGEMENTS

Thanks to Core Facility Center, Capital Medical University for providing platform support for this research.

FIGURE LEGENDS

Fig. 1 Palbociclib-resistant breast cancer cells express higher levels of PTEN neddylation modification.

A Representative image from immunohistochemical staining of PTEN and PTEN neddylation on K402 site in breast cancer and matched adjacent tissues. ImageJ was used to perform Semi-quantitative analysis. Data were analyzed using Student's *t*-test. Scale bars, 50 µm. PTEN and PTEN neddylation on K402 expression are shown as box plots. **B** Kaplan-Meier plot of overall

survival of patients with breast carcinomas. A log-rank test was used to show differences between groups (red indicates high PTEN or PTEN-N8 K402 expression; black indicates low PTEN or PTEN-N8 K402 expression). **C** Representative images from immunohistochemical staining of PTEN and neddylated PTEN on K402 site in different breast cancer subtypes and adjacent tissues are shown. Scale bars, 50 µm. Data were analyzed using the one-way ANOVA test. PTEN and PTEN neddylation on K402 expression are shown as box plots. **D** Representative images from immunohistochemical staining of PTEN and PTEN neddylation on K402 in preinvasive breast cancer tissues (n=6), palbociclib resistance breast cancer tissues (n=6) and adjacent tissues (n=4). Data were analyzed using Student's *t*-test. Scale bars, 100 µm. The protein level of PTEN and PTEN neddylation on K402 expression are shown as bar plots. **E-H** To establish the palbociclib-resistant mouse model, the palbociclib-resistant MMTV-*PyMT* mouse model was established by palbociclib (75 mg/kg) gavage daily for 50 days and the palbociclib-sensitive group were administered palbociclib (75 mg/kg) by gavage daily for 20 days. The three conditions were compared in the same time (n=5) (**E**). The breast tumors were removed and photographed (**F**). Tumors were isolated, volumes (**G**) and their weights (**H**) were measured. Data are presented as means ± S.D. **I-L** To establish the palbociclib-resistant mouse model, the palbociclib-resistant MMTV-*PyMT* mouse model was established by palbociclib (75 mg/kg or 100 mg/kg) gavage daily for 35 days and the palbociclib-sensitive group were administered palbociclib (75 mg/kg or 100 mg/kg) by gavage daily for 20 days (placebo group n=5; palbo-S-75mg/kg group n=5; palbo-R-75mg/kg group n=4; palbo-S-100mg/kg group n=5; palbo-R-100mg/kg group n=5). The three conditions were compared in the same time (**I**). The breast tumors were removed and photographed (**J**). Tumors were isolated, volumes (**K**) and their weights (**L**) were measured. Data are presented as means ± S.D. **M** Immunoblot analysis of anti-PTEN immunoprecipitate and cell lysates derived from spontaneous breast tumors of MMTV-*PyMT* mice in the indicated three groups. **N** Immunoblot analysis of anti-Nedd8 immunoprecipitate and cell lysates derived from

spontaneous breast tumors of MMTV-*PyMT* mice in the indicated three groups. **O** IC50 of

palbociclib-resistant breast tumor cells obtained from MMTV-*PyMT* mice (MMTV-*PyMT*+palbo-

S and MMTV-*PyMT*+palbo-R). The cells were seeded in 96-well plates treated with increasing

concentration (0.01 nM-1000 µM) of palbociclib and then analyzed for CCK8 assay. *P* values

were calculated by Student's *t*-test. (**A**), log-rank test (**B**) and one-way ANOVA test (**C**, **D**, **G**, **H**,

K, **L**). Error bars, ± S.D. ns, not significant, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Fig. 2 Upregulation of XIAP increases the PTEN neddylation in palbociclib-resistant breast

cancer.

A IC50 of MCF-7-PS (palbociclib sensitivity, PS) and MCF-7-PR (palbociclib resistance, PR)

cells. MCF-7-PR cells and MCF-7-PS cells were seeded in 96-well plates and then treated with

increasing concentrations (0.01 nM-1000 µM) of palbociclib and then analyzed for CCK8 assay.

B MCF-7-PS and MCF-7-PR cells are treated with palbociclib (0.5□µM, 24 h) or not. The

percentage of cells in G1/S-phase is depicted. Data are presented as the means ± S.D. (n=3). **C**

Colony formation assay was performed in the MCF-7-PS cells and MCF-7-PR cells and the cells

were treated with palbociclib (0.1□µM, 15 days). Data are presented as means ± S.D. Results are

from a representative experiment performed in triplicate. ImageJ was used to perform quantitative

analysis. **D** Immunoblot analysis of anti-PTEN immunoprecipitate from MCF-7-PS and MCF-7-

PR cells to detect PTEN neddylation *in vivo*. **E** Immunoblot analysis of anti-PTEN

immunoprecipitate from MCF-7 drug resistant cells including MCF-7-PR, MCF-7 sublines

resistant to doxorubicin (MCF-7-ADR), cisplatin (MCF-7-DDP) and paclitaxel (MCF-7-PTX)

cells. **F** MCF-7 cells and T-47D cells were treated with palbociclib (0.5 µM, 0 h/48 h/96 h). The

right panel was treated with palbociclib (0.5 µM, 96 h) and MLN4924 (1 µM, 12 h)

simultaneously. The cells were collected and lysed for immunoblot analysis with indicated

antibodies. **G** Immunoblot analysis of anti-PTEN immunoprecipitate from MCF-7 cells and T98G

cells. MCF-7 cells and T98G cells were respectively treated with palbociclib (0.5 µM, 0 h/48 h/96

h). **H** Western blot analysis of XIAP, PTEN, UBA3, Ubc12, Nedd8, NEDP1 in palbociclib sensitivity and resistance MCF-7 and T-47D cells. **I** Analysis of CDK2 and XIAP mRNA levels in MCF-7-PS/PR cells. **J** Analysis of CDK2 and XIAP mRNA levels in MMTV-*PyMT*-palbo-S/palbo-R tissues. **K, L** Immunoblot analysis of anti-PTEN immunoprecipitate and WCL from adaptative palbociclib resistance in MCF-7 (**K**) and T-47D (**L**). **M** Coimmunoprecipitation of XIAP after immunoprecipitation of the PTEN protein in palbociclib sensitivity and resistance MCF-7 cells. *P* values were calculated by one-way ANOVA test (**B, C**) and Student's *t*-test (**I, J**). Error bars, ± S.D. ns, not significant, $**P < 0.01$, $***P < 0.001$.

Fig. 3 PTEN neddylation enhances JUND protein stability.

A, B The bubble charts depict the top ranked pathway analyzed from the KEGG pathway database. Enriched KEGG pathways were identified, which included upregulated or downregulated genes modulated by the palbociclib resistance MCF-7 cells (**A**) and MCF-7 cells harboring a PTEN neddylation mutation (**B**). **C, D** GSEA plots of the palbociclib resistance MCF-7 cells (**C**) and PTEN neddylation deficiency MCF-7 cells (**D**). **E** AP-1 luciferase reporter plasmid was transfected in the palbociclib sensitivity and resistance MCF-7 cells. Reporter activity was assayed and represented as the means ± S.D. of three separate experiments. **F, G** Heat map of the palbociclib resistance MCF-7 cells (**F**) and PTEN neddylation deficiency MCF-7 cells (**G**). **H, I** Immunoblot of WCL from acquired palbociclib-sensitive or palbociclib-resistant cells. **J** Western blot analysis of lysates from MCF-7 cells treated for 24 to 96 hours with palbociclib (Palbo). Addition of fresh vehicle or drug every 24 hours over a 96-hour period. **K** The cells obtained from breast tumors of MMTV-*PyMT* mice were collected and lysed for immunoblot analysis. **L** Immunoblot of Nuclear (N) *vs* Cytoplasmic (C) fractionation from palbociclib sensitivity and resistance MCF-7 cells. **M** Representative images from immunohistochemical staining of JUND in palbociclib-resistant breast tumors (n=4) and matched adjacent tissues (n=3). ImageJ was used to perform Semi-quantitative analysis. Scale bars, 50 µm. *P* values were

calculated by Student's *t*-test. Error bars, ± S.D. ns, not significant, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$.

Fig. 4 PTEN neddylation hinders the ITCH-mediated ubiquitination degradation of JUND.

A Immunoprecipitation of JUND in the MCF-7-PS and MCF-7-PR cells and analysis using immunoblotting with the indicated antibodies. **B** Immunoprecipitation of Ub in the MCF-7-PS and MCF-7-PR cells and analysis using immunoblotting with the indicated antibodies. **C** Immunoblot of JUND and p-JUND in the MCF-7 sh*NC*, MCF-7 sh*PTEN*+PTEN-NLS and MCF-7 sh*PTEN*+PTEN-N8 cells. **D, E** Immunoprecipitation of PTEN (**D**) and JUND (**E**) in the MCF-7 sh*NC*, MCF-7 sh*PTEN*+PTEN-NLS and MCF-7 sh*PTEN*+PTEN-NLS and MCF-7 sh*PTEN*+PTEN-N8 cells. **F** Flag tagged-ITCH was transfected into the MCF-7-PS cells and MCF-7-PR cells and immunoblotted with the indicated antibodies. The cells were treated with palbociclib (0.5 μM for 48 h). **G** Si*ITCH* was transfected into the MCF-7-PS cells and MCF-7-PR cells and immunoblotted with the indicated antibodies. The cells were treated with palbociclib (0.5 μM for 48 h). **H** Immunoprecipitation of JUND in HEK293T cells transfected with plasmids of HA-tagged-Ub, Myc-tagged-ITCH and Myc-tagged-ITCH C830A and analysis using immunoblot analysis with the indicated antibodies. **I** Immunoprecipitation of JUND in HEK293T cells transfected with si*ITCH* and analysis using immunoblot analysis with the indicated antibodies. **J** Structure simulation of ITCH (yellow), JUND (pink) and PTEN or PTEN-Nedd8 (grey) by AlphaFold2.0. **K** MCF-7 sh*PTEN*+PTEN-WT and MCF-7 sh*PTEN*+PTEN-N8 cells transfected with plasmids for Flag-tagged ITCH and analyzed using immunoblot analysis with the indicated antibodies. **L, M** Flag tagged-PTEN or PTEN-Nedd8 plasmid was transfected into the MCF-7 cells and immunoblotted with the indicated antibodies. **N** Coimmunoprecipitation of ITCH and PTEN after immunoprecipitation of the JUND protein in palbociclib sensitivity and resistance MCF-7 cells.

Fig. 5 Echinacoside is identified as an inhibitor of PTEN neddylation.

A Immunoblot analysis of GST pull-downs or WCL from HEK293T cells transfected with indicated constructs. **B** Molecular docking of PTEN C2 domain (gray) and XIAP BIR2 domain (blue). The key amino acids (red) at the PTEN and XIAP binding sites are shown. **C** Immunoblot analysis of GST pull-downs or WCL from HEK293T cells transfected with indicated constructs. **D** The workflow of virtual drug screening. **E** Structural formula and molecular docking score of Echinacoside with PTEN. **F** Molecular docking of Echinacoside (ECH) with PTEN was conducted to estimate the binding affinity. **G** Representative images of SPR binding assay of ECH to PTEN. ECH exhibited dose-dependent binding to GST-tagged PTEN that was immobilized onto a CM5 sensor chip surface. **H** All hits were titrated at various serially diluted concentrations (0.0153 μM to 500 μM) into a fixed concentration of GST-PTEN C2 domain (150 nM). **I** Dose and temperature-dependent CETSA experiments were conducted to verify the interaction between Echinacoside (ECH) and PTEN (n=3). The samples were divided into seven equal parts and heated at temperatures of 45 °C, 48 °C, 51 °C, 58 °C, 65 °C, 69 °C, and 72 °C. The abundance of PTEN protein was quantified using ImageJ. **J** Immunoprecipitates of PTEN were obtained from MCF-7-PS and MCF-7-PR cells transfected with Myc-XIAP (amino acids 152-235) and treated with Echinacoside (200 μM for 24 h). **K** Immunoprecipitates of XIAP were obtained from MCF-7-PS and MCF-7-PR cells transfected with Myc-PTEN C2 domain and treated with Echinacoside (200 μM for 24 h). **L** Immunoblot analysis of GST pull-downs or WCL from HEK293T cells transfected with indicated constructs. The concentration of ECH was used from 20 μM to 200 μM for 24 h. **M** PTEN neddylation was attenuated by ECH *in vivo*. The cells were treated with ECH at a concentration of 200 μM for 24 h. Immunoblot analysis was performed using anti-Nedd8, anti-Ub, anti-XIAP, and anti-PTEN antibodies on immunoprecipitates and whole cell lysates (WCL) from MCF-7 cells. **N** *In vitro* neddylation of PTEN. Purified His-XIAP and GST-PTEN proteins were incubated with Nedd8, Nedd8-E1/E2. Reactions were performed as described in the materials and methods section. Samples were analyzed by western blot with indicated antibodies.

ECH (100 μg) was added and reacted for 1 h. **O** Immunoblot of WCL from MCF-7 PS and PR

cells for indicated proteins. The cells were treated with 200 μM ECH for 24 h. **P** *In vivo*

ubiquitination of PTEN. Immunoblot analysis of anti-JUND immunoprecipitate and WCL from

MCF-7-PS cells and MCF-7-PR cells. **Q** AP-1 luciferase reporter plasmid was transfected in the

palbociclib sensitivity and resistance MCF-7 cells. Reporter activity was assayed and represented

as the means ± S.D. of three separate experiments. The cells were treated with ECH (200 μM) for

24 h.

Fig. 6 ECH is dependent on PTEN neddylation to suppress tumorigenesis.

A-C CCK8 (**A**), colony formation (**B**) and cell invasion assay (**C**) were performed in the indicated

MCF-7 cells treated with ECH. **D-G** CCK8 (**D, E**), colony formation (**F**) and apoptosis assay (**G**)

were performed in the indicated MCF-7 cells treated with palbociclib. **H, I** CCK8 (**H**), and colony

formation (**I**) were performed in the indicated MCF-7 cells treated with ECH. **J** Immunoblot of

WCL from the sh*PTEN*+PTEN-WT/2KR MCF-7 cells treated by ECH (200 μM, 24 h) or not. **K**

AP-1 luciferase reporter plasmid was transfected in the indicated cells. Reporter activity was

assayed and represented as the means ± S.D. of three separate experiments. The cells were treated

with ECH (200 μM) for 24 h. **L, M** Identification of transcription targets of the JUND in the

sh*PTEN*-PTEN WT or neddylation deficient MCF-7 cells. Venn diagrams of overlapped

promoters bound by JUND (**L**). A bubble chart of the enriched KEGG pathways comprising the

6306 target genes (**M**). **N** qChIP analysis of the indicated genes in indicated MCF-7 cells. Results

are represented as fold change relative to control with GAPDH as a negative control. **O**

Cytometric bead array was performed to detect the indicated cytokines. Heat map was performed

to exhibit protein expression in the indicated cells. The cells were treated with ECH (200 μM) for

24 h. **P** The protein level of G-CSF, GM-CSF, IFNγ, CCL2, IL-8, VEGF, IL-1RA, IL-1β, IL-17

and CCL4 from the immunoassay array of the indicated MCF-7 cells are shown as bar chart. **Q**

G-CSF, IL-1β and IL-17 protein levels were measured in the indicated MCF-7 cells by ELISA

assay. *P* values were calculated by one-way ANOVA test (**A-C, E-I, K, N, P, Q**). Error bars, ±

S.D. ns, not significant, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$.

Fig. 7 Combination ECH enhances the sensitivity of palbociclib in breast cancer.

A-C The palbociclib-resistant mice (FVB/NJGpt background) model was established. The mice

were treated with ECH (20 mg/kg) after palbociclib resistance for 15 days. The breast tumors

were removed and photographed (n=3) (**A**). Tumors were isolated, then the weights (**B**) and their

volume (**C**) were measured. Data are presented as means ± S.D. **D** Representative images from

immunohistochemical staining of CD3, CD4, CD8, Ly6G and F4/80 in the indicated tumors.

ImageJ was used to perform Semi-quantitative analysis. Scale bars, 50 µm. **E** Primary breast

cancer cells from the indicated MMTV-*PyMT* mice were isolated and labeled with CD11b$^+$ and

Ly6G$^+$ following flow cytometry sorting. **F** CCK8 assay was performed in MCF-7 cells treated

with palbociclib (0.5 µM, 5 days) and ECH (200 µM, 5 days). **G, H** Representative images of

organoids derived from a luminal breast cancer patient (**G**). Echinacoside (ECH) (200 µM) and

palbociclib (0.5 µM) were administered either individually or in combination for 7 days. The

diameters of the organoids are presented as a scatter plot (n=5) (**H**). **I** Immunofluorescent of E-

Cadherin (green) and neddylated PTEN (K402) (red) in luminal breast cancer organoids. Data are

shown as means ± S.D. and analyzed by a one-way ANOVA test (**B-F, H**). ns, not significant, $*P$

< 0.05, $**P < 0.01$, $***P < 0.001$.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A MMTV-*PyMT*

palbo. (75 mg/kg)
ECH (20 mg/kg)

palbo-S
palbo-R
palbo-R+ECH

0   20   35 Days

placebo palbo-S palbo-R palbo-R+ECH   placebo palbo-S palbo-R palbo-R+ECH

MMTV-*PyMT*

B
Tumor Weight (g)
ns
***   ***

● MMTV-*PyMT*+placebo
■ MMTV-*PyMT*+palbo-S
▲ MMTV-*PyMT*+palbo-R
▼ MMTV-*PyMT*+palbo-R
  +ECH

C
Tumor Volume (cm³)
ns
ns
***
0 5 10 15 20 25 30 35 Days

● MMTV-*PyMT*+placebo
■ MMTV-*PyMT*+palbo-S
▲ MMTV-*PyMT*+palbo-R
▼ MMTV-*PyMT*+palbo-R+ECH

D

	CD3	CD4	CD8	Ly6G	F4/80
MMTV-*PyMT*+placebo					
MMTV-*PyMT*+palbo-S					
MMTV-*PyMT*+palbo-R					
MMTV-*PyMT*+palbo-R+ECH					

CD3 IOD/Area (%) ns ns
CD4 IOD/Area (%) ns ns
CD8 IOD/Area (%) ns ns
Ly6G IOD/Area (%) * *** ***
F4/80 IOD/Area (%) ns * *

□ MMTV-*PyMT*+placebo
■ MMTV-*PyMT*+palbo-S
■ MMTV-*PyMT*+palbo-R
■ MMTV-*PyMT*+palbo-R+ECH

E
placebo   palbo-S
4.56 %   1.07 %
palbo-R   palbo-R+ECH
7.99 %   1.68 %

CD11b (APC)
Ly6G (BV421)

% Ly6G⁺ CD11b⁺ of cells
***
**   **   ***

□ MMTV-*PyMT*+placebo
■ MMTV-*PyMT*+palbo-S
■ MMTV-*PyMT*+palbo-R
■ MMTV-*PyMT*+palbo-R+ECH

F MCF-7
OD. 450nm
*
***
0 2 4 6 Days

● DMSO
■ ECH
▲ palbo.
▼ palbo.+ECH

G
0   3   7 Days

DMSO
palbo.
ECH
ECH+palbo.

Luminal breast cancer patient

H
Organoids (μm)
***
***
***
**
1   7 Days

● DMSO
■ palbo.
▲ ECH
▼ palbo.+ECH

I

	E-Cad	PTEN-N8 (K402)	DAPI	Merge
DMSO				
ECH				
palbo.				
ECH+palbo.				