The novel ECM protein SNED1 mediates cell adhesion

via the RGD-binding integrins α5β1 and αvβ3

Dharma Pally[1], Nandini Kapoor[1], Alexandra Naba[1,2]

[1] Department of Physiology and Biophysics, University of Illinois Chicago, Illinois, 60612, USA

[2] University of Illinois Cancer Center, Chicago, Illinois, 60612, USA

Correspondence: anaba@uic.edu

Running title: SNED1 mediates cell adhesion

Keywords: Cell-ECM interactions, ECM receptors, Breast cancer metastasis, Neural crest cells, RGD motif, Matrisome

Word count: 2,951

ABSTRACT

The extracellular matrix (ECM) is a complex meshwork comprising over 100 proteins. It serves as an adhesive substrate for cells and, hence, plays critical roles in health and disease. We have recently identified a novel ECM protein, SNED1, and have found that it is required for neural crest cell migration and craniofacial morphogenesis during development and in breast cancer, where it is necessary for the metastatic dissemination of tumor cells. Interestingly, both processes involve the dynamic remodeling of cell-ECM adhesions via cell surface receptors. Sequence analysis revealed that SNED1 contains two amino acid motifs, RGD and LDV, known to bind integrins, the largest class of ECM receptors. We thus sought to investigate the role of SNED1 in cell adhesion. Here, we report that SNED1 mediates breast cancer and neural crest cell adhesion via its RGD motif. We further demonstrate that cell adhesion to SNED1 is mediated by the RGD integrins α5β1 and αvβ3. These findings are a first step toward identifying the signaling pathways activated downstream of the SNED1-integrin interactions guiding craniofacial morphogenesis and breast cancer metastasis.

SUMMARY STATEMENT

We report that the novel extracellular matrix protein SNED1 promotes the adhesion of breast cancer cells and neural crest cells via interaction with α5β1 and αvβ3 integrins, the first SNED1 receptors identified to date.

INTRODUCTION

The extracellular matrix (ECM), a fundamental component of multicellular organisms, is a complex 3-dimensional (3D) meshwork consisting of over a hundred proteins (Hynes and Naba, 2012; Naba, 2024). The primary function of the ECM is to serve as a substrate for cell adhesion (Hynes, 2009). The adhesion of cells to their surrounding ECM is mediated by cell-surface receptors and is critical for cell survival, as detachment from the ECM results in apoptotic cell death, known as anoikis (Frisch and Francis, 1994). In addition, cell-ECM interactions trigger molecular events that regulate a multitude of cellular phenotypes including migration (Dzamba and DeSimone, 2018; Pally and Naba, 2024), proliferation (Hynes, 2009), and differentiation (Walma and Yamada, 2020). As a result, alterations of cell-ECM adhesions and downstream signaling pathways lead to developmental defects (Rozario and DeSimone, 2010) and pathologies like cancer (Cox, 2021; Pickup et al., 2014) and fibrosis (Herrera et al., 2018). Yet, only a small subset of the hundreds of proteins comprising the matrisome is the focus of active investigations, and the mechanisms by which they interact with cells and guide cell phenotype are known for an even smaller subset.

One such understudied ECM protein is Sushi, Nidogen, and EGF like Domains 1 (SNED1). The murine gene *Sned1* was cloned two decades ago (Leimeister et al., 2004), however, it took ten years to identify its first function as a promoter of breast cancer metastasis (Naba et al., 2014). Beyond its role in breast cancer, we recently reported that *Sned1* is an essential gene, as knocking it out resulted in early neonatal lethality and severe craniofacial malformations (Barqué et al., 2021). We further showed that knocking out *Sned1* specifically from neural crest cells, the cell population that contributes to forming most craniofacial features (Mankarious and Goudy, 2010; Martik and Bronner, 2021; Trainor, 2005), was sufficient to recapitulate the craniofacial phenotype observed upon global *Sned1* deletion, demonstrating a new role for SNED1 in craniofacial morphogenesis (Barqué et al., 2021). However, as of today, the mechanisms through which SNED1 interacts with cells to mediate its phenotypes remain unknown. Of note, metastatic breast cancer cells and neural crest cells share common features (Gallik et al., 2017), including their ability to remodel their adhesions to acquire increased migratory potential (Doyle et al., 2022; Gallik et al., 2017; Mayor and Theveneau, 2013; Tucker et al., 1988). This process is, in part,

mediated by integrins, which are the main class of ECM receptors (Bökel and Brown, 2002;
Campbell and Humphries, 2011; Hood and Cheresh, 2002; Hynes, 2002; Kanchanawong and
Calderwood, 2023). For example, *in vivo* and *in vitro* experiments have demonstrated that β1
integrins at the surface of neural crest cells interact with the ECM and ECM proteins like
fibronectin to promote cell adhesion and subsequent migration (Alfandari et al., 2003; Duband et
al., 1991; Leonard and Taneyhill, 2020; Pietri et al., 2004; Testaz et al., 1999; Yang et al., 1993).
β1-containing integrin heterodimers expressed by breast cancer cells have been shown to interact
with the vascular ECM to promote extravasation during metastasis (Chen et al., 2016) and be
essential to every step of the metastatic cascade (Hamidi and Ivaska, 2018). Interestingly, SNED1
contains two putative integrin-binding motifs, an arginine-glycine-aspartic acid (RGD) triplet and
a leucine-aspartic acid-valine (LDV) triplet, known in other proteins to mediate cell-ECM
adhesion. We thus sought to determine whether SNED1 played a role in cell adhesion.

Here we report that SNED1 mediates the adhesion of breast cancer cells and neural crest cells, two
cell types of relevance to the *in-vivo* functions of SNED1. Using a combination of genetic and
pharmacological approaches, we further show that cell adhesion to SNED1 is mediated by its RGD
motif and the engagement of α5β1 and αvβ3 integrins. Our study is thus the first to report the
identification of SNED1 receptors and constitutes an important step toward the identification of
the biochemical signaling events leading to SNED1-dependent breast cancer metastasis and
craniofacial development.

RESULTS AND DISCUSSION

SNED1 mediates breast cancer and neural crest cell adhesion

We first sought to determine whether SNED1 could mediate cell adhesion. To do so, we seeded highly metastatic MDA-MB-231 'LM2' breast cancer cells (further termed LM2) or O9-1 neural crest cells on surfaces coated with increasing concentrations of purified human SNED1 (Vallet et al., 2021) (Fig S1A) or murine Sned1, respectively, and allowed cells to adhere for 30 min. Cell adhesion was assayed using a crystal-violet-based colorimetric assay and we found that SNED1 mediated the adhesion of LM2 breast cancer cells and O9-1 neural crest cells in a concentration-dependent manner with maximum adhesion observed at 10 μg/mL concentration (Fig 1A and B, respectively). At this concentration, SNED1 exerted the same adhesive properties as fibronectin toward these two cell populations (represented by 100% adhesion based on the standard curve). We further showed that the O9-1 cells, which are of murine origin, adhered in the same proportion to both murine Sned1 and human SNED1 (Fig S1B), perhaps not surprisingly, as the two orthologs share an 84% sequence identity (Barqué et al., 2021).

Sned1 also mediated the adhesion of immortalized mouse embryonic fibroblasts derived from *Sned1* knockout mice (*Sned1*[KO] iMEFs) to the same extent as the other two cell lines (Fig S1C). Altogether these experiments demonstrate that SNED1 mediates cell adhesion and exerts adhesive properties toward a panel of cell types.

The N-terminal region of SNED1 is sufficient to mediate cell adhesion

SNED1 is a modular protein composed of different domains, including a NIDO domain, a follistatin domain, and a sushi domain (also known as complement control protein or CCP domain), in addition to multiple repeats of EGF-like (EGF), calcium-binding EGF-like (EGF-Ca), and fibronectin type 3 (FN3) domains, all involved in protein-protein interactions (Fig 2A). To assess which region(s) of SNED1 mediate cell adhesion, we generated and purified three truncated forms of SNED1 (Fig 2A): the SNED1[1-751] form lacks the C-terminal region that comprises three FN3 domains, shown to mediate cell adhesion in other ECM proteins, and the most C-terminal EGF-like domains; the SNED1[1-530] fragment additionally lacks the sushi domain and EGF-like domains after the follistatin domain; the SNED1[1-260] construct encompasses the very N-terminal region, including a single NIDO domain, which is only present in four other proteins in the human

and mouse proteomes (nidogen-1 and nidogen-2, alpha-tectorin, and mucin-4) and its function

remains unknown (Barqué et al., 2021). These synthetic constructs were designed to include

adequate domain boundaries compatible with proper folding for the truncated proteins to be

secreted (Fig 2B).

Using these purified truncated proteins as substrates, we found that LM2 cells had a decreased

ability to adhere to SNED1$^{1-751}$ and SNED1$^{1-530}$ as compared to full-length SNED1 (1.4- and 1.75-

fold decrease, respectively; Fig 2C). Similarly, we observed that O9-1 cells had a decreased ability

to adhere to SNED1$^{1-751}$ and SNED1$^{1-530}$ compared to full-length SNED1 (1.58- and 1.7-fold

decrease, respectively; Fig 2D). Notably, although SNED1$^{1-530}$ further lacks the sushi domain and

four EGF-like domains, along with the domains absent in SNED1$^{1-751}$, it mediated cell adhesion

to the same extent as SNED1$^{1-751}$ for both cell lines (Fig 2C and D). Interestingly, LM2 and O9-1

cells could adhere to the shortest N-terminal SNED1$^{1-260}$ fragment in a similar proportion as to

full-length SNED1 (Fig 2C and D, respectively). Since, the SNED1$^{1-260}$ fragment is 6 times smaller

than full length SNED1, we thought to perform the same experiment but using similar molar

concentration (66.3 μM) rather than amount (10μg/mL), and obtained a similar result, namely that

LM2 cell adhesion on SNED1$^{1-260}$ is comparable to full length SNED1 (Fig S2). Altogether, these

results indicate that the adhesive property of SNED1 is primarily mediated by its N-terminal

region. These results also suggests that the three FN3 domains and the EGF-like domains lacking

in the SNED1$^{1-751}$ and SNED1$^{1-530}$ constructs are required for full length SNED1 to adopt a

conformation where the N-terminal adhesive site is fully accessible to cells, since their absence

resulted in decreased cell adhesion.

SNED1 mediates cell adhesion via its RGD motif

Analysis of the SNED1 sequence has revealed the presence of two putative integrin binding motifs:

RGD and LDV. These motifs were discovered in other ECM proteins, such as fibronectin and

thrombospondin 1, and interact with integrin heterodimers at the cell surface to mediate cell

adhesion (Lawler et al., 1988; Ruoslahti and Pierschbacher, 1987).

To determine whether the RGD and LDV motifs of SNED1 are required for cell adhesion, we

mutated these sites alone or in combination (p40D>E in the RGD motif; p311D>A in the LDV

motif; Fig 3A). Similar mutations in other ECM proteins have been shown to disrupt their

interaction with integrin heterodimers (Cherny et al., 1993; Pytela et al., 1985). We next expressed

these constructs in 293T cells and purified the corresponding secreted proteins from the

conditioned culture medium using affinity chromatography (Fig S3). Purified proteins were then

used to perform cell adhesion assays. We observed that LM2 cells showed a statistically significant

decrease in their ability to adhere to SNED1^{RGE} and SNED1$^{RGE/LAV}$ as compared to SNED1^{WT} (3-

and 2.5-fold decrease, respectively; Fig 3B). Similarly, O9-1 cells showed a significant decrease

in their ability to adhere to SNED1^{RGE} and SNED1$^{RGE/LAV}$ as compared to SNED1^{WT} (4- and 5.5-

fold decrease, respectively; Fig 3C). However, mutation of the LDV motif to LAV did not affect

cell adhesion (Fig 3B, C). These results demonstrate that the RGD but not the LDV motif in

SNED1 is required for cell adhesion.

Functional inhibition of RGD integrins significantly reduces breast cancer and neural crest

cell adhesion to SNED1

To complement this set of observations and determine whether integrins mediate adhesion to

SNED1, we performed experiments aimed at targeting the ability of integrins to interact with

SNED1. First, we performed adhesion assays in the presence of cyclic RGDfV (cRGDfV) peptide,

a peptide known to bind with high affinity to integrins that engage with the RGD motif of ECM

proteins (Aumailley et al., 1991). We observed reduced adhesion of both LM2 (Fig 4A) and O9-1

cells (Fig 4B) in a concentration-dependent manner in presence of cRGDfV as compared to

vehicle-treated cells. Since we previously showed that the N-terminal region of SNED1 was

sufficient to mediate cell adhesion and the RGD motif is located within this fragment, we evaluated

the ability of cells to adhere to SNED1$^{1-260}$ in presence of 10μM of cRGDfV peptide. We found

that this integrin inhibitor fully abrogated LM2 and O9-1 cell adhesion (Fig S4A, B). This result

suggests that the RGD motif at the N-terminal end is essential for the adhesive property of the

SNED1$^{1-260}$.

The cRGDfV peptide inhibits a panel of RGD-binding integrin heterodimers such as αvβ3, αvβ6,

α5β1, and αvβ5 (Kapp et al., 2017). Using *in silico* molecular modeling, we previously predicted

that SNED1 could potentially interact with 11 integrin subunits, α1, α4, α7, α10, α11, β1, β2,

β3, β4, β5, and β7 (Vallet et al., 2021) forming six functional heterodimers: α1β1, α4β1, α7β1,

α10β1, α11β1, and α4β7 (Hynes, 2002) including the RGD-binding integrin α5β1 (Fig 4C). We

thus sought to test whether cell adhesion to SNED1 was dependent on α5β1 integrin. We first

confirmed that LM2 and O9-1 cells expressed α5β1 integrin (Fig 4D). We then performed cell

adhesion assays in presence of functional blocking antibodies targeting β1 and α5 integrins and

demonstrated that the adhesion of LM2 cells (Fig 4E) and O9-1 cells (Fig 4F) to SNED1 was

significantly reduced in presence of antibodies targeting β1 integrin (52% and 41% decrease,

respectively, as compared to isotype controls) or α5 integrin (30% and 9% decrease, respectively,

as compared to isotype controls).

While cRGDfV inhibits α5β1 integrin, αvβ3 is more sensitive to inhibition by this peptide

(Gurrath et al., 1992; Pfaff et al., 1994). Since LM2 cells also express αvβ3 (Fig 4D), we sought

to test whether αvβ3, could act as a receptor for SNED1, although it was not predicted by our

modeling approach. Interestingly, we observed that the adhesion of LM2 cell was significantly

decreased in presence of an antibody targeting αvβ3 integrin (35.5% decrease as compared to

isotype control; Fig. 4E). This result is in line with our observation of decreased cell adhesion to

SNED1[RGE]. Altogether, these experiments identified α5β1 and αvβ3 integrin as the first SNED1

receptors.

It is worth noting that inhibition of α5, β1, or αvβ3 integrins did not fully abrogate cell adhesion

to SNED1, suggesting that there are likely additional integrin (such as the RGD-integrin αvβ1 or

the non-RGD integrin α1β1) and non-integrin SNED1 receptors at the surface of these cells.

Indeed, we have previously shown using i*n-silico* prediction, that SNED1 could interact with 55

transmembrane proteins, in addition to integrins, including the basal cell adhesion molecule

(BCAM) or dystroglycan 1 (DAG1), two known ECM receptors (Vallet et al., 2021). The

identification of these receptors at the surface of breast cancer cells and neural crest cells will be

the focus of future studies.

While our results demonstrate that the RGD motif mediates breast cancer and neural crest cell

adhesion, they also raise questions on the role of the LDV motif in SNED1. The LDV motif is

primarily recognized by α4β1 and α4β7 integrins that are mainly expressed by leukocytes. Integrin

α4 has been previously demonstrated to play a vital role in neural crest cell migration (Kil et al.,

1998) and cancer cell adhesion to the vascular endothelium (Taichman et al., 1991). Here, we show

that α4 is expressed by LM2 and O9-1 cells (Fig S5A); however, functional blocking of integrin

α4 did not affect the adhesion of LM2 cells to SNED1 (Fig S5B), in line with our observation that

these cells could adhere similarly to SNED1^{LAV} and SNED1^{WT}. This is also in line with the

observation that the two truncated forms of SNED1, SNED1$^{1-530}$ and SNED1$^{1-751}$, which contain

the LDV motif, are not sufficient to mediate cell adhesion. This opens the possibility that SNED1,

via its LDV motif, could engage other cell populations in other pathophysiological processes that

have yet to be discovered.

In addition to mediating cell adhesion, integrin heterodimers are involved in the early steps of

ECM assembly; for example, α5β1 integrin is critical for the initiation of fibronectin

fibrillogenesis (Mao and Schwarzbauer, 2005). We have previously shown that SNED1 forms

fibers in the ECM (Vallet et al., 2021), yet we do not know the mechanisms required for this

process. It will thus be interesting to determine if α5β1 integrin also mediates SNED1 fiber

assembly.

Finally, given the critical role integrin-ECM interactions play in driving pathological processes,

devising therapeutic strategies to prevent or disrupt these interactions is an active area of

investigation (Hamidi et al., 2016; Pang et al., 2023; Raab-Westphal et al., 2017). To date, seven

small molecule inhibitors and biologics have been successfully marketed, including monoclonal

antibodies specifically blocking α4β7 and α4β1 to treat inflammatory disorders such as ulcerative

colitis, Crohn's disease, and multiple sclerosis (Pang et al., 2023; Slack et al., 2022). Our study

has thus the potential to pave the way to devise novel therapeutic strategies aimed at targeting

SNED1-integrin interaction to prevent breast cancer metastasis.

MATERIALS AND METHODS

Plasmid constructs

The cDNA encoding full-length human SNED1 cloned into pCMV-XL5 was obtained from Origene (clone SC315884). 6X-His-tagged SNED1 previously described (Vallet et al., 2021) was subcloned into the bicistronic retroviral vector pMSCV-IRES-Hygromycin between the BglII and HpaI sites. FLAG-tagged SNED1 was previously described (Vallet et al., 2021).

Site-directed mutagenesis to generate SNED1[RGE] (c.C120A; p.D40E) and SNED1[LAV] (c.A932C; p.D311A) was performed using the QuikChange kit (Agilent #200519) following the manufacturer's instructions using the cDNA encoding SNED1 from Origene as a template. To obtain the double mutant SNED1[RGE/LAV] (c.C120A; p.D40E / c.A932C; p.D311A), we introduced the c.C120A mutation in the c.A932C mutant. These constructs were then subcloned into the bicistronic retroviral vector pMSCV-IRES-Hygromycin between the BglII and HpaI sites, and a 6X-Hix-tag was introduced by PCR in 3' (C-terminus of the protein).

Truncated forms of human SNED1 were subcloned by PCR to generate the following fragments: SNED1[1-530], encompassing the N-terminal region of SNED1 until the follistatin domain, and SNED1[1-751], encompassing the N-terminal region of SNED1 until the sushi domain (Fig 2A). These constructs were then subcloned into the bicistronic retroviral vector pMSCV-IRES-Hygromycin between the BglII and HpaI sites, and a FLAG tag (DYKDDDDK) was added at the C-terminus via PCR as previously described (Vallet et al., 2021). FLAG-tagged SNED1[1-260], encompassing the very N-terminal region of SNED1 was previously described (Vallet et al., 2021). The sequences of the primers used to introduce point mutations or generate truncated fragments of SNED1 are listed in Supplementary Table 1. All constructs were verified by sequencing.

Cell culture

Cell maintenance

Highly metastatic breast cancer cells, MDA-MB-231 'LM2' (termed LM2 in the manuscript), were kindly gifted by Dr. Joan Massagué (Memorial Sloan Kettering Cancer Center, New York, NY). LM2 cells, human embryonic kidney 293T cells (termed 293T in the manuscript) stably overexpressing different constructs of SNED1, and immortalized mouse embryonic fibroblasts isolated from *Sned1*[KO] mice (*Sned1*[KO] iMEFs) (Vallet et al., 2021) were cultured in Dulbecco's

Modified Eagle's medium (DMEM; Corning, #10-017-CV) supplemented with 10% fetal bovine

serum (FBS; Sigma, #F0926) and 2 mM glutamine (Corning, #25-005-CI); this formulation is

termed "complete medium" in this manuscript. The O9-1 mouse cranial neural crest cell line

(Millipore Sigma, #SCC049) was cultured per the manufacturer's instructions. Briefly, cells were

cultured on dishes coated with Matrigel® (Corning, #356234) prepared at 0.18 mg/mL in 1X

Dulbecco's Phosphate Buffered Saline (D-PBS) containing calcium and magnesium (Cytiva,

#SH30264.FS) in presence of complete ES cell medium containing 15% FBS and leukemia

inhibitory factor (Millipore Sigma, #ES-101-B) and supplemented with 25 ng/mL fibroblast

growth factor-2 (FGF-2, R&D systems, #233-FB). All cell lines were maintained at 37°C in a 5%

CO_2 humidified incubator.

Retrovirus production

293T cells were plated at ~ 30% confluency in a 6-well plate. Cells were transfected the following

day using a Lipofectamine 3000 (Invitrogen, #L3000-008) mixture containing 1 μg of a retroviral

vector with the construct of interest and 0.5 μg each of a packaging vector (pCL-Gag/Pol) and a

vector encoding the VSVG coat protein prepared in Opti-MEM™ (Gibco, #31985070). After 24h,

the transfection mixture was replaced with fresh complete medium, and cells were cultured for an

additional 24h, after which the conditioned medium containing viral particles was collected,

passed through 0.45 μ filter, and then either immediately used for cell transduction or stored at

-80°C for later use.

Generation of 293T cells stably expressing SNED1 constructs.

293T cells were seeded at ~30% confluency. The following day, cells were transduced with

undiluted viral particles-containing conditioned medium. 24h after transduction, the medium was

replaced with fresh complete medium, and cells were allowed to grow for another 24h before

selection with hygromycin (100 μg/mL). Once stable cell lines were established, we assessed the

production and secretion of recombinant proteins using immunoblotting of total cell extract (TCE)

and conditioned medium, respectively, using either an anti-SNED1, anti-His or anti-FLAG

antibody.

Protein purification

Condition medium collection

293T cells stably expressing constructs of interest were seeded in a 15 cm dish in complete medium and allowed to grow until reaching 100% confluency. The culture medium was aspirated, the monolayer was rinsed with 1X D-PBS containing calcium and magnesium, and the medium was replaced with serum-free DMEM medium supplemented with 2 mM glutamine for 48h. The serum-free conditioned medium (CM) containing secreted SNED1 was harvested, and cells were allowed to recover for 48h in complete medium before repeating the next cycle. Cells were discarded after five cycles. An EDTA-free protease inhibitor cocktail (0.067X final concentration; Thermo Scientific, #A32955) was added to the CM, and the CM was centrifuged at 4000 rpm for 10 min to remove any cell or cellular debris. Pre-cleared supernatants were collected and stored at -80°C until further processing.

Metal affinity purification of His-tagged SNED1 proteins

6X His-tagged SNED1 proteins (wild-type or integrin-binding mutants) were purified via immobilized metal affinity chromatography (IMAC) using an AKTA Pure system for fast protein liquid chromatography (FPLC) at the UIC Biophysics Core Facility. In brief, conditioned medium (CM) containing the secreted protein of interest was thawed at 4°C, concentrated using a 100-kDa protein concentrator with a polyethersulfone membrane (Thermo Scientific, #88537), buffer-exchanged against a binding buffer containing 20 mM Tris, 500 mM NaCl, 20 mM imidazole (pH 7.5), and filtered using a 0.2 μm filter. In parallel, a HisTrap HP column (column volume: 1 mL) was equilibrated with the binding buffer. The filtered CM was applied to the equilibrated HisTrap column to allow protein binding. The column was then washed with 20 column volumes of binding buffer and bound proteins were eluted with a buffer containing 20 mM Tris-HCl, 500 mM NaCl, 5 mM β-mercaptoethanol (pH 7.5), and a stepwise gradient of increasing imidazole concentration (12.5mM, 50mM, 125mM, 250mM, 375mM, and 500mM). The elution fractions were concentrated, and a buffer exchange was performed with HEPES Buffered Saline (HBS; 10 mM HEPES, 150 mM NaCl, pH 7.5).

Purification of FLAG-tagged SNED1 proteins

Since we could not achieve sufficient purity of His-tagged truncated forms of SNED1, we cloned

FLAG-tagged versions of these proteins for purification (Fig 2B) In brief, conditioned medium

containing FLAG-tagged full-length or truncated forms of SNED1 was thawed at 4°C overnight

and concentrated using protein concentrators with a 10kDa weight cut-off for the SNED1$^{1-260}$ and

SNED1$^{1-530}$ fragments (Thermo Scientific, #88535), a 30kDa cut-off for the SNED1$^{1-751}$ fragment

(Thermo Scientific, #88536), and a 100kDa cut-off for full-length SNED1 (Thermo Scientific,

#88537). An anti-FLAG resin (Sigma, #A220), containing monoclonal M2 anti-FLAG antibodies

coupled to agarose beads was washed with 20 column volumes of HBS twice. Concentrated CM

was applied to the resin and incubated overnight under constant rotation at 4°C to allow protein

binding. The following day, the unbound fraction was collected, and the resin was washed with 20

column volumes of HBS thrice. Bound FLAG-tagged proteins were eluted by competing with 200

μg/mL of FLAG peptide in HBS in a stepwise manner, resulting in 4 elution fractions. Protein

fractions were then pooled, and buffer-exchanged with HBS using protein concentrators of

appropriate molecular weight cut-off (see above) to remove the excess of unbound FLAG peptide.

Protein quantification and quality assessment

Purified proteins were quantified by measuring the absorbance at λ=280 nm using a NanoDrop

spectrophotometer. To assess their quality and purity, proteins were resolved by electrophoresis

on polyacrylamide gels. Gels were stained overnight using Coomassie-based AquaStain (Bulldog,

#AS001000) and imaged with a ChemiDoc MP™ imaging system (Bio-Rad).

Adhesion assay

Substrate coating

Adhesion assays were performed as described previously (Humphries, 1998). In brief, wells of a

96-well plate were coated with 50 μL of different concentrations, ranging from 0.1 μg/mL to 10

μg/mL, of purified human SNED1 proteins (full-length, fragments, or integrin-binding mutants)

or murine Sned1 (R&D systems, 9335-SN) for 180 min at 37°C. Fibronectin (5 μg/mL; Millipore,

#FC010) was used as a positive control. The adsorbed protein was immobilized with 0.5% (v/v)

glutaraldehyde for 15 min at room temperature (RT). To prevent cell adhesion to plastic, wells

were blocked using 200 μL of 10 mg/mL heat-denatured bovine albumin serum (BSA; Sigma,

#A9576). Uncoated wells blocked with BSA (10 mg/mL) alone were used as a negative control.

Cell seeding and crystal violet assay

Single-cell suspension of LM2 or O9-1 cells containing 5 x 10^5 cells/mL were prepared in

complete medium. 25,000 cells were added to each well and allowed to adhere for 30 min at 37°C.

Loosely attached or non-adherent cells were removed by gently washing the wells with 100 μL of

D-PBS^{++} thrice and fixed using 100 μL of 5% (v/v) glutaraldehyde for 30 minutes at RT. Cells

were stained with 100 μL of 0.1% (w/v) crystal violet in 200 mM 2-(N-morpholino)ethanesulfonic

acid (MES), pH 6.0 for 60 min at RT. After washing the excess of unbound crystal violet, the dye

was solubilized in 10% (v/v) acetic acid. Absorbance values were measured at $\lambda = 570$ nm using

a Bio-Tek Synergy HT microplate reader. Cell adhesion was determined by interpolating the

absorbance values from a standard curve that was generated by seeding cells at several dilutions

(10%-100%) from the single cell suspension on poly-L-lysine (0.01% w/v; Sigma, P4707) coated

wells and fixed directly by adding 5% (v/v) glutaraldehyde.

Integrin-blocking experiments were performed by seeding cells on a SNED1 substrate (10 μg/mL)

in presence of a cyclic RGDfV peptide (cRGDfV; Sigma, #SCP0111) at concentrations ranging

between 1.25 μM and 10 μM or in presence of 10 μg/mL of anti-β1, anti-α5, anti-α4, or anti-αvβ3

integrin-blocking antibodies (see Table S2 for detailed description of all antibodies used in this

study).

Immunoblotting

LM2 and O9-1 cells were cultured for three days in complete medium and lysed in 3x Laemmli

buffer (0.1875 M Tris-HCl, 6% SDS, 30% glycerol) containing 100 mM dithiothreitol. Cell lysates

were passed through a $26^{1/2}$-gauge needle to ensure complete cell lysis, and samples were heated

at 95ºC for 10 min. Lysates were resolved by gel electrophoresis on polyacrylamide gels at

constant current (20 mA for stacking, 25 mA for resolving). Proteins were transferred onto

nitrocellulose membranes at constant voltage (100 V) for 180 min at 4ºC. Membranes were

incubated in 5% (w/v) non-fat milk prepared in 1X PBS + 0.1% Tween-20 (PBST) for 60 min at

room temperature to prevent non-specific antibody binding and then incubated in the presence of

primary anti-β1 integrin, anti-α5 integrin, anti-αV integrin, anti-β3 integrin or anti-α4 integrin

antibodies (see Table S2 for details) in 5% (w/v) non-fat milk in PBST overnight at 4^{o}C.

Membranes were washed and incubated with horseradish peroxidase (HRP)-conjugated secondary

antibodies for 60 min at RT. Immunoreactive bands were detected using chemiluminescence

(Thermo Scientific, #E32109) and imaged with a ChemiDoc MP™ imaging system (Bio-Rad).

Data analysis and statistics

All experiments were performed with at least three technical replicates (n) and, unless noted

otherwise, repeated independently three times (N; biological replicates). Data is represented as a

mean±standard deviation from three biological replicates. Unpaired Student's two-tailed *t*-test

with Welch's correction or Welch and Brown-Forsythe one-way ANOVA with Dunnett's T3

correction for multiple comparisons was performed to measure statistical significance. Plots were

generated using PRISM (GraphPad).

ACKNOWLEDGEMENTS

We would like to thank Dr. Hyun Lee, director of the Biophysics Core facility at UIC, and her

team for their help with protein purification. We would also like to thank Dr. Sandra Pinho for

recommendations on anti-α4 integrin antibodies and all the members of the Naba lab for insightful

discussions.

COMPETING INTERESTS

The Naba laboratory holds a sponsored research agreement with Boehringer-Ingelheim for work

not related to the content of this manuscript. AN holds consulting agreements with AbbVie, XM

Therapeutics, and RA Capital.

FUNDING

This work was partly supported by the National Institute of General Medical Sciences of the

National Institutes of Health [R01CA232517], an award from the UIC Chancellor's Translational

Research Initiative, and a start-up fund from the Department of Physiology and Biophysics of the

University of Illinois Chicago to AN. NK was supported by an award from the UIC Liberal Arts

and Sciences Undergraduate Research Initiative (LASURI) and a UIC Honors College

Undergraduate Research Grant.

DATA AVAILABILITY

All relevant data can be found within the article and its supplementary information. Research

materials are available upon request to Dr. Naba.

REFERENCES

Alfandari, D., Cousin, H., Gaultier, A., Hoffstrom, B. G. and DeSimone, D. W. (2003).
Integrin α5β1 supports the migration of *Xenopus* cranial neural crest on fibronectin.
Developmental Biology **260**, 449–464.

Aumailley, M., Gurrath, M., Müller, G., Calvete, J., Timpl, R. and Kessler, H. (1991). Arg-
Gly-Asp constrained within cyclic pentapoptides Strong and selective inhibitors of cell
adhesion to vitronectin and laminin fragment P1. *FEBS Letters* **291**, 50–54.

Barqué, A., Jan, K., De La Fuente, E., Nicholas, C. L., Hynes, R. O. and Naba, A. (2021).
Knockout of the gene encoding the extracellular matrix protein SNED1 results in early
neonatal lethality and craniofacial malformations. *Developmental Dynamics* **250**, 274–
294.

Bökel, C. and Brown, N. H. (2002). Integrins in Development: Moving on, Responding to, and
Sticking to the Extracellular Matrix. *Developmental Cell* **3**, 311–321.

Campbell, I. D. and Humphries, M. J. (2011). Integrin Structure, Activation, and Interactions.
Cold Spring Harb Perspect Biol **3**, a004994.

Chen, M. B., Lamar, J. M., Li, R., Hynes, R. O. and Kamm, R. D. (2016). Elucidation of the
Roles of Tumor Integrin β1 in the Extravasation Stage of the Metastasis Cascade. *Cancer*
Research **76**, 2513–2524.

Cherny, R. C., Honan, M. A. and Thiagarajan, P. (1993). Site-directed mutagenesis of the
arginine-glycine-aspartic acid in vitronectin abolishes cell adhesion. *Journal of*
Biological Chemistry **268**, 9725–9729.

Cox, T. R. (2021). The matrix in cancer. *Nat Rev Cancer* **21**, 217–238.

Doyle, A. D., Nazari, S. S. and Yamada, K. M. (2022). Cell-extracellular matrix dynamics.
Phys Biol **19**,.

Duband, J.-L., Dufour, S., Yamada, S. S., Yamada, K. M. and Thiery, J. P. (1991). Neural
crest cell locomotion induced by antibodies to β1 integrins A tool for studying the roles of
substratum molecular avidity and density in migration. *Journal of Cell Science* **98**, 517–
532.

Dzamba, B. J. and DeSimone, D. W. (2018). Chapter Seven - Extracellular Matrix (ECM) and
the Sculpting of Embryonic Tissues. In *Current Topics in Developmental Biology* (ed.
Litscher, E. S.) and Wassarman, P. M.), pp. 245–274. Academic Press.

Frisch, S. M. and Francis, H. (1994). Disruption of epithelial cell-matrix interactions induces
apoptosis. *J Cell Biol* **124**, 619–626.

Gallik, K. L., Treffy, R. W., Nacke, L. M., Ahsan, K., Rocha, M., Green-Saxena, A. and
Saxena, A. (2017). Neural crest and cancer: Divergent travelers on similar paths.
Mechanisms of Development **148**, 89–99.

Gurrath, M., Müller, G., Kessler, H., Aumailley, M. and Timpl, R. (1992).
Conformation/activity studies of rationally designed potent anti-adhesive RGD peptides.
European Journal of Biochemistry **210**, 911–921.

Hamidi, H. and Ivaska, J. (2018). Every step of the way: integrins in cancer progression and
metastasis. *Nat Rev Cancer* **18**, 533–548.

Hamidi, H., Pietilä, M. and Ivaska, J. (2016). The complexity of integrins in cancer and new
scopes for therapeutic targeting. *Br. J. Cancer* **115**, 1017–1023.

Herrera, J., Henke, C. A. and Bitterman, P. B. (2018). Extracellular matrix as a driver of
progressive fibrosis. *J Clin Invest* **128**, 45–53.

Hood, J. D. and Cheresh, D. A. (2002). Role of integrins in cell invasion and migration. *Nature*
Reviews Cancer **2**, 91–100.

Humphries, M. J. (1998). Cell-Substrate Adhesion Assays. *Current Protocols in Cell Biology*
00, 9.1.1-9.1.11.

Hynes, R. O. (2002). Integrins: Bidirectional, Allosteric Signaling Machines. *Cell* **110**, 673–687.

Hynes, R. O. (2009). The Extracellular Matrix: Not Just Pretty Fibrils. *Science* **326**, 1216–1219.

Hynes, R. O. and Naba, A. (2012). Overview of the Matrisome—An Inventory of Extracellular
Matrix Constituents and Functions. *Cold Spring Harbor Perspectives in Biology* **4**,
a004903.

Kanchanawong, P. and Calderwood, D. A. (2023). Organization, dynamics and
mechanoregulation of integrin-mediated cell–ECM adhesions. *Nat Rev Mol Cell Biol* **24**,
142–161.

Kapp, T. G., Rechenmacher, F., Neubauer, S., Maltsev, O. V., Cavalcanti-Adam, E. A.,
Zarka, R., Reuning, U., Notni, J., Wester, H.-J., Mas-Moruno, C., et al. (2017). A
Comprehensive Evaluation of the Activity and Selectivity Profile of Ligands for RGD-
binding Integrins. *Sci Rep* **7**, 39805.

Kil, S. H., Krull, C. E., Cann, G., Clegg, D. and Bronner-Fraser, M. (1998). The α4Subunit
of Integrin Is Important for Neural Crest Cell Migration. *Developmental Biology* **202**, 29–
42.

Lawler, J., Weinstein, R. and Hynes, R. O. (1988). Cell attachment to thrombospondin: the
role of ARG-GLY-ASP, calcium, and integrin receptors. *J Cell Biol* **107**, 2351–2361.

Leimeister, C., Schumacher, N., Diez, H. and Gessler, M. (2004). Cloning and expression
analysis of the mouse stroma marker Snep encoding a novel nidogen domain protein.
Developmental Dynamics **230**, 371–377.

Leonard, C. E. and Taneyhill, L. A. (2020). The road best traveled: Neural crest migration
upon the extracellular matrix. *Semin Cell Dev Biol* **100**, 177–185.

Mankarious, L. A. and Goudy, S. L. (2010). Craniofacial and upper airway development.
Paediatric Respiratory Reviews **11**, 193–198.

Mao, Y. and Schwarzbauer, J. E. (2005). Fibronectin fibrillogenesis, a cell-mediated matrix
assembly process. *Matrix Biology* **24**, 389–399.

Martik, M. L. and Bronner, M. E. (2021). Riding the crest to get a head: neural crest evolution
in vertebrates. *Nat Rev Neurosci* **22**, 616–626.

Mayor, R. and Theveneau, E. (2013). The neural crest. *Development* **140**, 2247–2251.

Naba, A. (2024). Mechanisms of assembly and remodelling of the extracellular matrix. *Nat Rev
Mol Cell Biol*.

Naba, A., Clauser, K. R., Lamar, J. M., Carr, S. A. and Hynes, R. O. (2014). Extracellular
matrix signatures of human mammary carcinoma identify novel metastasis promoters.
eLife **3**, e01308.

Pally, D. and Naba, A. (2024). Extracellular matrix dynamics: A key regulator of cell migration
across length-scales and systems. *Curr Opin Cell Biol* **86**, 102309.

Pang, X., He, X., Qiu, Z., Zhang, H., Xie, R., Liu, Z., Gu, Y., Zhao, N., Xiang, Q. and Cui, Y.
(2023). Targeting integrin pathways: mechanisms and advances in therapy. *Signal
Transduction and Targeted Therapy* **8**, 1.

**Pfaff, M., Tangemann, K., Müller, B., Gurrath, M., Müller, G., Kessler, H., Timpl, R. and
Engel, J.** (1994). Selective recognition of cyclic RGD peptides of NMR defined
conformation by alpha IIb beta 3, alpha V beta 3, and alpha 5 beta 1 integrins. *Journal of
Biological Chemistry* **269**, 20233–20238.

Pickup, M. W., Mouw, J. K. and Weaver, V. M. (2014). The extracellular matrix modulates the
hallmarks of cancer. *EMBO reports* **15**, 1243–1253.

**Pietri, T., Eder, O., Breau, M. A., Topilko, P., Blanche, M., Brakebusch, C., Fässler, R.,
Thiery, J.-P. and Dufour, S.** (2004). Conditional β1-integrin gene deletion in neural crest
cells causes severe developmental alterations of the peripheral nervous system.
Development **131**, 3871–3883.

Pytela, R., Pierschbacher, M. D. and Ruoslahti, E. (1985). Identification and isolation of a 140
kd cell surface glycoprotein with properties expected of a fibronectin receptor. *Cell* **40**,
191–198.

Raab-Westphal, S., Marshall, J. F. and Goodman, S. L. (2017). Integrins as Therapeutic
Targets: Successes and Cancers. *Cancers* **9**, 110.

Rozario, T. and DeSimone, D. W. (2010). The extracellular matrix in development and
morphogenesis: A dynamic view. *Developmental Biology* **341**, 126–140.

Ruoslahti, E. and Pierschbacher, M. D. (1987). New perspectives in cell adhesion: RGD and
integrins. *Science* **238**, 491–497.

Slack, R. J., Macdonald, S. J. F., Roper, J. A., Jenkins, R. G. and Hatley, R. J. D. (2022).
Emerging therapeutic opportunities for integrin inhibitors. *Nature Reviews Drug*
Discovery **21**, 60–78.

Taichman, D. B., Cybulsky, M. I., Djaffar, I., Longenecker, B. M., Teixidó, J., Rice, G. E.,
Aruffo, A. and Bevilacqua, M. P. (1991). Tumor cell surface alpha 4 beta 1 integrin
mediates adhesion to vascular endothelium: demonstration of an interaction with the N-
terminal domains of INCAM-110/VCAM-1. *Cell Regulation* **2**, 347–355.

Testaz, S., Delannet, M. and Duband, J. (1999). Adhesion and migration of avian neural crest
cells on fibronectin require the cooperating activities of multiple integrins of the (beta)1
and (beta)3 families. *J Cell Sci* **112 (Pt 24)**, 4715–4728.

Trainor, P. A. (2005). Specification and patterning of neural crest cells during craniofacial
development. *Brain Behav Evol* **66**, 266–280.

Tucker, G. C., Duband, J. L., Dufour, S. and Thiery, J. P. (1988). Cell-adhesion and substrate-
adhesion molecules: their instructive roles in neural crest cell migration. *Development*
103 Suppl, 81–94.

Vallet, S. D., Davis, M. N., Barqué, A., Thahab, A. H., Ricard-Blum, S. and Naba, A. (2021).
Computational and experimental characterization of the novel ECM glycoprotein SNED1
and prediction of its interactome. *Biochemical Journal* **478**, 1413–1434.

Walma, D. A. C. and Yamada, K. M. (2020). The extracellular matrix in development.
Development **147**, dev175596.

Yang, J. T., Rayburn, H. and Hynes, R. O. (1993). Embryonic mesodermal defects in α5
integrin-deficient mice. *Development* **119**, 1093–1105.

562
563

564 **FIGURE LEGENDS**
565

Figure 1. SNED1 promotes breast cancer and neural crest cell adhesion

(A) Adhesion of MDA-MB-231 'LM2' breast cancer cells to purified human SNED1 is

concentration-dependent.

(B) Adhesion of O9-1 mouse neural crest cells to purified murine Sned1 is concentration

dependent. Data is represented as mean ± SD from three biological experiments.

Figure 2. The N-terminal region of SNED1 mediates cell adhesion

(A) Schematic showing FLAG-tagged full-length SNED1 and the different truncated forms of

SNED1 used in this study: SNED1$^{1\text{-}751}$ encompasses the N-terminal region until the sushi domain;

SNED1$^{1\text{-}530}$ encompasses the N-terminal region until the follistatin domain; SNED1$^{1\text{-}260}$

encompasses the N-terminal region until the NIDO domain.

(B) Coomassie-stained gel showing the purity of purified full-length and truncated forms of

SNED1.

(C-D) Adhesion of MDA-MB-231 'LM2' breast cancer cells (C) and O9-1 neural crest cells (D)

to full-length and truncated forms of SNED1. Data is represented as mean ± SD from at least two

biological experiments. Welch and Brown-Forsythe one-way ANOVA with Dunnett's T3

correction for multiple comparisons was performed to determine statistical significance. ns: non-

significant, *p<0.05, *** p<0.001.

Figure 3. The RGD motif in SNED1 is required to mediate cell adhesion

(A) Schematic of SNED1 showing the two putative integrin-binding motifs in SNED1 (^{38}RGD and

^{310}LDV) and the point mutations introduced to disrupt their integrin-binding activity

(^{38}RGD>RGE, ^{310}LDV>LAV).

(B, C) Adhesion of MDA-MB-231 'LM2' breast cancer cells (B) and O9-1 mouse neural crest

cells (C) to wild type or integrin-binding mutants (RGE, LAV or RGE/LAV) of SNED1. Data is

represented as mean ± SD from three biological experiments. Welch and Brown-Forsythe one-

way ANOVA with Dunnett's T3 correction for multiple comparisons was performed to determine

statistical significance. ns: non-significant, *p<0.05, ** p<0.01.

Figure 4. Functional blocking of integrins decreases cell adhesion to SNED1

(A-B) Adhesion of MDA-MB-231 'LM2' breast cancer cell (A) and O9-1 neural crest cells (B) to

SNED1 is inhibited in presence of increasing concentrations of cRGDfV peptide. Data is

represented as mean ± SD from three biological experiments. Welch and Brown-Forsythe one-

way ANOVA with Dunnett's T3 correction for multiple comparisons was performed to determine

statistical significance. *$p < 0.05$, **$p < 0.01$.

(C) Schematic showing the seven integrin heterodimers previously predicted to interact with

SNED1. Adapted from (Vallet et al., 2021).

(D) Immunoblots on total cell extract from MDA-MB-231 'LM2' and O9-1 cells shows β1

integrin, α5 integrin, αv integrin and β3 integrins expression.

(E) Adhesion of MDA-MB-231 'LM2' breast cancer cells to SNED1 is decreased in presence of

anti-β1, anti-α5 and anti-αVβ3 integrin-blocking antibodies.

(F) Adhesion of O9-1 mouse neural crest cells to SNED1 is decreased in presence of anti-β1 and

anti-α5 integrin blocking antibodies. Data is represented as mean ± SD from three biological

experiments. Unpaired Student's two-tailed t-test with Welch's correction was performed to

determine statistical significance. *$p < 0.05$, ** $p < 0.01$.

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. Cell adhesion on human SNED1 and murine Sned1

(A) Coomassie-blue-stained polyacrylamide gel showing the quality and purity of affinity-purified

full-length SNED1-His.

(B) Murine O9-1 neural crest cells adhere to the same extent to human SNED1 and murine Sned1.

Data is represented as mean ± SD from three biological experiments. Unpaired Student's two-

tailed t-test with Welch's correction was performed to test statistical significance. ns: non-

significant.

(C) Graph showing the adhesion of immortalized embryonic fibroblast cells isolated from *Sned1*

knockout mice (*Sned1*[KO] iMEF) on increasing concentrations of Sned1.

Figure S2. LM2 cell adhesion on equimolar concentration of full length SNED1 and

SNED1[1-260]

Graph showing LM2 cell adhesion on 66.3 μM of full length SNED1 and SNED1[1-260].

Figure S3. Purification of integrin-binding mutants of SNED1

Coomassie-blue-stained polyacrylamide gels showing the quality and purity of affinity-purified

His-tagged SNED1[RGE] (A), SNED1[LAV] (B), and SNED1[RGE/LAV] (C). The different bands

correspond to different levels of glycosylation of the proteins, as previously shown (Vallet et al.,

2021).

Figure S4. The RGD motif in SNED1[1-260] is sufficient to mediate cell adhesion

Adhesion of MDA-MB-231' LM2' breast cancer cells (A) and O9-1 neural crest cells (B) to the

N-terminal fragment of SNED1 (SNED1[1-260]) is significantly decreased in presence of the integrin-

binding cRGDfV peptide. Data is represented as mean ± SD from three biological experiments.

Unpaired Student's two-tailed t-test with Welch's correction was performed to determine

statistical significance. *p<0.05, ** p<0.01.

Figure S5. Functional blocking of integrin α4 does not affect breast cancer cell adhesion on

SNED1

(A) Immunoblot on total cell extract from MDA-MB-231' LM2' and O9-1 cells showing α4

integrin expression.

(B) Adhesion of MDA-MB-231' LM2' breast cancer cells to SNED1 is not altered in presence of

anti-α4 integrin-blocking antibody. Data is represented as mean ± SD from three biological

experiments. Unpaired Student's two-tailed t-test with Welch's correction was performed to test

statistical significance. ns: non-significant.

Figure S6. Immunoblot transparency

Uncropped immunoblots for β1 integrin, α5 integrin, αv integrin, β3 integrin, and α4 integrin.

Supplementary Table 1: List of primers

Primer	Sequence	Purpose
BglII-KOZAK-ATG-SNED1Hs_F	5' TCGAAGATCTGCCACCATGCGGCACGGCGTCGC 3'	Subcloning SNED1 into retroviral pMSCV-IRES-Hygro vector
HpaI-Stop-His-SNED1Hs_R	5'GATGTTAACTTAGTGGTGATGGTGATGATGAGATTTCTCCAGTGTCTGACTCTTACT 3'	Shuttling of SNED1-His WT, RGE, LAV, or GE/LAV) into retroviral pMSCV-IRES-Hygro vector
HpaI-Stop-FLAG-NFS_R	5' CCGTTAACTTACTTGTCGTCATCGTCTTTGTAGTCGCACTGGGGAGGCTCAC 3'	Subcloning SNED1$^{1-571}$ into retroviral pMSCV-IRES-Hygro vector with the addition of a C-terminal FLAG tag
HpaI-Stop-FLAG-NF_R	5' CCGTTAACTTACTTGTCGTCATCGTCTTTGTAGTCGACGCAGAGGTAGCTCCC 3'	Subcloning SNED1$^{1-530}$ into retroviral pMSCV-IRES-Hygro vector with the addition of a C-terminal FLAG tag
SNED1_c120a _F	5' CCGAGCGCGGCGAAGCCGTCACC 3'	Introduction of the c120>a (pG40E) point mutation to generate SNED1^{RGE}
SNED1_c120a _R	5' GGTGACGGCTTCGCCGCGCTCGG 3'	
SNED1_a932c_F	5' GGAGGTGCCACCTGGCCGTGAACGAATGTGC 3'	Introduction of the a932>c (pD311A) point mutation to generate SNED1^{LAV}
SNED1_a932c_R	5' GCACATTCGTTCACGGCCAGGTGGCACCTCC 3'	

656 **Supplementary Table 2: List of antibodies used for functional blocking and immunoblotting**

Antibody	Host species	Reactivity	Application	Concentration used (µg/mL)	Catalog #
Anti- β1 Integrin	Rat	Human	Functional blocking	10	Sigma, MABT821
Anti-α5 Integrin	Rat	Human	Functional blocking	10	Sigma, MABT820
Anti-α4 Integrin	Mouse	Human	Functional blocking	10	Sigma, MAB1383
Anti-β1 Integrin	Hamster	Mouse	Functional blocking	10	BD Biosciences, BDB555002
Anti-α5 Integrin	Rat	Mouse	Functional blocking	10	Biolegend, 103817
Anti-αvβ3 Integrin	Mouse	Human	Functional blocking	10	Sigma, MAB1976Z
Rat IgG	Rat	Isotype control	Functional blocking	10	Invitrogen, PI31903
Mouse IgG	Mouse	Isotype control	Functional blocking	10	Invitrogen, PI31933
Hamster IgM	Hamster	Isotype control	Functional blocking	10	Biolegend, 401014
Anti-SNED1	Rabbit	Human	Immunoblotting	1	Naba lab
Anti-His	Mouse	-	Immunoblotting	1	Sigma, SAB2702218
Anti-FLAG	Rabbit	-	Immunoblotting	1	Sigma, F7425
Anti-β1 Integrin sera	Rabbit	Human, Mouse	Immunoblotting	1:1000 dilution	In-house
Anti-α5 Integrin	Rabbit	Human, Mouse	Immunoblotting	0.217	Abcam, AB150361
Anti-α4 Integrin	Rabbit	Human, Mouse	Immunoblotting	0.5	Invitrogen, MA5-27947
Anti-αv Integrin	Rabbit	Human, Mouse	Immunoblotting	1	Invitrogen, MA5-32195
Anti-β3 Integrin	Rabbit	Human, Mouse	Immunoblotting	1	Invitrogen, MA5-32077

Figure 1. SNED1 promotes breast cancer and neural crest cell adhesion

A)

B)

Figure 2. The N-terminal region of SNED1 mediates cell adhesion

Figure 3. The RGD motif in SNED1 is required for cell adhesion

Figure 4: Functional blocking of integrins decreases cell adhesion to SNED1

Figure S1. Cell adhesion on human SNED1 and murine Sned1

Figure S2. LM2 cell adhesion on equal molar concentration of full length SNED1 and SNED1[1-260]

A)

MDA-MB-231 'LM2'

Figure S3. Purification of integrin binding mutants of SNED1

A) **B)** **C)**

Figure S4. The RGD motif in SNED1$^{1\text{-}260}$ is required for cell adhesion

Figure S5. Functional blocking of α4 integrin does not affect breast cancer cell adhesion to SNED1

A)

B)

Figure S6. Immunoblot transparency

Figure 4D
β1 integrin

Figure 4D
α5 integrin

Figure 4D
αv integrin

Figure 4D
β3 integrin

Figure S5A
α4 integrin