1

Histone Variant H3.3 Mediates DNA Repair and cGAS-STING Pathway Activation in

Telomere Dysfunction

Chiao-Ming Huang[1] and Liuh-Yow Chen[1,2,3,*]

[1] Molecular and Cell Biology, Taiwan International Graduate Program, Academia

Sinica and Graduate Institute of Life Sciences, National Defense Medical Center,

Taipei City, Taiwan

[2] Institute of Molecular Biology, Academia Sinica, Taipei, 11529, Taiwan

[3] Lead contact

[*] Correspondence: lyowchen@gate.sinica.edu.tw

ABSTRACT

The telomere damage response is a critical mechanism that regulates cellular senescence. Deprotected telomeres activate the cytosolic DNA sensing cGAS-STING pathway, leading to cellular senescence. Our previous studies revealed that extrachromosomal telomere repeats (ECTRs) activate the cGAS-STING pathway, and this process is attenuated by histone H3.3 depletion. However, the role of histone H3.3 in telomere deprotection-dependent cGAS-STING pathway activation remains unclear. Here, we discover that histone H3.3 is required for cGAS-STING pathway activation in cells with deprotected telomeres. Expression of the TRF2 dominant-negative mutant, TRF2ΔBΔM, induces telomere dysfunction in fibroblast cells, triggering cGAS-STING pathway activation and growth inhibition. Histone H3.3 depletion significantly reduces this activation, highlighting its critical role in linking telomere deprotection to the cGAS-STING-mediated innate immune signaling. Furthermore, we assess the signaling cascade in triggering telomeric DNA damaging response (DDR), and reveal that histone H3.3 regulates cGAS-STING signaling through controlling the telomeric DDR and repair pathways. Additionally, depletion of histone H3.3 chaperones, including ATRX, DAXX, and HIRA, inhibits telomere fusion and cGAS-STING pathway activation, underscoring the role of histone H3.3 in telomere maintenance and DDR. Collectively, our study establishes histone H3.3 as a key regulator of telomeric DDR and telomere dysfunction-induced cGAS-STING pathway activation, emphasizing the importance of this pathway in cellular senescence.

Keywords

Histone H3 Variant H3.3, Telomere Deprotection, Telomere DNA damage response

INTRODUCTION

Telomeres and their associated proteins are crucial for protecting chromosome ends. Deprotection of telomeres triggers a DNA damage response (DDR), leading to replicative senescence[1-3]. Among the telomere-binding proteins, telomeric repeat binding factor 2 (TERF2, TRF2) plays a critical role in preventing DDR at chromosome ends. Depletion of TRF2 from telomeres induces cellular senescence and chromosome end-to-end fusion[2], an event that is often found when telomeres are deprotected, particularly under conditions involving dominant-negative TRF2 expression[2]. Previous studies have demonstrated that the absence of wild-type TRF2 activates a DDR mediated by the ataxia telangiectasia mutated (ATM) pathway[2-4]. Upstream components of ATM, including the double-strand break (DSB) sensing complex, the meiotic recombination 11 (MRE11)-radiation sensitive 50 (RAD50)-Nijmegen breakage syndrome 1 (NBS1, NBN) (MRN) complex, play a crucial role in sensing TRF2-dependent telomere deprotection[5-7]. Conversely, downstream effector proteins of ATM, such as ligase IV (LIG4), and x-ray repair cross-complementing 4 (XRCC4), mediate the non-homologous end joining (NHEJ) signaling pathways induced by TRF2 depletion and subsequent telomere fusion[2,8,9]. Together, these findings establish a well-characterized ATM-dependent mechanism underlying telomere deprotection and the induction of cellular senescence.

Beyond ATM-dependent mechanisms, recent studies have implicated the cGAS-STING pathway in telomere dysfunction[10]. In this pathway, cyclic GMP-AMP (cGAMP) synthase (cGAS) produces cGAMP to activate stimulator of interferon response cGAMP interactor (STING) and downstream phosphorylation cascade involving

TANK-binding kinase 1 (TBK1), STING, and interferon regulatory factor 3 (IRF3),

which drives the innate immune response by promoting type I IFN transcription[11-19].

Studies demonstrate that telomere-damaging agents and defective telomere-binding

proteins can lead to micronuclei formation, thereby activating the cGAS-STING

pathway[10]. Furthermore, the mutated TRF2, TRF2ΔB induces the formation of ECTRs,

which released into the cytosol to trigger the cGAS-STING-mediated innate immune

response[20]. These findings suggest that telomere deprotection activates both ATM

dependent and cGAS-STING pathway, contributing to cellular senescence.

At telomeres, the specific histone variant, histone H3.3, is enriched and plays a crucial

role in telomere maintenance[21-23]. Histone H3.3 is deposited in a replication-

independent manner by the histone cell cycle regulator (HIRA) complex on

euchromatin and by the alpha-thalassemia/mental retardation syndrome X-linked

(ATRX)-death domain associated protein (DAXX) complex on heterochromatin,

centromeres, and telomeres[22]. While histone H3.3 is known to regulate the active

expression of genes[24-28], its function at telomeres remains unclear. Mutations in

histone H3.3, such as histone H3.3^{K27M} and histone H3.3^{G34V}, are associated with

glioblastoma and other telomerase-negative cancers that utilize the alternative

lengthening of telomeres (ALT) pathway[29-32]. ALT, a recombination-based telomere

elongation mechanism, is characterized by reduced ATRX expression[32-37] and

impaired cGAS-STING pathway activation[20]. Given the role of ATRX-DAXX in

depositing histone H3.3 at telomeres, mutations in ATRX, DAXX, and histone H3.3

may alter telomere dynamics and contribute to ALT-associated cancer

development[33,38-40]. These observations underscore histone H3.3's potential roles in

regulating telomere maintenance and its association with ALT mechanisms.

Previously, we reported that depletion of histone H3.3, ATRX, and DAXX impairs the

cGAS-STING pathway activated by ECTR[20]. Our findings indicated that histone H3.3

participated in the telomere-associated cGAS-STING pathway activation. Additionally,

histone H3.3 has been shown to participate in DNA damage repair[41], yet the

relationship of histone H3.3's roles in regulating DNA damage responses at telomeres

and activating the cGAS-STING pathway remains unclear. Here, we demonstrated a

role of histone H3.3 in telomere deprotection, the activation of the cGAS-STING

pathway, and deprotected telomere-induced cell growth inhibition.

RESULTS

TRF2ΔBΔM expression drives cGAS-mediated growth inhibition

To investigate the signaling pathway induced by the telomeric deprotection, we focused on exploring the downstream cellular response in cells lacking TRF2. We developed an inducible expression system using a Tet-On mechanism to express dominant-negative mutant TRF2ΔBΔM in telomerase-immortalized BJ human fibroblasts (BJ^t), under the control of a tetracycline-dependent promoter. TRF2ΔBΔM, a deletion mutant of TRF2 lacking its N-terminal basic domain and C-terminal DNA binding domain, disrupts the formation of a functional telomere shelterin complex by displacing endogenous TRF2, leading to telomere deprotection and fusion[2]. Upon adding doxycycline (DOX), we observed the expression of myc-tagged TRF2ΔBΔM (Fig. 1A) and the formation of telomere dysfunction-induced foci (TIFs), characterized by the colocalization of phosphorylated histone H2AX (γH2AX) with telomeres (Fig. 1B) in BJ^t cells.

In line with our aims, we observed the activation of cGAS-STING response following TRF2ΔBΔM induction. Notably, STING phosphorylation, a critical event for the cGAS-STING pathway that facilitates the phosphorylation of IRF3 to drive IFN expression[11,12], was detected on day 2 and peaked on day 3 post-DOX treatment, indicating that cGAS-STING pathway was activated (Fig. S1A). This activation correlated with the induction of TRF2ΔBΔM expression (Fig. S1A). Reverse-transcription quantitative polymerase chain reaction (RT-qPCR) revealed upregulated expression of IFN beta (IFNβ) mRNA after induced expression of TRF2ΔBΔM (Fig. S1B). These findings

suggest that induced expression of TRF2ΔBΔM drives downstream responses

including the STING phosphorylation and the transcription of IFNβ.

To determine whether cGAS plays a role in STING activation upon TRF2ΔBΔM

expression, we depleted it using small interfering RNAs (siRNAs) before DOX

induction. Depletion of cGAS inhibited STING phosphorylation and IFNβ transcription

in the TRF2ΔBΔM-expressing cells (Fig. 1C, D), supporting the conclusion that

TRF2ΔBΔM activates the cGAS-STING signaling cascade to promote IFNβ production.

These results are reminiscent of those from previous studies indicating that TRF2

inhibition activates the cGAS-STING pathway[10,20].

Furthermore, previous reports have described telomere deprotection induced by

TRF2ΔBΔM expression as a trigger for cell growth arrest and senescence[2].

Consistently, we also observed that DOX-induced TRF2ΔBΔM expression in BJ[t] cells

led to significantly reduced cell growth (Fig. 1E). Since cGAS-STING pathway

activation induces IFNβ expression and affects cell proliferation[20], we hypothesized

that the cell growth defects associated with TRF2ΔBΔM expression might be driven

by cGAS-STING pathway activation. To test this possibility, we transfected BJ[t] cells

with siRNAs to silence cGAS expression and monitored their subsequent cell growth.

Our results demonstrated that cGAS depletion partially rescued the cell proliferation

defects of BJ[t] cells expressing TRF2ΔBΔM (Fig. 1F), indicating that induction of the

cGAS-STING pathway contributes to TRF2ΔBΔM-mediated cell growth defects. Thus,

the cGAS-STING pathway exerts a fundamental role in mediating cellular growth

impairment due to telomere deprotection.

Histone H3.3 regulates the cGAS-STING-pathway-mediated cell growth inhibition upon telomere dysfunction

Previously, we showed that histone H3.3 and its chaperone, the ATRX-DAXX complex, promote the activation of cGAS-STING pathway[20]. This activation occurs in response to the accumulation of ECTRs following artificially induced telomere excision by another TRF2 mutant, TRF2ΔB, which lacks only the N-terminal basic domain[20]. This finding demonstrates that the release of telomeric DNA into the cytosol, a phenomenon commonly observed in ALT cancers[42-44], activates cGAS-STING pathway. Given that ectopic expression of either TRF2ΔB or TRF2ΔBΔM elicits TRF2 dysfunction, we sought to investigate if histone H3.3 is involved in the cGAS-STING pathway activation driven by TRF2ΔBΔM-induced telomere deprotection. To do so, we used siRNAs to deplete histone H3.3 from BJ^t cells and then examined TRF2ΔBΔM-induced STING phosphorylation and IFNβ production. Our results revealed that histone H3.3 depletion significantly reduced STING phosphorylation and IFNβ expression in BJ^t cells expressing TRF2ΔBΔM (Fig. 2A, B). Furthermore, siRNA-mediated histone H3.3 silencing alleviated the phenotype of defective cell proliferation associated with TRF2ΔBΔM (Fig. 2C). Importantly, the phosphorylation of STING can be rescued by ectopically expressing the siRNA-insensitive H3.3 (Fig. 2D). Collectively, our findings suggest that histone H3.3 mediates cGAS-STING pathway activation and the subsequent cell growth impairment arising from telomere deprotection.

Telomere fusion by non-homologous end joining (NHEJ) promotes cGAS-STING pathway activation

Telomere deprotection resulting from TRF2 deficiency leads to telomere fusion via the

NHEJ pathway[2,9,45], which in turn generates micronuclei[10,46] and activates the cGAS-

STING pathway[10]. While some studies debate whether micronucleus formation

directly triggers cGAS-STING pathway activation[47], we hypothesized that cGAS-

STING pathway activation following TRF2ΔBΔM expression may result from telomere

fusion. To test this hypothesis, we used siRNAs to silence the expression of the LIG4

and XRCC4, two key proteins that mediate telomere fusion via the NHEJ mechanism[9].

We found that depletion of LIG4 or XRCC4 in BJ^t cells reduced TRF2ΔBΔM-induced

telomere fusion, as evidenced by the diminished apparent convergence of telomere

signals at chromosome ends (Fig. 3A, B). Furthermore, silencing LIG4 or XRCC4

decreased the formation of abnormal nuclear structures, including nucleoplasmic

bridges, nuclear buds, and micronuclei (Fig. 3C, D). Notably, LIG4 or XRCC4 depletion

also inhibited STING phosphorylation and IFNβ mRNA induction in BJ^t cells

expressing TRF2ΔBΔM (Fig. 4). These results suggest that activation of the cGAS-

STING pathway in response to TRF2ΔBΔM expression may be attributable to NHEJ-

mediated fusion of deprotected telomeres.

Histone H3.3 mediates DDR and telomere fusion for cGAS-STING pathway

activation

As shown above, histone H3.3 plays a role in activating the cGAS-STING pathway in

BJ^t cells with deprotected telomeres (Fig. 2). We hypothesized that histone H3.3

contributes to telomere fusion, which in turn influences cGAS-STING pathway

activation. To examine this possibility, we silenced histone H3.3 expression in BJ^t cells

by siRNAs and analyzed subsequent telomere fusion and nuclear structural

abnormalities following TRF2ΔBΔM induction for two days. Our results demonstrate

that depletion of histone H3.3 significantly reduced the number of telomere fusion

events and partially reduced the formation of micronuclei, nuclear buds, and

nucleoplasmic bridges typically induced by TRF2ΔBΔM expression (Fig. 3). These

findings indicate that histone H3.3 is required for NHEJ-mediated telomere fusion

following telomere deprotection. Our results reveal that chromosomal and nuclear

abnormalities occur on the second day after inducting TRF2 mutant expression.

Notably, this precedes the peak cGAS-STING response, which occurs on the third day

post-induction of TRF2ΔBΔM expression (Fig. 3, S1). These observations suggest that

histone H3.3 orchestrates the downstream response to TRF2ΔBΔM-induced telomere

deprotection, including telomere fusion, nuclear abnormalities, and subsequent cGAS-

STING pathway activation. Since depletion of cGAS did not rescue TRF2ΔBΔM-

mediated telomere fusion or nuclear abnormalities (Fig. 3), cGAS does not appear to

directly participate in DNA repair at deprotected telomeres but rather functions

downstream of telomere fusion and nuclear abnormalities. Thus, our results suggest

that histone H3.3 mediates cGAS-STING activation by promoting NHEJ-mediated

repair of deprotected telomeres.

Next, we investigated if histone H3.3 regulates the telomere DDR and thereby controls

NHEJ for subsequent cGAS-STING activation. As evidenced above, expression of

TRF2ΔBΔM elicited moderate TIF formation (Fig. 1B, 5A-C), indicating that telomere

deprotection can trigger a DDR. To determine whether histone H3.3 regulates the

telomeric DDR, we treated BJ^t cells with siRNAs to deplete the expression of histone

H3.3. Interestingly, we observed that depletion of histone H3.3 also reduced TIF

formation in the TRF2ΔBΔM-expressing cells (Fig. 5A-C), supporting that the role of

histone H3.3 in regulating the DDR at deprotected telomeres. These findings

underscore the role of histone H3.3 in regulating the TRF2ΔBΔM-induced telomeric

DDR.

Since NBS1 is a key component of the MRN complex, we hypothesized that its

depletion would impair telomeric DDR. The results showed that TRF2ΔBΔM-induced

TIF formation was abolished upon NBS1 depletion (Fig. 5A-C), consistent with the

MRN complex playing a role in sensing telomere deprotection to activate a DDR[48]. To

investigate whether the MRN complex regulates cGAS-STING activation, we

measured IFNβ production in cells treated with the MRN inhibitor, Mirin[49]. Treatment

with 25 μM Mirin significantly reduced STING phosphorylation and IFNβ transcription

in TRF2ΔBΔM-expressing cells, confirming that MRN is essential for cGAS-STING

activation following telomere deprotection (Fig. 5D, E). Our observations further

demonstrate that the cellular machinery responsible for telomeric DDR also modulates

the activation of cGAS-STING pathway. Collectively, these results indicate that

histone H3.3 and MRN complex facilitate DDR activation and NHEJ at deprotected

telomeres, thereby priming cells for subsequent cGAS-STING pathway activation.

Histone H3.3's chaperones are required for TRF2ΔBΔM-driven cGAS-STING

pathway activation, telomere fusion, and TIF formation

Histone chaperones, including ATRX-DAXX and HIRA, have been shown to mediate

the deposition of histone H3.3 on telomeres[22,50]. Anti-silencing function 1 (ASF1) binds

histone H3.3-H4 dimers and transfers them to HIRA[50]. To investigate if deposition of

histone H3.3 on telomeres is important for regulating telomere deprotection-induced

cGAS-STING activation, we used siRNAs to deplete individual histone H3.3

chaperone proteins, including ASF1, HIRA, ATRX, and DAXX, from BJ^t cells

expressing TRF2ΔBΔM. Consistently, depletion of ATRX, DAXX, or HIRA reduced the

telomeric distribution of histone H3.3, as evidence by decreased telomeric signals in

histone H3.3 chromatin immunoprecipitation (ChIP) assays following siRNA-mediated

depletion of ATRX, DAXX, or HIRA siRNAs (Fig. S2). STING phosphorylation and

IFNβ expression were reduced upon TRF2ΔBΔM expression in the absence of ATRX

or DAXX (Fig. 6A, 6C). Additionally, depletion of HIRA or ASF1 significantly reduced

STING phosphorylation and IFNβ expression TRF2ΔBΔM-expressing cells (Fig. 6B,

6C). Together, these observations indicated that ATRX, DAXX, HIRA, and ASF1 are

all required to activate the cGAS-STING response caused by telomere deprotection.

To determine how these histone chaperones regulate downstream responses to

telomere deprotection, we assessed telomeric DDR and telomere fusion in cells

depleted of ATRX, DAXX, or HIRA. Depletion of ATRX, DAXX, or HIRA reduced the

percentages of fused chromatids and TIF formation in TRF2ΔBΔM-expressing BJ^t

cells (Fig. 6D, S3), indicating that proper telomeric distribution of histone H3.3 is

essential for DDR activation and telomere fusion following telomere deprotection.

Thus, histone H3.3 localization at telomeres is essential for DDR activation, telomere

fusion, and cGAS-STING pathway activation.

DISCUSSION

Our results reveal a role for histone H3.3 in mediating telomere deprotection-induced activation of the cGAS-STING pathway. Although histone H3.3 deposition at telomeres has been previously documented, its role in telomere dynamics and the DNA damage response remains poorly understood. Notably, mutations in *H3F3A*, the gene encoding histone H3.3, are strongly associated with glioblastoma, a tumor that predominantly relies on the ALT mechanism[32,51]. Our previous work demonstrated that ALT cancer cells lack a functional cGAS-STING-mediated intracellular DNA sensing mechanism and that histone H3.3 and its chaperone contribute to regulating this pathway[20]. Here, we show that histone H3.3 contributes to promote telomeric DDR and downstream response, including chromosomal end fusion and activation of cGAS-STING pathway (Fig. 3, 5). These findings suggest that histone H3.3 contributes to ALT cancer-associated telomere dysfunctions.

Recent studies have shown that glioblastoma-associated histone H3.3 variants, such as histone H3.3[K27M] and histone H3.3[G34V], impair chromosome segregation, trigger genetic instability[29], and regulate ALT-associated features[52]. These findings suggest that ALT-associated histone H3.3 mutations may suppress the DDR, promote genetic instability, and facilitate error-prone genome doubling and rapid cell cycle progression. These mutations may also suppress cGAS-STING activation, dampening the innate immune response and allowing cancer cells to thrive under genomic stress. However, further studies are needed to clarify the relationship between histone H3.3 mutations and cGAS-STING pathway activity in ALT cancer cells to better understand how these mutations contribute to ALT phenotypes.

ALT cancers maintain telomeres through a telomerase-independent recombination-

driven recombination. Given the link between histone H3.3 and ALT cancers[29-32], we

propose that its expression and telomeric deposition influence DNA repair pathway

choice, shifting the balance between NHEJ and homologous recombination (HR).

Histone H3.3 has been shown to function downstream of the PARP1-CHD2 machinery

to promote NHEJ at telomeres[41]. Similarly, the histone H3.3's chaperone molecule,

ATRX is active in both NHEJ[37] and HR[53]. In our observations, depletion of histone

H3.3 or its chaperones (ATRX, DAXX, and HIRA) reduces telomere fusion events (Fig.

3, 6) and impairs TIF formation (Fig. 5, S3), underscoring their importance in telomere

repair. Since ALT phenotypes depend on HR, we propose that mutations in histone

H3.3 or its chaperones disrupt NHEJ while enhancing HR, thereby supporting ALT

telomere maintenance. It will be important to establish in the future how histone H3.3

localization and chaperone interactions mediate this switch.

Telomere fusion has been proposed as a mechanism to counteract telomere

shortening[54,55]. However, our findings reveal that telomere fusion also activates the

cGAS-STING pathway, which may impair cell growth. This scenario is evidenced by

the impairment of cGAS-STING activation we detected upon depletion of the DNA

repair proteins LIG4 and XRCC4 (Fig 4). Furthermore, cGAS depletion rescues the

growth inhibition induced by telomere deprotection (Fig. 1E, F), suggesting that

telomere fusion promotes cellular senescence via cGAS-STING pathway activation.

Interestingly, our findings indicate that cGAS-STING activation occurs in a pulsatile

manner following telomere deprotection (Fig. S1). Thus, telomere fusion appears to

trigger rapid but transient activation of this pathway under telomeric stress. Notably,

transient activation of the cGAS-STING pathway by TRF2ΔBΔM expression is

sufficient to inhibit cell growth (Fig. 1E, F). These findings reveal a novel role for

telomere fusion in transient cGAS-STING pathway activation and cell growth

regulation. Future studies should explore the mechanisms underlying this temporal

control and its broader implications. Overall, our results highlight the essential role of

histone H3.3 in deprotected telomere-driven cGAS-STING signaling and provide new

insights into its underlying mechanisms.

RESOURCE AVAILABILITY

Lead contact

Requests for further information and resources should be directed to and will be fulfilled by the lead contact, Liuh-Yow Chen (lyowchen@gate.sinica.edu.tw).

Materials availability

This study did not generate new unique reagents.

Data and code availability

Original Western blot images and microscopy data reported in this paper will be shared by the lead contact. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

ACKNOWLEDGEMENT

We thank the Imaging Core of the Institute of Molecular Biology (IMB), Academia Sinica, for technical support. We also thank Nithila Joseph in our lab and John O'Brien for assistance with English editing. This project is supported by grants from Academia Sinica (AS-GCP-113-L02) and the Ministry of Science and Technology (111-2311-B-001-013).

AUTHOR CONTRIBUTIONS

Conceptualization, C.-M. H. and L.-Y. C.; methodology, C.-M. H.; analysis, C.-M. H.; investigation, C.-M. H.; resources, L.-Y. C.; writing – original draft, C.-M. H. and L.-Y. C.; writing – review and editing, C.-M. H. and L.-Y. C.; funding acquisition, L.-Y. C.;

361   supervision, C.-M. H. and L.-Y. C.

363   **DECLARATION OF INTERESTS**

364   The authors declare no competing interests.

FIGURES

Figure 1. Expression of TRF2ΔBΔM triggers cGAS-STING pathway activation

and cGAS-mediated growth inhibition.

BJ^t cells were separated into untreated and treatment groups, and then 25 ng/ml

doxycycline (DOX) was mixed into the culture medium of the treatment group to

induce expression of TRF2ΔBΔM. (**A**) Ectopic expression of myc-tagged TRF2ΔBΔM

and GAPDH and was determined by Western blotting. (**B**) Immunostaining for

γH2AX and TRF1 was performed on day 4 after DOX treatment. Representative

images stained for γH2AX (green) and TRF1 (red). White arrowheads indicate

representative colocalized signals of TRF1 and γH2AX. For silencing the expression

of cGAS, siRNAs were added twice, i.e., at 60 h and 12 h before inducing

TRF2ΔBΔM. (**C**) Protein levels of STING phosphorylated at residue S366 (pSTING),

STING, cGAS, myc, and actin were determined by Western blotting. (**D**) The

expression level of IFNβ mRNA was determined by RT-qPCR. qPCR experiments

represent the mean ±SD of three technical repeats. (**E**) Numbers of cells surviving

under the indicated conditions were determined by counting the DAPI signals in the

culture dishes. Cell numbers were normalized to day 1 after treatment for each

experiment. Data represent the mean ±SD of three biological repeats. (**F**) BJ[t] cells

inducibly expressing GFP or TRF2ΔBΔM and respective controls were treated with

or without 25 ng/ml DOX. Surviving cells were counted 4 days later. Statistical

analysis was conducted using one-way ANOVA. n.s. represents no significant

differences, * p-value < 0.05, *** p-value <0.001, and **** p-value < 0.0001.

Figure 2. Depletion of H3.3 impairs cGAS-STING pathway activation and rescues growth defects due to the expression of TRF2ΔBΔM.

(**A**) BJ^t cells stably expressing GFP or myc-tagged TRF2ΔBΔM were established. siRNAs were then added at two different times, i.e., at 60 h and 12 h before inducing TRF2ΔBΔM expression through doxycycline (DOX). Samples were cultured for 2.5 days to investigate the cGAS-STING signaling response. Protein levels of phosphorylated STING (pSTING), STING, H3.3, myc, and GAPDH were determined by Western blotting. (**B**) mRNA expression levels of IFNβ were determined by RT-qPCR. qPCR experiments represent the mean ±SD of three technical repeats. (**C**) Numbers of cells in the DOX+ or DOX- treatment groups were counted to evaluate cell growth 4 days later. Data represent the mean ±SD of three biological repeats. (**D**)

403   A siRNA-insensitive H3.3 construct was introduced into cells via viral vectors and

404   ectopically expressed. Protein levels of phosphorylated STING (pSTING), STING,

405   H3.3, myc, and GAPDH were then determined by Western blotting. Statistical analysis

406   was conducted using one-way ANOVA. n.s. represents no significant differences, * p-

407   value < 0.05, *** p-value <0.001, and **** p-value < 0.0001.

Figure 3. Depletion of H3.3 expression limits telomere-to-telomere fusion events and the formation of nuclear abnormalities.

BJ[t] cells were treated with siRNAs including siCtrl, siH3.3, sicGAS, siLIG4, and siXRCC4 at 60 h and 12 h before DOX treatment to induce expression of TRF2ΔBΔM for 24 h. Cells were arrested using 0.1 μg/ml Demecolcine for another 24 h, and then spread on a slide to visualize telomere-to-telomere fusion events by staining with TelC probes. (**A**) Representative images of telomere fusion events (white arrows). (**B**) Quantitative results for telomere fusion events were calculated from >2000 chromosomes for each sample. Data represent mean ±SD of three biological repeats. (**C**) Representative cells images of siRNA-treated cells on day 2 after DOX treatment. For each group, >1,000 nuclear images were acquired and counted to determine percentages of the indicated nuclear abnormalities. Representative images are shown for a nuclear bud (left), a nucleoplasmic bridge (middle), and a micronucleus (right). (**D**) Quantitative results for nuclear abnormalities. Data represent the mean ±SD of three technical repeats. Statistical analysis was conducted using one-way ANOVA. n.s. represents no significant differences, ** p-value < 0.01, *** p-value <0.001, and **** p-value < 0.0001.

Figure 4. Depletion of LIG4 and XRCC4 disrupts activation of the cGAS-STING

response.

BJ^t cells inducibly expressing myc-tagged TRF2ΔBΔM were treated with siRNAs at 60

h and 12 h before doxycycline (DOX) treatment. To correlate telomere fusion events

and nuclear abnormalities, treated cells were lysed on day 2 after DOX treatment. (**A**)

Protein levels of phosphorylated STING (pSTING), STING, H3.3, cGAS, LIG4, XRCC4,

myc, and GAPDH were determined by Western blotting. (**B**) mRNA expression levels

of IFNβ were determined by RT-qPCR. qPCR experiments represent the mean ±SD

of three technical repeats. Statistical analysis was conducted using one-way ANOVA.

**** p-value < 0.0001.

Figure 5. Depletion of H3.3 expression limits TIF formation.

BJ^t cells were treated with siRNAs including siCtrl, siH3.3, and siNBS1 at 60 h and 12 h before inducing stable expression of TRF2ΔBΔM or TPP1ΔRD through 2.5 ng/ml DOX treatment. (**A**) Immunostaining for γH2AX and TRF1 was performed on day 2 after DOX treatment. Representative images stained for γH2AX (green) and TRF1 (red). White arrowheads indicate representative colocalized signals of TRF1 and γH2AX, which are defined as TIFs, in the cells. (**B**) Numbers of TIFs per cell for the different treatment groups. Each dot represents the TIF count per cell. Red lines

indicate the median value for each group. The data represent three biological repeats,

with 70-120 cells counted in each repeat, and each experimental group represents

250-330 cells. (**C**) Percentages of cells harboring >5 TIFs. More than 100 cells were

quantified for each sample in each repeat. BJ^t cells inducibly expressing either

TRF2ΔBΔM were treated with 0, 5, or 25 μM of MRN complex inhibitor, Mirin. Total

RNA and protein samples are collected after doxycycline treatment for 2.5 days. (A)

Protein levels of phosphorylated STING (pSTING), STING, myc, and GAPDH were

determined by Western blotting. (B) Relative expression levels of IFNβ mRNA in each

sample group were determined by qPCR. qPCR experiments represent the mean ±SD

of three technical repeats. Data represent mean ±SD of three biological repeats.

Statistical analysis was conducted using one-way ANOVA. ** p-value < 0.01, *** p-

value <0.001, and **** p-value < 0.0001.

Figure 6. Depletion of H3.3 chaperones impairs cGAS-STING pathway activation.

BJ[t] cells were treated with siRNAs at 60 h and 12 h before inducing expression of myc-tagged TRF2ΔBΔM through DOX treatment. Samples were cultured for 2.5 days to investigate the cGAS-STING signaling response. (**A**, **B**) Protein levels of phosphorylated STING (pSTING), STING, ASF1, HIRA, ATRX, DAXX, H3.3, myc, and GAPDH were determined by Western blotting. (**C**) mRNA expression levels of IFNβ were determined by RT-qPCR. qPCR experiments represent the mean ±SD of three technical repeats. Another set of BJ[t] cells was treated with siRNAs including siCtrl, siH3.3, siATRX, siDAXX, siHIRA, and siLIG4 at 60 h and 12 h before DOX treatment to induce expression of TRF2ΔBΔM for 24 h. Cells were arrested using 0.1 μg/ml Demecolcine for another 24 h, and then spread on a slide to visualize telomere fusion events by staining with a TelC probe. Statistical analysis was conducted using one-way ANOVA. **** p-value < 0.0001. (**D**) Quantitative results for telomere fusion events

477   were calculated from >2000 chromosomes for each sample. Data represent the mean

478   ±SD of two sets of experiments.

REFERENCES

1. Lundblad, V., and Szostak, J.W. (1989). A mutant with a defect in telomere elongation leads to senescence in yeast. Cell *57*, 633-643. 10.1016/0092-8674(89)90132-3.
2. van Steensel, B., Smogorzewska, A., and de Lange, T. (1998). TRF2 protects human telomeres from end-to-end fusions. Cell *92*, 401-413. 10.1016/s0092-8674(00)80932-0.
3. Karlseder, J., Smogorzewska, A., and de Lange, T. (2002). Senescence induced by altered telomere state, not telomere loss. Science *295*, 2446-2449. 10.1126/science.1069523.
4. Karlseder, J., Broccoli, D., Dai, Y., Hardy, S., and de Lange, T. (1999). p53- and ATM-dependent apoptosis induced by telomeres lacking TRF2. Science *283*, 1321-1325. 10.1126/science.283.5406.1321.
5. Xu, Y., and Baltimore, D. (1996). Dual roles of ATM in the cellular response to radiation and in cell growth control. Genes Dev *10*, 2401-2410. 10.1101/gad.10.19.2401.
6. Carney, J.P., Maser, R.S., Olivares, H., Davis, E.M., Le Beau, M., Yates, J.R., 3rd, Hays, L., Morgan, W.F., and Petrini, J.H. (1998). The hMre11/hRad50 protein complex and Nijmegen breakage syndrome: linkage of double-strand break repair to the cellular DNA damage response. Cell *93*, 477-486. 10.1016/s0092-8674(00)81175-7.
7. Usui, T., Ogawa, H., and Petrini, J.H. (2001). A DNA damage response pathway controlled by Tel1 and the Mre11 complex. Mol Cell *7*, 1255-1266. 10.1016/s1097-2765(01)00270-2.
8. Celli, G.B., Denchi, E.L., and de Lange, T. (2006). Ku70 stimulates fusion of dysfunctional telomeres yet protects chromosome ends from homologous recombination. Nat Cell Biol *8*, 885-890. 10.1038/ncb1444.
9. Smogorzewska, A., Karlseder, J., Holtgreve-Grez, H., Jauch, A., and de Lange, T. (2002). DNA ligase IV-dependent NHEJ of deprotected mammalian telomeres in G1 and G2. Curr Biol *12*, 1635-1644. 10.1016/s0960-9822(02)01179-x.
10. Abdisalaam, S., Bhattacharya, S., Mukherjee, S., Sinha, D., Srinivasan, K., Zhu, M., Akbay, E.A., Sadek, H.A., Shay, J.W., and Asaithamby, A. (2020). Dysfunctional telomeres trigger cellular senescence mediated by cyclic GMP-AMP synthase. J Biol Chem *295*, 11144-11160. 10.1074/jbc.RA120.012962.
11. Hopfner, K.P., and Hornung, V. (2020). Molecular mechanisms and cellular functions of cGAS-STING signalling. Nat Rev Mol Cell Biol *21*, 501-521. 10.1038/s41580-020-0244-x.
12. Stetson, D.B., and Medzhitov, R. (2006). Recognition of cytosolic DNA activates an IRF3-dependent innate immune response. Immunity *24*, 93-103. 10.1016/j.immuni.2005.12.003.
13. Tanaka, Y., and Chen, Z.J. (2012). STING specifies IRF3 phosphorylation by TBK1 in the cytosolic DNA signaling pathway. Sci Signal *5*, ra20. 10.1126/scisignal.2002521.
14. Ablasser, A., Goldeck, M., Cavlar, T., Deimling, T., Witte, G., Rohl, I., Hopfner, K.P., Ludwig, J., and Hornung, V. (2013). cGAS produces a 2'-5'-linked cyclic dinucleotide second messenger that activates STING. Nature *498*, 380-384. 10.1038/nature12306.

15. Diner, E.J., Burdette, D.L., Wilson, S.C., Monroe, K.M., Kellenberger, C.A., Hyodo, M.,
Hayakawa, Y., Hammond, M.C., and Vance, R.E. (2013). The innate immune DNA
sensor cGAS produces a noncanonical cyclic dinucleotide that activates human
STING. Cell Rep *3*, 1355-1361. 10.1016/j.celrep.2013.05.009.
16. Mankan, A.K., Schmidt, T., Chauhan, D., Goldeck, M., Honing, K., Gaidt, M.,
Kubarenko, A.V., Andreeva, L., Hopfner, K.P., and Hornung, V. (2014). Cytosolic
RNA:DNA hybrids activate the cGAS-STING axis. EMBO J *33*, 2937-2946.
10.15252/embj.201488726.
17. Lam, E., Stein, S., and Falck-Pedersen, E. (2014). Adenovirus detection by the
cGAS/STING/TBK1 DNA sensing cascade. J Virol *88*, 974-981. 10.1128/JVI.02702-13.
18. Sun, L., Wu, J., Du, F., Chen, X., and Chen, Z.J. (2013). Cyclic GMP-AMP synthase is a
cytosolic DNA sensor that activates the type I interferon pathway. Science *339*, 786-
791. 10.1126/science.1232458.
19. Wu, J., Sun, L., Chen, X., Du, F., Shi, H., Chen, C., and Chen, Z.J. (2013). Cyclic GMP-
AMP is an endogenous second messenger in innate immune signaling by cytosolic
DNA. Science *339*, 826-830. 10.1126/science.1229963.
20. Chen, Y.A., Shen, Y.L., Hsia, H.Y., Tiang, Y.P., Sung, T.L., and Chen, L.Y. (2017).
Extrachromosomal telomere repeat DNA is linked to ALT development via cGAS-
STING DNA sensing pathway. Nat Struct Mol Biol *24*, 1124-1131.
10.1038/nsmb.3498.
21. Wong, L.H., McGhie, J.D., Sim, M., Anderson, M.A., Ahn, S., Hannan, R.D., George,
A.J., Morgan, K.A., Mann, J.R., and Choo, K.H. (2010). ATRX interacts with H3.3 in
maintaining telomere structural integrity in pluripotent embryonic stem cells.
Genome Res *20*, 351-360. 10.1101/gr.101477.109.
22. Goldberg, A.D., Banaszynski, L.A., Noh, K.M., Lewis, P.W., Elsaesser, S.J., Stadler, S.,
Dewell, S., Law, M., Guo, X., Li, X., et al. (2010). Distinct factors control histone
variant H3.3 localization at specific genomic regions. Cell *140*, 678-691.
10.1016/j.cell.2010.01.003.
23. Drane, P., Ouararhni, K., Depaux, A., Shuaib, M., and Hamiche, A. (2010). The death-
associated protein DAXX is a novel histone chaperone involved in the replication-
independent deposition of H3.3. Genes Dev *24*, 1253-1265. 10.1101/gad.566910.
24. Armache, A., Yang, S., Martinez de Paz, A., Robbins, L.E., Durmaz, C., Cheong, J.Q.,
Ravishankar, A., Daman, A.W., Ahimovic, D.J., Klevorn, T., et al. (2020). Histone H3.3
phosphorylation amplifies stimulation-induced transcription. Nature *583*, 852-857.
10.1038/s41586-020-2533-0.
25. Ishiuchi, T., Abe, S., Inoue, K., Yeung, W.K.A., Miki, Y., Ogura, A., and Sasaki, H.
(2021). Reprogramming of the histone H3.3 landscape in the early mouse embryo.
Nat Struct Mol Biol *28*, 38-49. 10.1038/s41594-020-00521-1.
26. Huang, X.W., Cheng, X.R., Wang, N., Zhang, Y.W., Liao, C., Jin, L.H., and Lei, L. (2018).
Histone variant H3.3 and its functions in reprogramming. Yi Chuan *40*, 186-196.
10.16288/j.yczz.17-233.
27. Loppin, B., and Berger, F. (2020). Histone Variants: The Nexus of Developmental
Decisions and Epigenetic Memory. Annu Rev Genet *54*, 121-149. 10.1146/annurev-
genet-022620-100039.
28. Alvarez-Astudillo, F., Garrido, D., Varas-Godoy, M., Gutierrez, J.L., Villanueva, R.A.,
and Loyola, A. (2021). The histone variant H3.3 regulates the transcription of the
hepatitis B virus. Ann Hepatol *21*, 100261. 10.1016/j.aohep.2020.09.005.

29.   Bockaj, I., Martini, T.E.I., de Camargo Magalhaes, E.S., Bakker, P.L., Meeuwsen-de
Boer, T.G.J., Armandari, I., Meuleman, S.L., Mondria, M.T., Stok, C., Kok, Y.P., et al.
(2021). The H3.3K27M oncohistone affects replication stress outcome and provokes
genomic instability in pediatric glioma. PLoS Genet *17*, e1009868.
10.1371/journal.pgen.1009868.
30.   Voon, H.P.J., Hii, L., Garvie, A., Udugama, M., Krug, B., Russo, C., Chueh, A.C., Daly,
R.J., Morey, A., Bell, T.D.M., et al. (2023). Pediatric glioma histone H3.3 K27M/G34R
mutations drive abnormalities in PML nuclear bodies. Genome Biol *24*, 284.
10.1186/s13059-023-03122-5.
31.   Haase, S., Banerjee, K., Mujeeb, A.A., Hartlage, C.S., Nunez, F.M., Nunez, F.J.,
Alghamri, M.S., Kadiyala, P., Carney, S., Barissi, M.N., et al. (2022). H3.3-G34
mutations impair DNA repair and promote cGAS/STING-mediated immune responses
in pediatric high-grade glioma models. J Clin Invest *132*. 10.1172/JCI154229.
32.   Schwartzentruber, J., Korshunov, A., Liu, X.Y., Jones, D.T., Pfaff, E., Jacob, K., Sturm,
D., Fontebasso, A.M., Quang, D.A., Tonjes, M., et al. (2012). Driver mutations in
histone H3.3 and chromatin remodelling genes in paediatric glioblastoma. Nature
482, 226-231. 10.1038/nature10833.
33.   Brosnan-Cashman, J.A., Yuan, M., Graham, M.K., Rizzo, A.J., Myers, K.M., Davis, C.,
Zhang, R., Esopi, D.M., Raabe, E.H., Eberhart, C.G., et al. (2018). ATRX loss induces
multiple hallmarks of the alternative lengthening of telomeres (ALT) phenotype in
human glioma cell lines in a cell line-specific manner. PLoS One *13*, e0204159.
10.1371/journal.pone.0204159.
34.   Heaphy, C.M., de Wilde, R.F., Jiao, Y., Klein, A.P., Edil, B.H., Shi, C., Bettegowda, C.,
Rodriguez, F.J., Eberhart, C.G., Hebbar, S., et al. (2011). Altered telomeres in tumors
with ATRX and DAXX mutations. Science *333*, 425. 10.1126/science.1207313.
35.   Ramamoorthy, M., and Smith, S. (2015). Loss of ATRX Suppresses Resolution of
Telomere Cohesion to Control Recombination in ALT Cancer Cells. Cancer Cell *28*,
357-369. 10.1016/j.ccell.2015.08.003.
36.   Jiao, Y., Killela, P.J., Reitman, Z.J., Rasheed, A.B., Heaphy, C.M., de Wilde, R.F.,
Rodriguez, F.J., Rosemberg, S., Oba-Shinjo, S.M., Nagahashi Marie, S.K., et al. (2012).
Frequent ATRX, CIC, FUBP1 and IDH1 mutations refine the classification of malignant
gliomas. Oncotarget *3*, 709-722. 10.18632/oncotarget.588.
37.   Koschmann, C., Calinescu, A.A., Nunez, F.J., Mackay, A., Fazal-Salom, J., Thomas, D.,
Mendez, F., Kamran, N., Dzaman, M., Mulpuri, L., et al. (2016). ATRX loss promotes
tumor growth and impairs nonhomologous end joining DNA repair in glioma. Sci
Transl Med *8*, 328ra328. 10.1126/scitranslmed.aac8228.
38.   Dyer, M.A., Qadeer, Z.A., Valle-Garcia, D., and Bernstein, E. (2017). ATRX and DAXX:
Mechanisms and Mutations. Cold Spring Harb Perspect Med *7*.
10.1101/cshperspect.a026567.
39.   Haase, S., Garcia-Fabiani, M.B., Carney, S., Altshuler, D., Nunez, F.J., Mendez, F.M.,
Nunez, F., Lowenstein, P.R., and Castro, M.G. (2018). Mutant ATRX: uncovering a
new therapeutic target for glioma. Expert Opin Ther Targets *22*, 599-613.
10.1080/14728222.2018.1487953.
40.   Shi, L., Wen, H., and Shi, X. (2017). The Histone Variant H3.3 in Transcriptional
Regulation and Human Disease. J Mol Biol *429*, 1934-1945.
10.1016/j.jmb.2016.11.019.

41. Luijsterburg, M.S., de Krijger, I., Wiegant, W.W., Shah, R.G., Smeenk, G., de Groot,
A.J.L., Pines, A., Vertegaal, A.C.O., Jacobs, J.J.L., Shah, G.M., and van Attikum, H.
(2016). PARP1 Links CHD2-Mediated Chromatin Expansion and H3.3 Deposition to
DNA Repair by Non-homologous End-Joining. Mol Cell *61*, 547-562.
10.1016/j.molcel.2016.01.019.
42. Cesare, A.J., and Griffith, J.D. (2004). Telomeric DNA in ALT cells is characterized by
free telomeric circles and heterogeneous t-loops. Mol Cell Biol *24*, 9948-9957.
10.1128/MCB.24.22.9948-9957.2004.
43. Nabetani, A., and Ishikawa, F. (2009). Unusual telomeric DNAs in human telomerase-
negative immortalized cells. Mol Cell Biol *29*, 703-713. 10.1128/MCB.00603-08.
44. Cesare, A.J., and Reddel, R.R. (2008). Telomere uncapping and alternative
lengthening of telomeres. Mech Ageing Dev *129*, 99-108.
10.1016/j.mad.2007.11.006.
45. Celli, G.B., and de Lange, T. (2005). DNA processing is not required for ATM-
mediated telomere damage response after TRF2 deletion. Nat Cell Biol *7*, 712-718.
10.1038/ncb1275.
46. Pampalona, J., Soler, D., Genesca, A., and Tusell, L. (2010). Telomere dysfunction and
chromosome structure modulate the contribution of individual chromosomes in
abnormal nuclear morphologies. Mutat Res *683*, 16-22.
10.1016/j.mrfmmm.2009.10.001.
47. Takaki, T., Millar, R., Hiley, C.T., and Boulton, S.J. (2024). Micronuclei induced by
radiation, replication stress, or chromosome segregation errors do not activate
cGAS-STING. Mol Cell *84*, 2203-2213 e2205. 10.1016/j.molcel.2024.04.017.
48. Dimitrova, N., and de Lange, T. (2009). Cell cycle-dependent role of MRN at
dysfunctional telomeres: ATM signaling-dependent induction of nonhomologous end
joining (NHEJ) in G1 and resection-mediated inhibition of NHEJ in G2. Mol Cell Biol
29, 5552-5563. 10.1128/MCB.00476-09.
49. Malaquin, N., Olivier, M.A., Martinez, A., Nadeau, S., Sawchyn, C., Coppe, J.P.,
Cardin, G., Mallette, F.A., Campisi, J., and Rodier, F. (2020). Non-canonical ATM/MRN
activities temporally define the senescence secretory program. EMBO Rep *21*,
e50718. 10.15252/embr.202050718.
50. Grover, P., Asa, J.S., and Campos, E.I. (2018). H3-H4 Histone Chaperone Pathways.
Annu Rev Genet *52*, 109-130. 10.1146/annurev-genet-120417-031547.
51. Sturm, D., Witt, H., Hovestadt, V., Khuong-Quang, D.A., Jones, D.T., Konermann, C.,
Pfaff, E., Tonjes, M., Sill, M., Bender, S., et al. (2012). Hotspot mutations in H3F3A
and IDH1 define distinct epigenetic and biological subgroups of glioblastoma. Cancer
Cell *22*, 425-437. 10.1016/j.ccr.2012.08.024.
52. Udugama, M., Hii, L., Garvie, A., Cervini, M., Vinod, B., Chan, F.L., Das, P.P., Mann,
J.R., Collas, P., Voon, H.P.J., and Wong, L.H. (2021). Mutations inhibiting KDM4B
drive ALT activation in ATRX-mutated glioblastomas. Nat Commun *12*, 2584.
10.1038/s41467-021-22543-z.
53. Elbakry, A., Juhasz, S., Chan, K.C., and Lobrich, M. (2021). ATRX and RECQ5 define
distinct homologous recombination subpathways. Proc Natl Acad Sci U S A *118*.
10.1073/pnas.2010370118.
54. Cleal, K., Norris, K., and Baird, D. (2018). Telomere Length Dynamics and the
Evolution of Cancer Genome Architecture. Int J Mol Sci *19*. 10.3390/ijms19020482.

55.   Capper, R., Britt-Compton, B., Tankimanova, M., Rowson, J., Letsolo, B., Man, S.,
666   Haughton, M., and Baird, D.M. (2007). The nature of telomere fusion and a definition
667   of the critical telomere length in human cells. Genes Dev *21*, 2495-2508.
668   10.1101/gad.439107.
669

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Phospho-STING (Ser366) (D7C3S)	Cell Signaling Technology	19781 RRID: AB_2737062
STING	R&D Systems	MAB7169 RRID: AB_1091940
cGAS	Cell Signaling Technology	15102 RRID: AB_2732795
H3.3	Milipore	09-838 RRID: AB_10845793
H3.3	Abcam	Ab176840 RRID: AB_2715502
myc	Sigma-Aldrich	M4439 RRID: AB_439694
LIG4	Proteintech	12695-1-AP RRID: AB_2136253
XRCC4	Proteintech	15817-1-AP RRID: AB_2257478
ATRX	Bethyl	A301-045A RRID: AB_2243144
DAXX	Bethyl	A301-353A RRID: AB_937792
HIRA	Milipore	04-1488 RRID: AB_1977097
ASF1	Santa Cruz Biotechnology	sc-53171 RRID: AB_2059662
Phosphor-Histone H2AX (Ser139)	Millipore	05-636 RRID: AB_309864
DAPI	Thermo Fisher Scientific	D1306
TRF1	GeneTex, customized	Customized, 90549
TRF2	GeneTex, customized	Customized, 90550
HRP-conjugated GAPDH	ProteinTech	HRP-60004 RRID: AB_2737588
HRP-conjugated beta-actin	ProteinTech	HRP-60008 RRID: AB_2819183
Mouse IgG Alexa Fluor 488	Thermo Fisher Scientific	A-11001 RRID: AB_2534069
Rabbit IgG Alexa Fluor 555	Thermo Fisher Scientific	A32727 RRID: AB_2633276
Chemicals, peptides, and recombinant proteins		
Mirin	Sigma-Aldrich	M9948
Doxycycline	TaKaRa	631311
Critical commercial assays		
Lipofectamine RNAiMAX Transfection Reagent	Invitrogen	13778075
TransIT-LT1	Mirus	MR-MIR2300
Mini-Rprotean TGX precast gels, 15 well	Bio-Rad	456-1096
ECL Western Blotting Detection kit	Millipore	NEL122001EA
Supersignal West Femto	Thermo	PIE 34096
NucleoSpin RNA, Mini kit for RNA purification	Macherey-Nagel	740955.50
iScript cDNA synthesis general supply	Bio-Rad	170-8891
ChIP Kit	Abcam	Ab500
Blocking reagent	Roche	10057177103

Experimental models: Cell lines		
BJ [Human fibroblast]	ATCC	RRID: CVCL-3653 CRL-2522
Oligonucleotides		
On-target plus siRNA reagent, siRNA sequences, see Table S1	Dharmacon	N/A
qRTPCR primers	This paper	Table S2
HPLC purified, >90% TMR-OO-ccctaaaccctaaaccctaaa probe	Panagene, customized	Customized
Software and algorithms		
ImageJ	ImageJ	http://imagej.net/ij/
Prism 9	GraphPad	http://www.graphpad.com/
Imaging systems		
Axio Imager Z1 System	Zeiss	N/A
ImageXpress Micro XLS System	Molecular Devices	N/A
DeltaVision Deconvolution Microscope	Image Solutions	N/A

Materials and Methods

Cell lines

Cell lines were obtained from ATCC and maintained under their recommended culturing conditions. BJ^t cells originated from BJ normal fibroblasts that stably express hTERT. The retroviral transformation was performed to generate cells that inducibly express green fluorescence protein (GFP), the TRF2 mutant TRF2ΔBΔM, or the TPP1 mutant TPP1ΔRD. Cells were tested for target protein expression by treating with a minimum of 25 ng/ml of doxycycline.

Gene silencing

The siRNAs used were On-target plus siRNA reagents (Dharmacon). The siRNA reagents were generated using the SMARTselection algorithm, with chemical modifications to eliminate off-target effects. Four siRNAs for each of H3F3A, H3F3B, cGAS, LIG4, XRCC4, ATRX, DAXX, HIRA, ASF1A, and ASF1B, as well as two siRNAs for NBS1, were employed. siRNAs were transfected into the cells using Lipofectamine RNAiMAX (Invitrogen) according to the manufacturer's instructions. Depletion of H3.3 and ASF1 was performed by mixing equal amounts of siRNAs targeting H3.3 and ASF1. MRN chemical inhibitors, Mirin (M9948, Sigma), was dissolved in DMSO as 10mM stock, and treated to the cell culture at the indicated concentrations.

Protein extraction, Western blots, and antibodies

Cell extracts were harvested using 1 X SDS sample buffer (62.5 mM Tris, pH 6.8, 10% glycerol, 2% SDS, 0.01% bromophenol blue, and 10% β-mercaptoethanol). Proteins were denatured by

heating the extract at 95 °C for 10 minutes. Proteins were separated by SDS-PAGE, and
transferred to a nitrocellulose membrane. The membranes were blocked with either Tris-
buffered saline (TBS) at pH 8.0 with 5% milk and 0.1% Tween-20 to determine phosphorylated
proteins or using Phosphate-buffered saline (PBS) at pH 8.0 with 5% milk and 0.1% Tween-20
to determine other proteins. Primary antibodies including phosphorylated STING (19781, Cell
Signaling), STING (MAB7169, R&D Systems), cGAS (15102, Cell Signaling), H3.3 (09-838,
Merkmillpore; ab176840, Abcam), myc (M4439, Sigma), LIG4 (12695-1-AP, Proteintech),
XRCC4 (15817-1-AP, Proteintech), ATRX (A301-045A, Bethyl), DAXX (A301-353A, Bethyl), HIRA
(04-1488, Millipore), and ASF1 (sc-53171, Santa Cruz Biotechnology) were used at 1:1000
dilution. Horseradish peroxidase-conjugated secondary antibodies were used at 1:2000
dilution to determine the proteins in the samples. As an internal control for gene expression,
HRP-conjugated GAPDH antibody (HRP60004, ProteinTech) and HRP-labeled ACTB (HRP-60008,
ProteinTech) were used at 1:5000 dilution. Either enhanced chemiluminescence (ECL) or
SuperSignal West Femto Maximum Sensitivity reagents (Thermo Fisher Scientific) were used
for signal detection.
714
715 ## RNA extraction and quantitative real-time PCR
716
Cells were lysed and harvested to extract total RNA using an RNA Mini Kit (Macherey-Nagel).
Reverse transcription was performed with iScript Reverse Transcriptase Supermix (Bio-Rad) to
generate cDNA for quantitative analyses. Quantitative real-time PCR was performed using
SYBR Green Select reagent (LifeTechnologies) to determine expression levels of target genes.
Quantitative PCR analysis was performed using a QuantStudio 12K Flex Real-Time PCR System
(ThermoFisher Scientific). The primer sequences are as follows: β-actin, 5′-
AGCACTGTGTTGGCGTACAG-3′ and 5′-TCCCTGGAGAAGAGCTACGA-3′; interferon β, 5′-
AAACTCATGAGCAGTCTGCA-3′ and 5′-AGGAGATCTTCAGTTTCGGAGG-3′. Data analysis was
performed by calculating the comparative Ct (ΔΔCt) of target genes normalized to an internal
control (β-actin).
727
728 ## Metaphase spread and telomere FISH on chromosomes
729
Analysis of telomere-to-telomere fusion was performed in DOX-treated cells subjected to
siRNA-mediated gene expression depletion. Cells were treated with DOX for 30 hours, and
reseeded in 0.1 μg/ml Demecolcine-containing medium. After 18 h of incubation, the floating
and attached cells were collected for further analysis. Cells were incubated with 75 mM KCl for
15 mins at 37 °C and routinely mixed at 5-minute intervals. Following KCl incubation, the cells
were fixed with fixing buffer (3:1 methanol: acetic acid) at 4 °C for more than 24 h. Fixed cells
were dropped on a glass slide from a height of 20 cm, heated at 75 °C for 1 min, and then dried
at room temperature overnight. The slides were stained according to a standard fluorescence
in situ hybridization (FISH) FISH protocol. In brief, the slides were fixed with 10% formalin for
2 mins, and then dehydrated by incubating them in increasing percentages of ethanol solution.
The slides were dried and incubated with a probe mixture (70% formamide, 0.5% blocking
reagent (Roche), 250 nM TMR-conjugated (CCCTAA)3-PNA probe, 10 mM Tris pH 7.4) at 80 °C
for 3 mins. The slides were hybridized at room temperature in the dark for 3 h. Serial wash
steps were then performed, including twice with Wash I (70% formamide, 10 mM Tris-HCl pH
7.5), and twice with Wash II (100 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.08% Tween-20). The
slides were counterstained with DAPI, dehydrated with ethanol, and mounted with

Fluoromount Aqueous Mounting Medium (Sigma). Images were acquired with a Zeiss Axio Imager Z1 System. Determination and quantitative analysis of telomere fusion events were performed by counting for 700 to 1500 chromosome pairs.

Immunofluorescence staining

Cells grown on coverslips were treated with permeabilization solution (20 mM Hepes pH 8.0, 50 mM NaCl, 3 mM MgCl2, 0.5% Triton-X 100, 300 mM sucrose), and fixed with 2% paraformaldehyde. The cells were permeabilized with 0.5% Triton-X 100, and then blocked with 2% fetal bovine serum (FBS) in phosphate-buffered saline. Primary antibodies including anti-TRF1 (90549, GeneTex) and anti-γH2AX (05-636, Millipore) were used. The secondary antibodies were Mouse IgG Alexa Fluor 488 (A-11001, Thermo Fisher Scientific) and Rabbit IgG Alexa Fluor 555 (A32727, Thermo Fisher Scientific). The cells were stained with DAPI, dehydrated, and mounted with ProLong Gold Antifade Mountant (Life Technologies). Images were acquired under a DeltaVision Deconvolution Microscope. Colocalization events were determined based on the overlapping of signals from different channels, and quantification was performed by counting for > 100 nuclei on one coverslip.

Chromatin immunoprecipitation (ChIP)

The chromatin immunoprecipitation protocol was adapted from the Abcam ChIP kit (Ab500). After depletion of gene expression by siRNAs for 4 days, the BJ^t cells were fixed with 1% formaldehyde for 10 min. The cross-linking process was quenched with glycine, and the cells were collected from the culture dish. Chromatin DNA was extracted from cells, and sheared into 200-500 bp. Sheared chromatin DNAs were immunoprecipitated with antibodies, including H3.3 (09-838, Merkmillpore) and TRF2 (90550, GeneTex) antibodies. Antibody-precipitated chromatin DNAs were enriched with protein A beads. The enriched chromatin DNAs were reversed crosslinking with protease and extracted from the precipitated DNA pellet. Precipitated DNAs were then loaded on the Hybond-XL membrane. The telomeric DNA from each sample was determined with α-P32-dCTP-labelled telomere probe.

Cell growth and quantification of nuclear abnormalities

Cell growth was determined by counting cells in the multi-well culture plates. Cells were plated in 12-well plates and fixed with 2% paraformaldehyde on the 1st-7th days post-treatment. The cells were stained with DAPI and imaged using an ImageXpress Micro XLS System (Molecular Devices). For each well, images in 64 fixed regions were taken to quantify the number of cells inside the well. These images were used to quantify nuclear abnormalities, including nucleoplasmic bridges, nuclear buds, and micronuclei. Such abnormalities were identified based on a deformed nuclear structure, and the quantification was performed by counting for >2000 nuclei in one sample.

Quantification and statistical analysis

With the help of GraphPad Prism, a one-way analysis of variance (ANOVA) was performed to assess the statistical differences among group means. Following a significant ANOVA result ($P<0.05$), post-hoc comparisons were conducted using Tukey's Honest Significant Difference

793   (HSD) test to determine pairwise difference between group means and to control for Type I
794   error due to multiple comparisons.
795