Neuropeptide oxytocin facilitates its own brain-to-periphery uptake by regulating blood

flow dynamics and permeability

Preethi Rajamannar[1,2], Oren Raz[3] and Gil Levkowitz[1,2*]

Departments of [1]Molecular Cell Biology, [2]Molecular Neuroscience, and [3]Physics of Complex

Systems, Weizmann Institute of Science, PO Box 26, Rehovot 7610001, Israel

*Correspondence should be addressed to

G.L. gil.levkowitz@weizmann.ac.il

Running title: Oxytocin regulates pituitary blood flow and permeability

Key words: neurohormones, oxytocin, arginine-vasopressin, neurohypophysis,

neuroendocrine, zebrafish

13
14

Abstract

The hypothalamo-neurohypophyseal system is an important neuroendocrine brain-to-blood

conduit through which the neurohormones oxytocin and arginine-vasopressin are released from

the brain into the general circulation to affect peripheral physiological functions such as salt

balance, metabolism and reproduction. However, the mechanism which executes fast and

efficient neurohormone release to the periphery remains unsolved. We show, using live imaging

in zebrafish, that a hyperosmotic physiological challenge elicits a local increase in

neurohypophyseal blood flow velocities and a change in capillary diameter, which is dictated

by the geometry of the hypophyseal vascular microcircuit. Genetic ablation of oxytocin

neurons and inhibition of oxytocin receptor signaling attenuated changes in capillary blood

flow and diameter. Optogenetic stimulation of oxytocin neurons resulted in an oxytocin

receptor-dependent increase in blood flow velocities. Lastly, both osmotic challenge and

oxytocin neuronal activation elicited a local rise in neurohypophyseal capillary permeability in

an oxytocin signaling-dependent manner. Our study demonstrates that physiologically elicited

changes in neurohypophyseal blood flow and permeability are regulated by oxytocin. We

propose that oxytocin-dependent neuro-vascular coupling facilitates its efficient uptake into

the blood circulation, suggesting a self-perpetuating mechanism of peripheral hormone

transfer.

INTRODUCTION

How the brain exerts control over peripheral functions is a fundamental question that has occupied researchers for over a century. In the early 1900s, Berta and Ernest Scharrer noticed that neurons residing in the hypothalamus of a small freshwater fish have properties similar to hormone-secreting endocrine cells (*1, 2*). This discovery served as an inspiration for Scharrers' "neurosecretion concept", arguing that hypothalamic neurons release packets of neuro-hormones from the brain into the general circulation by directly projecting their axons onto local blood vessels of the posterior pituitary gland, also known as the neurohypophysis. This brain-to-periphery neurosecretion machinery termed the hypothalamo-neurohypophyseal system (HNS), is evolutionarily conserved among all vertebrate species and is a major neuroendocrine system responsible for the secretion of the neuro-hormones oxytocin (OXT) and arginine-vasopressin (AVP) from hypothalamic axonal termini onto fenestrated (i.e., permeable) capillary network of the neurohypophysis to affect peripheral organs. In the periphery, AVP regulates water homeostasis by increasing the water permeability of the kidney, whereas OXT regulates labor and milk let-down by causing the contraction of the uterine smooth muscles and the myoepithelial cells of breast ducts, respectively(*3, 4*). In teleost fish species, OXT and AVP are involved in osmoregulation and regulating reproductive success(*5, 6*).

Despite the physiological importance of OXT and AVP, there is a gap in our understanding of their regulated secretion into the blood. The prevailing dogma maintains that in response to physiological demands, hypothalamic OXT and/or AVP neurons are stimulated, released from the axonal termini into the neurohypophysis, taken up by blood capillaries into the systemic circulation, and transported to target tissues (*7*). This process requires tight regulation between the secretory neurons, the release of the hormones, and their transport into the blood. However, it has been noted that a simplified stimulus–secretion coupling mechanism, wherein the release of the contents of vesicles upon neuronal activation (e.g. neurotransmitter release), is directly associated with its effect on the target neuron does not apply in the context of neuroendocrine effect on peripheral targets (*8*). Thus, while the measured firing rhythms of OXT and/or AVP neurons range within milliseconds or even seconds, it takes several minutes to detect them in the peripheral blood (*9*). Moreover, unlike the dense synaptic connection between neurons or the tightly arranged neurovascular units observed in the brain, HNS nerve endings do not form compact, juxtaposed connections with pituitary blood vessels. They rather form numerous

axonal varicosities packed with large dense core vesicles that release their content into the

neurohypophysis, wherein the secreted neurohormones are required to transverse the cell-

abundant pituitary parenchyma, the endothelial basement membrane, and then enter the

endothelial lumen (*10*). ***It is, therefore, highly plausible that an additional driving force is***

required to facilitate efficient coupling between neurohormone release and its vascular

uptake, leading to downstream effects on peripheral target tissues.

In the brain, synaptic activity increases local blood flow to meet the metabolic needs of a

firing neuron, a process termed neurovascular coupling. It has been suggested that endocrine

cells of the pituitary gland and pancreas are coordinated with blood vessels to generate

hormone pulses (*11, 12*). Thus, local blood flow in neuroendocrine interfaces may facilitate

efficient hormone uptake rather than just a means to cope with the increased demands for

oxygen.

Here, we hypothesized that in the neurohypophysis, activity-dependent changes in local

blood flow modulate peripheral neurohormone uptake. Using live imaging of the zebrafish

HNS, we monitored hypophyseal vascular dynamics in unanesthetized animals. We

demonstrate that the physiological activation of oxytocin neurons is directly coupled to changes

in blood flow velocity and capillary permeability. We further demonstrate that these

physiologically elicited vascular changes are regulated by oxytocin, suggesting that oxytocin

facilitates its own brain-to-periphery effects.

RESULTS

Physiological osmotic challenge increases local neurohypophyseal blood flow

To study neuro-vascular coupling in the HNS, we have devised a live-imaging platform

using the zebrafish model. We have previously established that the development, cell biology,

and anatomy of the HNS neurovascular interface are conserved between zebrafish and

mammals (*13-15*). The core components of larval zebrafish HNS neurovascular interface

comprised of elaborate oxytocinergic axonal varicosities, the hypophyseal artery (HyA), which

conveys blood flow into a capillary loop (HyC) and a venous hypophyseal outlet (HyV) (**Fig.**

1A). This relatively simple and discernible anatomical architecture allowed us to reproducibly

monitor vascular dynamics in the same locale within multiple unanesthetized zebrafish larvae.

To monitor blood flow, we employed the *Tg(l-fabp:DBP-EGFP)* reporter (*16, 17*) (**Fig. 1B,C**),

in which a serum albumin-like molecule is labeled, allowing visualization and quantification

of flow velocities calculated from displacement time kymographs of red blood cells (RBC)

(**Fig. S1B**). Capillary diameter was monitored using an endothelial-specific transgenic

Tg(kdrl:EGFP) reporter (**Fig. S1C**).

To examine whether physiological activation of the HNS altered hypophyseal blood flow,

we employed a hyperosmotic challenge known to stimulate OXT and AVP neurons (*18, 19*)

(**Fig. S1A**). When zebrafish larvae were acutely transferred from freshwater to hyperosmotic

solution equivalent to 50% seawater (17.5 ppt), we observed significant activation of OXT

neurons, displaying increased *c-fos* expression in their soma (**Fig. 1D,E**). This hyperosmotic

challenge induced a robust increase in blood flow (**Fig. 1F,G**) and variability of RBCs

velocities (**Fig. 1H**). The single-sided amplitude spectra, as computed by a Fast Fourier

transformation (FFT), indicated that the osmotic challenge induced an increase in the amplitude

of velocity fluctuations below 2Hz (**Fig. 1I**).

To ensure that the effect of osmotic challenge on blood flow was specific to the hypophyseal

capillary, we measured blood flow in the anterior mesencephalic central artery (AMCtA),

which is a well-discernable vascular brain structure located at a similar depth and of similar

diameter as the hypophyseal capillary loop (**Fig. 1A**). We found that osmotic challenge did not

affect the measured flow parameters of AMCtA vessel (**Fig. 1J-M**).

These results show that a physiological challenge, which activates OXT neurons elicits a

specific local increase in hypophyseal blood flow velocity.

118 The relationship between flow velocity and diameter is determined by HNS vascular

119 geometry

According to Poiseuille's law for viscous fluids, it was expected that local vasoconstriction

of a single capillary would increase flow resistance, resulting in decreased blood flow velocity

(*20*). However, we observed that the osmotic challenge, which caused an increase in velocity

was accompanied by a decrease in hypophyseal capillary diameter (**Fig. 2A-C**). This effect was

specific as the diameter of the adjacent AMCtA capillary did not change following the osmotic

challenge (**Fig. 2D-F**). Since it was shown that the geometry of the microvasculature network

in the brain plays a crucial role in determining blood flow behaviour (*21, 22*), we resolved the

relationship between blood flow velocity and capillary diameter considering the structure of the zebrafish hypophyseal vascular microcircuit. This structure is comprised of a stereotypic arterial inlet, capillary loop, and venous outlet; each having its corresponding length (L), and radius (r) (**Fig. 2G**). We expressed the capillary velocity (V_2) in terms of the vascular components' radii and lengths, according to Poiseuille's law, considering a range of relevant inlet-outlet total pressure differences (Δp) (**Fig. 2G and S1E,** for details, see '*Methods*' section). Plotting the resolved V_2 values as a function of capillary radii revealed that flow velocity is inversely proportional to an increase in capillary radii that are larger than 3 microns (**Fig. 2H**). This inverse proportionality between capillary velocity and diameter, aligned with our measured values of the hypophyseal capillary radii (3-6 μm **Fig. 2B**). Thus, in accordance with our velocity and radii measurements, we found that for basal velocity to increase after osmotic challenge within a range of relevant estimated inlet-outlet pressure differences (Δp), there must be either a small decrease in radii with a small increase in Δp or a large decrease in radii with no change in Δp (**Fig. 2H'**).

These results show that an increase in blood flow velocity can be caused by vasoconstriction given the geometry of the hypophyseal vascular microcircuit.

Hypophyseal blood flow is regulated by oxytocin signaling

We tested whether the observed neuronal and vascular responses to an osmotic challenge are coupled. This was demonstrated using the *Tg(oxt:Gal4;UAS:NTR-mCherry)* genetic system for conditional cell ablation (**Fig. 3A**). Specific death of OXT cells expressing the nitroreductase (NTR) enzyme was induced by the administration of the drug Nifurpirinol (NFP), which is converted to a cytotoxic agent by NTR (*23-25*). Treatment of these transgenic animals for 18 hours with NFP led to a significant decrease in OXT-positive hypophyseal axonal projections (**Fig. 3B,C**). When subjected to an osmotic challenge, control Gal4/UAS-negative larvae displayed increased blood flow velocity while their sibling larvae with ablated OXT neurons did not (**Fig. 3D,E**). The mean capillary blood flow velocity, amplitudes, and frequency spectra, and variability in velocity did not show a significant difference after the osmotic challenge of OXT ablated larvae (**Fig. 3F-I**). Notably, whereas the osmotic challenge caused vasoconstriction in non-ablated controls, it led to vasodilation of hypophyseal capillaries in larvae with ablated OXT neurons, presumably as a response to the absence of

OXT neurovascular innervation and low basal OXT secretion (**Fig. 3J-M**). The effects of this genetic OXT-ergic ablation were confined to the hypophyseal vasculature as they were not observed in the control AMCtA vessel (**Fig. 3N-P**).

In view of our observed effects of OXT neuron ablation, we hypothesized that OXT itself might regulate local hypophyseal blood flow velocity upon its release. We found that endothelial cells isolated from zebrafish pituitaries express mRNAs encoding the two known functional OXT receptors, *oxtra,* and *oxtrb,* with markedly higher expression levels of *oxtrb* (**Fig. S2**). We therefore examined whether the administration of a selective oxytocin receptor antagonist, L-368,899, which is known to inhibit both zebrafish OXT receptors (*26, 27*), affects hypophyseal blood flow. We found that larvae treated with L-368, 899, were refractory to the osmotic challenge in terms of challenge-induced changes in mean flow velocity parameters (**Fig. 4A-F**) and capillary diameter (**Fig. 4G-J**). Administration of L-368, 899 did not influence blood flow velocity in the extra-hypophyseal AMCtA blood vessel ruling out a general systemic effect of the antagonist (**Fig. 4K-M**).

Next, we tested whether direct activation of OXT neurons, rather than through osmotic challenge, is sufficient to modulate hypophyseal blood flow and diameter. To stimulate OXT neurons, we employed the CoChR-tdTomato channelrhodopsin variant, an efficient optogenetic tool for zebrafish (*28*). Injection of *UAS:CoChR-tdTomato* construct into the OXT specific driver line, *Tg(oxt:Gal4)* (*29*), resulted in specific expression of the CoChR-tdTomato protein in OXT neurons (**Fig. 5A,B**). Exposure of *Tg(oxt:Gal4;UAS:CoChR-tdTomato)* larvae to blue light resulted in a transient but significant increase in blood flow velocity, which peaked at approximately 30-50 seconds following light exposure (**Fig. 5D, E**). The control CoChR-tdTomato-negative siblings did not respond to blue light (**Fig. 5C, E**). Light-activated CoChR-positive larvae exhibited a marked increase in $\Delta V/V_0$ as compared to their CoChR-negative siblings (**Fig. 5F**). The responses of hypophyseal vasculature to optogenetic stimulation of OXT neurons were dependent on oxytocin receptor signalling as the above effects were significantly attenuated by the administration of the OXT receptor antagonist, L-368, 899 (**Fig. 5G-I**). In accordance with the vasoconstrictive response of the hypophyseal capillary to osmotic challenge, light activation of CoChR-positive, but not CoChR-negative, OXT neurons caused a reduction in mean capillary diameter across time (**Fig. 5J-M**). As in the case of the increased flow velocity, CoChR-induced vasoconstrictive response was inhibited by L-368, 899 (**Fig. 5N-P**). Notably, hypophyseal vascular responses (i.e., flow and diameter) to

optogenetic activation of OXT neurons were transient as compared to the hyperosmotic
challenge. Although this may imply the existence of additional mediators of the effects of
osmotic challenge on hypophyseal vascular dynamics, it could be due to the artificial activation
of OXT neurons through optogenetics, as opposed to a genuine physiological stimulus, which
activates these neurons.

These results show that local hypophyseal vascular responses triggered by a hyperosmotic
physiological challenge are coupled to OXT neuronal activation and are mediated by OXT
receptor(s).

Hypophyseal vascular permeability is regulated by OXT signaling

We hypothesized that OXT-mediated neurovascular-coupling may affect the vascular
passage of neurohormones and other proteins by regulating diffusion rate through the
fenestrated hypophyseal capillaries (*16*). We visualized and quantified hypophyseal capillary
permeability using the *Tg(l-fabp:DBP-EGFP)* reporter described above (**Fig. 6A**). We
performed fluorescent recovery after photo-bleaching (FRAP) followed by line scan imaging
comparing the fluorescent recovery of bleached and unbleached regions of interest positioned
on either side of the hypophyseal capillary (**Fig.6A', B**). Paired comparisons of permeability
before and after the challenge indicated that the average time taken to recover half the initial
fluorescent intensity ($T_{1/2}$) following an osmotic challenge was significantly shorter than in the
basal condition, indicating a higher rate of diffusion through the hypophyseal capillary (**Fig.
6C**).

We then investigated the inter-dependent relationship between blood flow velocity and
permeability. We have previously shown that Tgf-β signaling regulates hypophyseal vascular
permeability (*13*). We now show that albeit the observed attenuation of basal hypophyseal
permeability by the selective TGF-β receptor inhibitor, SB431542 (**Fig. 6D, E**), blocking Tgf-
β signaling did not affect either the basal blood flow velocity or the response to osmotic
challenge (**Fig. 6F**). We next examined whether decreasing blood flow affects permeability.
We observed that treatment of zebrafish larvae with the cholinergic receptor inhibitor alpha-
bungarotoxin (α-BTX) for 150 seconds led to a nearly 50% reduction in blood flow velocity
(**Fig. 6G-H**). This reduction in blood flow had no effect on hypophyseal vessel permeability
(**Fig. 6I**).

These results indicate that hypophyseal vascular permeability and blood flow velocity are
mutually independent.

To determine the involvement of OXT signaling in hypophyseal vascular permeability, we
performed the FRAP analysis in the presence of the OXT receptor antagonist. Basal
permeability was measured continuously at different time points following the administration
of the OXT receptor antagonist. A significant increase in $T_{1/2}$ was observed up to an hour after
the administration of the antagonist, indicating a cumulative decrease in protein diffusion rate
through the hypophyseal capillary (**Fig. 6J**). Furthermore, the permeability-enhancing effect
of OC was also attenuated in antagonist-treated larvae as compared to their untreated siblings
(**Fig. 6K**).

The involvement of OXT in modulating permeability was further determined by direct
optogenetic activation of OXT neurons combined with FRAP analysis. A significant decrease
in $T_{1/2}$ was measured in light-activated, CoChR-tdTomato-expressing larvae compared to their
inactivated basal state (**Fig. 6L**). The control CoChR-negative larvae showed no response to
light (**Fig. 6L**). Thus, direct activation of oxytocin neurons enhances hypophyseal
permeability.

Taken together, our data show that hyperosmotic physiological challenge elicits an increase
in local blood flow velocity and permeability of the hypophyseal vasculature, where OXT is
released into the peripheral circulation. These hypophyseal vascular responses are influenced
by the local vascular geometry and are coupled to OXT neuronal activation, which is mediated
by OXT receptors. We submit that OXT may function as a self-perpetuating hormone that
facilitates its own peripheral uptake by regulating blood flow dynamics.

DISCUSSION

Neurohormones are key factors interfacing between the brain and periphery, yet not much
is known about their behavior at the site of release, and in particular, the factors affecting their
uptake into the blood. We focus here on the evolutionarily conserved HNS, which is a canonical
neuroendocrine interface wherein neurohormones are directly released from the brain into the
blood. The data we have gathered indicates that when faced with a hyperosmotic physiological
challenge, there's a noticeable increase in hypophyseal blood flow velocity and vascular

permeability where OXT is released into the peripheral circulation. The local hypophyseal vascular responses to OXT neuronal activation are influenced by vascular geometry and mediated by OXT receptors. This leads us to propose that OXT may function as a self-perpetuating hormone, regulating its own peripheral uptake by influencing blood flow dynamics and permeability.

Neurovascular coupling in the HNS

Upon stimulation of OXT neurons, OXT is released from the nerve termini into the hypophyseal perivascular space and thereafter exerts effects on peripheral target tissues. However, there is a gap in our understanding of the exact events that bring about efficient transport of OXT to the periphery. Unlike the rapid, short-distanced post-synaptic actions of neurotransmitters in the brain, OXT must travel a long distance through the dense hypophyseal perivascular space and its concentration in the blood must increase rapidly in response to physiological demands. This suggests that additional factors may be required for an efficient coordination between the release of neurohormones and their peripheral effects. We hypothesized that activity-mediated axonal release needs to be coupled to blood flow to regulate an effective transfer of OXT into the systemic circulation. However, despite extensive understanding of the electrical activity of oxytocin (OXT) neurons, the role of local blood flow in coordinating neurohormone secretion remains unclear (*30, 31*). Lafont et al. inferred the existence of such a mechanism, demonstrating that the injection of growth hormone-releasing hormone (GHRH) leads to increased local blood flow, coinciding with the timing of the growth hormone (GH) pulse in the peripheral circulation (*32*). These authors discussed the possibility that flow dynamics of the pituitary microvasculature network may influence the pulsatile secretion of growth hormone (*32*).

We show here that an osmotic challenge stimulates OXT neuronal activity concomitant with a specific local increase in volumetric blood flow in the hypophyseal capillary abutting OXT axons termini. This result is in agreement with the reported increased neurohypophyseal blood flow in chronically water-deprived rats (*33, 34*). Others have recently shown a connection between osmotic challenge and changes in supraoptic nucleus (SON) blood flow (*35*). They reported that an acute systemic salt load activates AVP neurons, decreasing blood flow and causing temporary hypoxia, which intensifies the excitatory response of AVP neurons to the systemic hyperosmotic salt challenge (*35*).

Local regulation of organ perfusion rate relies on local changes in vascular resistance, which
is controlled by capillary radius (*36, 37*). Thus, local vasoconstriction of a single capillary,
obeying Poiseuille's flow, will increase resistance and decrease flow (*20*). Having said that, the
geometry of a particular microvasculature network, including factors such as capillary length,
branching patterns and positions of inlet-outlet, plays an important role in determining flow
velocity (*22, 38*). In the case of the specific configuration of the hypophyseal vascular circuit,
we found that blood flow velocity increases upon vasoconstriction. This result also fits our
theoretically resolved relationship between blood flow velocity and capillary diameter,
considering the specific architecture, lengths, and radii of the hypophyseal vascular
components.

Role of oxytocin signaling

Vascular endothelial cells were shown to express the OXT receptor and have been
implicated in the regulation of vascular tone and blood pressure in mammals and fish (*39-43*).
It was shown that OXT can exert both vasodilation and vasoconstriction via differential
activation of either endothelial nitric oxide or ERK signaling (*39*). We show here that, as in
other species, zebrafish OXT neurons are osmoresponsive, and the effect of osmotic challenge
on hypophyseal blood flow is mediated by OXT signaling. Similarly, increased osmolality,
which was induced by dehydration, led to an increased blood flow in the rat neurohypophysis
(*33, 34*). Using Brattleboro rats, Kapitola et al. concluded that the increase in hypophyseal
blood flow following dehydration could not be mediated by AVP as these rats cannot
synthesize this neurohormone (*34*). However, it is likely that OXT is accountable for the
observed dehydration effect on Brattleboro rat hypophysis. This is supported by our findings
that the impact of hyperosmotic challenge is mediated by OXT signaling, and that direct
optogenetic stimulation of OXT neurons influences both blood flow and diameter.

We show that endothelial cells express OXT receptors and that blocking OXT signaling
attenuates the induced changes in blood flow and capillary diameter. However, it is important
to note that other cell types in the neurohypophysis may also play a role in OXT-mediated
vascular response. For example, pericytes have been implicated in the regulation of blood flow
in the brain (*44-46*). Additionally, hypophyseal astroglial pituicytes, which receive synaptoid

connections from HNS axon termini, facilitate the release of OXT and AVP to the periphery
in response to parturition, lactation, and dehydration (*29, 47, 48*).

Hypophyseal vascular permeability

Although the HNS is considered part of the central nervous system, its vascular component
has unique properties that differ from neurovascular units in other brain areas. Unlike most
BBB-containing vessels of the CNS, neurohypophyseal capillaries are fenestrated, allowing bi-
directional transfer of HNS hormones and blood-borne substances between the brain and
circulation (*13, 16, 49*). Our research shows that protein diffusion through the hypophyseal
capillary increases after a hyperosmotic challenge. Although we show that the increase in
hypophyseal blood flow velocity and permeability appears to be independent of one other, both
occur in an oxytocin-dependent manner. Thus, blocking OXT signaling decreases hypophyseal
permeability and direct activation of OXT neurons enhances permeability.

Modulation of local blood flow has been linked to hormonal uptake in several instances. For
example, somatostatin causes vasoconstriction, reducing blood flow and increasing vascular
permeability of pial vessels of the parietal cortex (*50*). Blood flow in the median eminence,
which contains fenestrated capillaries, influences food intake by regulating the access of
ghrelin to the arcuate nucleus (*51, 52*). Finally, pericytes, which regulate blood flow (*45, 46*)
were shown to regulate vascular permeability to circulating leptin (*53*). We propose that in the
case of the neurohypophysis, oxytocin orchestrates an efficient neuroendocrine secretion
machinery by affecting both hypophyseal blood flow velocity and permeability, thereby
coupling its own release and transport into the blood.

Stimulus-secretion-uptake coupling

Based on the results we have presented, we propose the following paradigm, which we term
'*stimulus-secretion-uptake coupling*': When HNS neurons are stimulated, OXT is released into
the hypophyseal parenchyma and establishes a concentration gradient of OXT between the
parenchyma and the local capillary. The diffusion of OXT into the capillary tends to bring the
system closer to equilibrium, thus reducing the diffusion of oxytocin into the vessel. The
increase in local volumetric flow, which is induced by OXT, has a dual function: **1**) enhancing
local transport of OXT to the periphery and **2**) increasing diffusion of the secreted OXT into

the blood by maintaining a local oxytocin concentration gradient. This, together with the increased permeability, which is also evoked by OXT, ensures its efficient transfer into the peripheral circulation in response to physiological demands.

REFERENCES

1. E. Scharrer, Die lichtempfindlichkeit blinder elritzen.(Untersuchungen über das zwischenhirn der fische I.). *Zeitschrift für vergleichende Physiologie*, (1928).

2. E. Scharrer, B. Scharrer, Neurosecretion. *Physiological Reviews* **25**, 171-181 (1945).

3. C. A. Pearson, M. Placzek, Development of the medial hypothalamus: forming a functional hypothalamic-neurohypophyseal interface. *Curr Top Dev Biol* **106**, 49-88 (2013).

4. E. Wircer, S. Ben-Dor, G. Levkowitz, in *Molecular neuroendocrinology : from genome to physiology,* D. G. Murphy, Harold, Ed. (John Wiley & Sons, Ltd, Chichester, West Sussex, UK ; Hoboken, NJ, USA, 2016), pp. xiii, 467 pages.

5. J. Biran, J. Blechman, E. Wircer, G. Levkowitz, in *Model Animals in Neuroendocrinology: From Worm to Mouse to Man,* M. Ludwig, G. Levkowitz, Eds. (Wiley-Blackwell, 2018), pp. 101-103.

6. P. Rajamannar, I. Arokiadhas, G. Levkowitz, J. Biran, in *Neuroanatomy of Neuroendocrine Systems,* V. Grinevich, Á. Dobolyi, Eds. (Springer International Publishing, Cham, 2021), pp. 95-118.

7. G. Leng, M. Ludwig, Neurotransmitters and peptides: whispered secrets and public announcements. *J Physiol* **586**, 5625-5632 (2008).

8. P. Le Tissier *et al.*, An updated view of hypothalamic-vascular-pituitary unit function and plasticity. *Nat Rev Endocrinol* **13**, 257-267 (2017).

9. G. Leng, N. Sabatier, Measuring Oxytocin and Vasopressin: Bioassays, Immunoassays and Random Numbers. *J Neuroendocrinol* **28**, (2016).

10. G. I. Hatton, Pituicytes, glia and control of terminal secretion. *The Journal of experimental biology* **139**, 67-79 (1988).

11. M. Schaeffer, D. J. Hodson, C. Lafont, P. Mollard, Endocrine cells and blood vessels work in tandem to generate hormone pulses. *J Mol Endocrinol* **47**, R59-66 (2011).

12.   M. Schaeffer, D. J. Hodson, C. Lafont, P. Mollard, Functional importance of blood
flow dynamics and partial oxygen pressure in the anterior pituitary. *Eur J Neurosci*
32, 2087-2095 (2010).

13.   S. Anbalagan *et al.*, Pituicyte Cues Regulate the Development of Permeable Neuro-
Vascular Interfaces. *Dev Cell* **47**, 711-726 e715 (2018).

14.   A. Gutnick *et al.*, The hypothalamic neuropeptide oxytocin is required for formation
of the neurovascular interface of the pituitary. *Dev. Cell* **21**, 642-654 (2011).

15.   Q. Chen *et al.*, Neural plate progenitors give rise to both anterior and posterior
pituitary cells. *Dev Cell* **58**, 2652-2665 e2656 (2023).

16.   L. Gordon *et al.*, The fenestrae-associated protein Plvap regulates the rate of blood-
borne protein passage into the hypophysis. *Development* **146**, (2019).

17.   J. Xie, E. Farage, M. Sugimoto, B. Anand-Apte, A novel transgenic zebrafish model
for blood-brain and blood-retinal barrier development. *BMC developmental biology*
10, 76 (2010).

18.   G. Leng, J. A. Russell, The osmoresponsiveness of oxytocin and vasopressin
neurones: Mechanisms, allostasis and evolution. *J Neuroendocrinol* **31**, e12662
(2019).

19.   I. Neumann, R. Landgraf, L. Bauce, Q. J. Pittman, Osmotic responsiveness and cross
talk involving oxytocin, but not vasopressin or amino acids, between the supraoptic
nuclei in virgin and lactating rats. *J Neurosci* **15**, 3408-3417 (1995).

20.   S. P. Sutera, R. Skalak, The History of Poiseuille's Law. *Annual Review of Fluid*
Mechanics **25**, 1-20 (1993).

21.   P. Blinder *et al.*, The cortical angiome: an interconnected vascular network with
noncolumnar patterns of blood flow. *Nat Neurosci* **16**, 889-897 (2013).

22.   X. Ji *et al.*, Brain microvasculature has a common topology with local differences in
geometry that match metabolic load. *Neuron* **109**, 1168-1187 e1113 (2021).

23.   S. Curado, D. Y. Stainier, R. M. Anderson, Nitroreductase-mediated cell/tissue
ablation in zebrafish: a spatially and temporally controlled ablation method with
applications in developmental and regeneration studies. *Nat Protoc* **3**, 948-954
(2008).

24.   D. Bergemann *et al.*, Nifurpirinol: A more potent and reliable substrate compared to
metronidazole for nitroreductase-mediated cell ablations. *Wound Repair Regen* **26**,
238-244 (2018).

25. A. R. Nunes *et al.*, Developmental Effects of Oxytocin Neurons on Social Affiliation
and Processing of Social Information. *J Neurosci* **41**, 8742-8760 (2021).

26. F. F. Zimmermann, K. V. Gaspary, A. M. Siebel, C. D. Bonan, Oxytocin reversed MK-
801-induced social interaction and aggression deficits in zebrafish. *Behavioural Brain*
Research **311**, 368-374 (2016).

27. J. Landin *et al.*, Oxytocin Receptors Regulate Social Preference in Zebrafish. *Sci Rep*
10, 5435 (2020).

28. P. Antinucci *et al.*, A calibrated optogenetic toolbox of stable zebrafish opsin lines.
Elife **9**, (2020).

29. S. Anbalagan *et al.*, Robo2 regulates synaptic oxytocin content by affecting actin
dynamics. *Elife* **8**, (2019).

30. J. A. Russell, G. Leng, A. J. Douglas, The magnocellular oxytocin system, the fount
of maternity: adaptations in pregnancy. *Front Neuroendocrinol* **24**, 27-61 (2003).

31. D. A. Poulain, J. B. Wakerley, Electrophysiology of hypothalamic magnocellular
neurones secreting oxytocin and vasopressin. *Neuroscience* **7**, 773-808 (1982).

32. C. Lafont *et al.*, Cellular in vivo imaging reveals coordinated regulation of pituitary
microcirculation and GH cell network function. *Proc Natl Acad Sci U S A* **107**, 4465-
4470 (2010).

33. B. Lichardus, I. Albrecht, J. Ponec, L. Linhart, Water deprivation for 24 hours
increases selectively blood flow in posterior pituitary of conscious rats. *Endocrinol*
Exp **11**, 99-104 (1977).

34. J. Kapitola, H. Dlouha, J. Krecek, J. Zicha, The effect of dehydration on the
neurohypophyseal blood flow in rats with hereditary diabetes insipidus. *Experientia*
33, 1615-1616 (1977).

35. R. K. Roy *et al.*, Inverse neurovascular coupling contributes to positive feedback
excitation of vasopressin neurons during a systemic homeostatic challenge. *Cell Rep*
37, 109925 (2021).

36. A. R. Pries *et al.*, Resistance to blood flow in microvessels in vivo. *Circ Res* **75**, 904-
915 (1994).

37. R. E. Shipley, D. E. Gregg, THE EFFECT OF EXTERNAL CONSTRICTION OF A
BLOOD VESSEL ON BLOOD FLOW. *American Journal of Physiology-Legacy*
Content **141**, 289-296 (1944).

38.   P. Blinder, A. Y. Shih, C. Rafie, D. Kleinfeld, Topological basis for the robust

distribution of blood to rodent neocortex. *Proc Natl Acad Sci U S A* **107**, 12670-12675

(2010).

39.   Q. Xu *et al.*, Oxytocin-induced endothelial nitric oxide dependent vasorelaxation and

ERK1/2-mediated vasoconstriction in the rat aorta. *Korean J Physiol Pharmacol* **26**,

255-262 (2022).

40.   N. Japundzic-Zigon, M. Lozic, O. Sarenac, D. Murphy, Vasopressin & Oxytocin in

Control of the Cardiovascular System: An Updated Review. *Curr Neuropharmacol*

18, 14-33 (2020).

41.   M. Petersson, T. Lundeberg, K. Uvnas-Moberg, Oxytocin decreases blood pressure in

male but not in female spontaneously hypertensive rats. *J Auton Nerv Syst* **66**, 15-18

(1997).

42.   M. Jankowski, T. L. Broderick, J. Gutkowska, The Role of Oxytocin in

Cardiovascular Protection. *Front Psychol* **11**, 2139 (2020).

43.   L. Haraldsen, V. Soderstrom-Lauritzsen, G. E. Nilsson, Oxytocin stimulates cerebral

blood flow in rainbow trout (Oncorhynchus mykiss) through a nitric oxide dependent

mechanism. *Brain Res* **929**, 10-14 (2002).

44.   C. M. Peppiatt, C. Howarth, P. Mobbs, D. Attwell, Bidirectional control of CNS

capillary diameter by pericytes. *Nature* **443**, 700-704 (2006).

45.   C. N. Hall *et al.*, Capillary pericytes regulate cerebral blood flow in health and

disease. *Nature* **508**, 55-60 (2014).

46.   D. A. Hartmann *et al.*, Brain capillary pericytes exert a substantial but slow influence

on blood flow. *Nat Neurosci* **24**, 633-645 (2021).

47.   L. Rosso, J. M. Mienville, Pituicyte modulation of neurohormone output. *Glia* **57**,

235-243 (2009).

48.   J. G. Tasker, S. H. Oliet, J. S. Bains, C. H. Brown, J. E. Stern, Glial regulation of

neuronal function: from synapse to systems physiology. *J Neuroendocrinol* **24**, 566-

576 (2012).

49.   S. Miyata, New aspects in fenestrated capillary and tissue dynamics in the sensory

circumventricular organs of adult brains. *Front Neurosci* **9**, 390 (2015).

50.   J. B. Long, D. D. Rigamonti, K. Dosaka, J. M. Kraimer, A. Martinez-Arizala,

Somatostatin causes vasoconstriction, reduces blood flow and increases vascular

permeability in the rat central nervous system. *J Pharmacol Exp Ther* **260**, 1425-1432

(1992).

51.   N. Romano *et al.*, Median eminence blood flow influences food intake by regulating
472   ghrelin access to the metabolic brain. *JCI Insight* **8**, (2023).

473   52.   M. Schaeffer *et al.*, Rapid sensing of circulating ghrelin by hypothalamic appetite-
474   modifying neurons. *Proc. Natl. Acad. Sci. USA* **110**, 1512-1517 (2013).

475   53.   L. I. Butiaeva *et al.*, Leptin receptor-expressing pericytes mediate access of
476   hypothalamic feeding centers to circulating leptin. *Cell Metab* **33**, 1433-1448 e1435
477   (2021).

478   54.   G. K. Batchelor, in *An introduction to fluid dynamics*. (Cambridge University Press,
479   2000), chap. 4.

ACKNOWLEDGEMENTS

We thank Takashi Kawashima and Pablo Blinder for valuable discussions and advice; Roy
Hofi, Estar Regev and fish facility personnel. G.L. lab is supported by the Israel Science
Foundation (#349/21); Hedda, Alberto, and David Milman Baron Center for Research on the
Development of Neural Networks; Wolfe Family Center for Research on Neuroimmunology
and Neuromodulation; Sagol Center for Research on the Aging Brain; and Maurice and
Vivienne Wohl Biology Endowment and Foundation for Higher Education and Culture. P.R.
was supported by a research grant for a student fellowship from the Benoziyo Endowment
Fund for the Advancement of Science and by the Weizmann–CNRS Collaboration Program.
G.L. is an incumbent of the Elias Sourasky Professorial Chair.

AUTHOR CONTRIBUTIONS

492   Conceptualization, P.R. and G.L.; Methodology, P.R. Investigation, P.R., O.R. and G.L.;
493 Resources, P.R., Formal analysis, P.R. and O.R.; Writing-original draft, P.R. and G.L.;
494 Writing-review and editing, P.R., G.L., and O.R.; Visualization, P.R.; Funding acquisition,
495 G.L.; Project administration, P.R. and G.L.

MATERIALS AND METHODS

Husbandry and housing conditions of experimental animals

All experiments using zebrafish were approved by the Weizmann Institute's Institutional Animal Care and Use Committee (Application number 00590123-3 and 16970919-3). Zebrafish were maintained and bred using standard protocols and according to FELASA guidelines.

Transgenic animals

The following transgenes were used in this study: *Tg(l-fabp:DBP-eGFP)*,*Tg(kdrl:eGFP)*, *Tg(oxt:gal4-VP16; UAS:NTR-mCherry; l-fabp:DBP-eGFP/kdrl:eGFP)*, *Tg(oxt:gal4-VP16; UAS:CoChR-tdTomato; l-fabp:DBP-eGFP/kdrl:eGFP)*. Larvae that were to be imaged were kept in the Danieau's buffer with 0.003% PTU from 24hpf till the day of the experiment.

Organism/ Recombinant DNA	Allele number	ZFIN ID
Tg(kdrl:eGFP)	s843	ZDB-ALT-050916-14
Tg(l-fabp:DBP-EGFP)	lri500	ZDB-ALT-120118-1
Tg(oxt:Gal4; cmlc2:EGFP)	wz06	ZDB-ALT-171113-2
Tg(UAS:NTR-mCherry)	jh17	ZDB-ALT-110222-4
Tg(UAS:CoChR-tdTomato)	u332	ZDB-TGCONSTRCT-190226-2

In-vivo imaging of live zebrafish

Six days post fertilization (dpf) zebrafish larvae were immobilized with a 30s incubation with 1mg/ml alpha-bungarotoxin followed by a quick wash in E3 media and mounted in 0.8% low-melt agarose in a custom moulded 3cm plate filled with 2% agarose. The plate was filled with 3ml E3 media prior to imaging. Samples were excited using the Zeiss LSM7 Multi-Photon Microscope with a pulsed laser (Coherent Chameleon) tuned to 870nm set at a 4-6% laser intensity (max. 210mW) through a 20x water immersion Pln Apochromatic Lens. Emitted light was detected through the BiG detector after passage through a BP500-550 filter. Line scans were obtained by manually defining a single-pixel scan path (pixel scaling of 0.03um) as required which was then scanned for a pre-defined number of frames using the Galvanometric scanning head.

Osmotic challenge, cell ablation and pharmacological treatments

Zebrafish were osmotically challenged using 50% instant ocean salt (17.5 gr/l; Instant Ocean Sea Salt, Aquarium Systems) dissolved in E3 media (5.03mM NaCl, 0.17mM KCl, 0.33mM $CaCl_2.2H_2O$, 0.33mM $MgSO_4.7H_2O$). Oxytocin neurons were ablated by treating 5 dpf *Tg(oxt:gal4-VP16; UAS:NTR-mCherry)* larvae on the background of either *Tg(l-fabp:eGFP)* or *Tg(kdrl:eGFP)* with 5uM Nifurpirinol (NFP; LGC, 13411-16-0) for 18h at 28.5^0C in E3 media. Larvae where then washed for two hours in E3 media before live imaging. Controls for the experiment were *Tg(oxt:gal4-VP16; UAS:NTR-mCherry)* negative siblings which were also treated with NFP to account for the effects of the drug itself.

Oxytocin receptor signaling was attenuated by treating 6 dpf *Tg(l-fabp:eGFP)* or *Tg(kdrl:eGFP)* larvae with 100mM Oxtr antagonist L-368,899 (No.2641, Tocris Bioscience) dissolved in E3 media for 1hour at 28.5^0C followed by live imaging.

Tgfβ receptor inhibition was carried out by treating 4 dpf *Tg(l-fabp:eGFP)* or *Tg(kdrl:eGFP)* larvae with 25uM SB431542 (S4317, Sigma-Aldrich) dissolved in DMSO for 48h at 28.5^0C followed by live imaging.

To slow down blood flow in the brain/pituitary, 6 dpf *Tg(l-fabp:eGFP)* or *Tg(kdrl:eGFP)* larvae were incubated in a-bungarotoxin for 150 seconds followed by a wash and live imaging.

Optogenetic activation of oxytocin neurons

Adult *Tg(oxt:gal4-VP16)* were crossed with *Tg(l-fabp:eGFP)* or *Tg(kdrl:eGFP)* and the resultant embryos were injected with 40ng/ul of UAS:CoChR-tdTomato (Addgene; plasmid #124233) construct and 40ng/ml of Transposase mRNA at the single-cell stage. Larvae were visually sorted at 5dpf for strong expression of tdTomato in the oxytocin neurons under a fluorescent stereo microscope. Control larvae were siblings that were not injected with the UAS:CoChR-tdTomato construct. All experimental cohorts were grown in an incubator at 28.5^0C in the dark.

Hypophyseal capillary was imaged under basal conditions, with minimal light activation of the neurons, followed by 10 minutes of rest. Oxytocin neurons were then stimulated for 5 seconds with a 470nm Fiber-coupled LED (ThorLabs Inc,USA) through a fiber optic cannula (400um,

548   0.39NA; ThorLabs) placed near the head of the larvae at 1.5 mW/mm2, followed immediately

549   by live imaging of the hypophyseal capillary only.

Monitoring and analysis of blood flow velocity

Arbitrary line scans with a length of approx. 20 um along the vessel length were obtained at

133 Hz in the *Tg(l-fabp:DBP-eGFP)* transgenic line. These line scans were then combined into

a multi-frame time-displacement image (kymograph). Two signals were derived from two

different points on the displacement axis, and along the time axis. These signals were a

representation of the fluctuation in fluorescence intensities across time. Independently, each of

these signals gives a measure of the number of red blood cells (RBCs) flowing through the

cross-sectional area of the hypophyseal capillary per unit time. On comparing the two signals

from each RBC, taken at points x1 and x2, we get the displacement of the RBC within the time

frame between t1 and t2. Using these coordinates, the slope of each RBC was computed, and

converted to velocity (**Fig. S1C**). The individual RBC velocities in the given kymograph were

then plotted as a function of normalized displacement. With an optimal sampling rate of RBCs

per second, individual velocities of the RBCs across time were used to obtain a power density

spectrum to identify the frequencies at which the flow velocities oscillate. Changes between

basal and treated conditions were also depicted as a change in power of particular frequencies

or a shift in the frequency itself using a fast Fourier transform (FFT).

Capillary diameter measurements

To measure capillary diameter, line scans were taken perpendicular to vessel length at 133 Hz

in the *Tg(kdrl:eGFP)* line. The length of the line was approx. 40um, which covered both ends

of arms of the hypophyseal capillary. These line scans were then combined into a multi-frame

time-displacement image (kymograph), where each line perpendicular to the x-axis represents

a single diameter trace at the corresponding time (**Fig. S1D**). Signals derived from each line

corresponding to time values on the x-axis were plotted to obtain two Gaussians each

representing the left or right vessel wall. Inner and outer diameter measurements were obtained

by measuring the distance between the inner and outer Full-Width Half-Maxima (FWHM)

values of these Gaussians. This diameter measured is then converted from pixels to microns

and plotted as the final graph as the diameter changes across time for both basal and treated

conditions.

Vascular permeability assay

Permeability was visualized and quantified using the *Tg(l-fabp:DBP-EGFP)* reporter by employing fluorescence recovery after photobleaching (FRAP) to determine the average recovery half-time ($T_{1/2}$) of the extravasated DBP-EGFP signal. This was carried out by defining two circular regions of interest (ROIs), each of 5μm in diameter, on either side of the hypophyseal capillary. A line scan path (pixel size 0.06um) was drawn perpendicular to direction of flow, through the vessel, and through the centers of the two ROIs. Basal measurements of raw fluorescence were obtained, followed by bleaching the ROI in the area within the hypophyseal loop (**Fig. 6A**). Photobleaching was performed at 100% laser power (approx. 3500mW) for 2.2s at 870nm, followed by the acquisition of the recovery of fluorescence along the line scan path. Raw fluorescence was extracted directly by Zen and was exported to MATLAB to calculate $T_{1/2}$ values. $T_{1/2}$ was calculated by fitting an exponential curve to the recovery florescence and calculating the intercept on the time axis at the point of 50% of the maximal fluorescence. The unbleached ROI was used as a control for unintentional bleaching, and basal fluorescence was used to validate the maximal fluorescence values after recovery.

Whole-mount immuno-fluorescence, HCR mRNA in-situ hybridization

Six day-old larvae were fixed in 4% PFA, followed by PBST washes and overnight incubation with 100% Methanol at -20C. Larvae were washed with PBST (0.5% Tween-20 in 1x Phosphate Buffered Saline), incubated in blocking solution for 1 hour at room temperature, followed by overnight incubation at 4C with Guinea Pig anti-oxytocin antibody (Peninsula Laboratories International Inc., CA, USA, no. T-5021) in a blocking solution (5% goat serum in PBST). Larvae were washed with PBST, and then incubated for 4 hours at room temperature with Mouse anti-guinea pig Cy5 secondary antibody (Jackson ImmunoReasearch Laboratories, PA, USA) followed by PBST washes.

Hybridization chain reaction (HCR) in-situ v3.0 probes for *c-fos mRNA* were custom designed and synthesized by Molecular Instruments (USA). In-situ hybridization was performed according to Chen et al. (*15*) .

Larvae were then mounted in 75% glycerol and imaged with a 40x oil immersion lens using the Zeiss LSM800 confocal setup using the appropriate emission/excitation filters.

FACS Sorting and qRT-PCR

Five adult pituitaries of the *Tg(kdrl:eGFP)* transgenic line were dissected into Ca^{2+} free Ringer's solution (116 mM NaCl, 2.9 mM KCl, 5.0 mM HEPES, pH 7.2), followed by a 30 minute incubation in 1X TrypLE Express (Gibco, 12604-013). Pituitaries were then mashed onto a 40um cell stainer and flushed into an Eppendorf using 500ul suspension buffer (L15 media without phenol red (Gibco, 21083-027), 1% FBS, 0.5% pen-strep solution). 25ul of 20ug/ml DAPI was added, and the cells were sorted into three populations: all pituitary cells (GFP+ and GFP-), endothelial cells only (GFP+), and endothelial depleted (GFP-). populations. Cells were collected in a lysis buffer (0.3% NP-40, 0.1% FBS, 1% RNase Inhibitor) and converted into cDNA directly (Takara, #RR036A). qRT-PCR was performed with 200cells per well and three technical triplicates using the following gene-specific primers:

Gene	Forward Primer (5' -> 3')	Reverse Primer (5' -> 3')
B-actin	TCTTCACTCCCCTTGTTCACAA	TGCCAACCATCACTCCCTGA
kdrl	TCCCCTTACCCTGGCTTACA	GTGGGCCTCTGAGATGGTTC
pdgfrB	TTGTCGAGGCAACAGCGTAT	CTGCGAGCCGTTGATTTGAG
oxtra	TTCTCCGTGCAGATGTGGTC	TAGAGCGTGGAGCAGCAAAA
oxtrb	TACATCATCTGCTGGACGCC	TGCATGAGATCCCGGAACAG

Codes

All algorithms to measure velocity, diameter and permeability were custom written on MATLAB (Mathworks, USA) and are available on request.

Statistical analysis

All statistical analyses and graphing were performed using PRISM 10.2 (GraphPad Software Inc, San Diego, USA). Data tables were exported from MATLAB to Prism were required. The normality of the data sets were tested using the Shapiro-Wilk test, and parametric/non-parametric tests were carried out with either a Paired t-test, Mann-Whitney test or the Wilcoxon test for paired samples. All the tests were two-tailed analyses. Welch's ANOVA with Geisser-Greenhouse corrections or Kruskal-Wallis tests was carried out where appropriate with multiple comparisons (Dunnett's, Dunn's or Tukey's). Significance thresholds for all analyses were set at a 0.05 threshold and p-values were corrected for multiple comparisons were necessary. No data points were excluded as outliers in all analyses.

Theoretical resolution of hypophyseal vascular response

Expressing Poiseuille's law for fluid flow through a vessel in terms of flow velocity and vessel radius, we get:

$$V = \frac{r^2 \Delta p}{8\mu L}$$

where r is the radius of the vessel, V is the velocity, Δp is the pressure difference between the two ends of the vessel, μ is the viscosity, and L is the length of the vessel (*54*).

To account for the geometry of the vascular microcircuit, we reduced the hypophyseal vasculature as a function of their vascular resistances (**Fig. 2G**).

$$R_{tot} = R_1 + \frac{R_2^2}{2R_2}$$

$$\text{Total Flow}, Q_{tot} = \frac{\Delta p}{R_{tot}} = \frac{\Delta p}{R_1 + \frac{R_2}{2}}$$

where R_{tot}, R_1, and R_2 are the total resistance, arterial and capillary resistance respectively, and Q_{tot} is the total flow rate.

Pressure difference in the capillary, $\Delta p_2 = Q_{tot} R_{par}$

$$\Delta p_2 = \frac{\Delta p}{R_1 + \frac{R_2}{2}} \cdot \frac{R_2}{2}$$

$$V_2 = \frac{Q_2}{A_2} = \frac{\Delta p}{2R_1 + R_2} \cdot \frac{1}{\pi r_2^2}$$

where R_{par} is the total resistance on the parallel arms of the hypophyseal capillary

Breaking down the denominator and expressing resistance in terms of radius, we get:

$$2R_1 r_2^2 + R_2 r_2^2 = \frac{8\mu}{\pi}\left(\frac{2L_1 r_2^2}{r_1^4} + \frac{L_2}{r_2^2}\right)$$

where r_1 and r_2 are the radii of the artery and capillary, L_1 and L_2 are the lengths of the artery and capillary, respectively.

654   Substituting the above equation back in the original equation, and $L_1 / L_2 = 2.5L_2$
655   (experimentally derived):

$$V_2 = \frac{r_1^4 r_2^2 \Delta p}{8\mu L_2 (5r_2^4 + r_1^4)}$$

657   From this equation, given our experimental data, we derived a range of Δp values by pair
658   matching different combinations of mean velocity and radii from out experimental cohorts. A
659   velocity vs radius graph was then plotted using a range of radii with the above pre-defined sets
660   of values for each curve (**Fig. 2H**).

    LEGENDS TO FIGURES

Figure 1. Physiological osmotic challenge increases local hypophyseal blood flow

(**A**) Representative image and scheme showing the hypophyseal neurovascular interface of the

transgenic 6-day-old zebrafish larval *Tg(oxt:eGFP;kdrl:mCherry)*. Oxytocin axonal termini

and blood vessels are labelled in green and red respectively. (**A, right**) Schema depicting the

stereotypic anatomical structure of the hypophyseal artery (HyA), capillary (HyC) and vein

(HyV), as well as the anterior mesenchephalic central artery (AMCtA), which was used as a

control. Arrows indicate the positions of line scans imaging through and across the vessel were

employed to acquire changes in capillary diameter and red blood cell (RBC) velocity traces.

(**B**) Scheme showing the production of DBP-eGFP under the liver promoter, L-fabp, in the

Tg(l-fabp:DBP-eGFP) transgenic line labelling the serum. (**C**) Snap shot of the hypophyseal

capillary loop of the *Tg(l-fabp:DBP-eGFP)* transgenic larvae at 6dpf. Dark shadows are

created by the red blood cells (RBCs) on a background of DBP-eGFP labelled serum.

(**D-E**) Transgenic *Tg(oxt:eGFP)* zebrafish larvae with labeled oxytocin (OXT) neurons were

incubated with hyperosmotic solution equivalent to 50% seawater (17.5 ppt) followed by in-

situ hybridization of the activity-dependent *c-fos mRNA*. Representative image of a single z-

plane (**D**) showing increased *c-fos* expression in OXT neurons following the osmotic challenge

quantified (**E**) as the number of *c-fos* positive OXT neurons expressing cells (p <0.0001, Mann-

Whitney test, naïve n=9, OC=10). Scale bar, 20um.

(**F**) Representative blood flow traces in the HyC at basal conditions and following the osmotic

challenge (OC) which led to increased mean RBC velocities in the HyC (**G,** p=0.001, Paired t-

test, n=9) and higher variability of velocities measured by spread of individual velocities

around the mean (**H,** p=0.0039, Wilcoxon test, n=9). Single-sided amplitude spectra of the

velocities (**I**) indicate an increase in amplitudes for frequencies less than 2Hz following the

OC.

(**J-M**) Osmotic challenge induced no changes to the blood flow traces (**J**), mean velocity (**K,**

p=0.2296, Paired t-test, n=7), variability in velocities (**L**, p=0.2669, Paired t-test, n=7), or any

changes in the spectra (**M**) in the AMCtA which served as control for the specific effect of the

OC.

Figure 2. Hypophyseal vascular response is determined by it's geometry.

Changes in capillary diameter in the HyC (**A-C**) and AMCtA (**D-F**).

(**A-F**) Representative diameter traces of HyC (**A**) and AMCtA (**D**). Mean diameter of the capillary significantly decreases after OC (**B**, p=0.0156, Wilcoxon test, n=7), as further shown as a ratio of diameter change to basal diameter (**C**, p=0.0006, Mann-Whitney test, n=7).

(**D-F**) OC did not affect the mean diameter (**E**, p=0.3076, Paired t-test, n=7), or the ratio of diameter change to basal diameter (**F**, p=0.6900, Mann-Whitney test, n=7) of the AMCtA.

(**G**) The predicted relationships between hypophyseal diameter and velocity was theoretically resolved according to Poiseuille's law for viscous fluids by representing the individual components the hypophyseal vascular microcircuit having an inlet-outlet pressure differences (Δp). The corresponding resistance (R), length (L), and radius (r) of the hypophyseal artery (HyA) and capillary (HyC) are indicated. The resulted equation of hypophyseal flow velocity (V_2) as a function of the above parameters is shown (**Fig. 2G,** for details, see '*Methods*' section).

(**H**) A range of pressure differences, Δp, was determined by multiple random matches between velocity and radii values that were obtained experimentally. This range was plugged into the resolved V_2 equation resulting three curves indicating the relationship between velocity and radii of the hypophyseal capillary for a given Δp. The measured average velocities before and after osmotic challenge (OC) are indicated by dashed lines (**H**). The point of intersection between the corresponding basal and OC velocities and the curves for a given Δp are indicated by the green and red Xs (**H'**). A corresponding increase in velocity following an OC requires a small increase in pressure and vasoconstriction (solid line) or a large decrease in radius (**H'**, dashed line).

Figure 3. Ablation of oxytocin neurons attenuates vascular response to an osmotic challenge.

(**A**) Schema representing the transgenic *Tg(oxt:gal4;UAS:NTR-mCherry)* larvae, in which oxytocin neurons express the nitroreductase (NTR) enzyme which reduces the drug nifurpirinol (NFP) into a cytotoxic compound causing specific oxytocinergic cell death.

(**B-C**) *Tg(oxt:gal4;UAS:NTR-mCherry)* larvae were treated for 18 hours with 5uM NFP and immune-stained with and antibody directed to endogenous OXT protein. NTR- (control) and NTR+ (Ablated) siblings were treated with NFP, followed by anti-OXT antibody staining. Representative confocal z-stacks images show a reduction of OXT protein in neurohypophyseal axonal projections following NFP treatment of NTR-positive larvae (**B**). Scale bar, 10um. Quantification of OXT-immunoreactive fluorescence shows significantly

reduction of OXT in neurohypophyseal termini in the NTR+ cohort (**C**, p<0.0001, Mann-Whitney test, control n=19, treated n=19).

Blood flow velocity traces representing the osmotic challenge (OC) increase in hypophyseal blood flow velocity in control (**D**), but not OXT-ablated (**E**) larvae. (**F**) Mean velocity after OC was increased in NTR- larvae (p=0.0046, Paired t-test, n=7), but not in their NTR+ siblings (p=0.0947, Paired t-test, n=7). (**G, H**) Single sided amplitude spectra of the NTR+ cohort show no difference before and after OC as compared to the controls. Variability in velocity was increased after OC in NTR- cohort also (**I**, p=0.0081, Paired t-test, n=7), but not in their NTR+ siblings (**I**, p=0.5781, Wilcoxon test, n=7).

(**J-K**) Representative diameter traces before and after osmotic challenge showing capillary vasoconstriction of control NTR- larvae (**J**) compared to vasodilation of ablated NTR+ siblings (**K**).

(**L**) Following OC, the hypophyseal capillary of NTR- larvae showed a decrease in diameter (p= 0.0156, Wilcoxon test, n=7) while NTR+ larvae showed an increase in diameter (p=0.0045, Paired t-test, n=7). The ratio between diameter change to basal diameter after OC (**M**) was significantly different between the NTR- and NTR+ cohorts (p<0.0001, Dunn's multiple comparison, NTR- n=7, NTR+ n=7).

(**N-P**) Mean velocity (p=0.3941, Paired t-test, n=7), variability of velocity (p=0.6983, Paired t-test, n=7) and diameter (p=0.5249, Paired t-test, n=7) of the AMCtA did not change after OC in NTR+ larvae treated with NFP. The control NTR- larvae treated with NFP also did not show any difference in mean velocity (p=0.3095, Paired t-test, n=7), variability of velocity (p=0.5781, Wilcoxon test, n=7) and diameter (p=0.4430, Paired t-test, n=7) in the AMCtA after OC.

Figure 4. Hypophyseal blood flow is regulated by oxytocin signaling

(**A, B**) Representative hypophyseal blood flow velocity traces before and after osmotic challenge (OC). Larvae treated with the Oxtr antagonist (Oxtr ant.) L-368,899 show no change in response to OC as compared to their control sibling. Mean velocity (**C**) after OC in the oxtr ant- cohort show an increase (p=0.0033, Paired t-test, n=7), that is not observed in L-368,899 antagonist treated siblings (p=0.2509, Paired t-test, n=7).

(**D, E**) Single sided amplitude spectra show that while the frequency spectra of a control cohort were affected the OC, larvae treated with the Oxtr antagonist show no difference. Variability in velocities (**F**) was increased following OC (p=0.0213, Paired t-test, n=7), but not in Oxtr antagonist- treated siblings (p=0.7195, Paired t-test test, n=7).

(**G, H**) Representative diameter traces showing vasoconstriction following osmotic challenge of control sibling but not in larvae treated with Oxtr antagonist (**I**) No significant change in the mean capillary diameter of the Oxtr antagonist-treated larvae which were subjected to OC (p=0.3892, Paired t-test, n=7) compared to a decrease in diameter in the control cohort (p=0.0001, Paired t-test, n=7).

(**J**) The ratio between diameter change to basal diameter after OC showed a significant difference between the control and antagonist-treated cohorts (p=0.0010, Dunn's multiple comparison, Oxtr ant- n=7, Oxtr ant+ n=7).

(**K-M**) Mean velocity (p=0.2008, Paired t-test, n=7), variability of velocity (p=0.8040, Paired t-test, n=7) and diameter (p=0.8509, Paired t-test, n=7) of the AMCtA did not change after OC in Oxtr ant+ larvae. The control Oxtr ant- larvae also did not show any difference in mean velocity (p=0.8573, Paired t-test, n=7), variability of velocity (p=0.8533, Paired t-test, n=7) and diameter (p=0.9422, Paired t-test, n=7) in the AMCtA after OC.

Figure 5. Optogenetic stimulation of oxytocin neurons increases hypophyseal blood flow and is dependent on oxytocin signaling

(**A-B**) Scheme representing the Gal4-UAS system for expression of a blue light- activated channel rhodopsin (*UAS:CoChR-tdTomato*) in oxytocin (OXT) neurons using *Tg*(oxt:Gal) transgenic driver. (**B**) Representative single z-plane confocal image of oxytocin neurons expressing CoChR-tdTomato. Scale bar, 20um.

(**C, D**) Representative blood flow velocity traces before and after light stimulation in larvae expressing CoChR show an increase in blood flow velocity as compared to their control sibling.

(**E**) Blue light stimulation of CoChR+ larvae led to an increase in mean peak blood flow velocity (p=<0.0001, Paired t-test, n=7). Control CoChR- siblings were not responsive to the light stimulation (p=0.4688, Wilcoxon test, n=7).

(**F**) The ratio between velocity change to basal velocity after light stimulation of the CoChR+ cohort increased as compared to their CoChR- sibling cohort (p=0.0225, Dunn's multiple comparison, CoChR- n=7, CoChR+ n=7).

(**G**) Representative blood flow velocity traces before and after light stimulation in larvae expressing CoChR and treated with the Oxtr antagonist L-368,899 (Oxtr ant.).

(**H**) Mean velocity after light stimulation in the CoChR+ Oxtr ant+ larvae show an increase (p=0.0313, Wilcoxon test, n=7), which was also observed in their CoChR+ Oxtr ant- siblings (p=<0.0001, Paired t-test, n=7). (**I**) The ratio between velocity change to basal velocity after light stimulation in the vehicle-treated CoChR+ was significantly larger as compared to their

CoChR+ siblings treated with L-368,899 (p=0.0287, Dunn's multiple comparison, Oxtr ant-n=7, Oxtr ant+ n=7).

(**J-K**) Representative diameter traces before and after light stimulation in larvae expressing CoChR show an increase in blood flow velocity as compared to their control sibling.

(**L-M**) Mean diameter after light stimulation in the CoChR+ larvae show a decrease (p=0.0043, Paired t-test, n=7), that is not observed in their CoChR- siblings (p=0.3349, Paired t-test, n=7). The ratio between diameter change to basal diameter after light stimulation in the CoChR+ cohort is lower as compared to their CoChR- sibling cohort (p=0.0047, Dunn's multiple comparison, CoChR- n=7, CoChR+ n=7).

(**N**) Representative diameter trace before and after light stimulation in larvae expressing CoChR and treated with the Oxtr antagonist L-368,899.

(**O**) Light stimulation of CoChR+ larvae led to a decrease in mean capillary diameter (p=0.0010, Paired t-test, n=7). The capillary diameter of the antagonist-treated CoChR+ larvae did not respond to the optogenetic activation (p=0.6043, Paired t-test, n=7).

(**P**) The ratio between diameter change to basal diameter after light stimulation of the antagonist-treated CoChR+ larvae were significantly lower as compared to vehicle-treated CoChR+ sibling cohort (p=0.0060, Dunn's multiple comparison, Oxtr ant- n=7, Oxtr ant+ n=7).

Dashed lines in all representative traces indicate a LOWESS curve fitted to the data with a smoothing window of ten points.

Figure 6. Hypophyseal vascular permeability is regulated by oxytocin signaling

(**A**) Graphical representation of the permeable hypophyseal capillary visualized in the *Tg(l-fabp:DBP-eGFP)* larvae. (**A'**) Scheme depicting the hypophyseal capillary permeability assay in which fluorescent recovery after photobleaching (FRAP) of the extravasated DBP-EGFP signal the *Tg(l-fabp:DBP-EGFP)* reporter was measured in a region of interest (ROI) within the hypophyseal capillary loop (red circle) compared to a control ROI outside of the loop. (**B**) A kymograph representing a line scan across time showing the bleached and unbleached ROIs on either side of the hypophyseal capillary, which is marked by dashed lines. The bleaching time point is marked by an arrowhead. (**C**) Average FRAP recovery half-time $T_{1/2}$ was decreased after osmotic challenge (OC) (p<0.0001, Paired t-test, n=8).

(**D**) Kymographs representing FRAP traces of naïve (unchallenged) larva displaying vehicle-treated control larva and larva treated with 25 μM of the Tgfβ receptor inhibitor SB431542

(TgfβR in.). The bleaching time point is marked by an arrowhead. Higher intensity images of

the FRAP kymographs before and after photobleaching are shown on the right (**D'**).

(**E**) The basal $T_{1/2}$ values of SB431542-treated larvae were significantly higher than their

vehicle-treated siblings ($p<0.0001$, Dunnett's multiple comparison, TgfβR inh.- n=7, TgfβR

inh.+ n=7). Larvae treated with SB431542 exhibit a decrease in $T_{1/2}$ values after OC ($p=0.0002$,

Paired t-test, n=7) as do their vehicle-treated control larvae ($p=0.0012$, Paired t-test, n=7).

(**F**) Mean blood flow velocity values following OC was increased in larvae that were treated

with the TgfβR inhibitor ($p<0.0001$, Paired t-test, n=7) as do their vehicle-treated controls

($p=0.0007$, Paired t-test, n=7).

(**G**) Representative kymographs of larvae exhibiting basal versus reduced blood flow.

(**H**) Application of 1 mg/ml of α-bungarotoxin for 30 seconds immobilizes larvae without

affecting blood flow velocity (basal), whereas incubation with α-bungarotoxin for 150 seconds

reduced mean blood flow velocity values ($p<0.0001$, Paired t-test, n=13).

(**I**) Reduction in blood flow velocity did not affect capillary permeability as indicated by FRAP

measurements of $T_{1/2}$ values ($p=0.0854$, Paired t-test, n=13).

(**J**) Hypophyseal capillary permeability was monitored at 20 mins intervals before and after the

application of Oxtr antagonist. A significant increase in $T_{1/2}$ after 40 and 60 mins of antagonist

treatment ($p=0.8090$, Basal *vs*. 20min; $p=0.0004$, Basal vs 40min; $p<0.0001$; Basal *vs*. 60min,

Dunnett's multiple comparison, n=7).

(**K**) The increment in capillary permeability which is induced by OC ($p<0.0001$, Paired t-test,

n=7) is blocked in larvae which were treated with the Oxtr antagonist L-368,899, showing no

change in $T_{1/2}$ after OC ($p=0.6147$, Paired t-test, n=7). In accord with Fig. 6J above, basal $T_{1/2}$

values of L-368,899- treated larvae were significantly higher than their untreated siblings

($p<0.0001$, Dunnett's multiple comparison, Oxtr ant- n=7, Oxtr ant+ n=7).

(**L**) Optogenetic activation of OXT neurons was performed using oxt:Gal;*UAS:CoChR-*

tdTomato transgenic larvae, as described in the legend to Figure 5 following by FRAP-based

permeability measurements. Blue light stimulation of CoChR+ larvae led to a decrease in $T_{1/2}$

values ($p=0.0003$, Paired t-test, n=7). $T_{1/2}$ values were unaffected by light in CoChR- larvae

($p=0.2559$, Paired t-test, n=7).

Supplemental Fig. S1.

(**A**) Graphical depiction of experimental workflow where either osmotic challenge or

optogenetic stimulation is administered to the zebrafish larvae followed by live *in-vivo* awake

imaging.

(**B**) Kymograph shows red blood cells (RBCs) as black streaks on a background of serum

labelled with DBP-eGFP. Signals from each RBC, taken at points x1 and x2, denote the

displacement of the RBC within the time frame between t1 and t2. The velocity is given by the

tangential derived from these components. (**C**) A kymograph across the blood vessel where

each line perpendicular to the x-axis represents a single diameter trace at the corresponding

time. $d_1(t)$ and $d_2(t)$ represent the position of the inner walls of the vessel on the displacement

axis for a given time, t.

(**D**) Diameter of the hypophyseal artery shows no change after OC (p=0.1783, Paired t-test,

n=7).

Supplemental Fig. S2.

(**A**) Adult pituitary sorted for all cells, endothelia only, and endothelia depleted populations

show an enrichment of *oxtrb* mRNA in the endothelial population. *oxtra* mRNA was enriched

in the endothelial depleted population but was still observed in low levels in the endothelia

population. All comparisons of fold change are made with respect to the mRNA levels of the

genes in the entire pituitary.

Figure 1

Figure 2

Figure 3

Figure 5

Figure 6

A

Stimulation via
optogenetics or
osmotic challenge

In-vivo
awake imaging

B

 C

D

A