SNARE mimicry by the CD225 domain of IFITM3 enables regulation of homotypic late endosome fusion

Kazi Rahman[1], Isaiah Wilt[1], Abigail A. Jolley[1], Bhabadeb Chowdhury[1], Siddhartha A. K. Datta[1], Alex A. Compton[1,*]

[1]HIV Dynamics and Replication Program, Center for Cancer Research, National Cancer Institute, Frederick, Maryland

[*]Corresponding author: alex.compton@nih.gov

Abstract

The CD225/Dispanins superfamily consists of integral membrane proteins that regulate vesicular transport and membrane fusion events driving critical cellular functions, including glucose transport, neurotransmission, and antiviral immunity. However, how the conserved CD225 domain contributes to the diverse roles played by CD225 proteins during membrane trafficking was unknown. Here, we reveal that the CD225 domain contains a SNARE-like motif that enables interaction with cellular SNARE fusogens and modulation of SNARE complex formation. We show that CD225 member proline rich transmembrane protein 2 (PRRT2) encodes an R-SNARE-like motif that enables interaction with neuronal Q-SNARE proteins. Importantly, mutations in PRRT2 linked to neurological disease in humans reside within key structural features of the R-SNARE-like motif and cause loss of Q-SNARE binding, revealing that SNARE binding via this motif in PRRT2 is critical for its role in regulating neurotransmission in vivo. Another CD225 member, interferon-induced transmembrane protein 3 (IFITM3), provides cell-intrinsic protection against Influenza A virus infection and has been shown to accelerate the trafficking of incoming virions to lysosomes for degradation. Here we demonstrate that an R-SNARE-like motif in IFITM3 is predicted to adopt a VAMP-like fold and to enable interaction with endosomal Q-SNARE proteins involved in homotypic late endosome fusion. Specifically, IFITM3 binds to Q-SNARE syntaxin 7 in cells and in vitro, and mutations in IFITM3 that abrogate syntaxin 7 binding exhibit a loss of antiviral activity against Influenza A virus, indicating that this previously uncharacterized activity is central to the antiviral mechanism of IFITM3. IFITM3 deficiency in human cells resulted in enhanced co-immunoprecipitation between Q-SNARE syntaxin 7/8 and R-SNARE VAMP8, while IFITM3 overexpression inhibited this interaction. Furthermore, overexpression of IFITM3 WT, but not a mutant of IFITM3 defective for syntaxin 7/8 binding, disrupted assembly of the SNARE complex controlling homotypic late endosome fusion. Our findings invoke a mechanistic model whereby IFITM3 diverts incoming virus towards lysosomes by preventing homotypic late endosome fusion. Our identification of an R-SNARE-like motif in many members of the CD225 family suggests that SNARE modulation plays a previously unrecognized role in the diverse membrane trafficking functions performed by these proteins and provides fresh insight into how their loss of function results in disease or susceptibility to infection.

Introduction

Membrane fusion is a critical process enabling cargo transfer between membrane-bound compartments. The soluble NSF attachment protein receptor (SNARE) proteins are key mediators of membrane fusion and are involved in nearly every vesicular trafficking event in eukaryotic cells[1,2]. SNARE proteins contain a so-called SNARE motif consisting of heptad repeats of generally hydrophobic amino acids surrounding a central hydrophilic residue (R for arginine or Q for glutamine)[3]. The central residue is referred to as the "0" layer while the hydrophobic residues surrounding it are numbered relative to the "0" layer, from "-7" to "+8" (**Figure 1A**). These features enable the assembly of four parallel alpha-helices (one R-SNARE motif and three Q-SNARE motifs) into a coiled-coil trans-SNARE complex across two membranes. Progressive folding of the bundle towards a cis-SNARE complex, in which the SNARE proteins reside within the same membrane, drives membrane fusion[4]. Humans encode a minimum of 38 SNARE proteins which interact selectively to impart specificity and directionality to membrane fusion events between different organelles[5].

Among the best studied are the synaptic SNARE proteins that mediate the exocytosis of neurotransmitters from neurons[6]. Neurotransmitter release at neuronal synapses requires the fusion of synaptic vesicles with the presynaptic plasma membrane, and this requires the assembly of syntaxin 1A and SNAP-25 (in the presynaptic plasma membrane) with VAMP2 (in the synaptic vesicle membrane)[7,8]. The study of neurological diseases associated with impaired neurotransmission has led to the identification of cellular proteins that regulate synaptic SNARE assembly, such as proline-rich transmembrane protein 2 (PRRT2)[9]. Mutations in PRRT2 associated with a range of neurological disorders in humans (including paraoxysmal kinesigenic dyskinia (PKD), infantile epilepsy and convulsions, hemiplagic migraines, ataxia, and intellectual disability) indicated that PRRT2 plays important roles in neurotransmission[9]. Subsequently, it was shown that PRRT2 interacts with Q-SNAREs syntaxin 1A and SNAP-25 and disfavors their interaction with the R-SNARE VAMP2, thereby reducing synaptic SNARE assembly and neurotransmitter release[10-13]. However, a comprehensive understanding of the protein-protein interface enabling this regulatory function of PRRT2 was unknown. PRRT2 belongs to the CD225 superfamily, a group that includes trafficking regulator of GLUT4 (TRARG1/TUSC5) and interferon-induced transmembrane protein 3 (IFITM3)[14]. TUSC5 and IFITM3 regulate vesicular trafficking during glucose transport and virus infection, respectively, through incompletely understood mechanisms.

Here, we report the identification of an R-SNARE-like motif shared among many CD225 family members that enables SNARE binding and modulation of SNARE assemblies. Mutation of residues within the R-SNARE-like motif of PRRT2, TUSC5, and IFITM3 resulted in reduced Q-SNARE binding and loss of their respective functions, as demonstrated experimentally or by associations with human disease. In-depth analysis revealed that IFITM3 interacts with endosomal SNARE proteins and, as a result, inhibits the assembly of the SNARE complex mediating homotypic late endosome fusion. Our findings suggest that cells expressing IFITM3 prioritize the heterotypic fusion pathway whereby late endosomes fuse with lysosomes, resulting in endosomal cargo degradation. Our findings demonstrate that CD225 proteins regulate diverse membrane trafficking processes in different tissues by regulating cellular fusion machinery via a conserved R-SNARE-like motif.

Results

To investigate whether the known SNARE-binding activity exhibited by human PRRT2
may be attributed to the presence of a previously unrecognized SNARE-like motif, we performed
multiple protein sequence alignment between PRRT2 and a panel of Q- and R-SNARE proteins.
We found that PRRT2 encodes an arginine that aligns with the central arginines present within
canonical R-SNARE motifs, which are approximately 53 amino acids in length (**Figure 1A**).
Furthermore, like canonical R-SNARE proteins, PRRT2 contains regularly spaced, generally
hydrophobic residues extending on both sides of the central arginine. Therefore, PRRT2 contains
a putative R-SNARE motif, resembling that which is present in VAMP proteins. Importantly, we
also found this R-SNARE-like motif in additional members of the human CD225 protein family,
including IFITM1, IFITM2, IFITM3, IFITM5, IFITM10, TUSC5, although the identify and
number of hydrophobic layers surrounding the central arginine varied between members (**Figure
1**). The central arginine was lacking in a subset of CD225 family members (**Supplemental
Figure 1A**), suggesting either that the R-SNARE-like motif does not trace its origins to the
CD225 ancestor or that the motif was lost to decay in certain family members.
Next, since PRRT2 has been described to regulate the SNARE complex controlling
neurotransmission[10], we measured whether the R-SNARE-like motif present within PRRT2
confers it with the ability to bind neuronal SNARE proteins. In agreement with previous reports,
ectopic PRRT2 co-immunoprecipitated with ectopic Q-SNARE SNAP-25 (**Figure 1B**). Attesting
to a critical role played by the R-SNARE-like motif of PRRT2 in mediating this interaction,
mutation of the central arginine (R295Q) resulted in decreased co-immunoprecipitation between
PRRT2 and SNAP-25 (**Figure 1B**). Furthermore, mutation of one of the hydrophobic layers of
the R-SNARE-like motif of PRRT2 (A320V) also abrogated the interaction with SNAP-25, and a
G305W mutation that is adjacent to a hydrophobic layer and which inhibits PRRT2
homodimerization[15] also abrogated this interaction (**Figure 1B**). Meanwhile, these mutations in
PRRT2 did not affect the interaction between PRRT2 and another Q-SNARE in the neuronal
SNARE complex, syntaxin 1A (**Supplemental Figure 1B**). Importantly, all three mutations
(R295Q, G305W, and A320V) in PRRT2 occur naturally as polymorphisms in humans and are
linked to neurological dysfunction (references)[12,13,16]. Therefore, our results describing an R-
SNARE-like motif in human PRRT2 provide a mechanistic basis for the previously described
molecular interaction between PRRT2 and neuronal SNARE proteins, as well as an explanation
for how R295Q, G305W, and A320V cause loss of PRRT2 function in vivo and neurological
dysfunction.
TUSC5 is a CD225 family member expressed in adipose tissue and implicated in the
translocation of glucose transporter type 4 (GLUT4) to the cell surface, a process essential for
insulin-stimulated glucose transport[17,18]. It was previously reported that TUSC5 co-
immunoprecipitated with GLUT4 and SNARE proteins implicated in the vesicular trafficking of
GLUT4[19,20]. However, the basis for this interaction was unknown. Like the CD225 family
member PRRT2, TUSC5 also encodes a putative R-SNARE-like motif (**Figure 1A**).
Accordingly, we found that ectopic TUSC5 co-immunoprecipitated with ectopic syntaxin 4, a Q-
SNARE involved in GLUT4 translocation, while a G141W mutation in TUSC5 (analogous to
G305W in PRRT2) disrupted this interaction (**Supplemental Figure 1C**). Collectively, these
results highlight the presence of a R-SNARE-like motif in CD225 family members that enables
coordination of SNARE proteins involved in diverse membrane fusion processes.
The IFITM proteins IFITM1, IFITM2, and IFITM3 are among the best characterized
members of the CD225 protein family due to their established roles in cell-intrinsic defense
against diverse pathogenic viruses[21,22]. IFITM3 provides constitutive and interferon-induced

protection of cells from endocytic viruses such as Influenza A virus (IAV) by homing to virus-
containing endosomes and inhibiting virus-cell membrane fusion, effectively preventing the
delivery of the viral core into host cell cytosol (where virus replication begins)[23-26]. Moreover,
Ifitm3 knockout mice and disease-associated single-nucleotide polymorphisms in human *IFITM3*
revealed that IFITM3 prevents severe viral disease sequelae in vivo[27,28]. IFITM2 arose through
gene duplication of IFITM3 in the ancestor of great apes, and both IFITM2 and IFITM3 proteins
accumulate in membranes of the endolysosomal compartment[29]. In addition to preventing virus
escape from the endocytic compartment, it has been demonstrated that IFITM3 redirects
incoming virions to lysosomes for degradation[26,30,31]. However, the mechanism for this latter
effect by IFITM3 was unknown, as was its relevance to the antiviral activity exhibited by
IFITM3. Since IFITM3 is an endolysosomal protein, we assessed whether the R-SNARE-like
motif in IFITM3 (**Figure 1A**) conferred IFITM3 with the ability to bind and regulate
endolysosomal SNARE proteins.
Fusion between compartments of the endolysosomal system are controlled by SNARE
complexes consisting of the Q-SNAREs syntaxin 7 (STX7), syntaxin 8 (STX8), Vti1b, and one
of two R-SNAREs—VAMP8 or VAMP7 (**Figure 2A**). The STX7-STX8-Vti1b-VAMP8 complex
enables late endosome-late endosome (homotypic) fusion while the STX7-STX8-Vti1b-VAMP7
complex is responsible for late endosome-lysosome (heterotypic) fusion[32-35]. Endocytic viruses
like IAV exploit the endolysosomal network and the SNARE machinery found therein in order to
avoid lysosomal degradation and gain access to host cell cytosol, where virus replication is
initiated. VAMP8 and its role in late endosome homotypic fusion was previously reported to
promote IAV entry into cells. In contrast, VAMP7 inhibited IAV entry, presumably by favoring
the degradation of virus-laden late endosomes with lysosomes[36,37]. Therefore, we tested whether
IFITM3 and the R-SNARE-like motif found in its CD225 domain could influence the assembly
of SNARE complexes regulating the endolysosomal network.
The R-SNARE-like motif is conserved among the interferon-inducible members of the
human IFITM family that exhibit antiviral activity—IFITM1, IFITM2, and IFITM3 (**Figure 2A**).
In contrast to a canonical R-SNARE protein like VAMP8, the R-SNARE-like motif of IFITM1-3
lack two layers of hydrophobicity at the amino-terminal end of the motif. Instead, IFITM1-3
encode an amphipathic helix that was shown to mediate cholesterol binding activity, membrane
binding, and which is essential for antiviral activity against IAV[15,38-40] (**Figure 2A**). Interestingly,
the intact hydrophobic layers forming part of the R-SNARE-like motif downstream of the
amphipathic helix include residues that were previously shown to be important for antiviral
activity: a cysteine residue previously shown to be palmitoylated[41-43] and two phenylalanine
residues shown to be critical for antiviral function[26,44] (**Figure 2A**). Therefore, to provide further
evidence that the R-SNARE-like motif of IFITM3 contributes functionally to its antiviral
activity, we assessed how mutation of the central arginine (R85Q) or the two phenylalanines
(F75A/F78A) impacted antiviral activity against IAV. We also tested the effect of mutating
glycine 95 (G95L), since it is adjacent to a hydrophobic layer in the R-SNARE-like motif of
IFITM3, is homologous to glycine 305 in PRRT2 and glycine 141 in TUSC5 (**Figure 1B and**
Supplemental Figure 1C), and was previously shown by us to be critical for antiviral activity[15].
While IFITM3 WT reduced IAV infection by approximately 80% relative to Empty Vector, all
mutants tested exhibited a near-total loss of antiviral activity (**Figure 2B**) despite exhibiting
similar steady-state protein levels in transfected cells (**Supplemental Figure 2B**). These results
indicate that the antiviral activity of IFITM3 against IAV depends upon a previously
uncharacterized R-SNARE-like motif. Furthermore, our findings suggest that interactions with

SNARE machinery may comprise an essential but unrecognized component of IFITM3 function in cells.

Next, we addressed whether the R-SNARE-like motif of IFITM3 adopts a protein fold resembling that of a genuine R-SNARE. Since the atomic structure of IFITM3 is incompletely understood[45] and the influence of interaction partners on its structure has not been explored, we used the TM-align platform[46] to perform protein structure alignment between the R-SNARE-like motif of IFITM3 and the R-SNARE motif of VAMP8. The results supported that IFITM3 and VAMP8 adopt a highly similar alpha helical fold in the region spanning their respective R-SNARE motifs (TM-score = 0.76) (**Supplemental Figure 2A**). To further investigate the SNARE-like structural qualities of IFITM3, we used Alphafold[47,48] and the flexible protein structure alignment algorithm FATCAT[49] to model the R-SNARE-like motif of IFITM3 against a template consisting of the R-SNARE motif of VAMP8, which was previously crystallized in the context of the STX8-STX7-Vti1b-VAMP8 complex[50]. We found that the Alphafold model for the R-SNARE-like motif of IFITM3 exhibited a high degree of structural homology with the R-SNARE motif of VAMP8 (**Figure 2C**). Therefore, IFITM3 encodes an R-SNARE-like motif that is predicted to behave like an endosomal R-SNARE protein.

To determine whether the R-SNARE-like motif of IFITM3 confers it with the ability to bind endolysosomal SNARE proteins in cells, we overexpressed IFITM3 in combination with individual endolysosomal SNARE proteins in HEK293T cells and performed co-immunoprecipitation. We found that IFITM3 immunoprecipitated from transfected cells pulled down relatively high quantities of ectopic STX7, while other endolysosomal SNARE proteins were pulled down to a lesser, or negligible, extent (**Figure 2D**). The association between IFITM3 and STX7 was confirmed by reverse co-immunoprecipitation, whereby immunoprecipitated ectopic STX7 pulled down ectopic IFITM3 (**Supplemental Figure 2C**). Ectopic STX8 also co-immunoprecipitated with IFITM3 but to a lesser extent than STX7 (**Supplemental Figure 2C**). This selective co-immunoprecipitation between IFITM3 and STX7 was not due to elevated STX7 levels in transfected cells (**Figure 2D**). Importantly, mutants of IFITM3 that caused loss of antiviral function (F75A/F78A, R85Q, and G95L) (**Figure 2B**) diminished binding with STX7 and STX8 relative to WT (**Figure 2E and Supplemental Figure 2C**). IFITM3 F75/78A and R85Q exhibited somewhat decreased steady-state levels relative to WT, possibly attesting to the importance of the R-SNARE-like motif in maintaining proper folding and stability of the IFITM3 polypeptide (**Figure 2E**). These results demonstrate that an intact R-SNARE-like motif in IFITM3 enables it to bind to endosomal SNARE proteins, particular Q-SNAREs STX7 and STX8. Furthermore, there exists a functional requirement for the R-SNARE-like motif of IFITM3 during restriction of IAV infection.

We then assessed whether endogenous IFITM3 exhibits SNARE binding activity. Following transfection of ectopic STX7 into HeLa cells competent or deficient for IFITM3, we found that ectopic STX7 pulled down endogenous IFITM3 (**Figure 3A**). As a control, pull down of endogenous IFITM3 by ectopic STX7 was lost in *IFITM3* knockout cells (**Figure 3A**). *IFITM3* knockout was specific, as IFITM2 protein levels were unaffected, and did not influence the overall levels of endolysosomal SNARE proteins (**Supplemental Figure 3A**). Moreover, the presence or absence of IFITM3 did not affect the co-immunoprecipitation between STX7 and other Q-SNAREs STX8 and Vti1b (**Supplemental Figure 3B**). We were unable to detect or immunoprecipitate endogenous STX7 in HeLa cells (data not shown). However, we succeeded in immunoprecipitating endogenous STX8 and found that it pulled down endogenous IFITM3 (**Figure 3B**). To determine whether IFITM3 can bind STX7 directly, we produced and purified

recombinant IFITM3 (**Supplemental Figure 3C-D**) and incubated it with recombinant STX7
(**Supplemental Figure 3E**) *in vitro*. Immunoprecipitation of recombinant STX7 from the
reaction resulted in pull down of recombinant IFITM3, and this interaction was lost when
F75/78A or G95L mutations were introduced into IFITM3 (**Figure 3C**). Overall, we demonstrate
for the first time that IFITM3 interacts directly with endosomal Q-SNAREs via an R-SNARE-
like motif that is critical for antiviral activity.
Since STX7 is an essential part of the SNARE complexes controlling homotypic late
endosome fusion and heterotypic late endosome fusion, we determined whether IFITM3
regulates the assembly of the STX7-STX8-Vti1b-VAMP8 complex, the STX7-STX8-Vti1b-
VAMP7 complex, or both. STX7, VAMP8, and IFITM3 were ectopically co-expressed in
HEK293T and STX7 was immunoprecipitated to identify its interaction partners. As shown
above, STX7 pulled down IFITM3 WT, while the G95L mutation in IFITM3 abrogated this
interaction (**Figure 4A**). Importantly, STX7 pulled down a reduced amount of VAMP8 in the
presence of IFITM3 relative to Empty Vector. In contrast, IFITM3 G95L did not affect the
STX7-VAMP8 interaction (**Figure 4A**). We performed additional experiments whereby STX7,
VAMP7, and IFITM3 were ectopically co-expressed. Here, IFITM3 binding to STX7 did not
inhibit the interaction between STX7 and VAMP7. Instead, the STX7-VAMP7 interaction was
slightly elevated in the presence of IFITM3 WT, but not IFITM3 G95L (**Supplemental Figure
4A**). These results indicate that IFITM3 binding to STX7 selectively impairs the interaction
between STX7 and VAMP8. To address whether endogenous IFITM3 performs a similar
function, we immunoprecipitated endogenous STX8 from HeLa. As shown above, endogenous
IFITM3 pulls down with endogenous STX8. However, endogenous STX8 pulled down increased
quantities of VAMP8 in *IFITM3* knockout cells relative to WT cells (**Figure 4B**). These results
support a model whereby IFITM3 acts as a negative regulator of the interaction between STX7/8
and VAMP8. Since it was previously shown that endocytic viruses including IAV may co-opt the
STX7-STX8-Vti1b-VAMP8 SNARE complex to promote the virus entry process[36], we
performed co-immunoprecipitation experiments in cells that had been inoculated with IAV.
Interestingly, we noticed that virus addition resulted in enhanced co-immunoprecipitation
between endogenous STX8 and VAMP8 in HEK293T cells. However, in the presence of ectopic
IFITM3 WT, this co-immunoprecipitation was diminished, while co-immunoprecipitation
between endogenous STX8 and VAMP7 was enhanced (**Figure 4C**). The IFITM3 G95L mutant,
in contrast, co-immunoprecipitated with STX8 to a lesser extent than IFITM3 WT and only
marginally impacted the STX8-VAMP8 and STX8-VAMP7 interactions (**Figure 4C**). These
findings suggest that IFITM3 disfavors the STX7/8-VAMP8 interaction while favoring the
STX7/8-VAMP7 interaction. To test whether IFITM3 reduces the assembly of the entire STX7-
STX8-Vti1b-VAMP8 SNARE complex involved in homotypic late endosome fusion, we
ectopically co-expressed STX7, STX8, Vti1b, and VAMP8 in HEK293T cells. SNARE protein
assemblies have been previously characterized as SDS-resistant complexes in whole cell lysates
under mildly denaturing conditions[51]. We identified an SDS-resistant complex approximately
200 kD in size or greater by immunoblotting for VAMP8 (**Figure 4D**) or STX8 (**Supplemental
Figure 4B**). The complex was sensitive to heat, as high molecular weight bands disappeared
after boiling lysates (**Figure 4D and Supplemental Figure 4B**). Consistent with IFITM3 acting
as a negative regulator of complex formation, complex abundance was markedly reduced in cells
expressing IFITM3 WT (**Figure 4D and Supplemental Figure 4B**). In contrast, IFITM3 G95L
only marginally reduced complex formation (**Figure 4D**). Therefore, by binding STX7/8 via an

R-SNARE-like motif and reducing their interaction with VAMP8, IFITM3 impedes formation of
the STX7-STX8-Vti1b-VAMP8 complex involved in homotypic late endosome fusion.
Our results showing that IFITM3 uses an R-SNARE-like motif to inhibit homotypic late
endosome fusion while maintaining, and possibly amplifying, the heterotypic late endosome-
lysosome fusion pathway may explain how IFITM3 redirects incoming virions towards
lysosomes for degradation[26,30,31]. To test the functional association between SNARE modulation
and lysosomal targeting of virions, we examined early steps of IAV entry into cells using
confocal immunofluorescence microscopy. In HEK293T cells stably expressing IFITM3 WT or
IFITM3 G95L, which differ with respect to STX7 binding potential (**Figure 4A**), IAV was added
to cells at a multiplicity of infection of 1 and incubated on ice for 40 mins. Cells were fed warm
medium containing Lysotracker, to visualize acidic lysosomes, and placed at 37°C to enable
synchronized internalization of virus particles. Four hours later, cells were washed, fixed and
immunostained with an antibody against IAV nucleoprotein (NP), which detects incoming,
virion-associated NP as well as de novo synthesized NP. In cells expressing Empty Vector,
nuclear NP accumulation was apparent at 4 hours post-inoculation in roughly 20% of cells,
which signals the initiation of virus replication following successful entry of the virus (i.e. escape
from the endocytic compartment) (**Figure 5A-B**). In cells expressing IFITM3 WT, nuclear NP
accumulation was virtually absent, consistent with strong inhibition of cellular entry of IAV as
reported previously[30]. In contrast, IFITM3 G95L did not reduce nuclear NP accumulation, in
agreement with a loss of antiviral activity exhibited by this mutant[15] (**Figure 5A-B**).
Interestingly, we observed a negative correlation between the extent of nuclear NP staining and
the intensity of Lysotracker staining in cells. IFITM3 WT expression resulted in intensification
of Lysotracker fluorescence, consistent with expansion of the lysosomal compartment as
described previously[30]. However, expression of IFITM3 G95L did not (**Figure 5C-D**). In cells
expressing IFITM3 G95L, most NP protein was localized to the nucleus at four hours post-
inoculation. However, some instances of extra-nuclear, bright NP puncta associated with the
Lysotracker-positive compartment could be observed, suggesting that a small proportion of
incoming IAV virions were trafficked to lysosomes in IFITM3 G95L-expressing cells (**Figure
5D**). However, the occurrence of bright NP puncta colocalizing with Lysotracker-positive
vesicles was enhanced in cells expressing IFITM3 WT (**Figure 5D and Supplemental Figure
5**). In IFITM3 WT cells, NP was found to colocalize with Lysotracker to a greater extent than in
cells expressing IFITM3 G95L as measured by Pearson's Correlation Coefficient (0.389 in WT
versus 0.023 in G95L) (**Figure 5D**). These results support the notion that inhibition of homotypic
late endosome fusion and promotion of heterotypic late endosome fusion by IFITM3 underlies
its ability to promote trafficking of IAV towards lysosomes. Altogether, this body of work
demonstrates that endolysosomal SNARE modulation is a key aspect of the mechanism by which
IFITM3 protects cells from invading viruses.
311
312 **Discussion**
313
Our identification of an R-SNARE-like motif in human CD225 proteins suggests that
these proteins act as SNARE mimics to regulate SNARE-mediated membrane fusion events.
Even though there are CD225 members that lack the polar arginine or glutamine in the central
layer characteristic of a canonical SNARE motif, they encode layered hydrophobicity which may
still afford some degree of SNARE modulation (**Supplemental Figure 1A**). For sample, Vti1a is
SNARE protein involved in neurotransmission despite encoding an aspartic acid in the central

layer[52] (**Figure 1A**). Moreover, other well-characterized SNARE proteins, like STX8, VAMP8,
VAMP7, and Vti1b lack a full complement of hydrophobic layers in their respective SNARE
motifs (**Figure 1A**). Since the CD225 family is poorly characterized and contains many proteins
with currently unknown functions[14], our discovery of built-in SNARE binding potential within
the CD225 domain may serve as a catalyst for uncovering the physiological roles they perform.
As such, we provide important insight into the birth and functional expansion of the CD225
domain over the course of evolutionary time.

A limited number of SNARE-like proteins functioning as SNARE mimics have been
described before. Two of the best characterized R-SNARE-like proteins in humans are Tomosyn
and Amisyn, which adopt VAMP-like protein folds and regulate the same neuronal SNARE
complex that PRRT2 does[53-55]. Furthermore, microbial proteins containing SNARE-like
sequence and functionality have been discovered in prokaryotes and viruses[56-60]. Interestingly,
phylogenetic evidence suggests that the CD225 domain was introduced into eukaryotes
following multiple horizontal transmission events from prokaryotes[61]. Therefore, it is possible
that the SNARE-like motif of CD225 traces its beginnings to a similar, ancient motif in
microbes.

IFITM3 was arguably the most completely understood member of the CD225 family, but
our finding that it acts as a SNARE assembly modulator significantly boosts our understanding
of its key cellular functions. The closely related proteins, IFITM1, IFITM2, IFITM5, and
IFITM10, are also likely to interact with SNARE proteins since they all share an R-SNARE-like
motif (**Figure 1A**). Interestingly, one of the most significant series of findings with regards to
IFITM function was the characterization of a conserved amphipathic alpha helix in its amino
terminus. This helix was shown to be essential for the antiviral activity of IFITM3[38] and
necessary for the ability of IFITM3 to reduce cellular membrane fluidity[15]. Following our
discovery here, we now know that this amphipathic helix is immediately adjacent to, or part of,
an R-SNARE-like motif (**Figure 2A**). Interestingly, a SNARE protein in yeast known as Spo20
also contains an amphipathic alpha helix amino terminal to its SNARE motif, resembling the
structural organization of IFITM3[62]. Moreover, another group of SNARE binding proteins
known as complexins are also bestowed with an amphipathic helix, and they use it to induce
membrane curvature and stabilize fusion pores produced by SNARE complexes[63]. Therefore, it
is possible that IFITM3 employs its amphipathic helix in conjunction with its R-SNARE-like
motif to sculpt membranes at precisely the sites where endolysosomal SNARE assemblies are
triggering membrane fusion. Other structural features lend further support to our case that
IFITM3 is a SNARE-like protein. Some SNARE proteins encode a single transmembrane
domain to directly anchor into vesicular or target membranes[1], and IFITM3 is similar in that
regard[45]. Additionally, palmitoylation is a post-translational modification of cysteine residues
that promotes membrane localization, and IFITM3 and various SNARE proteins are known to be
palmitoylated[43,64-67].

Notably, the relatively limited structural information previously available for IFITM3 did
not indicate the presence of a continuous alpha helix reminiscent of a SNARE motif. In isolation,
the region comprising the R-SNARE-like sequence motif of IFITM3 was previously thought to
form an amphipathic helix followed by a disordered, intracellular loop[38,45]. The structure of
another CD225 protein, PRRT2, was posited to contain a similarly disordered, intracellular loop
as well[68]. The typical SNARE fold characteristic of a SNARE motif may only form in the
context of a coiled-coil assembly of multiple SNARE alpha helices (trans-SNARE complex).
Our description of IFITM3 as a SNARE-like protein capable of interacting with endosomal

SNARE proteins suggests that multiple protein folds and membrane topologies of IFITM3 may
be possible, depending on the presence of interacting partners (such as STX7). Thus, attempts to
structurally characterize IFITM3 in complex with STX7 other endosomal SNAREs may provide
substantial structural insight that has been lacking.
Regarding the structural interface between IFITM3 and endosomal SNARE proteins, the
stoichiometry of the interactions remains unclear. We previously characterized a GxxxG motif
present in CD225 proteins that enables homomultimerization and which is essential to the
functions of PRRT2 and IFITM3[15]. Here we show that mutation of the GxxxG motif in PRRT2
(G305W), TUSC5 (G141W), or IFITM3 (G95L) caused loss of binding with Q-SNAREs SNAP-
25, syntaxin 4, or syntaxin 7, respectively (**Figure 1B, Supplemental Figure 1C, and**
Supplemental Figure 2C). These glycines reside adjacent to the +3 hydrophobic layer (**Figure**
1A) and may disrupt the structural integrity of the R-SNARE-like motif. Alternatively, these
findings may signal a requirement for CD225 proteins to multimerize prior to gaining SNARE
binding potential. Another possibility is that the GxxxG motif in CD225 proteins not only
controls homomultimerization but also impacts binding with other proteins including SNAREs.
What is more certain is that this newfound ability of IFITM3 to bind endosomal SNARE
proteins and inhibit homotypic late endosome fusion may provide a leap in our understanding of
how IFITM3 inhibits the cellular entry of a diverse spectrum of enveloped and non-enveloped
viruses[21]. By showing that IFITM3 inhibits homotypic late endosome fusion while reinforcing
heterotypic late endosome fusion, whereby late endosomes fuse with lysosomes, we provide an
explanation for how IFITM3 overexpression causes expansion of the endolysosomal
compartment and degradation of virions within it[30]. Taken together with previous findings, we
propose a revised model for the molecular mechanism of IFITM3-mediated antiviral activity that
is two-pronged: 1) IFITM3 traps incoming virions in late endosomes by inhibiting membrane
fusion between the viral and cellular lipid bilayers; and 2) IFITM3 promotes the fusion of late
endosomes with lysosomes to facilitate the degradation of trapped viral cargo. Promotion of late
endosome-lysosome fusion by IFITM3 is further supported by evidence showing that IFITM3
accelerates the turnover of cellular cargo in lysosomes, namely the Epidermal Growth Factor
Receptor (EGFR)[31]. Importantly, the fate of EGFR in cells is also controlled by the STX7-STX8-
Vti1b-VAMP7/8 SNARE machinery[69].
In addition to blocking infection through inhibition of virus entry into cells, IFITM3 also
performs an antiviral function in cells that are already infected—it has been shown to interact
with and inhibit the function of viral fusogens (known as viral envelope glycoproteins) found in
virions, which in turn reduces in infectivity of said virions[70-73]. Remarkably, viral fusogens share
structural and functional similarities to SNARE proteins, including heptad repeats and the
formation of coiled-coil assemblies to drive membrane merger[74,75]. Therefore, our description of
an R-SNARE-like motif in IFITM3 may reveal the molecular basis by which IFITM3 interacts
with both cellular and viral fusion machinery to control virus infections at multiple levels.
404

405 **Acknowledgements**
406 We would like to thank Alexis Schirling for assisting in molecular cloning of SNARE proteins,
407 and Guoli Shi for assisting in confocal microscopy immunofluorescence. This work was
408 supported by funding from the Intramural Research Program, National Institutes of Health.

409
410 **Declaration of interests**
411 The authors declare no competing interests.

412
413 **Materials and Methods**
414

Tissue culture, cell lines, and expression constructs
HEK293T (CRL-3216) and HeLa (CCL-2) were purchased from ATCC and cultivated at
37°C and 5% CO_2 in DMEM (Gibco) complemented with 10% fetal bovine serum (Hyclone)
and 1% penicillin-streptomycin (Gibco). HeLa *IFITM3* KO cells were generated as previously
described[76]. pQCXIP retroviral vectors encoding human IFITM3 (tagged with FLAG on amino
terminus), PRRT2 (tagged with FLAG on carboxy terminus), or TUSC5 (tagged with FLAG on
the amino terminus) were previously described[15] or synthesized by Integrated DNA Technologies
(IDT) as gBlock fragments and cloned into pQCXIP using BamH1 and EcoR1 restriction
enzymes. Mutations were introduced via QuikChange Lightning Site-Directed Mutagenesis
(Agilent). HEK293T stably expressing pQCXIP-Empty Vector, pQCXIP-IFITM3 WT, or
pQCXIP-IFITM3 G95L were described previously[15]. pcDNA3.1 encoding SNARE proteins
(SNAP25, Syntaxin 1A, Syntaxin 4, Syntaxin 7, Syntaxin 8, Vti1b, VAMP8, and VAMP7)
tagged with HA or myc at the amino terminus were synthesized by IDT as gBlock fragments and
cloned into pcDNA3.1 using BamH1 and EcoR1 restriction enyzmes.
429

Antibodies, immunoblotting, and immunoprecipitations
Whole cell lysis was performed in a buffer containing 50 mM Tris-HCl pH 7.4, 1.0%
TritonX-100 (Sigma), 150 mM NaCl, and 1 mM EDTA. Immunoprecipitation was performed as
follows: antibodies were loaded with Dynabeads (Invitrogen) for 1 hr at 4°C while rotating;
Dynabead-conjugated antibodies were added to whole cell lysates for 3 hr at 4°C while rotating
to capture antibody-antigen complexes; immunoprecipitated fractions were isolated using the
DynaMag-2 magnet (Invitrogen) and washed three times with lysis buffer prior to addition of
NuPage LDS Sample Buffer (Invitrogen) and NuPage Sample Reducing Agent (Invitrogen); the
immunoprecipitated and non-immunoprecipitated fractions were heat denatured at 100°C for 5
min; samples were loaded into Criterion XT 12% Bis-Tris polyacrylamide gels (Bio-Rad) for
SDS-PAGE using NuPAGE MES SDS Running Buffer (Invitrogen). Proteins were transferred to
Amersham Protran Nitrocellulose Membrane (GE Healthcare), and membranes were blocked
with Intercept Blocking Buffer-PBS (Li-COR) prior to incubation with primary and secondary
antibodies. The following antibodies were used in this study: anti-STX8 (110-083; Synaptic
Systems); anti-STX7 (110-072; Synaptic Systems), anti-Vti1b (164-002; Synaptic Systems),
anti-VAMP8 (104-302; Synaptic Systems), anti-VAMP7 (ab36195; Abcam), anti-HA (901514,
clone 16B12; Biolegend), anti-HA (ab9110; Abcam (this was used for immunoprecipitation)),
anti-myc (C3956, Sigma), anti-FLAG M2 (F1804; Sigma), anti-IFITM3 (ab109429; Abcam),
anti-IFITM2/3 (66081-1-Ig, Proteintech), and anti-actin (SC-47778; Santa Cruz Biotechnology).
Goat anti-mouse and goat anti-rabbit secondary antibodies conjugated to DyLight 680 or 800
(Li-COR) and the Li-COR Odyssey CLx imaging system were used for generating immunoblots,
Membrane images were analyzed using ImageStudioLite (Li-COR).
452

Sequence retrieval and protein alignments
Protein sequences for CD225 proteins, Q-SNAREs, and R-SNAREs were retrieved from
UniProt. Multi-sequence alignments were performed with MEGA11 and manually trimmed to
the 53-residue SNARE motif. UniProt numbers for SNAREs are as follows: syntaxin1, Q16623;
syntaxin4, Q12846; syntaxin 6, O43752; syntaxin7, O15400; syntaxin8, Q9UNK0; SNAP23,

O00161; Vti1a, Q96AJ9; Vti1b, Q9UEU0; VAMP1, P23763; VAMP2, P63027; VAMP3,
Q15836; VAMP5, O95183; VAMP7, P51809; VAMP8, Q9BV40; Ykt6, O15498; Syn8p,
P31377; Tomosyn/STXBP5, Q5T5C0; Amisyn/STXBP6, Q8NFX7.
461
Recombinant IFITM3 protein production
Recombinant human IFITM3 protein (WT, F65/68A, and G95L) was produced by
cloning the nucleotide sequence appended with a Strep-II tag at the amino terminus into
pET3XC and transforming it into Rosetta 2 DE3 pLysS Competent Cells (Sigma). A single
colony of transformed cells was grown in LB overnight at 37°C. One mL of the culture was
expanded in 200 mL 2YT medium (Sigma) at 37°C until the optical density reached 0.4, and the
temperature was lowered to 18°C for 1 hr before induction with 0.4 mM IPTG (Tecknova) for 20
hr. Pelleted cells were frozen at -80°C before proceeding to protein purification. Thawed cells
were resuspended in ice cold lysis buffer (at 10% w/v) consisting of 20 mM Tris-HCl pH 7.6,
100 mM NaCl, 1 mM DTT, 0.5 mg/ml lysozyme (Sigma), 1 mM PMSF (Sigma), 0.025% (v/v)
Triton X-100 (Sigma), and 1X Complete Protease Inhibitors (Roche). Sample was stirred at 4°C
for 20 min before applying 2 freeze-thaw cycles by transferring the sample container between
dry ice and a room temperature water bath. The thawed lysate was stirred at room temperature
for 15 min after adding 5 U/mL benzonase (Sigma) prior to pulse sonication for 30 s at 4°C.
Sample was clarified by centrifugation at 5000 g for 10 min and the supernatant was further
centrifuged at 40000 g for 60 min. The pellet was thoroughly resuspended in the lysis buffer
(modified to include 1 mM EDTA and excluding lysozyme) and centrifuged at 40000 g for 60
min. This process was then performed a second time. The pellet was extracted twice with 20 mM
Tris-HCl pH 7.6, 100 mM NaCl, 20% glycerol, 2% (v/v) Anapoe-X-100 (Anatrace), 1 mM DTT
(Gold Bio), and 0.1 mM AEBSF (Santa Cruz Biotechnology) at 4°C while stirring for 4 hr each.
The protein in the pellet was extracted with 20 mM Tris-HCl pH 7.6, 0.6% Sarkosyl (w/v)
(Sigma), and 2 mM DTT. Strep-II-fused IFITM3 was purified by affinity chromatography using
Strep-Tactin Sepharose Resin (IBA Lifesciences). Protein was >90% pure as determined by
SDS-PAGE and Coomassie blue staining (**Supplemental Figure 3D**). Protein concentration was
determined by Bio-Rad Protein Assay and stored at a concentration of 15-30 mg/mL at -80°C. In
the final reaction mixture of purified, recombinant IFITM3, the Sarkosyl concentration was
0.00002%.
489
In vitro protein-protein co-immunoprecipitation
Two µg of recombinant Strep-II tagged IFITM3 WT or F75/78A or G95L were incubated
with 2 µg of recombinant His-tagged STX7 (IHUSTX7RN6HISLY50UG; Innovative Research)
in 0.3 mL of binding buffer (20 mM HEPES pH 7.4, 150 mM NaCl, 0.2% TritonX-100, and
0.1% BSA) overnight at 4°C. The samples were incubated with 1 ug of IFITM3 antibody for 1
hr. The antibody-antigen complex was captured with 15 uL of Dynabeads (Invitrogen). After
washing the samples 3 times with binding buffer (lacking BSA), proteins captured by beads
using the DynaMag-2 were solubilized in 20 uL of 1X NuPAGE LDS Sample Buffer and 1X
NuPAGE Sample Reducing Reagent (Invitrogen), and heat denatured at 100°C for 5 min. SDS-
PAGE and immunoblotting were performed and proteins were detected as described above.
500
Structural modeling of the R-SNARE-like motif of IFITM3
The R-SNARE-like motif of IFITM3 (residues 55-113) and the R-SNARE motif of
VAMP8 (residues 13-65) were compared by predictive protein structure alignment and

comparison algorithm TM-align[46]. A TM score of 0.75847 was recorded, indicating that the two
regions are likely to adopt the same alpha helical protein fold. Structural comparison between the
R-SNARE-like motif of IFITM3 and the R-SNARE motif of VAMP8, which was previously
crystallized as part of the STX7-STX8-Vti1b-VAMP8 trans-SNARE complex (PDB: 1GL2)[50],
was performed using Alphafold version 3[77] with assistance from the Colabfold server[48]. A
structure for the R-SNARE-like motif of IFITM3 was predicted using the R-SNARE motif of
VAMP8 as a template. The Alphafold structure was then aligned with that of VAMP8 using the
FATCAT server[49]. The aligned structures were then visualized and reoriented for figure
presentation using ChimeraX[78].

Influenza A Virus infection
Influenza A Virus [A/PR/8/34 (PR8), H1N1] supplied as clarified allantoic fluid was
purchased from Charles River Laboratories. Infectious virus titers were calculated using a flow
cytometry-based method in HEK293T cells[79], and infections were performed as follows:
HEK293T cells (transiently transfected with 1.5 μg pQCXIP-Empty Vector, pQCXIP-FLAG-
IFITM3 WT, or pQCXIP-FLAG-IFITM3 G95L were seeded in 24-well plates (50,000 per well)
overnight and overlaid with virus diluted in 225 μL of complete DMEM (multiplicity of infection
= 0.01) for approximately 18 hr. Cells were washed with 1X D-PBS (Gibco), detached with
Trypsin-EDTA (Gibco), fixed/permeabilized with Cytofix/Cytoperm (BD), immunostained with
anti-IAV NP (ab20343; Abcam), and analyzed on a LSRFortessa flow cytometer (BD).

Influenza A Virus cellular entry assay
An 8-well μ-Slide dish (80826; Ibidi) was coated with 3 mg/mL PureCol (Advanced
Biomatrix) for 1 hr at room temperature and then allowed to dry overnight. HEK293T stably
expressing pQCXIP-Empty Vector, pQCXIP-FLAG-IFITM3 WT, or pQCXIP-FLAG-IFITM3
G95L were seeded at 20,000 cells per well in DMEM completed with 10% fetal bovine serum
and 1% penicillin-streptomycin and incubated at 37°C overnight. Cells were placed on ice for 5
min and then remained on ice while medium was removed and replaced with 120 uL complete
DMEM containing Influenza A virus PR8 (multiplicity of infection = 1). Cells mock ionoculated
with PBS served as a negative control. Inoculated cells were incubated on ice for 40 min to allow
for virus adhesion. Medium was removed, cells were washed twice with D-PBS, and medium
was replaced with 150 uL complete DMEM containing 50 nM Lysotracker Deep Red (L12492;
Invitrogen) and incubated at 37°C for 4 hr. Medium was removed, cells were washed twice with
D-PBS, and cells were fixed/permeabilized with Cytofix/Cytoperm (BD) for 10 min at room
temperature. Cells were washed twice with PermWash buffer (BD) and blocked with Intercept
Blocking Buffer (PBS) (Li-COR) for 1 hr at room temperature. Cells were immunostained with
anti-NP (BE0159; InvivoMab) at a 1:200 dilution in Intercept Antibody Diluent (PBS) (Li-COR)
for 1 hr at room temperature. Cells were washed twice in PermWash buffer and goat anti-mouse
IgG (H+L) Alexa Fluor 888 secondary antibody (Invitrogen) at 1:300 dilution in Intercept
Antibody Diluent (PBS) was added for 1 hr at room temperature. Cells were washed twice in
PermWash Buffer and Hoechst 33342 (H3570; Invitrogen) was added for 5 min at room
temperature. Cells were washed once in PermWash Buffer and overlaid with 200 uL D-PBS for
image acquisition on a Leica Stellaris 5 confocal microscope using the 63X oil objective.
Confocal Z stacks of 20-30 optical sections were acquired for each field of view, and multiple
fields were imaged per condition. Images were extracted and subjected to analysis in Fiji. To
quantify Lysotracker staining, the Corrected Total Cell Fluorescence (CTCF) was measured for

the middle Z slice of the entire field. To quantify nuclear NP accumulation, colocalization
between NP and Hoechst was measured for the middle Z slick of each cell in the field and a ratio
of NP+ nuclei: total nuclei was calculated.
553
554 *Statistics*
555   Statistical differences were determined with one-way ANOVA test corrected for multiple
556 comparisons, using an alpha cut-off of 0.05. *P* values are indicated in the figure captions.
557
558 **Figure Legends**
559
Figure 1: CD225 proteins contain a motif resembling an R-SNARE motif.
(A) A selection of SNARE proteins from yeast and mammals were aligned with proteins with
SNARE-like motifs, and a 53-residue stretch of sequence is shown. Proteins are grouped by Q-
SNARE, R-SNARE, or R-SNARE-like, and CD225 proteins are labeled as a subset of the R-
SNARE-like proteins. Residues in red text are the central polar arginine or glutamine residues
characteristic of canonical SNARE proteins and are labeled as the central "0" layer. Surrounding
heptad repeats of hydrophobic residues are labeled from "-7" to "+8" in relation to the "0" layer,
and sites highlighted in yellow correspond to hydrophobic residues. Underlined residues are
those found in PRRT2 that are associated with neurological dysfunction. (B) Left: HEK293T
cells were co-transfected with SNAP25-HA and either PRRT2-FLAG (WT, R295Q, G305W, or
A320V) or Empty Vector. SDS-PAGE and immunoblotting were performed with anti-HA and
anti-FLAG in whole cell lysates. Anti-actin was used as loading control. Right: From co-
transfected cells, SNAP25-HA was immunoprecipitated with anti-HA followed by SDS-PAGE
and immunoblotting with anti-HA and anti-FLAG. Immunoglobulin chain was used as loading
control. Numbers and tick marks left of blots indicate position and size (in kilodaltons) of protein
standard in ladder. The FLAG/HA ratio was calculated for each lane and made relative to WT
(set to 100%). Immunoblots were performed independently three times, and a representative
example is shown. Sc; Saccharomyces cerevisiae. Ig; immunoglobulin. IP; immunoprecipitation.
EV; empty vector. WT; wild-type.
579
Supplemental Figure 1:
(A) Human CD225 proteins were aligned and trimmed to a 53-residue stretch of sequence
covering the SNARE motif. Proteins are grouped by whether they contain an arginine at the
central "0" layer (R-SNARE-like) or not. Surrounding heptad repeats of hydrophobic residues
are labeled from "-7" to "+8" in relation to the "0" layer, and sites highlighted in yellow
correspond to hydrophobic residues. (B) Left: HEK293T cells were co-transfected with STX1A-
HA and either PRRT2-FLAG (WT, R295Q, G305W, or A320V) or Empty Vector. SDS-PAGE
and immunoblotting were performed with anti-HA and anti-FLAG in whole cell lysates. Anti-
actin was used as loading control. Right: From co-transfected cells, STX1A-HA was
immunoprecipitated with anti-HA followed by SDS-PAGE and immunoblotting with anti-HA
and anti-FLAG. Immunoglobulin chain was used as loading control. (C) Left: HEK293T cells
were co-transfected with STX4-HA and either TUSC5-FLAG (WT or G141W) or Empty Vector.
SDS-PAGE and immunoblotting were performed with anti-HA and anti-FLAG in whole cell
lysates. Anti-actin was used as loading control. Right: From co-transfected cells, STX4-HA was
immunoprecipitated with anti-HA followed by SDS-PAGE and immunoblotting with anti-HA
and anti-FLAG. Immunoglobulin chain was used as loading control. Numbers and tick marks left

of blots indicate position and size (in kilodaltons) of protein standard in ladder. Immunoblots
were performed independently three times, and a representative example is shown. Ig;
immunoglobulin. IP; immunoprecipitation. EV; Empty Vector. WT; wild-type.
599
Figure 2: IFITM3 contains an R-SNARE-like motif and interacts with ectopic endosomal
Q-SNARE STX7.
(A) Human endosomal SNARE proteins were aligned with aligned with human IFITM proteins,
and a 53-residue stretch of sequence is shown. Residues in red text are the central polar arginine
or glutamine residues characteristic of canonical SNARE proteins and are labeled as the central
"0" layer. Surrounding heptad repeats of hydrophobic residues are labeled from "-7" to "+8" in
relation to the "0" layer, and sites highlighted in yellow correspond to hydrophobic residues.
Black underlined residues in IFITM3 which were mutated for functional experiments. Green
underlined residues in IFITM3 correspond to the previously described amphipathic alpha helix.
(B) HEK293T cells were transfected with IFITM3-FLAG (WT, F75/78A, R85Q, or G95L) and,
48 hours later, challenged with IAV (A/PR/8/34 (PR8), H1N1) at a multiplicity of infection of
0.10. 18 hours post-infection, cells were fixed, permeabilized, immunostained with anti-FLAG
and anti-NP, and analyzed by flow cytometry. The percentage of NP+ cells was measured as a
fraction of FLAG+ cells and made relative to Empty Vector (set to 100%). Statistical differences
between WT and F75/78A, R85Q, or G95L were measured by one-way ANOVA test (*, $p <$
0.0001). (C) Structural prediction of the R-SNARE-like motif of IFITM3 was performed with
Alphafold and FATCAT. Residues 55-113 of IFITM3 were modeled against a template of the R-
SNARE motif of VAMP8 which was previously crystallized as part of the STX7-STX8-Vti1b-
VAMP8 trans-SNARE complex (PDB: 1GL2). (D) HEK293T cells were co-transfected with
IFITM3-FLAG or Empty Vector and a construct encoding a single SNARE protein: STX7-HA,
STX8-HA, Vti1b-HA, VAMP8-HA, or VAMP7-myc. SDS-PAGE and immunoblotting were
performed with anti-HA, anti-myc, and anti-FLAG in whole cell lysates. Anti-actin was used as
loading control. Right: From co-transfected cells, IFITM3-FLAG was immunoprecipitated with
anti-FLAG followed by SDS-PAGE and immunoblotting with anti-HA, anti-myc, and anti-
FLAG. Immunoglobulin chain was used as loading control. Co-transfection of Empty Vector and
STX7-HA was used as a negative control for immunoprecipitation. (E) HEK293T cells were co-
transfected with STX7-HA and either IFITM3-FLAG (WT, F75/78A, or R85Q) or Empty Vector.
SDS-PAGE and immunoblotting were performed with anti-HA and anti-FLAG in whole cell
lysates. Anti-actin was used as loading control. Right: From co-transfected cells, IFITM3-FLAG
was immunoprecipitated with anti-FLAG followed by SDS-PAGE and immunoblotting with
anti-FLAG and anti-HA. Immunoglobulin chain was used as loading control. Co-transfection of
Empty Vector and STX7-HA was used as a negative control for immunoprecipitation. The
HA/FLAG ratio was calculated for each lane and made relative to WT (set to 100%). Numbers
and tick marks left of blots indicate position and size (in kilodaltons) of protein standard in
ladder. Immunoblots were performed independently three times, and a representative example is
shown. IAV; Influenza A virus. Ig; immunoglobulin. IP; immunoprecipitation. EV; Empty
Vector. WT; wild-type.
637
638 **Supplemental Figure 2:**
639 (A) The R-SNARE-like motif of IFITM3 and the R-SNARE motif of VAMP8 (53 residues each)
640 were compared by predictive protein structure alignment and comparison algorithm TM-align. A
641 TM score of 0.75847 was recorded, indicating that the two regions are likely to adopt the same

alpha helical protein fold. (B) HEK293T cells were co-transfected with STX8-HA and either Empty Vector or IFITM3-FLAG (WT, F75/78A, or G95L). SDS-PAGE and immunoblotting were performed with anti-HA and anti-FLAG in whole cell lysates. Anti-actin was used as loading control. (C) From co-transfected cells, STX8-HA, STX7-HA, or Vti1b-HA were immunoprecipitated with anti-HA followed by SDS-PAGE and immunoblotting with anti-HA and anti-FLAG. Immunoglobulin chain was used as loading control. Co-transfection of Empty Vector and STX8-HA was used as a negative control for immunoprecipitation. The FLAG/HA ratio was calculated for each lane and made relative to WT (set to 100%). Numbers and tick marks left of blots indicate position and size (in kilodaltons) of protein standard in ladder. Immunoblots were performed independently three times, and a representative example is shown. Ig; immunoglobulin. IP; immunoprecipitation. EV; Empty Vector. WT; wild-type.

Figure 3: Endogenous IFITM3 interacts with endosomal Q-SNAREs in cells and in vitro via an R-SNARE-like motif.
(A) HeLa cells (WT or *IFITM3* KO) were transfected with STX7-HA. SDS-PAGE and immunoblotting were performed with anti-HA and anti-IFITM3 in whole cell lysates. Anti-actin was used as loading control. Right: STX7-HA was immunoprecipitated with anti-HA antibody followed by SDS-PAGE and immunoblotting with anti-HA and anti-IFITM3. Immunoglobulin chain was used as loading control. An isotype matched antibody was used as a control for immunoprecipitation. (B) Endogenous STX8 was immunoprecipitated from HeLa (WT or *IFITM3* KO) with anti-STX8 followed by SDS-PAGE and immunoblotting with anti-STX8 and anti-IFITM3. Immunoglobulin chain was used as loading control. An isotype matched antibody was used as a control for immunoprecipitation. (C) Recombinant STX7 and recombinant IFITM3 (WT, F75/78A, or G95L) were mixed together in vitro and reaction inputs were visualized by SDS-PAGE and immunoblotting with anti-STX7 and anti-IFITM3. Right: recombinant STX7 was immunoprecipitated with anti-STX7 followed by SDS-PAGE and immunoblotting with anti-STX7 and anti-IFITM3. Immunoglobulin chain was used as loading control. Numbers and tick marks left of blots indicate position and size (in kilodaltons) of protein standard in ladder. Immunoblots were performed independently three times, and a representative example is shown. Ig; immunoglobulin. IP; immunoprecipitation. EV; Empty Vector. WT; wild-type. r; recombinant.

Supplemental Figure 3:
(A) HeLa cells (WT or *IFITM3* KO) were subjected to whole cell lysis, SDS-PAGE, and immunoblotting with anti-IFITM2/3, anti-STX8, anti-Vti1b, anti-VAMP8, anti-VAMP7, and anti-actin. Red and green arrows indicate IFITM3 and IFITM2, respectively, recognized by the anti-IFITM2/3 antibody. (B) STX7-HA was transfected into HeLa (WT or *IFITM3* KO) and immunoprecipitated with anti-HA followed by SDS-PAGE and immunoblotting with anti-HA, anti-IFITM3, anti-STX8, and anti-Vti1b. Immunoglobulin chain was used as loading control. An isotype matched antibody was used as a control for immunoprecipitation. (C) Purified recombinant IFITM3 protein (WT, F75/78A, or G95L) of varying inputs (6, 12 or 18 µg) were subjected to SDS-PAGE and visualized by Coomassie stain. (D) 20 µg of purified recombinant IFITM3 protein (WT, F75/78A, or G95L) were subjected to SDS-PAGE and immunoblotting with anti-IFITM3. (E) Recombinant STX7 was subjected to SDS-PAGE and immunoblotting with anti-STX7. Numbers and tick marks left of blots indicate position and size (in kilodaltons) of protein standard in ladder. Immunoblots were performed independently three times, and a

688   representative example is shown. Ig; immunoglobulin. IP; immunoprecipitation. EV; Empty
689   Vector. WT; wild-type. r; recombinant.
690

Figure 4: IFITM3 regulates endosomal trans-SNARE assembly by inhibiting the
interaction between STX7/8 and VAMP8.
(A) HEK293T cells stably expressing Empty Vector, IFITM3 WT-FLAG, or IFITM3 G95L-
FLAG were co-transfected with STX7-HA and VAMP8-myc. Whole cell lysates were subjected
to SDS-PAGE and immunoblotting with anti-HA, anti-FLAG, and anti-myc. Actin was used as a
loading control. Right: STX7-HA was immunoprecipitated with anti-HA followed by SDS-
PAGE and immunoblotting with anti-HA, anti-FLAG, and anti-myc. The myc/HA ratio was
calculated for each lane and made relative to Empty Vector (set to 100%). (B) Endogenous STX8
was immunoprecipitated from HeLa cells (WT or *IFITM3* KO) with anti-STX8 followed by
SDS-PAGE   and   immunoblotting   with   anti-STX8,   anti-IFITM3,   and   anti-VAMP8.
Immunoglobulin chain was used as loading control. The VAMP8/STX8 ratio was calculated for
each lane and made relative to HeLa WT (set to 100%). An isotype matched antibody was used
as a control for immunoprecipitation. (C) HEK293T cells stably expressing Empty Vector,
IFITM3 WT-FLAG, or IFITM3 G95L-FLAG were inoculated with IAV (+) or medium (-) for 18
hours and subjected to whole cell lysis. Endogenous STX8 was immunoprecipitated with anti-
STX8 followed by SDS-PAGE and immunoblotting with anti-STX8, anti-FLAG, anti-VAMP8,
and anti-VAMP7. The VAMP8/STX8 and VAMP7/STX8 ratios was calculated for each lane and
made relative to Empty Vector, no virus (set to 100%). (D) HEK293T stably expressing Empty
Vector, IFITM3 WT-FLAG, or IFITM3 G95L-FLAG were co-transfected with STX7-HA,
STX8-HA, Vti1b-HA, and VAMP8-Myc. Following whole cell lysis, samples were either boiled
at 100°C (+) or not (-). SDS-PAGE and immunoblotting was performed with anti-myc and anti-
FLAG. Actin was used as loading control. Numbers and tick marks left of blots indicate position
and size (in kilodaltons) of protein standard in ladder. Immunoblots were performed
independently three times, and a representative example is shown. Ig; immunoglobulin. IP;
immunoprecipitation. EV; Empty Vector. WT; wild-type.
716

Supplemental Figure 4:
(A) HEK293T cells stably expressing Empty Vector, IFITM3 WT-FLAG, or IFITM3 G95L-
FLAG were co-transfected with STX7-HA and VAMP7-myc. Whole cell lysates were subjected
to SDS-PAGE and immunoblotting with anti-HA, anti-FLAG, and anti-myc. Actin was used as a
loading control. Right: STX7-HA was immunoprecipitated with anti-HA followed by SDS-
PAGE and immunoblotting with anti-HA, anti-FLAG, and anti-myc. The myc/HA ratio was
calculated for each lane and made relative to Empty Vector (set to 100%). (B) HEK293T stably
expressing Empty Vector or IFITM3 WT-FLAG were co-transfected with STX7-HA, STX8-HA,
Vti1b-HA, and VAMP8-Myc. Following whole cell lysis, samples were either boiled at 100°C
(+) or not (-). SDS-PAGE and immunoblotting was performed with anti-STX8 and anti-FLAG.
Actin was used as loading control. Numbers and tick marks left of blots indicate position and
size (in kilodaltons) of protein standard in ladder. Immunoblots were performed independently
three times, and a representative example is shown. IIP; immunoprecipitation. EV; Empty
Vector. WT; wild-type.
731

732   **Figure 5: IFITM3 traffics incoming IAV to lysosomes in a G95-dependent manner.**

(A) HEK293T cells stably expressing Empty Vector, IFITM3 WT-FLAG, or IFITM3 G95L-FLAG inoculated with IAV at a multiplicity of infection of 1 and incubated on ice for 40 minutes. Cells were washed, fresh medium containing Lysotracker Deep Red (50 nM) was added, and cells were placed at 37°C for 4 hours. Four hours later, cells were fixed and immunostained with anti-NP antibody and Hoechst and analyzed by confocal immunofluorescence microscopy. Mock-inoculated (PBS) Empty Vector cells were used as a negative control. Scale bar = 15 microns. Medial Z-slices representative of each condition are shown. (B) The percentage of NP+ nuclei was quantified in Empty Vector, IFITM3 WT-FLAG, or IFITM3 G95L-FLAG cells. Symbols represent fields of view containing approximately 15-30 total cells and mean percentage plus standard error is shown. (C) The Corrected Total Cell Fluorescence (CTCF) intensity of Lysotracker staining was quantified in Empty Vector, IFITM3 WT-FLAG, or IFITM3 G95L-FLAG cells. Symbols represent fields of view containing approximately 15-30 total cells and mean CTCF intensity plus standard error is shown. Statistical differences between the indicated conditions were determined by one-way ANOVA (*, $p < 0.005$). (D) Examination of NP signal in relation to Lysotracker staining in HEK293T stabling expressing IFITM3 WT-FLAG or IFITM3 G95L-FLAG. White boxes indicate fields of view selected for detailed analysis. Right: detailed view of three different Z slices from confocal stack. White boxes indicate fields of view selected for further detailed analysis. White arrows indicate bright NP puncta that colocalize with Lysotracker or which tightly appose a Lysotracker+ compartment. Colocalization between NP and Lysotracker was measured by calculating the Pearson's Correlation Coefficient using Imaris software. Coefficients were calculated from medial Z-slices from three fields of view containing 5-15 cells per condition and presented as means plus standard deviation. Scale bar = 15 microns. EV; Empty Vector. WT; wild-type. hpi; hours post-inoculation. CTCF; Corrected Total Cell Fluorescence. PCC; Pearson's Correlation Coefficient.

Supplemental Figure 5:
An additional example of IAV trafficking to lysosomes in cells expressing IFITM3 WT. A succession of Z slices from the confocal stack are shown. White arrows indicate bright NP puncta that colocalize with Lysotracker or which tightly appose a Lysotracker+ compartment. Scale bar = 5 microns.

References

1   Jahn, R. & Scheller, R. H. SNAREs — engines for membrane fusion. *Nature reviews. Molecular cell biology* **7**, 631-643 (2006). https://doi.org/10.1038/nrm2002
2   Hong, W. SNAREs and traffic. *Biochim Biophys Acta* **1744**, 493-517 (2005).
3   Fasshauer, D., Sutton, R. B., Brunger, A. T. & Jahn, R. Conserved structural features of the synaptic fusion complex: SNARE proteins reclassified as Q- and R-SNAREs. *Proc Natl Acad Sci U S A* **95**, 15781-15786 (1998). https://doi.org/10.1073/pnas.95.26.15781
4   Han, J., Pluhackova, K. & Bockmann, R. A. The Multifaceted Role of SNARE Proteins in Membrane Fusion. *Front Physiol* **8**, 5 (2017). https://doi.org/10.3389/fphys.2017.00005
5   Kloepper, T. H., Kienle, C. N. & Fasshauer, D. An elaborate classification of SNARE proteins sheds light on the conservation of the eukaryotic endomembrane system. *Mol Biol Cell* **18**, 3463-3471 (2007). https://doi.org/10.1091/mbc.e07-03-0193

779   6   Sudhof, T. C. The synaptic vesicle cycle. *Annu Rev Neurosci* **27**, 509-547 (2004).
780   https://doi.org/10.1146/annurev.neuro.26.041002.131412

781   7   Sutton, R. B., Fasshauer, D., Jahn, R. & Brunger, A. T. Crystal structure of a SNARE
782   complex involved in synaptic exocytosis at 2.4 A resolution. *Nature* **395**, 347-353 (1998).
783   https://doi.org/10.1038/26412

784   8   Xiao, W., Poirier, M. A., Bennett, M. K. & Shin, Y. K. The neuronal t-SNARE complex is
785   a parallel four-helix bundle. *Nat Struct Biol* **8**, 308-311 (2001).
786   https://doi.org/10.1038/86174

787   9   Ebrahimi-Fakhari, D., Saffari, A., Westenberger, A. & Klein, C. The evolving spectrum of
788   PRRT2-associated paroxysmal diseases. *Brain* **138**, 3476-3495 (2015).
789   https://doi.org/10.1093/brain/awv317

790   10   Coleman, J. *et al.* PRRT2 Regulates Synaptic Fusion by Directly Modulating SNARE
791   Complex Assembly. *Cell Reports* **22**, 820-831 (2018).
792   https://doi.org/10.1016/j.celrep.2017.12.056

793   11   Valente, P. *et al.* PRRT2 Is a Key Component of the Ca2+-Dependent Neurotransmitter
794   Release Machinery. *Cell Reports* **15**, 117-131 (2016).
795   https://doi.org/10.1016/j.celrep.2016.03.005

796   12   Ma, H. *et al.* A PRRT2 variant in a Chinese family with paroxysmal kinesigenic
797   dyskinesia and benign familial infantile seizures results in loss of interaction with
798   STX1B. *Epilepsia* **59**, 1621-1630 (2018). https://doi.org/10.1111/epi.14511

799   13   Tan, G. H. *et al.* PRRT2 deficiency induces paroxysmal kinesigenic dyskinesia by
800   regulating synaptic transmission in cerebellum. *Cell Res* **28**, 90-110 (2018).
801   https://doi.org/10.1038/cr.2017.128

802   14   Coomer, C. A., Rahman, K. & Compton, A. A. CD225 Proteins: A Family Portrait of
803   Fusion Regulators. *Trends in Genetics* **7**, 1-4 (2021).
804   https://doi.org/10.1016/j.tig.2021.01.004

805   15   Rahman, K. *et al.* Homology-guided identification of a conserved motif linking the
806   antiviral functions of IFITM3 to its oligomeric state. *eLife* **9**, 975-925 (2020).
807   https://doi.org/10.7554/eLife.58537

808   16   Becker, F. *et al.* PRRT2-related disorders: further PKD and ICCA cases and review of the
809   literature. *J Neurol* **260**, 1234-1244 (2013). https://doi.org/10.1007/s00415-012-6777-y

810   17   Stöckli, J., Fazakerley, D. J. & James, D. E. GLUT4 exocytosis. *Journal of Cell Science*
811   **124**, 4147-4159 (2011). https://doi.org/10.1242/jcs.097063

812   18   St-Denis, J. F. & Cushman, S. W. Role of SNARE's in the GLUT4 translocation response
813   to insulin in adipose cells and muscle. *J Basic Clin Physiol Pharmacol* **9**, 153-165
814   (1998). https://doi.org/10.1515/jbcpp.1998.9.2-4.153

815   19   Beaton, N. *et al.* TUSC5 regulates insulin-mediated adipose tissue glucose uptake by
816   modulation of GLUT4 recycling. *MOLMET* **4**, 795-810 (2015).
817   https://doi.org/10.1016/j.molmet.2015.08.003

818   20   Fazakerley, D. J. *et al.* Proteomic Analysis of GLUT4 Storage Vesicles Reveals Tumor
819   Suppressor Candidate 5 (TUSC5) as a Novel Regulator of Insulin Action in Adipocytes. *J*
820   *Biol Chem* **290**, 23528-23542 (2015). https://doi.org/10.1074/jbc.M115.657361

821   21   Majdoul, S. & Compton, A. A. Lessons in self-defence: inhibition of virus entry by
822   intrinsic immunity. *Nature Reviews Immunology* (2021). https://doi.org/10.1038/s41577-
823   021-00626-8

22 Shi, G., Schwartz, O. & Compton, A. A. More than meets the I: the diverse antiviral and
cellular functions of interferon-induced transmembrane proteins. *Retrovirology* **14**, 1-11
(2017). https://doi.org/10.1186/s12977-017-0377-y
23 Desai, T. M. *et al.* IFITM3 restricts influenza A virus entry by blocking the formation of
fusion pores following virus-endosome hemifusion. *PLoS Pathogens* **10**, e1004048
(2014). https://doi.org/10.1371/journal.ppat.1004048
24 Klein, S. *et al.* IFITM3 blocks influenza virus entry by sorting lipids and stabilizing
hemifusion. *Cell Host Microbe* **31**, 616-633 e620 (2023).
https://doi.org/10.1016/j.chom.2023.03.005
25 Li, K. *et al.* IFITM proteins restrict viral membrane hemifusion. *PLoS Pathogens* **9**,
e1003124 (2013). https://doi.org/10.1371/journal.ppat.1003124
26 Suddala, K. C. *et al.* Interferon-induced transmembrane protein 3 blocks fusion of
sensitive but not resistant viruses by partitioning into virus-carrying endosomes. *PLoS*
Pathogens **15**, e1007532-1007535 (2019). https://doi.org/10.1371/journal.ppat.1007532
27 Everitt, A. R. *et al.* IFITM3 restricts the morbidity and mortality associated with
influenza. *Nature* **484**, 519-523 (2012). https://doi.org/10.1038/nature10921
28 Bailey, C. C., Huang, I. C., Kam, C. & Farzan, M. Ifitm3 limits the severity of acute
influenza in mice. *PLoS Pathogens* **8**, e1002909 (2012).
https://doi.org/papers3://publication/doi/10.1371/journal.ppat.1002909
29 Compton, A. A. *et al.* Natural mutations in IFITM3 modulate post-translational regulation
and toggle antiviral specificity. *EMBO reports* **17**, 1657-1671 (2016).
https://doi.org/10.15252/embr.201642771
30 Feeley, E. M. *et al.* IFITM3 inhibits influenza A virus infection by preventing cytosolic
entry. *PLoS Pathogens* **7**, e1002337 (2011). https://doi.org/10.1371/journal.ppat.1002337
31 Spence, J. S. *et al.* IFITM3 directly engages and shuttles incoming virus particles to
lysosomes. *Nature chemical biology* **1**, 1-15 (2018). https://doi.org/10.1038/s41589-018-
0213-2
32 Pryor, P. R. *et al.* Combinatorial SNARE complexes with VAMP7 or VAMP8 define
different late endocytic fusion events. *EMBO reports* **5**, 590-595 (2004).
https://doi.org/10.1038/sj.embor.7400150
33 Mullock, B. M. *et al.* Syntaxin 7 is localized to late endosome compartments, associates
with Vamp 8, and Is required for late endosome-lysosome fusion. *Mol Biol Cell* **11**, 3137-
3153 (2000). https://doi.org/10.1091/mbc.11.9.3137
34 Advani, R. J. *et al.* VAMP-7 mediates vesicular transport from endosomes to lysosomes.
J Cell Biol **146**, 765-776 (1999). https://doi.org/10.1083/jcb.146.4.765
35 Prekeris, R., Yang, B., Oorschot, V., Klumperman, J. & Scheller, R. H. Differential roles
of syntaxin 7 and syntaxin 8 in endosomal trafficking. *Molecular biology of the cell* **10**,
3891-3908 (1999). https://doi.org/10.1091/mbc.10.11.3891
36 Pirooz, S. D. *et al.* UVRAG is required for virus entry through combinatorial interaction
with the class C-Vps complex and SNAREs. *Proceedings of the National Academy of*
Sciences of the United States of America **111**, 2716-2721 (2014).
https://doi.org/10.1073/pnas.1320629111
37 Pirooz, S. *et al.* Viruses customize autophagy protein for efficient viral entry. *Autophagy*
10, 1355-1356 (2014). https://doi.org/10.4161/auto.29075

868	38	Chesarino, N. M. *et al.* IFITM3 requires an amphipathic helix for antiviral activity.
869		*EMBO reports* **18**, e201744100-201744112 (2017).
870		https://doi.org/10.15252/embr.201744100
871	39	Guo, X. *et al.* Interferon-Induced Transmembrane Protein 3 Blocks Fusion of Diverse
872		Enveloped Viruses by Altering Mechanical Properties of Cell Membranes. *ACS nano*
873		(2021). https://doi.org/10.1021/acsnano.0c10567
874	40	Rahman, K. *et al.* Cholesterol Binds the Amphipathic Helix of IFITM3 and Regulates
875		Antiviral Activity. *J Mol Biol* **434**, 167759 (2022).
876		https://doi.org/10.1016/j.jmb.2022.167759
877	41	Yount, J. S. *et al.* Palmitoylome profiling reveals S-palmitoylation-dependent antiviral
878		activity of IFITM3. *Nature chemical biology* **6**, 610-614 (2010).
879		https://doi.org/10.1038/nchembio.405
880	42	Chesarino, N. M., McMichael, T. M. & Yount, J. S. Regulation of the trafficking and
881		antiviral activity of IFITM3 by post-translational modifications. *Future Microbiology* **9**,
882		1151-1163 (2014). https://doi.org/10.2217/fmb.14.65
883	43	Hach, J. C., McMichael, T., Chesarino, N. M. & Yount, J. S. Palmitoylation on Conserved
884		and Nonconserved Cysteines of Murine IFITM1 Regulates Its Stability and Anti-
885		Influenza A Virus Activity. *Journal of Virology* **87**, 9923-9927 (2013).
886		https://doi.org/10.1128/JVI.00621-13
887	44	John, S. P. *et al.* The CD225 domain of IFITM3 is required for both IFITM protein
888		association and inhibition of influenza A virus and dengue virus replication. *Journal of*
889		*Virology* **87**, 7837-7852 (2013). https://doi.org/10.1128/JVI.00481-13
890	45	Ling, S. *et al.* Combined approaches of EPR and NMR illustrate only one transmembrane
891		helix in the human IFITM3. *Scientific Reports* **6**, 1-8 (2016).
892		https://doi.org/10.1038/srep24029
893	46	Zhang, Y. & Skolnick, J. TM-align: a protein structure alignment algorithm based on the
894		TM-score. *Nucleic Acids Res* **33**, 2302-2309 (2005). https://doi.org/10.1093/nar/gki524
895	47	Varadi, M. *et al.* AlphaFold Protein Structure Database in 2024: providing structure
896		coverage for over 214 million protein sequences. *Nucleic Acids Res* **52**, D368-D375
897		(2024). https://doi.org/10.1093/nar/gkad1011
898	48	Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**,
899		679-682 (2022). https://doi.org/10.1038/s41592-022-01488-1
900	49	Li, Z., Jaroszewski, L., Iyer, M., Sedova, M. & Godzik, A. FATCAT 2.0: towards a better
901		understanding of the structural diversity of proteins. *Nucleic Acids Res* **48**, W60-W64
902		(2020). https://doi.org/10.1093/nar/gkaa443
903	50	Antonin, W., Fasshauer, D., Becker, S., Jahn, R. & Schneider, T. R. Crystal structure of
904		the endosomal SNARE complex reveals common structural principles of all SNAREs.
905		*Nat Struct Biol* **9**, 107-111 (2002). https://doi.org/10.1038/nsb746
906	51	Kubista, H., Edelbauer, H. & Boehm, S. Evidence for structural and functional diversity
907		among SDS-resistant SNARE complexes in neuroendocrine cells. *J Cell Sci* **117**, 955-966
908		(2004). https://doi.org/10.1242/jcs.00941
909	52	Tang, F. *et al.* The Role of Vti1a in Biological Functions and Its Possible Role in Nervous
910		System Disorders. *Front Mol Neurosci* **15**, 918664 (2022).
911		https://doi.org/10.3389/fnmol.2022.918664

53 Masuda, E. S., Huang, B. C., Fisher, J. M., Luo, Y. & Scheller, R. H. Tomosyn binds t-
SNARE proteins via a VAMP-like coiled coil. *Neuron* **21**, 479-480 (1998).
https://doi.org/10.1016/s0896-6273(00)80559-0
54 Scales, S. J., Hesser, B. A., Masuda, E. S. & Scheller, R. H. Amisyn, a novel syntaxin-
binding protein that may regulate SNARE complex assembly. *J Biol Chem* **277**, 28271-
28279 (2002). https://doi.org/10.1074/jbc.M204929200
55 Hatsuzawa, K., Lang, T., Fasshauer, D., Bruns, D. & Jahn, R. The R-SNARE motif of
tomosyn forms SNARE core complexes with syntaxin 1 and SNAP-25 and down-
regulates exocytosis. *J Biol Chem* **278**, 31159-31166 (2003).
https://doi.org/10.1074/jbc.M305500200
56 Shi, X., Halder, P., Yavuz, H., Jahn, R. & Shuman, H. A. Direct targeting of membrane
fusion by SNARE mimicry: Convergent evolution of Legionella effectors. *Proc Natl
Acad Sci U S A* **113**, 8807-8812 (2016). https://doi.org/10.1073/pnas.1608755113
57 Delevoye, C. *et al.* SNARE protein mimicry by an intracellular bacterium. *PLoS Pathog*
4, e1000022 (2008). https://doi.org/10.1371/journal.ppat.1000022
58 Paumet, F. *et al.* Intracellular Bacteria Encode Inhibitory SNARE-Like Proteins. *PLoS
ONE* **4**, e7375-7379 (2009). https://doi.org/10.1371/journal.pone.0007375
59 King, N. P. *et al.* Soluble NSF attachment protein receptor molecular mimicry by a
Legionella pneumophila Dot/Icm effector. *Cell Microbiol* **17**, 767-784 (2015).
https://doi.org/10.1111/cmi.12405
60 Wesolowski, J. & Paumet, F. SNARE motif: a common motif used by pathogens to
manipulate membrane fusion. *Virulence* **1**, 319-324 (2010).
https://doi.org/10.4161/viru.1.4.12195
61 Sällman Almén, M., Bringeland, N., Fredriksson, R. & Schiöth, H. B. The Dispanins: A
Novel Gene Family of Ancient Origin That Contains 14 Human Members. *PLoS ONE* **7**,
e31961 (2012). https://doi.org/10.1371/journal.pone.0031961
62 Nakanishi, H., de los Santos, P. & Neiman, A. M. Positive and negative regulation of a
SNARE protein by control of intracellular localization. *Mol Biol Cell* **15**, 1802-1815
(2004). https://doi.org/10.1091/mbc.e03-11-0798
63 Courtney, K. C. *et al.* The complexin C-terminal amphipathic helix stabilizes the fusion
pore open state by sculpting membranes. *Nat Struct Mol Biol* **29**, 97-107 (2022).
https://doi.org/10.1038/s41594-021-00716-0
64 Veit, M., Sollner, T. H. & Rothman, J. E. Multiple palmitoylation of synaptotagmin and
the t-SNARE SNAP-25. *FEBS Lett* **385**, 119-123 (1996). https://doi.org/10.1016/0014-
5793(96)00362-6
65 Fukasawa, M., Varlamov, O., Eng, W. S., Sollner, T. H. & Rothman, J. E. Localization
and activity of the SNARE Ykt6 determined by its regulatory domain and palmitoylation.
Proc Natl Acad Sci U S A **101**, 4815-4820 (2004).
https://doi.org/10.1073/pnas.0401183101
66 Yount, J. S., Karssemeijer, R. A. & Hang, H. C. S-palmitoylation and ubiquitination
differentially regulate interferon-induced transmembrane protein 3 (IFITM3)-mediated
resistance to influenza virus. *Journal of Biological Chemistry* **287**, 19631-19641 (2012).
https://doi.org/10.1074/jbc.M112.362095
67 Prekeris, R., Klumperman, J. & Scheller, R. H. Syntaxin 11 is an atypical SNARE
abundant in the immune system. *Eur J Cell Biol* **79**, 771-780 (2000).
https://doi.org/10.1078/0171-9335-00109

68   Rossi, P. *et al.* A Novel Topology of Proline-rich Transmembrane Protein 2 (PRRT2).
959   *Journal of Biological Chemistry* **291**, 6111-6123 (2016).
960   https://doi.org/10.1074/jbc.M115.683888

961   69   Futter, C. E., Pearse, A., Hewlett, L. J. & Hopkins, C. R. Multivesicular endosomes
962   containing internalized EGF-EGF receptor complexes mature and then fuse directly with
963   lysosomes. *J Cell Biol* **132**, 1011-1023 (1996). https://doi.org/10.1083/jcb.132.6.1011

964   70   Compton, A. A. *et al.* IFITM Proteins Incorporated into HIV-1 Virions Impair Viral
965   Fusion and Spread. *Cell Host & Microbe* **16**, 736-747 (2014).
966   https://doi.org/10.1016/j.chom.2014.11.001

967   71   Tartour, K. *et al.* IFITM proteins are incorporated onto HIV-1 virion particles and
968   negatively imprint their infectivity. *Retrovirology* **11**, 103 (2014).
969   https://doi.org/papers3://publication/doi/10.1186/s12977-014-0103-y

970   72   Yu, J. *et al.* IFITM Proteins Restrict HIV-1 Infection by Antagonizing the Envelope
971   Glycoprotein. *Cell Reports* **13**, 1-13 (2015). https://doi.org/10.1016/j.celrep.2015.08.055

972   73   Wang, Y. *et al.* The V3 Loop of HIV-1 Env Determines Viral Susceptibility to IFITM3
973   Impairment of Viral Infectivity. *Journal of Virology* **91**, e02441-02416-02417 (2017).
974   https://doi.org/10.1128/JVI.02441-16

975   74   Skehel, J. J. & Wiley, D. C. Coiled coils in both intracellular vesicle and viral membrane
976   fusion. *Cell* **95**, 871-874 (1998). https://doi.org/10.1016/s0092-8674(00)81710-9

977   75   Sollner, T. H. Intracellular and viral membrane fusion: a uniting mechanism. *Curr Opin*
978   *Cell Biol* **16**, 429-435 (2004). https://doi.org/10.1016/j.ceb.2004.06.015

979   76   Shi, G., Ozog, S., Torbett, B. E. & Compton, A. A. mTOR inhibitors lower an intrinsic
980   barrier to virus infection mediated by IFITM3. *Proceedings of the National Academy of*
981   *Sciences of the United States of America* **115**, E10069-E10078 (2018).
982   https://doi.org/10.1073/pnas.1811892115

983   77   Jumper, J. & Hassabis, D. Protein structure predictions to atomic accuracy with
984   AlphaFold. *Nat Methods* **19**, 11-12 (2022). https://doi.org/10.1038/s41592-021-01362-6

985   78   Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci*
986   **32**, e4792 (2023). https://doi.org/10.1002/pro.4792

987   79   Grigorov, B., Rabilloud, J., Lawrence, P. & Gerlier, D. Rapid Titration of Measles and
988   Other Viruses: Optimization with Determination of Replication Cycle Length. *PLoS ONE*
989   **6**, e24135-24112 (2011). https://doi.org/10.1371/journal.pone.0024135

Figure 1

A

| | -7 | -6 | -5 | -4 | -3 | -2 | -1 | 0 | +1 | +2 | +3 | +4 | +5 | +6 | +7 | +8 | |

Syntaxin 1a   IIKLENSIRELHDMFMDMAMLVESQGEMIDRIEYNVEHAVDYVERAVSDTKKA
Syntaxin 4   IQQLERSIRELHDIFTFLATEVEMQGEMINRIEKNILSSADYVERGQEHVKTA
Syntaxin 6   LELVSGSIGVLKNMSQRIGGELEEQAVMLEDFSHELESTQSRLDNVMKKLAKV
Syntaxin 7   IRQLEADIMDINEIFKDLGMMIHEQGDVIDSIEANVENAEVHVQQANQQLSRA
Syntaxin 8   LDALSSIISRQKQMGQEIGNELDEQNEIIDDLANLVENTDEKLRNETRRVNMV
Vti1a   LEAGYQIAVETEQIGQEMLENLSHDREKIQRARERLRETDANLGKSSRILTGM
Vti1b   IERSHRIATETDQIGSEIIEELGEQRDQLERTKSRLVNTSENLSKSRKILRSM
SNAP-25   TRRMLQLVEESKDAGIRTLVMLDEQGEQLERIEEGMDQINKDMKEAEKNLTDL
Syn8P (Sc)   LGALSQSIGRTHDISLDLNNEIVSQNDSLLVDLENLIDNNGRNLNRASRSMHG

VAMP1   LQQTQAQVEEVVDIIRVNVDKVLERDQKLSELDDRADALQAGASQFESSAAKL
VAMP2   LQQTQAQVDEVVDIMRVNVDKVLERDQKLSELDDRADALQAGASQFETSAAKL
VAMP3   LQQTQNQVDEVVDIMRVNVDKVLERDQKLSELDDRADALQAGASQFETSAAKL
VAMP5   LERCQQQANEVTEIMRNNFGKVLERGVKLAELQQRSDQLLDMSSTFNKTTQNL
VAMP7   VMETQAQVDELKGIMVRNIDLVAQRGERLELLIDKTENLVDSSVTFKTTSRNL
VAMP8   VRNLQSEVEGVKNIMTQNVERILARRGENLEHLRNKTEDLEATSEHFKTTSQKV
YKT6   MTKVQAELDETKIILHNTMESLLERGEKLDDLVSKSEVLGTQSKAFYKTARKQ

Tomosyn   IEGVKGAASGVVGELARARLALDERGKLGDLEERTAAMLSSAESFSKHAHEI
Amisyn   NSILHSAADSVTSAVQKASQALNERGERLGRAEEKTEDLKNSAQQFAETAHKL
IFITM1   SLFNTLFLNWCCLGFIAFAYSVKSRDRKMVGDVTGAQAYASTAKCLNIWALIL
IFITM2   SLFNTLFMNTCCLGFIAFAYSVKSRDRKMVGDVTGAQAYASTAKCLNIWALIL
IFITM3   55 SLFNTLFMNPCCLGFIAFAYSVKSRDRKMVGDVTGAQAYASTAKCLNIWALIL 113
TUSC5   108 LAVVACFCPVWPLNLIPLIISIMSRSSMQQGNVDGARRLGRLARLLSITLIIM 159
PRRT2   271 LAILSCFCPMWPVNIVAFAYAVMSRNSLQQGDVDGAQRLGRVAKLLSIVALVG 323

Q-SNARE

R-SNARE

R-SNARE-like CD225

B

SNAP25-HA
PRRT2-FLAG
EV   WT   R295Q G305W A320V

35 —   HA
50 —   FLAG
40 —   Actin

Whole Cell Lysates

SNAP25-HA
PRRT2-FLAG
EV   WT   R295Q G305W A320V

35 —   HA
50 —   FLAG
        100   29   <1   <1   FLAG / HA
35 —   Ig chain

Anti-HA IP

Figure 2

A

	-7	-6	-5	-4	-3	-2	-1	0	+1	+2	+3	+4	+5	+6	+7	+8

Syntaxin 7   IRQLEADIMDINEIFKDLGMMIHEQGDVIDSIEANVENAEVHVQQANQQLSRA
Syntaxin 8   LDALSSIISRQKQMGQEIGNELEDQNEIIDDLANLVENTDEKLRNETRRVNMV
Vti1b   IERSHRIATETDQIGSEIIEELGEQRDQLERTKSRLVNTSENLSKSRKILRSM
VAMP8   VRNLQSEVEGVKNIMTQNVERILARGENLEHLRNKTEDLEATSEHFKTTSQKV
VAMP7   VMETQAQVDELKGIMVRNIDLVAQRGERLELLIDKTENLVDSSVTFKTTSRNL

IFITM1   SLFNTLFLNMCCLGFIAFAYSVKSRDRKMVGDVTGAQAYASTAKCLNIMALIL
IFITM2   SLFNTLFMNTCCLGFIAFAYSVKSRDRKMVGDVTGAQAYASTAKCLNIMALIL
IFITM3   55 SLFNTLFMNPCCLGFIAFAYSVKSRDRKMVGDVTGAQAYASTAKCLNIWALIL 113

B

IAV Infection
(Relative to EV)

EV
WT
F75/78A
R85Q
G95L

C

VAMP8

IFITM3

Overlay

VAMP8 STX8 STX7
Vti1b

D

Whole Cell Lysates

EV | IFITM3-FLAG
STX7-HA, STX7-HA, STX8-HA, Vti1b-HA, VAMP8-HA, VAMP7-Myc

HA/Myc
FLAG
Actin
40 —
15 —
40 —

Anti-FLAG IP

EV | IFITM3-FLAG
STX7-HA, STX7-HA, STX8-HA, Vti1b-HA, VAMP8-HA, VAMP7-Myc

HA/Myc
FLAG
Ig chain
40 —
15 —
35 —

E

Whole Cell Lysates

STX7-HA
IFITM3-FLAG
EV | WT | F75A, F78A | R85Q

FLAG
HA
Actin
15 —
40 —
40 —

Anti-FLAG IP

STX7-HA
IFITM3-FLAG
EV | WT | F75A, F78A | R85Q

FLAG
HA
HA / FLAG: 100 29 37
Ig chain
15 —
40 —
35 —

Figure 3

Figure 4

Figure 5

NP / Lyso PCC = 0.023 -/+ 0.052

NP / Lyso PCC = 0.389 -/+ 0.096