Renal cancer cells acquire immune surface protein through trogocytosis and horizontal gene transfer

Haley Q. Marcarian[1,2], Anutr Sivakoses[1,2], Anika M. Arias[1], Olivia C. Ihedioha[2], Benjamin R. Lee[3], Maria C. Bishop[4], and Alfred L.M. Bothwell[1,2,5]

[1]Cancer Biology Graduate Program, University of Arizona, Tucson, AZ, 85724.

[2]Department of Pathology, Microbiology, and Immunology, University of Nebraska Medical Center, Omaha, NE, 68106.

[3]Department of Urology, University of Arizona, Tucson, AZ, 85724.

[4]College of Medicine, University of Arizona, Tucson, AZ, 85724.

[5]Department of Immunobiology, Yale University School of Medicine, New Haven, CT, 06520.

*Correspondence to:

Alfred L.M. Bothwell

albothwell@unmc.edu, (402)-559-8050

505 S 45th St. Omaha, NE 68106, 4.12.321

Keywords: Trogocytosis, horizontal gene transfer, cancer, tumor-immune microenvironment

SIGNIFICANCE STATEMENT

We have identified trogocytosis as a mechanism by which human clear cell renal

carcinoma tumors acquire lymphocyte surface protein expression from tumor infiltrating

immune cells. In addition to the transfer of membrane fragments, we have provided

evidence to show that genomic DNA is transferred from a normal immune cell to a tumor

cell during trogocytosis. This process alters the transcriptome of cancer cells such that

they express significantly more mRNA for immune proteins such as the lymphocyte

marker CD45 compared to tumor cells that have not undergone trogocytosis. This study

provides an in-depth analysis of the interactions between cancer cells and tumor

infiltrating lymphocytes, and how these interactions alter the development of human

tumors.

ABSTRACT

Trogocytosis is an underappreciated phenomenon that shapes the immune

microenvironment surrounding many types of solid tumors. The consequences of

membrane-bound proteins being deposited from a donor immune cell to a recipient

cancer cell via trogocytosis are still unclear. Here, we report that human clear cell renal

carcinoma tumors stably express the lymphoid markers CD45, CD56, CD14, and CD16.

Flow cytometry performed on fresh kidney tumors revealed consistent CD45 expression

on tumor cells, as well as varying levels of the other markers mentioned previously. These

results were consistent with our immunofluorescent analysis, which also revealed

colocalization of lymphoid markers with carbonic anhydrase 9 (CAIX), a standard kidney

tumor marker. RNA analysis showed a significant upregulation of genes typically

associated with immune cells in tumor cells following trogocytosis. Finally, we show

evidence of chromosomal DNA being transferred from immune cells to tumor cells during

trogocytosis. This horizontal gene transfer has transcriptional consequences in the

recipient tumor cell, resulting in a fusion phenotype that expressed both immune and

cancer specific proteins. This work demonstrates a novel mechanism by which tumor cell

protein expression is altered through the acquisition of surface membrane fragments and

genomic DNA from infiltrating lymphocytes. These results alter the way in which we

understand tumor-immune cell interactions and may reveal new insights into the

mechanisms by which tumors develop. Additionally, further studies into trogocytosis will

help push the field towards the next generation of immunotherapies and biomarkers for

treating renal cell carcinoma and other types of cancers.

54   INTRODUCTION

Trogocytosis is the transfer of membrane fragments from a donor cell to an

acceptor cell [1]. In mammalian cells, this process was first described on CD8$^+$ T cells, in

which the CD8$^+$ T cells were shown to take fragments of the plasma membrane from

certain antigen presenting cells (APCs) during the formation of the immunological

synapse [2]. Since then, trogocytosis has been observed in many different types of

immune cells, including CD4$^+$ T cells [3], Natural Killer (NK) cells [4], and macrophages

[5]. Previously, we have shown that trogocytosis plays an important role in the acquisition

of immune regulatory molecules by colon cancer cells from infiltrating lymphocytes [6].

This resulted in increased immune regulator molecules such as CTLA4, PDL1, Tim3,

VISTA, LAG3, CD38, CD80, CD86, MHC Class II, and PD-L1 being presented on the

surface of trogocytic tumor cells compared to non-trogocytic tumor cells. Trogocytosis

was also observed with numerous lymphocyte populations including macrophages, dendritic cells, NK cells, and monocytes[6]. Along with the transfer of surface proteins, genomic DNA PCR analysis also demonstrates the transfer of DNA from donor immune cells to mouse CRC cells, implying that tumor cells can display an altered transcriptome because of trogocytosis. Trogocytosis has also been implicated in other types of disease such as Hodgkin lymphoma and renal cell carcinoma[7, 8]. In mouse models of leukemia, trogocytosis promotes the transfer of target antigens from cancer cells to T cells, which induces T cell exhaustion and fratricide [9]. T cell exhaustion and fratricide emphasize the observation of relapse in patients treated with CAR T cell therapies. However, the extent to which trogocytosis occurs in tumors as well as the implications it has in shaping the tumor-immune microenvironment have not been fully elucidated.

Horizontal gene transfer (HGT) is a process that has been widely studied in the field of microbiology. HGT between different species of bacteria has been shown to grant survival advantages to the recipient microbe, such as resistance to antibiotics[10]. However, the concept of horizontal gene transfer between mammalian cells is largely uninvestigated. Recently, it has been shown that RPE1 cells, an immortalized non-transformed pigmented epithelium line can integrate a genomically encoded fluorescent reporter from a breast cancer cell line expressing that reporter when the two lines are cocultured[11]. Interestingly, physical contact was required between the two cell lines for HGT to occur. This indicates that HGT could occur concurrently with trogocytosis. To our knowledge, this is the first evidence of HGT between mammalian cells, however the mechanism behind such a transfer remains unknown. The consequences of this transfer, particularly in the context of cancer development and progression, remain uninvestigated.

In this study, we detected the presence of the immune markers CD14, CD16,

CD56, and CD45 on tumor cells from renal cell carcinoma (RCC) patients. Previously,

these markers have not been identified on RCC cells or other solid tumors. We suggest

they were acquired through trogocytosis of infiltrating immune cells as these tumors

develop. Other markers, including CD19 and CD11b were also tested and determined to

not be significant markers of trogocytosis. These results were determined through

quantification of immunofluorescent staining and flow cytometry. RNA analysis further

showed trogocytosis between RCC cells and infiltrating lymphocytes resulted in an

altered phenotype of tumor cells that expressed both tumor-specific and immune-specific

genes. NanoString analysis of trogocytic tumor cells from human RCC tumors showed

that trogocytic cells display rampant expression of immune cell-associated and tumor-

associated genes. Finally, in vitro analysis of EdU and a genomically encoded GFP tag

transfer from T cells to RCC cells showed that genomic DNA is transferred during the

process of trogocytosis.

RESULTS

Identification and Quantification of Tumor Cells Presenting both Tumor and Lymphocyte

Markers in Human RCC Samples

Macrophages have been recently implicated in abating the effects of immune

checkpoint inhibitors (ICI) in treating RCC through the process of trogocytosis with cancer

cells [8]. Given that RCC is an immunogenic cancer with high levels of immune infiltration,

we hypothesized that other types of immune cells may play an important role in shaping

the tumor microenvironment through trogocytosis. We obtained paraffin embedded slides

from 21 human RCC tumors of varying stages to determine whether markers of

trogocytosis could be detected. Positive staining for carbonic anhydrase 9 (CAIX) was

used to define tumor cells as it is a well-characterized marker of hypoxia and is commonly

upregulated in RCC cells [12].

To identify trogocytic tumor cells, slides were stained with CAIX and various

immune markers of interest. Fluorescent imaging revealed significant populations of cells

displaying both CAIX and CD14, CD16, CD56, or CD45RA (Figure 1A). Imaging of normal

kidney tissue revealed no expression of these markers by healthy kidney epithelial cells

(Supplemental Figure 1A). These results revealed that macrophages, monocytes, NK

cells, and CD45RA$^+$ T cells are likely the most active participants of trogocytosis with RCC

cells. The RA isoform of CD45 is typically expressed on naïve CD4$^+$ and CD8$^+$ T cells,

until it is converted into the RO form upon the formation of memory cells [13]. This implies

that naïve T cells could be more trogocytic than mature memory T cells in RCC.

To compare the percentage of tumor cells displaying signs of trogocytosis, we

developed an image quantification program that records the percentage of individual cells'

surface area occupied by a trogocytic marker based on IF images (Supplemental Figure

1B). Cutoffs were established to define a tumor cell as trogo$^+$ or trogo$^-$ based on what

estimated percentage of their surface area was positive for a given immune marker.

Through this analysis, we found that of the 21 patient tumor images, all displayed

significant signs of trogocytosis between immune cells and tumor cells (Figure 1 B,C).

The average percentage of trogo$^+$ RCC cells per tumor was between 15-20%, with some

patient tumors as high as 50% trogo$^+$ for certain markers.

Characterization of Trogocytic RCC Tumors by Flow Cytometry

Next, we further characterized fresh human RCC tumors for markers of trogocytosis via flow cytometry. The IF analysis previously described has certain limitations that make distinguishing trogocytic staining from background signal difficult. Flow cytometry was chosen to provide a more accurate depiction of the trogocytic properties of RCC tumor cells. Fresh tumor samples were acquired from the University of Arizona biobanking facility and immediately dissociated and stained for markers of interest, namely CAIX, CD45, CD56, CD16 and CD14. Samples were evaluated within 12 hours of receiving them to ensure maximum viability of the cells.

Tumor cells were discriminated from infiltrating lymphocytes and non-tumor tissue gating on CAIX^{hi} cells. Size discrimination was also used to determine that the CAIX^{hi} population was independent from lymphocyte populations present in the tumor (Figure 2C). After establishing the tumor cells as distinct from all other cell types present in the tissue, we further characterized both the tumor and lymphocyte populations using a panel of trogocytic markers (CD14, CD16, CD56, and CD45). Analysis showed significant levels of expression of all four markers by fresh human ccRCC tumor cells (Figure 2C,D, supplementary figures 3A,4).

CD45 is an established marker for all nucleated cells of hematopoietic lineage[14]. CD45 labeling revealed widespread expression on the surface of the CAIX^{hi} (tumor) population (Figure 2C,D). Our data suggest that the majority of tumor cells in RCC tumors have expression of immune markers, and that this property is detectable by flow cytometry.

155  Given the high percentage of tumor cells exhibiting trogocytic markers, we then

156 decided to focus on characterizing cells that express multiple immune proteins. Our data

157 revealed that the vast majority of CAIX^{hi} cells were positive for multiple immune cell

158 markers (Figure 2D). Therefore, it is possible that RCC tumor cells undergo multiple

159 trogocytic events that allow them to "steal" the ability to express these markers from

160 multiple types of immune cells. The exact identity of these immune cells is not yet fully

161 elucidated and is an important field of study for future investigations.

162  To demonstrate the transfer of CD45 transfer from a lymphocyte to a tumor cell in

163 vitro we performed several coculture assays between RCC cell lines and Jurkat T cells or

164 primary CD3$^+$ T cells isolated from healthy donor PBMC. Cell lines were cocultured for 16

165 hours and then CD45 transfer was assessed via flow cytometry. Our results reveal that

166 there was a significant increase in CD45 labeling on RCC cells that had been cocultured

167 with primary T cells compared to RCC cells in monoculture (Figure 2A,B). We also

168 observed a significant increase in CD45 expression by RCC cell in cocultures done with

169 Jurkat T cells (Supplementary Figure 2A-C). This transfer was inhibited when a transwell

170 barrier was placed between the two cell types in coculture, indicating that physical contact

171 is necessary (Supplemental Figure 2D-E).

172 *RCC Tumor cells Express RNA for Immune Surface Proteins*

173  Given the abundance of immune antigens being expressed by RCC tumor cells,

174 we decided to identify the source of this protein expression. Upwards of 98% of tumor

175 cells in human RCC tumors were positive for one or more trogocytic protein when

176 characterized by flow cytometry (Figure 2C). The current definition of trogocytosis

177 describes the transfer of membrane fragments from one cell to another, however it

seemed unlikely that simple membrane transfer from lymphocytes to tumor cells could

account for the relatively high quantities of trogocytic protein present on the surface of

RCC cells. Therefore, we sought to identify other possible sources of this protein

expression by tumor cells. More specifically, we attempted to determine whether tumor

cells were capable of self-sufficient transcription of immune cell-specific proteins after

trogocytosis occurred. To investigate this, we cocultured Jurkat T cells with four kidney

carcinoma cell lines. We performed fluorescence activated cell sorting (FACS) to isolate

RCC cells from the coculture prior to harvesting RNA to ensure there was no

contamination of T cells in the cells being analyzed. Using qRT-PCR analysis, we

identified a significant induction of CD45 gene expression in RCC cells post coculture

(Figure 3C).

Next, we isolated trogocytic (T, $CAIX^{hi}CD45^{hi}$), non-trogocytic (NT,

$CAIX^{hi}CD45^{low}$), and lymphocyte (L, $CAIX^{low}CD45^{hi}$) populations from human RCC tumor

samples via FACS sterile sorting (Figure 3A). We then ran an nCounter® PanCancer

Immune Profiling analysis on the sorted cells using Nanostring technology to identify key

transcriptional differences between T and NT cells. Our analysis showed that trogocytic

tumor cells exhibit a "fusion" phenotype, defined as expressing both tumor and immune-

specific genes (Figure 3B). There was significantly higher expression of the trogocytic

markers CD45, CD14, and CD16 in trogocytic tumor cell populations compared to non-

trogocytic tumor cell populations (Figure 3D). We also identified new potential trogocytic

markers such as CD4 and CD53 on the trogocytic tumor samples (Figure 3B,

Supplementary Table 1).

RCC Cell Lines Acquire Genomic DNA from Primary T Cells

In order to determine the mechanism by which RCC tumor cells gained immune protein

expression, we investigated the possibility of horizontal gene transfer (HGT) occurring

during trogocytosis. Primary human T cells isolated from a healthy donor were incubated

with EdU, a thymidine analog that is readily incorporated into the DNA of dividing cells

[15]. EdU-labeled T cells were then cocultured overnight with RCC cell lines (ACHN,

A498, Caki-1, and 786O cells). Following coculture, the cells were then analyzed via flow

cytometry to determine whether we could detect a transfer of EdU-labeled DNA from the

T cells to the RCC cell lines. Results showed that between 1-8% of cocultured RCC cells

contained EdU labeled DNA (Figure 4C). In contrast, monocultured RCC cell lines and

cocultures done with a transwell control separating the T cells from the RCC cells did not

show any evidence of DNA transfer.

Additionally, we generated a primary T cell line expressing GFP-tagged histone H2B as

a secondary method of tracking DNA transfer. GFP-H2B T cells were cocultured overnight

with Caki-1 cells and then analyzed via flow cytometry. Analysis showed that there was

a small population of GFP-positive Caki-1 cells that was not present in monocultured

Caki-1 cells (Figure 4A).

Finally, we performed super-resolution structured illumination microscopy to gain insight

into the physical properties of the transferred material. We discovered CD45[+] intracellular

bodies present in RCC cells post coculture with primary human T cells (Fig. 4D,E).

Additionally, some of these "trogosomes" contained EdU labeled DNA (Fig. 4D,E). The

presence of these trogosomes could be indicative of a mechanism by which lymphocyte

DNA is transferred through CD45[+] lymphocyte membrane bubbles into the intracellular

space of RCC cells.

DISCUSSION

In this study, we showed that RCC tumor cells express significant levels of lymphoid proteins, such as CD45RA, CD14, CD16, and CD56. We have demonstrated that this expression is obtained through trogocytosis and subsequent horizontal gene transfer occurring between a tumor cell and lymphocyte. Importantly, this phenomenon is widespread, with over 90% of fresh RCC tumor cells showing high levels of trogocytic protein expression when analyzed via flow cytometry. Additionally, RNA extraction and analysis showed that tumor cells exhibit an altered transcriptome after trogocytosis occurs. Namely, we observed an increase in gene expression for genes that are typically only expressed by cells of hematopoietic lineage. Finally, we demonstrated that genomic DNA can be transferred during trogocytosis from a lymphocyte to a tumor cell resulting in an altered tumor cell phenotype. Taken together, this data demonstrates that RCC tumor cells acquire the ability to express large quantities of hematopoietic surface antigens following a trogocytic event with infiltrating lymphocytes.

Using immunofluorescent analysis of RCC slides, we developed a computer algorithm to quantify the percentage of tumor cells that express proteins found on lymphocytes. This study revealed consistent expression of the markers CD14, CD16, CD56, and CD45RA in a cohort of 21 human patient tumors when analyzed via immunofluorescence microscopy (IF). Despite limitations in the sensitivity of this method, analyzing 21 tumors originating from different stages of disease revealed no significant difference in trogocytic antigen expression at any stage of tumor development. Additionally, of the four trogocytic antigens we analyzed, there were no significant changes in the specific type of antigen being expressed at any stage. These data imply that trogocytosis occurs at an early stage

during tumor development, and that the trogocytic phenotype of RCC tumor cells are maintained as disease progresses. Currently, it is unknown whether this is due to repeated trogocytic interactions with infiltrating lymphocytes, acquisition of lymphocytic DNA that promotes expression of these antigens, or some combination of the two.

Analysis of fresh tumors by flow cytometry corroborated the results we obtained via IF. Flow cytometry revealed much higher levels of trogocytic marker expression than previously anticipated, with over 90% of tumor cells in all RCC tumors tested being positive for at least one lymphocytic marker. To our knowledge, this is the first time these lymphocytic proteins have been identified on any solid tumor and could shift the paradigm of how we understand a tumor's interactions with its microenvironment.

Our investigation into horizontal gene transfer between RCC cells and T cells has shown that it is possible for tumor cells to acquire DNA from lymphocytes that they encounter. We have shown that the transfer of both EdU labeled DNA and GFP-tagged histones is observed after coculturing tumor cells and immune cells *in vitro*. Currently, it is unknown whether this transfer confers greater survival capabilities or a more aggressive phenotype to tumor cells that have undergone horizontal gene transfer. Additional studies are needed to fully characterize this process and its role in disease progression. Our data also suggests that membrane transfer can occur independently of horizontal gene transfer, implying that there are multiple mechanisms that regulate these processes.

Trogocytosis has been shown to promote the acquisition of the immune check point receptor PD-1 by natural killer cells in mouse models of leukemia, resulting in suppressed NK cells antitumor immunity [16]. Additionally, colorectal cancer cells have

been shown to obtain other immune regulatory molecules such as CTLA4 through

trogocytosis with infiltrating lymphocytes [6]. These results suggest that trogocytosis is a

mechanism by which tumors evade elimination by the immune system, either by acquiring

regulatory molecules themselves, or by indirectly benefiting from certain immune cells

negatively regulating themselves. In the context of RCC, it is unknown how tumor cells

use the acquisition of immune cell surface protein and DNA to promote their survival.

However, given the frequent expression of these markers on RCC tumors from all stages

of disease, we believe trogocytosis confers a survival advantage to tumor cells. More

research is needed to determine whether this survival advantage is selected for by the

tumor to evade the immune system, or some other unknown mechanism.

It is possible that trogocytosis and the effect it has on the phenotype of tumor cells

may be exploited to improve cancer therapies. The expression of lymphoid antigens on

cancer cells may constitute effective targets of antibody drug conjugates (ADCs). Further

profiling of the markers that are transferred during this process may reveal unique protein

expression signatures by trogocytic tumors that are not otherwise found in healthy cells.

In addition to their use as new drug targets, these markers can also potentially be used

as novel biomarkers to help better inform treatment strategies for clear cell renal

carcinoma patients. Further research in the field of trogocytosis is critical for developing

the next generation of immunotherapies.

ACKNOWLEDGEMENTS

We acknowledge use of the University of Nebraska Medical Center - UNMC Advanced

Microscopy Core Facility, RRID:SCR_022467, P20 GM103427 (NIGMS, NE-INEBRE),

P30 GM106397 (NIGMS, NCS), P20GM130447 (NIGMS, CoNDA), P30 CA036727 (NCI,

Buffett Cancer Center), S10RR02730 (NIH), S10OD030486 (NIH), Nebraska Research

Initiative, UNMC Vice Chancellor for Research Office. Research reported in this

publication was supported by the National Cancer Institute of the National Institutes of

Health under award number P30 CA023074. The UNMC Flow Cytometry Research

Facility is administrated through the Office of the Vice Chancellor for Research and

supported by state funds from the Nebraska Research Initiative (NRI) and The Fred and

Pamela Buffett Cancer Center's National Cancer Institute Cancer Support Grant (P30

CA036727). Major instrumentation has been provided by the Office of the Vice

Chancellor for Research, The University of Nebraska Foundation, the Nebraska Banker's

Fund, and by the NIH-NCRR Shared Instrument Program. Special thanks to J.C. Fitch for

assistance with flow cytometry analysis. Studies were supported by funding from the

National Cancer Institute (T32CA009213-44) to A. Sivakoses and 7R01AI137060-06

awarded to A.L.M. Bothwell.

USE OF HUMAN MATERIALS

All human materials were acquired through the University of Arizona Tissue Acquisition

and Cellular/Molecular Analysis Shared Resource. Patient materials were obtained and

processed in accordance with University of Arizona tissue acquisition policy.

DATA AVAILABILITY

Source code for immunofluorescent analysis available for use at bit.Ly/TrogoTracker.

MATERIALS AND METHODS

Immunofluorescent staining assays

The following primary antibodies were used in immunofluorescent experiments: anti-human CAIX (R&D Biosystems, AF2188), CD45RA (Sino Biological Inc., 102580-T08), CD56 (ProteinTech®, 14255-1-AP), CD68 (Sino Biological Inc., 11192-T26), CD16 (ProteinTech®, 16559-1-AP), CD14 (R&D Biosystems, BAF383). Paraffin-embedded tissue slides were obtained from Tissue Acquisition and Cellular/Molecular Analysis (TACMASR) core facility at the University of Arizona. Slides were deparaffinized for 9min in Xylenes. They were then washed in decreasing concentrations of ethanol solutions at 1min intervals for 5min. The slides were then washed in MilliQH$_2$O for 10 minutes. Then, antigen retrieval was performed using boiling Antigen Unmasking Solution (Vector Laboratories, H-3300) for 10min and left to cool for an additional 20min. Slides were blocked with 20% donkey serum/1X DPBS-0.1% Tween-20 for 30min. Primary antibody solutions were prepared according to their recommended dilutions. Slides were incubated with primary antibodies overnight at 4°C and washed with 1X DPBS-0.1% Tween-20 for 5min. Samples were quenched using Invitrogen ReadyProbes™ Tissue Autofluorescence Quenching Kit for 3-5min. Secondary anti-donkey antibodies from Invitrogen were applied and incubated for 1 hour at room temperature in the dark. Samples were quenched an additional time and stained with Hoescht nuclear stain for

10min. Coverslips were mounted on the stained slides using Fluoromount-G™

(Invitrogen, 00-4958-02). Slides were imaged using the Echo Inc. Revolution fluorescent

microscope.

Flow cytometry assays

The following conjugated antibodies were used in flow cytometric experiments: CD45

(clone 2D1, BioLegend, 368532), CD16 (clone CB16, Invitrogen, 17-0168-42), CD14

(clone 61D3, Invitrogen, 414-0149-42), CD45RO (clone UCHL1, Invitrogen, 12-0457-42),

CD45RA (clone HI100, Invitrogen, 17-0458-42), CD44 (clone IM7, Invitrogen, 404-0441-

82), CAIX (R&D Biosystems, FAB2188G), CD56(clone CMSSB, Invitrogen, 12-0567-42),

and CD68 (clone REA886, Miltenyi Biotec, 130-114-463). Fresh tumor tissue was

acquired through the Tissue Acquisition and Cellular/Molecular Analysis (TACMASR)

core facility at the University of Arizona. Tumor samples were finely minced into 1mm^2

sections and washed with DPBS. They were then suspended in a solution of 1mg/ml

Collagenase Type 4 (Worthington, LS0004186) and incubated at 37°C for 30min, being

shaken every 3min. The reaction was stopped using DMEM media with 5% fetal bovine

serum. Cell suspensions were then filtered through a 40μm mesh filter and resuspended

in 100μl of DPBS. Cell viability was stained with Ghost Dye™ Violet 450 (Tonbo

Biosciences, 13-0863-T100). Human BD Fc Block™ (BD Pharmingen™, 564220) was

used at a concentration of 5μl/10^6 cells for 10min at room temperature. Fluorescent

antibody staining was done at the recommended concentrations of each antibody for

30min at room temperature in the dark. Cell suspensions were then washed using DPBS

and resuspended in 1X RBC lyse/fix solution (eBioscience™ 00-5333-57) for 40min.

Samples were then washed in DPBS and analyzed using the BD FACSCanto™ II system.

Data was analyzed using FlowJo v10.8.1.

Immunofluorescent image quantification

Images used for analysis were taken at 40X (NA 1.40) on an Echo Revolution LED

microscope. Images were exported in separate channels into ImageJ (FIJI) for

quantification. Nuclei were identified using the StarDist2D plugin [17], then a Voronoi

algorithm was applied to estimate the boundaries of individual cells. The tumor marker

channel was used to subtract all non-tumor cells from the Voronoi cell map. The resulting

regions of interest (ROIs) were then saved to be used as a map for all tumor cells in each

image. The channel containing the trogocytic marker of interest was then run through a

background subtraction algorithm and the resulting signal was normalized using the

CLAHE histogram normalization plugin. The channel was then converted to a binary

image and the ROIs previously described were applied. The total surface area of each

cell was measured to give a two-dimensional representation of the cellular surface area

being occupied by a trogocytic marker. Cells with trogocytic markers occupying over 55%

and less than 90% of their total 2D surface area were considered trogocytic. These cutoffs

were determined to exclude autofluorescence, other sources of background signal, and

infiltrating immune cells from quantification. We believe this method of quantification is

quite conservative and may exclude low expressing trogocytic tumor cells, implying that

the true percentage of tumor cells that are trogocytic may be much higher than this

analysis estimates. This could account for the difference between the IF quantification

and flow cytometry results.

Cell Culture

Human primary T cells were obtained from the Elutriation Core Facility at the University

of Nebraska Medical Center. T cell populations were isolated using the Miltenyi Biotec

Pan T Cell Isolation Kit (130-096-535). T cells were cultured in X-VIVO™ 15 Serum-Free

Hematopoietic Cell Medium (Lonza, 02-053Q) supplemented with 20ng/mL recombinant

human IL-2 (R&D, BT-002-010) once a week and 7ul/mL ImmunoCult™ Human

CD3/CD28/CD2 T Cell Activator (Stem Cell Technologies, 10970) every other week. RCC

cell lines (Caki-1, ACHN, A498, 786O) were obtained from Dr. George Sutphin's

laboratory at the University of Arizona. RCC cell lines were cultured based on the

recommendations made by ATCC.

Lentiviral Transduction of Primary T Cells

GFP-H2B lentiviral vector was obtained from Dr. Ghassan Mouneimne at the University

of Arizona. Lentivirus was generated in 239T cells, and the viral supernatant was

harvested for primary T cell infection. T cells were spun for 30 minutes at 3000 rpm with

the virus. Lentivirus was removed the following day after incubation and GFP+ T cells

were isolated using FACS selection 2 weeks after transduction.

Coculture Assays

RCC lines (ACHN, A498, Caki-1, 786-O) were plated a minimum of 4 hours prior to

coculture. T cells were added to the culture at a 1:1 ratio and incubated overnight.

Cocultures were harvested for experimental purposes 16 hours post addition of the T

cells.

Confocal Microscopy of EdU Transfer

Images were taken on a Zeiss 710 Laser Scanning Confocal microscope at 40X (NA = 1.30). Images shown are max intensity projections of 60 Z-stacked images. FIJI smoothing algorithm was applied to phalloidin imaging for the purpose of minimizing the quality loss of actin labeling as a result of the EdU staining process.

REFERNCES

1. Joly, E., *What is trogocytosis and what is its purpose?* Nature Immunology, 2003. **4**(9): p. 815.
2. Arnold, P.Y., D.K. Davidian, and M.D. Mannie, *Antigen presentation by T cells: T cell receptor ligation promotes antigen acquisition from professional antigen-presenting cells.* European Journal of Immunology, 1997. **27**(12): p. 3198-3205.
3. Osborne, D.G. and S.A. Wetzel, *Trogocytosis results in sustained intracellular signaling in CD4(+) T cells.* J Immunol, 2012. **189**(10): p. 4728-39.
4. Nakayama, M., et al., *Natural killer (NK)-dendritic cell interactions generate MHC class II-dressed NK cells that regulate CD4+ T cells.* Proc Natl Acad Sci U S A, 2011. **108**(45): p. 18360-5.
5. Sarvari, A.K., et al., *Interaction of differentiated human adipocytes with macrophages leads to trogocytosis and selective IL-6 secretion.* Cell Death Dis, 2015. **6**(1): p. e1613.
6. Shin, J.H., et al., *Colon cancer cells acquire immune regulatory molecules from tumor-infiltrating lymphocytes by trogocytosis.* Proc Natl Acad Sci U S A, 2021. **118**(48).
7. Zeng, Q. and H. Schwarz, *The role of trogocytosis in immune surveillance of Hodgkin lymphoma.* Oncoimmunology, 2020. **9**(1): p. 1781334.
8. Park, H.R., et al., *Blockade of CD47 enhances the antitumor effect of macrophages in renal cell carcinoma through trogocytosis.* Sci Rep, 2022. **12**(1): p. 12546.
9. Hamieh, M., et al., *CAR T cell trogocytosis and cooperative killing regulate tumour antigen escape.* Nature, 2019. **568**(7750): p. 112-116.
10. Smillie, C.S., et al., *Ecology drives a global network of gene exchange connecting the human microbiome.* Nature, 2011. **480**(7376): p. 241-4.
11. Xu, Q., et al., *Cell type-specific intercellular gene transfer in mammalian cells via transient cell entrapment.* Cell Discov, 2022. **8**(1): p. 20.
12. Courcier, J., et al., *Carbonic Anhydrase IX in Renal Cell Carcinoma, Implications for Disease Management.* Int J Mol Sci, 2020. **21**(19).
13. Prince, H.E., *Phenotypic comparison of the three populations of human lymphocytes defined by CD45RO and CD45RA expression.* Cellular Immunology, 1992. **145**(2): p. 254-262.
14. Woodford-Thomas, T., *The leukocyte common antigen, CD45 and other protein tyrosine phosphatases in hematopoietic cells.* Cell Biology 1993. **4**: p. 409-418.
15. *<salic-mitchison-2008-a-chemical-method-for-fast-and-sensitive-detection-of-dna-synthesis-in-vivo.pdf>.*

440   16.   Hasim, M.S., *When killers become thieves: Trogocytosed PD-1 inhibits NK cells in cancer.* Science
441   Advances, 2022. **8**(15).
442   17.   Schmidt, U., et al., *Cell Detection with Star-Convex Polygons*, in *Medical Image Computing and*
443   *Computer Assisted Intervention – MICCAI 2018*. 2018. p. 265-273.
444

FIGURE 3

Figure 3. RNA expression analysis of immune cell-specific genes by trogocytic RCC cells in human ccRCC tumors. **(A)** Representative gating strategy used to identify lymphocyte (L), trogocytic tumor (T), and non-trogocytic tumor (NT) cell populations in human ccRCC tumors when sorting cells via FACS. L cells were identified as CD45high CAIX^{low}, trogocytic tumor cells were identified as CD45^{high} CAIX^{high}, and non-trogocytic tumor cells were identified as CD45^{low} CAIX^{high}. **(B)** NanoString® gene expression analysis for human ccRCC tumor. Populations were sorted from fresh human ccRCC tumors via FACS according to the gating strategy described in Figure 3A. Data shows Log2 RNA expression counts for select immune and cancer specific genes. **(C)** qRT-PCR analysis

of the change in CD45 expression in RCC cell lines post coculture with Jurkat T cells. Cells were sorted via FACS prior to analysis to isolate RCC cells. Graph depicts Log2 CD45 gene expression fold change in cocultured cells relative to monoculture. (Mean Ct values (n=4): Caki-1 coculture 28.297, monoculture 31.174, ACHN coculture 29.131, monoculture 33.575, 786O coculture 28.501, monoculture 36.759, A498 coculture 29.071, monoculture 33.385). (D) Comparison of Log2 gene expression counts of trogocytosis markers between trogocytic and non-trogocytic tumor cell populations isolated from human RCC tumors. Data was analyzed using multiple T tests (p value < 0.01 = **, p value < 0.001 = ***).

FIGURE 4

Figure 4. DNA is transferred from primary T cells to RCC cell lines. **(A)** Flow cytometry

results of Caki-1 cells that were cocultured with GFP-histone H2B primary T cells. Data

shows the percentage of GFP-H2B⁺ Caki-1 cells in coculture relative to monocultured

Caki-1 cells. **(B)** Mean fluorescent intensity of GFP-H2B expression of cocultured Caki-1 cells compared to monoculture (p value = 0.0058, unpaired T test was used to analyze the data). **(C)** Flow cytometry analysis of RCC cell lines cocultured with EdU-labeled primary human T cells. Plots show the percentage of RCC cells that are EdU$^+$ and CD45$^+$ post 16-hour coculture. **(D)** Image taken from a 10:1 coculture of EdU-labeled primary human T cells with ACHN cells. Left panels depict an enhanced view of an EdU$^+$ CD45$^+$ trogosome within the intracellular space of an ACHN cell. Right panels depict an enhanced view of an EdU$^+$ CD45$^+$ T cell for comparison purposes. **(E)** Representative images of trogosomes in three RCC cell lines (top to bottom, A498, 786O, ACHN). Boxes in the merged images of middle and bottom panels indicate the region being enhanced in the split color panels to the right of the merged image. Arrows in the merged zoom panels indicate the location of a trogosome. All images taken on a Zeiss 710 laser scanning confocal, 40X (NA 1.30).

Figure 1. Trogocytic RCC tumor cells are detected and quantified using immunofluorescent staining. **(A)** Immunofluorescent microscopy of human RCC tumors. Trogocytic markers (CD14, CD56, CD45RA, CD16), CAIX, and Hoechst are displayed as green, red, and blue respectively. Arrows indicate trogocytic tumor cells. **(B)**

Quantification of average percentage of trogocytic tumor cells for 21 human RCC tumors.

(C) Quantification of average percentage of trogocytic tumor cells in 21 human RCC tumors organized by tumor stage. Methodology of quantification described in Methods and Materials.

Figure 2. RCC cells acquire expression of immune surface proteins through contact with lymphocytes. **(A)** Flow cytometry analysis of CD45 expression by RCC cell lines post coculture with primary human T cells. RCC cells were cocultured with T cells for 16 hours prior to analysis. Data shows an increase in CD45 expression relative to RCC cells cultured in isolation (monoculture). **(B)** Mean fluorescent intensity of CD45 expression by cocultured RCC cells relative to monoculture. **(C)** Representative flow cytometry analysis of fresh human ccRCC tumors. Tumor cells were identified as CAIX^{hi} and lymphocytes and other cells present in the tumor microenvironment were identified as CAIX^{low}. Data shows expression of four immune cell surface proteins (CD14, CD16, CD56, CD45) by

ccRCC and non-ccRCC cells isolated from this tumor. **(D)** Percentage of tumor cells from 4 human ccRCC tumors that expressed CD14, CD16, CD56, and CD45 based on flow cytometry analysis. Tumor cells were identified via CAIX^{hi} labeling as represented in Figure 2C.