1
2 **Engineering the plant intracellular immune receptor Sr50 to restore recognition of the AvrSr50 escape mutant**
3

4   Kyungyong Seong[1], Wei Wei[1], Brandon Vega[1], Amanda Dee[1], Griselda Ramirez-Bernardino[1], Rakesh Kumar[1],
5   Lorena Parra[1] and Ksenia Krasileva[1,2,*]
6

7   [1] Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA.

8   [2] Innovative Genomics Institute, University of California, Berkeley, CA 94704

9   *To whom correspondence should be addressed: kseniak@berkeley.edu

10
11 **Abstract**

12
Sr50, an intracellular nucleotide-binding leucine-rich repeat receptor (NLR), confers resistance of wheat against stem
rust caused by the fungal pathogen *Puccinia graminis* f. sp. *tritici*. The receptor recognizes the pathogen effector
AvrSr50 through its C-terminal leucine-rich repeat domain, initiating a localized cell death immune response.
However, this immunity is compromised by mutations in the effector, as in the escape mutant AvrSr50^{QCMJC}, which
evades Sr50 detection. In this study, we employed iterative computational structural analyses and site-directed
mutagenesis for rational engineering of Sr50 to gain recognition of AvrSr50^{QCMJC}. Following an initial structural
hypothesis driven by molecular docking, we identified the Sr50^{K711D} single mutant, which induces an intermediate
immune response against AvrSr50^{QCMJC} without losing recognition against AvrSr50. Increasing gene expression with
a stronger promoter enabled the mutant to elicit a robust response, indicating weak effector recognition can be
complemented by enhanced receptor expression. Further structural refinements led to the creation of five double
mutants and two triple mutants with dual recognition of AvrSr50 and AvrSr50^{QCMJC} with greater immune response
intensities than Sr50^{K711D} against the escape mutant. All effective mutations against AvrSr50^{QCMJC} required the K711D
substitution, indicating that multiple solutions exist for gain of recognition, but the path to reach these mutations may
be confined. Furthermore, this single substitution alters the prediction of AlphaFold 2, allowing it to model the
complex structure of Sr50^{K711D} and AvrSr50 that match our final structural hypothesis. Collectively, our study outlines
a framework for rational engineering of NLR systems to overcome pathogen escape mutations and provides datasets
for future computational models for NLR resurrection.
30
31

Introduction

Plant intracellular immunity relies on nucleotide-binding leucine-rich repeat receptors (NLRs) that monitor and detect
pathogen activities within cells (Jones and Dangl 2006). Upon direct recognition of pathogen effector molecules,
NLRs trigger immune responses, typically culminating in localized cell death known as a hypersensitive response
(HR) (Dodds and Rathjen 2010). The ability of NLRs to impede pathogen proliferation has driven researchers and
breeders to identify functional NLRs and deploy them in genetic protection strategies for important crops (Dangl et
al. 2013; Arora et al. 2019). Notably, Sr50, originally identified in rye, has been effectively utilized to protect wheat
(*Triticum aestivum*) against stem rust disease caused by the fungal pathogen *Puccinia graminis* f. sp. *tritici* by directly
recognizing its cognate effector AvrSr50 (Mago et al. 2015).

Pathogens counteract plant defenses through evolution of their effectors, enabling them to evade NLR-mediated
immune responses (Möller and Stukenbrock 2017; Sánchez-Vallet et al. 2018). For example, the *P. graminis* f. sp.
tritici isolate QCMJC secretes a variant of AvrSr50 (hereafter referred to as AvrSr50^{QCMJC}) that escapes Sr50
recognition with a single substitution, Q121K, on the protein surface (Chen et al. 2017; Ortiz et al. 2022). This escape
mutant compromises plant immunity and subsequently poses threats to genetically uniform crops. However, restoring
the effectiveness of NLRs remains a significant challenge, as the mutational landscape to be explored is extensive
(Tamborski and Krasileva 2020; Zdrzałek et al. 2023).

Key structural biology techniques, such as cryogenic electron microscopy (Cryo-EM) and crystallography, have been
instrumental in deciphering the complex binding mechanisms between NLRs and effectors, guiding subsequent
bioengineering efforts (Wang et al. 2019a, 2019b; Ma et al. 2020; Martin et al. 2020; Cesari et al. 2022; Förderer et
al. 2022; Zhao et al. 2022; Contreras et al. 2023; Liu et al. 2023; Maidment et al. 2023; Selvaraj et al. 2023; Kourelis
et al. 2024; Lawson et al. 2024; Madhuprakash et al. 2024). Yet, dependency on specialized expertise, experimental
challenges in protein purification and extensive resources limit applicability of crystallography and Cryo-EM
techniques to the full spectrum of NLR-effector pairs. Computational structure prediction has emerged as a potential
alternative to address this issue. AlphaFold 2 (AF2) and 3 (AF3) have been used to understand NLR oligomerization
and to elucidate the interaction between MLA3, an NLR from barely, and its effector Pwl2 (Evans et al. 2021; Jumper
et al. 2021; Abramson et al. 2024; Gómez De La Cruz et al. 2024, 2024; Madhuprakash et al. 2024). However, their
low accuracy was challenged by a recently solved Cryo-EM structure of MLA13 and its effector Avr$_{A13}$-1 (Lawson et
al. 2024). In such circumstances where molecular interactions between most NLRs and effectors cannot be fully
provided, engineering NLRs requires development of new rational engineering approaches that can be successful with
incomplete information.

Despite the inherent difficulties, significant strides have been made in NLR engineering. Advancements include
engineering small integrated domains (IDs) found in a subset of NLRs as platforms for effector binding (Kroj et al.
2016; Sarris et al. 2016; Baggs et al. 2017). Particularly, rational design strategies focusing on heavy-metal associated
domains (HMAs), guided by crystallography structures, have shown promise by altering or transferring effector
recognition specificity (De La Concepcion et al. 2019; Cesari et al. 2022; Bentham et al. 2023; Maidment et al. 2023).
Conversely, engineering the C-terminal leucine-rich repeat (LRR) domain, postulated to participate in effector binding
across most NLRs, remains a challenging pursuit. Most studies resort to gain-of-function random mutagenesis to
counter escape effector mutants (Farnham and Baulcombe 2006; Harris et al. 2013; Segretin et al. 2014; Huang et al.
2021). In our recent endeavor, we harnessed the natural sequence diversity of NLRs, pinpointing and targeting rapidly
evolving, highly variable (hv) residues to switch recognition specificity between two related NLRs (Prigozhin and
Krasileva 2021; Tamborski et al. 2023). Nevertheless, receptor-centric random mutagenesis and engineering
approaches do not consider how the effectors are bound to the receptors and overlook the residual interplay between
them. When effectors mutate, therefore, numerous receptor residues need to be re-screened to resurrect the NLRs.
Consequently, there is a pressing need for methodologies that can illuminate the conformation of NLR and effector as
a part of the rational LRR design.

In this study, we engineer Sr50 to recognize the escape mutant AvrSr50^{QCMJC}. We employ a strategic combination of
site-directed mutagenesis, molecular docking and structural analyses to iteratively infer and refine the heterodimeric
structure of Sr50 and AvrSr50. Guided by predictive models through molecular docking simulations, we introduced
mutations to charged or polar residues of Sr50 and AvrSr50, which likely govern ligand-receptor specificity, to disrupt
effector recognition or mediate recognition escape, respectively (Fig. 1A). We then tested their complementary
AvrSr50 and Sr50 mutants in *Nicotiana benthamiana* to experimentally corroborate residual proximity and leveraged
it to refine our structural hypotheses with ColabDock (Feng et al. 2023). In this iterative process, we generated single,
double, and triple mutants of Sr50 that induced variable levels of cell death against AvrSr50^{QCMJC}. Interestingly, all

successful mutation combinations required the K711D substitution to result in gain of AvrSr50[QCMJC] recognition.
Furthermore, although AF2-Multimer could not predict the structure of Sr50 and AvrSr50, it could model the complex
structure of Sr50[K711D] bound to AvrSr50 and AvrSr50[QCMJC]. This predicted AF2 model showed good agreement with
our final structural hypothesis, indicating that the outcomes from different approaches converged. Collectively, our
exploration extended beyond engineering solutions, providing not only valuable insights of the Sr50 and AvrSr50
interaction, but also training datasets and rational design strategies that can be expanded to other systems.

91
Figure 1. Initial structural hypotheses derived from molecular docking simulations and gain of AvrSr50[QCMJC]
recognition

A. A generalized workflow of structure-guided NLR engineering. **B.** A predicted pose of Sr50 and AvrSr50 by molecular docking
simulations. Some parts of a loop between β2–β3, mostly including an unstructured region, are removed from AvrSr50 for
visualization (positions 42-66). The local environment around AvrSr50's Q121 is visualized. **C**. The distribution of selected amino
acids across the surface of Sr50's leucine-rich repeat (LRR) domain. **D** and **E**. The qualitative and quantitative cell death phenotypes

on *Nicotiana benthamiana*. The optical density (OD_{600}) was set to 0.3 for receptors and effectors, respectively, and the receptors were expressed under pRPP13. Indicated receptor and effector pairs were co-infiltrated. The cell death phenotypes were recorded at three days post infiltration (dpi). **E**. The mode intensity of the infiltrated spots was measured with imageJ across the given number of biological replicates in bold. The statistics were calculated with one-way ANOVA followed by a post-hoc Tukey HSD (Honestly Significant Difference) test. The compact letters indicate significant differences between groups. **F**. Representative Western blot for the wild type Sr50 and the Sr50^{K711D} mutant under pRPP13 or p35S. **G** and **H**. The qualitative and quantitative cell death phenotypes on *N. benthamiana*. The OD_{600} was set to 0.1 for receptors and 0.3 for effectors, respectively, and the receptors were expressed under p35S. The cell death phenotypes were recorded at 2 dpi. **H**. The mode intensity of the infiltrated spots was measured with imageJ across the given number of biological replicates in bold. The statistics were calculated with one-way ANOVA followed by a post-hoc Tukey HSD test.

Results

Initial structural hypothesis generation with molecular docking

To formulate our initial structural hypothesis, we attempted modeling the heterodimeric complex of Sr50 and AvrSr50 with AF2-Multimer and AF3. However, the reported accuracy was low, indicating uncertainty in the prediction (Fig. S1). We alternatively turned to molecular docking algorithms to generate a set of initial structural hypotheses based on specific criteria drawn from the Sr35 resistosome structure and the assumption that AvrSr50's Q121K directly participates in binding through its interaction with Sr50's LRR residues (Förderer et al. 2022; Ortiz et al. 2022; Zhao et al. 2022). Our intention was to diversify the initial hypotheses and gather more experimental data to either support or disprove these models (Fig. 1A).

The molecular docking simulations produced three distinct models (Fig. 1B; Fig. S2). We prioritized two of these poses that aligned with our simplified assumption: the Q121K substitution in AvrSr50^{QCMJC} results in repulsive interactions with an amino acid side chain in Sr50, which impedes effector recognition. In the first model (Fig. 1B), AvrSr50's Q121 was proximal to two positively charged residues of Sr50, R688 on LRR 8 and K711 on LRR 9. An alternative model (Fig. S2A) positioned Q121 in the vicinity of K824 on LRR 13 and R904 on LRR 16. These four charged residues form a neighboring cluster of tryptophan residues across LRRs 11 to 13, which might offer affinity for effector binding (Fig. 1C). We hypothesized that the two selected regions may determine specificity for AvrSr50, and modifying one of these two interfaces could recover the interaction with AvrSr50^{QCMJC}.

Sr50^{K711D} induces cell death against AvrSr50^{QCMJC} without losing the interaction with AvrSr50

To test the initial hypotheses about incompatible side chain interactions between Sr50 and AvrSr50^{QCMJC}, we introduced aspartic and glutamic acid into R688, K711, K824 and R904 of Sr50. These specific substitutions were selected based on our simplified assumption to re-establish complementary interactions with Q121K of AvrSr50^{QCMJC} through ionic or hydrogen bonds. Subsequently, we infiltrated *N. benthamiana* with *Agrobacterium* carrying Sr50 single mutants and AvrSr50 or AvrSr50^{QCMJC}. The expression of the receptors was driven by a native promoter of NLR RPP13 to avoid autoactivity (Tamborski et al. 2023). The aspartic and glutamic acid substitutions at K824 and R904 led to the loss of HR against AvrSr50; however, none of these receptor mutants could mediate AvrSr50^{QCMJC}-dependent cell death (Fig. S3). The mutations on Sr50's R688 neither completely abolished HR against AvrSr50 nor led to the gain of AvrSr50^{QCMJC} recognition (Fig. S3). Sr50^{K711D} and Sr50^{K711E} did not compromise the receptor's ability to recognize AvrSr50 (Fig. 1D and 1E). While Sr50^{K711E} could not restore HR against AvrSr50^{QCMJC}, Sr50^{K711D} induced AvrSr50^{QCMJC}-dependent cell death. Compared to Sr50 and AvrSr50, the restored HR was weaker in intensity, likely indicating suboptimal interactions of Sr50^{K711D} towards AvrSr50^{QCMJC}. Nonetheless, these experiments suggested that Sr50^{K711D} gained recognition against AvrSr50^{QCMJC} without losing the interaction with AvrSr50.

Weak recognition of Sr50^{K711D} can be complemented by strong gene expression

To determine whether the increased HR of Sr50^{K711D} in response to AvrSr50^{QCMJC} was due to difference in protein
abundance, we estimated relative quantity of Sr50 and Sr50^{K711D} following their co-infiltration with AvrSr50 and
AvrSr50^{QCMJC}, respectively, using Western blot (Fig. 1F). Unexpectedly, we consistently observed lower protein levels
of Sr50^{K711D} compared to Sr50 under the RPP13 promoter. We postulated that assessment of the magnitude of cell
deaths may be confounded by the difference in protein abundance. To drive receptor abundance to comparable levels,
we instead used the constitutive 35S promoter from the Cauliflower Mosaic Virus.

Under the 35S promoter, the protein level of Sr50^{K711D} was slightly greater than Sr50 after co-infiltrations with
AvrSr50^{QCMJC} and AvrSr50, respectively (Fig. 1F). In consistency, the average mode intensity of HR appeared higher
for p35S::Sr50^{K711D} than p35S::Sr50 at two days post-infiltration , when the receptors were co-infiltrated with AvrSr50
(Fig. 1G and 1H). Notably, the HR induced by p35S::Sr50^{K711D} and AvrSr50^{QCMJC} was similar in magnitude to
p35S::Sr50 and AvrSr50 (Fig. 1G and 1H). This suggested that the Sr50^{K711D} mutant does recognize AvrSr50^{QCMJC}
and that potentially weak recognition of Sr50^{K711D} was complemented by strong expression driven by the 35S promoter
for enhanced immune responses. Although altering the promoter is a viable engineering strategy, we continued our
pursuit with the weaker RPP13 promoter to further enhance the recognition of Sr50^{K711D} through additional amino
acid mutations.

Sr50^{K711D} does not recognize the AvrSr50^{Q121K} single mutant

To generate Sr50 mutants capable of inducing robust HR against AvrSr50^{QCMJC} under the RPP13 promoter, we aimed
to refine our structural model to better guide targeted mutagenesis. The observed gain of function in Sr50^{K711D} was
not direct confirmation of the existing structural hypothesis (Fig. 1B). The mature forms of AvrSr50 and AvrSr50^{QCMJC}
have ten substitutions, including the Q121K substitution which has been shown to mediate recognition escape (Fig.
2A; Fig. S4) (Ortiz et al. 2022). It was therefore crucial to determine whether Sr50^{K711D} could induce HR against the
AvrSr50^{Q121K} single mutant. As reported previously (Ortiz et al. 2022), reverting K121 of AvrSr50^{QCMJC} to glutamine
as in AvrSr50 was sufficient to restore the HR for Sr50 and Sr50^{K711D} (Fig. 2B and 2C) However, co-infiltration of
Sr50^{K711D} and AvrSr50^{Q121K} did not lead to cell death. This possibly suggested that the interaction between Sr50's
K711D and AvrSr50's Q121K may not be one-to-one, and there might be other effector and receptor residues involved
in interaction.

Mutations in AvrSr50's N124 alter the interaction with Sr50 but may not be associated with Sr50's K711

The lack of HR between Sr50^{K711D} and AvrSr50^{Q121K} prompted a reassessment of other factors potentially influencing
the interaction. Notably, AvrSr50^{QCMJC} carries another adjacent substitution, N124T, along the terminal alpha helix
(Fig. 2A). We postulated that this adjacent mutation might contribute to the differential responses of Sr50^{K711D} towards
AvrSr50^{QCMJC} and AvrSr50^{Q121K}.

Additional experiments with the AvrSr50^{N124T} single mutant revealed that this substitution alone did not significantly
impact the interaction with Sr50 and Sr50^{K711D} (Fig. 2B and 2C). However, co-infiltration of Sr50^{K711D} and the
AvrSr50$^{Q121K/N124T}$ double mutant partially restored the cell death phenotype. Consistently, the presence of N124 in
AvrSr50$^{QCMJC\ K121Q}$ tended to attenuate the intensity of HR, compared to the AvrSr50$^{QCMJC\ K121Q}$ single mutant. This
suggested that the amino acid variations at position 124 of AvrSr50 alter the interaction with the receptor.

176
Figure 2. Mutations in AvrSr50's Q121 and N124 alter the interaction with Sr50

A. Variable positions between AvrSr50 and AvrSr50^{QCMJC}. Amino acid variations between the two proteins are indicated in the
predicted structure of AvrSr50 without its mature form. **B** and **C**. The qualitative and quantitative cell death phenotypes on
Nicotiana benthamiana. The optical density (OD_{600}) was set to 0.3 for receptors and effectors, respectively, and the receptors were
expressed under pRPP13. Indicated receptor and effector pairs were co-infiltrated. The cell death phenotypes were recorded at
three days post infiltration. **B**. The mode intensity of the infiltrated spots was measured with imageJ across the given number of
biological replicates in bold. The statistics were calculated with one-way ANOVA followed by a post-hoc Tukey Honestly
Significant Difference test. The compact letters indicate significant differences between groups.

To gain further mechanistic insights, we substituted AvrSr50's N124 to two very distinct amino acids, alanine and
tryptophan. Unexpectedly, the AvrSr50$^{Q121K/N124W}$ double mutant increased HR when co-infiltrated with Sr50^{K711D},
compared to AvrSr50^{Q121K}. Yet, this enhanced HR was also observed in the co-infiltration with Sr50^{K711A} and Sr50^{K711E}
(Fig. S5). This likely indicated that AvrSr50's N124 may not form specific interactions with Sr50's K711D and may
be simply located in an environment that favors a bulky hydrophobic amino acid. In accordance with this postulation,
AvrSr50^{N124A} diminished cell death, and AvrSr50$^{Q121K/N124A}$ failed to enhance HR (Fig. 2B and 2C). The phenotypic
changes mediated by AvrSr50 N124 mutants without specific association with the K711D substitution suggested that
while AvrSr50's Q121K and N124T contact the LRR domain of Sr50, they may not be interacting with Sr50's K711D.

Sr50's K824 and R904 form an additional contact with AvrSr50

To improve our structural model, we decided to explore another possible contact point between the receptor and the
effector from the initial structural hypothesis (Fig. 1A and 1B). Our previous experiments supported that the upper
part of AvrSr50's terminal alpha helix is involved in the interaction with Sr50 (Fig. 2). Additionally, our initial

screening showed that mutating K824 or R904 in Sr50 to negatively charged amino acids resulted in the complete or
near loss of AvrSr50 recognition (Fig. S3). We hypothesized that Sr50's K824 and R904 interact with negatively
charged side chains of AvrSr50.

200
Figure 3. Sr50's K824 and R904 in terminal leucine-rich repeats form a critical effector binding interface

A and **B.** The predicted structure of AvrSr50. The orientation of the structure is identical in both panels. **A**. Negatively charged
amino acids are indicated in red, as well as Q90 and Q121 in purple. **B**. The estimated electrostatic potential is mapped to the
surface of AvrSr50. **C** and **D**. The qualitative and quantitative cell death phenotypes on *Nicotiana benthamiana*. The optical density
(OD_{600}) was set to 0.3 for receptors and effectors, respectively, and the receptors were expressed under pRPP13. Indicated receptor
and effector pairs were co-infiltrated. The cell death phenotypes were recorded at three days post infiltration. **D**. The mode intensity
of the infiltrated spots was measured with imageJ across the given number of biological replicates in bold. The statistics were
calculated with one-way ANOVA followed by a post-hoc Tukey Honestly Significant Difference test. The compact letters indicate

significant differences between groups. **E**. The predicted poses of Sr50 and AvrSr50 in Model II. Some parts of a loop between
β2–β3, mostly including an unstructured region, are removed from AvrSr50 for visualization (positions 42-66). **F** and **G**. The
superposition of AvrSr50 in Model I and II. The entire complex structures of the two models were superposed to fix the coordinates
of Sr50 consistent. The AvrSr50 structures are then visualized. **G**. Only the terminal alpha helix of AvrSr50 is displayed from the
superposed AvrSr50 structures. **H**. The local environment around Sr50's K824 and R904 in Model II.

Positioning AvrSr50's alpha helix adjacent to the LRR domain of Sr50 as in the initial structural hypothesis (Fig. 1B)
and mapping the locations of negatively charged amino acids (Fig. 3A), we identified two aspartic acids and three
glutamic acids that create a surface with negative electrostatic potential and may face the LRR domain (Fig. 3B).
Under the postulation that some of these residues could restore interactions with Sr50^{K824E} and Sr50^{R904D}, we mutated
each of these residues to lysine. Two exceptions were AvrSr50's E35, which is internally bound to K28 and K79
possibly for protein stability (Fig. S6), and D119 closely located to Q121 and unlikely to contact the two receptor
residues in our current hypothesis (Fig. 1B).

Consequently, we created AvrSr50^{D30K}, AvrSr50^{E115K} and AvrSr50^{E117K} and co-infiltrated them with Sr50^{K824E} or
Sr50^{R904D} (Fig. 3C and 3D). AvrSr50^{D30K} neither disrupted the interaction with Sr50 nor restored the cell death
phenotypes of the Sr50 mutants. AvrSr50^{E115K} induced some HR against Sr50^{R904D}, the magnitude of which is slightly
higher than Sr50^{R904D} and AvrSr50. Notably, AvrSr50^{E117K} induced strong cell death against both Sr50^{K824E} and
Sr50^{R904D}. This suggested that AvrSr50's E117 is possibly proximal to Sr50's K824 and R904.

Refining structural hypotheses with experimental constraints

The identification of the contact point between Sr50's K824 and R904 and AvrSr50's E117 enabled updating our
structural hypothesis (Fig. 1A). We used ColabDock to constrain four pairs of residues from Sr50 and AvrSr50 and
infer a new structural model. Two pairs—K824 and E117, as well as R904 and E117—were derived from the
experimental data (Fig. 3C and 3D), and the other pairs—K711 and Q121, as well as K711 and N124—originated
from Model I, as we could not completely reject the initial structural hypothesis (Fig. 1B).

Our refined structural hypothesis, Model II, displayed notable changes in the position of AvrSr50 compared to Model
I (Fig. 3E and 3F). In this model, ColabDock distorted the structure of AvrSr50's terminal alpha helix to accommodate
the specified restraints (Fig. 3G). Inspecting the structure closely, we found that it would be impossible to physically
satisfy the constraints: the distance between AvrSr50's E117 and Q121 (Fig. 3A) is much smaller than the distance
between Sr50's K711 and K824 (Fig. 1C). This possibly suggested that not all the constraints driven from Model I are
correct, as our experiments hinted (Fig. 2). Sr50's K711 may indeed interact with some other residues than AvrSr50's
Q121 and N124, provided that the interaction between Sr50's K824 and R904 and AvrSr50's E117 was much more
plausible (Fig. 3C). Simulating molecular docking without the two constraints from Model I placed AvrSr50 at the
edge of the upper loop of LRRs (Fig. S7). This violated our experimental observations that suggest the involvement
of Sr50's K711 in the interactions (Fig. 1 and 2) and potentially indicated that ColabDock did not have algorithmic
power to predict the correct conformation without enough restraints. We therefore proceeded with Model II as our
next structural hypothesis.

Additional AvrSr50 residues contact Sr50's K824 and R904

As the structure of AvrSr50 became repositioned in Model II, additional charged or polar effector residues could
potentially interact with Sr50^{K824E} and Sr50^{R904D}: Q90 and D119 (Fig. 3H). Unlike Model I, AvrSr50' D119 lied much
closer to these receptor residues. We evaluated the involvement of the two effector residues in interactions by co-
infiltrating AvrSr50^{Q90K} and AvrSr50^{D119K} with Sr50^{K824E} and Sr50^{R904D}. AvrSr50^{Q90K} could induce cell death against

Sr50^{R904D} but not Sr50^{K824E} (Fig. 3C and 3D). Furthermore, AvrSr50$^{Q90K/E117K}$ double mutant triggered robust HR

against the Sr50$^{K824E/R904E}$ double mutant, a response not observed with any of the single effector mutants. This

supported the AvrSr50's Q90 would interact with Sr50's R904. Similarly, AvrSr50^{D119K} could strongly restore the

abolished HR of Sr50^{K824E} and induced relatively weaker HR against Sr50^{R904D} and Sr50$^{K824E/R904E}$. Notably,

AvrSr50^{D119K} was the only effector mutant that abolished the interaction with Sr50 among the tested mutants at this

interface. This potentially suggested that AvrSr50's D119 may be critical for recognition, interacting with Sr50's

K824 and positioned closely to R904.

Positively charged residues on the terminal LRR leads to auto-activity

Following the revised hypothesis (Fig. 3H), we aimed to validate additional interaction between another pair of

residues: Sr50's E938 and AvrSr50's R80. However, Sr50^{E938K} showed strong auto-activity, making it challenging to

discern phenotypic changes upon co-infiltration with effector mutants (Fig. S8). Subsequently, we attempted Sr50^{T940K}

and Sr50^{T940R} mutants, but they also triggered severe auto-activity (Fig. S8). These outcomes potentially suggested

that the very terminal LRR unit might play a role in stabilizing inactive Sr50, and the introduction of positively charged

residues in these positions might interfere with Sr50's stability.

AvrSr50's alpha helix enriched with positively charged residues are positioned towards the inner concave of
the LRR domain

Model II led to the identification of interacting receptor and effector residues. However, it was not a plausible

biological model with structural distortion (Fig. 3G). Some effector side chains, such as R128, were clashing into the

receptor backbones, creating physically impossible conformation (Fig. 4A). Nevertheless, we noted potential

electrostatic complementarity in this region. Near Sr50's K711 are negatively charged amino acids, D589, D618,

D641 and D643 (Fig. 4A and 4B). Although some of these residues, such as D589 and D618, contact the NB-ARC

domain potentially for interdomain stabilization and may not be available, the others could participate in the interaction

with the effector (Fig. 4B). Notably, AvrSr50 contains H125, R128, R129, H131 and R132 at the end of the terminal

alpha helix (Fig. 4C), which creates strong positive electrostatic potential (Fig. 4D) and potentially lies around the

negatively charged amino acids of Sr50 (Fig. 4A). In particular, R128 lies on the same plane as E117, D119, Q121

and N124 shown to alter the interaction with Sr50 mutants (Fig. 4C).

Based on these observations, we postulated that AvrSr50's R128 would interact with negatively charged amino acids

of Sr50, potentially D641 and D643 that are adjacent (Fig. 4A). We created AvrSr50^{R128E} and AvrSr50^{R129E} for

comparison and co-infiltrated them with Sr50^{D641R} or Sr50^{D643K}. Co-infiltration of Sr50^{D641R} or Sr50^{D643K} with AvrSr50

indicated that the mutations at these positions attenuate AvrSr50 recognition (Fig. 4E and 4F). AvrSr50^{R128E} generated

a variable phenotype from no cell death to strong HR in co-infiltration with Sr50. Nevertheless, AvrSr50^{R128E} could

clearly restore cell death for the Sr50^{D643K} mutant and increase the level of HR against the Sr50^{D641R} mutant. This

suggested that AvrSr50's R128 is possibly proximal to Sr50's D643, as indicated in our model.

Refining structural hypotheses with experimental constraints and AlphaFold

The collection of potential interacting receptor-effector residues was used as restraints to update our structural

hypothesis (Fig. 1A). These included D643 and R128, K824 and E117 as well as D119, and R904 and Q90, E117 as

well as D119 (Fig. 3 and 4). Compared to Model II, Model III slightly repositioned the AvrSr50 structure (Fig. 4G

and 4I), better fitting the terminal alpha helix of AvrSr50 into the groove of the LRR domain.

Figure 4. AvrSr50's alpha helix enriched with positively charged residues are positioned towards the inner concave of the LRR domain

A. The local environment around Sr50's K711 in Model II. Colored in green are aspartic and glutamic acids. **B**. The distribution of selected amino acids across the surface of Sr50's leucine-rich repeat (LRR) domain. In bottom, a portion of the NB-ARC domain that contacts LRR is included. **C** and **D.** The predicted structure of AvrSr50. The orientation of the structure is identical in both panels. **C**. Positively charged amino acids are indicated in blue, negatively charged E117 and D119 in red, and polar Q90, Q121 and N124 in purple. **D.** The estimated electrostatic potential is mapped to the surface of AvrSr50. **E** and **F**. The qualitative and quantitative cell death phenotypes on *Nicotiana benthamiana*. The optical density (OD$_{600}$) was set to 0.3 for receptors and effectors, respectively, and the receptors were expressed under pRPP13. Indicated receptor and effector pairs were co-infiltrated. The cell death phenotypes were recorded at three days post infiltration. **F**. The mode intensity of the infiltrated spots was measured with

imageJ across the given number of biological replicates in bold. The statistics were calculated with one-way ANOVA followed by
a post-hoc Tukey Honestly Significant Difference test. The compact letters indicate significant differences between groups. **G** and
H. The predicted poses of Sr50 and AvrSr50 in Model III and IV. Some parts of a loop between β2–β3, mostly including an
unstructured region, are removed from AvrSr50 for visualization (positions 42-66). **I** and **J**. The superposition of AvrSr50. The
entire complex structures of the two models were superposed to fix the coordinates of Sr50 consistent. The AvrSr50 structures are
then visualized. **K**. The local environment around Sr50's K711 in Model IV.

Although we had some confidence in Model III, we were uncertain whether ColabDock had not introduced any
unexpected alterations in the structures, such as backbone clashes in Model II (Fig. 4G). We used ColabFold to
eliminate abnormal structural features and remodel the flexible loop structures by providing Model III as a structural
template. This refined model, Model IV, was nearly identical to Model III, with only slight alterations in the
positioning of AvrSr50 relative to the LRR domain of Sr50 (Fig. 4H and 4J). In this model, the terminal alpha helix
of AvrSr50 enriched with positively charged side chains was positioned closely to Sr50's D643 and K711, although
AvrSr50's R128 was not oriented to form direct interaction with Sr50's D643 (Fig. 4K)

Multiple Sr50^{K711D} double mutants enhance cell death against AvrSr50^{QCMJC}

Unlike Model I (Fig. 1B), Model IV placed AvrSr50's Q121 near the terminal LRRs of Sr50—LRRs 13 and 14 in
particular (Fig. 5A). We decided to introduce additional mutations to Sr50^{K711D}, targeting W822, E847 and N849 on
these LRRs. These residues were mutated to negatively charged aspartic and glutamic acids, as well as polar
asparagine and glutamine, except for N849Q. Among tested double mutants, Sr50$^{K711D/W822N}$, Sr50$^{K711D/W822Q}$,
Sr50$^{K711D/E847D}$, Sr50$^{K711D/N849D}$ and Sr50$^{K711D/N849E}$ could induce HR against AvrSr50^{QCMJC} (Fig. 5B and 5C). The
average mode intensity of these double mutants was clearly greater than Sr50^{K711D} and AvrSr50^{QCMJC}, collectively
suggesting that based on our final structural model, we could successfully restore the recognition against the escape
mutant. Moreover, the recognition against AvrSr50 was not severely attenuated by these additional mutations (Fig.
5B and 5C).

The Sr50 K711D mutation leads to synergistic impacts and is required for the engineered mutants to recognize AvrSr50^{QCMJC}

To gain more insights into the effect of the second layer of mutations introduced to Sr50^{K711D}, we created Sr50^{W822N},
Sr50^{W822Q}, Sr50^{E847D}, Sr50^{N849D} and Sr50^{N849E} single mutants and examined their abilities to induce HR against
AvrSr50 and AvrSr50^{QCMJC} (Fig. 5D and S9). Sr50^{W822N} and Sr50^{W822Q} single mutants led to detectable HR when co-
infiltrated with AvrSr50^{QCMJC} (Fig. 5E); however, a comparable level of autoactivity was induced, likely indicating
that the observed HR cannot be specifically associated with AvrSr50^{QCMJC} recognition. When the K711D mutation is
additionally introduced, the resulting Sr50$^{K711D/W822N}$ and Sr50$^{K711D/W822Q}$ double mutants were no longer autoactive,
while inducing stronger HR against AvrSr50^{QCMJC} (Fig. 5C). The E847D substitution led to the complete loss of the
receptor's ability to recognize AvrSr50 (Fig. 5D). Sr50^{N849D} and Sr50^{N849E} could still recognize AvrSr50 but did not
gain the ability to cause cell death against AvrSr50^{QCMJC}. In all these cases, the phenotypes of these single mutants
were altered when the K711D mutation was introduced, with gain of AvrSr50^{QCMJC} recognition and without losing
ability to interact with AvrSr50 (Fig. 5B).

The phenotypic discrepancy between the single mutants and the double mutants was unexpected, as we mostly
postulated that mutation effects are likely additive, if the two mutations happen distantly. To gain more insights into
the interplay between substitutions, we generated all combinations of single to triple mutants that include K711D,
W822N, E847D and N849E and quantified the magnitude of HR when the receptor was co-infiltrated with infiltration
buffer, AvrSr50 or AvrSr50^{QCMJC} (Fig. 5B, 5D, 5F and S9). Two triple mutants, Sr50$^{K711D/W822N/E847D}$ and

Sr50$^{K711D/W822N/N849E}$, could induce robust HR against AvrSr50^{QCMJC}. In particular, the magnitude of HR between
Sr50$^{K711D/W822N/E847D}$ and AvrSr50^{QCMJC} was comparable to Sr50 and AvrSr50, despite the recognition of AvrSr50
becoming weaker (Fig. 5C). Sr50$^{K711D/E847D/N849E}$ could not recognize AvrSr50^{QCMJC}, even though Sr50$^{K711D/N849E}$ and
Sr50$^{K711D/E847D}$ did, possibly suggesting that the two negatively charged side chains of E847D and N849E may be
interfering with each other, hindering the recognition. No other combinations of mutations could lead to robust gain
of AvrSr50^{QCMJC} recognition. All mutants that induced HR specifically towards AvrSr50^{QCMJC} required the K711D
substitution.

346
Figure 5. Engineered double and triple Sr50 mutants induce robust HR against AvrSr50^{QCMJC}

A. The local environment around AvrSr50's Q121 in Model IV. **B** to **F**. The qualitative and quantitative cell death phenotypes on
Nicotiana benthamiana. The optical density (OD_{600}) was set to 0.3 for receptors and effectors, respectively, and the receptors were
expressed under pRPP13. Indicated receptor and effector pairs were co-infiltrated. The cell death phenotypes were recorded at
three days post infiltration. **C** and **E**. The mode intensity of the infiltrated spots was measured with imageJ across the given number
of biological replicates in bold. The statistics were calculated with one-way ANOVA followed by a post-hoc Tukey Honestly
Significant Difference test. The compact letters indicate significant differences between groups.

Engineered Sr50 alleles overlaps with a patch of hvLRRs is essential for effector binding, yet they are distinct
from natural variation

We previously curated a set of 89 NLRs belonging to the MLA family sourced from Pooideae species (Tamborski et
al. 2023). Of these, 16 members were identified as part of the Sr50 homologous group based on their close
phylogenetic distances to Sr50. To characterize receptor residues involved in effector recognition, we quantified
sequence variations using normalized Shannon entropy and mapped these values onto the predicted Sr50 structure.
This revealed that a central effector binding site identified in our experiments is indeed associated with highly variable
LRR (hvLRR) residues (Prigozhin and Krasileva 2021; Tamborski et al. 2023). Notably, a patch of hvLRRs with the
greatest Shannon Entropy were distributed at the terminal LRRs above the central beta sheets (Fig. 6A) that includes
K824 and R904 shown as critical specificity-determinants. These regions also contained W822, E847 and N849, which
were altered to enhance the AvrSr50^{QCMJC} recognition.

Furthermore, in proximity to the hvLRR patch are three central tryptophan residues—W771, W799, and W822—
which we initially hypothesized to be essential for affinity towards AvrSr50 through hydrophobic interactions (Fig.
1C and 6A). We observed that W799, closest to the hvLRR patch, is essential for AvrSr50 recognition and cannot be
replaced by tyrosine or phenylalanine, while they can substitute W771 (Fig. S10). Our structural hypothesis suggests
that this hvLRR patch is crucial for interaction with the central binding site of AvrSr50, particularly involving residues
D119 and Q121, the mutations of which were shown to induce recognition escape (Fig. 2 and 3). This potentially
probes the rapid co-evolution at the interaction interface between the receptor and effector.

Figure 6. The structural and evolutionary feature of mutagenized receptor residues

A. Inner beta strands of leucine-rich repeat (LRR) domain of Sr50. The residue is colored based on normalized Shannon entropy
scores, which indicate variability of homologous sequences within the Sr50 family. The score ranges from 0 (no sequence
variability) to 1 (complete variability). The score was capped at 0.75 for visualization. The positions chosen for mutagenesis and

the phenotypes of single mutants are indicated. The mutations indicated with asterisks lead to different phenotypes when K711D
mutation is introduced together. **B**. The frequency of amino acids within the Sr50 family in the given homologous positions. The
mutations introduced in this study are colored brown.

In contrast, D641 and D643 were non-hvLRRs, with D641 displaying perfect sequence conservation among the
homologous sequences and glycine being the predominant amino acid at position 643 (Fig. 7B). This conservation
may be attributed to their proximity to the NB-ARC domain, where incompatible mutations might also disrupt
interdomain interactions.

Within the 16 members of the Sr50 homologous group, the mutations we introduced to create functional Sr50 double
and triple mutants occurred rarely (Fig. 7B). Aspartic acid appeared in 6.25% of instances at position 711, asparagine
and glutamine in no instances at position 822, aspartic acid in 12.5% at position 847, and glutamic acid in no instances
at position 849. Aspartic acid was relatively frequent at position 849 with 30% of instances. No homologous
sequences, however, had the combinations of amino acids shown to induce HR against AvrSr50^{QCMJC}. Collectively,
although our structure-guided rational design approaches targeted the region of the receptor in which the greatest
natural variation occurs, our engineered receptor alleles were distinct from known natural variations (Fig. 5 and 6B).

Amino acid substitutions in input sequences alter the behavior of AlphaFold 2, and the K711D substitution
enables prediction of the complex structure of Sr50 and AvrSr50

We initially observed that AF2-Multimer was unable to accurately predict the structure of Sr50 and AvrSr50
complexes (Fig. 7A and S1). We attempted modeling the complex structure of Sr50$^{K711D/W822N/E847D}$ and AvrSr50^{QCMJC},
as the engineered receptor mutant could induce strong HR against the effector. Although AF2 assigned low accuracy
to the prediction (Fig. 7B), AF2 predicted a complex structure of Sr50$^{K711D/W822N/E847D}$ and AvrSr50^{QCMJC} that nearly
matched model IV (Fig. 7A). Interestingly, the predicted heterodimeric complex of Sr50$^{K711D/W822N/E847D}$ and AvrSr50
also showed great similarity.

To examine the sequence variation that switched the behavior of AF2, we modeled the protein complex structure of
Sr50^{K711D}, Sr50^{W822N} and Sr50^{E847D} with AvrSr50. Only the K711D substitution, but not W822N and E847
substitutions, could alter the outcome of AF2, enabling the prediction of the Sr50^{K711D} and AvrSr50 complex that
agreed with Model IV (Fig. 7A and S11). This observation was consistent for Sr50^{K711D} and AvrSr50^{QCMJC} (Fig.7A).
These results collectively indicated that the single amino substitution, K711D, not only altered the biological response
of the receptor to the effector (Fig. 1) but was also sufficient to change the outcome of AF2 prediction.

In a recent study, Gómez De La Cruz et al. modeled the structure of AvrSr50 and Sr50^{3BI} with AF2 to predict the
potential binding site of AvrSr50 (2024). Sr50^{3BI} contains a scratch of 25 amino acids at positions 920 to 950
transferred from barley NLR MLA3 that recognizes its effector Pwl2. Although Sr50 and Sr50^{3BI} differ only by 12
amino acids, AF2 could produce a high confidence model for Sr50^{3BI} and AvrSr50. Under the identical conditions
used to predict Sr50^{K711D} and AvrSr50, we modeled the complex structure of Sr50^{3BI} and AvrSr50 and could obtain
the heterodimeric complex structure as previously reported (Gómez De La Cruz et al. 2024) (Fig. 7A). This model
showed good agreement with Model IV as well as the AF models of Sr50^{K711D} and Sr50$^{K711D/W822N/E847D}$ complexes
(Fig. 7A and 7C). However, AF2 failed to predict the structure of Sr50^{3BI} and AvrSr50^{QCMJC} (Fig. 7A). Notably, the
predicted accuracy of the complex structure (iptm) and the predicted aligned errors showed high confidence for Sr50^{3BI}
and AvrSr50, as reported by the authors (Gómez De La Cruz et al. 2024). Despite similar topologies, AF2 did not
report comparable confidence for Sr50^{K711D} or Sr50$^{K711D/W822N/E847D}$ and AvrSr50 or AvrSr50^{QCMJC} (Fig. 7B). These
results indicated that slight changes in the input sequences can alter the behavior and outcome of AF2 prediction; This

potentially suggests that iterative in silico mutagenesis of input sequences could aid the prediction of NLR-effector
complexes.

421
Figure 7. AlphaFold 2 predicts the complex structure of Sr50 and AvrSr50

A. The structural superposition of AvrSr50 in AlphaFold 2 (AF2) models compared to Model IV. The predicted protein complex
structures were superposed against Model IV to keep the coordinates of Sr50 consistent. The AvrSr50 structures were then
visualized. **B**. The confidence scores of AF2 models. The ptm and iptm scores, as well as the predicted aligned error (PAE) plots

are shown for the best model (Elfmann and Stülke 2023). The colored boxes correspond to the predicted model specified in A. **C**.
The comparison of protein complex structures with DockQ scores. The DockQ scores can be divided into incorrect (0 < DockQ <
0.23), acceptable quality (0.23 ≤ DockQ < 0.49), medium quality (0.49 ≤ DockQ < 0.80) and high quality (0.80 ≤ DockQ ≤ 1). The
heterodimeric complexes of Sr35 and AvrSr50 solved by two different groups using the Cryo-EM structures produce a DockQ
score of 0.76 (PDB: 7XC2 and 7XE0) (Förderer et al. 2022; Zhao et al. 2022). The DockQ score was capped at 0.76 for comparison.
The loop between β2–β3, mostly including an unstructured region, is excluded (positions 42-66). **D**. The comparison of structural
hypotheses to Model IV and AF structures.

Discussion

Experimentally determined NLR-effector complex structures can aid in developing engineering strategies against
evolving effectors, but they may not always guarantee success. These structures provide static snapshots, while
proteins are inherently dynamic. Predicting the impact of new effector mutations on the interaction with NLRs and
host disease phenotypes remains challenging even with reliable structures. Therefore, accumulating evolutionary and
experimental data is crucial for effective engineering solutions. Our results highlight that iterative computational
modeling approaches together with experimental determination of structural constraints is an effective strategy for
engineering the plant intracellular immune receptor to restore recognition of the escape effector mutants. Below, we
outline main lessons and suggestions from our work that could be used for guiding experimental resurrection of other
NLRs as well as for developing machine learning models.

Reducing search space for efficient and rational mutagenesis-based approaches

Our approach operates without experimentally determined receptor-effector complex structures and evolutionary data
from receptors known to recognize their escape effector mutants. In this circumstance, exploring all mutational
landscapes through experiments is currently impossible due to high order of information complexity. To bypass this
problem, we simplify candidate residue selection and mutagenesis process, relying on the central biochemical
principle of protein-protein interactions: while hydrophobic interactions provide the main energy for ligand binding,
it is the charged residues that determine ligand specificity through ionic or hydrogen bonds. Our experiments
demonstrated that altering the specificity determinants to amino acids with their opposite charges can be a practical
approach to confirm the interactions between receptor and effector residues and their proximity.

A recent study demonstrated that the single L902S substitution led to gain of recognition of Avr_{A13}-1 in MLA7
(Lawson et al. 2024). Similar solutions may exist for Sr50 to acquire resistance to $AvrSr50^{QCMJC}$, but identifying them
through our methodology can be challenging. In similar contexts, if the effector mutants escape through steric clashes
and changes in affinity-determinants, engineering solutions would be more difficult to obtain. Nevertheless, our work
effectively elucidates mutation effects on the NLR-effector interactions across the LRR domain, creates engineering
solutions and generates structural hypotheses, which can be useful to solve current and future problems against
evolving effectors.

All models are wrong, but some are useful

Our workflow to predict NLR-effector interactions showed that, as of today, modeling-based engineering approaches
need to be iterative. We used three iterations of experimentally constrained modeling and a round of computational
refinement to derive the final solution, Model IV. Although the actual accuracy of the model can only be evaluated
when Cryo-EM structure becomes available, this model showed great similarity to independently derived AF models
(Fig. 7). When we compared all modeling iterations to Model IV and other AF models to evaluate their accuracy, we
noted that our initial structural hypothesis (Model I) was completely incorrect with the DockQ scores between 0.11

and 0.15 (Fig. 7D). However, in each iteration of model refinement, the structural hypothesis continued to improve,
suggesting that although these models were imperfect, they were useful to derive the next hypothesis.

The main aim of our structural hypotheses is to refine and streamline the extensive and random screening processes
into more targeted approaches by providing structural contexts and hypotheses. This approach does not aim to
precisely depict molecular interactions or replace experimentally determined structures but to offer practical solutions
in the absence of experimentally determined or accurately predicted NLR-effector complex structures. This
methodology can be expanded to other NLR-effector systems, providing a useful framework for addressing evolving
effector challenges.

Hacking AlphaFold

Our study initially began with the observation that AF struggles to predict the correct topology of receptor-effector
complexes, as evidenced by the recent Cryo-EM structure of MLA13 and its effector AVR_{A13}-1 (Lawson et al. 2024).
Surprisingly, AF2 produced high-confidence models for MLA3 and its effector Pwl2, as well as for Sr50[3BI] and
AvrSr50 (Gómez De La Cruz et al. 2024). Moreover, a single amino acid change in Sr50 rectified the previously
unsuccessful prediction of Sr50 and AvrSr50 complexes, resulting in models that match our final structural hypothesis
(Fig. 7). AF2 is not specifically trained to predict the impact of small amino acid mutations in protein folding.
However, our study, along with the previous research, suggests that small changes in input sequences can significantly
alter the outcome of the complex structure prediction. This observation hints at the potential for in silico screening of
receptor-effector complexes, provided we can uncover methods to effectively 'hack' AF2 to our advantage.

Figure 8. The initial mutation significantly influences subsequent evolutionary opportunities

A. Phenotypes of engineered Sr50 mutants along possible mutational paths. The circles are composed of three layers, which indicate
the strength of hypersensitive responses (HR) of the given receptors towards buffer (autoactivity), AvrSr50 and AvrSr50[QCMJC] co-
infiltrations. Each wild type Sr50 accumulates a single mutation per event. The paths highlight additional mutations towards
enhanced HR against AvrSr50[QCMJC] or reduced autoactivity. **B**. The impact of the initial mutations in subsequent mutational
opportunities. The initial mutation can significantly influence subsequent evolutionary opportunities, either limiting or expanding
them, even though the overall paths may lead to an identical set of mutations.

The order of mutations impacts the emergence of functional NLRs against escape effector mutants

The order of mutations and resulting phenotypic changes provide insights into the natural evolution of functional
NLRs (Fig. 8). Consider a scenario where wheat accumulates mutations in the interaction with *P. graminis* QCMJC
in the field. Substitutions like E847D, which abolish the interaction with effectors, or N849E, which do not lead to

phenotypic changes, may appear first (Fig. 8A; I and II). Individuals with these mutations would likely be selected
against, preventing them from acquiring a second mutation, such as K711D, even though the double mutations can
ensure resistance against AvrSr50^{QCMJC}. To resurrect resistance through multiple mutations, NLRs would initially
need to develop weak to intermediate resistance, as seen in Sr50^{K711D}, or some level of autoactivity at the cost of
development, as observed in Sr50^{W822N} (Fig. 8A; III and IV). These early mutations could lead to additional mutations
that either enhance HR or reduce autoactivity in regulatory regions or protein-coding sequences. Notably, the
emergence of K711D as an initial mutation opens more avenues for acquiring robust resistance compared to other
tested mutations (Fig. 8A). This suggests that multiple solutions exist to achieve gain of recognition towards escape
mutants, but the path to these solutions may be restricted (Fig. 8B). In other words, the initial mutation can significantly
influence subsequent evolutionary opportunities, either preventing or expanding them, even though the overall paths
may lead to an identical set of final mutations.

We postulate that Sr50's K711 only had a supplementary role in the initial interactions between Sr50 and AvrSr50.
The mutations at position 711 did not significantly impact the AvrSr50 recognition (Fig. 1 and S5). However, when
K121 of AvrSr50^{QCMJC} disrupts the interaction with the hvLRRs patch around Sr50's K824 and R904 (Fig. 3 and Fig.
6), the K711D substitution in Sr50 would likely facilitates additional interactions with the alpha helix of AvrSr50^{QCMJC}
enriched with positively charged arginines (Fig. 4). This interaction may be subtle as the HR between Sr50^{K711D} and
AvrSr50^{QCMJC} was only weak to intermediate (Fig. 1 and 2); however, this may be essential for the upper alpha helix
of AvrSr50^{QCMJC} to align properly with the terminal LRRs of Sr50. This alignment potentially allows for more stable
binding to the receptor, thereby enabling the engineered receptor mutants to recognize the escape mutant only in the
presence of the K711D substitution (Fig. 5).

Expanding to other NLR-effector systems

Our data, derived from 6,000 quantifications from infiltrations, provides robust representation of a wide spectrum of
biological responses between cognate NLR and effector variants (Table S1). This dataset can be useful in correlating
biological responses and in silico prediction, serving as an experimentally derived training dataset for the future
computational or machine learning approaches. Expanding recognition specificity to other paralogs and sequence-
unrelated structurally similar (SUSS) effectors is the next challenge (Seong and Krasileva 2021, 2023). While AvrSr50
forms a SUSS effector family, no other members are yet known to interact with NLRs. Alternatively, MLA receptors
from barley and their cognate effectors from powdery mildew, *Blumeria graminis*, can be more compelling systems
(Saur et al. 2019). Not only do MLAs share close evolutionary relationships with Sr50 (Tamborski et al. 2023), but
also they recognize sequence-divergent or SUSS effectors originating from an extensively expanded RNAse-like
protein family with many experimentally determined structures (Pennington et al. 2019; Cao et al. 2023; Seong and
Krasileva 2023; Lawson et al. 2024). Leveraging preexisting rich experimental data can aid in inferring initial
structural hypotheses and predicting the molecular interactions between MLAs and their effectors. Elucidating the
complex structures of diverse Sr and MLA receptors and their effectors can reveal how plants evolved NLRs to detect
SUSS effectors, and through the framework of this study, we may guide experimental designs and engineering
solutions for other NLR-effector systems.

Methods

Vectors and mutagenesis

We used the previously generated constructs (Tamborski et al. 2023). The receptor constructs carried *Sr50* under the
pRPP13 promoter and the *tNos* terminator, or the *p35S* promoter and the *t35S* terminator. The effector constructs
contained *AvrSr50* and *AvrSr50^{QCMJC}* with the *p35S* promoter and *t35S* terminator. All additional mutations were

introduced with QuikChange Lightning Site-directed Mutagenesis Kits from Agilent. We followed the standard
protocol but reduced the volume of all reagents by four. All primers and mutants used in this study are available in
Table S2.
540

Transformation and mutation confirmation

542   XL10-Gold ultracompetent cells were transformed, following the standard protocol given in QuikChange Lightning
543   Site-directed Mutagenesis Kits. Plasmids were extracted from a liquid culture inoculated with a single colony,
544   following the standard protocol of The QIAprep Spin Miniprep Kit. The desired mutation was confirmed with Sanger
545   sequencing.
546

547   About 40 µL of *Agrobacterium tumefaciens* GV3101:pMP90 was mixed with 100 ng of the mutagenized plasmid in
548   a 1.5 ml plastic tube and transferred to an electroporation cuvette. After an electric pulse, the cells were transferred
549   back to the tube with 250 µL of LB media and shaken at 28°C and 250 rpm for two hours. The liquid culture was
550   plated on LB agar containing Carbenicillin, Rifampicin and Gentamicin, and the plates were incubated at 28°C for
551   two days. A single colony was picked and transferred to a liquid LB medium and grown for a day.
552

We additionally confirmed the mutations in the transformed *Agrobacterium*. Each mutant was replated from its
glycerol stock on LB agar containing Carbenicillin, Rifampicin and Gentamicin and grown at 28°C for two days.
Colonies were scraped from the plates, resuspended in 10 µL of water and incubated at 98°C for 10 minutes. The cells
were centrifuged for one minute, and target regions with introduced mutations were amplified, following the PCR
Protocol for Phusion® High-Fidelity DNA Polymerase (M0530) or for repliQa HiFi ToughMix. The mutation was
confirmed with Sanger sequencing.
559

Agrobacterium*-mediated transient gene expression in *N. benthamiana

The liquid cultures containing *Agrobacterium* transformants were centrifuged at 6,400g for five minutes, and the
pellets were resuspended in infiltration medium composed of deionized water, 10 mM MES (pH 5.6), 10 mM $MgCl_2$,
150 µM acetosyringone. The optical density (OD_{600}) of each transformant was re-adjusted to 0.6. The transformants
carrying receptors and effectors were mixed to adjust the OD_{600} of the receptors to 0.1 (for p35S) or 0.3 (for *pRPP13*).
The OD_{600} of the effectors was set to 0.3. *N. benthamiana*'s 4-5 weeks old leaves were inoculated with the suspension
using blunt syringes. The phenotypes were recorded at 2 dpi for the *p35S* promoter and at 3 dpi for the *pRPP13*
promoter.
568

Cell death quantification and statistics

The infiltrated *N. benthamiana* leaves were imaged with the ChemiDoc MP Imaging System and Image Lab v5.2.1
(https://www.bio-rad.com). Following the previous publication (Landeo Villanueva et al. 2021), we used green epi-
illumination with a filter set to 605/50. Exposure time was set to 0.5 seconds. To quantify the cell death, we manually
selected the treated area and measured the mode of intensity with ImageJ v2.14.0 (Rueden et al. 2017). The statistics
were computed in R v4.1.3 (Ihaka and Gentleman 1996). One-way ANOVA was performed to compare significant
differences between the sample means, and post-hoc Tukey's honest significance test was followed for pairwise
comparisons.
577

Protein extraction and co-immunoprecipitation assays

Six *N. benthamiana* leaf discs of 0.8 cm in diameter were collected at 1 dpi after *Agrobacterium*-mediated infiltrations.
The samples were frozen in liquid nitrogen and ground with a bead beater at 1,500 Hz for 1 min with two 3.2 mm
stainless beads. Protein was extracted with 300 µL of the 2x Laemmli sample buffer (Biorad) with 5% β-
mercaptoethanol. Samples were boiled at 95°C for 5 min and centrifuged at maximum speed for 10 min at 4 °C.

Supernatants were transferred into fresh tubes for SDS PAGE analysis. 10L of protein extraction was separated on
15% Mini-PROTEAN® TGX™ Precast Protein Gels (Biorad, 15-well), transferred to PVDF membrane (Biorad) at
300 mA for 70 min. Immunoblotting was performed using rat HRP-conjugated α-HA (monoclonal 3F10, Roche) and
subsequently chemiluminescent substrate SuperSignal™ West Pico PLUS (Thermo Scientific™). Total protein was
stained using Ponceau S.
588

Protein structure prediction and visualization

The initial Sr50 and AvrSr50 structures were predicted by AlphaFold v2.2.2 (Jumper et al. 2021), with the full
database, available homologous templates and model_preset set to monomer. Protein complexes were predicted with
ColabFold v1.5.2 that relies on AlphaFold v2.3.1, with alphafold2_multimer_v3 and template_model set to pdb100,
as well as AlphaFold 3 (Evans et al. 2021; Mirdita et al. 2022; Abramson et al. 2024). We used customized ColabDock
(Feng et al. 2023) through Google Colab to obtain Models II and III. Pairwise constraints derived from the experiments
were provided. Among the five models produced by ColabDock, the best structure (1st_best) was used for the
analyses. The side chains of this structure were relaxed with Amber, a module within ColabFold.

The AF2 models, which were compared to Model IV, were generated with ColabFold v1.5.5. The structure of
AvrSr50^{QCMJC} was submitted as a template (PDB:7MQQ) (Ortiz et al. 2022). All loop regions were excluded from the
initial PDB file to allow ColabFold to remodel the flexible loop structures. Five models were predicted with the
alphafold2_multimer_v3 model, and the best structure was relaxed. The default parameters were used, except for
num_recycles set to 24 and pair_mode changed to unpaired. We used PyMOL v2.5.2 for protein structure
superposition and visualization (The PyMOL Molecular Graphics System). The analysis of the electrostatic potential
relied on the APBS plugin in PyMOL (Jurrus et al. 2018).

Molecular docking and initial model selection

The best Sr50 and AvrSr50 monomer models were submitted to ZDOCK, HDOCK and ClusPro web servers as a
receptor and a ligand, respectively (Pierce et al. 2014; Kozakov et al. 2017; Yan et al. 2020). From each server, 100
models were obtained, and each model was evaluated for the following criteria, based on the backbone distances (C_β
or C_α of glycine). First, all Sr50 residues in the coiled-coil and NB-ARC domains (positions 1 to 520) are not within
8 Å of AvrSr50. Second, AvrSr50 should touch the NB-ARC latch–a loop structure exposed from the NB-ARC
domain–, forming close contact with LRRs (positions 492-499). Specifically, Sr50's E494 is within 12 Å of AvrSr50.
This residue was chosen as the predicted Sr50 structure suggested that its sidechain points toward the putative effector
binding site surrounded by the concave of LRR units. The distance cut-off was relaxed based on our assumption that
the interaction between the NB-ARC latch and the effectors occur through long side chains. Third, AvrSr50's Q121
is within 8 Å of Sr50's LRR domain (Ortiz et al. 2022; Tamborski et al. 2023). Fourth, at least 12 hvLRR residues on
the inner β-strands or upper loops of Sr50 are within 8 Å of AvrSr50 (Prigozhin and Krasileva 2021). Then, the effector
poses of the models that satisfied these criteria were clustered with the RMSD (root mean square deviation) cut-off of
3.0. We required at least two predictors to produce similar poses, despite their differing scoring functions and
underlying algorithms. A representative conformation was chosen from each cluster based on the model ranking.

Protein structure prediction and visualization

The initial Sr50 and AvrSr50 structures were predicted by AlphaFold v2.2.2 (Jumper et al. 2021), with the full
database, available homologous templates and model_preset set to monomer. Protein complexes were predicted with
ColabFold v1.5.2 that relies on AlphaFold v2.3.1, with alphafold2_multimer_v3 and template_model set to pdb100,
as well as AlphaFold 3 (Evans et al. 2021; Mirdita et al. 2022; Abramson et al. 2024)..We used customized ColabDock
(Feng et al. 2023) through Google Colab. Among the five models produced by ColabDock, the best structure that
satisfied all provided restraints was selected. To further refine this model, we ran AlphaFold through ColabFold with
the model given as a customized template. The best prediction was relaxed with amber.

The AF2 models, which were compared to Model IV, were generated with ColabFold v1.5.5. The structure of AvrSr50^{QCMJC} was submitted as a template (PDB:7MQQ) (Ortiz et al. 2022). All loop regions were excluded from the initial PDB file to allow ColabFold to remodel the flexible loop structures. Five models were predicted with the alphafold2_multimer_v3 model, and the best structure was relaxed. The default parameters were used, except for num_recycles set to 24 and pair_mode changed to unpaired. We used PyMOL v2.5.2 for protein structure superposition and visualization. The analysis of the electrostatic potential relied on the APBS plugin in PyMOL (Jurrus et al. 2018). Structural similarities between complex structures were quantified with DockQ v2.1.1 (Basu and Wallner 2016).

Evolutionary analyses

The multiple sequence alignment of MLA family members and the normalized Shannon entropy for the Sr50 homologous group were obtained from the previous study and analyzed with the identical workflow (Tamborski et al. 2023).

Data availability

All scripts and command lines used for computational analyses are available and recorded in Github: https://github.com/s-kyungyong/Sr50_AvrSr50/. All input, intermediate and output data were deposited in Zenodo: https://zenodo.org/doi/10.5281/zenodo.13205869

Acknowledgements

We thank China Lunde for phenotyping, and Sarah Weber for attempting transient gene expression assays, and Chandler Sutherland for the critical review of the manuscript. K.S. is supported by the Berkeley BioEnginuity Fellowship. K.V.K. has been supported by the Gordon and Betty Moore Foundation (grant number: 8802) as well as the joint funding from the Foundation for Food and Agriculture and 2Blades (CA19-SS-0000000046), the Innovative Genomics Institute and the National Institute of Health Director's Award (1DP2AT011967-01).

Contributions

K.S. conceptualized and designed the project. K.S. wrote the manuscript with edits from K.V.K and W.W. W.W. performed Western blot. K.S. performed computational analyses. K.S., W.W., B.V., A.D., G.R., R.K. and L.P. conducted experiments. K.V.K. supervised the research.

Competing interests

The authors declare no competing interests.

References

Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L, Ballard AJ, Bambrick J, et al. Accurate structure prediction of biomolecular interactions with AlphaFold 3. Nature. 2024. https://doi.org/10.1038/s41586-024-07487-w

Arora S, Steuernagel B, Gaurav K, Chandramohan S, Long Y, Matny O, Johnson R, Enk J, Periyannan S, Singh N, et al. Resistance gene cloning from a wild crop relative by sequence capture and association genetics. Nat Biotechnol. 2019:**37**(2):139–143. https://doi.org/10.1038/s41587-018-0007-9

Baggs E, Dagdas G, and Krasileva K. NLR diversity, helpers and integrated domains: making sense of the NLR IDentity. Curr Opin Plant Biol. 2017:**38**:59–67. https://doi.org/10.1016/j.pbi.2017.04.012

Basu S and Wallner B. DockQ: A Quality Measure for Protein-Protein Docking Models. PLOS ONE. 2016:**11**(8):e0161879. https://doi.org/10.1371/journal.pone.0161879

Bentham AR, De La Concepcion JC, Benjumea JV, Kourelis J, Jones S, Mendel M, Stubbs J, Stevenson CEM,

Maidment JHR, Youles M, et al. Allelic compatibility in plant immune receptors facilitates engineering of
new effector recognition specificities. Plant Cell. 2023:**35**(10):3809–3827.
https://doi.org/10.1093/plcell/koad204

Cao Y, Kümmel F, Logemann E, Gebauer JM, Lawson AW, Yu D, Uthoff M, Keller B, Jirschitzka J, Baumann
U, et al. Structural polymorphisms within a common powdery mildew effector scaffold as a driver of
coevolution with cereal immune receptors. Proc Natl Acad Sci. 2023:**120**(32):e2307604120.
https://doi.org/10.1073/pnas.2307604120

Cesari S, Xi Y, Declerck N, Chalvon V, Mammri L, Pugnière M, Henriquet C, De Guillen K, Chochois V,
Padilla A, et al. New recognition specificity in a plant immune receptor by molecular engineering of its
integrated domain. Nat Commun. 2022:**13**(1):1524. https://doi.org/10.1038/s41467-022-29196-6

Chen J, Upadhyaya NM, Ortiz D, Sperschneider J, Li F, Bouton C, Breen S, Dong C, Xu B, Zhang X, et al.
Loss of *AvrSr50* by somatic exchange in stem rust leads to virulence for *Sr50* resistance in wheat. Science.
2017:**358**(6370):1607–1610. https://doi.org/10.1126/science.aao4810

Contreras MP, Pai H, Selvaraj M, Toghani A, Lawson DM, Tumtas Y, Duggan C, Yuen ELH, Stevenson CEM,
Harant A, et al. Resurrection of plant disease resistance proteins via helper NLR bioengineering. Sci Adv.
2023:**9**(18):eadg3861. https://doi.org/10.1126/sciadv.adg3861

Dangl JL, Horvath DM, and Staskawicz BJ. Pivoting the Plant Immune System from Dissection to Deployment.
Science. 2013:**341**(6147):746–751. https://doi.org/10.1126/science.1236011

De La Concepcion JC, Franceschetti M, MacLean D, Terauchi R, Kamoun S, and Banfield MJ. Protein
engineering expands the effector recognition profile of a rice NLR immune receptor. eLife. 2019:**8**:e47713.
https://doi.org/10.7554/eLife.47713

Dodds PN and Rathjen JP. Plant immunity: towards an integrated view of plant–pathogen interactions. Nat Rev
Genet. 2010:**11**(8):539–548. https://doi.org/10.1038/nrg2812

Elfmann C and Stülke J. PAE viewer: a webserver for the interactive visualization of the predicted aligned error for
multimer structure predictions and crosslinks. Nucleic Acids Res. 2023:**51**(W1):W404–W410.
https://doi.org/10.1093/nar/gkad350

Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, Žídek A, Bates R, Blackwell S, Yim J, et al.
Protein complex prediction with AlphaFold-Multimer (Bioinformatics).
https://doi.org/10.1101/2021.10.04.463034

Farnham G and Baulcombe DC. Artificial evolution extends the spectrum of viruses that are targeted by a disease-
resistance gene from potato. Proc Natl Acad Sci. 2006:**103**(49):18828–18833.
https://doi.org/10.1073/pnas.0605777103

Feng S, Chen Z, Zhang C, Xie Y, Ovchinnikov S, Gao Y, and Liu S. ColabDock: inverting AlphaFold structure
prediction model for protein-protein docking with experimental restraints (Bioinformatics).
https://doi.org/10.1101/2023.07.04.547599

Förderer A, Li E, Lawson AW, Deng Y, Sun Y, Logemann E, Zhang X, Wen J, Han Z, Chang J, et al. A wheat
resistosome defines common principles of immune receptor channels. Nature. 2022:**610**(7932):532–539.
https://doi.org/10.1038/s41586-022-05231-w

Gómez De La Cruz D, Zdrzałek R, Banfield MJ, Talbot NJ, and Moscou MJ. Molecular mimicry of a pathogen
virulence target by a plant immune receptor. 2024. https://doi.org/10.1101/2024.07.26.605320

Harris CJ, Slootweg EJ, Goverse A, and Baulcombe DC. Stepwise artificial evolution of a plant disease resistance
gene. Proc Natl Acad Sci. 2013:**110**(52):21189–21194. https://doi.org/10.1073/pnas.1311134110

Huang H, Huang S, Li J, Wang H, Zhao Y, Feng M, Dai J, Wang T, Zhu M, and Tao X. Stepwise artificial
evolution of an Sw-5b immune receptor extends its resistance spectrum against resistance-breaking isolates
of *Tomato spotted wilt virus*. Plant Biotechnol J. 2021:**19**(11):2164–2176. https://doi.org/10.1111/pbi.13641

Ihaka R and Gentleman R. R: A Language for Data Analysis and Graphics. J Comput Graph Stat. 1996:**5**(3):299–
314. https://doi.org/10.1080/10618600.1996.10474713

Jones JDG and Dangl JL. The plant immune system. Nature. 2006:**444**(7117):323–329.
https://doi.org/10.1038/nature05286

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A,
Potapenko A, et al. Highly accurate protein structure prediction with AlphaFold. Nature.
2021:**596**(7873):583–589. https://doi.org/10.1038/s41586-021-03819-2

Jurrus E, Engel D, Star K, Monson K, Brandi J, Felberg LE, Brookes DH, Wilson L, Chen J, Liles K, et al.
Improvements to the APBS biomolecular solvation software suite. Protein Sci. 2018:**27**(1):112–128.
https://doi.org/10.1002/pro.3280

Kourelis J, Schuster M, Demir F, Mattinson O, Krauter S, Kahlon PS, O'Grady R, Royston S, Bravo-Cazar

AL, Mooney BC, et al. Bioengineering secreted proteases converts divergent Rcr3 orthologs and paralogs
into extracellular immune co-receptors. Plant Cell. 2024:koae183. https://doi.org/10.1093/plcell/koae183

Kozakov D, Hall DR, Xia B, Porter KA, Padhorny D, Yueh C, Beglov D, and Vajda S. The ClusPro web server
for protein–protein docking. Nat Protoc. 2017:**12**(2):255–278. https://doi.org/10.1038/nprot.2016.169

Kroj T, Chanclud E, Michel-Romiti C, Grand X, and Morel J. Integration of decoy domains derived from protein
targets of pathogen effectors into plant immune receptors is widespread. New Phytol. 2016:**210**(2):618–626.
https://doi.org/10.1111/nph.13869

Landeo Villanueva S, Malvestiti MC, Van Ieperen W, Joosten MHAJ, and Van Kan JAL. Red light imaging for
programmed cell death visualization and quantification in plant–pathogen interactions. Mol Plant Pathol.
2021:**22**(3):361–372. https://doi.org/10.1111/mpp.13027

**Lawson AW, Flores-Ibarra A, Cao Y, An C, Neumann U, Gunkel M, Saur IML, Chai J, Behrmann E, and
Schulze-Lefert P**. The barley MLA13-AVR $_{A13}$ heterodimer reveals principles for immunoreceptor
recognition of RNase-like powdery mildew effectors. 2024. https://doi.org/10.1101/2024.07.14.603419

Liu F, Yang Z, Wang C, Martin R, Qiao W, Carette JE, Luan S, Nogales E, and Staskawicz B. The activated
plant NRC4 immune receptor forms a hexameric resistosome (Plant Biology).
https://doi.org/10.1101/2023.12.18.571367

Ma S, Lapin D, Liu L, Sun Y, Song W, Zhang X, Logemann E, Yu D, Wang J, Jirschitzka J, et al. Direct
pathogen-induced assembly of an NLR immune receptor complex to form a holoenzyme. Science.
2020:**370**(6521):eabe3069. https://doi.org/10.1126/science.abe3069

**Madhuprakash J, Toghani A, Contreras MP, Posbeyikian A, Richardson J, Kourelis J, Bozkurt TO, Webster
M, and Kamoun S**. A disease resistance protein triggers oligomerization of its NLR helper into a hexameric
resistosome to mediate innate immunity. 2024. https://doi.org/10.1101/2024.06.18.599586

**Mago R, Zhang P, Vautrin S, Šimková H, Bansal U, Luo M-C, Rouse M, Karaoglu H, Periyannan S, Kolmer
J, et al.** The wheat Sr50 gene reveals rich diversity at a cereal disease resistance locus. Nat Plants.
2015:**1**(12):15186. https://doi.org/10.1038/nplants.2015.186

**Maidment JH, Shimizu M, Bentham AR, Vera S, Franceschetti M, Longya A, Stevenson CE, De La Concepcion
JC, Białas A, Kamoun S, et al.** Effector target-guided engineering of an integrated domain expands the
disease resistance profile of a rice NLR immune receptor. eLife. 2023:**12**:e81123.
https://doi.org/10.7554/eLife.81123

Martin R, Qi T, Zhang H, Liu F, King M, Toth C, Nogales E, and Staskawicz BJ. Structure of the activated ROQ1
resistosome directly recognizing the pathogen effector XopQ. Science. 2020:**370**(6521):eabd9993.
https://doi.org/10.1126/science.abd9993

Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, and Steinegger M. ColabFold: making protein folding
accessible to all. Nat Methods. 2022:**19**(6):679–682. https://doi.org/10.1038/s41592-022-01488-1

Möller M and Stukenbrock EH. Evolution and genome architecture in fungal plant pathogens. Nat Rev Microbiol.
2017:**15**(12):756–771. https://doi.org/10.1038/nrmicro.2017.76

**Ortiz D, Chen J, Outram MA, Saur IML, Upadhyaya NM, Mago R, Ericsson DJ, Cesari S, Chen C, Williams
SJ, et al.** The stem rust effector protein AvrSr50 escapes Sr50 recognition by a substitution in a single
surface-exposed residue. New Phytol. 2022:**234**(2):592–606. https://doi.org/10.1111/nph.18011

**Pennington HG, Jones R, Kwon S, Bonciani G, Thieron H, Chandler T, Luong P, Morgan SN, Przydacz M,
Bozkurt T, et al.** The fungal ribonuclease-like effector protein CSEP0064/BEC1054 represses plant
immunity and interferes with degradation of host ribosomal RNA. PLOS Pathog. 2019:**15**(3):e1007620.
https://doi.org/10.1371/journal.ppat.1007620

Pierce BG, Wiehe K, Hwang H, Kim B-H, Vreven T, and Weng Z. ZDOCK server: interactive docking prediction
of protein–protein complexes and symmetric multimers. Bioinformatics. 2014:**30**(12):1771–1773.
https://doi.org/10.1093/bioinformatics/btu097

Prigozhin DM and Krasileva KV. Analysis of intraspecies diversity reveals a subset of highly variable plant immune
receptors and predicts their binding sites. Plant Cell. 2021:**33**(4):998–1015.
https://doi.org/10.1093/plcell/koab013

Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, and Eliceiri KW. ImageJ2: ImageJ
for the next generation of scientific image data. BMC Bioinformatics. 2017:**18**(1):529.
https://doi.org/10.1186/s12859-017-1934-z

Sánchez-Vallet A, Fouché S, Fudal I, Hartmann FE, Soyer JL, Tellier A, and Croll D. The Genome Biology of
Effector Gene Evolution in Filamentous Plant Pathogens. Annu Rev Phytopathol. 2018:**56**(1):21–40.
https://doi.org/10.1146/annurev-phyto-080516-035303

Sarris PF, Cevik V, Dagdas G, Jones JDG, and Krasileva KV. Comparative analysis of plant immune receptor

architectures uncovers host proteins likely targeted by pathogens. BMC Biol. 2016:**14**(1):8.
https://doi.org/10.1186/s12915-016-0228-7

Saur IM, Bauer S, Kracher B, Lu X, Franzeskakis L, Müller MC, Sabelleck B, Kümmel F, Panstruga R,
Maekawa T, et al. Multiple pairs of allelic MLA immune receptor-powdery mildew AVRA effectors argue
for a direct recognition mechanism. eLife. 2019:**8**:e44471. https://doi.org/10.7554/eLife.44471

Segretin ME, Pais M, Franceschetti M, Chaparro-Garcia A, Bos JIB, Banfield MJ, and Kamoun S. Single
Amino Acid Mutations in the Potato Immune Receptor R3a Expand Response to *Phytophthora* Effectors.
Mol Plant-Microbe Interactions®. 2014:**27**(7):624–637. https://doi.org/10.1094/MPMI-02-14-0040-R

Selvaraj M, Toghani A, Pai H, Sugihara Y, Kourelis J, Yuen ELH, Ibrahim T, Zhao H, Xie R, Maqbool A, et
al. Activation of plant immunity through conversion of a helper NLR homodimer into a resistosome (Plant
Biology). https://doi.org/10.1101/2023.12.17.572070

Seong K and Krasileva KV. Computational Structural Genomics Unravels Common Folds and Novel Families in
the Secretome of Fungal Phytopathogen *Magnaporthe oryzae*. Mol Plant-Microbe Interactions®.
2021:**34**(11):1267–1280. https://doi.org/10.1094/MPMI-03-21-0071-R

Seong K and Krasileva KV. Prediction of effector protein structures from fungal phytopathogens enables
evolutionary analyses. Nat Microbiol. 2023:**8**(1):174–187. https://doi.org/10.1038/s41564-022-01287-6

Tamborski J and Krasileva KV. Evolution of Plant NLRs: From Natural History to Precise Modifications. Annu
Rev Plant Biol. 2020:**71**(1):355–378. https://doi.org/10.1146/annurev-arplant-081519-035901

Tamborski J, Seong K, Liu F, Staskawicz BJ, and Krasileva KV. Altering Specificity and Autoactivity of Plant
Immune Receptors Sr33 and Sr50 Via a Rational Engineering Approach. Mol Plant-Microbe Interactions®.
2023:**36**(7):434–446. https://doi.org/10.1094/MPMI-07-22-0154-R

The PyMOL Molecular Graphics System.

Wang J, Hu M, Wang J, Qi J, Han Z, Wang G, Qi Y, Wang H-W, Zhou J-M, and Chai J. Reconstitution and
structure of a plant NLR resistosome conferring immunity. Science. 2019a:**364**(6435):eaav5870.
https://doi.org/10.1126/science.aav5870

Wang J, Wang J, Hu M, Wu S, Qi J, Wang G, Han Z, Qi Y, Gao N, Wang H-W, et al. Ligand-triggered allosteric
ADP release primes a plant NLR complex. Science. 2019b:**364**(6435):eaav5868.
https://doi.org/10.1126/science.aav5868

Yan Y, Tao H, He J, and Huang S-Y. The HDOCK server for integrated protein–protein docking. Nat Protoc.
2020:**15**(5):1829–1852. https://doi.org/10.1038/s41596-020-0312-x

Zdrzałek R, Stone C, De La Concepcion JC, Banfield MJ, and Bentham AR. Pathways to engineering plant
intracellular NLR immune receptors. Curr Opin Plant Biol. 2023:**74**:102380.
https://doi.org/10.1016/j.pbi.2023.102380

Zhao Y-B, Liu M-X, Chen T-T, Ma X, Li Z-K, Zheng Z, Zheng S-R, Chen L, Li Y-Z, Tang L-R, et al. Pathogen
effector AvrSr35 triggers Sr35 resistosome assembly via a direct recognition mechanism. Sci Adv.
2022:**8**(36):eabq5108. https://doi.org/10.1126/sciadv.abq5108