Short Report

Overcoming cryo-EM map anisotropy reveals ALK-cytokine assemblies with distinct stoichiometries

Jan Felix[1,2*], Steven De Munck[1,2], J. Fernando Bazan[3], Savvas N. Savvides[1,2*]

[1]VIB-UGent Center for Inflammation Research, 9052 Ghent, Belgium.

[2] Unit for Structural Biology, Department of Biochemistry and Microbiology, Ghent University, 9052 Ghent, Belgium.

[3] ħ bioconsulting llc, Stillwater, MN, USA.

*Correspondence: savvas.savvides@ugent.be ; jan.felix@ugent.be

1 Abstract

Activation of Anaplastic lymphoma kinase (ALK) and leukocyte tyrosine kinase (LTK) by their
cognate cytokines ALKAL2 and ALKAL1 play important roles in development, metabolism, and
cancer. Recent structural studies revealed ALK/LTK-cytokine assemblies with distinct
stoichiometries. Structures of ALK-ALKAL2 and LTK-ALKAL1 complexes with 2:1 stoichiometry
determined by X-ray crystallography (De Munck *et al.* Nature, 2021) contrasted the 2:2 ALK-ALKAL2
complexes determined by cryo-EM (Reshetnyak *et al.,* Nature, 2021) and X-ray crystallography (Li
et al., Nature, 2021). Here, we show based on reanalysis of the cryo-EM data deposited in EMPIAR-
10930 that over half of the ALK-ALKAL2 particles in the dataset are classified into 2D classes obeying
a 2:1 stoichiometry besides the originally reported structure displaying 2:2 stoichiometry. Unlike
particles representing the 2:2 ALK-ALKAL2 complex, particles for the 2:1 ALK-ALKAL2 complex
suffer severely from preferred orientations that resulted in cryo-EM maps displaying strong
anisotropy. Here, we show that extensive particle orientation rebalancing in cryoSPARC followed
by 3D refinement with Blush regularization in RELION constitutes an effective strategy for avoiding
map artefacts relating to preferred particle orientations and report a 3D reconstruction of the 2:1
ALK-ALKAL2 complex to 3.2 Å resolution from EMPIAR-10930. This new cryo-EM structure together
with the crystal structures of ALK-ALKAL2 and LTK-ALKAL1 complexes with 2:1 stoichiometry
reconciles a common receptor dimerization mode for ALK and LTK and provides direct evidence for
the presence of an ALK-ALKAL2 complex with 2:1 stoichiometry next to the reported 2:2
stoichiometric assembly in the EMPIAR-10930 dataset. Finally, our analysis emphasizes the
importance of public deposition of raw cryo-EM data to allow reanalysis and interpretation.

Main Text

Introduction & Results

Anaplastic lymphoma kinase (ALK) and the related leukocyte tyrosine kinase (LTK) are receptor tyrosine kinases (RTKs) that are activated by binding their cognate cytokines, ALKAL1 and ALKAL2 (also called FAM150A/B or AUGβ/α)[1-3]. Their signaling outputs elicit a range of pleiotropic activities in development and metabolism, and their dysregulated intracellular kinases are keenly pursued as therapeutic drug targets in various cancers[4]. However, the field had long been hampered by a paucity of insights into the intricate folds of ALK and LTK and their cytokine-bound complexes. A triad of recent studies[5-7] have aimed to provide the missing structural and mechanistic framework for the cytokine-mediated activation of ALK family receptors but diverged in their proposed assemblies.

Our study by De Munck *et al.*[5] provided representative assemblies for the entire ALK family by reporting crystal structures of both the ligand-binding extracellular regions of human ALK and LTK (ALK$_{TG}$/LTK$_{TG}$) in complex with ALKAL2 and ALKAL1, respectively (PDB entries 7nwz and 7nx0). These structures revealed how a single cytokine molecule nucleates homodimeric receptor assembly via distinct site 1 and 2 interaction interfaces, creating a key site 3 receptor-receptor contact (**Supplementary Figure 1A**). In contrast, Li *et al.*[6] and Reshetnyak *et al.*[7] have focused only on ALK-ALKAL2 complexes comprising the extracellular ligand-binding fragment of ALK and the membrane-proximal EGFL domains (ALK$_{TG-EGFL}$), and reported assemblies obeying a 2:2 stoichiometry obtained via X-ray crystallography using an ALK$_{TG-EGFL}$-ALKAL2 fusion construct (PDB entry 7LS0) or separate ALK$_{TG-EGFL}$ and full-length ALKAL2 via electron cryo-microscopy (cryo-EM, PDB entry 7n00), respectively (**Supplementary Figure 1B**). In this side-by-side architecture, the main cytokine-receptor interface corresponds to the high affinity site 1 also reported by De Munck *et al.*[5]. An additional ALK-ALKAL2' interaction, covering 250 Å^2 of buried surface area was proposed to be responsible for bridging the two 1:1 ALK-ALKAL2 subcomplexes into an assembly that could support signaling. Intrigued by the 2:2 stoichiometry differing from our 2:1 stoichiometric crystal structures of truncated ALK$_{TG}$ in complex with ALKAL2 (and LTK$_{TG}$ with ALKAL1), we closely examined the cryo-EM data and analysis reported by Reshetnyak *et al.*[7] as deposited in the EMPIAR data base (entry EMPIAR-10930).

At the outset, we identified a number of 3D-classes (Reshetnyak *et al.*, Supplementary Figure 4 in reference [7], classes 1, 3, 7, 8 and 9) resembling the compact 2:1 ALK$_{TG}$/LTK$_{TG}$-cytokine complexes we had reported[5]. To investigate this apparent particle heterogeneity, we leveraged the raw cryo-EM data and uncleaned particle stack as deposited by Reshetnyak *et al.* (access code EMPIAR-10930). An initial data processing strategy following the methods described in Reshetnyak *et al.*, including 8x binning of the data (**Materials and Methods**) revealed that 6 of the 10 most populated classes after initial 2D classification do not correspond to classes shown in the final 2D classification reported in Reshetnyak *et al.* (**Figure 1A**). This prompted us to pursue extensive data processing (**Materials and Methods, Supplementary Figure 2**) of particles present in both sets of 2D classes, namely the ones that correspond to clear 2-fold symmetry (**Figure 1A, orange squares**) and the ones that lack apparent symmetry (**Figure 1A, blue squares**). However, the final refined map of the C1 symmetric particles displayed severe preferential particle orientations leading to a map that is smeared along the Z-direction, a common problem in cryo-EM requiring sample and data collection optimization. Although the resulting map clearly displays two ALK$_{TG-EGFL}$ extracellular domains and one ALKAL2 copy in a similar 2:1 assembly as observed in the crystal structure of the ALK$_{TG}$-ALKAL2

complex (**Supplementary Figure 2**), artefacts resulting from severe anisotropy hampered confident
model building and prohibited detailed structural interpretations.

To overcome the apparent anisotropy presented in the 3D reconstructions towards a 2:1 $ALK_{TG-EGFL}$–ALKAL2 complex, we reasoned that combining exhaustive automatic particle orientation
rebalancing (available in cryoSPARC[8], v4.5) to exclude particles from over-populated direction
bins, followed by 3D refinement with Blush regularization[9] in RELION v5, might result in more
isotropic 3D reconstructions. Blush regularization uses a pre-trained denoising convolutional neural
network on pairs of half-set reconstructions as a smoothness prior and is shown to remove
artifactual densities resulting from uneven angular distributions or streaky features in overfitted
maps[9]. This new regularization approach to resolving vexing cryo-EM structural data was
successfully employed to reveal the distinct binding modes of the anti-CRISPR repressor Aca2 with
DNA and RNA[10].

Starting from a set of selected 3D classes (1.4 million particles) after 3D hetero-refinement
(**Materials and Methods**), we divided these 3D classes in two sets based on the absence or presence
of apparent twofold symmetry. Such grouping resulted in 3D classes resembling either the C1-
symmetric 2:1 ALK_{TG}-ALKAL2 crystal structure reported by De Munck *et al*.[5] (876.806 particles,
~60% of the total), or the deposited C2-symmetric 2:2 $ALK_{TG-EGFL}$-ALKAL2 cryo-EM map reported by
Reshetnyak *et al.*[7] (523.467 particles, ~40% of the total) (**Figure 2A**).

Firstly, we wondered whether our approach would lead to a cryo-EM map of similar resolution and
informational content for the C2 symmetric $ALK_{TG-EGFL}$-ALKAL2 complex with 2:2 stoichiometry as
reported by Reshetnyak *et al.*[7]. Indeed, further data processing in cryoSPARC resulted in a 2.3 Å
resolution map for the 2:2 $ALK_{TG-EGFL}$-ALKAL2 complex (**Materials and Methods**). The resulting map
has a comparable resolution as the deposited map by Reshetnyak *et al. (*EMD-24095, 2.3 Å) and,
importantly, does not contain any particles from 2D classes corresponding to a 2:1 $ALK_{TG-EGFL}$-
ALKAL2 stoichiometry (**Figure 2A**). We note that, similarly as the cryoSPARC refined map reported
in Reshetnyak *et* al., map density for the EGFL domains is less defined than for the RELION refined
map obtained by Reshetnyak *et al.* at slightly lower resolution.

We then pursued analysis of the particle set with apparent resemblance to $ALK_{TG-EGFL}$-ALKAL2
complexes with 2:1 stoichiometry (**Materials and Methods**). The combined use of Orientation
Rebalancing in CryoSPARC and 3D refinement in RELION v5 with enabled Blush regularization,
resulted in a massively improved final map at 3.2 Å resolution (**Figure 1**, **Figure 2A**, **Supplementary
Table 1**) displaying an impressive improvement in isotropy, as evidenced by the 10-fold increase in
the cFAR score (**Figure 2D**). Most importantly, post-processing in RELION yielded a sharpened map
with clear main chain connectivity and density for most side chains, allowing complete model
building of a 2:1 $ALK_{TG-EGFL}$-ALKAL2 complex that includes the membrane-proximal EGFL-like
domains of ALK. While the majority of the 2:1 $ALK_{TG-EGFL}$-ALKAL2 cryo-EM map has a local resolution
to better than 3.5 Å resolution (**Figure 1F**), the C-terminal tips of the EGFL domains are only
resolved at 4 – 4.5 Å resolution pointing to their inherent flexibility. We next wondered whether
3D refinement with Blush regularization would be sufficient to obtain an isotropic map from the
particle set without Orientation Rebalancing. Starting from the same particle set before the first
Orientation Rebalancing step, followed by *Ab-Initio* model generation (3 classes), hetero-
refinement and a final 3D refinement with Blush Regularization of the highest resolution class,
resulted in a map at higher resolution (2.7 Å) but with lower isotropy (cFAR score of 0.1) than the
map obtained by combining Orientation Rebalancing and 3D refinement with Blush regularization

(Figure 3). We note that 3D refinement with Blush regularization of the non-Orientation
Rebalanced particle set (cFAR score of 0.1) is superior in improving map isotropy than NU-
refinement without or with Orientation Rebalancing, which yields maps with cFAR scores of 0.01
and 0.8 respectively (Figure 3). Thus, our analysis demonstrates that the combination of extensive
Orientation Rebalancing and 3D refinement with Blush regularization may serve as a suitable
strategy to overcome severe preferred particle orientations *In Silico*, albeit at the cost of some
moderate loss in resolution.

Discussion

The new structure of ALK$_{TG\text{-}EGFL}$-ALKAL2 with 2:1 stoichiometry presented here is a critical
contribution to the present collection of ALK/LTK-cytokine complexes revealed by X-ray
crystallography and cryo-EM. The elucidation of an ALK$_{TG\text{-}EGFL}$-ALKAL2 complex with 2:1
stoichiometry based on a dataset that was initially reported to only support ALK$_{TG\text{-}EGFL}$-ALKAL2
complexes with 2:2 stoichiometry underscores the power of modern cryo-EM to harness the
conformational and stoichiometric diversity harbored in protein samples. Furthermore, our
analysis emphasizes the importance of public deposition of raw (cryo-EM) data to allow reanalysis
and interpretation. Most importantly, though, this new structure is now the most complete ALK-
ALKAL2 complex with 2:1 stoichiometry to date and agrees closely with the ALK$_{TG}$-ALKAL2 with 2:1
stoichiometry reported by De Munck et al.[5] featuring site 1 (ALK-ALKAL2), site 2 (ALK'-ALKAL2),
and site 3 (ALK-ALK') interaction interfaces (Figure 1). Accordingly, the root-mean-square deviation
(r.m.s.d.) between these two structures is 1.3 Å over 675 aligned Cα-atoms. The site 2 interface
present in the 2:1 ALK:ALKAL2 complex, but not in the 2:2 ALK-ALKAL2 complex, involves the α1
helix of ALKAL2 with a key histidine residue (H100) interacting with S737, Y739 and L760 of ALK'
(Figure 1E). Furthermore, the C-termini of the EGFL domains of the two ALK copies in the 2:1 ALK$_{TG\text{-}}$
$_{EGFL}$-ALKAL2 complex obtained from EMPIAR-10930 are spaced ~30 Å apart (Figure 1), a distance
that would allow facile juxtapositioning of receptor chains and TM helices as already illustrated for
many RTK—cytokine assemblies featuring cytokine-mediated receptor dimerization such as KIT–
SCF[11, 12], CSF-1R–CSF-1[13], Flt3–Flt3R[14], VEGF-VEGFR[15], PDGF-PDGFR[16], EGF–EGFR[17],
HER2–HER3–NRG1β[18] and Insulin bound to insulin receptor (IR) family receptors[19, 20]. The
presence of two distinct stoichiometries of the ALK-ALKAL2 complex among the particles imaged
in the EMPIAR-10930 dataset is intriguing and their role in ALK signaling is currently unclear. One
hypothesis could be that increasing local cytokine concentrations could shift an ALK-ALKAL2
assembly from a 2:1 to a 2:2 stoichiometry, resulting in an increase in the distance between the
membrane proximal EGFL domains from 30 Å tot 90 Å, possibly leading to different signaling
outcomes(Supplementary Figure 3). Interestingly, designed ligands for erythropoietin receptor
aimed at tuning the distance between receptor monomers upon dimerization result in modulation
of erythropoietin receptor signaling and phosphorylation efficiency of downstream adaptors[21].
Finally, structural, biophysical and cellular studies, including full-length ALK-ALKAL2 complexes, will
be required to fully define the possible roles of distinct ALK/LTK-cytokine stoichiometries.

Materials and Methods

Reanalysis of cryo-EM data as reported in EMPIAR-10930

Initial data analysis was carried out in CryoSPARC[8] v3.3.1, and additional steps were performed in either cryoSPARC[8] v4.5 or RELION[22, 23] v5, specifically for the use of Orientation Rebalancing ('Rebalance Orientations' job) and 3D refinement including Blush regularization[9] respectively. Dose-weighted and motion-corrected micrographs were downloaded from https://www.ebi.ac.uk/empiar/EMPIAR-10930/ and were CTF-corrected using patch CTF estimation. After importing the uncleaned particle stack (~18 million particles) from EMPIAR-10930 and extracting the corresponding particles (8x binned), several iterative cycles of reference-free 2D classification were performed to get rid of 'junk' particles, resulting in a 'cleaned' set comprising ~11.8 million particles. This number of particles approaches the total number of particles (~12.4 million) used by Reshetnyak *et al*. Particle unbinning followed by reference-free 2D classification revealed that almost 2/3 of the ten most populated classes do not correspond to classes shown in the final 2D classification represented in Reshetnyak *et al.* (**Figure 1a**). *Ab initio* models generated on a selected subset of most populated classes (4x binned, 1.5 million particles, 3 *Ab Initio* classes, C1 symmetry) followed by a round of heterorefinement revealed one model that clearly corresponds to the 2:2 ALK–ALKAL2 structure reported by Reshetnyak *et al.,* as well as a model resembling the 2:1 ALK–ALKAL2 structure presented in De Munck *et al.* (**Figure 1b & Supplementary Figure 2**). Next, the three *Ab Initio* classes were used together with the cleaned dataset of ~11.8 million particles (unbinned), as input for heterorefinement (C1, 9 classes). Class 1 and 2 of this heterorefinement were grouped together, and selected particles (~1.2 million) were used for 3D classification without alignment (8 classes). Class 5 after 3D classification was selected (334,654 particles) and used as an input for a final Non-Uniform (NU) refinement[24], While the final map shows the presence of two ALK copies and one ALKAL2 copy (with its characteristic three central α-helices), the map suffers from the severe preferential particle orientations of the 2:1 ALK–ALKAL2 particles, as evidenced by the paucity of different views in the 2D classes as well as the Azimuth versus Elevation plot shown in Supplementary Figure 2. Indeed, Orientation Diagnostics, as implemented in the newest version of cryoSPARC[8] (v4.5), pointed to extreme anisotropy corresponding to a cFAR score of 0.03 (**Supplementary Figure 2**). Accordingly, the anisotropy present in the obtained NU-refined map of the 2:1 ALK$_{TG-EGFL}$–ALKAL2 complex prohibited further model building and refinement.

Next, starting from the heterorefinement of 11.849.911 particles, highest resolution 3D classes were grouped in two pools based on the presence or absence of twofold symmetry. Further processing of the particles with twofold symmetry (523.467 particles) included one additional round of 2D classification to remove remaining junk classes, *ab-Initio* model generation (2 classes), hetero-refinement and 3D NU-refinement of the highest resolution class while applying C2 symmetry, and resulted in a 2.3 Å resolution map for the 2:2 ALK$_{TG-EGFL}$-ALKAL2 complex (**Figure 2**). Further processing of the particles without apparent symmetry (876.806 particles) by 2D classification showed that 2D classes mainly consisted of side views of the 2:1 ALK-ALKAL2 complex, with a lack of discernible top views (**Figure 2**). We next used Orientation Rebalancing in cryoSPARC v4.5 to trim overpopulated views in the particle stack using a rebalance percentile threshold of 60, resulting in 599.954 remaining particles. These particles were used as an input for an Ab Initio refinement job (C1, 3 classes) followed by heterorefinement (C1, 3 classes). The highest resolution 3D class (class 1) was used for NU-refinement, followed by another round of Orientation Rebalancing, now using a more stringent rebalance percentile threshold of 5. Retained particles

(142.986) where used for a second NU-refinement and subsequently exported to RELION v5 for a
final round of 3D refinement with enabled Blush regularization. A final 2D classification was
performed on the final particle stack (142.986 particles) to demonstrate the appearance of new
side and top views not seen in previous 2D classification runs (**Figure 2**). Orientation Diagnostics
jobs were run on obtained intermediate 3D reconstructions as well as the final map to monitor
improvements in map isotropy. The final obtained map has a Gold-Standard Fourier Shell
Correlation (FSC) resolution of 3.2 Å at the 0.143 threshold (**Figure 1, Supplementary Table 1**), and
was post-processed either in RELION using a sharpening B-factor of -100 Å^2, or by using
deepEMhancer[25].
208

Model Building & Refinement

Two copies of ALK$_{TG-EGFL}$ and one copy of ALKAL2 were extracted from PDB ID 7n00, and rigid-body
fitted in the sharpened map of the 2:1 ALK$_{TG-EGFL}$-ALKAL2 complex (B-factor: -100 Å^2) using USCF
Chimera[26]. The resulting rigid-body fitted model was subjected to a cycle of manual building in
Coot[27], aided by both the regularly sharpened map and a map sharpened using
DeepEMhancer[25]. Next, the manually-built and corrected model was real-space refined against
the regularly sharpened map in Phenix v1.19.2-4158[28], using global minimization, local grid
search, atomic displacement parameter (ADP) refinement, secondary structure restraints and
Ramachandran restraints. Several cycles of manual building in Coot were followed by real-space
refinement in Phenix. After the last cycle of manual building in Coot, a nonbonded-weight
parameter of 200 was used during refinement in Phenix. The resulting model was subjected to a
final refinement against the two half-maps in REFMAC-Servalcat[29] run within CCP-EM suite
v1.6[30], with automatic weighting applied. The final model has a map-model FSC of 3.20 Å at the
0.5 threshold, calculated using the unfiltered, unsharpened full map (**Figure 1)**. A summary of cryo-
EM data collection, processing, refinement and validation statistics can be found in **Supplementary**
Table 1.
225

Data Availability

The cryo-EM map and accompanying structural model for the 2:1 ALKTG-EGFL-ALKAL2 complex
have been deposited in the EMDB/PDB with accession codes: EMD-51087/9g5i. The particle .star
file obtained after 3D refinement with Blush Regularization in RELION 5 is added to this manuscript
as a Supplementary Data file, and can be used to extract particles after running CTF estimation on
the Motion Corrected and Dose Weighted (*_DW.mrc) micrographs available from EMPIAR-10930
(https://www.ebi.ac.uk/empiar/EMPIAR-10930/).

References

1. Guan, J., et al., *FAM150A and FAM150B are activating ligands for anaplastic lymphoma kinase.* Elife, 2015. **4**: p. e09811.
2. Reshetnyak, A.V., et al., *Identification of a biologically active fragment of ALK and LTK-Ligand 2 (augmentor-alpha).* Proc Natl Acad Sci U S A, 2018. **115**(33): p. 8340-8345.
3. Reshetnyak, A.V., et al., *Augmentor alpha and beta (FAM150) are ligands of the receptor tyrosine kinases ALK and LTK: Hierarchy and specificity of ligand-receptor interactions.* Proc Natl Acad Sci U S A, 2015. **112**(52): p. 15862-7.
4. Hallberg, B. and R.H. Palmer, *Mechanistic insight into ALK receptor tyrosine kinase in human cancer biology.* Nat Rev Cancer, 2013. **13**(10): p. 685-700.
5. De Munck, S., et al., *Structural basis of cytokine-mediated activation of ALK family receptors.* Nature, 2021. **600**(7887): p. 143-147.
6. Li, T., et al., *Structural basis for ligand reception by anaplastic lymphoma kinase.* Nature, 2021. **600**(7887): p. 148-152.
7. Reshetnyak, A.V., et al., *Mechanism for the activation of the anaplastic lymphoma kinase receptor.* Nature, 2021. **600**(7887): p. 153-157.
8. Punjani, A., et al., *cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination.* Nat Methods, 2017. **14**(3): p. 290-296.
9. Kimanius, D., et al., *Data-driven regularization lowers the size barrier of cryo-EM structure determination.* Nat Methods, 2024.
10. Birkholz, N., et al., *Phage anti-CRISPR control by an RNA- and DNA-binding helix-turn-helix protein.* Nature, 2024. **631**(8021): p. 670-677.
11. Yuzawa, S., et al., *Structural basis for activation of the receptor tyrosine kinase KIT by stem cell factor.* Cell, 2007. **130**(2): p. 323-34.
12. Krimmer, S.G., et al., *Cryo-EM analyses of KIT and oncogenic mutants reveal structural oncogenic plasticity and a target for therapeutic intervention.* Proc Natl Acad Sci U S A, 2023. **120**(13): p. e2300054120.
13. Felix, J., et al., *Structure and Assembly Mechanism of the Signaling Complex Mediated by Human CSF-1.* Structure, 2015. **23**(9): p. 1621-1631.
14. Verstraete, K., et al., *Structural insights into the extracellular assembly of the hematopoietic Flt3 signaling complex.* Blood, 2011. **118**(1): p. 60-8.
15. Markovic-Mueller, S., et al., *Structure of the Full-length VEGFR-1 Extracellular Domain in Complex with VEGF-A.* Structure, 2017. **25**(2): p. 341-352.
16. Chen, P.H., V. Unger, and X. He, *Structure of Full-Length Human PDGFRbeta Bound to Its Activating Ligand PDGF-B as Determined by Negative-Stain Electron Microscopy.* J Mol Biol, 2015. **427**(24): p. 3921-34.
17. Huang, Y., et al., *A molecular mechanism for the generation of ligand-dependent differential outputs by the epidermal growth factor receptor.* Elife, 2021. **10**.
18. Diwanji, D., et al., *Structures of the HER2-HER3-NRG1beta complex reveal a dynamic dimer interface.* Nature, 2021. **600**(7888): p. 339-343.
19. Uchikawa, E., et al., *Activation mechanism of the insulin receptor revealed by cryo-EM structure of the fully liganded receptor-ligand complex.* Elife, 2019. **8**.
20. Wang, L., et al., *Structural basis of the alkaline pH-dependent activation of insulin receptor-related receptor.* Nat Struct Mol Biol, 2023. **30**(5): p. 661-669.

21.   Mohan, K., et al., *Topological control of cytokine receptor signaling induces differential effects in hematopoiesis.* Science, 2019. **364**(6442).

22.   Scheres, S.H., *RELION: implementation of a Bayesian approach to cryo-EM structure determination.* J Struct Biol, 2012. **180**(3): p. 519-30.

23.   Kimanius, D., et al., *New tools for automated cryo-EM single-particle analysis in RELION-4.0.* Biochem J, 2021. **478**(24): p. 4169-4185.

24.   Punjani, A., H. Zhang, and D.J. Fleet, *Non-uniform refinement: adaptive regularization improves single-particle cryo-EM reconstruction.* Nat Methods, 2020. **17**(12): p. 1214-1221.

25.   Sanchez-Garcia, R., et al., *DeepEMhancer: a deep learning solution for cryo-EM volume post-processing.* Commun Biol, 2021. **4**(1): p. 874.

26.   Pettersen, E.F., et al., *UCSF Chimera--a visualization system for exploratory research and analysis.* J Comput Chem, 2004. **25**(13): p. 1605-12.

27.   Emsley, P. and K. Cowtan, *Coot: model-building tools for molecular graphics.* Acta Crystallogr D Biol Crystallogr, 2004. **60**(Pt 12 Pt 1): p. 2126-32.

28.   Liebschner, D., et al., *Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix.* Acta Crystallogr D Struct Biol, 2019. **75**(Pt 10): p. 861-877.

29.   Yamashita, K., et al., *Cryo-EM single-particle structure refinement and map calculation using Servalcat.* Acta Crystallogr D Struct Biol, 2021. **77**(Pt 10): p. 1282-1291.

30.   Burnley, T., C.M. Palmer, and M. Winn, *Recent developments in the CCP-EM software suite.* Acta Crystallogr D Struct Biol, 2017. **73**(Pt 6): p. 469-477.

31.   Chen, S., et al., *High-resolution noise substitution to measure overfitting and validate resolution in 3D structure determination by single particle electron cryomicroscopy.* Ultramicroscopy, 2013. **135**: p. 24-35.

Acknowledgments

S.N.S. acknowledges research support from the FWO (grant number G0B4918N) and the Flanders Institute for Biotechnology (VIB, grant number C0101).

Author contributions

J.F and S.N.S. wrote the manuscript with contributions from all authors. J.F performed analysis and processing of the deposited EMPIAR-10930 cryo-EM dataset with contributions from S.D.M. J.F. and S.D.M. performed structural analyses with contributions from S.N.S. and J.F.B.

Competing interests

S.D.M. and S.N.S. are listed as co-inventors on a patent filing EP 21184332.1 with the European Patent Office.

327
328
329   Figure 1: Reanalysis of cryo-EM data from EMPIAR entry EMPIAR-10930 deposited by Reshetnyak *et al.,*
330   *Nature* (2021). **(A)** 10 most populated 2D class averages after performing 2D classification on 11.8 million
331   selected particles. Classes corresponding to 2:1 ALK–ALKAL2 particles are highlighted in blue, while classes
332   corresponding to 2:2 ALK–ALKAL2 particles are highlighted in orange. **(B)** *Ab initio* model generation in
333   cryoSPARC[8] results in classes corresponding to both a 2:1 ALK-–ALKAL2 and 2:2 ALK–ALKAL2 assembly.

Hetero-refinement of the 2:1 and 2:2 ALK–ALKAL2 classes results in the two models displaying 2:1 and 2:2 ALK–ALKAL2 assemblies, respectively. The ALK–ALKAL2 complexes with 2:1 (PDB 7nwz) and 2:2 (PDB 7n00) stoichiometries are shown fitted inside the maps, with ALK colored pink/grey and ALKAL2 colored blue/purple. **(C)** 3D refinement with Blush regularization[9] in RELION v5 of a trimmed particle set after extensive automatic rebalancing of particle orientations as implemented in cryoSPARC 4.5 (see also Materials and Methods & Supplementary Table 1). A front view is shown of a map sharpened in RELION using a B-factor of -100 Å^2, with ALK colored pink/grey and ALKAL2 colored blue. Front, side and top views are shown of a transparent map, sharpened using deepEMhancer, with fitted structural model. **(D & E)** Zoomed-in insets illustrating the quality of map density corresponding to the two major interaction interfaces (site 1 & 2) in the ALK:ALKAL2 complex. Key interacting residues are annotated and shown as sticks. **(F)** Local Resolution Estimation performed in RELION v5 of the final map for the 2:1 ALK-ALKAL 2 complex. **(G)** Gold-standard Fourier Shell Correlation (FSC) plot corresponding to the final map of the 2:1 ALK:ALKAL2 complex. The corrected FSC (purple lines) is calculated after performing correction by noise substitution[31], and the resolution at FSC=0.143 is annotated via a dotted purple line. A map-to model FSC curve calculated using the fitted model and an unsharpened, unfiltered full map is shown (grey line) with the resolution at FSC=0.5 annotated via a dotted grey line.

352
Figure 2: Final processing pipeline of cryo-EM dataset EMPIAR-10930. (A) Final processing steps were
performed in CryoSPARC[8] v4.5 or RELION[22, 23] v5. Post-processing was performed in RELION or using

deepEMhancer[25]. **(B & C)** Gold-standard Fourier Shell Correlation (FSC) plots corresponding to the final
maps of the 2:1 **(B)** and 2:2 **(C)** ALK:ALKAL2 complexes. The corrected FSCs (purple lines) are calculated after
performing correction by noise substitution[31], and the resolution at FSC=0.143 is annotated via dotted
purple lines. **(D & E)** conical FSC summary plots corresponding to the final maps of the 2:1 **(D)** and 2:2 **(E)**
ALK:ALKAL2 complexes, generated via 'Orientation Diagnostics' in cryoSPARC v4.5. **(F)** Azimuth plot showing
the distribution of orientations over Azimuth (X-axis) and Elevation (Y-axis) angles for the particle set
corresponding to the final map of the 2:1 ALK-ALKAL2 complex, obtained after Orientation Rebalancing in
cryoSPARC v4.5 and 3D refinement with Blush regularization[23] in RELION v5. **(G)** Display of the 20 most
populated classes after 2D classification of the final particle stack corresponding to the 2:1 ALK-ALKAL2
complex. New side & top views not observed in initial 2D classification runs are shown as blue squares.
365

Figure 3: Comparison of map (an)isotropy following data processing strategies with or without Orientation Rebalancing and/or 3D refinement with enabled Blush Regularization. **(A)** Processing workflows (1 to 4) of cryo-EM dataset EMPIAR-10930 starting from 876.806 selected particles as outlined in Methods. Workflow 1 does not include Orientation Rebalancing (CryoSPARC v4.5) nor 3D refinement with Blush regularization (RELION v5). Workflow 2 is the same as 1 but with a final 3D refinement with Blush regularization in RELION v5 instead of NU-refinement in CryoSPARC v4.5. Workflow 3 utilizes Orientation Rebalancing in CryoSPARC v4.5 followed by NU-refinement in CryoSPARC, while workflow 4 combines Orientation Rebalancing in CryoSPARC with 3D refinement with Blush regularization in RELION. **(B)** Conical FSC summary plots for the

375   final 3D refinements obtained in processing workflows 1, 2, 3 and 4, generated via 'Orientation Diagnostics'
376   in cryoSPARC v4.5.

378   Supplementary Information

Supplementary Figure 1: Comparison of 2:1 ALK/LTK and 2:2 ALK receptor-ligand complexes. (a) Cartoon
views of structures of ALK-ALKAL2 and LTK-ALKAL1 complexes with 2:1 stoichiometry as determined by De
Munck *et al.*[5] (PDB entries 7nwz and 7nx0) and **(b)** of ALK-ALKAL2 with 2:2 stoichiometry as determined by
Reshetnyak *et al.*[7] and Li *et al.*[6] (PDB entries 7n00 and 7ls0). Putative positions of the EGFL domains
absent in **(a)** are shown as dotted outlines.

Supplementary Figure 2: Initial processing workflow of cryo-EM dataset EMPIAR-10930. **(A)** All initial processing steps were performed in CryoSPARC[8] v3.3.1. **(B)** Gold-standard Fourier Shell Correlation (FSC) plot corresponding to the final map obtained after non-uniform refinement (C1 symmetry). Curves are shown after applying no mask (blue), a loose mask (green), or a tight mask (red) to the two half maps before calculating FSCs. The corrected FSC (purple) is calculated using the tight mask with correction by noise substitution[31], and the resolution at FSC=0.143 is annotated via a dotted purple line. **(C)** conical FSC summary plot generated via 'Orientation Diagnostics' in cryoSPARC v4.5. **(D)** Left: Azimuth plot showing the distribution of orientations over Azimuth (X-axis) and Elevation (Y-axis) angles for the particle set corresponding to the NU-refined map shown in (A). Right: Plot showing Relative Signal amount vs. Viewing Direction.

399 **Supplementary Figure 3: Side-by-side comparison of 2:1 and 2:2 ALK-ALKAL2 complexes present in the**
400 **EMPIAR-10930 cryo-EM dataset.** Models with pdb code 9g5i (this study) and 7n00 (Reshetnyak *et al.*, 2021)
401 are shown as cartoons with ALK/ALK' colored pink/grey and ALKAL2 colored blue in the 2:1 ALK-ALKAL2
402 complex, or blue/purple in the 2:2 ALK-ALKAL2 complex. The distance between the C-termini of membrane-
403 proximal EGFL domains is annotated.

 Supplementary Table 1. Cryo-EM data collection, processing, refinement and validation statistics

	2:1 ALK$_{TG-EGFL}$-ALKAL2
Data Collection (EMPIAR-10930, Reshetnyak *et al.* 2021[7])	
Magnification	15,000
Voltage (kV)	300
Electron exposure (e$^-$/Å^2)	88.4
Defocus range (μm)	0.8 – 1.8
Pixel size (Å)	0.826
Micrographs	13.618
Data Processing (This study)	
Symmetry imposed	C1
Initial particle images (no.)	18,053,705
Final particle images (no.)	142,986
Map resolution (Å)	3.2
FSC threshold	0.143
Map resolution range (Å)	3.0 – 5.5
Refinement (This study)	
Initial model used (PDB code)	7N00
Model resolution (Å)	3.2
FSC threshold	0.5
Map sharpening B factor (Å^2)	-100
Model composition	
Non-hydrogen atoms	5,252
Protein residues	732
B factors (Å^2)	
Protein	165.45
R.m.s. deviations	
Bond lengths (Å)	0.009
Bond angles (°)	1.193
Validation	
MolProbity score	1.86
Clashscore	9.85
Poor rotamers (%)	0.00
CaBLAM outliers (%)	2.64
CC (mask/volume) sharpened, unsharpened	0.76/0.71, 0.80/0.88
EMRinger (sharpened/unsharpened)	2.10/1.51
Ramachandran plot	
Favored (%)	95.04
Allowed (%)	4.96
Disallowed (%)	0.00
Ramachandran Z-score	
Whole	-1.92
Helix	-1.17
Sheet	-1.29
Loop	-1.25