Spatial Transcriptomic Analysis Reveals HDAC Inhibition Modulates Microglial Dynamics to Protect Against Ischemic Stroke in Mice

Kevin Jayaraj[1], Ritesh Kumar[1], Sukanya Shyamasundar[1], Thiruma V Arumugam[2], Jai S Polepalli[1], S Thameem Dheen[1]

[1] Department of Anatomy, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117594

[2] Centre for Cardiovascular Biology and Disease Research, Department of Microbiology, Anatomy, Physiology and Pharmacology, School of Agriculture, Biomedicine and Environment, La Trobe University, Bundoora, Victoria, Australia

Correspondence: S Thameem Dheen   E-mail: antstd@nus.edu.sg

Co-Correspondence: Jai S Polepalli   E-mail: jpolepalli@nus.edu.sg

Key Words

Abstract:

Ischemic stroke significantly contributes to global morbidity and disability through a cascade of neurological responses. Microglia, the immune modulators within the brain, exhibit dual roles in exacerbating and ameliorating ischemic injury through neuroinflammatory and neuroprotective roles, respectively. Despite emerging insights into microglia's role in neuronal support, the potential of epigenetic intervention to modulate microglial activity remains largely unexplored. We have previously shown that sodium butyrate, a histone deacetylase inhibitor (HDACi) epigenetically regulates inflammatory response of microglia after ischemic stroke and this study was aimed to characterize the transcriptomic profiles of microglia and their spatial distribution in the stroke brain followed by HDACi administration. We hypothesized that the administration of HDACi epigenetically modulates microglial activation and a region-specific microglial phenotype in the stroke brain, shifting their phenotype from neurotoxic to neuroprotective and facilitating neuronal repair and recovery in the ischemic penumbra. Utilizing a rodent model of middle cerebral artery occlusion (MCAo), spatial transcriptomics and 3D morphometric reconstruction techniques were employed to investigate microglial responses in critical penumbral regions, such as the hippocampus, thalamus, cortex and striatum following HDACi administration. We found that HDACi significantly altered the microglial transcriptomic landscape involving biological pathways of neuroinflammation, neuroprotection and phagocytosis as well as morphological phenotype, promoting a shift towards reparative, neurotrophic profiles within the ischemic penumbra. These changes were also associated with enhanced neuronal survival and reduced neuroinflammation in specific regions in the ischemic brain. By elucidating the mechanisms through which HDAC inhibition influences microglial function, our findings propose therapeutic avenues for neuroprotection and rehabilitation in ischemic stroke, and possibly other neurodegenerative conditions that involve microglia-mediated neuroinflammation.

Introduction

Ischemic stroke is one of the primary causes of long-term disability and mortality worldwide, presenting a significant public health challenge by exacerbating both health and economic burdens on society (Donkor, 2018; Feigin *et al.*, 2022). Focal ischemic stroke which is characterized by abrupt cessation of blood supply to a specific region in the brain, leads to hypoxia (a deprivation of oxygen), triggering a cascade of deleterious cellular and molecular events such as excitotoxicity, neuroinflammation, and apoptosis. During reperfusion, restoration of blood flow to the infarcted tissue further aggravates tissue injury (Traystman, 2003; Mergenthaler, Dirnagl and Kunz, 2022). Complex, multicellular pathways and interactions dictate mechanisms that mediate neuronal damage post-ischemic injury and highlight the need for novel therapeutic strategies for enhancing recovery and clinical outcomes (Zhang *et al.*, 2019; Wang, Leak and Cao, 2022).

Microglia, resident immune cells of the brain are at the forefront of the body's response to ischemic injury. They exhibit a spectrum of phenotypic changes (Dubbelaar *et al.*, 2018; Wendimu and Hooks, 2022) and transition dynamically between conventional pro-inflammatory (M1) and anti-inflammatory (M2) states depending on spatial and temporal contexts after ischemia. In the stroke brain, microglia initially clear debris and release pro-inflammatory cytokines like TNF-α and interleukins, triggering neuroinflammatory responses crucial for initiating repair but turn deleterious if it progresses without biological checkpoints controlling the response (Wang, Leak and Cao, 2022). Transitioning to an anti-inflammatory phenotype, microglia release cytokines and neurotrophins promoting tissue repair and limiting ongoing damage (Gupta *et al.*, 2018; Zhang *et al.*, 2021; Charlton *et al.*, 2023). The balance between these microglial states and their heterogenous expression profiles are crucial for the prognosis of stroke. Positive modulation of microglial activation is essential in ameliorating neurodegeneration and reperfusion mediated injury, suggesting that targeted interventions aimed at microglia could offer significant therapeutic benefits (Nagy *et al.*, 2020).

Epigenetic mechanisms, such as DNA methylation, histone modification, and RNA-
associated silencing, play pivotal roles in regulating microglial function and response to
ischemic stroke. Histone deacetylases (HDACs) are key enzymes involved in removing
acetyl groups from histones, leading to reduced gene expression that has been explored to
regulate progress of pathology and neuroinflammation in stroke and other inflammatory
conditions (Adcock, 2007; Fessler *et al.*, 2013; Patnala *et al.*, 2017). In the context of
ischemic stroke, such epigenetic modifications influence the phenotypic switching of
microglia between pro-inflammatory and anti-inflammatory states, thus affecting the overall
inflammatory milieu and recovery process (Saw *et al.*, 2020; Qiu, Xu and Zhan, 2021).

Sodium butyrate, a HDAC inhibitor (HDACi), has emerged as a promising epigenetic therapy
for ischemic stroke (Sun *et al.*, 2015). By modulating the epigenetic landscape, HDACi can
shift microglial activation towards neuroprotective states, reducing neuroinflammation and
promoting neuroprotection. This therapeutic potential is further supported by studies
demonstrating its ability to enhance the expression of neurotrophic factors and modulate key
signaling pathways involved in cell survival, neurogenesis, and synaptic plasticity (Patnala *et*
al., 2017; Ziemka-Nalecz *et al.*, 2017; Saw *et al.*, 2020; Li *et al.*, 2024; Xu *et al.*, 2024).

Our research leverages advanced methodologies, including high-resolution imaging, 3D
microglial reconstruction, and spatial transcriptomics, to delve deeper into the effects of
HDACi on microglial dynamics and their transcriptional profile in the stroke brain. By
exploring biological networks relevant to microglia-neuron interaction and the impact of
epigenetic modulation on microglial function, we aim to uncover novel therapeutic targets
that could significantly enhance the functional outcomes in ischemic stroke therapy (Qiu, Xu
and Zhan, 2021; Zhang *et al.*, 2022)

Given the intricate interplay between microglia, neurons, and other glial cells in the ischemic
brain, understanding the precise molecular pathways including those affected by epigenetic
modifications induced by microglial activity, is critical for developing targeted therapies for
stroke. The application of HDAC inhibitors like sodium butyrate offers a unique approach to

modulating the brain's innate immune response, steering it towards pathways that support
neuroprotection, repair, and regeneration. Such insights are crucial for developing targeted
interventions which harness the therapeutic potential of microglial modulation in the context
of ischemic stroke.

The present study demonstrates that HDACi treatment not only reduces infarct volume in
stroke brain but also influences microglial phenotypes towards reparative states, facilitating
neuronal survival and mitigating neuroinflammation. Through analysis incorporating cutting-
edge spatial transcriptomic techniques, we obtain nuanced understanding of microglial roles
in stroke pathophysiology and recovery, paving the way for targeted therapies that can
capitalize on the therapeutic promise of microglial modulation.

Methods

Ethics Statement

Adult male C57Bl/6J mice, 12-15☐weeks old were acquired (InVivos Pte Ltd, Singapore)
and housed under controlled temperature, humidity, light and dark cycles at the NUS
Comparative Medicine (CM) Vivarium. Mice were provided access to standard food pellets
and water ad libitum. Animal work was performed in accordance with the Institutional Animal
Care and Use Committee (IACUC) through protocol NUS/IACUC/R19-0596 approved for the
experiments performed with animal(s) and tissues in this study.

Surgical Ischemic Stroke Modelling and HDACi treatment in Mice

MCAo modelling was performed as previously described (Patnala *et al.*, 2017) using adult
male C57BL/6J mice (12-15 weeks old, 26-29g). The mice were anesthetized with
isofluorane in an induction chamber. Ophthalmic lubricant (Duratears 3.5g, Alcon, USA) was
applied to prevent corneal drying, and body temperature was monitored and maintained at
37.0 ± 0.5°C with a heating pad. The neck of the mice was shaved, disinfected with 70%

ethanol and povidone-iodine solution, and a midline incision was made to expose the

common carotid artery (CCA), internal carotid artery (ICA), and external carotid artery (ECA).

The CCA and ICA were clamped, and the ECA was cauterized and retracted. A Doccol 6-0

monofilament nylon suture (catalog no.: 602123PK10, Doccol Corporation, USA), (diameter

of 0.21-0.23 mm), was introduced through the ECA and advanced into the ICA to occlude the

MCA. Occlusion was maintained for 60 minutes, after which the suture was withdrawn to

allow reperfusion. The ECA incision was cauterized, and the microvascular clips were

removed to restore blood flow through the CCA and ICA. The surgical site was sutured, and

the animals were allowed to recover from anaesthesia under observation. In sham group

mice, surgical reflection of muscles to reveal CCA bifurcation was performed without

disturbing the vasculature to affect cerebral blood flow. Sodium butyrate (HDACi) was

administered intraperitoneally (I.P.) to the mice (Cat. no. B5887-1G, Sigma-Aldrich) at a dose

of 300 mg/kg bodyweight. Two doses of NaBu were administered, 1-hour prior ischemia and

6-hours post reperfusionas described previously (Patnala *et al.*, 2017). Mice were housed a

temperature-controlled environment with a 12-hour light/dark cycle and provided food and

water ad libitum. Motor deficit after surgical modelling was evaluated through preliminary

neurological deficit scoring (pNDS). Severity of behavioural deficit and rescue was plotted for

each functional parameter of circling, latency to move home cage length to enrichment shell,

spasticity/paralysis and allodynia (n=3 in Sham and MCAo and n=4 in MCAo+HDACi

groups) (Garcia *et al.*, 1995; Li *et al.*, 2000). Animals were regularly monitored for adverse

effects or complications, ensuring compliance with guidelines for the care and use of

laboratory animals, aimed at minimizing suffering and the number of animals used. 24 hours

post-stroke induction, mice were euthanized with an intraperitoneal overdose of sodium

pentobarbital (150 mg/kg) and absence of pedal and corneal reflexes was confirmed prior to

harvest of brain tissue. For histological and immunofluorescence analyses, mice were

perfused with ice-cold 4% paraformaldehyde (PFA) and post-fixed at 4°C for 24 hours prior

to histological processing. For spatial transcriptomic analyses, brain tissue was processed

as per 10x Genomics Visium fresh frozen tissue preparation guide CG000240 - Rev E.

TTC Staining of mouse brain tissue and evaluation of tissue damage

For 2,3,5-Triphenyltetrazolium chloride (TTC) staining, fresh brain tissue was sectioned into 2 mm coronal slices using a brain matrix. The brain slices were submerged in cold PBS to maintain tissue viability. Freshly prepared 2% solution of TTC (Cat. no. T8877, Sigma-Aldrich) was added to brain sections in a 24-well plate and incubated at 37°C for 30 minutes in a dark, humidified incubator. Viable tissue stained red, while infarcted areas remained pale or white. Post-incubation, sections were rinsed with PBS and transferred to a clear, flat transparent sheet for digital imaging on a flatbed scanner (CanoScan LiDe 400, Canon). Image processing and analysis of TTC-stained brain sections was performed by thresholding scanned images to greyscale and extrapolating pixel values (Goldlust *et al.*, 1996).Scanned images were loaded and processed using R packages to calculate and distinguish viable (non-infarcted) and non-viable (infarcted) area in brain tissue. Viable and non-viable areas were calculated in square millimeters based on pixel counts and known image resolution. Statistical analysis, including t-tests and ANOVA, was performed to compare the groups (n=3).. The results were then imported into Microsoft Excel for visualization.

Histological Processing of Mouse Brain Tissue Sections

Harvested brain tissues were post-fixed in 4% paraformaldehyde and dehydrated through a graded series of ethanol baths and clearene. Dehydrated and cleared samples were then incubated in molten paraffin wax, embedded and allowed to solidify at room temperature. Solidified wax blocks were mounted onto a microtome, trimmed, and sectioned at 7μm thickness. Sections were floated in a warm water bath, transferred onto microscope slides, and stained with toluidine blue. Routine staining protocols were performed involving deparaffinization in clearene, rehydration through descending ethanol series, staining in toluidine blue followed by rinsing, dehydrating ethanol series and tissue clearing prior to coverslipping slides with mounting medium (S302380-2, Dako North America Inc.). For

cryosectioning, PFA-fixed brain tissues were cryoprotected in 30% sucrose until fully

saturated, embedded in OCT compound, and rapidly frozen using dry ice or pre-cooled

isopentane. Frozen OCT-embedded tissue blocks were sectioned at -20°C to -25°C in a

cryostat (Leica Instruments GmbH, Germany), with sections cut at 30 to 50 microns,

collected onto chilled slides, and stored at -80°C in sealed containers with desiccants until

further use.

Immunofluorescence and Morphometric analysis

Mouse brain tissues were fixed with 4% paraformaldehyde, cryoprotected in 30% sucrose

and embedded in OCT compound. Mouse brains were sectioned at 30 or 50 micron

thickness on a cryostat (Leica Instruments GmbH, Germany). The sections were

permeabilized with 0.3% Triton X-100   and non-specific binding sites were blocked by

incubating sections in 5% normal donkey serum (D9663-10ML, Sigma-Aldrich) or 1% bovine

serum albumin (126615-25MLCN, Merck). The sections were then incubated with primary

antibodies (antibody list in table below) diluted in blocking solution, overnight at 4°C. After

rinsing with PBS, sections were incubated with fluorescence-conjugated secondary

antibodies for 1 hour at room temperature, protected from light. Following this, sections were

rinsed with PBS and mounted with anti-fade fluorescence mounting medium (ab104135,

Abcam). The stained sections were imaged using a confocal microscope (FluoView 1000

and FluoView 3000, Olympus Corporation, Japan) at 10x, 20x, 60x and 100x objectives. Z-

stacks were captured at 60x and 100x objectives with a z-plane of 20-30µm with z-sections

of 0.5µm. Microglia-labelled confocal, Z-stacked imaging datasets were imported into

IMARIS v9.5 software for 3D reconstruction and analysis of microglial morphology (n=3).

The 'Surpass' view was used to inspect the z-stack, and a 3D volume rendering of the z-

stack was created using the 'Volume' button. For morphometric evaluation and analysis, the

'Surfaces' function was employed by selecting the appropriate marker channel (IBA1 for

microglia), setting the 'Surface Detail Level' and 'Background Subtraction', and using the

'Thresholding' slider to separate cells from the background. The surface creation was

finalized to generate surfaces encompassing the 3D structures of the labelled cells.

Morphometric analysis was exported in the 'Statistics' tab, quantifying parameters such as

volume and surface area. Images and videos of the 3D reconstructions were exported using

the snapshot and animation features in IMARIS v9.5. Comparisons of these parameters

between groups (Sham, MCAo, and HDACi-treated) were conducted to evaluate the effects

of ischemic stroke and HDACi treatment on microglial morphology.

Primary Antibodies

Antibody	Cat no.	Dilution	Source	Host	Clonality	Application
IBA1	ab48004	1:450	Abcam PLC, Cambridge, UK	Goat	Polyclonal	IF
NeuN	ab177487	1:600	Abcam PLC, Cambridge, UK	Rabbit	Monoclonal	IF
NeuN	ab177487	1:100	Abcam PLC, Cambridge, UK	Rabbit	Monoclonal	Spatial Transcriptomics
GFAP	ab4674	1:70	Abcam PLC, Cambridge, UK	Chicken	Polyclonal	Spatial Transcriptomics
H3K27me3	9733	1:400	Cell Signaling Technology, MA, USA	Rabbit	Monoclonal	IF
H3K4me3	9751	1:400	Cell Signaling Technology, MA, USA	Rabbit	Monoclonal	IF

216 **Secondary Antibodies**

Antibody	Cat no.	Dilution	Source	Host	Conjugate	Application
Donkey Anti-Goat (Alexa Fluor 488)	ab150133	1:600	Abcam PLC, Cambridge, UK	Donkey	Alexa Fluor 488	IF
Donkey Anti-Rabbit (Alexa Fluor 647)	ab150063	1:600	Abcam PLC, Cambridge, UK	Donkey	Alexa Fluor 647	IF
Donkey Anti-Rabbit (Alexa Fluor 647)	ab150063	1:100	Abcam PLC, Cambridge, UK	Donkey	Alexa Fluor 647	Spatial Transcriptomics
Goat Anti-Chicken (Alexa Fluor 488)	ab150169	1:100	Abcam PLC, Cambridge, UK	Goat	Alexa Fluor 488	Spatial Transcriptomics
DAPI	D9542-10MG	1:2000	Sigma-Aldrich			Nuclear staining (spatial transcriptomics)
Hoechst 33342	R37605	2drops/ml	Invitrogen			Nuclear staining (IF)

Spatial Transcriptomics Pipeline

Fresh-frozen brain tissue samples from the three groups (n=3) were sectioned at 10 μm

thickness using a cryostat (Leica Instruments GmbH, Germany). Tissue optimization was

performed with TO slides (part no. 3000394, 10x Genomics, California, USA) and reagents

as per 10x Genomics user guide CG000238-RevE and involved testing permeabilization

durations, with six minutes selected based on tissue morphology preservation and RNA

quality, assessed with a Bioanalyzer (Bioanalyzer 2100 system, Agilent, Santa Clara, CA,

USA) (Figure 4 Supplementary Figure 1, 2). The sections were placed on Visium Spatial

Gene Expression slides (part no. 2000233, 10x Genomics, California, USA) ensuring precise

alignment and adhesion to capture areas for spatial mapping. Immunofluorescence staining

(as per 10x Genomics user guide CG000312-RevD) was performed using antibodies against

neuronal nuclei (NeuN) and glial fibrillary acidic protein (GFAP), with DAPI for nuclear

staining, to correlate spatial gene expression data with specific cell types and anatomical

features. Antibody concentrations and incubation times were optimized through preliminary

titration experiments as advised by 10x standard protocols, to ensure specificity and

maximal signal-to-noise ratios. Following immunofluorescence and tissue preparation, in situ

reverse transcription on the Visium Spatial slides was performed to convert localized mRNA

into cDNA while retaining spatial coordinates using unique molecular identifiers (UMIs) and

spatial barcodes. Library preparation, conducted according to Visium protocols (10x

Genomics user guide CG000239-RevF), included second-strand synthesis, adaptor ligation,

and amplification steps. Libraries were quantified and validated for quality using Novogene's

(NovogeneAIT Genomics, Singapore Pte Ltd) sequencing pipeline for their Illumina platform,

to ensure suitability for high-throughput sequencing. Sequencing was performed to a depth

of 50 million reads to cover the complexity of the brain transcriptome and ensure spatial

resolution of gene expression patterns. Sequenced spatial libraries were processed using

10X Genomics SpaceRanger 2.0.1, aligning reads to the reference genome and attributing

them to specific spatial locations on tissue sections based on their spatial barcodes. Spatial

transcriptomic data was exported from Loupe Browser 6.0 into Microsoft Excel for

consolidation, upload and visualization; systematically organizing metrics such as Log2 Fold

Change (Log2FC) values for genes of interest across different tissue sections and

experimental conditions. Data wrappers such as venny (Oliveros, 2007), S.R. Plot (Tang *et*

al., 2023) and heatmapper.ca (Babicki *et al.*, 2016) were utilized for downstream data

processing and visualization. The Kyoto Encyclopedia of Genes and Genomes (KEGG), the

Database for Annotation, Visualization, and Integrated Discovery (DAVID), and Gene

Ontology (GO) profiler were employed to enrich and categorize gene expression data within

biological pathways, functional annotations, and cellular component. Data was visualized

through S.R Plot (Tang *et al.*, 2023) with clusterprofiler and pathview packages in the form of

enrichment bubble and ontology bar plots. Venny, an interactive tool for comparing lists

(Oliveros, 2007) was utilized to generate comparative lists of upregulated, downregulated

and rescued genes across the biological groups with spatial segregation of microglial

transcriptomes. Heatmaps were generated using Heatmapper.ca (Babicki *et al.*, 2016) to

visualize Log2FC data exported from MS Excel organized into selected pathways from

enrichment analyses, providing a graphical representation of gene expression variations

across different brain regions and treatment groups.

Statistical analyses

For infarct area evaluation from TTC stained brain slices, one-way ANOVA

(F-statistic=32625.11, p=2.74 $\times$ 10^{-9}) and independent two-sample t-tests ($<1 \times 10^{-6}$) were

performed on the calculated non-viable (infarcted) areas from all three biological groups

(n=3). The high F-statistic value confirms that the variability between group means is much

larger than the variability within groups while the t-tests confirm the findings from one-way

ANOVA. Similarly, independent two-sample t-tests revealed significant differences (<0.05) in

IMARIS morphometry data focusing on microglial surface area; microglial volume exhibited

significant differences (<0.005) as depicted in data between Sham and MCAo, as well as

between MCAo and MCAo+HDACi groups. Spatial transcriptomics data processed using

proprietary 10x software, SpaceRanger 2.0.1 and Loupe Browser 6.0 utilizes a statistical

framework based on sSeq for differential expression analysis (Yu, Huber and Vitek, 2013),

Moran's I for spatial enrichment (Getis, 2007) and Benjamini-Hochberg correction for

controlling false discovery rates (p<0.05) natively in the transcriptomics pipeline as

described in 10x Genomics repositories. Online data wrappers were fed downstream data

exported from Loupe Browser 6.0 by filtering for genes limited to the selected pathways

within the microglial/neuronal spatiotranscriptomes by MS Excel's VLOOKUP function and

data was visualized in the form of heatmaps (heatmapper.ca), enrichment bubble plots,

combined Gene Ontology bar plots, CNETplots for KEGG pathways-targets enriched (S.R.

Plot) and Venn diagrams for dysregulated gene expression across microglial

spatiotranscriptomes (Venny).

Results

HDACi treatment reduces ischemic infarct volume in the brain and enhances
preliminary locomotor function of mice post-ischemic insult.

HDACi was administered intra-peritoneally to mice undergoing 1-hour ischemia and 24-hour

reperfusion periods in two doses. The first dose at 1-hour pre-ischemic insult to prime cells

and the subsequent dose post 6-hours of ischemic insult to effectively influence reperfusion

mediated injury. As reported earlier, MCAo was performed for 1h and reperfusion post-

ischemia was done for 24h prior to euthanasia (Fig.1A) (Patnala et al., 2017). Neurological

viability and infarct volume were assessed through toluidine blue (Nissl) staining and TTC

(2,3,5-Triphenyltetrazolium Chloride) (Fig.1B-C). From the staining, the extent of ischemic

penumbra at brain regions of interest (ROI)'s such as hippocampus, striatum, thalamus and

the cortex were identified to proceed with assessment of microglia-neuron interaction in

regions (Fig.1B). Evaluation of basic locomotor activity in mice prior to euthanasia (Fig.1D,

Fig.1 Supplementary Video 1) showed circling contralateral to the infarct, and reduced

mobility in the MCAo groups, while HDACi-treated mice did not exhibit circling and showed

reduced latency to move within the home cage towards enrichment shell. Substantive

restoration of viable neuronal soma was observed (Fig.1E-F) 24-hour after administration

HDACi. Taken together, the findings suggest a) successful modelling of ischemia in the

mouse brain; along with identification of regions of interest (ROIs) (hippocampus, thalamus,

striatum and cortex) in the ischemic penumbra, presenting as potential areas to target

reperfusion mediated neuronal damage. b) Gross evaluation of viable brain tissue and

preliminary behavioural performance of HDACi administration after MCAo showcases a

potential restorative effect, substantiating a need for further assessment of microglial

contribution to ischemia-reperfusion mediated neuroinflammation as well as neuroprotection.

HDACi ameliorates neurodegeneration in brain nuclei after ischemic stroke and

inhibits microglial aggregation.

The hippocampal CA3-dentate gyrus are particularly vulnerable regions of ischemic insult

(Einenkel and Salameh, 2024). Histological evaluation of toluidine blue stained sections of

the hippocampal CA3-dentate gyrus for neuronal and Nissl body aggregation, reveals a

degree of neuroprotection after HDACi administration (Fig.1E); preservation of Nissl bodies

correlates with enhanced neuronal survival (Smith et al., 2012), substantiating that HDACi

confers neuroprotection in vulnerable regions of the mouse brain. Confocal images (Fig.1F)

showing immunofluorescence expression of NeuN and IBA1 in neurons and microglia

respectively, in the cortex surrounding infarct periphery indicate recruitment of microglia in

stroke brain while there was a reduction in recruitment and alteration of microglial

morphology in stroke brain following HDACi treatment; implying a tempered inflammatory

response that mitigates further neuronal damage by fostering a more conducive environment

for recovery (Lee et al., 2014).

HDACi treatment inhibits ischemia-induced microglial phenotype in mouse brain

Microglial cells that are ramified with slender processes in the sham mouse brain exhibit

amoeboid and reactive phenotypes in the stroke brain (Fig.2A). Upon HDACi administration,

microglial phenotypes shift towards a sham spectrum ramified phenotype with long and

slender processes (Fig.2A). Quantitative morphometric analysis by 3D-reconstruction of

thick tissue sections generated using IMARIS image analysis software (v9.5) reveals that

microglial surface area decreases while intracellular volume is increased in the stroke brain

compared to sham brain microglia (Fig.2B-C). However, HDACi treatment reverses the

stroke microglial phenotype closer to sham mice, suggesting that HDACi inhibits ischemia-

induced phenotypic changes in microglia (Fig.2B-C). The distinct microglial morphology

observed in this study represents transcriptomic profiles that may influence their role in

inflammatory response and neuroprotection through the course of pathophysiology after

ischemic stroke.

339   HDACi inhibits ischemia-induced increase in nuclear expression of histone marks

340   H3K27me3 and H3K4me3 in microglia

Phenotypic heterogeneity of microglia in pathological conditions is regulated by epigenetics,

and various signaling molecules and transcription factors (Wang *et al.*, 2021). We have

previously shown that acetylation marks such as H3K9ac regulate the inflammatory

response of microglia in the stroke brain (Patnala *et al.*, 2017). In the present study, we have

analyzed the expression pattern of methylation marks such as H3K27me3 and H3K4me3 to

understand molecular mechanisms contributing to microglial phenotypic heterogeneity in the

ischemic brain. The nuclear expression of H3K27me3 (Fig.3A-B) and H3K4me3 (Fig.3C)

appeared to be remarkably elevated in microglia of the stroke brain. Upon HDACi

administration, microglia exhibit a more ramified phenotype with inhibition in nuclear

expression of H3K27me3 and H3K4me3, similar to expression patterns of sham brain

microglia suggesting that microglial phenotypic heterogeneity is influenced by histone mark

status. Interestingly, we also found neuronal cells in the hippocampal CA2-CA3 gyri (Fig.3A-

B) exhibiting an inverse H3k27me3 expression pattern to microglia, suggesting an epigenetic

feedback mechanism across microglia-neuron regulatory network of histone status that

warrants further study.

357 Mapping of spatial transcriptome of microglia in ischemic mouse brain following

358 HDACi administration

The spatial transcriptome profile of microglia in response to ischemic injury and HDACi

administration was analyzed using the 10x Genomics Visium (FF) platform. Fresh frozen

brain slices were immunolabelled and imaged for neurons (NeuN, red) and astroglia (GFAP,

green) along with DAPI (blue) on GEX Visium slides prior to library preparation for the spatial

overlay (Fig.4A-B). Immunoflourescence imaging and proprietary Loupe Browser v6.0

cluster overlay with key ROIs, namely the hippocampus, thalamus, striatum, and cortex were

marked for post processing (Fig.4A). Stroke brain tissue reveals a loss of cellular

heterogeneity, and distinct organizational patterns present in sham brain tissue, suggesting

the disruption of cellular microenvironments and signaling networks upon ischemic insult

(Fig.4B-C). Preliminarily, the spatial transcriptomic data revealed 19 distinct cell populations

based on marker gene expression across the three experimental groups (Fig.4F) with a

Uniform Manifold Approximation and Projection (UMAP) plot demonstrating the spatial

distribution of these populations alongside the spatial overlay (Fig.4E, 4C). To specifically

isolate the microglial transcriptome, a sub-clustering approach through feature selection and

thresholding in loupe browser v6.5.0 was employed (Fig.4D) by leveraging a list of microglia

signature genes identified in recent studies (Gerrits *et al.*, 2020; Quintana *et al.*, 2022). This

allowed for the extraction of microglia-enriched spots from brain co-ordinates to generate

distinct microglial spatiotranscriptomic datasets within the four penumbra ROIs.

378 HDACi alters microglial spatiotranscriptome profiles following ischemic stroke

Differential expression analysis and pathway enrichment studies were carried out to

investigate microglial spatiotranscriptomics and their functional implications. The anatomic

regions of interest on the overlay of spatiotranscriptomic spots, focusing on the ischemic

penumbra has been demarcated using loupe browser v6.0 (Fig.5A). Four-set Venn diagrams

were drawn to identify region-specific, unique and commonly upregulated (Fig.5B)

/downregulated (Fig.5C) /rescued (Fig.5D) microglial genes across the ischemic penumbra

in the hippocampus, thalamus, striatum and cortex. Among the upregulated microglial genes

across the four regions of interest, 22% of genes (3544) was found to be commonly

upregulated after HDACi administration (Fig.5B). On the other hand, only 1.5% (190) of

microglial genes was found to be commonly downregulated across the four regions of

interest (Fig.5C). Additionally, the datasets of differentially expressed genes (both

up/downregulated) were processed to obtain microglial genes that were modified or rescued

(closer to sham expression levels) (Fig.5D) following HDACi treatment.

The dysregulated microglial genes were subjected to KEGG pathway enrichment (Fig 5E)

and Gene Ontology (GO) (Fig.5F) analyses. Pathway analysis yields enrichment of key

functional pathways pivotal for neuroinflammatory response, mediating neurotrophic support,

and facilitating phagocytosis (Fig.5E) alongside other pathways that may be relevant to

microglial responses after ischemic stroke (Fig.5 Supplementary Figure 1). Regulatory

signaling pathways such as PI3K-Akt signaling (PIK3CA, PIK3CB, AKT1, AKT2, AKT3,

PTEN), FC gamma R-mediated phagocytosis (FCGR1, FCGR3, FCGR4, SYK, and

PIP5K1C), MAPK signaling (DUSP1, DUSP4, DUSP6, ERK1-2, cJUN, UCP4), TNF

signaling (TNF, NFKB1, RelA), and TGFβ signaling (TGFBR1, TGFBR2, SMAD2, SMAD3,

SMAD4) pathways shed light on the transformative influence of epigenetic regulators on

microglial functionality in the post-stroke brain environment. The genes from these pathways

were found to be modified closer to sham expression levels in each region of interest within

the transcriptome of microglia after HDACi administration.

Thus, HDACi treatment results in a significant alteration of the microglial transcriptomic

landscape, with a spatial context. Pathway-enrichment (Fig.6 Supplementary File 1) and GO

analyses conducted for the microglial genes of hippocampus, thalamus, striatum, and cortex

(Fig.6 Supplementary Figure 1A-1B) revealed that HDACi alters, and rescues expression of

several key pathways involved in neuroprotection, inflammatory response and phagocytosis

after ischemic stroke (Fig.6A,6B,6C,6D). These analyses of microglial spatiotranscriptomes

shed light on the emphasis of these pathways describing their big-picture and region-specific

roles after ischemic insult. Heatmaps visualize the expression pattern of rescued transcripts

across the functionally relevant pathways, alongside radar plots delineating the top 50

rescued targets from these pathways in each microglial spatiotranscriptome

(Fig.6A,6B,6C,6D). Confocal microscopy of microglia in the hippocampus (from Fig.2B), can

be corroborated with microglial spatiotranscriptome to draw parallels between HDACi-

modified microglial transcriptome <> microglial phenotypic heterogeneity in the hippocampus

(Fig 6A <> Fig.2B).

The spatiotranscriptomic profile of microglia across groups emphasizes the impact of HDACi

on microglial gene expression involved in neuroprotection. We also analyzed if HDACi

treatment improves neuroprotection by influencing the neuronal spatiotranscriptome in the

penumbra following ischemic stroke. The HDACi- modified spatiotranscriptome of penumbral

neurons reveal a distinct inverse expression pattern across groups in enriched pathways

(Fig.7B,7C), between ischemia-affected and HDACi treated mouse brains (Fig.7A) with

broader implications for neuroprotection, synaptic remodeling, and plasticity. This suggests a

granular modulatory effect of HDACi that is potentially influenced by the local

microenvironment and neuronal signaling to remodel microglial expression profiles.

The findings highlight the spatial and molecular heterogeneity of microglial responses to

ischemic injury and HDAC inhibition, emphasizing the therapeutic capabilities of epigenetic

modulation in ischemic stroke. The identified pathways and rescued genes provide a robust

framework for further exploration of microglial-neuron interactions and the development of

targeted therapies.

Discussion

Recent advancements in spatial transcriptomics analysis have enriched our ability to dissect

stroke pathophysiology by providing spatial resolution to gene expression patterns in

neurons and glial cells including microglia. In the current study, integration of spatial

transcriptomics with advanced imaging and reconstruction techniques has offered a

comprehensive view of microglial dynamics in the stroke brain. The MCAo model

recapitulates human physiology to facilitate investigating microglial responses within the

ischemic brain (Chen *et al.*, 2009; Ansari *et al.*, 2011), in particular the penumbra regions

including the hippocampus, thalamus, striatum and cortex. In the MCAo model, HDACi

treatment improves sensory-locomotor deficits inflicted on mice post-ischemic stroke as

observed in certain behavioural paradigms prior to euthanasia. An extended battery of

behavioural tests were not incorporated in the experimental design due to the 24-hour time

point that was optimal for capturing a snapshot of reperfusion-mediated neuroinflammation

in the stroke brain. As functional improvements upon HDAC inhibition, particularly sodium

butyrate has been described previously in rodent MCAo models (Zhou *et al.*, 2021), this

study was designed to observe the microglial transcriptome specifically during reperfusion in

the brain's ischemic penumbra. Microglial activation in the stroke brain appears to influence

the neuronal transcriptome, which is substantiated by the dysregulated transcriptional

responses observed in microglia and neurons of the stroke brain. This altered microglial

response, which is regulated by various signalling molecules and epigenetic mechanisms,

can be ameliorated through epigenetic intervention as observed in the present study.

The Influence of Transcription Factors on phenotypic heterogeneity of Microglia

Microglial cells are dynamic and exhibit region- and time-dependent phenotypic

heterogeneity in response to neuropathological conditions such as neuroinflammation and

neurodegeneration including ischemic stroke (Hickman *et al.*, 2018; Li *et al.*, 2020). In the

current study, differential expression of microglial spatiotranscriptomes associated with
phenotypic heterogeneity of microglia in response to ischemic stroke, particularly under the
modulation of HDAC inhibition was observed. This variation in microglial phenotypes
elucidates the interplay of transcription factors, cell-specific receptors and microglia-neuron
interaction; outlining the complexity behind microglial polarization such as neuroprotective or
neurotoxic phenotypes in the post-stroke brain (Olah *et al.*, 2011; Haroon, Miller and
Sanacora, 2016; Dubbelaar *et al.*, 2018).

In our microglial spatiotranscriptomic analysis, ischemic stroke was found to alter key
signalling molecular pathways involved in neurotrophic support, phagocytic activity and
neuroinflammation including PI3K-Akt signaling, FC gamma R-mediated phagocytosis,
MAPK signaling, TNF signaling, and TGFβ signaling pathways and transcription factors
regulating immune response such as NF-κB, STAT1/STAT3, Nrf, and PPAR family (Yang *et*
al., 2018; Simpson and Oliver, 2020; Li *et al.*, 2024). Epigenetic mechanisms such as post-
translational histone modifications provide an additional layer of regulation to elicit cell-
specific and stimuli specific immune responses (Allis and Jenuwein, 2016) in the stroke
brain.

Post-translational histone modifications determine basic transcriptional rates and the
magnitude of gene expression (Bannister and Kouzarides, 2011). Among the two major
histone modifications namely histone acetylation and histone methylation, histone
acetylation is associated with increased accessibility to transcriptional machinery leading to
increased gene expression (Marmorstein and Zhou, 2014), whereas histone methylation is
more complex in regulation of gene expression through repression. Different histone
modification patterns of specific marks at different sites have been shown to play distinct
biological roles; this prompted us to analyze the spatial transcriptome of microglia distributed
in infarct penumbra of the stroke brain upon HDACi administration to understand the
epigenetic control of microglia-mediated neuropathological outcomes in the stroke brain.

Modification in histone acetylation and deacetylation status, is maintained by histone
acetyltransferase (HATs) and histone deacetylase (HDACs) enzymes respectively (Yang and
Seto, 2007). Several studies have shown that inhibition of HDAC exerts immunomodulatory
effects, specifically suppressing pro-inflammation and inducing anti-inflammatory signaling
pathways (Adcock, 2007; Kulthinee *et al.*, 2022). We have previously demonstrated the
HDACi-specific immunosuppression of pro-inflammatory mediators (TNFα and NOS2) and
upregulation of IL10, an anti-inflammatory mediator in microglia in ischemic stroke mice
model exerting neuroprotection (Patnala *et al.*, 2017).

Dynamic regulation of repressive histone methylation marks, H3K27me3 and H3K4me3
within microglia play a significant role in their response to ischemic injury and subsequent
recovery processes (Cheray & Joseph, 2018; Tang et al., 2014). In the present study,
markedly elevated expression of these markers were observed in microglia that are
distributed in the penumbra of ischemic brain. The increased nuclear expression of these
histone marks in microglia appears to reflect a response to the ischemic environment-
induced transcriptional activity which may be associated with changes in migration,
phagocytosis and phenotypic heterogeneity inferred through the spatial transcriptome of
microglia. Remarkably, this ischemia-induced expression of H3K27me3 and H3K4me3 was
found to be reverted to near sham levels in microglia upon HDACi treatment, suggesting
epigenetic regulation of H3K27me3 and H3K4me3 downstream pathways that influence
recovery mechanisms.

Further HDACi-induced suppression of H3K27me3 in microglia of ischemic brain may be
associated with activation of its upstream targets such as Enhancer of Zeste Homolog2
(EZH2) (Suppl.tab.1_averaged_MicroglialLog2FC_10X.xslx) a catalytic subunit of the
Polycomb Repressive Complex 2 (PRC2) (Ayata *et al.*, 2018) which has been shown to play
important roles in the epigenetic regulation of phagocytic clearance and pruning of synaptic
activity in microglia (Hong, Dissing-Olesen and Stevens, 2016; Ayata *et al.*, 2018). Further,
since the PRC2 complex, including EZH2 has been shown to regulate TNFα and TGFβ
signalling pathways (Penas and Navarro, 2018; Chen *et al.*, 2019), the increase in EZH2

expression post HDACi treatment in microglia may be responsible for altered expression of

both inflammatory and neurotrophic regulators such as TNFα and TGFβ signalling observed

in the current study.

Epigenetic Regulation over Pathway-Specific Microglial transcriptomes involved in

Neuroprotection, Neuroinflammation and Phagocytosis in the Stroke Brain

Impact of HDAC inhibition on microglia in stroke brain suggests spatially defined

transcriptomic changes as the results highlight the spatial heterogeneity of microglial

transcriptomes within various regions of the stroke brain. Notably this was evident as

pathways crucial to neuroprotection in microglia of stroke brain after HDACi treatment exhibit

a transcriptomic signature similar to that of the sham brain.

The TGF-β signaling pathway, central to immune modulation and cell differentiation,

displayed marked upregulation of TGFβ receptors and SMAD family genes (e.g., TGFβR1,

TGFβR2, SMAD 2-4) in microglia of stroke brain following HDACi treatment. In TGFβ

signaling, TGFβR-Smad family complexes regulate the expression of TGFβ target genes

(Chandra *et al.*, 2012; Taylor *et al.*, 2017; Du *et al.*, 2018) (Lin *et al.*, 2017; Zöller *et al.*, 2018;

Zhu *et al.*, 2023); the surge of TGF-β signaling pathway genes suggests an amplified anti-

inflammatory and regenerative response in microglia, potentially enhancing their interactions

with neurons to support tissue repair and functional recovery in stroke brain following HDACi

treatment. HDACi has been found to upregulate many other anti-inflammatory genes

including STAT3 and PPARγ indicating an induction of anti-inflammatory response of

microglia in the stroke brain (Patnala et al., 2017).

Further, HDACi treatment restored expression of several other TGF-β signaling pathway

molecules including BMP and activin receptor membrane-bound inhibitor (*BAMBI*) which

interferes   with   TGFβ   or   BMP   pathway   transduction   by   forming   heterodimers

with *TGFβRI/BMPRI* (Chen *et al.*, 2023). The involvement of *BAMBI*, and various BMPs

highlights the diverse role of TGF-β signaling through microglia, influencing not only immune

responses but also neurogenesis, and synaptic plasticity (Meyers and Kessler, 2017).

It has been well established that microglia contribute to neuroinflammation through

excessive secretion of proinflammatory mediators in the stroke brain (Jayaraj *et al.*, 2019).

The activation of proinflammatory mediators, is regulated by mitogen-activated protein

kinases (MAPK) comprising of extracellular signal regulated kinase (ERK), c-Jun N-terminal

kinases (JNK), and p38 and nuclear factor kappa beta (NF-κβ) gene expression (Kaminska,

Mota and Pizzi, 2016). HDAC inhibition has been shown to modulate the p38 MAPK-MK2

signaling cascade, regulating microglial activation and cytokine production (Kaminska *et al.*,

2009; Kaminska, Mota and Pizzi, 2016). In the present study, many of the MAPK signalling

pathway genes were found to be altered in microglia following ischemia and HDACi

treatment. Further, in microglia of stroke brain the downregulation of c-Jun upon HDACi

administration appears to modulate the neuroinflammatory response, promoting

neuroprotection (Waetzig *et al.*, 2005). This downregulation of cJun could be mediated by

HDACi-induced activation of MKP-1 (DUSP1) which has been shown to inactivate MAP

Kinases (Eljaschewitsch *et al.*, 2006). Overall, HDACi appears to shift microglial activation

towards neuroprotective states, reducing neuroinflammation and promoting neuroprotection

through epigenetic modifications.

Further, HDACi-induced upregulation of PI3K-AKT pathway in microglia of stroke brain

appears to associated with neuroprotection since we have previously demonstrated that

microglial PI3K-AKT-BDNF pathway regulates memory and learning in neurons via long term

potentiation (LTP) (Saw *et al.*, 2020). In addition, HDACi treatment seems to potentiate an

array of neuroprotective signaling cascades such as through the upregulation of microglial

TrkB, a neurotrophin receptor tyrosine kinase which is a potent regulator of hippocampal LTP

(Minichiello, 2009).

Microglia play a crucial role in clearing debris from the stroke microenvironment through

phagocytosis to enhance neuroprotection (Wang, Leak and Cao, 2022). This was evident as

expression of FC-γ receptor-mediated phagocytosis pathway genes such as FCGR1,

FCGR3, FCGR4, SYK, and PIP5K1C was found to be upregulated in microglia of stroke

brain in the current study. Interestingly, expression of these genes was further upregulated

upon HDACi treatment indicating that HDACi enhances microglial response to ischemia and

capacity for clearing cellular debris, limiting secondary neurodegeneration and inflammation

(Qiao, Liu and Qie, 2023). Overall, HDACi-mediated phagocytic activity of microglia

potentially ameliorates exacerbated neuroinflammation in the penumbra of the stroke brain

(Lai, Zhang and Wang, 2014; Belov Kirdajova *et al.*, 2020). Spatially, these transcripts in

hippocampal microglia emphasize their role in mitigating damage and promoting recovery in

regions critical for cognitive functions.

The microglial transcriptomic changes induced by HDACi in different penumbral regions of

the stroke brain showcase a shift in microglial functionality from a proinflammatory

phenotype to a phenotype which improves brain repair and regeneration by enhancing

phagocytosis and reducing neuroinflammation.

This study also shows that HDACi altered the neuronal spatiotranscriptome to promote

resilience and plasticity within the ischemic brain environment (Ganai, Ramadoss and

Mahadevan, 2016; Kitahara *et al.*, 2021). The inverse expression patterns in neurotrophic

and neuroprotective pathways of these neurons in sham and stroke brains suggest broader

implications for brain plasticity through upstream epigenetic mechanisms aimed at

counterbalancing ischemia-associated excitotoxicity and neuroinflammatory response to

promote tissue repair and reduce neuronal susceptibility to reperfusion injury (Delcuve, Khan

and Davie, 2012; Kitahara *et al.*, 2021).

In conclusion, our high-throughput spatio-transcriptomic analysis unravels the mechanisms

by which a) microglia contribute to ischemic pathology with region-specificity and b) HDACi

ameliorates reperfusion mediated ischemic-injury to neurons through modulation of

microglial transcriptome. These findings emphasize an interplay between epigenetic

regulation and spatial transcriptomic changes altering communication networks between

microglia and neurons. Understanding the molecular mechanisms by which HDACi induced

phenotypic changes together with altered spatial transcriptomic landscape in microglia of

stroke brain would provide insight into development of potential microglia-mediated

therapeutic strategies for brain recovery post ischemia.

Funding

This study was funded by the Ministry of Education, Singapore (MOE Tier2, A-0002277-00-00 and MOE Tier1, A-8001684-00-00 awarded to Professor S T Dheen).

Acknowledgements

This article has been published on a preprint server: bioRxiv 2024.08.08.607139; doi: https://doi.org/10.1101/2024.08.08.607139

Components of Figure 1 have been created with BioRender.com.

Author Contribution

Data acquisition: **KJ, RK**. Data analysis and interpretation: **KJ, ST.** Study design and conceptualization: **ST.** Writing of the original draft: **KJ, ST.** Reviewing & editing of manuscript: **all authors.** Supervision: **ST.**

Data Availability:

High-throughput, next-generation sequencing data from spatial transcriptomic libraries in this study is available at NCBI Sequence Read Archive (SRA) under BioProject PRJNA1144233.

Conflict of Interest Declaration

The authors declare that they have conflicts of interest related to the study.

Ethics Approval

All experimental procedures performed *in vivo* were approved by the National University of Singapore Animal Care and Use Committee, under protocol NUS/IACUC/R19-0596.

Bibliography

1. Adcock, I. M. (2007) 'HDAC inhibitors as anti-inflammatory agents', *British Journal of Pharmacology*. Wiley, 150(7), p. 829. doi: 10.1038/SJ.BJP.0707166.

2. Allis, C. D. and Jenuwein, T. (2016) 'The molecular hallmarks of epigenetic control', *Nature Reviews Genetics*. Nature Publishing Group, 17(8), pp. 487–500. doi: 10.1038/nrg.2016.59.

3. Ansari, S. *et al.* (2011) 'Intraluminal middle cerebral artery occlusion (MCAO) model for ischemic stroke with laser doppler flowmetry guidance in mice', *Journal of Visualized Experiments*, (51), pp. 8–11. doi: 10.3791/2879.

4. Ayata, P. *et al.* (2018) 'Epigenetic regulation of brain region-specific microglia clearance activity', *Nature Neuroscience*. Nature Publishing Group, 21(8), pp. 1049–1060. doi: 10.1038/s41593-018-0192-3.

5. Babicki, S. *et al.* (2016) 'Heatmapper: web-enabled heat mapping for all', *Nucleic acids research*. Nucleic Acids Res, 44(W1), pp. W147–W153. doi: 10.1093/NAR/GKW419.

6. Bannister, A. J. and Kouzarides, T. (2011) 'Regulation of chromatin by histone modifications', *Cell Research 2011 21:3*. Nature Publishing Group, 21(3), pp. 381–395. doi: 10.1038/cr.2011.22.

7. Belov Kirdajova, D. *et al.* (2020) 'Ischemia-Triggered Glutamate Excitotoxicity From the Perspective of Glial Cells', *Frontiers in Cellular Neuroscience*. Frontiers Media SA, 14. doi: 10.3389/FNCEL.2020.00051.

8. Chandra, M. *et al.* (2012) 'Nuclear Translocation of Type I Transforming Growth Factor β Receptor Confers a Novel Function in RNA Processing', *Molecular and Cellular Biology*. Taylor & Francis, 32(12), p. 2183. doi: 10.1128/MCB.00320-12.

9. Charlton, T. *et al.* (2023) 'Brain-derived neurotrophic factor (BDNF) has direct anti-inflammatory effects on microglia', *Frontiers in cellular neuroscience*. Front Cell Neurosci, 17. doi: 10.3389/FNCEL.2023.1188672.

10. Chen, J. *et al.* (2009) *Animal models of acute neurological injuries, Animal Models of*

Acute Neurological Injuries. doi: 10.1007/978-1-60327-185-1.

11. Chen, J. *et al.* (2019) 'EZH2 inhibitor DZNep modulates microglial activation and

protects against ischaemic brain injury after experimental stroke', *European Journal*

of Pharmacology, 857. doi: 10.1016/j.ejphar.2019.172452.

12. Chen, X. *et al.* (2023) 'BMP and activin receptor membrane bound inhibitor: BAMBI

has multiple roles in gene expression and diseases (Review)', *Experimental and*

Therapeutic Medicine, 27(1). doi: 10.3892/etm.2023.12316.

13. Cheray, M. and Joseph, B. (2018) 'Epigenetics control microglia plasticity', *Frontiers*

in Cellular Neuroscience, 12(August), pp. 1–13. doi: 10.3389/fncel.2018.00243.

14. Delcuve, G. P., Khan, D. H. and Davie, J. R. (2012) 'Roles of histone deacetylases in

epigenetic regulation: Emerging paradigms from studies with inhibitors', *Clinical*

Epigenetics. BioMed Central, 4(1), pp. 1–13. doi: 10.1186/1868-7083-4-

5/FIGURES/2.

15. Donkor, E. S. (2018) 'Stroke in the 21st Century: A Snapshot of the Burden,

Epidemiology, and Quality of Life', *Stroke Research and Treatment.* Hindawi Limited,

2018. doi: 10.1155/2018/3238165.

16. Du, X. *et al.* (2018) 'SMAD4 feedback regulates the canonical TGF-β signaling

pathway to control granulosa cell apoptosis', *Cell Death & Disease 2018 9:2.* Nature

Publishing Group, 9(2), pp. 1–12. doi: 10.1038/s41419-017-0205-2.

17. Dubbelaar, M. L. *et al.* (2018) 'The Kaleidoscope of Microglial Phenotypes', *Frontiers*

in immunology, 9(July), p. 1753. doi: 10.3389/fimmu.2018.01753.

18. Einenkel, A. M. and Salameh, A. (2024) 'Selective vulnerability of hippocampal CA1

and CA3 pyramidal cells: What are possible pathomechanisms and should more

attention be paid to the CA3 region in future studies?', *Journal of Neuroscience*

Research. John Wiley & Sons, Ltd, 102(1), p. e25276. doi: 10.1002/JNR.25276.

19. Eljaschewitsch, E. *et al.* (2006) 'The endocannabinoid anandamide protects neurons

during CNS inflammation by induction of MKP-1 in microglial cells', *Neuron.* Neuron,

49(1), pp. 67–79. doi: 10.1016/J.NEURON.2005.11.027.

20. Feigin, V. L. *et al.* (2022) 'World Stroke Organization (WSO): Global Stroke Fact

Sheet 2022', *International Journal of Stroke*. SAGE Publications Inc., 17(1), pp. 18–

29. doi: 10.1177/17474930211065917.

21. Fessler, E. *et al.* (2013) 'Potential Roles of HDAC Inhibitors in Mitigating Ischemia-

induced Brain Damage and Facilitating Endogenous Regeneration and Recovery',

Current Pharmaceutical Design, 19(28), pp. 5105–5120. doi:

10.2174/1381612811319280009.

22. Ganai, S. A., Ramadoss, M. and Mahadevan, V. (2016) 'Histone Deacetylase

(HDAC) Inhibitors - Emerging Roles in Neuronal Memory, Learning, Synaptic

Plasticity and Neural Regeneration', *Current Neuropharmacology*. Bentham Science

Publishers, 14(1), p. 55. doi: 10.2174/1570159X13666151021111609.

23. Garcia, J. H. *et al.* (1995) 'Neurological deficit and extent of neuronal necrosis

attributable to middle cerebral artery occlusion in rats: Statistical validation', *Stroke*.

Lippincott Williams and Wilkins, 26(4), pp. 627–635. doi:

10.1161/01.STR.26.4.627/ASSET/D6F51A65-684F-4409-A4B6-

F5F58099E815/ASSETS/GRAPHIC/HS0451886005.GIF.

24. Gerrits, E. *et al.* (2020) 'Transcriptional profiling of microglia; current state of the art

and future perspectives', *GLIA*, 68(4). doi: 10.1002/glia.23767.

25. Getis, A. (2007) 'Reflections on spatial autocorrelation', *Regional Science and Urban

Economics*. North-Holland, 37(4), pp. 491–496. doi:

10.1016/J.REGSCIURBECO.2007.04.005.

26. Goldlust, E. J. *et al.* (1996) 'Automated measurement of infarct size with scanned

images of triphenyltetrazolium chloride-stained rat brains', *Stroke*. Lippincott Williams

and Wilkins, 27(9), pp. 1657–1662. doi:

10.1161/01.STR.27.9.1657/ASSET/67D2E47D-FF57-447E-A968-

A4899AD60E1E/ASSETS/GRAPHIC/0011F5.JPEG.

27. Gupta, N. *et al.* (2018) 'Recent progress in therapeutic strategies for microglia-

mediated neuroinflammation in neuropathologies', *Expert Opinion on Therapeutic*

Targets. Taylor & Francis, 22(9), pp. 765–781. doi:

10.1080/14728222.2018.1515917.

28. Haroon, E., Miller, A. H. and Sanacora, G. (2016) 'Inflammation, Glutamate, and Glia:

A Trio of Trouble in Mood Disorders', *Neuropsychopharmacology 2017 42:1*. Nature

Publishing Group, 42(1), pp. 193–215. doi: 10.1038/npp.2016.199.

29. Hickman, S. *et al.* (2018) 'Microglia in neurodegeneration', *Nature Neuroscience*.

Springer US, 21(10), pp. 1359–1369. doi: 10.1038/s41593-018-0242-x.

30. Hong, S., Dissing-Olesen, L. and Stevens, B. (2016) 'New insights on the role of

microglia in synaptic pruning in health and disease', *Current Opinion in Neurobiology*.

Elsevier Current Trends, 36, pp. 128–134. doi: 10.1016/J.CONB.2015.12.004.

31. Jayaraj, R. L. *et al.* (2019) 'Neuroinflammation: Friend and foe for ischemic stroke',

Journal of Neuroinflammation. BioMed Central Ltd., pp. 1–24. doi: 10.1186/s12974-

019-1516-2.

32. Kaminska, B. *et al.* (2009) 'MAPK Signal Transduction Underlying Brain Inflammation

and Gliosis as Therapeutic Target', *The Anatomical Record: Advances in Integrative*

Anatomy and Evolutionary Biology. John Wiley & Sons, Ltd, 292(12), pp. 1902–1913.

doi: 10.1002/AR.21047.

33. Kaminska, B., Mota, M. and Pizzi, M. (2016) 'Signal transduction and epigenetic

mechanisms in the control of microglia activation during neuroinflammation',

Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease. Elsevier, 1862(3),

pp. 339–351. doi: 10.1016/J.BBADIS.2015.10.026.

34. Kitahara, M. *et al.* (2021) 'Exercise and pharmacological inhibition of histone

deacetylase improves cognitive function accompanied by an increase of gene

expressions crucial for neuronal plasticity in the hippocampus', *Neuroscience Letters*.

Elsevier, 749, p. 135749. doi: 10.1016/J.NEULET.2021.135749.

35. Kulthinee, S. *et al.* (2022) 'Critical Functions of Histone Deacetylases (HDACs) in

Modulating Inflammation Associated with Cardiovascular Diseases',

Pathophysiology. Multidisciplinary Digital Publishing Institute (MDPI), 29(3), p. 471.

doi: 10.3390/PATHOPHYSIOLOGY29030038.

36. Lai, T. W., Zhang, S. and Wang, Y. T. (2014) 'Excitotoxicity and stroke: Identifying

novel targets for neuroprotection', *Progress in Neurobiology*. Pergamon, 115(C), pp.

157–188. doi: 10.1016/J.PNEUROBIO.2013.11.006.

37. Li, L. *et al.* (2020) 'Potential microglia-based interventions for stroke', *CNS*

Neuroscience & Therapeutics. Blackwell Publishing Ltd, 26(3), pp. 288–296. doi:

10.1111/cns.13291.

38. Li, Y. *et al.* (2000) 'Intrastriatal transplantation of bone marrow nonhematopoietic

cells improves functional recovery after stroke in adult mice', *Journal of Cerebral*

Blood Flow and Metabolism. Lippincott Williams and Wilkins, 20(9), pp. 1311–1319.

doi: 10.1097/00004647-200009000-

00006/ASSET/IMAGES/LARGE/10.1097_00004647-200009000-00006-FIG2.JPEG.

39. Li, Y. *et al.* (2024) 'Sodium butyrate alleviates lead-induced neuroinflammation and

improves cognitive and memory impairment through the ACSS2/H3K9ac/BDNF

pathway', *Environment International*. Pergamon, 184, p. 108479. doi:

10.1016/J.ENVINT.2024.108479.

40. Lin, E. *et al.* (2017) 'Transforming growth factor-β signaling pathway-associated

genes SMAD2 and TGFBR2 are implicated in metabolic syndrome in a Taiwanese

population', *Scientific reports*. Sci Rep, 7(1). doi: 10.1038/S41598-017-14025-4.

41. Marmorstein, R. and Zhou, M. M. (2014) 'Writers and Readers of Histone Acetylation:

Structure, Mechanism, and Inhibition', *Cold Spring Harbor Perspectives in Biology*.

Cold Spring Harbor Laboratory Press, 6(7). doi: 10.1101/CSHPERSPECT.A018762.

42. Mergenthaler, P., Dirnagl, U. and Kunz, A. (2022) 'Ischemic Stroke: Basic

Pathophysiology and Clinical Implication', *Neuroscience in the 21st Century: From*

Basic to Clinical: Third Edition. Springer International Publishing, pp. 3807–3827. doi:

10.1007/978-3-030-88832-9_97/COVER.

43. Meyers, E. A. and Kessler, J. A. (2017) 'TGF-β Family Signaling in Neural and

Neuronal Differentiation, Development, and Function', *Cold Spring Harbor*

Perspectives in Biology. Cold Spring Harbor Laboratory Press, 9(8). doi:

10.1101/CSHPERSPECT.A022244.

44. Minichiello, L. (2009) 'Long-term potentiation Synaptic plasticity TrkB signalling

pathways in LTP and learning', *Nature Reviews Neuroscience*, 10.

45. Nagy, E. E. *et al.* (2020) 'Neuroinflammation and microglia/macrophage phenotype

modulate the molecular background of post-stroke depression: A literature review',

Experimental and Therapeutic Medicine. Spandidos Publications, 20(3), p. 2510. doi:

10.3892/ETM.2020.8933.

46. Olah, M. *et al.* (2011) 'Microglia Phenotype Diversity', *CNS & Neurological Disorders

- Drug Targets*, 10(1), pp. 108–118. doi: 10.2174/187152711794488575.

47. Oliveros, J. C. (2007) *Venny. An Interactive Tool for Comparing Lists with Venn's

Diagrams. - References - Scientific Research Publishing*, *Csic*.

48. Patnala, R. *et al.* (2017) 'HDAC Inhibitor Sodium Butyrate-Mediated Epigenetic

Regulation Enhances Neuroprotective Function of Microglia During Ischemic Stroke',

Molecular Neurobiology, 54(8), pp. 6391–6411. doi: 10.1007/s12035-016-0149-z.

49. Penas, C. and Navarro, X. (2018) 'Epigenetic modifications associated to

neuroinflammation and neuropathic pain after neural trauma', *Frontiers in Cellular

Neuroscience*. Frontiers Media S.A., p. 158. doi: 10.3389/fncel.2018.00158.

50. Qiao, C., Liu, Z. and Qie, S. (2023) 'The Implications of Microglial Regulation in

Neuroplasticity-Dependent Stroke Recovery', *Biomolecules 2023, Vol. 13, Page 571*.

Multidisciplinary Digital Publishing Institute, 13(3), p. 571. doi:

10.3390/BIOM13030571.

51. Qiu, M., Xu, E. and Zhan, L. (2021) 'Epigenetic Regulations of Microglia/Macrophage

Polarization in Ischemic Stroke', *Frontiers in Molecular Neuroscience*. Frontiers

Media S.A., 14, p. 697416. doi: 10.3389/FNMOL.2021.697416/BIBTEX.

52. Quintana, J. F. *et al.* (2022) 'Single cell and spatial transcriptomic analyses reveal

microglia-plasma cell crosstalk in the brain during Trypanosoma brucei infection',

Nature Communications, 13(1). doi: 10.1038/s41467-022-33542-z.

53. Saw, G. *et al.* (2020) 'Epigenetic regulation of microglial phosphatidylinositol 3-kinase

pathway involved in long-term potentiation and synaptic plasticity in rats', *Glia*, 68(3),

pp. 656–669. doi: 10.1002/glia.23748.

54. Simpson, D. S. A. and Oliver, P. L. (2020) 'ROS Generation in Microglia:

Understanding Oxidative Stress and Inflammation in Neurodegenerative Disease',

Antioxidants. MDPI AG, 9(8), p. 743. doi: 10.3390/antiox9080743.

55. Sun, J. *et al.* (2015) 'Neuroprotective Effect of Sodium Butyrate against Cerebral

Ischemia/Reperfusion Injury in Mice', *BioMed Research International*. Hindawi

Publishing Corporation, 2015. doi: 10.1155/2015/395895.

56. Tang, D. *et al.* (2023) 'SRplot: A free online platform for data visualization and

graphing', *PLOS ONE*. Public Library of Science, 18(11), p. e0294236. doi:

10.1371/JOURNAL.PONE.0294236.

57. Tang, Y. *et al.* (2014) 'Jmjd3 is essential for the epigenetic modulation of microglia

phenotypes in the immune pathogenesis of Parkinson's disease', *Cell Death and*

Differentiation. Nature Publishing Group, 21(3), pp. 369–380. doi:

10.1038/cdd.2013.159.

58. Taylor, R. A. *et al.* (2017) 'TGF-β1 modulates microglial phenotype and promotes

recovery after intracerebral hemorrhage', *Journal of Clinical Investigation*, 127(1).

doi: 10.1172/JCI88647.

59. Traystman, R. J. (2003) 'Animal models of focal and global cerebral ischemia', *ILAR*

Journal. Institute for Laboratory Animal Research, 44(2), pp. 85–95. doi:

10.1093/ILAR.44.2.85.

60. Waetzig, V. *et al.* (2005) 'c-Jun N-terminal kinases (JNKs) mediate pro-inflammatory

actions of microglia', *Glia*. Glia, 50(3), pp. 235–246. doi: 10.1002/GLIA.20173.

61. Wang, L. *et al.* (2021) 'Epigenetic Modulation of Microglia Function and Phenotypes

in Neurodegenerative Diseases', *Neural Plasticity*, 2021. doi: 10.1155/2021/9912686.

62. Wang, Y., Leak, R. K. and Cao, G. (2022) 'Microglia-mediated neuroinflammation

and neuroplasticity after stroke', *Frontiers in Cellular Neuroscience.* Frontiers Media

S.A., 16, p. 980722. doi: 10.3389/fncel.2022.980722.

63. Wendimu, M. Y. and Hooks, S. B. (2022) 'Microglia Phenotypes in Aging and Neurodegenerative Diseases', *Cells*. Multidisciplinary Digital Publishing Institute (MDPI), 11(13). doi: 10.3390/CELLS11132091.

64. Xu, Y. *et al.* (2024) 'Sodium butyrate activates the KATP channels to regulate the mechanism of Parkinson's disease microglia model inflammation', *Immunity, Inflammation and Disease*. John Wiley & Sons, Ltd, 12(3), p. e1194. doi: 10.1002/IID3.1194.

65. Yang, S. J. *et al.* (2018) 'The NLRP3 Inflammasome: An Important Driver of Neuroinflammation in Hemorrhagic Stroke', *Cellular and Molecular Neurobiology*. Springer US, 38(3), pp. 595–603. doi: 10.1007/s10571-017-0526-9.

66. Yang, X. J. and Seto, E. (2007) 'HATs and HDACs: From structure, function and regulation to novel strategies for therapy and prevention', *Oncogene*. doi: 10.1038/sj.onc.1210599.

67. Yu, D., Huber, W. and Vitek, O. (2013) 'Shrinkage estimation of dispersion in Negative Binomial models for RNA-seq experiments with small sample size', *Bioinformatics*. Oxford Academic, 29(10), pp. 1275–1282. doi: 10.1093/BIOINFORMATICS/BTT143.

68. Zhang, F. *et al.* (2022) 'Regulation of N6-methyladenosine (m6A) RNA methylation in microglia-mediated inflammation and ischemic stroke', *Frontiers in Cellular Neuroscience*. Frontiers Media S.A., 16, p. 955222. doi: 10.3389/FNCEL.2022.955222/BIBTEX.

69. Zhang, H. *et al.* (2019) 'Isosteviol Sodium Protects against Ischemic Stroke by Modulating Microglia/Macrophage Polarization via Disruption of GAS5/miR-146a-5p sponge', *Scientific Reports 2019 9:1*. Nature Publishing Group, 9(1), pp. 1–16. doi: 10.1038/s41598-019-48759-0.

70. Zhang, W. *et al.* (2021) 'Microglia-associated neuroinflammation is a potential therapeutic target for ischemic stroke', *Neural Regeneration Research*. Wolters

854  Kluwer Medknow Publications, 16(1), p. 6. doi: 10.4103/1673-5374.286954.

855 71. Zhou, Z. *et al.* (2021) 'Sodium butyrate attenuated neuronal apoptosis via

856  GPR41/Gβγ/PI3K/Aktpathway after MCAO in rats', *Journal of Cerebral Blood Flow &*

857  *Metabolism*. SAGE Publications, 41(2), p. 267. doi: 10.1177/0271678X20910533.

858 72. Zhu, K. *et al.* (2023) 'Disrupting microglial TGF-β signaling triggers region-specific

859  pathology in the spinal cord', *bioRxiv*. Cold Spring Harbor Laboratory, p.

860  2023.04.24.538074. doi: 10.1101/2023.04.24.538074.

861 73. Ziemka-Nalecz, M. *et al.* (2017) 'Sodium Butyrate, a Histone Deacetylase Inhibitor,

862  Exhibits Neuroprotective/Neurogenic Effects in a Rat Model of Neonatal Hypoxia-

863  Ischemia', *Molecular Neurobiology*, 54(7), pp. 5300–5318. doi: 10.1007/s12035-016-

864  0049-2.

865 74. Zöller, T. *et al.* (2018) 'Silencing of TGFβ signalling in microglia results in impaired

866  homeostasis', *Nature Communications 2018 9:1*. Nature Publishing Group, 9(1), pp.

867  1–13. doi: 10.1038/s41467-018-06224-y.

Figure Legend:

Figure 1: Middle cerebral artery occlusion and gross neurological status upon HDACi

treatment.

(A) Graphical pipeline describing ischemic stroke modelling in the rodent brain, ischemia-

associated microglia and phenotypes. (B) Toluidine blue (nissl) staining of brain tissue

sections with MCAo and Sham showcasing infarct extent and anatomical regions of interest

(ROI) covered under the penumbra. (C) Epigenetically mediated protection through HDACi

attenuates infarct volume 24hours post ischemic insult (TTC or 2,3,5-Triphenyltetrazolium

Chloride assessment of brain slices exhibit substantial recovery of viable brain tissue upon

NaBu administration) (n=3 in each experimental group). (D) Behavioural assessment of

ischemic affect through neurological deficits occurring through stroke demonstrates

enhanced locomotor activity upon HDACi administration with reduced circling, spastic

paralysis, allodynia and locomotor latency (to reach enrichment shell within home cage) after

HDACi treatment (n=3 in Sham and MCAo and n=4 in MCAo+HDACi groups). X axis

represents behavioural paradigm tested. Y axis represents scoring achieved for each group -

Circling: [no = 0, yes = 1], Latency: [1 to 8 seconds to reach enrichment shell] Spasticity and

Paralysis: [No spasticity or paralysis = 0, Spasticity = 1, Spastic paralysis = 2] Allodynia: [no

allodynia or vocalization = 0, allodynia sensitivity = 1, allodynia+vocalization = 2]. (E)

Toluidine blue stained mouse brain sections of dentate gyrus neurons in the Hippocampus

exhibit variance in infarct intensity and neuronal damage post-ischemic stroke with HDACi

administration. (F) Immunostained confocal images of NeuN (channel AF 674, false-colored

green) and IBA1 (channel AF488, false-colored red) appear to display preserved neuronal

organization in the penumbral cortex and reduced microglial aggregation at the infarct

periphery upon HDAC inhibition. Nucleus is stained with Hoechst 33342 (blue).

Figure 1 Supplementary Figure 1

TTC (2,3,5-Triphenyltetrazolium Chloride) treated mouse brain sections across biological

groups showcasing extent of infarct through viable (stained, red) tissue and non-viable

(unstained, white) tissue. (n=3 in each experimental group).

Figure 1 Supplementary Video 1

Assessment of ischemic stroke affect through preliminary neurological deficit scoring

(pNDS); video capture of each biological group showcases mild-moderate immobility and

circling behaviour in the MCAo group while sham and HDACi-treated groups exhibit reduced

to nil locomotor defects such as circling and difficulty reaching enrichment shell within home

cage.

Figure 2. Microglial morphometric assessment through surface reconstruction with

IMARIS 9.5

(A) High-resolution confocal images of Microglia stained with IBA1 (channel AF 488, green)

reveal diverse morphological phenotypes between biological groups (n=3 in each

experimental group). (B) Individual microglial cells were segmented and processed across

groups to obtain surface area and cellular volume morphometric data. 3D-reconstruction of

microglial surfaces spotlights microglia with ramified and long, slender processes in HDACi

inhibitor administered mouse brain reminiscent of sham brain microglia while MCAo brain

microglia are amoeboid with substantial depth across the z-stack. (C) Microglial surface area

and intracellular volume are inverse between stroke and epigenetically treated groups,

presenting a phenotypic reversal of microglia in the penumbra.

Figure 2 Supplementary Video 1

Video capture of IMARIS 9.5 reconstructions (30μm thick, z-stacked sections), showcasing surface creation and segmentation of labelled microglial cells (IBA1, AF488, false-colored red) that are processed to obtain morphometric data such as surface area and cellular volume to evaluate phenotypic status of microglia in the stroke brain after HDACi treatment (n=3 in each experimental group)

Figure 3. HDAC inhibition and histone mark status in microglia after ischemic stroke

(A-B) Hyper-intense expression of H3K27me3 (channel 647, false-colored green) in MCAo brain microglia (IBA1, channel 488, false-colored red) appear to be rescued towards sham expression levels in hippocampal microglia from HDACi-treated stroke brains. Potential shuttling of H3K27me3 suggests upstream involvement of the PRC2 complex and EZH2, components mediating H3K27me3-related gene silencing upon ischemic insult. (C) Similar to H3K27me3, H3K4me3 (channel 647, false-colored green) expression levels in sham and HDACi-treated stroke brain tissue showcase similar expression patterns with stroke elevating expression in microglial nuclei suggesting neuron-microglia regulatory networks activated by HDACi administration.

Figure 3 Supplementary Figure 1

(A,B,C) Tilescan (10x) images of complete brain sections across biological groups; (D,E,F) showcasing hippocampal expression of H3K27me3 (AF647, false-colored green) in microglia stained with IBA1 (AF488, false-colored red) that is upregulated upon ischemic stroke and ameliorated upon HDACi-treatment.

Figure 3 Supplementary Video 1

Video capture of (30μm thick, z-stacked sections) microglia stained with IBA1 (AF488, false-

colored red) and H3K27me3 (AF647, false-colored green) across all three biological groups,

showcasing elevation of histone mark expression and presence within the nuclei of

microglial cells.

Figure 4. Spatial transcriptome pipeline: ROIs, population clusters and microglial

spatiotranscriptomic signature (Visium Fresh Frozen)

(A) Immunofluorescence (IF) image and spatial (spot) overlay (B) of Visium fresh frozen

sections and marker gene population clusters (C) (n=3 in each experimental group). (B) 10x

Genomics proprietary Loupe Browser 6.0's default transcriptome overlay represented as k-

clusters and population identities; (C) the distinct population clusters appear to lose

organization upon ischemic insult. (D) Microglial spots enriched through microglia signature

genes adapted from (Gerrits *et al.*, 2020; Quintana *et al.*, 2022) performed through Loupe

browser 6.0's feature selection and thresholding to output microglia-specific spatial

transcriptomes segmented based on regions on interest in the tissue section. (E) Heatmap

of total spatial library with cell signature markers for processed brain samples exhibiting 19

population identities. (F) Uniform Manifold Approximation and Projection (UMAP) dimension

reduction of all cell populations in the dataset.

Figure 4 Supplementary Figure 1

(A,B) Tissue optimization timepoint experiments for brain samples with 10x Genomics

Visium-TO slides for evaluation of optimum cDNA permeabilization time points (10x UG

CG000238-RevE) prior to proceeding with gene expression and library preparation of Visium

fresh-frozen spatial libraries. 6 minutes of permeabilization was selected based on

optimization time points evaluated. (C) Antibody titre evaluation for IF-staining of spatial

tissue sections (10x UG CG000312-RevD) identified concentration of antibodies (table in

methodology).

Figure 4 Supplementary Figure 2

Permeabilized Visium fresh-frozen slides and quality control (QC) of RNA integrity checks

and pre-cDNA amplification/ cleanup for visium libraries prior to preparation and sequencing.

Figure 5. Spatial segmentation of microglial transcriptomes: expression signatures

across the ischemic penumbra and enriched pathways (KEGG)

(A) Grouping and segmentation of Penumbra regions of interest in *LoupeBrowser* for spatial

demarcation of microglia after ischemic stroke. (B, C, D) Four-set Venn diagrams of unique

and overlapping HDACi-Upregulated (B), Downregulated (C) and Rescued (D) Genes,

elucidate the homogenous and heterogenous marks in microglia across brain regions.

Pathway (Kegg) Enrichment (E) and Gene Ontology (F) analysis of genes dysregulated

homogenously across the four microglial spatiotranscriptomes are processed and showcase

potential microglial pathways of interest to analyze and target after ischemic stroke. (n=3 for

each experimental group)

Figure 5 Supplementary Figure 1

Alternative enriched (KEGG) pathways from commonly dysregulated genes of microglial

spatiotranscriptomes across all 4 ROIs (hippocampus, thalamus, striatum, cortex). These

pathways present additional targets and mechanisms that may be dysregulated throughout

penumbral microglia after ischemic stroke and HDACi-administration.

Figure 6. Microglial spatiotranscriptome expression of enriched, key biological pathways after HDACi administration in ischemic stroke.

Heatmaps of microglial spatiotranscriptomes of each ROI, hippocampal (A), thalamic (I), striatal (J), cortical (K) microglia exhibit differential profiles of selected biological pathways that were enriched and involved in ischemic injury across the three biological groups (n=3 for each experimental group). Modified and rescued (closer to sham expression) genes within identified pathways crucial for neuroprotective, neuroinflammatory and phagocytosis mediation responses are represented, and top 50 targets are represented in the radar plot for each penumbra ROI.

Figure 6 Supplementary Figure 1A

Pathways enriched in HDACi-rescued, region-specific microglial spatiotranscriptomes (hippocampus-A, thalamus-B, striatum-C, cortex-D). Evaluation of pathways rescued by HDACi after ischemic injury, highlights pathways crucial for neuroprotective, neuroinflammatory and phagocytosis mediation responses.

Figure 6 Supplementary Figure 1B

Gene Ontology analyses of region-specific, microglial spatiotranscriptomes (hippocampus-A, thalamus-B, striatum-C, cortex-D) exhibit differential GO terms in respective compartments (BP, CC and MF) involved in HDACi-mediated rescue after ischemic injury in microglia.

Figure 6 Supplementary Figure 2

Heatmaps representing all dysregulated genes within selected biological pathways (HDACi

vs MCAo Log2fc difference>1.0, Euclidean distance metric with hierarchical clustering

through complete linkage) in each penumbra microglial spatiotranscriptomes (hippocampus-

A, thalamus-B, striatum-C, cortex-D). Microglial heterogeneity in expression patterns of

these pathways further highlights the specificity of microglial responses to ischemic stroke

and HDACi therapy based on spatial contexts within the mouse brain.

1023

Figure 6 Supplementary Figure 3

Circular Network Plot (cnetplot) of HDACi-rescued pathways and targets in each microglial

spatiotranscriptome (hippocampus-A, thalamus-B, striatum-C, cortex-D) post-ischemic

stroke. Genes (nodes) are positioned around the plot, sized by log2 fold change (log2FC)

and colored from red (upregulated) to blue (downregulated). Pathway nodes are centrally

located, sized by the number of associated genes, and color-coded by functional categories.

Key pathways identified in the study including "MAPK signaling," "PI3K-Akt signaling," and

"Fc gamma R-mediated phagocytosis." are depicted with lines indicating gene-pathway

associations, illustrating the molecular interactions affected by HDACi treatment.

1033

Figure 7. Penumbra Neuronal Spatiotranscriptome

(A-B) Spatiotranscriptome of penumbral neurons showcases enriched neuronal pathways

relevant for survival through HDACi-mediated rescue after ischemic stroke. (C) Distinct,

inverse expression profiles after HDACi treatment can be observed, indicating context-

dependent microglia-neuron interaction upon ischemic stroke and epigenetic regulation

through HDACi. (n=3 for each experimental group)

1040

1041 **Figure 2. Supplementary File 1**

1042   Morphometry data table consisting of IMARIS *Surfaces* feature metrics (Volume, Surface
1043   Area of microglial cells)

1044

1045

1046   **Figure 6. Supplementary File 1**

1047   Enriched Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways of interest from
1048   individual microglial spatiotranscriptomes.

A

Mouse Model → Unilateral Middle Cerebral Artery Occlusion → Infarct extent coronal section of mouse brain → Spatial transcriptomic analysis

B

Sham | MCAo | MCAo+HDACi

Cortex
Hippocampus
Thalamus
Striatum
1mm

C

Infarct Extent Reduction after HDACi

Sham | MCAo | MCAo+HDACi

Infarct Area (mm²)

Viable Area (mm²) | Non-Viable Area (mm²)

*** pvalue < 0.001

D

HDACi-Mediated Recovery of Behavioural Deficits Upon Stroke

Sham | MCAo | MCAo+HDACi

Circling | Spasticity & Paralysis | Allodynia | Latency

* pvalue < 0.005

E

Sham | MCAo | MCAo+HDACi

25µm

F

Sham | MCAo | MCAo+HDACi

50µm

NeuN **IBA1** **HOECHST 33342**

Figure 2

Figure 3

Figure 4

HDACi Dysregulated and Rescued Genes across microglial spatiotranscriptomes

A

B — Striatum, Thalamus, Hippocampus, Cortex Venn diagram

80 (0.5%), 62 (0.4%), 253 (1.6%), 59 (0.4%), 214 (1.3%), 316 (2%), 229 (1.4%), 178 (1.1%), 535 (3.3%), 232 (1.4%), 3544 (22%), 328 (2%), 2021 (12.5%), 4310 (26.7%), 3764 (23.3%)

Upregulated microglial transcripts upon HDACi administration after MCAo

C — Striatum, Thalamus, Hippocampus, Cortex Venn diagram

2277 (18%), 4209 (33.3%), 195 (1.5%), 3561 (28.2%), 255 (2%), 168 (1.3%), 502 (4%), 300 (2.4%), 244 (1.9%), 297 (2.3%), 190 (1.5%), 266 (2.1%), 73 (0.6%), 64 (0.5%), 49 (0.4%)

Downregulated microglial transcripts upon HDACi administration after MCAo

D — Striatum, Thalamus, Hippocampus, Cortex Venn diagram

2093 (16.3%), 4284 (33.4%), 201 (1.6%), 3647 (28.5%), 308 (2.4%), 223 (1.7%), 541 (4.2%), 330 (2.6%), 190 (1.5%), 318 (2.5%), 203 (1.6%), 230 (1.8%), 107 (0.8%), 75 (0.6%), 63 (0.5%)

Rescued microglial transcripts upon HDACi administration after MCAo

E

Microglial pathways enriched in commonly dysregulated transcripts across all ROIs after HDACi administration in the ischemic penumbra

Focal adhesion
Rap1 signaling pathway
Purine metabolism
PI3K-Akt signaling pathway
Hippo signaling pathway
AMPK signaling pathway
MAPK signaling pathway
Ras signaling pathway
Neurotrophin signaling pathway
Sphingolipid signaling pathway
ErbB signaling pathway
Autophagy - other
T cell receptor signaling pathway
p53 signaling pathway
IL-17 signaling pathway

- log₁₀(pvalue): $-\log_{10}(pvalue)$ 1.5 to 3.5
count: 20, 40, 60

GeneRatio: 10/1399, 18/1399, 21/1399, 24/1399, 30/1399, 32/1399, 36/1399, 39/1399, 51/1399, 52/1399, 53/1399, 63/1399, 78/1399

F

GO: BP, CC, MF - commonly dysregulated spatiotranscriptome across ROIs of HDACi microglia in the ischemic penumbra

BP / CC / MF

Negative regulation of phosphorylation
Negative regulation of protein phosphorylation
tRNA processing
Small GTPase mediated signal transduction
Negative regulation of cell migration
RNA modification
DNA repair
Negative regulation of cellular component movement
Negative regulation of cell motility
Mitotic cell cycle phase transition
Lamellipodium
Cell leading edge
Lateral plasma membrane
Basolateral plasma membrane
Cell projection membrane
Ciliary basal body
Actin cytoskeleton
Membrane raft
Membrane microdomain
Membrane region
Methyltransferase activity
Transferase activity, transferring one-carbon groups
Ubiquitin-like protein ligase binding
Lyase activity
Amide binding
PDZ domain binding
Catalytic activity, acting on RNA
Ribonucleoside binding
Catalytic activity, acting on a tRNA
Nucleoside binding

Enrichment Score: 0.0, 2.5, 5.0, 7.5, 10.0

A Sham   MCAo   MCAo+HDACi

250μm   250μm   250μm

NeuN   Gfap   Dapi

B

Pathway enrichment in HDACi-treated Neuronal transcriptome within then Penumbra

PI3K-Akt signaling pathway
MAPK signaling pathway
Focal adhesion
Apoptosis
Notch signaling pathway
Ras signaling pathway
TGF-beta signaling pathway
Rap1 signaling pathway
Neurotrophin signaling pathway
mTOR signaling pathway
VEGF signaling pathway
FoxO signaling pathway
TNF signaling pathway
HIF-1 signaling pathway
AMPK signaling pathway
Hippo signaling pathway
Autophagy - animal
Calcium signaling pathway
Toll-like receptor signaling pathway
Long-term potentiation
Glutamatergic synapse
NF-kappa B signaling pathway
Serotonergic synapse
p53 signaling pathway
JAK-STAT signaling pathway
Long-term depression
GABAergic synapse
IL-17 signaling pathway
Fc gamma R-mediated phagocytosis
cAMP signaling pathway

Gene Ratio

pvalue
0.015
0.010
0.005

Gene Count

C

HDACi-altered gene expression in Penumbral Neurons

-1 0 1
Row Z-Score

SHAM   MCAo   MCAo+HDACi

TGF-Beta Signaling

PI3K-AKT Signaling

MAPK Signaling

Notch Signaling

Apoptosis