Spatial Transcriptomic Analysis Reveals HDAC Inhibition Modulates Microglial Dynamics to Protect Against Ischemic Stroke in Mice

Kevin Jayaraj[1], Ritesh Kumar[1], Sukanya Shyamasundar[1] Jai S Polepalli[1], Thiruma V Arumugam[2], S Thameem Dheen[1]

[1] Department of Anatomy, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117594

[2] Centre for Cardiovascular Biology and Disease Research, Department of Microbiology, Anatomy, Physiology and Pharmacology, School of Agriculture, Biomedicine and Environment, La Trobe University, Bundoora, Victoria, Australia

Correspondence: S Thameem Dheen   E-mail: antstd@nus.edu.sg

Co-Correspondence: Jai S Polepalli   E-mail: jpolepalli@nus.edu.sg

Key Words

Abstract:

Ischemic stroke significantly contributes to global morbidity and disability through a cascade of neurological responses. Microglia, the immune modulators within the brain, exhibit dual roles in exacerbating and ameliorating ischemic injury through neuroinflammatory and neuroprotective roles, respectively. Despite emerging insights into microglia's role in neuronal support, the potential of epigenetic intervention to modulate microglial activity remains largely unexplored. We have previously shown that sodium butyrate, a histone deacetylase inhibitor (HDACi) epigenetically regulates inflammatory response of microglia after ischemic stroke and this study was aimed to characterize the transcriptomic profiles of microglia and their spatial distribution in the stroke brain followed by HDACi administration. We hypothesized that the administration of HDACi epigenetically modulates microglial activation and a region-specific microglial phenotype in the stroke brain, shifting their phenotype from neurotoxic to neuroprotective and facilitating neuronal repair and recovery in the ischemic penumbra. Utilizing a rodent model of middle cerebral artery occlusion (MCAo), spatial transcriptomics and 3D morphometric reconstruction techniques were employed to investigate microglial responses in critical penumbral regions, such as the hippocampus, thalamus, cortex and striatum following HDACi administration. We found that HDACi significantly altered the microglial transcriptomic landscape involving biological pathways of neuroinflammation, neuroprotection and phagocytosis as well as morphological phenotype, promoting a shift towards reparative, neurotrophic profiles within the ischemic penumbra. These changes were also associated with enhanced neuronal survival and reduced neuroinflammation in specific regions in the ischemic brain. By elucidating the mechanisms through which HDAC inhibition influences microglial function, our findings propose therapeutic avenues for neuroprotection and rehabilitation in ischemic stroke, and possibly other neurodegenerative conditions that involve microglia-mediated neuroinflammation.

Introduction

Ischemic stroke is one of the primary causes of long-term disability and mortality worldwide, presenting a significant public health challenge by exacerbating both health and economic burdens on society (Donkor, 2018; Feigin *et al.*, 2022). Focal ischemic stroke which is characterized by abrupt cessation of blood supply to a specific region in the brain, leads to hypoxia (a deprivation of oxygen), triggering a cascade of deleterious cellular and molecular events such as excitotoxicity, neuroinflammation, and apoptosis. During reperfusion, restoration of blood flow to the infarcted tissue further aggravates tissue injury (Traystman, 2003; Mergenthaler, Dirnagl and Kunz, 2022). Complex, multicellular pathways and interactions dictate mechanisms that mediate neuronal damage post-ischemic injury and highlight the need for novel therapeutic strategies for enhancing recovery and clinical outcomes (Zhang *et al.*, 2019; Wang, Leak and Cao, 2022).

Microglia, resident immune cells of the brain are at the forefront of the body's response to ischemic injury. They exhibit a spectrum of phenotypic changes (Dubbelaar *et al.*, 2018; Wendimu and Hooks, 2022) and transition dynamically between conventional pro-inflammatory (M1) and anti-inflammatory (M2) states depending on spatial and temporal contexts after ischemia. In the stroke brain, microglia initially clear debris and release pro-inflammatory cytokines like TNF-α and interleukins, triggering neuroinflammatory responses crucial for initiating repair but turn deleterious if it progresses without biological checkpoints controlling the response (Wang, Leak and Cao, 2022). Transitioning to an anti-inflammatory phenotype, microglia release cytokines and neurotrophins promoting tissue repair and limiting ongoing damage (Gupta *et al.*, 2018; Zhang *et al.*, 2021; Charlton *et al.*, 2023). The balance between these microglial states and their heterogenous expression profiles are crucial for the prognosis of stroke. Positive modulation of microglial activation is essential in ameliorating neurodegeneration and reperfusion mediated injury, suggesting that targeted interventions aimed at microglia could offer significant therapeutic benefits (Nagy *et al.*, 2020).

Epigenetic mechanisms, such as DNA methylation, histone modification, and RNA-associated

silencing, play pivotal roles in regulating microglial function and response to ischemic stroke.

Histone deacetylases (HDACs) are key enzymes involved in removing acetyl groups from

histones, leading to reduced gene expression that has been explored to regulate progress of

pathology and neuroinflammation in stroke and other inflammatory conditions (Adcock, 2007;

Fessler *et al.*, 2013; Patnala *et al.*, 2017). In the context of ischemic stroke, such epigenetic

modifications influence the phenotypic switching of microglia between pro-inflammatory and

anti-inflammatory states, thus affecting the overall inflammatory milieu and recovery process

(Saw *et al.*, 2020; Qiu, Xu and Zhan, 2021).

Sodium butyrate, a HDAC inhibitor (HDACi), has emerged as a promising epigenetic therapy

for ischemic stroke (Sun *et al.*, 2015). By modulating the epigenetic landscape, HDACi can

shift microglial activation towards neuroprotective states, reducing neuroinflammation and

promoting neuroprotection. This therapeutic potential is further supported by studies

demonstrating its ability to enhance the expression of neurotrophic factors and modulate key

signaling pathways involved in cell survival, neurogenesis, and synaptic plasticity (Patnala *et*

al., 2017; Ziemka-Nalecz *et al.*, 2017; Saw *et al.*, 2020; Li *et al.*, 2024; Xu *et al.*, 2024).

Our research leverages advanced methodologies, including high-resolution imaging, 3D

microglial reconstruction, and spatial transcriptomics, to delve deeper into the effects of HDACi

on microglial dynamics and their transcriptional profile in the stroke brain. By exploring

biological networks relevant to microglia-neuron interaction and the impact of epigenetic

modulation on microglial function, we aim to uncover novel therapeutic targets that could

significantly enhance the functional outcomes in ischemic stroke therapy (Qiu, Xu and Zhan,

2021; Zhang *et al.*, 2022)

Given the intricate interplay between microglia, neurons, and other glial cells in the ischemic

brain, understanding the precise molecular pathways including those affected by epigenetic

modifications induced by microglial activity, is critical for developing targeted therapies for

stroke. The application of HDAC inhibitors like sodium butyrate offers a unique approach to

modulating the brain's innate immune response, steering it towards pathways that support

neuroprotection, repair, and regeneration. Such insights are crucial for developing targeted

interventions which harness the therapeutic potential of microglial modulation in the context of

ischemic stroke.

The present study demonstrates that HDACi treatment not only reduces infarct volume in

stroke brain but also influences microglial phenotypes towards reparative states, facilitating

neuronal survival and mitigating neuroinflammation. Through analysis incorporating cutting-

edge spatial transcriptomic techniques, we obtain nuanced understanding of microglial roles

in stroke pathophysiology and recovery, paving the way for targeted therapies that can

capitalize on the therapeutic promise of microglial modulation.

Methods

Ethics Statement

Adult male C57Bl/6J mice, 12-15 weeks old were acquired (InVivos Pte Ltd, Singapore) and

housed under controlled temperature, humidity, light and dark cycles at the NUS Comparative

Medicine (CM) Vivarium. Mice were provided access to standard food pellets and water ad

libitum. Animal work was performed in accordance with the Institutional Animal Care and Use

Committee (IACUC) through protocol NUS/IACUC/R19-0596 approved for the experiments

performed with animal(s) and tissues in this study.

Surgical Ischemic Stroke Modelling and HDACi treatment in Mice

MCAo modelling was performed as previously described (Patnala *et al.*, 2017) using adult

male C57BL/6J mice (12-15 weeks old, 26-29g). The mice were anesthetized with isofluorane

in an induction chamber. Ophthalmic lubricant (Duratears 3.5g, Alcon, USA) was applied to

prevent corneal drying, and body temperature was monitored and maintained at 37.0 ± 0.5°C

with a heating pad. The neck of the mice was shaved, disinfected with 70% ethanol and

povidone-iodine solution, and a midline incision was made to expose the common carotid

artery (CCA), internal carotid artery (ICA), and external carotid artery (ECA). The CCA and

ICA were clamped, and the ECA was cauterized and retracted. A Doccol 6-0 monofilament

nylon suture (catalog no.: 602123PK10, Doccol Corporation, USA), (diameter of 0.21-0.23

mm), was introduced through the ECA and advanced into the ICA to occlude the MCA.

Occlusion was maintained for 60 minutes, after which the suture was withdrawn to allow

reperfusion. The ECA incision was cauterized, and the microvascular clips were removed to

restore blood flow through the CCA and ICA. The surgical site was sutured, and the animals

were allowed to recover from anaesthesia under observation. In sham group mice, surgical

reflection of muscles to reveal CCA bifurcation was performed without disturbing the

vasculature to affect cerebral blood flow. Sodium butyrate (HDACi) was administered

intraperitoneally (I.P.) to the mice (Cat. no. B5887-1G, Sigma-Aldrich) at a dose of 300 mg/kg

bodyweight. Two doses of NaBu were administered, 1-hour prior ischemia and 6-hours post

reperfusionas described previously (Patnala *et al.*, 2017). Mice were housed a temperature-

controlled environment with a 12-hour light/dark cycle and provided food and water ad libitum.

Motor deficit after surgical modelling was evaluated through preliminary neurological deficit

scoring (pNDS). Severity of motor affect through was graded on a scale of 0 to 4 (normal score

0, maximal deficit score 4) (n=4) (Garcia *et al.*, 1995; Li *et al.*, 2000) . Animals were regularly

monitored for adverse effects or complications, ensuring compliance with guidelines for the

care and use of laboratory animals, aimed at minimizing suffering and the number of animals

used. 24 hours post-stroke induction, mice were euthanized with an intraperitoneal overdose

of sodium pentobarbital (150 mg/kg) and absence of pedal and corneal reflexes was confirmed

prior to harvest of brain tissue. For histological and immunofluorescence analyses, mice were

perfused with ice-cold 4% paraformaldehyde (PFA) and post-fixed at 4°C for 24 hours prior to

histological processing. For spatial transcriptomic analyses, brain tissue was processed as

per 10x Genomics Visium fresh frozen tissue preparation guide CG000240 - Rev E.

TTC Staining of mouse brain tissue and evaluation of tissue damage

For 2,3,5-Triphenyltetrazolium chloride (TTC) staining, fresh brain tissue was sectioned into 2

mm coronal slices using a brain matrix. The brain slices were submerged in cold PBS to

maintain tissue viability. Freshly prepared 2% solution of TTC (Cat. no. T8877, Sigma-Aldrich)

was added to brain sections in a 24-well plate and incubated at 37°C for 30 minutes in a dark,

humidified incubator. Viable tissue stained red, while infarcted areas remained pale or white.

Post-incubation, sections were rinsed with PBS and transferred to a clear, flat transparent

sheet for digital imaging on a flatbed scanner (CanoScan LiDe 400, Canon). Image processing

and analysis of TTC-stained brain sections was performed by thresholding scanned images

to greyscale and extrapolating pixel values (Goldlust *et al.*, 1996).Scanned images were

loaded and processed using R packages to calculate and distinguish viable (non-infarcted)

and non-viable (infarcted) area in brain tissue. Viable and non-viable areas were calculated in

square millimeters based on pixel counts and known image resolution. Statistical analysis,

including t-tests and ANOVA, was performed to compare the groups (n=3).. The results were

then imported into Microsoft Excel for visualization.

Histological Processing of Mouse Brain Tissue Sections

Harvested brain tissues were post-fixed in 4% paraformaldehyde and dehydrated through a

graded series of ethanol baths and clearene. Dehydrated and cleared samples were then

incubated in molten paraffin wax, embedded and allowed to solidify at room temperature.

Solidified wax blocks were mounted onto a microtome, trimmed, and sectioned at 7μm

thickness. Sections were floated in a warm water bath, transferred onto microscope slides,

and stained with toluidine blue. Routine staining protocols were performed involving

deparaffinization in clearene, rehydration through descending ethanol series, staining in

toluidine blue followed by rinsing, dehydrating ethanol series and tissue clearing prior to

coverslipping slides with mounting medium (S302380-2, Dako North America Inc.). For

cryosectioning, PFA-fixed brain tissues were cryoprotected in 30% sucrose until fully

saturated, embedded in OCT compound, and rapidly frozen using dry ice or pre-cooled

isopentane. Frozen OCT-embedded tissue blocks were sectioned at -20°C to -25°C in a

cryostat (Leica Instruments GmbH, Germany), with sections cut at 30 to 50 microns, collected

onto chilled slides, and stored at -80°C in sealed containers with desiccants until further use.

Immunofluorescence and Morphometric analysis

Mouse brain tissues were fixed with 4% paraformaldehyde, cryoprotected in 30% sucrose and

embedded in OCT compound. Mouse brains were sectioned at 30 or 50 micron thickness on

a cryostat (Leica Instruments GmbH, Germany). The sections were permeabilized with 0.3%

Triton X-100   and non-specific binding sites were blocked by incubating sections in 5% normal

donkey serum (D9663-10ML, Sigma-Aldrich) or 1% bovine serum albumin (126615-25MLCN,

Merck). The sections were then incubated with primary antibodies (antibody list in table below)

diluted in blocking solution, overnight at 4°C. After rinsing with PBS, sections were incubated

with fluorescence-conjugated secondary antibodies for 1 hour at room temperature, protected

from light. Following this, sections were rinsed with PBS and mounted with anti-fade

fluorescence mounting medium (ab104135, Abcam). The stained sections were imaged using

a confocal microscope (FluoView 1000 and FluoView 3000, Olympus Corporation, Japan) at

10x, 20x, 60x and 100x objectives. Z-stacks were captured at 60x and 100x objectives with a

z-plane of 20-30µm with z-sections of 0.5µm. Microglia-labelled confocal, Z-stacked imaging

datasets were imported into IMARIS v9.5 software for 3D reconstruction and analysis of

microglial morphology (n=3). The 'Surpass' view was used to inspect the z-stack, and a 3D

volume rendering of the z-stack was created using the 'Volume' button. For morphometric

evaluation and analysis, the 'Surfaces' function was employed by selecting the appropriate

marker channel (IBA1 for microglia), setting the 'Surface Detail Level' and 'Background

Subtraction', and using the 'Thresholding' slider to separate cells from the background. The

surface creation was finalized to generate surfaces encompassing the 3D structures of the

labelled cells. Morphometric analysis was exported in the 'Statistics' tab, quantifying

parameters such as volume and surface area. Images and videos of the 3D reconstructions

were exported using the snapshot and animation features in IMARIS v9.5. Comparisons of

these parameters between groups (Sham, MCAo, and HDACi-treated) were conducted to

evaluate the effects of ischemic stroke and HDACi treatment on microglial morphology.

Primary Antibodies

Antibody	Cat no.	Dilution	Source	Host	Clonality	Application
IBA1	ab48004	1:450	Abcam PLC, Cambridge, UK	Goat	Polyclonal	IF
NeuN	ab177487	1:600	Abcam PLC, Cambridge, UK	Rabbit	Monoclonal	IF
NeuN	ab177487	1:100	Abcam PLC, Cambridge, UK	Rabbit	Monoclonal	Spatial Transcriptomics
GFAP	ab4674	1:70	Abcam PLC, Cambridge, UK	Chicken	Polyclonal	Spatial Transcriptomics
H3K27me3	9733	1:400	Cell Signaling Technology, MA, USA	Rabbit	Monoclonal	IF
H3K4me3	9751	1:400	Cell Signaling Technology, MA, USA	Rabbit	Monoclonal	IF

210
211 **Secondary Antibodies**

Antibody	Cat no.	Dilution	Source	Host	Conjugate	Application
Donkey Anti-Goat (Alexa Fluor 488)	ab150133	1:600	Abcam PLC, Cambridge, UK	Donkey	Alexa Fluor 488	IF
Donkey Anti-Rabbit (Alexa Fluor 647)	ab150063	1:600	Abcam PLC, Cambridge, UK	Donkey	Alexa Fluor 647	IF
Donkey Anti-Rabbit (Alexa Fluor 647)	ab150063	1:100	Abcam PLC, Cambridge, UK	Donkey	Alexa Fluor 647	Spatial Transcriptomics
Goat Anti-Chicken (Alexa Fluor 488)	ab150169	1:100	Abcam PLC, Cambridge, UK	Goat	Alexa Fluor 488	Spatial Transcriptomics
DAPI	D9542-10MG	1:2000	Sigma-Aldrich			Nuclear staining (spatial transcriptomics)
Hoechst 33342	R37605	2drops/ml	Invitrogen			Nuclear staining (IF)

Spatial Transcriptomics Pipeline

Fresh-frozen brain tissue samples from the three groups (n=3) were sectioned at 10 µm

thickness using a cryostat (Leica Instruments GmbH, Germany). Tissue optimization was

performed with TO slides (part no. 3000394, 10x Genomics, California, USA) and reagents as

per 10x Genomics user guide CG000238-RevE and involved testing permeabilization

durations, with six minutes selected based on tissue morphology preservation and RNA

quality, assessed with a Bioanalyzer (Bioanalyzer 2100 system, Agilent, Santa Clara, CA,

USA) (Figure 4 Supplementary Figure 1, 2). The sections were placed on Visium Spatial Gene

Expression slides (part no. 2000233, 10x Genomics, California, USA) ensuring precise

alignment and adhesion to capture areas for spatial mapping. Immunofluorescence staining

(as per 10x Genomics user guide CG000312-RevD) was performed using antibodies against

neuronal nuclei (NeuN) and glial fibrillary acidic protein (GFAP), with DAPI for nuclear staining,

to correlate spatial gene expression data with specific cell types and anatomical features.

Antibody concentrations and incubation times were optimized through preliminary titration

experiments as advised by 10x standard protocols, to ensure specificity and maximal signal-

to-noise ratios. Following immunofluorescence and tissue preparation, in situ reverse

transcription on the Visium Spatial slides was performed to convert localized mRNA into cDNA

while retaining spatial coordinates using unique molecular identifiers (UMIs) and spatial

barcodes. Library preparation, conducted according to Visium protocols (10x Genomics user

guide CG000239-RevF), included second-strand synthesis, adaptor ligation, and amplification

steps. Libraries were quantified and validated for quality using Novogene's (NovogeneAIT

Genomics, Singapore Pte Ltd) sequencing pipeline for their Illumina platform, to ensure

suitability for high-throughput sequencing. Sequencing was performed to a depth of 50 million

reads to cover the complexity of the brain transcriptome and ensure spatial resolution of gene

expression patterns. Sequenced spatial libraries were processed using 10X Genomics

SpaceRanger 2.0.1, aligning reads to the reference genome and attributing them to specific

spatial locations on tissue sections based on their spatial barcodes. Spatial transcriptomic

data was exported from Loupe Browser 6.0 into Microsoft Excel for consolidation, upload and

visualization; systematically organizing metrics such as Log2 Fold Change (Log2FC) values

for genes of interest across different tissue sections and experimental conditions. Data

wrappers such as venny (Oliveros, 2007), S.R. Plot (Tang *et al.*, 2023) and heatmapper.ca

(Babicki *et al.*, 2016) were utilized for downstream data processing and visualization. The

Kyoto Encyclopedia of Genes and Genomes (KEGG), the Database for Annotation,

Visualization, and Integrated Discovery (DAVID), and Gene Ontology (GO) profiler were

employed to enrich and categorize gene expression data within biological pathways, functional

annotations, and cellular component. Data was visualized through S.R Plot (Tang *et al.*, 2023)

with clusterprofiler and pathview packages in the form of enrichment bubble and ontology bar

plots. Venny, an interactive tool for comparing lists (Oliveros, 2007) was utilized to generate

comparative lists of upregulated, downregulated and rescued genes across the biological

groups with spatial segregation of microglial transcriptomes. Heatmaps were generated using

Heatmapper.ca (Babicki *et al.*, 2016) to visualize Log2FC data exported from MS Excel

organized into selected pathways from enrichment analyses, providing a graphical

representation of gene expression variations across different brain regions and treatment

groups.

Statistical analyses

For infarct area evaluation from TTC stained brain slices, one-way ANOVA

(F-statistic=32625.11, p=2.74 $\times$ 10^{-9}) and independent two-sample t-tests ($<1\times10^{-6}$) were

performed on the calculated non-viable (infarcted) areas from all three biological groups (n=3).

The high F-statistic value confirms that the variability between group means is much larger

than the variability within groups while the t-tests confirm the findings from one-way ANOVA.

Similarly, independent two-sample t-tests revealed significant differences (<0.05) in IMARIS

morphometry data focusing on microglial surface area; microglial volume exhibited significant

differences (<0.005) as depicted in data between Sham and MCAo, as well as between MCAo

and MCAo+HDACi groups. Spatial transcriptomics data processed using proprietary 10x

software, SpaceRanger 2.0.1 and Loupe Browser 6.0 utilizes a statistical framework based on

sSeq for differential expression analysis (Yu, Huber and Vitek, 2013), Moran's I for spatial

enrichment (Getis, 2007) and Benjamini-Hochberg correction for controlling false discovery

rates (p<0.05) natively in the transcriptomics pipeline as described in 10x Genomics

repositories. Online data wrappers were fed downstream data exported from Loupe Browser

6.0 by filtering for genes limited to the selected pathways within the microglial/neuronal

spatiotranscriptomes by MS Excel's VLOOKUP function and data was visualized in the form

of heatmaps (heatmapper.ca), enrichment bubble plots, combined Gene Ontology bar plots,

CNETplots for KEGG pathways-targets enriched (S.R. Plot) and Venn diagrams for

dysregulated gene expression across microglial spatiotranscriptomes (Venny).

Results

HDACi treatment reduces ischemic infarct volume in the brain and enhances

preliminary locomotor function of mice post-ischemic insult.

HDACi was administered intra-peritoneally to mice undergoing 1-hour ischemia and 24-hour

reperfusion periods in two doses. The first dose at 1-hour pre-ischemic insult to prime cells

and the subsequent dose post 6-hours of ischemic insult to effectively influence reperfusion

mediated injury. As reported earlier, MCAo was performed for 1h and reperfusion post-

ischemia was done for 24h prior to euthanasia (Fig.1A) (Patnala et al., 2017). Neurological

viability and infarct volume were assessed through toluidine blue (Nissl) staining and TTC

(2,3,5-Triphenyltetrazolium Chloride) (Fig.1B-C). From the staining, the extent of ischemic

penumbra at brain regions of interest (ROI)'s such as hippocampus, striatum, thalamus and

the cortex were identified to proceed with assessment of microglia-neuron interaction in

regions (Fig.1B). Evaluation of basic locomotor activity in mice prior to euthanasia (Fig.1D,

Fig.1 Supplementary Video 1) showed circling contralateral to the infarct, and reduced mobility

in the MCAo groups, while HDACi-treated mice did not exhibit circling and showed reduced

latency to move within the home cage towards enrichment shell. Substantive restoration of

viable neuronal soma was observed (Fig.1E-F) 24-hour after administration HDACi. Taken

together, the findings suggest a) successful modelling of ischemia in the mouse brain; along

with identification of regions of interest (ROIs) (hippocampus, thalamus, striatum and cortex)

in the ischemic penumbra, presenting as potential areas to target reperfusion mediated

neuronal damage. b) Gross evaluation of viable brain tissue and preliminary behavioural

performance of HDACi administration after MCAo showcases a potential restorative effect,

substantiating a need for further assessment of microglial contribution to ischemia-reperfusion

mediated neuroinflammation as well as neuroprotection.

HDACi ameliorates neurodegeneration in brain nuclei after ischemic stroke and inhibits

microglial aggregation.

The hippocampal CA3-dentate gyrus are particularly vulnerable regions of ischemic insult

(Einenkel and Salameh, 2024). Histological evaluation of toluidine blue stained sections of the

hippocampal CA3-dentate gyrus for neuronal and Nissl body aggregation, reveals a degree of

neuroprotection after HDACi administration (Fig.1E); preservation of Nissl bodies correlates

with enhanced neuronal survival (Smith et al., 2012), substantiating that HDACi confers

neuroprotection in vulnerable regions of the mouse brain. Confocal images (Fig.1F) showing

immunofluorescence expression of NeuN and IBA1 in neurons and microglia respectively, in

the cortex surrounding infarct periphery indicate recruitment of microglia in stroke brain while

there was a reduction in recruitment and alteration of microglial morphology in stroke brain

following HDACi treatment; implying a tempered inflammatory response that mitigates further

neuronal damage by fostering a more conducive environment for recovery (Lee et al., 2014).

HDACi treatment inhibits ischemia-induced microglial phenotype in mouse brain

Microglial cells that are ramified with slender processes in the sham mouse brain exhibit

amoeboid and reactive phenotypes in the stroke brain (Fig.2A). Upon HDACi administration,

microglial phenotypes shift towards a sham spectrum ramified phenotype with long and

slender processes (Fig.2A). Quantitative morphometric analysis by 3D-reconstruction of thick

tissue sections generated using IMARIS image analysis software (v9.5) reveals that microglial

surface area decreases while intracellular volume is increased in the stroke brain compared

to sham brain microglia (Fig.2B-C). However, HDACi treatment reverses the stroke microglial

phenotype closer to sham mice, suggesting that HDACi inhibits ischemia-induced phenotypic

changes in microglia (Fig.2B-C). The distinct microglial morphology observed in this study

represents transcriptomic profiles that may influence their role in inflammatory response and

neuroprotection through the course of pathophysiology after ischemic stroke.

HDACi inhibits ischemia-induced increase in nuclear expression of histone marks

H3K27me3 and H3K4me3 in microglia

Phenotypic heterogeneity of microglia in pathological conditions is regulated by epigenetics,

and various signaling molecules and transcription factors (Wang *et al.*, 2021). We have

previously shown that acetylation marks such as H3K9ac regulate the inflammatory response

of microglia in the stroke brain (Patnala *et al.*, 2017). In the present study, we have analyzed

the expression pattern of methylation marks such as H3K27me3 and H3K4me3 to understand

molecular mechanisms contributing to microglial phenotypic heterogeneity in the ischemic

brain. The nuclear expression of H3K27me3 (Fig.3A-B) and H3K4me3 (Fig.3C) appeared to

be remarkably elevated in microglia of the stroke brain. Upon HDACi administration, microglia

exhibit a more ramified phenotype with inhibition in nuclear expression of H3K27me3 and

H3K4me3, similar to expression patterns of sham brain microglia suggesting that microglial

phenotypic heterogeneity is influenced by histone mark status. Interestingly, we also found

neuronal cells in the hippocampal CA2-CA3 gyri (Fig.3A-B) exhibiting an inverse H3k27me3

expression pattern to microglia, suggesting an epigenetic feedback mechanism across

microglia-neuron regulatory network of histone status that warrants further study.

Mapping of spatial transcriptome of microglia in ischemic mouse brain following HDACi

administration

The spatial transcriptome profile of microglia in response to ischemic injury and HDACi

administration was analyzed using the 10x Genomics Visium (FF) platform. Fresh frozen brain

slices were immunolabelled and imaged for neurons (NeuN, red) and astroglia (GFAP, green)

along with DAPI (blue) on GEX Visium slides prior to library preparation for the spatial overlay

(Fig.4A-B). Immunoflourescence imaging and proprietary Loupe Browser v6.0 cluster overlay

with key ROIs, namely the hippocampus, thalamus, striatum, and cortex were marked for post

processing (Fig.4A). Stroke brain tissue reveals a loss of cellular heterogeneity, and distinct

organizational patterns present in sham brain tissue, suggesting the disruption of cellular

microenvironments and signaling networks upon ischemic insult (Fig.4B-C). Preliminarily, the

spatial transcriptomic data revealed 19 distinct cell populations based on marker gene

expression across the three experimental groups (Fig.4F) with a Uniform Manifold

Approximation and Projection (UMAP) plot demonstrating the spatial distribution of these

populations alongside the spatial overlay (Fig.4E, 4C). To specifically isolate the microglial

transcriptome, a sub-clustering approach through feature selection and thresholding in loupe

browser v6.5.0 was employed (Fig.4D) by leveraging a list of microglia signature genes

identified in recent studies (Gerrits *et al.*, 2020; Quintana *et al.*, 2022). This allowed for the

extraction of microglia-enriched spots from brain co-ordinates to generate distinct microglial

spatiotranscriptomic datasets within the four penumbra ROIs.

369   HDACi alters microglial spatiotranscriptome profiles following ischemic stroke

Differential expression analysis and pathway enrichment studies were carried out to

investigate microglial spatiotranscriptomics and their functional implications. The anatomic

regions of interest on the overlay of spatiotranscriptomic spots, focusing on the ischemic

penumbra has been demarcated using loupe browser v6.0 (Fig.5A). Four-set Venn diagrams

were drawn to identify region-specific, unique and commonly upregulated (Fig.5B)

/downregulated (Fig.5C) /rescued (Fig.5D) microglial genes across the ischemic penumbra in

the hippocampus, thalamus, striatum and cortex. Among the upregulated microglial genes

across the four regions of interest, 22% of genes (3544) was found to be commonly

upregulated after HDACi administration (Fig.5B). On the other hand, only 1.5% (190) of

microglial genes was found to be commonly downregulated across the four regions of interest

(Fig.5C). Additionally, the datasets of differentially expressed genes (both up/downregulated)

were processed to obtain microglial genes that were rescued closer to sham expression levels

(Fig.5D) following HDACi treatment.

The dysregulated microglial genes were subjected to KEGG pathway enrichment (Fig 5E) and

Gene Ontology (GO) (Fig.5F) analyses. Pathway analysis yields enrichment of key functional

pathways pivotal for neuroinflammatory response, mediating neurotrophic support, and

facilitating phagocytosis (Fig.5E) alongside other pathways that may be relevant to microglial

responses after ischemic stroke (Fig.5 Supplementary Figure 1). Regulatory signaling

pathways such as PI3K-Akt signaling (PIK3CA, PIK3CB, AKT1, AKT2, AKT3, PTEN), FC

gamma R-mediated phagocytosis (FCGR1, FCGR3, FCGR4, SYK, and PIP5K1C), MAPK

signaling (DUSP1, DUSP4, DUSP6, ERK1-2, cJUN, UCP4), TNF signaling (TNF, NFKB1,

RelA), and TGFβ signaling (TGFBR1, TGFBR2, SMAD2, SMAD3, SMAD4) pathways shed

light on the transformative influence of epigenetic regulators on microglial functionality in the

post-stroke brain environment. The genes from these pathways were found to be rescued in

each region of interest within the transcriptome of microglia after HDACi administration.

Thus, HDACi treatment results in a significant alteration of the microglial transcriptomic

landscape, with a spatial context. Pathway-enrichment (Fig.6 Supplementary File 1) and GO

analyses conducted for the microglial genes of hippocampus, thalamus, striatum, and cortex

(Fig.6 Supplementary Figure 1A-1B) revealed that HDACi alters, and rescues expression of

several key pathways involved in neuroprotection, inflammatory response and phagocytosis

after ischemic stroke (Fig.6A,6B,6C,6D). These analyses of microglial spatiotranscriptomes

shed light on the emphasis of these pathways describing their big-picture and region-specific

roles after ischemic insult. Heatmaps visualize the expression pattern of rescued transcripts

across the functionally relevant pathways, alongside radar plots delineating the top 50 rescued

targets from these pathways in each microglial spatiotranscriptome (Fig.6A,6B,6C,6D).

Confocal microscopy of microglia in the hippocampus (from Fig.2B), can be corroborated with

microglial spatiotranscriptome to draw parallels between HDACi-rescued microglial

transcriptome <> microglial phenotypic heterogeneity in the hippocampus (Fig 6A <> Fig.2B).

The spatiotranscriptomic profile of microglia across groups emphasizes the impact of HDACi

on microglial gene expression involved in neuroprotection. We also analyzed if HDACi

treatment improves neuroprotection by influencing the neuronal spatiotranscriptome in the

penumbra following ischemic stroke. The rescued spatiotranscriptome of penumbral neurons

reveal a distinct inverse expression pattern across groups in enriched pathways (Fig.7B,7C),

between ischemia-affected and HDACi treated mouse brains (Fig.7A) with broader

implications for neuroprotection, synaptic remodeling, and plasticity. This suggests a granular

modulatory effect of HDACi that is potentially influenced by the local microenvironment and

neuronal signaling to remodel microglial expression profiles.

The findings highlight the spatial and molecular heterogeneity of microglial responses to

ischemic injury and HDAC inhibition, emphasizing the therapeutic capabilities of epigenetic

modulation in ischemic stroke. The identified pathways and rescued genes provide a robust

framework for further exploration of microglial-neuron interactions and the development of

targeted therapies.

Discussion

Recent advancements in spatial transcriptomics analysis have enriched our ability to dissect

stroke pathophysiology by providing spatial resolution to gene expression patterns in neurons

and glial cells including microglia. In the current study, integration of spatial transcriptomics

with advanced imaging and reconstruction techniques has offered a comprehensive view of

microglial dynamics in the stroke brain. The MCAo model recapitulates human physiology to

facilitate investigating microglial responses within the ischemic brain (Chen *et al.*, 2009; Ansari

et al., 2011), in particular the penumbra regions including the hippocampus, thalamus, striatum

and cortex. Microglial activation in the stroke brain appears to influence the neuronal

transcriptome, which is substantiated by the dysregulated transcriptional responses observed

in microglia and neurons of the stroke brain. This altered microglial response, which is

regulated by various signalling molecules and epigenetic mechanisms, can be ameliorated

through epigenetic intervention as observed in the present study.

The Influence of Transcription Factors on phenotypic heterogeneity of Microglia

Microglial cells are dynamic and exhibit region- and time-dependent phenotypic heterogeneity

in response to neuropathological conditions such as neuroinflammation and

neurodegeneration including ischemic stroke (Hickman *et al.*, 2018; Li *et al.*, 2020). In the

current study, differential expression of microglial spatiotranscriptomes associated with

phenotypic heterogeneity of microglia in response to ischemic stroke, particularly under the

modulation of HDAC inhibition was observed. This variation in microglial phenotypes

elucidates the interplay of transcription factors, cell-specific receptors and microglia-neuron

interaction; outlining the complexity behind microglial polarization such as neuroprotective or

neurotoxic phenotypes in the post-stroke brain (Olah *et al.*, 2011; Haroon, Miller and

Sanacora, 2016; Dubbelaar *et al.*, 2018).

In our microglial spatiotranscriptomic analysis, ischemic stroke was found to alter key

signalling molecular pathways involved in neurotrophic support, phagocytic activity and

neuroinflammation including PI3K-Akt signaling, FC gamma R-mediated phagocytosis, MAPK

signaling, TNF signaling, and TGFβ signaling pathways and transcription factors regulating

immune response such as NF-κB, STAT1/STAT3, Nrf, and PPAR family (Yang *et al.*, 2018;

Simpson and Oliver, 2020; Li *et al.*, 2024). Epigenetic mechanisms such as post-translational

histone modifications provide an additional layer of regulation to elicit cell-specific and stimuli

specific immune responses (Allis and Jenuwein, 2016) in the stroke brain.

Post-translational histone modifications determine basic transcriptional rates and the

magnitude of gene expression (Bannister and Kouzarides, 2011). Among the two major

histone modifications namely histone acetylation and histone methylation, histone acetylation

is associated with increased accessibility to transcriptional machinery leading to increased

gene expression (Marmorstein and Zhou, 2014), whereas histone methylation is more

complex in regulation of gene expression through repression. Different histone modification

patterns of specific marks at different sites have been shown to play distinct biological roles;

this prompted us to analyze the spatial transcriptome of microglia distributed in infarct

penumbra of the stroke brain upon HDACi administration to understand the epigenetic control

of microglia-mediated neuropathological outcomes in the stroke brain.

Modification in histone acetylation and deacetylation status, is maintained by histone

acetyltransferase (HATs) and histone deacetylase (HDACs) enzymes respectively (Yang and

Seto, 2007). Several studies have shown that inhibition of HDAC exerts immunomodulatory

effects, specifically suppressing pro-inflammation and inducing anti-inflammatory signaling

pathways (Adcock, 2007; Kulthinee *et al.*, 2022). We have previously demonstrated the

HDACi-specific immunosuppression of pro-inflammatory mediators (TNFα and NOS2) and

upregulation of IL10, an anti-inflammatory mediator in microglia in ischemic stroke mice model

exerting neuroprotection (Patnala *et al.*, 2017).

Dynamic regulation of repressive histone methylation marks, H3K27me3 and H3K4me3 within

microglia play a significant role in their response to ischemic injury and subsequent recovery

processes (Cheray & Joseph, 2018; Tang et al., 2014). In the present study, markedly elevated

expression of these markers were observed in microglia that are distributed in the penumbra

of ischemic brain. The increased nuclear expression of these histone marks in microglia

appears to reflect a response to the ischemic environment-induced transcriptional activity

which may be associated with changes in migration, phagocytosis and phenotypic

heterogeneity inferred through the spatial transcriptome of microglia. Remarkably, this

ischemia-induced expression of H3K27me3 and H3K4me3 was found to be reverted to near

sham levels in microglia upon HDACi treatment, suggesting epigenetic regulation of

H3K27me3 and H3K4me3 downstream pathways that influence recovery mechanisms.

Further HDACi-induced suppression of H3K27me3 in microglia of ischemic brain may be

associated with activation of its upstream targets such as Enhancer of Zeste Homolog2

(EZH2) (Suppl.tab.1_averaged_MicroglialLog2FC_10X.xslx) a catalytic subunit of the

Polycomb Repressive Complex 2 (PRC2) (Ayata *et al.*, 2018) which has been shown to play

important roles in the epigenetic regulation of phagocytic clearance and pruning of synaptic

activity in microglia (Hong, Dissing-Olesen and Stevens, 2016; Ayata *et al.*, 2018). Further,

since the PRC2 complex, including EZH2 has been shown to regulate TNFα and TGFβ

signalling pathways (Penas and Navarro, 2018; Chen *et al.*, 2019), the increase in EZH2

expression post HDACi treatment in microglia may be responsible for altered expression of

both inflammatory and neurotrophic regulators such as TNFα and TGFβ signalling observed

in the current study.

Epigenetic Regulation over Pathway-Specific Microglial transcriptomes involved in

Neuroprotection, Neuroinflammation and Phagocytosis in the Stroke Brain

Impact of HDAC inhibition on microglia in stroke brain suggests spatially defined

transcriptomic changes as the results highlight the spatial heterogeneity of microglial

transcriptomes within various regions of the stroke brain. Notably this was evident as pathways

crucial to neuroprotection in microglia of stroke brain after HDACi treatment exhibit a

transcriptomic signature similar to that of the sham brain.

The TGF-β signaling pathway, central to immune modulation and cell differentiation, displayed

marked upregulation of TGFβ receptors and SMAD family genes (e.g., TGFβR1, TGFβR2,

SMAD 2-4) in microglia of stroke brain following HDACi treatment. In TGFβ signaling, TGFβR-

Smad family complexes regulate the expression of TGFβ target genes (Chandra *et al.*, 2012;

Taylor *et al.*, 2017; Du *et al.*, 2018) (Lin *et al.*, 2017; Zöller *et al.*, 2018; Zhu *et al.*, 2023); the

surge of TGF-β signaling pathway genes suggests an amplified anti-inflammatory and

regenerative response in microglia, potentially enhancing their interactions with neurons to

support tissue repair and functional recovery in stroke brain following HDACi treatment. HDACi

has been found to upregulate many other anti-inflammatory genes including STAT3 and
PPARγ indicating an induction of anti-inflammatory response of microglia in the stroke brain
(Patnala et al., 2017).

Further, HDACi treatment restored expression of several other TGF-β signaling pathway
molecules including BMP and activin receptor membrane-bound inhibitor (*BAMBI*) which
interferes with TGFβ or BMP pathway transduction by forming heterodimers
with *TGFβRI/BMPRI* (Chen *et al.*, 2023). The involvement of *BAMBI*, and various BMPs
highlights the diverse role of TGF-β signaling through microglia, influencing not only immune
responses but also neurogenesis, and synaptic plasticity (Meyers and Kessler, 2017).

It has been well established that microglia contribute to neuroinflammation through excessive
secretion of proinflammatory mediators in the stroke brain (Jayaraj *et al.*, 2019). The activation
of proinflammatory mediators, is regulated by mitogen-activated protein kinases (MAPK)
comprising of extracellular signal regulated kinase (ERK), c-Jun N-terminal kinases (JNK),
and p38 and nuclear factor kappa beta (NF-κβ) gene expression (Kaminska, Mota and Pizzi,
2016). HDAC inhibition has been shown to modulate the p38 MAPK-MK2 signaling cascade,
regulating microglial activation and cytokine production (Kaminska *et al.*, 2009; Kaminska,
Mota and Pizzi, 2016). In the present study, many of the MAPK signalling pathway genes were
found to be altered in microglia following ischemia and HDACi treatment. Further, in microglia
of stroke brain the downregulation of c-Jun upon HDACi administration appears to modulate
the neuroinflammatory response, promoting neuroprotection (Waetzig *et al.*, 2005). This
downregulation of cJun could be mediated by HDACi-induced activation of MKP-1 (DUSP1)
which has been shown to inactivate MAP Kinases (Eljaschewitsch *et al.*, 2006). Overall,
HDACi appears to shift microglial activation towards neuroprotective states, reducing
neuroinflammation and promoting neuroprotection through epigenetic modifications.

Further, HDACi-induced upregulation of PI3K-AKT pathway in microglia of stroke brain

appears to associated with neuroprotection since we have previously demonstrated that

microglial PI3K-AKT-BDNF pathway regulates memory and learning in neurons via long term

potentiation (LTP) (Saw *et al.*, 2020). In addition, HDACi treatment seems to potentiate an

array of neuroprotective signaling cascades such as through the upregulation of microglial

TrkB, a neurotrophin receptor tyrosine kinase which is a potent regulator of hippocampal LTP

(Minichiello, 2009).

Microglia play a crucial role in clearing debris from the stroke microenvironment through

phagocytosis to enhance neuroprotection (Wang, Leak and Cao, 2022). This was evident as

expression of FC-γ receptor-mediated phagocytosis pathway genes such as FCGR1, FCGR3,

FCGR4, SYK, and PIP5K1C was found to be upregulated in microglia of stroke brain in the

current study. Interestingly, expression of these genes was further upregulated upon HDACi

treatment indicating that HDACi enhances microglial response to ischemia and capacity for

clearing cellular debris, limiting secondary neurodegeneration and inflammation (Qiao, Liu and

Qie, 2023). Overall, HDACi-mediated phagocytic activity of microglia potentially ameliorates

exacerbated neuroinflammation in the penumbra of the stroke brain (Lai, Zhang and Wang,

2014; Belov Kirdajova *et al.*, 2020). Spatially, these transcripts in hippocampal microglia

emphasize their role in mitigating damage and promoting recovery in regions critical for

cognitive functions.

The microglial transcriptomic changes induced by HDACi in different penumbral regions of the

stroke brain showcase a shift in microglial functionality from a proinflammatory phenotype to

a phenotype which improves brain repair and regeneration by enhancing phagocytosis and

reducing neuroinflammation.

This study also shows that HDACi altered the neuronal spatiotranscriptome to promote

resilience and plasticity within the ischemic brain environment (Ganai, Ramadoss and

Mahadevan, 2016; Kitahara *et al.*, 2021). The inverse expression patterns in neurotrophic and

neuroprotective pathways of these neurons in sham and stroke brains suggest broader

implications for brain plasticity through upstream epigenetic mechanisms aimed at

counterbalancing ischemia-associated excitotoxicity and neuroinflammatory response to

promote tissue repair and reduce neuronal susceptibility to reperfusion injury (Delcuve, Khan

and Davie, 2012; Kitahara *et al.*, 2021).

In conclusion, our high-throughput spatio-transcriptomic analysis unravels the mechanisms by

which a) microglia contribute to ischemic pathology with region-specificity and b) HDACi

ameliorates reperfusion mediated ischemic-injury to neurons through modulation of microglial

transcriptome. These findings emphasize an interplay between epigenetic regulation and

spatial transcriptomic changes altering communication networks between microglia and

neurons. Understanding the molecular mechanisms by which HDACi induced phenotypic

changes together with altered spatial transcriptomic landscape in microglia of stroke brain

would provide insight into development of potential microglia-mediated therapeutic strategies

for brain recovery post ischemia.

Funding

This study was funded by the Ministry of Education, Singapore (MOE Tier2, A-0002277-00-00 awarded to Professor S T Dheen).

Acknowledgements

Components of Figure 1 have been created with BioRender.com.

Author Contribution

Data acquisition: **KJ, RK**. Data analysis and interpretation: **KJ, ST.** Study design and conceptualization: **ST.** Writing of the original draft: **KJ, ST.** Reviewing & editing of manuscript: **all authors.** Supervision: **ST.**

Data Availability:

High-throughput, next-generation sequencing data from spatial transcriptomic libraries in this study have been submitted to the NCBI Sequence Read Archive (SRA) under BioProject PRJNA1144233.

Conflict of Interest Declaration

The authors declare that they have conflicts of interest related to the study.

Ethics Approval

All experimental procedures performed *in vivo* were approved by the National University of Singapore Animal Care and Use Committee.

Bibliography

1. Adcock, I. M. (2007) 'HDAC inhibitors as anti-inflammatory agents', *British Journal of Pharmacology*. Wiley, 150(7), p. 829. doi: 10.1038/SJ.BJP.0707166.

2. Allis, C. D. and Jenuwein, T. (2016) 'The molecular hallmarks of epigenetic control', *Nature Reviews Genetics*. Nature Publishing Group, 17(8), pp. 487–500. doi: 10.1038/nrg.2016.59.

3. Ansari, S. *et al.* (2011) 'Intraluminal middle cerebral artery occlusion (MCAO) model for ischemic stroke with laser doppler flowmetry guidance in mice', *Journal of Visualized Experiments*, (51), pp. 8–11. doi: 10.3791/2879.

4. Ayata, P. *et al.* (2018) 'Epigenetic regulation of brain region-specific microglia clearance activity', *Nature Neuroscience*. Nature Publishing Group, 21(8), pp. 1049–1060. doi: 10.1038/s41593-018-0192-3.

5. Babicki, S. *et al.* (2016) 'Heatmapper: web-enabled heat mapping for all', *Nucleic acids research*. Nucleic Acids Res, 44(W1), pp. W147–W153. doi: 10.1093/NAR/GKW419.

6. Bannister, A. J. and Kouzarides, T. (2011) 'Regulation of chromatin by histone modifications', *Cell Research 2011 21:3*. Nature Publishing Group, 21(3), pp. 381–395. doi: 10.1038/cr.2011.22.

7. Belov Kirdajova, D. *et al.* (2020) 'Ischemia-Triggered Glutamate Excitotoxicity From the Perspective of Glial Cells', *Frontiers in Cellular Neuroscience*. Frontiers Media SA, 14. doi: 10.3389/FNCEL.2020.00051.

8. Chandra, M. *et al.* (2012) 'Nuclear Translocation of Type I Transforming Growth Factor β Receptor Confers a Novel Function in RNA Processing', *Molecular and Cellular Biology*. Taylor & Francis, 32(12), p. 2183. doi: 10.1128/MCB.00320-12.

9. Charlton, T. *et al.* (2023) 'Brain-derived neurotrophic factor (BDNF) has direct anti-inflammatory effects on microglia', *Frontiers in cellular neuroscience*. Front Cell Neurosci, 17. doi: 10.3389/FNCEL.2023.1188672.

10. Chen, J. *et al.* (2009) *Animal models of acute neurological injuries*, *Animal Models of Acute Neurological Injuries*. doi: 10.1007/978-1-60327-185-1.

11. Chen, J. *et al.* (2019) 'EZH2 inhibitor DZNep modulates microglial activation and protects against ischaemic brain injury after experimental stroke', *European Journal of Pharmacology*, 857. doi: 10.1016/j.ejphar.2019.172452.

12. Chen, X. *et al.* (2023) 'BMP and activin receptor membrane bound inhibitor: BAMBI has multiple roles in gene expression and diseases (Review)', *Experimental and Therapeutic Medicine*, 27(1). doi: 10.3892/etm.2023.12316.

13. Cheray, M. and Joseph, B. (2018) 'Epigenetics control microglia plasticity', *Frontiers in Cellular Neuroscience*, 12(August), pp. 1–13. doi: 10.3389/fncel.2018.00243.

14. Delcuve, G. P., Khan, D. H. and Davie, J. R. (2012) 'Roles of histone deacetylases in epigenetic regulation: Emerging paradigms from studies with inhibitors', *Clinical Epigenetics*. BioMed Central, 4(1), pp. 1–13. doi: 10.1186/1868-7083-4-5/FIGURES/2.

15. Donkor, E. S. (2018) 'Stroke in the 21st Century: A Snapshot of the Burden, Epidemiology, and Quality of Life', *Stroke Research and Treatment*. Hindawi Limited, 2018. doi: 10.1155/2018/3238165.

16. Du, X. *et al.* (2018) 'SMAD4 feedback regulates the canonical TGF-β signaling pathway to control granulosa cell apoptosis', *Cell Death & Disease 2018 9:2*. Nature Publishing Group, 9(2), pp. 1–12. doi: 10.1038/s41419-017-0205-2.

17. Dubbelaar, M. L. *et al.* (2018) 'The Kaleidoscope of Microglial Phenotypes', *Frontiers in immunology*, 9(July), p. 1753. doi: 10.3389/fimmu.2018.01753.

18. Einenkel, A. M. and Salameh, A. (2024) 'Selective vulnerability of hippocampal CA1 and CA3 pyramidal cells: What are possible pathomechanisms and should more attention be paid to the CA3 region in future studies?', *Journal of Neuroscience Research*. John Wiley & Sons, Ltd, 102(1), p. e25276. doi: 10.1002/JNR.25276.

19. Eljaschewitsch, E. *et al.* (2006) 'The endocannabinoid anandamide protects neurons during CNS inflammation by induction of MKP-1 in microglial cells', *Neuron*. Neuron, 49(1), pp. 67–79. doi: 10.1016/J.NEURON.2005.11.027.

20. Feigin, V. L. *et al.* (2022) 'World Stroke Organization (WSO): Global Stroke Fact Sheet 2022', *International Journal of Stroke*. SAGE Publications Inc., 17(1), pp. 18–29. doi:

10.1177/17474930211065917.

21. Fessler, E. *et al.* (2013) 'Potential Roles of HDAC Inhibitors in Mitigating Ischemia-
induced Brain Damage and Facilitating Endogenous Regeneration and Recovery',
Current Pharmaceutical Design, 19(28), pp. 5105–5120. doi:
10.2174/1381612811319280009.

22. Ganai, S. A., Ramadoss, M. and Mahadevan, V. (2016) 'Histone Deacetylase (HDAC)
Inhibitors - Emerging Roles in Neuronal Memory, Learning, Synaptic Plasticity and
Neural Regeneration', *Current Neuropharmacology*. Bentham Science Publishers,
14(1), p. 55. doi: 10.2174/1570159X13666151021111609.

23. Garcia, J. H. *et al.* (1995) 'Neurological deficit and extent of neuronal necrosis
attributable to middle cerebral artery occlusion in rats: Statistical validation', *Stroke*.
Lippincott Williams and Wilkins, 26(4), pp. 627–635. doi:
10.1161/01.STR.26.4.627/ASSET/D6F51A65-684F-4409-A4B6-
F5F58099E815/ASSETS/GRAPHIC/HS0451886005.GIF.

24. Gerrits, E. *et al.* (2020) 'Transcriptional profiling of microglia; current state of the art
and future perspectives', *GLIA*, 68(4). doi: 10.1002/glia.23767.

25. Getis, A. (2007) 'Reflections on spatial autocorrelation', *Regional Science and Urban
Economics*. North-Holland, 37(4), pp. 491–496. doi:
10.1016/J.REGSCIURBECO.2007.04.005.

26. Goldlust, E. J. *et al.* (1996) 'Automated measurement of infarct size with scanned
images of triphenyltetrazolium chloride-stained rat brains', *Stroke*. Lippincott Williams
and Wilkins, 27(9), pp. 1657–1662. doi:
10.1161/01.STR.27.9.1657/ASSET/67D2E47D-FF57-447E-A968-
A4899AD60E1E/ASSETS/GRAPHIC/0011F5.JPEG.

27. Gupta, N. *et al.* (2018) 'Recent progress in therapeutic strategies for microglia-
mediated neuroinflammation in neuropathologies', *Expert Opinion on Therapeutic
Targets*. Taylor & Francis, 22(9), pp. 765–781. doi: 10.1080/14728222.2018.1515917.

28. Haroon, E., Miller, A. H. and Sanacora, G. (2016) 'Inflammation, Glutamate, and Glia:

A Trio of Trouble in Mood Disorders', *Neuropsychopharmacology 2017 42:1*. Nature Publishing Group, 42(1), pp. 193–215. doi: 10.1038/npp.2016.199.

29. Hickman, S. *et al.* (2018) 'Microglia in neurodegeneration', *Nature Neuroscience*. Springer US, 21(10), pp. 1359–1369. doi: 10.1038/s41593-018-0242-x.

30. Hong, S., Dissing-Olesen, L. and Stevens, B. (2016) 'New insights on the role of microglia in synaptic pruning in health and disease', *Current Opinion in Neurobiology*. Elsevier Current Trends, 36, pp. 128–134. doi: 10.1016/J.CONB.2015.12.004.

31. Jayaraj, R. L. *et al.* (2019) 'Neuroinflammation: Friend and foe for ischemic stroke', *Journal of Neuroinflammation*. BioMed Central Ltd., pp. 1–24. doi: 10.1186/s12974-019-1516-2.

32. Kaminska, B. *et al.* (2009) 'MAPK Signal Transduction Underlying Brain Inflammation and Gliosis as Therapeutic Target', *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology*. John Wiley & Sons, Ltd, 292(12), pp. 1902–1913. doi: 10.1002/AR.21047.

33. Kaminska, B., Mota, M. and Pizzi, M. (2016) 'Signal transduction and epigenetic mechanisms in the control of microglia activation during neuroinflammation', *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*. Elsevier, 1862(3), pp. 339–351. doi: 10.1016/J.BBADIS.2015.10.026.

34. Kitahara, M. *et al.* (2021) 'Exercise and pharmacological inhibition of histone deacetylase improves cognitive function accompanied by an increase of gene expressions crucial for neuronal plasticity in the hippocampus', *Neuroscience Letters*. Elsevier, 749, p. 135749. doi: 10.1016/J.NEULET.2021.135749.

35. Kulthinee, S. *et al.* (2022) 'Critical Functions of Histone Deacetylases (HDACs) in Modulating Inflammation Associated with Cardiovascular Diseases', *Pathophysiology*. Multidisciplinary Digital Publishing Institute (MDPI), 29(3), p. 471. doi: 10.3390/PATHOPHYSIOLOGY29030038.

36. Lai, T. W., Zhang, S. and Wang, Y. T. (2014) 'Excitotoxicity and stroke: Identifying novel targets for neuroprotection', *Progress in Neurobiology*. Pergamon, 115(C), pp.

157–188. doi: 10.1016/J.PNEUROBIO.2013.11.006.

37. Li, L. *et al.* (2020) 'Potential microglia-based interventions for stroke', *CNS Neuroscience & Therapeutics*. Blackwell Publishing Ltd, 26(3), pp. 288–296. doi: 10.1111/cns.13291.

38. Li, Y. *et al.* (2000) 'Intrastriatal transplantation of bone marrow nonhematopoietic cells improves functional recovery after stroke in adult mice', *Journal of Cerebral Blood Flow and Metabolism*. Lippincott Williams and Wilkins, 20(9), pp. 1311–1319. doi: 10.1097/00004647-200009000-00006/ASSET/IMAGES/LARGE/10.1097_00004647-200009000-00006-FIG2.JPEG.

39. Li, Y. *et al.* (2024) 'Sodium butyrate alleviates lead-induced neuroinflammation and improves cognitive and memory impairment through the ACSS2/H3K9ac/BDNF pathway', *Environment International*. Pergamon, 184, p. 108479. doi: 10.1016/J.ENVINT.2024.108479.

40. Lin, E. *et al.* (2017) 'Transforming growth factor-β signaling pathway-associated genes SMAD2 and TGFBR2 are implicated in metabolic syndrome in a Taiwanese population', *Scientific reports*. Sci Rep, 7(1). doi: 10.1038/S41598-017-14025-4.

41. Marmorstein, R. and Zhou, M. M. (2014) 'Writers and Readers of Histone Acetylation: Structure, Mechanism, and Inhibition', *Cold Spring Harbor Perspectives in Biology*. Cold Spring Harbor Laboratory Press, 6(7). doi: 10.1101/CSHPERSPECT.A018762.

42. Mergenthaler, P., Dirnagl, U. and Kunz, A. (2022) 'Ischemic Stroke: Basic Pathophysiology and Clinical Implication', *Neuroscience in the 21st Century: From Basic to Clinical: Third Edition*. Springer International Publishing, pp. 3807–3827. doi: 10.1007/978-3-030-88832-9_97/COVER.

43. Meyers, E. A. and Kessler, J. A. (2017) 'TGF-β Family Signaling in Neural and Neuronal Differentiation, Development, and Function', *Cold Spring Harbor Perspectives in Biology*. Cold Spring Harbor Laboratory Press, 9(8). doi: 10.1101/CSHPERSPECT.A022244.

44. Minichiello, L. (2009) 'Long-term potentiation Synaptic plasticity TrkB signalling

pathways in LTP and learning', *Nature Reviews Neuroscience*, 10.

45. Nagy, E. E. *et al.* (2020) 'Neuroinflammation and microglia/macrophage phenotype modulate the molecular background of post-stroke depression: A literature review', *Experimental and Therapeutic Medicine*. Spandidos Publications, 20(3), p. 2510. doi: 10.3892/ETM.2020.8933.

46. Olah, M. *et al.* (2011) 'Microglia Phenotype Diversity', *CNS & Neurological Disorders - Drug Targets*, 10(1), pp. 108–118. doi: 10.2174/187152711794488575.

47. Oliveros, J. C. (2007) *Venny. An Interactive Tool for Comparing Lists with Venn's Diagrams. - References - Scientific Research Publishing*, *Csic*.

48. Patnala, R. *et al.* (2017) 'HDAC Inhibitor Sodium Butyrate-Mediated Epigenetic Regulation Enhances Neuroprotective Function of Microglia During Ischemic Stroke', *Molecular Neurobiology*, 54(8), pp. 6391–6411. doi: 10.1007/s12035-016-0149-z.

49. Penas, C. and Navarro, X. (2018) 'Epigenetic modifications associated to neuroinflammation and neuropathic pain after neural trauma', *Frontiers in Cellular Neuroscience*. Frontiers Media S.A., p. 158. doi: 10.3389/fncel.2018.00158.

50. Qiao, C., Liu, Z. and Qie, S. (2023) 'The Implications of Microglial Regulation in Neuroplasticity-Dependent Stroke Recovery', *Biomolecules 2023, Vol. 13, Page 571*. Multidisciplinary Digital Publishing Institute, 13(3), p. 571. doi: 10.3390/BIOM13030571.

51. Qiu, M., Xu, E. and Zhan, L. (2021) 'Epigenetic Regulations of Microglia/Macrophage Polarization in Ischemic Stroke', *Frontiers in Molecular Neuroscience*. Frontiers Media S.A., 14, p. 697416. doi: 10.3389/FNMOL.2021.697416/BIBTEX.

52. Quintana, J. F. *et al.* (2022) 'Single cell and spatial transcriptomic analyses reveal microglia-plasma cell crosstalk in the brain during Trypanosoma brucei infection', *Nature Communications*, 13(1). doi: 10.1038/s41467-022-33542-z.

53. Saw, G. *et al.* (2020) 'Epigenetic regulation of microglial phosphatidylinositol 3-kinase pathway involved in long-term potentiation and synaptic plasticity in rats', *Glia*, 68(3), pp. 656–669. doi: 10.1002/glia.23748.

54. Simpson, D. S. A. and Oliver, P. L. (2020) 'ROS Generation in Microglia: Understanding Oxidative Stress and Inflammation in Neurodegenerative Disease', *Antioxidants*. MDPI AG, 9(8), p. 743. doi: 10.3390/antiox9080743.

55. Sun, J. *et al.* (2015) 'Neuroprotective Effect of Sodium Butyrate against Cerebral Ischemia/Reperfusion Injury in Mice', *BioMed Research International*. Hindawi Publishing Corporation, 2015. doi: 10.1155/2015/395895.

56. Tang, D. *et al.* (2023) 'SRplot: A free online platform for data visualization and graphing', *PLOS ONE*. Public Library of Science, 18(11), p. e0294236. doi: 10.1371/JOURNAL.PONE.0294236.

57. Tang, Y. *et al.* (2014) 'Jmjd3 is essential for the epigenetic modulation of microglia phenotypes in the immune pathogenesis of Parkinson's disease', *Cell Death and Differentiation*. Nature Publishing Group, 21(3), pp. 369–380. doi: 10.1038/cdd.2013.159.

58. Taylor, R. A. *et al.* (2017) 'TGF-β1 modulates microglial phenotype and promotes recovery after intracerebral hemorrhage', *Journal of Clinical Investigation*, 127(1). doi: 10.1172/JCI88647.

59. Traystman, R. J. (2003) 'Animal models of focal and global cerebral ischemia', *ILAR Journal*. Institute for Laboratory Animal Research, 44(2), pp. 85–95. doi: 10.1093/ILAR.44.2.85.

60. Waetzig, V. *et al.* (2005) 'c-Jun N-terminal kinases (JNKs) mediate pro-inflammatory actions of microglia', *Glia*. Glia, 50(3), pp. 235–246. doi: 10.1002/GLIA.20173.

61. Wang, L. *et al.* (2021) 'Epigenetic Modulation of Microglia Function and Phenotypes in Neurodegenerative Diseases', *Neural Plasticity*, 2021. doi: 10.1155/2021/9912686.

62. Wang, Y., Leak, R. K. and Cao, G. (2022) 'Microglia-mediated neuroinflammation and neuroplasticity after stroke', *Frontiers in Cellular Neuroscience*. Frontiers Media S.A., 16, p. 980722. doi: 10.3389/FNCEL.2022.980722/BIBTEX.

63. Wendimu, M. Y. and Hooks, S. B. (2022) 'Microglia Phenotypes in Aging and Neurodegenerative Diseases', *Cells*. Multidisciplinary Digital Publishing Institute

808   (MDPI), 11(13). doi: 10.3390/CELLS11132091.

64. Xu, Y. *et al.* (2024) 'Sodium butyrate activates the KATP channels to regulate the mechanism of Parkinson's disease microglia model inflammation', *Immunity, Inflammation and Disease*. John Wiley & Sons, Ltd, 12(3), p. e1194. doi: 10.1002/IID3.1194.

65. Yang, S. J. *et al.* (2018) 'The NLRP3 Inflammasome: An Important Driver of Neuroinflammation in Hemorrhagic Stroke', *Cellular and Molecular Neurobiology*. Springer US, 38(3), pp. 595–603. doi: 10.1007/s10571-017-0526-9.

66. Yang, X. J. and Seto, E. (2007) 'HATs and HDACs: From structure, function and regulation to novel strategies for therapy and prevention', *Oncogene*. doi: 10.1038/sj.onc.1210599.

67. Yu, D., Huber, W. and Vitek, O. (2013) 'Shrinkage estimation of dispersion in Negative Binomial models for RNA-seq experiments with small sample size', *Bioinformatics*. Oxford Academic, 29(10), pp. 1275–1282. doi: 10.1093/BIOINFORMATICS/BTT143.

68. Zhang, F. *et al.* (2022) 'Regulation of N6-methyladenosine (m6A) RNA methylation in microglia-mediated inflammation and ischemic stroke', *Frontiers in Cellular Neuroscience*. Frontiers Media S.A., 16, p. 955222. doi: 10.3389/FNCEL.2022.955222/BIBTEX.

69. Zhang, H. *et al.* (2019) 'Isosteviol Sodium Protects against Ischemic Stroke by Modulating Microglia/Macrophage Polarization via Disruption of GAS5/miR-146a-5p sponge', *Scientific Reports 2019 9:1*. Nature Publishing Group, 9(1), pp. 1–16. doi: 10.1038/s41598-019-48759-0.

70. Zhang, W. *et al.* (2021) 'Microglia-associated neuroinflammation is a potential therapeutic target for ischemic stroke', *Neural Regeneration Research*. Wolters Kluwer Medknow Publications, 16(1), p. 6. doi: 10.4103/1673-5374.286954.

71. Zhu, K. *et al.* (2023) 'Disrupting microglial TGF-β signaling triggers region-specific pathology in the spinal cord', *bioRxiv*. Cold Spring Harbor Laboratory, p. 2023.04.24.538074. doi: 10.1101/2023.04.24.538074.

72. Ziemka-Nalecz, M. *et al.* (2017) 'Sodium Butyrate, a Histone Deacetylase Inhibitor,

Exhibits Neuroprotective/Neurogenic Effects in a Rat Model of Neonatal Hypoxia-

Ischemia', *Molecular Neurobiology*, 54(7), pp. 5300–5318. doi: 10.1007/s12035-016-

0049-2.

73. Zöller, T. *et al.* (2018) 'Silencing of TGFβ signalling in microglia results in impaired

homeostasis', *Nature Communications 2018 9:1*. Nature Publishing Group, 9(1), pp.

1–13. doi: 10.1038/s41467-018-06224-y.

Figure Legend:

Figure 1: Middle cerebral artery occlusion and gross neurological status upon HDACi
treatment.

(A) Graphical pipeline describing ischemic stroke modelling in the rodent brain, ischemia-
associated microglia and phenotypes. (B) Toluidine blue (nissl) staining of brain tissue
sections with MCAo and Sham showcasing infarct extent and anatomical regions of interest
(ROI) covered under the penumbra. (C) Epigenetically mediated protection through HDACi
attenuates infarct volume 24hours post ischemic insult (TTC or 2,3,5-Triphenyltetrazolium
Chloride assessment of brain slices exhibit substantial recovery of viable brain tissue upon
NaBu administration) (n=3 in each experimental group). (D) Qualitative assessment of
ischemic affect through preliminary neurological deficit scoring (pNDS) demonstrates
enhanced locomotor activity without immobility or circling and reduced locomotor latency
reduction (to reach enrichment shell within home cage) after HDACi treatment (n=4 in each
experimental group). (E) Toluidine blue stained mouse brain sections of dentate gyrus neurons
in the Hippocampus exhibit variance in infarct intensity and neuronal damage post-ischemic
stroke with HDACi administration. (F) Immunostained confocal images of NeuN (channel AF
674, false-colored green) and IBA1 (channel AF488, false-colored red) appear to display
preserved neuronal organization in the penumbral cortex and reduced microglial aggregation
at the infarct periphery upon HDAC inhibition. Nucleus is stained with Hoechst 33342 (blue).

Figure 1 Supplementary Figure 1

TTC (2,3,5-Triphenyltetrazolium Chloride) treated mouse brain sections across biological
groups showcasing extent of infarct through viable (stained, red) tissue and non-viable
(unstained, white) tissue. (n=3 in each experimental group).

Figure 1 Supplementary Video 1

Assessment of ischemic stroke affect through preliminary neurological deficit scoring (pNDS); video capture of each biological group showcases mild-moderate immobility and circling behaviour in the MCAo group while sham and HDACi-treated groups exhibit reduced to nil locomotor defects such as circling and difficulty reaching enrichment shell within home cage.

Figure 2. Microglial morphometric assessment through surface reconstruction with IMARIS 9.5

(A) High-resolution confocal images of Microglia stained with IBA1 (channel AF 488, green) reveal diverse morphological phenotypes between biological groups (n=3 in each experimental group). (B) Individual microglial cells were segmented and processed across groups to obtain surface area and cellular volume morphometric data. 3D-reconstruction of microglial surfaces spotlights microglia with ramified and long, slender processes in HDACi inhibitor administered mouse brain reminiscent of sham brain microglia while MCAo brain microglia are amoeboid with substantial depth across the z-stack. (C) Microglial surface area and intracellular volume are inverse between stroke and epigenetically treated groups, presenting a phenotypic reversal of microglia in the penumbra.

Figure 2 Supplementary Video 1

Video capture of IMARIS 9.5 reconstructions (30μm thick, z-stacked sections), showcasing surface creation and segmentation of labelled microglial cells (IBA1, AF488, false-colored red) that are processed to obtain morphometric data such as surface area and cellular volume to evaluate phenotypic status of microglia in the stroke brain after HDACi treatment (n=3 in each experimental group)

Figure 3. HDAC inhibition and histone mark status in microglia after ischemic stroke

(A-B) Hyper-intense expression of H3K27me3 (channel 647, false-colored green) in MCAo

brain microglia (IBA1, channel 488, false-colored red) appear to be rescued towards sham

expression levels in hippocampal microglia from HDACi-treated stroke brains. Potential

shuttling of H3K27me3 suggests upstream involvement of the PRC2 complex and EZH2,

components mediating H3K27me3-related gene silencing upon ischemic insult. (C) Similar to

H3K27me3, H3K4me3 (channel 647, false-colored green) expression levels in sham and

HDACi-treated stroke brain tissue showcase similar expression patterns with stroke elevating

expression in microglial nuclei suggesting neuron-microglia regulatory networks activated by

HDACi administration.

Figure 3 Supplementary Figure 1

(A,B,C) Tilescan (10x) images of complete brain sections across biological groups; (D,E,F)

showcasing hippocampal expression of H3K27me3 (AF647, false-colored green) in microglia

stained with IBA1 (AF488, false-colored red) that is upregulated upon ischemic stroke and

ameliorated upon HDACi-treatment.

Figure 3 Supplementary Video 1

Video capture of (30μm thick, z-stacked sections) microglia stained with IBA1 (AF488, false-

colored red) and H3K27me3 (AF647, false-colored green) across all three biological groups,

showcasing elevation of histone mark expression and presence within the nuclei of microglial

cells.

Figure 4. Spatial transcriptome pipeline: regions of interest, population clusters and microglial spatiotranscriptomic signature with the Visium Fresh Frozen platform (10x Genomics)

(A) Immunofluorescence (IF) image and spatial (spot) overlay (B) of Visium fresh frozen sections and marker gene population clusters (C) (n=3 in each experimental group). (B) 10x Genomics proprietary Loupe Browser 6.0's default transcriptome overlay represented as k-clusters and population identities; (C) the distinct population clusters appear to lose organization upon ischemic insult. (D) Microglial spots enriched through microglia signature genes adapted from (Gerrits *et al.*, 2020; Quintana *et al.*, 2022) performed through Loupe browser 6.0's feature selection and thresholding to output microglia-specific spatial transcriptomes segmented based on regions on interest in the tissue section. (E) Heatmap of total spatial library with cell signature markers for processed brain samples exhibiting 19 population identities. (F) Uniform Manifold Approximation and Projection (UMAP) dimension reduction of all cell populations in the dataset.

Figure 4 Supplementary Figure 1

(A,B) Tissue optimization timepoint experiments for brain samples with 10x Genomics Visium-TO slides for evaluation of optimum cDNA permeabilization time points (10x UG CG000238-RevE) prior to proceeding with gene expression and library preparation of Visium fresh-frozen spatial libraries. 6 minutes of permeabilization was selected based on optimization time points evaluated. (C) Antibody titre evaluation for IF-staining of spatial tissue sections (10x UG CG000312-RevD) identified concentration of antibodies (table in methodology).

Figure 4 Supplementary Figure 2

Permeabilized Visium fresh-frozen slides and quality control (QC) of RNA integrity checks and

pre-cDNA amplification/ cleanup for visium libraries prior to preparation and sequencing.

Figure 5. Spatial segmentation of microglial transcriptomes was performed and

expression signatures across the ischemic penumbra to obtain functionally relevant

enriched pathways (KEGG).

(A) Grouping and segmentation of Penumbra regions of interest in *LoupeBrowser* for spatial

demarcation of microglia after ischemic stroke. (B, C, D) Four-set Venn diagrams of unique

and overlapping HDACi-Upregulated (B), Downregulated (C) and Rescued (D) Genes,

elucidate the homogenous and heterogenous marks in microglia across brain regions.

Pathway (Kegg) Enrichment (E) and Gene Ontology (F) analysis of genes dysregulated

homogenously across the four microglial spatiotranscriptomes are processed and showcase

potential microglial pathways of interest to analyze and target after ischemic stroke. (n=3 for

each experimental group)

Figure 5 Supplementary Figure 1

Alternative enriched (KEGG) pathways from commonly dysregulated genes of microglial

spatiotranscriptomes across all 4 ROIs (hippocampus, thalamus, striatum, cortex). These

pathways present additional targets and mechanisms that may be dysregulated throughout

penumbral microglia after ischemic stroke and HDACi-administration.

Figure 6. Microglial spatiotranscriptome expression of enriched, key biological

pathways after HDACi administration in ischemic stroke.

Heatmaps of microglial spatiotranscriptomes of each ROI, hippocampal (A), thalamic (I),

striatal (J), cortical (K) microglia exhibit differential profiles of selected biological pathways that

were enriched and involved in ischemic injury across the three biological groups (n=3 for each

experimental group). Rescued genes within identified pathways crucial for neuroprotective,

neuroinflammatory and phagocytosis mediation responses are represented, and top 50

targets represented, in the radar plot for each penumbra ROI.

Figure 6 Supplementary Figure 1A

Pathways enriched in HDACi-rescued, region-specific microglial spatiotranscriptomes

(hippocampus-A, thalamus-B, striatum-C, cortex-D). Evaluation of pathways rescued by

HDACi after ischemic injury, highlights pathways crucial for neuroprotective,

neuroinflammatory and phagocytosis mediation responses.

Figure 6 Supplementary Figure 1B

Gene Ontology analyses of region-specific, microglial spatiotranscriptomes (hippocampus-A,

thalamus-B, striatum-C, cortex-D) exhibit differential GO terms in respective compartments

(BP, CC and MF) involved in HDACi-mediated rescue after ischemic injury in microglia.

Figure 6 Supplementary Figure 2

Heatmaps representing all dysregulated genes within selected biological pathways (HDACi vs

MCAo Log2fc difference>1.0, Euclidean distance metric with hierarchical clustering through

complete linkage) in each penumbra microglial spatiotranscriptomes (hippocampus-A,

thalamus-B, striatum-C, cortex-D). Microglial heterogeneity in expression patterns of these

pathways further highlights the specificity of microglial responses to ischemic stroke and

HDACi therapy based on spatial contexts within the mouse brain.

Figure 6 Supplementary Figure 3

Circular Network Plot (cnetplot) of HDACi-rescued pathways and targets in each microglial spatiotranscriptome (hippocampus-A, thalamus-B, striatum-C, cortex-D) post-ischemic stroke. Genes (nodes) are positioned around the plot, sized by log2 fold change (log2FC) and colored from red (upregulated) to blue (downregulated). Pathway nodes are centrally located, sized by the number of associated genes, and color-coded by functional categories. Key pathways identified in the study including "MAPK signaling," "PI3K-Akt signaling," and "Fc gamma R-mediated phagocytosis." are depicted with lines indicating gene-pathway associations, illustrating the molecular interactions affected by HDACi treatment.

1000

Figure 7. Penumbra Neuronal Spatiotranscriptome

(A-B) Spatiotranscriptome of penumbral neurons showcases enriched neuronal pathways relevant for survival through HDACi-mediated rescue after ischemic stroke. (C) Distinct, inverse expression profiles after HDACi treatment can be observed, indicating context-dependent microglia-neuron interaction upon ischemic stroke and epigenetic regulation through HDACi. (n=3 for each experimental group)

1007

Figure 2. Supplementary File 1

Morphometry data table consisting of IMARIS *Surfaces* feature metrics (Volume, Surface Area of microglial cells)

1011

1012

Figure 6. Supplementary File 1

Enriched Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways of interest from individual microglial spatiotranscriptomes.

A

Mouse Model → Unilateral Middle Cerebral Artery Occlusion → Infarct extent coronal section of mouse brain → Spatial transcriptomic analysis

B

Sham | MCAo | MCAo+HDACi

Cortex
Hippocampus
Thalamus
Striatum
1mm

C

Infarct Extent Reduction after HDACi

* P < 0.001

Area (mm2) — SHAM, MCAo, MCAo_HDACi

■ Avg Non-Viable Area (mm²) ■ Avg Viable Area (mm²)

D

pNDS

Deficit Score

— Sham — MCAo — HDACi

E

Sham | MCAo | MCAo+HDACi

25µm

F

Sham | MCAo | MCAo+HDACi

50µm

NeuN   IBA1   HOECHST 33342

A

Sham   MCAo   MCAo+HDACi

10 µm   10 µm   10 µm

B

Sham   MCAo   MCAo+HDACi

C

Microglial Surface Area (µm2)

Sham

MCAo

MCAo+HDACi

* P < 0.05

0   1000   2000   3000   4000   5000

Intracellular Volume (µm3)

Sham

MCAo

MCAo+HDACi

* P < 0.05

0   500   1000   1500   2000   2500   3000

A

HOECHST33342   H3K27me3   IBA1

Sham   MCAo   MCAo+HDACi

B

Sham   MCAo   MCAo+HDACi

20μm   20μm   20μm

C

HOECHST33342   H3K4me3   IBA1

Sham   MCAo   MCAo+HDACi

20μm   20μm   20μm

Microglial signature genes
(Quintana et al., 2022)
(Gerrits et al., 2020)

P2ry12	Tmem119
Sall1	Trem2
Trem119	P2ry12
Hexb	Mertk
Cx3cr1	Pros1
Fcrls	Aif1
Gpr43	Siglech
Cx3cr1	

HDACi Dysregulated and Rescued Genes across microglial spatiotranscriptomes

B **Upregulated** microglial transcripts upon HDACi administration after MCAo

C **Downregulated** microglial transcripts upon HDACi administration after MCAo

D **Rescued** microglial transcripts upon HDACi administration after MCAo

E Microglial pathways enriched in commonly dysregulated transcripts across all ROIs after HDACi administration in the ischemic penumbra

F GO: BP, CC, MF - commonly dysregulated spatiotranscriptome across ROIs of HDACi microglia in the ischemic penumbra

A

Sham   MCAo   MCAo+HDACi

250μm   250μm   250μm

NeuN **Gfap** **Dapi**

B

Pathway enrichment in HDACi-rescued Neuronal transcriptome within then Penumbra

PI3K-Akt signaling pathway
MAPK signaling pathway
Focal adhesion
Apoptosis
Notch signaling pathway
Ras signaling pathway
TGF-beta signaling pathway
Rap1 signaling pathway
Neurotrophin signaling pathway
mTOR signaling pathway
VEGF signaling pathway
FoxO signaling pathway
TNF signaling pathway
HIF-1 signaling pathway
AMPK signaling pathway
Hippo signaling pathway
Autophagy - animal
Calcium signaling pathway
Toll-like receptor signaling pathway
Long-term potentiation
Glutamatergic synapse
NF-kappa B signaling pathway
Serotonergic synapse
p53 signaling pathway
JAK-STAT signaling pathway
Long-term depression
GABAergic synapse
IL-17 signaling pathway
Fc gamma R-mediated phagocytosis
cAMP signaling pathway

Gene Ratio

pvalue
0.015
0.010
0.005

Gene Count

C

Rescued transcripts in Penumbral Neurons

Row Z-Score
-1 0 1

SHAM   MCAo   MCAo+HDACi

TGF-Beta Signaling

PI3K-AKT Signaling

MAPK Signaling

Notch Signaling

Apoptosis

Figure 1 Supplementary Figure 1

TTC (2,3,5-Triphenyltetrazolium Chloride)

Figure 3 Supplementary Figure 1

HOECHST33342   H3K27me3   IBA1

Figure 4 Supplementary Figure 1

A

B

C

Figure 4 Supplementary Figure 2

GEX1-FF slide tissues transcriptome permeabilized

GEX1-FF slide tissues IF Overlay for spatial transcriptome browser

Microglia through SpaceRanger overlay -> Microglia+Neuron or Microglia+Astrocyte transcript analysis

Pre-cDNA **Amplification & Cleanup** [QC]

Final **Visium Libraries** [QC]

▪Neurons (NeuN)   ▪Astrocytes (GFAP)   ▪Nuclei (DAPI)

Figure 5 Supplementary Figure 1

IL-17 Signaling | Autophagy | Hippo Signaling | Neurotrophin Signaling pathways

Figure 6 Supplementary Figure 1A

A   Pathways enriched in HDACi-rescued Hippocampal microglial transcriptome

B   Pathways enriched in HDACi-rescued Thalamic microglial transcriptome

C   Pathways enriched in HDACi-rescued Striatal microglial transcriptome

D   Pathways enriched in HDACi-rescued Cortical microglial transcriptome

Figure 6 Supplementary Figure 1B

A

Gene Ontology of HDACi-rescued
Hippocampal microglial transcriptome

B

Gene Ontology of HDACi-rescued
Thalamic microglial transcriptome

C

Gene Ontology of HDACi-rescued
Striatal microglial transcriptome

D

Gene Ontology of HDACi-rescued
Cortical microglial transcriptome

Figure 6 Supplementary Figure 2

Figure 6 Supplementary Figure 3

A CNETPLOT: HDACi-rescued hippocampal microglia KEGG pathway-transcript hits

B CNETPLOT: HDACi-rescued thalamic microglia KEGG pathway-transcript hits

C CNETPLOT: HDACi-rescued striatal microglia KEGG pathway-transcript hits

D CNETPLOT: HDACi-rescued cortical microglia KEGG pathway-transcript hits