Potent LILRB1 D1D2-containing antibodies inhibit RIFIN-mediated immune evasions

Yizhuo Wang[1,2, #], Hengfang Tang[1, #], Wanxue Wang[2,3], Ming Li[2], Chenchen Zhu[1,2], Han Dai[1], Hongxin Zhao[1,4,*], Bo Wu[2,*], Junfeng Wang[1,2,*]

[1]High Magnetic Field Laboratory, Key Laboratory of High Magnetic Field and Ion Beam Physical Biology, Hefei Institutes of Physical Science, Chinese Academy of Sciences, Hefei, Anhui 230031, China.

[2]University of Science and Technology of China, Hefei, Anhui 230026, China.

[3]Department of Anatomy, School of Basic Medical Sciences, Anhui Medical University, Hefei, China.

[4]Hefei China Science Longwood Biological Technology Co., Ltd. Hefei, Anhui 230088, China.

[#] Those authors contributed equally.

*Corresponding authors. Email: zhx@hmfl.ac.cn (Hongxin Zhao), wubo@hmfl.ac.cn (Bo Wu), junfeng@hmfl.ac.cn (Junfeng Wang).

Abstract

   Variant surface antigens of *Plasmodium falciparum*, including RIFIN, play a key role in malaria pathogenesis and promote immune evasion by binding to immunoinhibitory receptors such as LILRB1. Recently, receptor-containing antibodies have been identified in malaria-exposed individuals and reveal a novel antibody mechanism for inhibiting immune evasions in *Plasmodium falciparum*. Previous studies have identified several LAIR1- and LILRB1 D3D4-containing antibodies. However, no antibodies containing LILRB1-D1D2 have been identified, although some RIFINs interact with LILRB1-D1D2 to mimic the binding mode of β2-microglobulin, a major histocompatibility complex (MHC) class I molecules. In this study, we propose an in vitro strategy for generating receptor-containing antibodies through structure-based affinity maturation. Using this strategy, we successfully generated a LILRB1 D1D2-containing antibody, D1D2.v-IgG, which effectively blocked the specific binding of RIFIN#1 (from PF3D7_1254800) to LILRB1-D1D2 domains. Furthermore, we developed NK-biAb, a D1D2.v-IgG-based bispecific antibody targeting RIFIN#1 and NKG2D receptors. Both antibodies showed promising results in enhancing NK cell-mediated cytotoxicity against RIFIN#1-expressing K562 cells, with NK-biAb being more efficacious. The present strategy can be generalized for the development of antibodies against Plasmodium-host interactions, thus contributing to the advancement of malaria treatments and vaccines.

Keywords: *Plasmodium falciparum*, immune evasion, RIFIN, receptor-containing antibody

Introduction

Malaria is one of the most deadly infectious diseases affecting human health, causing nearly half a million deaths each year. The disease is caused by *Plasmodium* infections, of which *Plasmodium falciparum* (*P. falciparum*) is the deadliest species[1,2]. The emergence and spread of drug-resistant parasites threaten global malaria control and raise the demand for novel malaria treatments and vaccines[3-5].

The life cycle of *P. falciparum* is complex, comprising multiple developmental stages from the mosquito vector to the human host. Of these, the blood stage is responsible for the clinical symptoms of malaria[6,7]. During this stage, *P. falciparum* invades and replicates in erythrocytes and also expresses a number of variable surface antigens on infected erythrocytes (IEs). Blood-stage parasites are the main target of acquired immunity. However, *P. falciparum* employs multiple strategies to evade immune responses by interacting with host receptors through variable surface antigens, which is thought to contribute to the slow development of naturally acquired immunity against malaria. PfEMP1 and RIFIN are two of the three major multigene familiy proteins expressed on the surface of IEs. PfEMP1 proteins were found to interact with various human endothelial receptors and mediate the adhesion of the IEs to blood vessels and tissues surfaces, thereby helping *P. falciparum* to evade splenic clearance. RIFINs are the largest family of variable surface antigens of *P. falciparum*[7,8]. Some RIFINs have been shown to mediate immunosuppression by binding to and activating inhibitory immune receptors on myeloid and lymphoid cells , such as leukocyte-associated immunoglobulin-like receptor 1 (LAIR1) and leukocyte immunoglobulin-like receptor B1 (LILRB1)[9-12] (Fig. 1a). Theoretically, blocking those surface antigen-receptor interactions, thereby preventing the immune escape of *Plasmodium*, maybe a promising strategy for development of effective malaria treatments and vaccines[13,14]. Notably, LAIR1- and LILRB1-containing antibodies have recently been found in

malaria-infected individuals, suggesting that the human immune system employs this

Figure 1. Immune evasion of *P.falciparum* and design of novel receptor-containing antibody

(**a**) Mechanisms of immune evasion of *P. falciparum* and structures of RIFIN and immunoinhibitory

receptors**.** Different RIFIN isoforms bind to their corresponding receptor domains and transduce

immunosuppressive signals (PDB: 6ZDX, 7KFK and 7F9N). (**b**) Receptor-containing antibodies in

inhibiting immune evasion of *P. falciparum* and identification of LILRB1- and LAIR1-containing

antibodies. All reported LILRB1-containing antibodies only have LILRB1-D3D4 domains and no

antibodies that contain LILRB1-D1D2 domains were found. (**c**) Two representative receptor-

containing antibody. The left one is LILRB1 D3D4-containing antibody MDB1 (PDB:7KHF) and

the right one is LAIR1-containing antibody MGD21 (PDB: 7JZ4). (**d**) Strategy for generating high-

affinity LILRB1 D1D2-containing antibodies. The affinity-matured LILRB1-D1D2 variant was

generated through structure-based affinity maturation in vitro and then the LILRB1-D1D2 variant

was grafted into the framework of MDB1 by replacing LILRB1-D3D4 domains.

strategy to counteract immune evasions of malaria parasites[15-17] (Fig. 1b). The LAIR1-

containing antibodies reveal specific mutations in the LAIR1 insert, resulting in high

affinity toward RIFINs[16]. Unlike the LAIR1-containing antibodies, the LILRB1-

containing antibodies reveal no mutation in LILRB1 inserts and all LILRB1 inserts are

LILRB1-D3D4 domains or just LILRB1-D3 domain within the variable-constant (VH-

CH1) elbow region[15] (Fig. 1c).   Some RIFINs bind to LILRB1-D1D2, mimicking the

binding pattern between LILRB1-D1D2 and MHC class I. However, no antibodies

containing the LILRB1-D1D2 structural domain have been identified, probably due to

the low affinity of RIFINs for LILRB1-D1D2. In addition, the formation of receptor-

containing antibodies in vivo is time-consuming, especially during the in vivo affinity

maturation process, and the maturation site of receptor-containing antibodies is

uncontrollable.

Here, we propose a strategy for designing novel receptor-containing antibodies.

Our strategy is to obtain a high-affinity receptor fragment variant by structure-based

affinity maturation in vitro [18-20] , and then graft the affinity-matured receptor fragment

variants into the framework of the human anti-malaria natural antibody MDB1 by

replacing the LILRB1-D3D4 structural domain in MDB1 (Figure 1d). Using this

strategy, a novel LILRB1 D1D2-containing antibody, D1D2.v-IgG, was successfully

generated. D1D2.v-IgG binds to RIFIN with 110-fold higher affinity than LILRB.

Biochemical and cellular assays demonstrated that D1D2.v-IgG blocked the RIFIN#1-

LILRB1 interaction, and significantly enganced the NK cell-mediated cytotoxicity

against RIFIN#1-expressing K562 cells. In addition, based on the D1D2.v-IgG, a bi-

specific antibody NK-biAb was engineered by incorporating the natural killer group 2

D (NKG2D) receptor-specific scFv KYK2.0[21-23] at its C-terminus, mimicking CAR-

NK therapy[24,25]. Notably, NK-biAb exhibits an enhanced capacity to augment NK cell-

mediated cytotoxicity compared to D1D2.v-IgG. Our work provides a novel approach

to the design of antibodies targeting the interaction between surface antigens of

Plasmodium variants and host receptors, which may contribute to the development of

malaria therapeutics and vaccines.

Results

Structure-guided engineering of a high affinity LILRB1 D1D2 variant

RIFIN#1 binds to the extracellular domain (ECD) of LILRB1 through its variable

region with a moderate affinity (K_d =570 ± 130 nM). LILRB1-ECD comprises four

immunoglobulin-like domains (D1-D4), and its complex structure with RIFIN#1

suggests that the interaction is mediated by the D1D2 domain[9]. In order to obtain a

high-affinity variant capable of competitively antagonizing LILRB1-RIFIN#1

interaction, we performed structure-guided directed evolution of the LILRB1-D1D2

domain by yeast surface display. A library of LILRB1 D1D2 mutations was designed

using the structure of the LILRB1-RIFIN#1 complex as a guide. As shown in Figure

2a, ten LILRB1 D1D2 residues (K42, E68, G97, A98, Y99, I100, Q125, V126, F128,

E184) interfacing with RIFIN#1 were altered based on the structural information by

randomized PCR. The resulting D1D2 mutant library contained $2x10^7$ transformants.

The library was then displayed on the yeast surface, stained with biotinylated RIFIN ,

and screened to isolate high-affinity binders (Fig. 2b and Fig. S1). In total, eight round

of sorting were performed (Fig. S2). Three rounds of sorting with decreasing

concentrations of biotinylated RIFIN tetramers (2 μM, 1 μM and 0.3 μM) (Fig. S3) and

three rounds of sorting with decreasing concentrations of biotinylated RIFIN monomers

(0.3 μM, 0.1μM and 0.02 μM) (Fig. S4) were performed for binder enrichment. For

kinetic sorting, enriched library was screened using 0.02 μM biotinylated RIFIN

monomers in the presence of approximately a 50-fold excess unlabeled competition

RIFIN, with a duration of 2 h and 4 h respectively (Fig. S5). After completion of all

eight rounds of sorting, yeast cells were transplanted to obtain individual clones. DNA

sequencing results indicate that a single D1D2 mutant clone (named D1D2.v)

dominates this enrichment process, and this consensus variant contains a total of nine

mutations: K42M, G97S, A98T, Y99F, I100T, Q125E, V126I, F128Q, E184H (Fig. 2c

and Fig. S6).

To gain insight into the mechanisms of D1D2.v affinity enhancement, we obtained

a structural model of D1D2.v binding to RIFIN#1 using SWISS-MODEL (Fig. 2d).

Comparative analysis with the crystal structure of the LILRB1-RIFIN#1 complex show

that D1D2.v binds to RIFIN#1 in essentially the same manner as its wild-type

counterpart; however, some differences are observed at the binding interface. As shown

in Figure 2d, the Q125E mutation establishes a new salt bridge with R233 on RIFIN#1.

In addition, the G97S, A98T, and I100T mutations appears to enhance the strength of

their H-bonds with RIFIN#1 based on analysis of the H-bond distance.

Figure 2. Yeast display and affinity maturation of D1D2 domain

(**a**) (Left) LILRB1 models based on LILRB1-RIFIN#1 complex structure (PDB:6ZDX), where

RIFIN#1 is in orange color, LILRB1-D1D2 is in cyan color, and the randomized positions are

depicted by red balls; (Right) Table of random positions of the LILRB1-D1D2 interface library with

possible amino acid variants. (**b**) Flow cytometry results for the yeast stained with biotinylated

RIFIN#1 in tetramer, monomer and competition selection strategy. (**c**) Mutation frequencies (Above)

and consensus D1D2.v sequence (Below). (**d**) Structural modeling and interface analysis. The

sequences of D1D2.v and RIFIN#1 were imported into the SWISS-MODEL server for initial model

generation based on the coordinates of PDB code. The G97S, A98T, I100T and Q125E mutations

in D1D2.v increases H-bonding strength targeting RIFIN#1, while a salt-bridge forms between

E125 on D1D2.v and R233 on RIFIN#1.

Design and biochemical characterization of a D1D2.v-containing antibody that

antagonizes RIFIN#1-LILRB1 interaction.

Recently, several natural LILRB1-containing monoclonal antibodies were

identified in malaria-infected individuals, such as MDB1, in which the LILRB1 D3D4

structural domain was inserted into its variable-conatant(VH-CH1) elbow region[15,16].

We then replaced the LILRB1-D3D4 insertion domain in MDB1 with D1D2.v to

generate a D1D2-containing antibody, named D1D2.v-IgG (Fig. 3a). To express as the

secreted protein, the codon-optimized antibody sequence was genetically linked to a N-

terminal Trp2 signal peptide sequence within a pTT5 expression vector. The

recombinant antibody was transiently expressed in Expi293F cells and purified from

culture supernatant by protein A affinity chromatography. The purity was confirmed by

SDS-PAGE analysis under both non-reducing and reducing conditions. To evaluate the

RIFIN#1-binding activity of this recombinant antibody, we generated the Fab fragment

by enzymatically removing the Fc tag and detected its binding with RIFIN#1 by size-

exclusion chromatography (SEC) assay. The results showed that Fab bound strongly to

RIFIN#1, forming a tight complex (Figure 3b). Next, we quantified the binding affinity

of D1D2.v-IgG to RIFIN#1 using a protein-based ELISA assay with purified LILRB1

ECD as a control. As shown in Fig. 3c, the dissociation constant (K_d) for LILRB1 ECD

binding to RIFIN#1 was approximately 650 nM, which is consistent with the previously

reported affinity determined using surface plasmon resonance (SPR). In contrast, the

affinity of D1D2.v-IgG binding to RIFIN#1 was significantly higher, with a K_d of 6

nM, a 110-fold increase (Fig. 3c). To assess the ability of D1D2.v-IgG to antagonize

the binding of LILRB1 to RIFIN#1, we performed a competitive ELISA assay. The

results showed that D1D2.v-IgG effectively inhibited the binding of LILRB1 to

RIFIN#1 in a dose-dependent manner with a half-maximal inhibitory concentration

(IC_{50}) of 20 nM (Fig. 3d).

Fig.3: Molecular design and biochemical characterization of the D1D2.v-IgG

(a) Molecular structure and cDNA frame of LILRB1 D1D2-containing IgG. The inserted LILRB1-

D1 domain is colored in dark blue and LILRB1-D2 domain is colored in light blue, which are located

between heavy variable chain (VH) and heavy constant chain (CH). The antibody heavy chain and

Fc domain are depicted in dark grey and the light chain is depicted in light grey. **(b)** Purification

and functional validation of the antigen binding fragment (Fab) of D1D2.v-IgG. SDS-PAGE (Left)

show the purity and molecular weight of D1D2.v-IgG before protease digestion. SDS-PAGE (Right)

show the purity and molecular weight of D1D2.v-Fab. Size-exclusion chromatography (SEC)

chromatograms show the homogeneous Fab-RIFIN complex have a UV_{280} absorbance at around 13

mL. **(c)** Titration curves of wild-type LILRB1 (ECD) (dark green) and D1D2.v-IgG (red) binding

to RIFIN in ELISA assay. Data are mean ± SD. **(d)** The representative competition binding curve

using wild type LILRB1 (ECD) show D1D2.v-IgG antagonizes the RIFIN-LILRB1 (ECD)

interaction effectively. Data represent mean ± SD of three technical replicates.

D1D2.v-IgG significantly enhances NK92 cytolytic activity against RIFIN#1-

expressing K562 cells in vitro

Given that RIFIN#1 binding to LILRB1 inhibits NK cell responses, we

subsequently investigated whether D1D2.v-IgG could enhance NK cell-mediated

cytotoxicity by blocking the interaction between LILRB1 and RIFIN#1. To assess this,

we performed NK92 cell cytotoxicity assays on K562 cells stably transfected with full

length RIFIN#1 (Fig. S7). Flow cytometry verified the expression level of LILRB1 on

NK92 cells (Fig. 4a). The RIFIN#1-expressing K562 cells were co-cultured with NK92

cells at two different effector to target (E:T) ratios of 2.5 and 12.5 in the presence or

absence of D1D2.v-IgG. Flow cytometry analysis confirmed the specific binding of

purified D1D2.v-IgG to the RIFIN#1-expresssing K562 cells (Fig. 4b). The

cytotoxicity against K562 cells expressing RIFIN#1 was 5.8% in the absence of

D1D2.v-IgG when the E:T ratio was 2.5, and increased significantly to 13.8% when

D1D2.v-IgG at a concentration of 100 nM was added. Similarly, when the E:T ratio

was 12.5, the cell killing efficiency was 35% in the absence of D1D2.v-IgG, but

increased significantly to 61% when D1D2.v-IgG at a concentration of 100 nM was

added (Fig. 4c). In addition, we observed that D1D2.v-IgG significantly increased the

secretion of perforin and granzyme B from NK92 cells (Fig. 3d-e). These findings

clearly indicate that D1D2.v-IgG can effectively stimulate NK-cell cytotoxicity against

RIFIN#1-expressing K562 cells.

Fig.4: D1D2.v-IgG enhanced NK cell cytotoxicity against RIFIN-expressing K562 cell

(**a**) Detection of the LILRB1 on NK92 cells by flow cytometry. The red curve shows the binding of

LILRB1 antibody to LILRB1 on NK92 cells, while the blue curve informs the control sample of

NK92 cells. (**b**) Detection of the purified D1D2.v-IgG bound to RIFIN#1-expressing K562 cells by

flow cytometry. The red curve shows the binding of D1D2.v-IgG to full-length RIFIN#1 on stably-

transfected K562 cells, while the cyan curve informs the blank sample of K562 cells. (**c**) The results

for the NK92 cytotoxicity in the presence or absence of D1D2.v-IgG. RIFIN#1-expressing K562

cells were cocultured with NK92 cells and incubated with 100 nM control human IgG (hIgG) (Grey,

n = 3) or 100 nM D1D2.v-IgG (Yellow, n = 3). **(d,e)** Detections of the perforin (e) and granzyme B

(f) releases in the presence or absence of D1D2.v-IgG. RIFIN#1-expressing K562 cells were

cocultured with NK92 cells (E: T ratio = 1: 1) and incubated with 100 nM control hIgG (Grey, n =

3) or 100 nM D1D2.v-IgG (Orange, n = 3). After the incubation for 24h, the perforin and granzyme

B levels were determined by ELISA. Statistical test was t test and statistical significance was

analyzed between D1D2.v-IgG and hIgG. *P<0.05; **P<0.01; ***P<0.001; ****P<0.0001.

Generation of a bispecific antibody targeting RIFIN#1 and NKG2D

Next, we developed a bispecific antibody that recruits NK cells to RIFIN#1-

expressing cells using the D1D2.v-IgG as a basis. The redirection of NK cells requires

a highly specific and high-affinity 'compass' (Fig. 5a). To fulfill this requirement, we

chosen an engineered scFv called KYK2.0, which exhibits a nanomolar affinity towards

NKG2D expressed on all types of NK cells. By fusing the scFv form of KYK2.0 to the

C terminal of the Fc fragment of D1D2.v-IgG via a flexible GS linker, we generated a

bispecific antibody (bsAb), named NK-biAb (Fig. 5b). The resulting bsAb was then

transiently expressed and purified from Expi293F cell culture supernatants by affinity

chromatography. Integrity and purity of NK-biAb were analyzed by SDS-PAGE under

reducing and non-reducing conditions (Fig. S8a). As exprected, this purified bispecific

antibody demonstrated the ability to bind to both NK cells and RIFIN#1-expressing

K562 cells in flow cytometry analysis (Fig. S8b-c).

Fig.5: Design and characterization of NK-biAb

(a) Schematic of NK cell redirections by NK-biAb. Scfv form of KYK2.0 (scfv-KYK2.0), an

NKG2D activating antibody was fused to C terminus of D1D2.v-IgG. When recognizing and

blocking immune escape response of *P. falciparum* via D1D2.v-Fab domains, scfv-KYK2.0 domain

can also bind to NKG2D receptors, redirect NK cells to *P. falciparum* and trigger NK cell activities.

scFv-KYK2.0 is colored in light orange and NKG2D receptor is depicted as a light blue dart.

(b) cDNA frame of NK-biAb. cDNA of D1D2.v-IgG part is the same as previously described. The

heavy variable chain (VH) and light variable chain (VL) of KYK2.0 depicted in orange are fused to

C terminus of D1D2.v-IgG by a GS linker. **(c)** NK92 cytotoxicity assay in the presence of NK-biAb.

RIFIN#1-expressing K562 cells were cocultured with NK92 cells and incubated with 100 nM

control human IgG (hIgG) or 50 nM NK-biAb (Cyan, n =3) and 100 nM NK-biAb (Red, n =3).

Statistical tests were two-way ANOVA and statistical significance was analyzed between NK-biAb

and hIgG. **(d,e)** Detections of the perforin (d) and granzyme B (e) releases in the presence or absence

of NK-biAb. RIFIN#1-expressing K562 cells were cocultured with NK92 cells (E: T ratio = 1: 1)

and incubated with 100 nM control hIgG (Grey, n = 3), 50 nM NK-biAb (Cyan, n =3) and 100 nM

NK-biAb (Red, n = 3). After the incubation for 24h, the perforin and granzyme B levels were

determined by ELISA. Statistical tests were one-way ANOVA and statistical significance was

analyzed between NK-biAb and hIgG. **(f)** Flow detection results for the coculture of NK92 and

RIFIN#1-expressing K562 for 24 h at E: T = 1: 1 in the presence of 100 nM hIgG (Left), 50 nM

NK-biAb (Middle) and 100 nM NK-biAb (Right), showing the RIFIN#1-expressing K562 cells

killed by NK92 cells. The data in the second quadrant stands for the lysed RIFIN#1-expressing

K562 cells. *P<0.05; **P<0.01; ***P<0.001; ****P<0.0001.

NK-biAb exhibits an enhanced ability to trigger NK92 cytotoxicity against RIFIN#1-expressing K562 cells.

Finally, we evaluated the effect of NK-biAb on NK cell cytotoxicity against

RIFIN#1-expressing K562 cells. As shown in figure, the ability of this bispecific

antibody to trigger NK cytotoxicity was significantly enhanced compared to the

monospecific antibody D1D2.v-IgG. When treatment with NK-biAb at a concentration

of 100 nM, there was a remarkable enhancement in cytotoxic efficiency from 13.8% to

26.1% and from 61%-93.6%, at E/T ratios of 2.5 and 12.5 respectively, compared to

treatment with D1D2.v-IgG under the same conditions (Fig. 5c). Compared with

D1D2.v-IgG treatment, NK92 cells secreted more perforin and granzyme B in the

presence of 50 nM and 100 nM NK-biAb (Fig. 5d-5e). Flow cytometry results also

verified that NK-biAb can potently stimulate NK-cell cytotoxicity against RIFIN#1-

expressing K562 cells (Fig. 5f).

Discussion

Upon infection, *P. falciparum* utilizes variant surface antigens to engage with

receptors on the human host, thereby facilitating immune evasion and ensuring their

survival in the host[7,8]. Recent studies have identified several receptor-containing

antibodies in malaria-exposed individuals that block surface antigen-receptor

interactions. Functional studies have shown that these antibodies hold potential as

therapeutic agents and for incorporation into malaria vaccine design[15-17]. For instance,

MGD21 antibody has been shown to promote NK cell-dependent cleavage of IEs and

to inhibit the growth of *P. falciparum*[17]. However, reported receptor-containing

antibodies only have either LAIR1 domain or LILRB1-D3D4 domains, and no

antibodies containing LILRB1-D1D2 domains have been discovered. The RIFIN

family is divided into different species, some of which interact with LILRB1-D1D2

domains rather than LILRB1-D3D4 domain or LAIR1 to transduce immunoinhibitory

signals. Therefore, the use of LILRB1 D1D2-containing antibodies for the treatment of

malaria is of great importance.

In this study, we propose an in vitro strategy for designing novel receptor-

containing antibodies. Our approach utilizes structure-guided affinity maturation to

obtain receptor fragment variants with enhanced binding capacity, while using the

framework of a human natural antibody against malaria as a template for antibody

generation[15]. Due to the weak affinity between RIFIN and LILRB1-D1D2, no relevant

D1D2-containing antibody has been reported. We used this strategy to artificially create

high-affinity D1D2-containing antibodies, filling a gap in this field. As an illustrative

example, we successfully prepared D1D2.v-IgG antibody, which has 110-fold higher

affinity towards RIFIN1# than LILRB1 and can effectively block the interaction

between RIFIN1# and LILRB1. Cellular experiments demonstrated that D1D2.v-IgG

significantly enhanced NK cell-mediated cytotoxicity against K562 cells expressing

RIFIN#1. In addition, on the basis of D1D2.v-IgG, we developed a bispecific antibody

targeting both RIFIN#1 and the NKG2D receptor, which showed enhanced ability to

trigger NK cell-mediated cytotoxicity compared to D1D2.v-IgG. Bispecific antibodies

have been shown to be a promising avenue for immunotherapeutic interventions in a

variety of malignancies[21,22,24,26]. Our data emphasize the potential application of these

agents in combating malaria.

In conclusion, our study provides a strategy for the design of receptor-containing

antibodies, offering potential applications in the development of malaria therapeutics

and vaccines. The increasing availability of resolved structures of Plasmodium surface

antigens binding to host receptors[9,11,15,17] will facilitates the development of

corresponding antibodies using our strategy.

Materials and Methods

Antigen purifications

The variable region of RIFIN#1 (PF3D7_1254800, 165-274 aa) with an N-

terminal 6×His tag was cloned into a pET28a vector. The recombinant vector was

transformed into E.coli BL21 (gold), and the cells were induced with 1 mM IPTG at a

OD_{600} of 0.7. Protein was expressed as inclusion bodies and denatured in a guanidine

buffer (6 M Guanidine hydrochloride, 20 mM Tris pH 8.0, and 300 mM NaCl) for 4 h.

The inclusion bodies were then purified by Ni-NTA affinity chromatography and eluted

by the guanidine buffer containing additional 500 mM imidazole. The denatured

inclusion bodied were retrieved by dialyzing against a buffer (20 mM Tris pH 8.0, 300

mM NaCl) and resolubilized in the guanidine buffer. Refolding was performed by the

quick dilution of inclusion bodies into a buffer (1 M L-arginine, 20 mM Tris pH 8.5,

100 mM NaH$_2$PO$_4$, 2 mM ethylenediaminetetraacetic acid, 0.1 mM

phenylmethylsulfonyl fluoride, 3 mM reduced glutathione and 0.3 mM oxidized

glutathione) until a final protein concentration of 0.1 mg/mL-0.2 mg/mL. The refold

RIFIN#1 was further polished on a Superdex 75 Increase column in a PBS (pH 7.2)

buffer. The ECD of human LILRB1 gene with a C-terminus His tag was cloned into a

pTT5 vector, and the vector was then PEI-transfected into Expi293F cells (Procell,

China). After a post-transfection for 96 h, the supernatant was collected and the protein

was purified with a Ni-NTA affinity chromatography. The purified LILRB1-ECD was

further exchanged into PBS (pH 7.2) buffer on a Superdex 200 Increase column.

For biotinylation, the RIFIN#1 and LILRB1-ECD proteins were cultured with

NHS-pEG12-biotin in a PBS buffer (pH 7.2). The culture reactions were quenched with

a Tris buffer after 8 h incubation. The free NHS-pEG12-biotin reagents were splitted

from the biotin-labeled proteins by a size-exclusion chromatography.

EBY100 vector construction and LILRB1-D1D2 wild type display

To avoid weird display stuff, pYD1 vector was reconstructed by fusing Agar 1/2

domain to the C-terminus of displayed targets. Meanwhile, the detection tags of pYD1

vector were replaced by a N-terminal FLAG, a C-terminal V5-tag and a C-terminal His

tag. D1D2 wild type was co-electroporated with a precut reconstructed pYD1, while

yeast cells were induced with SGCAA. The signals of RIFIN#1 binding to the displayed

D1D2 wild type were detected by FCM.

Constructing the library of D1D2 variants

Based on the crystal structure of RIFIN#1-LILRB1 complex, a site-directed

library was generated by the several rounds of overlapping PCR reactions for

randomizing the binding residues of LILRB1-D1D2. The randomizing primers are as

shown in Supplementary Table 1.The overlapping PCR reactions were conducted under

20 cycles. The external primers for the amplification of LILRB1-D1D2 library on a

large scale introduce a 70 bp homologous sequence to pYD1 vector. The LILRB1-

D1D2 library were co-transformed into EBY100 with a precut pYD1 vector at a

library/vector ratio of 3: 1. Yeast transformants were recovered in a YPD/Sorbitol broth

and amplified in SDCAA, resulting in a library containing 2×10^7 transformants.

MACS selections of the naive library

Transformed yeast were induced in SGCAA medium at 20°C for 48 h. For the 1st

round selection, 4×10^8 cells referred to as 20× library size were pellet and washed with

a PBE buffer (PBS pH 7.2 supplemented with 0.5% BSA). Firstly, the cells were

resuspended in 2 mL of the PBE buffer containing 2 μM biotinylated RIFIN that had

been pre-coated with Streptavidin-Alexa Fluor 647 (Invitrogen, USA), followed by an

incubation at 4°C for 2 h. Secondly, the stained cells were pelleted by centrifugation

(3500 g, 5 min) , washed twice with 2 mL PBE, and resuspended in PBE containing

paramagnetic anti-Alexa Fluor 647 microbeads (Miltenyi Biotech, German). Thirdly, a

magnetically-labeled yeast cells were enriched on LS MACS column (Miltenyi Biotech,

German), and the enriched yeast cells were eluted with 5-10 mL PBE. Finally, the

selection outputs were pellet and amplified in SDCAA for next round selection. Five

additional selections were performed by successively decreasing concentrations of

biotinylated RIFIN#1 from 1 μM to 0.02 μM. In each round selection, the yeast cells

were first labeled with SAV-Alexa Fluor 647 after incubating with biotinylated

RIFIN#1 for 2 h and washed once with PBE, then coated with 100 μL of anti-Alexa

Fluor 647 microbeads for 30 min and washed once, and finally separated by LS MACS

column.

MACS competition selections

The MACS tetramer/monomer selection outputs were further performed

competition selections to get ultrahigh affinity variants. The induced tetramer/monomer

selection outputs were cultured with 0.02 µM biotinylated RIFIN#1 at 4°C for 2 h. Then

the cells were pellet, washed once with PBE and competed with 1 µM unlabeled

RIFIN#1 for 2 h. After the competition process, the cells were stained with

Streptavidin-Alexa Fluor 647 and coated with anti-Alexa Fluor 647 microbeads. MACS

sorting was also performed using LS column. By now, the 1st round competition was

finished, and the competition outputs were amplified and induced for next round

competition selection. In the 2nd round competition selection, the competing time of

unlabeled RIFIN was increased to 4 h, and the final selection outputs were titrated on

SDCAA plates for plasmid extraction and sequencing.

Preparation of D1D2.v-IgG

D1D2.v were inserted between MDB1-VH and MDB1-CH, and cloned into pTT5

vector as IgG4 forms. MDB1 light chain was also cloned to pTT5 vector. The DNAs

of MDB1 light chains and D1D2.v-IgG heavy chains were co-transfected into

Expi293F cells using PEI transfection reagents. After the transfection for 96 h, the

culture supernatant was separated by centrifugation (4000 rpm, 10 min) and purified

with protein A resin. The purified D1D2.v-IgG was further polished on a Superdex 200

Increase column in a PBS (pH 7.4) buffer.

Characterization of D1D2.v-IgG

Protein-based ELISA assays were performed to determine the affinity of D1D2.v-

IgG to RIFIN#1 in comparison to that of LILRB1-ECD. First, RIFIN#1 (5 µg/mL) was

immobilized on 96-well ELISA plates and incubated with D1D2.v-IgG of different

concentrations. Then, the plates were washed with PBST for 3 times and cultured with

the anti-human IgG-HRP at 1: 5000 dilution. Finally, the wells were washed with PBST,

developed with TMB for 15 min, and quenched with 2 M HCl. The absorbance at 450

nm was determined by a microplate reader.

For the protein-competition ELISA assay, RIFIN#1 (5 μg/mL) was immobilized

on 96-well ELISA plates, and cocultured with D1D2.v-IgG of different concentrations

and a competing biotinylated LILRB1 (ECD) of 1-1.5× K_d concentration, where K_d is

the dissociation constant. Then the plates were washed with PBST for 3 times and

cultured with HRP-conjugated streptavidin at 1: 2000 dilution. The absorbance at 450

nm was recorded by a microplate reader.

Purifying Fab fragment of D1D2.v-IgG and its functional validation

D1D2.v-IgG was digested by His-tagged HRV-3c protease overnight and

subjected to a negative protein A and Ni-NTA column purification. The 5-fold excess

of refold RIFIN#1 were co-cultured with the Fab fragment of D1D2.v-IgG and the

resulting complex was further polished on a Superdex 200 Increase column. The ratio

of RIFIN#1 binding to the Fab fragment of D1D2.v-IgG was assessed by SDS-PAGE.

Preparation of NK-biAb and its functional validation

scFv-KYK2.0 was fused to the C-terminus of D1D2.v-IgG heavy chain to generate

NK-biAb (M_r = 260 kDa). The DNAs of MDB1 light chains and NK-biAb heavy chains

were co-transfected into Expi293F cells using PEI transfection reagents. After the

transfection for 96 h, the culture supernatant was collected by centrifugation (4000 rpm,

10 min) and purified on a protein A column. The purified protein was further polished

on a Superdex 200 Increase column in a PBS (pH 7.4) buffer. To validate the functions

of NK-biAb to attach NK92 cells and bind NKG2D, the NK92 cells pre-stimulated with

IL-2 were cultured with NK-biAb at 37°C for 2 h, and stained with anti-human IgG-PE

for the detection of surface NKG2D.

Stable RIFIN#1-expressing K562 cell line production

K562 cells (Procell, China) were cultured at 37°C in DMEM with 10% FBS and

5% CO_2. The cDNA of RIFIN#1 with its natural signal peptide was cloned into pLVX-

ZsGreen plasmid. Lentivirus were produced by HEK-293T (Procell, China) for

infecting K562 cells. The infected K562 cells were enriched by the selection using 1

μg/mL puromycin, to obtain the stable RIFIN#1-expressing K562 cell line.

Evaluation of NK cell cytotoxicity

The RIFIN#1-expressing K562 cells were incubated at 37°C for 1 h with D1D2.v-

IgG or NK-biAb. The NK92 cells in the same volume as RIFIN#1-expressing K562

cells were then cocultured for 4 h at 37°C and 5% CO_2 with the incubated RIFIN#1-

expressing K562 cells at the E: T ratios of 2.5: 1 and 12.5: 1. The final mixtures contain

either 100 nM D1D2.v-IgG or 50 nM/100 nM NK-biAb. The killing efficiency (E) was

calculated from the absorbance (A) of CCK8 at 450 nm: $E\% = [1-(A_S-A_{EC})/A_{TC}]\times100\%$,

where A_S is for the mixture of effector and target cells, and A_{EC} and A_{TC} are, respectively,

for effector cells and target cells.

Cytokine measurement and FCM detection

10^5 NK92 cells were cocultured for 24 h with 10^5 RIFIN#1-expressing K562 cells

in a 96-well plate in the presence of either 100 nM D1D2.v-IgG or 50 nM/100 nM NK-

biAb. Perforin and granzyme B release were detected from the supernatants of the

cocultured mixture by protein-based ELISA assay kits (Beyotime, China). The mixtures

of NK92 and K562-RIFIN#1 cells were stained for 1 h at 4°C (controlled by ice) with

Annexin V-APC in 100 μL PBS containing 0.5% BSA. The stained cells were detected

by flow cytometry, with the data analyzed by Flowjo software.

Reference

1. World Malaria Report 2023 [M]. World Health Organization (2023).

2. Burton, D. R. Antibodies, viruses and vaccines. *Nature Reviews Immunology* **2**, 706-

713 (2002).

3. Haldar, K., Bhattacharjee, S. & Safeukui, I. Drug resistance in *Plasmodium*. *Nature*

Reviews Microbiology **16**, 156-170 (2018).

4. Burton, D. R. What Are the Most Powerful Immunogen Design Vaccine Strategies?

Reverse Vaccinology 2.0 Shows Great Promise. *Cold Spring Harb Perspect Biol*

9(2017).

5. Wells, T. & Donini, C. Monoclonal Antibodies for Malaria. *New England Journal of*

Medicine **387**, 462-465 (2022).

6. Nilsson, S. K., Childs, L. M., Buckee, C. & Marti, M. Targeting Human Transmission

Biology for Malaria Elimination. *PLOS Pathogens* **11**, e1004871(2015).

7. Sakoguchi, A. & Arase, H. Mechanisms for Host Immune Evasion Mediated by

Plasmodium falciparum-Infected Erythrocyte Surface Antigens. *Frontiers in*

Immunology **13** (2022).

8. Chan, J. A., Fowkes, F. J. & Beeson, J. G. Surface antigens of *Plasmodium*

falciparum-infected erythrocytes as immune targets and malaria vaccine candidates.

Cell Mol Life Sci **71**, 3633-3657(2014).

9. Harrison, T. E. *et al.* Structural basis for RIFIN-mediated activation of LILRB1 in

malaria. *Nature* **587**, 309-312 (2020).

10. Saito, F. *et al.* Immune evasion of *Plasmodium falciparum* by RIFIN via inhibitory

receptors. *Nature* **552**, 101-105 (2017).

11. Xie, Y. *et al.* Structural basis of malarial parasite RIFIN-mediated immune escape

against LAIR1. *Cell Rep* **36**, 109600 (2021).

12. Willcox, B. E., Thomas, L. M. & Bjorkman, P. J. Crystal structure of HLA-A2

bound to LIR-1, a host and viral major histocompatibility complex receptor. *Nature*

Immunology **4**, 913-919 (2003).

13. Wahl, I. & Wardemann, H. How to induce protective humoral immunity against

Plasmodium falciparum circumsporozoite protein. *Journal of Experimental Medicine*

219 (2022).

14. Cockburn, I. A. & Seder, R. A. Malaria prevention: from immunological concepts

to effective vaccines and protective antibodies. *Nature Immunology* **19**, 1199-1211,

doi:10.1038/s41590-018-0228-6 (2018).

15. Chen, Y. *et al.* Structural basis of malaria RIFIN binding by LILRB1-containing

antibodies. *Nature* **592**, 639-643 (2021).

16. Tan, J. *et al.* A LAIR1 insertion generates broadly reactive antibodies against

malaria variant antigens. *Nature* **529**, 105-109 (2016).

17. Xu, K. *et al.* Structural basis of LAIR1 targeting by polymorphic *Plasmodium*

RIFINs. *Nature Communications* **12**, 4226 (2021).

18. Chao, G. *et al.* Isolating and engineering human antibodies using yeast surface

display. *Nature Protocols* **1**, 755-768(2006).

19. Weiskopf, K. *et al.* Engineered SIRPα variants as immunotherapeutic adjuvants to

anticancer antibodies. *Science* **341**, 88-91(2013).

20. Gonzalez-Perez, D. *et al.* Affinity-matured DLL4 ligands as broad-spectrum

modulators of Notch signaling. *Nature Chemical Biology* **19**, 9-17(2023).

21. Kwong, K. Y., Baskar, S., Zhang, H., Mackall, C. L. & Rader, C. Generation,

affinity maturation, and characterization of a human anti-human NKG2D monoclonal

antibody with dual antagonistic and agonistic activity. *J Mol Biol* **384**, 1143-

1156(2008).

22. Zhang, C. *et al.* Bispecific antibody-mediated redirection of NKG2D-CAR natural

killer cells facilitates dual targeting and enhances antitumor activity. *J Immunother*

Cancer **9** (2021).

23. Zhao, C. *et al.* A new approach to produce IgG (4)-like bispecific antibodies. *Sci*

Rep **11**, 18630 (2021).

24. Wolf, N. K., Kissiov, D. U. & Raulet, D. H. Roles of natural killer cells in immunity

to cancer, and applications to immunotherapy. *Nature Reviews Immunology* **23**, 90-105

(2023).

25. Albinger, N., Hartmann, J. & Ullrich, E. Current status and perspective of CAR-T

and CAR-NK cell therapy trials in Germany. *Gene Therapy* **28**, 513-527 (2021).

26. Leaver-Fay, A. *et al.* Computationally Designed Bispecific Antibodies using

Negative State Repertoires. *Structure* **24**, 641-651 (2016).

Acknowledgements: Funding was provided by the National Key R&D Program of

China (Grant No. 2023YFA1607500), the National Natural Science Foundation of

China (Grant No. 32371362), and the Anhui Major Basic Research Projects (Grant No.

2023z04020016).

Author contributions: Y.W and H.T contributed equally to this work. Y.W initiated

the project, performed the library construction and selections, prepared antigen samples

and performed biochemical assays. H.T performed the FCM detections and conducted

the stable cell line production, cytotoxicity assays and ELISA assays. W.W performed

the antibody purifications and functional validations. L.M helped to prepare

biotinylated sample preparations. C.Z helped to perform biochemical assays and FCM

detections. Y.W. and B.W analyzed the data, interpreted the results. J.W, H.D and B.W

prepared the manuscript. J.W., B.W and H.Z supervised the project. All authors read

and edited the manuscript.

Competing interests: The authors declare no competing interests.

Data and materials availability: All data and materials are available within the paper

and its supplementary materials. Extra data are available from the corresponding author

(J. Wang) upon reasonable request.

List of Supplementary Materials

Fig S1 to S8 for multiple supplementary figures

Table S1 for supplementary table

570