1   **Potent LILRB1 D1D2-containing antibodies inhibit RIFIN-mediated**

2   **immune evasions**

Yizhuo Wang[1,2, #], Hengfang Tang[1, #], Wanxue Wang[2,3], Ming Li[2], Chenchen Zhu[1,2],

Han Dai[1], Hongxin Zhao[1,4,*], Bo Wu[2,*], Junfeng Wang[1,2,*]

[1]High Magnetic Field Laboratory, Key Laboratory of High Magnetic Field and Ion

Beam Physical Biology, Hefei Institutes of Physical Science, Chinese Academy of

Sciences, Hefei, Anhui 230031, China.

[2]University of Science and Technology of China, Hefei, Anhui 230026, China.

[3]Department of Anatomy, School of Basic Medical Sciences, Anhui Medical University,

Hefei, China.

[4]Hefei China Science Longwood Biological Technology Co., Ltd. Hefei, Anhui 230088,

China.

[#] Those authors contributed equally.

*Corresponding authors. Email: zhx@hmfl.ac.cn (Hongxin Zhao), wubo@hmfl.ac.cn

(Bo Wu), junfeng@hmfl.ac.cn (Junfeng Wang).

Abstract

Variant surface antigens of *Plasmodium falciparum*, including RIFIN, play

a pivotal role in malaria pathogenesis and facilitate immune evasion by binding

to immunoinhibitory receptors such as LILRB1. Recently, receptor-containing

antibodies have been discovered in malaria-exposed individuals and uncover a novel

antibody mechanism in inhibiting immune evasions of *Plasmodium falciparum*.

Previous studies have identified several LAIR1- and LILRB1 D3D4-containing

antibodies. However, no antibodies containing LILRB1-D1D2 have been identified,

even though some RIFINs interact with LILRB1-D1D2. In this study, we propose a in

vitro strategy for the generation of this type of antibodies by employing structure-based

affinity maturation. Using this strategy, we successfully generated D1D2.v-IgG, an

antibody that effectively blocks the specific binding of RIFIN#1 (from

PF3D7_1254800) to LILRB1. Furthermore, we developed NK-biAb, a bispecific

antibody targeting RIFIN#1 and the NKG2D receptor based on D1D2.v-IgG. Both

antibodies demonstrate promising results in augmenting NK cell-mediated cytotoxicity

against RIFIN#1-expressing K562 cells, with NK-biAb exhibiting superior efficacy.

The present strategy could be generally used for developing antibodies against

the malarial parasite-host interactions, thereby facilitating advancements in malaria

treatments and vaccines.

Keywords: *Plasmodium falciparum*, immune evasion, RIFIN, receptor-containing

antibody

Introduction

Malaria is one of the most deadly infectious diseases that affect human health, leading to nearly half a million deaths annually. The disease is caused by infection with *Plasmodium parasites*, with *Plasmodium falciparum* (*P. falciparum*) being the most lethal species[1,2]. The emergence and dissemination of drug-resistant parasites threaten malaria control and elimination globally and raise the demand for new malaria treatments and vaccines[3-5].

The life cycle of *P. falciparum* is complex, consisting of multiple developmental stages from the mosquito vector to the human host. Among them, the blood stage is responsible for the clinical symptoms of malaria[6,7]. During this stage, *P. falciparum* invades erythrocytes and replicates within them, while also expressing a number of variable surface antigens on infected erythrocytes (IEs). Blood-stage parasites are the main target of acquired immunity. However, *P. falciparum* employs multiple strategies to evade immune responses by interacting with host receptors through variable surface antigens, which is believed to contribute to the slow development of naturally acquired immunity against malaria. PfEMP1 and RIFIN are two of the three major multigene familiy proteins expressed on the surface of IEs. PfEMP1 proteins have been found to interact with various human endothelial receptors to mediate the adhesion of the IEs to the surfaces of blood vessels and tissues, thus facilitating *P. falciparum* to escape splenic clearance. RIFINs comprise the largest family of variable surface antigens in *P. falciparum*[7,8]. Some RIFINs have been shown to mediate immune suppression by binding to and activating inhibitory immune receptors on myeloid and lymphoid cells , such as leukocyte-associated immunoglobulin-like receptor 1 (LAIR1) and leukocyte immunoglobulin-like receptor B1 (LILRB1)[9-12] (Fig. 1a). In theory, blocking those surface antigen-receptor interactions, thereby preventing the immune escape of malaria parasites maybe a promising strategy to develop effective malaria treatments and vaccines[13,14]. Notably, the recent identification of the LAIR1- and LILRB1-containing antibodies in malaria-exposed individuals suggests that the human immune system

Figure 1. Immune evasion of *P.falciparum* and design of novel receptor-containing antibody

(**a**) The mechanism of immune evasion of *P. falciparum* and structures of RIFIN and immunoinhibitory receptors. Different RIFIN subtypes bind to their corresponding receptor domains to transduce immunoinhibitory signals (PDB: 6ZDX, 7KFK and 7F9N). (**b**) Receptor-containing antibodies in inhibiting immune evasion of *P. falciparum* and identification of LILRB1- and LAIR1-containing antibodies. All reported LILRB1-containing antibodies only have LILRB1-D3D4 domains and no antibodies that contain LILRB1-D1D2 domains were found. (**c**) Two representative receptor-containing antibody. The left one is LILRB1 D3D4-containing antibody MDB1 (PDB:7KHF) and the right one is LAIR1-containing antibody MGD21 (PDB: 7JZ4). (**d**) Strategy for generating high-affinity LILRB1 D1D2-containing antibodies. Affinity-matured

LILRB1-D1D2 variant is generated through in vitro structure-based affinity maturation and then the

LILRB1-D1D2 variant is grafted into the framework of MDB1 by replacing LILRB1-D3D4

domains.

employs this strategy to combat immune evasions of malaria parasites[15-17] (Fig. 1b).

LAIR1-containing antibodies reveal specific mutations inside LAIR1 inserts, resulting

in high affinity toward RIFINs[16]. Different from LAIR1-containing antibodies,

LILRB1-containing antibodies reveal no mutation in LILRB1 inserts and all LILRB1

inserts are LILRB1-D3D4 domains or just LILRB1-D3 domain within the variable-

constant (VH-CH1) elbow region[15] (Fig. 1c). Antibodies that contain LILRB1-D1D2

domain have not been identified, which might be due to the low affinity between RIFIN

and LILRB1-D1D2. Moreover the in vivo formation of receptor-containing antibody is

time-consuming; in particular, the maturation sites of the receptor-containing antibody

during in vivo affinity-maturation are uncontrollable.

Here, we propose a strategy for the design of novel receptor-containing antibodies.

Our strategy features obtaining a high-affinity receptor fragment variant through in

vitro structure-based affinity maturation[18-20] and then the affinity-matured receptor

fragment variant is grafted into the framework of a human natural antibody MDB1[15]

against malaria by replacing LILRB1-D3D4 domains in MDB1 (Figure 1d). Using this

strategy, a novel LILRB1 D1D2-containing antibody, D1D2.v-IgG, was successfully

generated. The binding affinity of D1D2.v-IgG to RIFIN is about 110-fold higher than

that of LILRB1. Biochemical and cellular assays have demonstrated that this antibody

can effectively block the RIFIN#1-LILRB1 interaction, and significantly augment the

NK cell-mediated cytotoxicity against RIFIN#1-expressing K562 cells. Furthermore,

based on the D1D2.v-IgG, a bi-specific antibody NK-biAb was engineered by

incorporating the natural killer group 2 D (NKG2D) receptor-specific scFv KYK2.0[21-

23] at its C-terminus, mimicking CAR-NK therapy[24,25]. It is worth noting that this bi-

specific antibody exhibits an enhanced capacity to augment NK cell-mediated

cytotoxicity compared to D1D2.v-IgG. Our work provides a novel antibody design

approach targeting the interaction between malaria parasite variant surface antigens and

host receptors, which could be useful for development malaria treatments and vaccines.

Results

Structure-guided engineering of a high affinity LILRB1 D1D2 variant

RIFIN#1, via its variable region binds to the extracellular domain (ECD) of

LILRB1 with a moderate affinity (K_d =570 ± 130 nM). LILRB1-ECD comprises four

immunoglobulin-like domains (D1-D4), and its complex structure with RIFIN#1

demonstrates that the interaction is mediated by the D1D2 domain[9]. In order to obtain

a high-affinity agent capable of competitively antagonizing LILRB1-RIFIN#1

interaction, we carried out a structure-guided directed evolution of the LILRB1-D1D2

domain by yeast surface display. The LILRB1-RIFIN#1 complex structure was used as

a guide for designing LILRB1 D1D2 mutation library. As shown in Fig. 2a, based on

the structural information, ten LILRB1 D1D2 residues in the interface with RIFIN#1

(K42, E68, G97, A98, Y99, I100, Q125, V126, F128, E184) were varied by random

PCR. The resulting D1D2 mutation library contained $2×10^7$ transformants. The library

was then displayed on the yeast surface and stained by biotinylated RIFIN and subjected

to selections in order to isolate high-affinity binders (Fig. 2b and Fig. S1). A total of

eight rounds of sorting were conducted (Fig. S2). Following three rounds of sorting

with decreasing concentrations of biotinylated RIFIN tetramers (2 μM, 1 μM and 0.3

μM) (Fig. S3) and three rounds of sorting with decreasing concentrations of biotinylated

RIFIN monomers (0.3 μM, 0.1μM and 0.02 μM) (Fig. S4), two rounds of kinetic

competition sorting were performed using 0.02 μM biotinylated RIFIN monomers in

the presence of approximately a 50-fold excess unlabeled competition RIFIN, with a

duration of 2 h and 4 h respectively (Fig. S5). After completing all eight rounds of

sorting, yeast cells were plated to obtain single clones. DNA sequencing shows that a

single D1D2 mutant clone (named D1D2.v) dominates this enrichment process, and

this consensus variant contains 9 total mutations: K42M, G97S, A98T, Y99F, I100T,

Q125E, V126I, F128Q, E184H (Fig. 2c and Fig. S6).

To gain insight into the mechanisms underlying affinity enhancement in D1D2.v,

we utilized SWISS-MODEL to obtain a structural model of D1D2.v binding to

RIFIN#1 (Fig. 2d). A comparative analysis with the crystal structure of LILRB1-

RIFIN#1 complex suggested that D1D2.v bound to RIFIN#1 in a manner essentially

identical with its wild-type counterpart; however, some distinctions were observed at

the binding interface. As shown in Fig. 2d, the Q125E mutation establishes a new salt

bridge with R233 on RIFIN#1. In addition, it appears that the G97S, A98T and I100T

mutations enhance their H-bonding strength towards RIFIN#1 based on analysis of H-

bonding distances.

Figure 2. Yeast display and affinity maturation of D1D2 domain

(**a**) (Left) LILRB1 models based on LILRB1-RIFIN#1 complex structure (PDB:6ZDX), where

RIFIN#1 is in orange color, LILRB1-D1D2 is in cyan color, and the randomized positions are

depicted by red balls; (Right) tabulated randomized positions of the LILRB1-D1D2 interface library

with possible amino acid variants. (**b**) Flow cytometry results for the yeast stained with biotinylated

RIFIN#1 in tetramer, monomer and competition selection methods. (**c**) The frequency of mutations

(Above) and consensus D1D2.v sequence (Below). (**d**) Structural modeling and interface analysis.

The sequences of D1D2.v and RIFIN#1 were imported into the SWISS-MODEL server for initial

model generation based on the coordinates of PDB code**.** G97S, A98T, I100T and Q125E mutations

in D1D2.v increases H-bonding strength targeting RIFIN#1, while a salt-bridge forms between

E125 on D1D2.v and R233 on RIFIN#1.

Design and biochemical characterization of a D1D2.v-containing antibody that

antagonizes RIFIN#1-LILRB1 interaction.

Recently, several natural LILRB1-containing monoclonal antibodies have been

discovered in individuals exposed to malaria, such as MDB1, wherein the LILRB1

D3D4 domain is inserted within its variable-constant (VH-CH1) elbow region[15,16]. We

then generated a D1D2.v-containing antibody, named D1D2.v-IgG, by substituting the

LILRB1-D3D4 insertion in MDB1 with D1D2.v (Fig. 3a). For expression as a secreted

protein, the codon-optimized antibody sequence was genetically linked to a N-terminal

Trp2 signal peptide sequence within a pTT5 expression vector. The recombinant

antibody was transiently expressed in Expi293F cells and purified from culture

supernatant by protein A affinity chromatography. The purity was confirmed by SDS-

PAGE analysis under both non-reducing and reducing conditions. To assess the

RIFIN#1-binding activity of this recombinant antibody, we initially generated the Fab

fragment by enzymatically removing the Fc tag and test its binding with RIFIN#1 by

size-exclusion chromatography (SEC) assay. The results indicate a robust binding

between the Fab and RIFIN#1, forming a tight complex (Fig. 3b). Next, we quantified

the binding affinity of D1D2.v-IgG to RIFIN#1 using a protein-based ELISA assay,

employing the purified LILRB1 ECD as a control. As shown in Fig. 3c, the dissociation

constant (K_d) for the binding of LILRB1 ECD to RIFIN#1 is approximately 650 nM,

which is consistent with the previously reported affinity determined using surface

plasmon resonance (SPR). In stark contrast, D1D2.v-IgG exhibits a significantly higher

affinity towards RIFIN#1, with a K_d of 6 nM representing a 110-fold improvement (Fig.

3c). To assess the ability of D1D2.v-IgG to antagonize LILRB1 binding to RIFIN#1,

we conducted a competitive ELISA assay. The results revealed that this antibody

effectively inhibited LILRB1 binding to RIFIN#1 in a dose-dependent manner with a

half-maximal inhibitory concentration (IC50) of 20 nM (Fig. 3d).

Fig.3: Molecular design and biochemical characterization of the D1D2 (V)-IgG

(a) Molecular structure and cDNA frame of D1D2 domain containing IgG. Inserted LILRB1-D1

domain was colored in dark blue and LILRB1-D2 domain was colored in light blue, which were

located between heavy variable chain (VH) and heavy constant chain (CH). Antibody heavy chain

and Fc domain were depicted in dark grey and light chain was depicted in light grey. **(b)** Purification

and functional validation of the antigen binding fragment (Fab) of D1D2.v-IgG. SDS-PAGE (Left)

show the purity and molecular weight of D1D2.v-IgG before protease digestion. SDS-PAGE (Right)

show the purity and molecular weight of D1D2.v-Fab. Size-exclusion chromatography (SEC)

chromatograms show the homogeneous Fab-RIFIN complex have a UV_{280} absorbance at around 13

mL. **(c)** Titration curves of wild-type LILRB1 (ECD) (dark green) and D1D2.v-IgG (red) binding

to RIFIN in ELISA assay. Data are mean ± SD. **(d)** The representative competition binding curve

using wild type LILRB1 (ECD) showed D1D2.v-IgG antagonized the RIFIN-LILRB1 (ECD)

interaction effectively. Data represent mean ± SD of three technical replicates.

D1D2.v-IgG significantly enhances NK92 cytolytic activity against RIFIN#1-expressing K562 cells in vitro

Given that the binding of RIFIN#1 to LILRB1 leads to the suppression of NK cell

response, we subsequently investigated whether D1D2.v-IgG could enhance NK cell-

mediated cytotoxicity by blocking the interaction between LILRB1 and RIFIN#1. To

assess this, we conducted NK92 cell cytotoxicity assays on K562 cells stably

transfected with full length RIFIN#1 (Fig. S7). The expression level of LILRB1on

NK92 cells was validated by flow cytometry (Fig. 4a). The RIFIN#1-expressing K562

cells were co-cultured with NK92 cells at two different effector to target (E:T) ratios of

2.5 and 12.5 in the presence or absence of D1D2.v-IgG. Flow cytometry analysis

confirmed the specific binding of purified D1D2.v-IgG to the RIFIN#1-expresssing

K562 cells (Fig. 4b). At an E:T ratio of 2.5, the observed cytotoxicity against RIFIN#1-

expresssing K562 cells was 5.8% in the absence of D1D2.v-IgG, which significantly

increased to 13.8% upon addition of D1D2.v-IgG at concentrations of 100 nM.

Similarly, at an E:T ratio of 12.5, the cell killing efficiency was 35% in the absence of

D1D2.v-IgG, but remarkably rose to 61% when treated with D1D2.v-IgG at

concentrations of 100 nM (Fig. 4c). Moreover, we observed that D1D2.v-IgG

significantly augmented the secretion of perforin and granzyme B from NK92 cells (Fig.

3d-e). These findings clearly demonstrate that D1D2.v-IgG can effectively stimulate

NK-cell cytotoxicity against RIFIN#1-expressing K562 cells.

Fig.4: D1D2.v-IgG enhanced NK cell cytotoxicity against RIFIN-expressing K562 cell

(**a**) Detection of the LILRB1 on NK92 cells by flow cytometry. The red curve shows the binding of

LILRB1 antibody to LILRB1 on NK92 cells, while the blue curve informs the control sample of

NK92 cells. (**b**) Detection of the purified D1D2.v-IgG bound to RIFIN#1-expressing K562 cells by

flow cytometry. The red curve shows the binding of D1D2.v-IgG to full-length RIFIN#1 on stably-

transfected K562 cells, while the cyan curve informs the blank sample of K562 cells. (**c**) The results

for the NK92 cytotoxicity in the presence or absence of D1D2.v-IgG. RIFIN#1-expressing K562

cells were cocultured with NK92 cells and incubated with 100 nM control human IgG (hIgG) (Grey,

n = 3) or 100 nM D1D2.v-IgG (Yellow, n = 3). (**d,e**) Detections of the perforin (e) and granzyme B

(f) releases in the presence or absence of D1D2.v-IgG. RIFIN#1-expressing K562 cells were

cocultured with NK92 cells (E: T ratio = 1: 1) and incubated with 100 nM control hIgG (Grey, n =

3) or 100 nM D1D2.v-IgG (Orange, n = 3). After the incubation for 24h, the perforin and granzyme

B levels were determined by ELISA. Statistical test was t test and statistical significance was

analyzed between D1D2.v-IgG and hIgG. *P<0.05; **P<0.01; ***P<0.001; ****P<0.0001.

Generation of a bispecific antibody targeting RIFIN#1 and NKG2D

Next, we aim to develop a bispecific antibody capable of recruiting NK cells to

RIFIN#1-expressing cells using the D1D2.v-IgG as a basis. The redirection of NK cells

necessitates a highly specific and high-affinity 'compass' (Fig. 5a). To fulfill this

requirement, we have chosen an engineered scFv known as KYK2.0, which exhibits a

nanomolar affinity towards NKG2D expressed on all types of NK cells. By fusing the

scFv form of KYK2.0 to the C terminal of the Fc fragment of D1D2.v-IgG through a

flexible GS linker, we have generated a bispecific antibody (bsAb), named NK-biAb

(Fig. 5b). The resulting bsAb was then transiently expressed and purified from

Expi293F cell culture supernatants via affinity chromatography. Integrity and purity of

NK-biAb were analyzed by SDS-PAGE under reducing and non-reducing conditions

(Fig. S8a). As anticipated, this purified bispecific antibody demonstrated the ability to

bind to both NK cells and RIFIN#1-expressing K562 cells in flow cytometry analysis

(Fig. S8b-c).

Fig.5: Design and characterization of NK-biAb

(a) Schematic of NK cell redirections by NK-biAb. Scfv form of KYK2.0 (scfv-KYK2.0), an

NKG2D activating antibody was fused to C terminus of D1D2.v-IgG. When recognizing and

blocking immune escape response of *P. falciparum* via D1D2.v-Fab domains, scfv-KYK2.0 domain

can also bind to NKG2D receptors, redirect NK cells to *P. falciparum* and trigger NK cell activities.

scFv-KYK2.0 was colored in light orange and NKG2D receptor was depicted as a light blue dart.

(b) cDNA frame of NK-biAb. cDNA of D1D2.v-IgG part is the same as previously described. The

heavy variable chain (VH) and light variable chain (VL) of KYK2.0 depicted in orange are fused to

C terminus of D1D2.v-IgG by a GS linker. **(c)** NK92 cytotoxicity assay in the presence of NK-biAb.

RIFIN#1-expressing K562 cells were cocultured with NK92 cells and incubated with 100 nM

control human IgG (hIgG) or 50 nM NK-biAb (Cyan, n =3) and 100 nM NK-biAb (Red, n =3).

Statistical tests were two-way ANOVA and statistical significance was analyzed between NK-biAb

and hIgG. **(d,e)** Detections of the perforin (d) and granzyme B (e) releases in the presence or absence

of NK-biAb. RIFIN#1-expressing K562 cells were cocultured with NK92 cells (E: T ratio = 1: 1)

and incubated with 100 nM control hIgG (Grey, n = 3), 50 nM NK-biAb (Cyan, n =3) and 100 nM

NK-biAb (Red, n = 3). After the incubation for 24h, the perforin and granzyme B levels were

determined by ELISA. Statistical tests were one-way ANOVA and statistical significance was

analyzed between NK-biAb and hIgG. **(f)** Flow detection results for the coculture of NK92 and

RIFIN#1-expressing K562 for 24 h at E: T = 1: 1 in the presence of 100 nM hIgG (Left), 50 nM

NK-biAb (Middle) and 100 nM NK-biAb (Right), showing the RIFIN#1-expressing K562 cells

killed by NK92 cells. The data in the second quadrant stands for the lysed RIFIN#1-expressing

K562 cells. *P<0.05; **P<0.01; ***P<0.001; ****P<0.0001.

NK-biAb exhibits an enhanced ability to trigger NK92 cytotoxicity against

RIFIN#1-expressing K562 cells.

Finally, we evaluated the impact of NK-biAb on NK cell cytotoxicity against

RIFIN#1-expressing K562 cells. As shown in figure, this bispecific antibody exhibited

a significantly enhanced ability to trigger NK cytotoxicity compared to the

monospecific antibody D1D2.v-IgG. When treatment with a concentration of 100 nM

NK-biAb, there was a remarkable enhancement in cytotoxic efficiency from 13.8% to

26.1% and from 61%-93.6%, again at E/T ratios of 2.5 and 12.5 respectively, compared

to treatment with D1D2.v-IgG under identical conditions (Fig. 5c). In the presence of

50 nM and 100 nM NK-biAb, NK92 cells secreted more perforin and granzyme B

compared to treatment with D1D2.v-IgG (Fig. 5d-5e). Flow cytometry results also

validated that NK-biAb can potently stimulate NK-cell cytotoxicity against RIFIN#1-

expressing K562 cells (Fig. 5f).

Discussion

Upon infection, *P. falciparum* employs variant surface antigens to engage with

receptors on the human host, thereby facilitating immune evasion and ensuring their

survival within the host organism[7,8]. Recent investigations have identified several

receptor-containing antibodies in individuals exposed to malaria that possess the

capability to impede surface antigen-receptor interactions. Functional studies have

demonstrated that these antibodies hold potential as therapeutic agents and for

incorporation into malaria vaccine design[15-17]. For instance, MGD21 antibody has been

shown to promote NK cell-dependent lysis of IEs and inhibit *P. falciparum* growth[17].

In this study, we propose a in vitro strategy for the design of novel receptor-

containing antibodies. Our approach utilizes structure-guided affinity maturation to

obtain a receptor fragment variant with enhanced binding capability, while leveraging

the framework of a human natural antibody against malaria as a template for generating

an antibody[15]. Due to the weak affinity between RIFIN and LILRB1-D1D2, no related

D1D2-containing antibody has been reported. We used this strategy to artificially create

high-affinity D1D2-containing antibodies that fill a gap in this field. As an illustrative

example, we have successfully generated D1D2.v-IgG, an antibody that exhibits a 110-

fold higher affinity towards RIFIN1# than LILRB1 and is capable of effectively

blocking the interaction between RIFIN1# and LILRB1. Cellular assays demonstrated

that this antibody significantly enhances NK cell-mediated cytotoxicity against K562

cells expressing RIFIN#1. Furthermore, based on D1D2.v-IgG, we have developed a

bispecific antibody that targets both RIFIN#1 and the NKG2D receptor, which

demonstrates superior ability to enhance NK cell-mediated cytotoxicity compared to

D1D2.v-IgG. Bispecific antibodies have been shown to be a promising avenue for

immunotherapeutic interventions against diverse malignancies[21,22,24,26]. Our data

underscore the potential application of these agents in combating malaria.

In summary, our study provides a strategy for designing receptor-containing

antibodies, offering potential applications in the development of malaria therapeutics

and vaccines. The increasing availability of resolved structures of Plasmodium surface

antigens binding to host receptors[9,11,15,17] will facilitates the development of

corresponding antibodies using our strategy.

Materials and Methods

Antigen purifications

The variable region of RIFIN#1 (PF3D7_1254800, 165-274 aa) with an N-

terminal 6×His tag was cloned into a pET28a vector. The recombinant vector was

transformed into E.coli BL21 (gold), and the cells were induced with 1 mM IPTG at a

OD_{600} of 0.7. Protein was expressed as inclusion bodies and denatured in a guanidine

buffer (6 M Guanidine hydrochloride, 20 mM Tris pH 8.0, and 300 mM NaCl) for 4 h.

The inclusion bodies were then purified by Ni-NTA affinity chromatography and eluted

by the guanidine buffer containing additional 500 mM imidazole. The denatured

inclusion bodied were retrieved by dialyzing against a buffer (20 mM Tris pH 8.0, 300

mM NaCl) and resolubilized in the guanidine buffer. Refolding was performed by the

quick dilution of inclusion bodies into a buffer (1 M L-arginine, 20 mM Tris pH 8.5,

100 mM NaH_2PO_4, 2 mM ethylenediaminetetraacetic acid, 0.1 mM

phenylmethylsulfonyl fluoride, 3 mM reduced glutathione and 0.3 mM oxidized

glutathione) until a final protein concentration of 0.1 mg/mL-0.2 mg/mL. The refold

RIFIN#1 was further polished on a Superdex 75 Increase column in a PBS (pH 7.2)

buffer. The ECD of human LILRB1 gene with a C-terminus His tag was cloned into a

pTT5 vector, and the vector was then PEI-transfected into Expi293F cells (Procell,

China). After a post-transfection for 96 h, the supernatant was collected and the protein

was purified with a Ni-NTA affinity chromatography. The purified LILRB1-ECD was

further exchanged into PBS (pH 7.2) buffer on a Superdex 200 Increase column.

For biotinylation, the RIFIN#1 and LILRB1-ECD proteins were cultured with

NHS-pEG12-biotin in a PBS buffer (pH 7.2). The culture reactions were quenched with

a Tris buffer after 8 h incubation. The free NHS-pEG12-biotin reagents were splitted

from the biotin-labeled proteins by a size-exclusion chromatography.

EBY100 vector construction and LILRB1-D1D2 wild type display

To avoid weird display stuff, pYD1 vector was reconstructed by fusing Agar 1/2

domain to the C-terminus of displayed targets. Meanwhile, the detection tags of pYD1

vector were replaced by a N-terminal FLAG, a C-terminal V5-tag and a C-terminal His

tag. D1D2 wild type was co-electroporated with a precut reconstructed pYD1, while

yeast cells were induced with SGCAA. The signals of RIFIN#1 binding to the displayed

D1D2 wild type were detected by FCM.

Constructing the library of D1D2 variants

Based on the crystal structure of RIFIN#1-LILRB1 complex, a site-directed

library was generated by the several rounds of overlapping PCR reactions for

randomizing the binding residues of LILRB1-D1D2. The randomizing primers are as

shown in Supplementary Table 1.The overlapping PCR reactions were conducted under

20 cycles. The external primers for the amplification of LILRB1-D1D2 library on a

large scale introduce a 70 bp homologous sequence to pYD1 vector. The LILRB1-

D1D2 library were co-transformed into EBY100 with a precut pYD1 vector at a

library/vector ratio of 3: 1. Yeast transformants were recovered in a YPD/Sorbitol broth

and amplified in SDCAA, resulting in a library containing 2×10^7 transformants.

MACS selections of the naive library

Transformed yeast were induced in SGCAA medium at 20°C for 48 h. For the 1st

round selection, 4×10^8 cells referred to as 20× library size were pellet and washed with

a PBE buffer (PBS pH 7.2 supplemented with 0.5% BSA). Firstly, the cells were

resuspended in 2 mL of the PBE buffer containing 2 μM biotinylated RIFIN that had

been pre-coated with Streptavidin-Alexa Fluor 647 (Invitrogen, USA), followed by an

incubation at 4°C for 2 h. Secondly, the stained cells were pelleted by centrifugation

(3500 g, 5 min) , washed twice with 2 mL PBE, and resuspended in PBE containing

paramagnetic anti-Alexa Fluor 647 microbeads (Miltenyi Biotech, German). Thirdly, a

magnetically-labeled yeast cells were enriched on LS MACS column (Miltenyi Biotech,

German), and the enriched yeast cells were eluted with 5-10 mL PBE. Finally, the

selection outputs were pellet and amplified in SDCAA for next round selection. Five

additional selections were performed by successively decreasing concentrations of

biotinylated RIFIN#1 from 1 μM to 0.02 μM. In each round selection, the yeast cells

were first labeled with SAV-Alexa Fluor 647 after incubating with biotinylated

RIFIN#1 for 2 h and washed once with PBE, then coated with 100 μL of anti-Alexa

Fluor 647 microbeads for 30 min and washed once, and finally separated by LS MACS

column.

MACS competition selections

The MACS tetramer/monomer selection outputs were further performed

competition selections to get ultrahigh affinity variants. The induced tetramer/monomer

selection outputs were cultured with 0.02 μM biotinylated RIFIN#1 at 4°C for 2 h. Then

the cells were pellet, washed once with PBE and competed with 1 μM unlabeled

RIFIN#1 for 2 h. After the competition process, the cells were stained with

Streptavidin-Alexa Fluor 647 and coated with anti-Alexa Fluor 647 microbeads. MACS

sorting was also performed using LS column. By now, the 1st round competition was

finished, and the competition outputs were amplified and induced for next round

402   competition selection. In the 2nd round competition selection, the competing time of

403   unlabeled RIFIN was increased to 4 h, and the final selection outputs were titrated on

404   SDCAA plates for plasmid extraction and sequencing.

Preparation of D1D2.v-IgG

D1D2.v were inserted between MDB1-VH and MDB1-CH, and cloned into pTT5

vector as IgG4 forms. MDB1 light chain was also cloned to pTT5 vector. The DNAs

of MDB1 light chains and D1D2.v-IgG heavy chains were co-transfected into

Expi293F cells using PEI transfection reagents. After the transfection for 96 h, the

culture supernatant was separated by centrifugation (4000 rpm, 10 min) and purified

with protein A resin. The purified D1D2.v-IgG was further polished on a Superdex 200

Increase column in a PBS (pH 7.4) buffer.

Characterization of D1D2.v-IgG

Protein-based ELISA assays were performed to determine the affinity of D1D2.v-

IgG to RIFIN#1 in comparison to that of LILRB1-ECD. First, RIFIN#1 (5 µg/mL) was

immobilized on 96-well ELISA plates and incubated with D1D2.v-IgG of different

concentrations. Then, the plates were washed with PBST for 3 times and cultured with

the anti-human IgG-HRP at 1: 5000 dilution. Finally, the wells were washed with PBST,

developed with TMB for 15 min, and quenched with 2 M HCl. The absorbance at 450

nm was determined by a microplate reader.

For the protein-competition ELISA assay, RIFIN#1 (5 µg/mL) was immobilized

on 96-well ELISA plates, and cocultured with D1D2.v-IgG of different concentrations

and a competing biotinylated LILRB1 (ECD) of 1-1.5× K_d concentration, where K_d is

the dissociation constant. Then the plates were washed with PBST for 3 times and

cultured with HRP-conjugated streptavidin at 1: 2000 dilution. The absorbance at 450

nm was recorded by a microplate reader.

Purifying Fab fragment of D1D2.v-IgG and its functional validation

D1D2.v-IgG was digested by His-tagged HRV-3c protease overnight and subjected to a negative protein A and Ni-NTA column purification. The 5-fold excess of refold RIFIN#1 were co-cultured with the Fab fragment of D1D2.v-IgG and the resulting complex was further polished on a Superdex 200 Increase column. The ratio of RIFIN#1 binding to the Fab fragment of D1D2.v-IgG was assessed by SDS-PAGE.

Preparation of NK-biAb and its functional validation

scFv-KYK2.0 was fused to the C-terminus of D1D2.v-IgG heavy chain to generate NK-biAb ($M_r = 260$ kDa). The DNAs of MDB1 light chains and NK-biAb heavy chains were co-transfected into Expi293F cells using PEI transfection reagents. After the transfection for 96 h, the culture supernatant was collected by centrifugation (4000 rpm, 10 min) and purified on a protein A column. The purified protein was further polished on a Superdex 200 Increase column in a PBS (pH 7.4) buffer. To validate the functions of NK-biAb to attach NK92 cells and bind NKG2D, the NK92 cells pre-stimulated with IL-2 were cultured with NK-biAb at 37°C for 2 h, and stained with anti-human IgG-PE for the detection of surface NKG2D.

Stable RIFIN#1-expressing K562 cell line production

K562 cells (Procell, China) were cultured at 37°C in DMEM with 10% FBS and 5% CO_2. The cDNA of RIFIN#1 with its natural signal peptide was cloned into pLVX-ZsGreen plasmid. Lentivirus were produced by HEK-293T (Procell, China) for infecting K562 cells. The infected K562 cells were enriched by the selection using 1 μg/mL puromycin, to obtain the stable RIFIN#1-expressing K562 cell line.

Evaluation of NK cell cytotoxicity

The RIFIN#1-expressing K562 cells were incubated at 37°C for 1 h with D1D2.v-IgG or NK-biAb. The NK92 cells in the same volume as RIFIN#1-expressing K562

cells were then cocultured for 4 h at 37°C and 5% CO_2 with the incubated RIFIN#1-expressing K562 cells at the E: T ratios of 2.5: 1 and 12.5: 1. The final mixtures contain either 100 nM D1D2.v-IgG or 50 nM/100 nM NK-biAb. The killing efficiency (E) was calculated from the absorbance (A) of CCK8 at 450 nm: $E\% = [1-(A_S-A_{EC})/A_{TC}]\times100\%$, where A_S is for the mixture of effector and target cells, and A_{EC} and A_{TC} are, respectively, for effector cells and target cells.

Cytokine measurement and FCM detection

10^5 NK92 cells were cocultured for 24 h with 10^5 RIFIN#1-expressing K562 cells in a 96-well plate in the presence of either 100 nM D1D2.v-IgG or 50 nM/100 nM NK-biAb. Perforin and granzyme B release were detected from the supernatants of the cocultured mixture by protein-based ELISA assay kits (Beyotime, China). The mixtures of NK92 and K562-RIFIN#1 cells were stained for 1 h at ca. 4ºC (controlled by ice) with Annexin V-APC in 100 µL PBS containing 0.5% BSA. The stained cells were detected by flow cytometry, with the data analyzed by Flowjo software.

Reference

1. Organization., W. H. World Malaria Report *World Malaria Report* (2016).

2. Burton, D. R. Antibodies, viruses and vaccines. *Nature Reviews Immunology* **2**, 706-713 (2002).

3. Haldar, K., Bhattacharjee, S. & Safeukui, I. Drug resistance in *Plasmodium*. *Nature Reviews Microbiology* **16**, 156-170 (2018).

4. Burton, D. R. What Are the Most Powerful Immunogen Design Vaccine Strategies? Reverse Vaccinology 2.0 Shows Great Promise. *Cold Spring Harb Perspect Biol* **9**(2017).

5. Wells, T. & Donini, C. Monoclonal Antibodies for Malaria. *New England Journal of*

Medicine **387**, 462-465 (2022).

6. Nilsson, S. K., Childs, L. M., Buckee, C. & Marti, M. Targeting Human Transmission

Biology for Malaria Elimination. *PLOS Pathogens* **11**, e1004871(2015).

7. Sakoguchi, A. & Arase, H. Mechanisms for Host Immune Evasion Mediated by

Plasmodium falciparum-Infected Erythrocyte Surface Antigens. *Frontiers in*

Immunology **13** (2022).

8. Chan, J. A., Fowkes, F. J. & Beeson, J. G. Surface antigens of *Plasmodium*

falciparum-infected erythrocytes as immune targets and malaria vaccine candidates.

Cell Mol Life Sci **71**, 3633-3657(2014).

9. Harrison, T. E. *et al.* Structural basis for RIFIN-mediated activation of LILRB1 in

malaria. *Nature* **587**, 309-312 (2020).

10. Saito, F. *et al.* Immune evasion of *Plasmodium falciparum* by RIFIN via inhibitory

receptors. *Nature* **552**, 101-105 (2017).

11. Xie, Y. *et al.* Structural basis of malarial parasite RIFIN-mediated immune escape

against LAIR1. *Cell Rep* **36**, 109600 (2021).

12. Willcox, B. E., Thomas, L. M. & Bjorkman, P. J. Crystal structure of HLA-A2

bound to LIR-1, a host and viral major histocompatibility complex receptor. *Nature*

Immunology **4**, 913-919 (2003).

13. Wahl, I. & Wardemann, H. How to induce protective humoral immunity against

Plasmodium falciparum circumsporozoite protein. *Journal of Experimental Medicine*

219 (2022).

14. Cockburn, I. A. & Seder, R. A. Malaria prevention: from immunological concepts

to effective vaccines and protective antibodies. *Nature Immunology* **19**, 1199-1211,

doi:10.1038/s41590-018-0228-6 (2018).

15. Chen, Y. *et al.* Structural basis of malaria RIFIN binding by LILRB1-containing

antibodies. *Nature* **592**, 639-643 (2021).

16. Tan, J. *et al.* A LAIR1 insertion generates broadly reactive antibodies against

malaria variant antigens. *Nature* **529**, 105-109 (2016).

17. Xu, K. *et al.* Structural basis of LAIR1 targeting by polymorphic *Plasmodium*

RIFINs. *Nature Communications* **12**, 4226 (2021).

18. Chao, G. *et al.* Isolating and engineering human antibodies using yeast surface

display. *Nature Protocols* **1**, 755-768(2006).

19. Weiskopf, K. *et al.* Engineered SIRPα variants as immunotherapeutic adjuvants to

anticancer antibodies. *Science* **341**, 88-91(2013).

20. Gonzalez-Perez, D. *et al.* Affinity-matured DLL4 ligands as broad-spectrum

modulators of Notch signaling. *Nature Chemical Biology* **19**, 9-17(2023).

21. Kwong, K. Y., Baskar, S., Zhang, H., Mackall, C. L. & Rader, C. Generation,

affinity maturation, and characterization of a human anti-human NKG2D monoclonal

antibody with dual antagonistic and agonistic activity. *J Mol Biol* **384**, 1143-

1156(2008).

22. Zhang, C. *et al.* Bispecific antibody-mediated redirection of NKG2D-CAR natural

killer cells facilitates dual targeting and enhances antitumor activity. *J Immunother*

Cancer **9** (2021).

23. Zhao, C. *et al.* A new approach to produce IgG (4)-like bispecific antibodies. *Sci*

Rep **11**, 18630 (2021).

24. Wolf, N. K., Kissiov, D. U. & Raulet, D. H. Roles of natural killer cells in immunity

to cancer, and applications to immunotherapy. *Nature Reviews Immunology* **23**, 90-105

(2023).

25. Albinger, N., Hartmann, J. & Ullrich, E. Current status and perspective of CAR-T

and CAR-NK cell therapy trials in Germany. *Gene Therapy* **28**, 513-527 (2021).

26. Leaver-Fay, A. *et al.* Computationally Designed Bispecific Antibodies using

Negative State Repertoires. *Structure* **24**, 641-651 (2016).

Acknowledgements: Funding was provided by the National Key R&D Program of China (Grant No. 2023YFA1607500), the National Natural Science Foundation of China (Grant No. 32371362), and the Anhui Major Basic Research Projects (Grant No. 2023z04020016).

Author contributions: Y.W and H.T contributed equally to this work. Y.W initiated the project, performed the library construction and selections, prepared antigen samples and performed biochemical assays. H.T performed the FCM detections and conducted the stable cell line production, cytotoxicity assays and ELISA assays. W.W performed the antibody purifications and functional validations. L.M helped to prepare biotinylated sample preparations. C.Z helped to perform biochemical assays and FCM detections. Y.W. and B.W analyzed the data, interpreted the results. J.W, H.D and B.W prepared the manuscript. J.W., B.W and H.Z supervised the project. All authors read and edited the manuscript.

Competing interests: The authors declare no competing interests.

Data and materials availability: All data and materials are available within the paper and its supplementary materials. Extra data are available from the corresponding author (J. Wang) upon reasonable request.

List of Supplementary Materials
Fig S1 to S8 for multiple supplementary figures
Table S1 for supplementary table