Complex genetic determination of male-fertility restoration in the gynodioecious snail

Physa acuta

Elpida Skarlou [1,2] **& Fanny Laugier** [1,2]**, Kévin Béthune** [1]**, Timothée Chenin** [1]**, Jean-Marc**

Donnay [1]**, Céline Froissard** [1]**, Patrice David** [1]

[1] CEFE, CNRS, Univ Montpellier, EPHE, IRD, Montpellier, France

[2] co-first authors

Abstract

Male fertility in plants is often controlled by the interaction between mitochondrial and

nuclear genes. Some mitotypes confer cytoplasmic male sterility (CMS), making the

individual male-sterile, unless the nuclear background contains alleles called restorers, that

suppress the effects of CMS and restore the hermaphroditic condition. Restorers in cultivated

crops are often alleles with strong and dominant effect, but in wild plants, data often suggest

more complex systems. Here, we characterized the inheritance and specificity of restoration

in a new CMS model, the freshwater snail *Physa acuta*. We explored two different

populations (i) a naive population i.e., without contact CMS in the past 80 generations, (ii)

a non-naive population, where CMS is present and largely restored. We found that

restoration potential was indeed heritable in both contexts and specific to a CMS type.

However our results suggest that restoration of male fertility could be of a different nature

depending on population history: in naïve populations not coevolved with CMS the

background variation may include alleles that happen to act as weak quantitative modifiers

of the penetrance of CMS, while in populations coevolved with CMS, selection may have

favored, when such variants were available, the emergence of strong alleles with a dominant

effect.

Key words: cytoplasmic male sterility, restoration, genetic determination, *Physa acuta*

Introduction

Gynodioecy is a sexual polymorphism, common in angiosperms (Dufay et al., 2014; Caruso et al., 2016), whereby hermaphrodites coexist with male-sterile individuals (i.e., functionally females, Saumitou-Laprade et al., 1994). Gynodioecy represents a well-known example of genetic conflict (Burt & Trivers, 2006) because sex determination in gynodioecious systems is often cyto-nuclear. In that case, cytoplasmic male sterility (CMS) genes, usually associated with the mitochondrial genome, suppress pollen production and render an individual functionally female unless CMS is counteracted by one or more nuclear genes that restore the male function (Cosmides & Tooby, 1981; Saumitou-Laprade et al., 1994; Werren & Beukeboom, 1998). An individual is phenotypically female if it carries CMS without the proper nuclear gene known as a restorer. It is a hermaphrodite if it carries either (i) a male-fertile cytoplasm, or (ii) a CMS cytoplasm paired with the proper nuclear restorer. CMS has been documented in 140 species of angiosperms (Laser & Lersten, 1972), and twenty-eight CMS genes from 13 crop species have been identified (Chen & Liu, 2014).

It appears that cyto-nuclear gynodieocious systems often contain multiple forms of CMS (e.g. *Oryza sativa, Zea mays*, see Chen and Liu 2014 for a review), each requiring its own mode of restoration (A. de Haan et al., 1997; Charlesworth & Laporte, 1998; van Damme et al., 2004). Fourteen restorer genes (*Rf*) from nine crops have been isolated (see Table 1 Kim & Zhang, 2018), with the majority of them encoding PPR (Pentatricopeptide Proteins Repeat). However, *Rf* genes encoding for other proteins have been discovered, including peptidase in sugar beet (Yamamoto et al., 2008; Matsuhira et al., 2012) and aldehyde dehydrogenase in maize (Cui et al., 1996). In most cases, a *Rf* gene is specific to one CMS, whereas a CMS gene can be affected by multiple *Rf* genes (see Table 1 Chen and Liu 2014). For example, in *Z. mays* the CMS-T is restored by both *Rf1* and *Rf2* (Cui et al., 1996; Dewey et al., 1987; Dill et al., 1997) and in *Plantago coronopus* at least five restorer alleles seem to be involved in the restoration of CMS (Koelewijn & Van Damme, 1995). Cases in which a gene restores more than one CMS mitotype are rare. Such an example is found in the common rice, where *Rf5* and *Rf6* (initially identified in CMS-HL individuals) are also able to restore CMS-BT mitotype (Huang et al., 2012, 2015; Kim & Zhang, 2018).

The maintenance of cyto-nuclear polymorphism has been extensively modelled (e.g. (Charlesworth & Ganders, 1979; Couvet et al., 1998; Gouyon et al., 1991 ; Bailey et al., 2003 ; Dufaÿ et al., 2007) and theoretical studies have shown that the stability of this polymorphism depends on a variety of factors, including the positive pleiotropic effects

of CMS alleles (i.e., seed fitness advantage of females, or "female advantage", Lewis, 1941) and negative pleiotropic effects of restorer alleles ("cost of restoration", Charlesworth & Ganders, 1979; Delannay et al., 1981; Frank, 1989; Gouyon et al., 1991). Models often assume that restoration of male fertility is achieved through one allele (but see Frank 1989) and a variety of assumptions regarding its dominance of restoration have been modeled (Delph et al., 2007). The majority of those models considered the presence of dominant restorer alleles with only a couple a cases using a recessive inheritance (Ross & Gregorius, 1985). Regarding empirical studies, most of the crop species exhibited dominant restorer alleles (see Table 2 Delph 2007 for a review but see also *Plantago lanceolata*, van Damme, 1984 ; de Haan et al., 1997). However, in natural populations the restoration of male fertility suggests a more complex genetic determination (Charlesworth & Laporte, 1998; Koelewijn, 2003; Touzet et al., 2004). Indeed, several nuclear loci may act in an additive manner (Charlesworth & Laporte, 1998; Dufay et al., 2008) and/or through a threshold effect (Ehlers et al., 2005). For example, Koelewijn & van Damme (1995) suggested that at least five restorer loci were involved in restoring male sterile cytoplasms of *Plantago coronopus*, and Dudle et al., (2001)found that sex determination in *Lobelia siphilitica* seemed to vary among cytoplasms with one CMS gene being restored by a single dominant allele, while restoration of other CMS could only be explained by the action of several nuclear loci and/or epistatic effects. Inspired by those examples, Elhers et. al (2005) developed a threshold model for sex-determination in gynodioecious species and showed that treating restoration of CMS as a quantitative trait can be an alternative way of explaining complicated patterns of restoration in CMS species. Following on those steps, Bailey & Delph (2007) described a model of cyto-nuclear controlled gynodioecy, in which sex was determined by two CMS types each one interacting with multiple restorer alleles. The used assumptions reflected several well-studied cases of gynodioecy, such as, *P. coronopus, P. lanceolata, S. vulgaris*, and *T. vulgaris* and showed that females can be maintained in the population (Belhassen et al., 1991; de Haan et al., 1997; Koelewijn, 2003; Mccauley & Olson, 2003).

A mix of several determinisms is an interesting situation to consider. Dominant and specific restorers acting on one or a few CMS mitotypes, found in present-day plant populations, are strong-effect alleles that may result from a history of coevolutionary arms-race with their specific CMS gene (e.g. Sloan et al., 2012), or at least from the selective sweep of a favorable mutant (Bergero et al., 2019). When a new CMS type first appears, it is unlikely that such strong-effect specific restorer alleles are already ready and waiting.

However, the standing variation may contain genotypes that happen to reduce the penetrance of CMS, not necessarily in a way that fully restores male-fertility, and not necessarily in a specific way. Such a background variation may contribute to the cases of complex inheritance in plants, but is expected to be more visible when there are not yet specific large-effect restorers segregating in the population, a relatively rare condition (refs).

As illustrated above, CMS is well understood theoretically and prevails in plants but is exceedingly rare among animals (Vellnow et al., 2017). However, CMS has been recently reported in one animal: *Physa acuta*, a freshwater snail (David et al., 2022). In this species, two mitochondrial types conferring male sterility, D and K, have been discovered so far, both of them showing an extreme molecular divergence from one another, as well as from normal, fertile cytotypes (collectively called N), at all mitochondrial genes. The D mitochondrial type was discovered first, but evidence for corresponding nuclear restoration is still lacking (David et al. 2022). Recently however, the second CMS mitotype K revealed a different situation, with evidence that most K individuals captured in natural populations have a male-fertile phenotype, i.e., there exist genes that restore male fertility in K-mitotype snails, while male-sterility reappears when the K mitotype is introgressed into a "naïve" laboratory genetic background maintained without contact with CMS (Laugier et al., 2024).

In this study, we explored the inheritance and specificity of restoration in *Physa acuta*. We aimed to characterize the variation in restoration for the K mitotype, specifically to test if the data are consistent with the existence of dominant-restorer alleles in natural populations where K is present. We also tested whether these restorers are specific to K or also have an effect on male-fertility of the D mitotype. Finally, we were interested in the determination of male-fertility in our laboratory population, where the K-mitotype, although mostly male-sterile, still maintains a fraction of male-fertile individuals. This population may represent a truly "naïve" state, i.e. the state of a population in which CMS would be newly introduced and confronted to standing variation; or it may have inherited, from many generations ago, the same restorer alleles as in natural populations where K is present, but these alleles would now be at a low frequency.

To achieve these objectives, we phenotyped CMS K- and D-mitotype individuals within two different nuclear backgrounds (Lyon and Montpellier). The Lyon population has been in contact with both K and D, while the Montpellier has not. We used two different approaches to reveal genetic variation in restoration potential within Lyon and Montpellier population: we produced populations selected against and for restoration, and we analysed inbred lines. This approach enables us to explore restoration in various contexts and

characterize it accordingly. Answers to these questions will provide insight into the CMS evolutionary dynamics and, for the first time, explore genetic variation for CMS restoration in animals.

Material and methods

Study system

Physa acuta is a cosmopolitan, hermaphroditic, predominantly outcrossing, freshwater snail, very common in natural freshwaters worldwide, and easy to raise in the laboratory. The generation time is about 6-8 weeks in the laboratory at 25°C ; we maintain this snail in standard conditions (25°C, 12/12 photoperiod) using ground water changed at least once a week, and grinded boiled lettuce ad libitum as food. Cytoplasmic male sterility has been first described by David et al. (2022) in this species, later completed by Laugier et al. (2024). Briefly two very divergent mitogenomes, named D and K, have been found to coexist with the normal (N) type in wild populations near Lyon (France). While N individuals are regular hermaphrodites, D and K are associated with male-sterile phenotypes, unable to play the male role in sexual interactions and to sire offspring. Nearly all the D (92%, David et al. 2022) individuals are male-sterile. In contrast, nuclear backgrounds restoring male-fertility of K snails are found in natural populations where K is present in Lyon (France). However the same K mitotype, when introgressed into the nuclear background of a population from Montpellier (hereafter « naive », maintained for >80 generations in the laboratory without contact with CMS), produces around 70% male-sterile phenotypes (Laugier et al., 2024). The nature of the ~30% residual fertility is still hypothetical, it could be either that restorer genes are segregating at low frequency in the naive population (despite the absence of K mitotypes) ; or that the penetrance of CMS is incomplete, always leaving a fraction of individuals with nonzero male fitness.

General structure of the experiments

Our objective was to characterize, within both the naive (Montpellier) and the non-naive (Lyon) backgrounds, genetic variation for the potential to restore male fertility in CMS mitotypes. We did this in two ways (i) first, we tested whether we could select for high versus low restoration potential within the naive population (ii) second, we produced inbred lines from both populations and characterized genetic variance in restoration potential among lines (trying, in particular, to find lines fixed for either restorer or maintainer alleles). In both

cases, lines were of the N mitotype, and their restoration potential was evaluated by crossing them with either a K or a D snail, and evaluating the male fertility of the offspring. In these crosses, the paternal strains were N while the maternal ones were K or D. As mentioned above, all (or nearly all) our D individuals are un-restored, mostly male-sterile individuals. However, two K strains are available, one with the original background from Lyon (with high restoration potential and mostly male-fertile, noted KF) and one with the naive background from Montpellier (low restoration potential, mostly male-sterile, noted KS). We used both types to be in position to observe the effects of potential restorer alleles provided by the paternal strain either in a predominantly homozygous (cross with KF) or heterozygous (cross with KS) form.

Source populations and their history in the laboratory

The Montpellier population has been founded in 2007 by a mix of ten natural populations sampled around Montpellier, and maintained in the laboratory, in high numbers, since then (Figure 1). The albino stock was separated from the main population in the first few laboratory generations (see Noël et al., 2016). It has been maintained as an autonomous large outbred population in aquaria to this day (estimated age in 2023: 96 generations) and is entirely of the N mitotype and male-fertile. Albinism is coded by a recessive allele, and albino mothers produce pigmented babies when inseminated by a wild type snail. More recently, a new type of albinotic mutants appeared within the Montpellier stock, with a slightly different phenotype (see Figure S1 and S1). After some crosses (see S1) we were able to establish it as an independent outbred stock (called alb-2) that complements with the first albino stock (now called alb-1): alb-1 and alb-2 are mutated at different loci, and crosses between them produce 100% pigmented individuals, as well as crosses between any of the two and pigmented homozygous snails.

The Lyon populations have been founded from two populations sampled in 2019 near Lyon (Irigny and Erevan, see Laugier et al. 2024) mixed together. They initially contained a mix of K and N mitotypes, the former with mostly male-fertile phenotypes (see above) and all pigmented (wild-type). After mass-mating, we collected progenies from individual mothers that were typed by PCR as either K or N; and the K and N progenies were pooled into two separate tanks, to initiate pure-K and pure-N stocks (Figure 1). Each stock was kept as a separate population (hereafter KF and NF populations) and maintained as a large outbred population to this day (in 2023, KF and NF were at generation 20 after foundation). Each of the outbred populations (alb-1, alb-2, NF, KF) was maintained throughout as a

metapopulation of 5 aquaria, each with 100-150 snails at reproductive stage, freely mating
(« mass-mating »), with regular exchange of ~10% individuals among aquaria; the eggs are
separated from the parents to hatch as soon as there are enough on the aquarium walls, and
a new cohort is started. Aquaria are fed once to twice a week depending on life stage.

A last source of snails for our experiments is a set of lines in which the D or the K mitotypes
have been introgressed into the naive Montpellier alb-1 background (David et al 2022,
Laugier et al 2024). Briefly this introgression takes place in sets of « matrilines » each
founded by a D or K ancestor from Lyon (wild population) and propagated by maternal
descent. At each generation each snail from a matriline is inseminated by an individual taken
at random from the alb-1 stock so that over generations the genome of the founder is
progressively replaced by the alb-1 background. At the time of the experiment the D
matrilines were at the 41th generation and the K matrilines at the 20th generation, their
genome can be considered as totally replaced. Matrilines are maintained pigmented and
heterozygous to distinguish them from their albino mates: at each generation we obtain a
mix of albino and pigmented progeny, of which we keep only the latter to propagate the
matriline. For the experiment however it was practical to use the albino progeny. Both K and
D matrilines are mostly male-sterile (as mentioned above); for this reason we will refer to
them as KS and DS respectively (by contrast, KF snails are restored and mostly male-fertile).

Figure 1: **Source populations and their history in the laboratory since extraction from
natural populations (indicated by dashed arrows, with year of sampling).** Thick lines

indicate propagation in large outbred populations in aquaria. Fine lines indicate propagation

in matrilines, with insemination at each generation by sperm donors from another population

(dotted arrows). The percentages of each mitotype (K, D or N) are indicated for the natural

populations at sampling (from David et al.2022, Laugier et al. 2024) and for populations

kept in the laboratory. The number of generations elapsed since extraction from natural

populations in late 2022 – early 2023 (when most of the experiment took place) is indicated

as G_{XX} below each stock. The numbers of independent matrilines are 12 for KS and 11 for

DS. Populations from which inbred lines were extracted are indicated by a star (*). The

coordinates of the populations of origin can be found in Noël et al. 2016 (Montpellier), David

et al. 2022 (Lyon, Autoroute), and Laugier et al. 2024 (Lyon, Irigny and Erevan)

Selection for high and low frequency of restoration of the K mitotype

The aim of this selection was to mimick the arrival of a CMS mutant in a naive population

and test whether there was variation from which restoration potential could be rapidly

selected in such a situation. This was envisaged for the K mitotype because the penetrance

of the male-sterile phenotype in KS individuals (mitotype K introgressed into the naive alb-

1 background) is around 60-70% ; if the residual fertility is genetic it can increase or decrease

by selection. On the other hand, the residual fertility of DS snails is too low to start a similar

selection.

The selection protocols are detailed in supplementary S2. In short, to select against

restoration, we kept the (N-mitotype) alb-1 sperm donors that, when crossed with a KS snail,

produced only male-sterile K daughters. The selected sperm donors where then grouped and

constituted the first generation of selection for low-frequency-of-restoration (LFR). We re-

used offspring of these individuals as sperm donors for a second round of selection. The LFR

individuals used here are therefore considered to be at their second generation of selection

against restoration. The selection for high frequency of restoration (HFR) is simpler because

selecting for fertility is easier than for sterility: we simply kept the offspring sired by male-

fertile KS individuals at each generation. The HFR population was selected for two

successive generations, as the LFR. Expected results of such a selection in the simplest

scenario (fully penetrant, monodominant restorers) are given in S2.

Constitution of inbred lines

Inbred lines allow in principle to fix restoring and non-restoring genotypes in the

homozygous form. We created inbred lines from N-mitotype ancestors originating from

either the Lyon or the Montpellier laboratory populations (more lines were made from Montpellier where restoration is supposed to be rare). Each ancestor produced a line by at least two generations of selfing (enforced by letting a virgin snail reproduce in isolation) and single-individual descent. Lines are not fully homozygous at the second generation (there remains ¼ of the initial heterozygosity) but we had to find a compromise with the time required and with the potential loss of lines by inbreeding depression over generations. In this experiment we used 9 lines from Lyon and 40 from Montpellier.

Crosses

In total we tested the restoration potential of focal individuals from two selection lines (HFR and LFR) and from 49 inbred lines of two origins: Montpellier (naive population= and Lyon (experienced population). To that end these focal individuals were used to inseminate three types of individuals: DS, KS and KF. The resulting progeny has the focal individual as father and DS, KS or KF mother. In each progeny we grew one to three G1 offspring until maturity and tested them for male fertility. If the father transmits restorer alleles, these alleles are present in the tested offspring either in a mostly heterozygous form (mothers DS or KS, with absent or rare restoration), or in a mostly homozygous form (mothers KF, with a high rate of restoration). The total number of progenies tested was 1508, with the repartition indicated in Table 1.

Table 1: Sample sizes by category. For each combination of paternal and maternal types, we provide N_1/N_2, where N_1 is the number of G_1 individuals successfully assessed for male fertility, and N_2 is the number of G_0 mothers (each mother produced 1 or 2 G_1 individual).

		Maternal			
		KS	KF	DS	Total
	Lyon inbred lines	164/82	36/21	65/47	265/150
	Montpellier inbred lines	350/231	220/130	211/168	781/529
	Total inbred lines	524/313	256/140	276/215	1056/668
Paternal	HFR selected population	52/31	49/27	32/22	133/80
	LFR selected population	248/105	39/20	32/25	319/150
	Total selected populations	300/136	88/47	64/47	452/230

 Assessment of male sterility

The assessment of male sterility of an individual is made by pairing it to a virgin partner for

three days, letting the latter lay for three days in isolation, and observing the progeny. The

studied individual is said to be male-sterile if either the partner does not lay any egg (failure

to stimulate egg-laying, which is a normal outcome of insemination) or it lays eggs but the

juveniles are 100% self-fertilized. Partners of male-fertile individuals in this situation

normally produce several tens of eggs and juveniles are nearly all outcrossed. A small

number of individuals had an intermediate status: outcrossed individuals were obtained but

in abnormally low numbers (<10) ; in this case we recorded them as semi-sterile. In the

analyses we grouped these together with fully male-sterile individuals (without appreciable

consequence on the results). It is, in any case, important to realize that the male-fertility test

is not perfect: even male-fertile individuals (with fertile mitotype) occasionally fail to

inseminate their partner during pairing, with a probability <10% in our lab conditions.

We used body pigmentation to recognise outcrossed offspring from self-fertilized

ones. Depending on the paternal pigmentation genotype, we used either albino alb-1 or alb-

2 partners in such a way that any pigmented offspring they produced must have been sired

by the tested individuals. We therefore simply counted the number of pigmented offspring

in the progeny after incubating clutches for 13 days (we stopped counting at 20 and noted

20+ if all offspring were pigmented, and counted exhaustively all pigmented and albino

offspring otherwise, including those that were well-developed though not yet hatched, if

any). In some cases (heterozygous individuals tested with an alb-1 partner) outcrossed

offspring were expected to be 50% pigmented, 50% albino instead of 100% pigmented; in

this case we set the threshold for semi-sterility to 5 pigmented (i.e. the expectation for 10

outcrossed offspring) instead of 10 (see above).

We tested male fertility in G1 individuals and in their DS, KS, KF mothers too. Many

of the latter were pigmented and mated with albino N-mitotype fathers; thus the father could

act as a partner for testing the male-sterility of the mother, and we kept the father's progeny

to that end. In case the father was a pigmented individual and/or the mother an albino, this

was not possible so we provided a second partner to the mother, after she had laid eggs, and

this second partner was alb-2 so that outcrossed offspring would be recognised (alb-1 and

alb-2 are complementary lines and produce pigmented babies upon cross-fertilization).

Because of the size of the experiment, it was spread over nearly two years from 2022 to early

2024, with four phases treated as temporal blocks.

Statistical analyses

We analysed separately the differences between the selected populations (HFR and LFR) on one hand, and the variation among inbred lines on the other hand. The male-fertility status of the G_1 individuals from selected lines was treated by GLMMs as a binary 0/1 variable. The main model included the maternal type (KF, KS or DS), the selected line (HFR or LFR) and their interaction as fixed effects, and the identity of the mother and the block as random effects. We also made more detailed models including the maternal status (fertile or sterile) in the fixed effects. The male-fertility status of the G_1 individuals from inbred lines was treated in a similar way, except that the fixed effects included: the geographic origin of the inbred line (Montpellier versus Lyon), the maternal type (KF, KS or DS) and their interaction; and the random effects included the identity of the inbred line, in interaction with maternal type, in addition to block and mother identity. All tests were performed by likelihood-ratio chi-squares, comparing models with and without the effect of interest with all other effects kept. The models were run using the `lme4` package in R (Bates et al., 2015).

Predictions

The selected lines inform on the nature of the variation in male-fertility observed when CMS is introduced into a naive background (the Montpellier laboratory population), i.e. why is there a residual male fertility in K individuals introgressed into this background ? If there exist restorer alleles with large effects, we expect the selection protocol to be efficient, and the frequency of male-fertility of K individuals sired by HFR fathers should exceed that of their congeners with LFR fathers. Depending on dominance, this effect should be more visible in different circumstances. A dominant allele transmitted by fathers makes a difference when the mother does not herself bring the same allele (in our case, with male-sterile mothers, and/or with mothers from the KS strain). A recessive allele, on the contrary, would have more effects when the mother is male-fertile and/or from the KF strain. Finally, the specificity of restoration with respect to the mitotype can be tested by checking whether the LFR/HFR difference, if any, affects male-fertility when the mother is of the D mitotype.

The inbred lines allow to document the nature and specificity of restoration in both the Montpellier and Lyon populations. Based on previous studies (Laugier et al. 2014) we expect that K-mitotype individuals sired by Lyon lines (a largely restored population) will have more male-fertile offspring than those sired by Montpellier. We expect, if restoration has a simple genetic basis, that some lines will be homozygous for restorers and others for maintainer alleles, generating high variation in male-fertility among lines, either when

crossed with KS / male-sterile mothers (if restorers are dominant) or when crossed with KF

/ male-fertile mothers (if recessive). Similar effects could be observed upon crossing with

DS if there exist restorers efficient against CMS brought by the D mitotype; restoration may

be specific (in that case, the same lines should restore fertility in both D and K) or not.

Results

Male-sterility in the maternal generation

The phenotypes of the mothers used to test the restoration potential in inbred lines and

selected (HFR/LFR) lines were distributed in the same way and conformed to expectations

based on previous studies (see Material and methods): mothers with CMS introgressed into

the alb-1 Montpellier background were mostly male-sterile with (in total) 5.7% of male-

fertile individuals among D-mitotype mothers (DS) and 27.2% in K-mitotype ones (KS). In

contrast nearly all (92.3%) mothers of the KF type were male-fertile. In both datasets the

proportion of male-fertile individuals differed very significantly among types (Binomial

GLMMs, χ^2_{1df} =119.9, and 36.6 respectively for the inbred lines and selected lines datasets,

both $p < 0.001$).

Figure 2: Proportions of individuals with male-sterile, male-fertile and male-semi-

sterile phenotype in the maternal generation. Proportion are represented for both datasets

as a function of maternal type DS: D-mitotype introgressed into the alb-1 background from

Montpellier; KF : K-mitotype within its native background (Lyon) ; KS : K-mitotype

introgressed into the alb-1 background from Montpellier. The two datasets are presented

separately because they were made at different times.

Male sterility in the offspring generation: LFR versus HFR

The proportions of male-fertile offspring closely mirror those observed in their mothers (Figure 3, compare with Figure 2), and the maternal type (DS, KS, or KF) accordingly has a very significant effect on the probability of being male-fertile (p <0.001, Table 2). The paternal line (HFR or HFR) has a weakly significant effect (p = 0.039, Table 2), and this effect is perceptible only when the mother has a K mitotype introgressed into the alb-1 Montpellier background (KS mothers, Table 2). In this category the proportion of male-fertile offspring is slightly higher when the father comes from the LFR line than when the father comes from the HFR line.

The effect of the maternal phenotype (i.e. whether the mother was a male-sterile or a male-fertile snail) cannot be studied in the DS and KF categories because of insufficient numbers (only one DS individual with a male-fertile mother; only two KF individuals with a male-sterile mother). In the KS category however, numbers are more balanced (98 individuals with male-fertile mothers, versus 202 with male-sterile or -semisterile mothers), and the maternal phenotype partially predicts the offspring's one. Indeed, male-fertile mothers tend to have more male-fertile offspring, whatever the paternal line; the influence of the paternal line remains significant, and in the same direction as previously, when the maternal phenotype is taken into account (GLMM: effect of paternal line χ^2_{1df} = 4.65, p = 0.031 ; effect of maternal phenotype χ^2_{1df} = 4.48, P = 0.034 ; interaction χ^2_{1df} = 0.15, p = 0.70 ; Figure 3).

Note that full-sibs (offspring sharing the same mother and father) tend to have similar phenotypes as attested by significant effects of mother identity (Table 2) on top of what is already explained by the fixed effect of maternal type. This mother effect is not completely predicted by maternal phenotype (among KS mothers) as it remains approximately the same in the model with the maternal phenotype included as a fixed effect (χ^2_{1df} = 4.80, variance = 0.72, p = 0.014). Note that the mother effect is also, surprisingly, significant within the KF category (Table 2) ; this is because among the few (four) individuals with KF mother and HFR father observed to be male-sterile, two have the same mother which is unlikely by chance.

Table 2: Generalised linear mixed models on the binomial male-fertility variable (fertile versus sterile or semi-sterile). LRT tests are given for fixed and random factors (χ^2_{df} and corresponding p-values). For random factors we also provide estimates of variance

(v, on logit scale) and *p*-values account for the one-sided alternative (variances cannot be

negative). Note that the very high « maternal ID » variance in the KF category is an effect

of the logit scale when proportions are close to 0 or 100% (here 92% individuals are fertile):

in these conditions a very large variance in the logit scale makes a small variance in actual

proportions. The block random factor was omitted in this table because in all models the

estimated variance was 0.00. *p*-values are bolded when <0.05.

Data	Fixed effects			random effects
	maternal type (DS, KF, KS)	paternal type (HFR, LFR)	Interaction	maternal ID
All	$\chi^2_{df2} = 64.36$ *p* = **1.06 10⁻¹⁴**	$\chi^2_{df1} = 4.25$ *p* = **0.039**	$\chi^2_{df2} = 0.91$ *p* = 0.634	$\chi^2_{df1} = 6.79$, v = 1.01 *p* = **0.0045**
only DS		$\chi^2_{df1} = 0.356$ *p* = 0.551		$\chi^2_{df1} = 0.00$, v = 0.00 *p* = 0.500
only KF		$\chi^2_{df1} = 0.001$ *p* = 0.970		$\chi^2_{df1} = 10.07$, v = 91.72 *p* = **0.008**
only KS		$\chi^2_{df1} = 4.23$, *p* = **0.039**		$\chi^2_{df1} = 6.79$, v = 0.76 *p* = **0.010**

418   **Figure 3: Proportions of individuals with male-sterile, male-fertile and male-semi-**

419   **sterile phenotype**. Left: proportions of male-fertile and male-sterile phenotypes as a

420   function of paternal selection line (HFR versus LFR), maternal type (DS, KF, KS). Right:

421   detailed view of proportions in offspring of KS mothers, as a function of paternal line and

maternal phenotype (male-fertile versus not male-fertile). Similar divisions were not considered in the DS and KF category because of the low numbers of male-fertile and non male-fertile mothers respectively.

Male-sterility in the offspring generation: inbred lines

As previously, maternal types (DS, KS or KF) induced large contrasts in the proportion of male-sterility, with DS being predominantly male-sterile, KF predominantly male-fertile, and KS in an intermediate situation (Table 3, Figure 4). Only in the KS category, the origin of the paternal inbred line made a difference: paternal lines from Montpellier (laboratory population entirely of the N mitotype) had largely male-sterile offspring, while paternal lines from Lyon (where K and D were initially found) had more than 50% male-fertile offspring. In these KS offspring, the variance explained by paternal lines was significant (Table 3), on top of what was already explained by the geographic origin, and the among-lines variance, when estimated separately, was higher in the Lyon population compared to Montpellier (Table 3). This can be visualised by representing BLUPs (Best Linear Unbiased Predictors) associated with paternal lines in the linear models: the Montpellier inbred lines span a range of male-fertility between 0 and 50% while the Lyon lines cover essentially the whole range (0-100%) (Figure 4). The apparent continuity of the distribution does not seem to be an artefact of the GLMM (which assumes an underlying normal distribution on the logit scale for random effects) as it is also observed when plotting the raw percentages of male-fertile individuals (supplementary Figure S4). In contrast with KS, the variance among paternal inbred lines is estimated to be zero in the KF category. However it is significant in the DS category, in which the distribution of BLUPs spans the 0-50% interval with an apparently random distribution of lines from Montpellier and Lyon. The predicted male-fertility of offspring of an inbred line in a DS context is not significantly correlated to the same line in a KS context (Spearman $\rho = 0.22$, NS, Figure 4).

Models including the maternal phenotype (male-fertile or not) in the predictors, in addition to paternal population of origin, showed no significant effect in KS ($\chi^2_{1df} = 0.018$, $p = 0.895$), and a weakly significant effect in the DS context ($\chi^2_{1df} = 4.37$, $p = 0.037$) where male-fertile mothers had more male-fertile daughters than male-sterile mothers (see plots in S3). The number of male-sterile mothers was insufficient to test a difference in the KF category.

Table 3: Generalised linear mixed models on the binomial male-fertility variable
(fertile versus sterile or semi-sterile), inbred lines dataset. Legend as in Table 2. The
block random factor was omitted because the estimated variance was 0.00. [1] the test gives
the significance of the variance explained by the paternal line, in interaction with maternal
type. Because of the significance of the interaction, the variances (ni logit scale) explained
by paternal line are examined within each maternal type separately. In each case we also
provide separate estimates of variance for the inbred lines derived from Montpellier and
Lyon populations.

Data	Fixed effects			Random effects	
	maternal type (DS, KF, KS)	paternal origin (Lyon, Montpellier)	interaction	maternal ID	paternal line
all	$\chi^2_{df2} = 80.61$ $p < 10^{-15}$	$\chi^2_{df1} = 9.73$ $p = 0.002$	$\chi^2_{df2} =$ 5.37 $p = 0.068$	$\chi^2_{df1} = 7.16$ v = 0.92 $p = 0.0037$	$\chi^2_{df5} = 16.75$ $p = 0.005$ [1]
only DS		$\chi^2_{df1} = 1.439$ $p = 0.203$		$\chi^2_{df1} = 0.18$ v = 0.29 $p = 0.387$	$\chi^2_{df1} = 6.58$, v = 0.80 $p = 0.005$ (in Montpellier: v= 0.77, in Lyon : v=0.79)
only KF		$\chi^2_{df1} = 0.28$, $p = 0.597$		$\chi^2_{df1} = 0.89$ v = 1.32 $p = 0.174$	$\chi^2_{df1} = 0.00$, v = 0.00, $p = 0.5$ (in Montpellier: v = 0.00, in Lyon : v = 0.00)
only KS		$\chi^2_{df1} = 14.84$, $p = 0.0001$		$\chi^2_{df1} = 6.54$, v = 1.00 , $p = 0.005$	$\chi^2_{df1} = 32.4$ v = 1.58 , $p = 6.2\ 10^{-9}$ (in Montpellier: v = 1.45, in Lyon : v= 1.97)

Figure 4: Proportions of male-fertile and male-sterile phenotypes as a function of the origin of the paternal inbred line (Lyon or Montpellier) and the maternal type (DS, KF, KS).

Figure 5: Proportions of male-fertile individuals in offspring sired by various inbred lines from Montpellier (red) and Lyon (blue), in a DS maternal context (left) or KS maternal context (middle). The estimates are extracted from binomial GLMMs (BLUPs), de-logited to put them back into natural scale, and presented with 95% confidence intervals. The lines are arranged by increasing male-fertility, and separated by origin in the KS context, as the population of origin has a significant effect in this context (in DS context, this is not the case). Horizontal lines provide mean proportions of male-fertile individuals estimated by

the model, over all paternal lines (DS context) or separately for each population of origin

(KS context). Note that there are more paternal lines tested in KS than in DS context. The

performance of lines tested in the two contexts are plotted together in the right panel (each

dot is a paternal inbred line). The correlation is not significant (Spearman's ρ = 0.22, *p* =

0.184).

Discussion

This experiment was designed to characterize the restoration in *P. acuta* using two different

nuclear backgrounds (i) a naive background, originated from Montpellier, which has never

been in contact with N individuals at least during the previous 90 generations, (ii) a non-

naive background, originated from Lyon, which has been in contact with the K and the D

mitotypes in its recent history. Our results show that the two nuclear backgrounds had

different genetic determination and that restoration, when present, was mitotype-specific, as

in most plant examples.

Male fertility is restored by two different systems in Physa acuta

In the Montpellier background the K mitotype induced male sterility with an average

penetrance close to 70%. Results from the selection experiment point to a significant but

limited heritability of this penetrance, as (i) within the KS stock (Montpellier background),

male-fertile mothers produced slightly more male-fertile daughters than male-sterile

mothers, regardless of the father, and (i) fathers from the two selected populations, both

derived from the same background after two generations, also differed in their restoration

potential. However the latter difference was counter prediction as the lines selected to have

a low restoration frequency (LFR) sired more male-fertile offspring than the lines assumed

to have a high restoration frequency (HFR). This means that selection on the LFR and the

HFR lines was ineffective, and has probably been overwhelmed by drift. This outcome is

unexpected if restoration depends on fully-penetrant, dominant alleles, in which case large

differences among LFR and HFR, in the direction of selection, would be expected (see

supplement S2). However, if genetic variation is not large, and is expressed as a quantitative

degree of penetrance (rather than binary 100% sterile or 100% fertile), the male-fertility

phenotype of a few offspring becomes an imprecise indicator of the breeding value of each

individual, lowering the efficiency of our selection process relative to drift, especially with

the low numbers of individuals that passed the selection criteria. The distribution of male

fertility in offspring of inbred lines derived from the Montpellier population strengthens this

view, as overall, we have no convincing indication that any of the lines produced more than

50% male-fertile daughters, and the distribution seemed continuous between 0 and 50%,

though mostly concentrated near 0.

From these results, we concluded that male-fertility of the K mitotype in the

Montpellier nuclear background was expressed as a quantitative trait with a small amount of

genetic variance that can be transmitted both by fathers and mothers, and some nongenetic

variance (i. e. the same genotype can be male-fertile or male-sterile, with a genetically

determined proportion, as in the threshold trait model Ehlers et al., 2005). In addition, we

did observe cases of partial male-sterility ("semi-steriles") although their low number did

not allow a separate analysis, suggesting that binary classification is a simplification of true

phenotypic distributions. Indeed, in case of polygenic determination of restoration,

individuals carrying CMS may rarely be fully restored and male fitness may vary

quantitatively among restored CMS hermaphrodites (e.g. Dufay et al., 2008). This result is

similar to previous studies on non-crop species that rejected Mendelian models of restoration

(e.g. *Thymus vulgaris,* Belhassen et al., 1991; Charlesworth & Laporte, 1998; *Lobelia*

siphilitica, Dudle et al., 2001). Ehlers et al. (2005) suggested that scenarios considering

restoration of male fertility as a threshold trait may fit these data better than multilocus

Mendelian models while producing overall similar expectations as single-allele evolutionary

models regarding sex-ratio dynamics.

Results obtained from the crosses with KF mothers (i.e. with a background from

Lyon) suggested that male-fertility of the K-mitotype has a very different genetic basis in

the original Lyon population (a population where K naturally occurs) compared to the

laboratory Montpellier population. The results are consistent with one or a few dominant

restorer alleles acting on the K mitotype fixed or segregating at very high frequency in the

Lyon nuclear background. Indeed, mothers from the KF population were nearly all (>90%)

male-fertile and their daughters too, even when their father came from the Montpellier

population (HFR, LFR or Montpellier inbred lines) where there is no or rare restoration (as

discussed above), as expected from transmission of dominant restorers by mothers.

Reciprocally, mothers from Montpellier, although predominantly male-sterile themselves,

produced high frequencies of male-fertile daughters when fertilized by fathers from inbred

lines derived from the Lyon population. It is interesting to note however, that if dominant

restorers were fixed in Lyon, we would expect all these inbred lines to restore fertility with

high efficiency ; this is the case of only three of the 9 lines (75 to 100% fertility) , the rest

being distributed in the 25%-75% range, with an overall estimated mean of 62%. The

contrast in male-fertility frequencies in Montpellier-Lyon crosses, depending on whether the father or the mother comes from the Lyon population, is likely to reflect the recent history of the Lyon stock in the laboratory. The KF mothers come from a 100% K population maintained autonomously for 20 generations, during which all sperm donors must have been of the K-mitotype. Therefore the frequency of any restorer of male-fertility was positively selected, and maintained high, or perhaps increased, compared to the natural population. On the other hand, fathers from inbred lines from Lyon were derived from a population that, although founded from the same natural population, was maintained in parallel with 100% N mitotype. In these conditions, restorers were silent (in the sense that all individuals were male-fertile anyway) and if they were costly, may have decreased in frequency. This hypothesis of a divergent selection on restorers depending on mitotype frequencies will however have to be checked by looking at more replicates than just one population of each type. Interestingly, the inbred lines from Lyon do not show a strikingly bimodal distribution, suggesting that several loci with cumulative effects may be involved in restoration.

Our results highlighted two different restoration systems in *P. acuta*. In plants, restorer alleles with different effects have been reported. For example, in *Plantago* genus, multiple independent restoration systems were identified with both dominant and recessive effects sometimes involving more than one locus (Koelewijn & Van Damme, 1995; E. van Damme, 1984). In *Lobelia siphilitica* one CMS gene is restored by a single dominant allele, while restoration of other CMS is controlled by several loci and/or epistatic effects (Dudle et al. 2001). In these examples, one restoration system corresponds to one CMS type, however in *P. acuta* we found that two restoration systems acting together on the same CMS type K.

The two restoration systems are specific of the K mitotype

The D-mitotype individuals were on average 90 – 100% male-sterile irrespective of the nuclear background. The low proportion of male-fertile individuals was consistent with the previous study of David et al. (2022) which found 8% of male-fertile D individuals in the original population. Here, we found no evidence for a dominant restorer with large penetrance; however we observed the existence of genetic variation for penetrance among inbred lines from both Montpellier and Lyon, without evidence for a difference between the two origins (we would expect a higher rate of male-fertility in Lyon if this variation has been specifically maintained by selection for restoring male fertility, as the Montpellier population had no recent contact with the D mitotype). This lack of difference between

Montpellier and Lyon, and the fact that the lines with the highest male-fertility rate in the

context of the D mitotype were not necessarily the ones that efficiently restore fertility in the

K-mitotype, suggest that the two types of variation act in a mitotype-specific manner. The

specificity of the restorers has been also observed in plants where different CMS genes can

coexist within a population each with its own specific nuclear restorer alleles (e.g.

van Damme & van Delden, 1982; Manicacci et al., 1997).

In the wild the D mitotype seems rare as it was found only in one population and at

low frequency (14%, David et al. 2022). Thus, suppression of the male sterility may not be

frequent enough for a restorer to be selected. The high evolutionary rate detected in the D

mitotype (David et al. 2022) could result from an arms race with a restorer gene which is

now extinct or rare, or has lost its ability to restore fertility following a recent evolution of

the D mitochondrial genome itself. According to the frequency-dependent model, as soon as

a CMS gene is present in a population, there is a strong selection for any nuclear mutation

that counteracts the phenotypic effects of CMS. Restorers should spread rapidly and if they

have no or little fitness cost, the polymorphism is transient. However, nuclear restoration is

expected to be rare under the hypothesis of a high cost of restoration (Gouyon et al., 1991;

Dufay et al., 2007). Founder effects causing a reduction of the nuclear diversity could also

lead to the local absence of certain nuclear restorers in a recent founded population (e.g.

Manicacci et al. 1996; Laporte et al., 2001). Therefore, we can hypothesize that the evolution

of restorers specific to the D mitotype might be constrained by a high cost or founder effects.

Two stages in the maintenance of cyto-nuclear polymorphism

Most models have studied the maintenance of the cytoplasmic and nuclear genetic variation

in gynodioecious populations assuming strong and dominant restorer alleles at a few loci.

Such alleles are indeed the most likely to be under strong selection and undergo selective

sweeps in populations invaded by new CMS genes. Depending on their associated costs,

they may either reach fixation ("hard-sweep") or stabilize at intermediate or cyclically

variable frequencies ("soft-sweep") together with their corresponding CMS cytotypes

(Gouyon et al., 1991; Dufaÿ et al., 2007). Gynodioecious populations observed today are

mostly in the polymorphic state and the restorers found indeed include large-effect alleles

with dominant action (e.g. de Haan et al., 1997; Dufay et al., 2008, 2009). In addition this

action is usually specific to each cytotype, revealing either a arms-race coevolutionary

history specific to each cytotype-restorer pair, and/or the fact that each new CMS cytotype

has a mode of action that differs so strongly from previous ones that existing restorers are

unlikely to neutralize it. In *Physa acuta,* the K mitotype, and the restorers present in the natural population where it was found, are likely to be in this situation, as we found restoration to be strong (though possibly distributed into several loci), dominant (same effect in the heterozygous state in daughters, than in homozygous state in mothers) and specific (no cross-mitotype effect).

However, while such situations are expected to be frequent in stable gynodioecious populations, they are probably not representative of initial stages when a new CMS mutant appears in a "naïve" population or species, where no selection has ever taken place on nuclear genes to suppress its effect. According to Lewis (1941), the spread of CMS in the population will in this case become limited only by pollen limitation when females become so abundant that many of them lose their ovules by lack of pollen to fertilize them. Our results suggest however a way by which such populations may lower the impacts of pollen (or, in the case of *Physa*, sperm) limitation: CMS may not be fully penetrant, leaving a residual production of male gametes that is at least sufficient to fertilize the eggs of mates that have not had previous contacts with male-fertile congeners. It is likely that even in a naïve population, pre-existing quantitative genetic variation includes variants that act on this penetrance. This probably corresponds to the situation of the K-mitotype when introgressed into the "naïve" Montpellier population, and of the D mitotype in both the Montpellier and Lyon populations. Here, we mean by "naïve" that it has either never had any history of selection in presence of the K and D mitotypes, or that the contact is too recent for large-effects restorers to have emerged, or too ancient for them to have persisted if they ever existed. The background variation acting on incomplete penetrance of CMS could produce a reservoir of variation within the population, by which male fertility can start to increase, until the appearance of complete penetrant restorers.

The emergence of restoration could therefore often take place in two stages: a first stage with predominantly quantitative, weak modifiers of CMS penetrance, and a second stage when dominant, strong restorer alleles appear and are selected at one or several loci, ending up with either fixation or a mixed determinism of male-fertility. Although theoretical models generally consider a simple genetic determination for restoration (but see Frank, 1989; Bailey & Delph, 2007), empirical studies on wild plants have often rejected this assumption as mentioned above (e.g. Charlesworth & Laporte, 1998; Koelewijn, 2003). A complex determination of restoration implies that within a population, CMS individuals may be partially restored (i. e. male fertility may be more of a continuous than binary 0/1

distribution) and this could affect the degree of selection for restorer alleles. Then, the
positive selection of restorers could slow down and ultimately modify the conditions of
maintenance of cytonuclear polymorphism, thereby favouring the maintenance of females
in gynodioecious populations (Bailey & Delph, 2007).

Conclusion

This study shows that as in plants, the genetic determination of male-sterility in *Physa acuta*
is complex and suggests that it encompasses two different layers: one of weak, quantitative
modifiers of the penetrance, not necessarily coevolved with CMS, the other of strong-effect
restorer alleles with dominant action, that were selected in the presence of CMS. Moreover,
the way the phenotype is measured could influence our perception of the restoration
potential. Indeed, the male fertility status was evaluated in the absence of competition as
individuals were always paired with virgin mates; the restoration of the male fertility might
be considered less efficient when measured in a population with competition for the male
function. Thus, it will be interesting to evaluate male fitness in competition, e.g. when
partners are not virgin, and have been first paired with a male-fertile partner before being
paired with focal individuals. The genetic determinism of the restoration may influence the
maintenance of the cyto-nuclear polymorphism by acting on the strength of the restoration
cost. This could be evaluated by looking at the correlations between the penetrance of the
nuclear backgrounds and the cost of restoration on male and/or female fitness. However,
studies performed in plants suggest that costs are difficult to detect (but see de Haan et al.,
1997; Bailey, 2002; Dufay et al., 2008). Finally, because our results suggest that the
restoration of the male function is a quantitative trait, identification of QTLs for male-
fertility restoration would be necessary.

References

Bailey, M. F. (2002). A Cost of Restoration of Male Fertility in a Gynodioecious Species, Lobelia Siphilitica. *Evolution*, *56*(11), 2178-2186. https://doi.org/10.1111/j.0014-3820.2002.tb00142.x

Bailey, M. F., & Delph, L. F. (2007). Sex-Ratio Evolution in Nuclear-Cytoplasmic Gynodioecy When Restoration Is a Threshold Trait. *Genetics*, *176*(4), 2465-2476. https://doi.org/10.1534/genetics.107.076554

Bailey, M. F., Delph, L. F., & Lively, C. M. (2003). Modeling Gynodioecy : Novel Scenarios for Maintaining Polymorphism. *The American Naturalist*, *161*(5), 762-776. https://doi.org/10.1086/374803

Bates, D., Kliegl, R., Vasishth, S., & Baayen, H. (2015). *Parsimonious Mixed Models* (arXiv:1506.04967). arXiv. http://arxiv.org/abs/1506.04967

Belhassen, E., Dommée, B., Atlan, A., Gouyon, P. H., Pomente, D., Assouad, M. W., & Couvet, D. (1991). Complex determination of male sterility in Thymus vulgaris L. : Genetic and molecular analysis. *Theoretical and Applied Genetics*, *82*(2), 137-143. https://doi.org/10.1007/BF00226204

Bergero, R., Levsen, N., Wolff, K., & Charlesworth, D. (2019). Arms races with mitochondrial genome soft sweeps in a gynodioecious plant, Plantago lanceolata. *Molecular Ecology*, *28*(11), 2772-2785. https://doi.org/10.1111/mec.15121

Burt, A., & Trivers, R. (2006). *Genes in conflict* (Belknap Press of Harvard University Press, Boston, MA.).

Caruso, C. M., Eisen, K., & Case, A. L. (2016). An Angiosperm-Wide Analysis of the Correlates of Gynodioecy. *International Journal of Plant Sciences*, *177*(2), 115-121. https://doi.org/10.1086/684260

Charlesworth, D., & Ganders, F. R. (1979). The population genetics of gynodioecy with cytoplasmic-genic male-sterility. *Heredity*, *43*(2), 213-218. https://doi.org/10.1038/hdy.1979.76

Charlesworth, D., & Laporte, V. (1998). Male-Sterility Polymorphism of Silene vulgaris : Analysis of Genetic Data From Two Populations and Comparison With Thymus vulgaris | Genetics | Oxford Academic. *The Genetics Society of America*. https://academic.oup.com/genetics/article/150/3/1267/6034628?login=false

Chen, L., & Liu, Y.-G. (2014). Male Sterility and Fertility Restoration in Crops. *Annual Review of Plant Biology*, *65*(Volume 65, 2014), 579-606. https://doi.org/10.1146/annurev-arplant-050213-040119

Cosmides, L., & Tooby, J. (1981). Cytoplasmic inheritance and intragenomic conflict.
Journal of Theoretical Biology, *89*(1), 83-129. https://doi.org/10.1016/0022-
5193(81)90181-8

Couvet, D., Ronce, O., & Gliddon, C. (1998). The Maintenance of Nucleocytoplasmic
Polymorphism in a Metapopulation : The Case of Gynodioecy. *The American*
Naturalist, *152*(1), 59-70. https://doi.org/10.1086/286149

Cui, X., Wise, R. P., & Schnable, P. S. (1996). The *rf2* Nuclear Restorer Gene of Male-
Sterile T-Cytoplasm Maize. *Science*, *272*(5266), 1334-1336.
https://doi.org/10.1126/science.272.5266.1334

David, P., Degletagne, C., Saclier, N., Jennan, A., Jarne, P., Plénet, S., Konecny, L.,
François, C., Guéguen, L., Garcia, N., Lefébure, T., & Luquet, E. (2022). Extreme
mitochondrial DNA divergence underlies genetic conflict over sex determination.
Current Biology, *32*(10), 2325-2333.e6. https://doi.org/10.1016/j.cub.2022.04.014

de Haan, A. A., Koelewijn, H. P., Hundscheid, M. P. J., & Van Damme, J. M. M. (1997).
The Dynamics of Gynodioecy in Plantago lanceolatu L. II. Mode of Action and
Frequencies of Restorer Alleles. *Genetics*, *147*(3), 1317-1328.
https://doi.org/10.1093/genetics/147.3.1317

de Haan, A., Hundscheid, M., & van Hinsberg, A. (1997). Effects of CMS types and
restorer alleles on plant performance in Plantago lanceolata L. : An indication for
cost of restoration—De Haan—1997—Journal of Evolutionary Biology—Wiley
Online Library. *Journal of Evolutionary Biology*, *10*, 803-820.
https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1420-9101.1997.10050803.x

Delannay, X., Gouyon, P. H., & Valdeyron, G. (1981). MATHEMATICAL STUDY OF
THE EVOLUTION OF GYNODIOECY WITH CYTOPLASMIC INHERITANCE
UNDER THE EFFECT OF A NUCLEAR RESTORER GENE. *Genetics*, *99*(1),
169B - 181. https://doi.org/10.1093/genetics/99.1.169a

Delph, L. F., Touzet, P., & Bailey, M. F. (2007). Merging theory and mechanism in studies
of gynodioecy. *Trends in Ecology & Evolution*, *22*(1), 17-24.
https://doi.org/10.1016/j.tree.2006.09.013

Dewey, R. E., Timothy, D. H., & Levings, C. S. (1987). A mitochondrial protein
associated with cytoplasmic male sterility in the T cytoplasm of maize.
Proceedings of the National Academy of Sciences, *84*(15), 5374-5378.
https://doi.org/10.1073/pnas.84.15.5374

Dill, C. L., Wise, R. P., & Schnable, P. S. (1997). Rf8 and Rf* Mediate Unique T-urf13-
Transcript Accumulation, Revealing a Conserved Motif Associated With RNA
Processing and Restoration of Pollen Fertility in T-Cytoplasm Maize. *Genetics*,
147(3), 1367-1379. https://doi.org/10.1093/genetics/147.3.1367

Dudle, D. A., Mutikainen, P., & Delph, L. F. (2001). Genetics of sex determination in the
gynodioecious species Lobelia siphilitica : Evidence from two populations.
Heredity, *86*(3), 265-276. https://doi.org/10.1046/j.1365-2540.2001.00833.x

Dufay, M., Champelovier, P., Käfer, J., Henry, J. P., Mousset, S., & Marais, G. A. B.
(2014). An angiosperm-wide analysis of the gynodioecy–dioecy pathway. *Annals*
of Botany, *114*(3), 539-548. https://doi.org/10.1093/aob/mcu134

Dufay, M., Cuguen, J., Arnaud, J.-F., & Touzet, P. (2009). Sex Ratio Variation Among
Gynodioecious Populations of Sea Beet : Can It Be Explained by Negative
Frequency-Dependent Selection? *Evolution*, *63*(6), 1483-1497.
https://doi.org/10.1111/j.1558-5646.2009.00653.x

Dufaÿ, M., Touzet, P., Maurice, S., & Cuguen, J. (2007). Modelling the maintenance of
male-fertile cytoplasm in a gynodioecious population. *Heredity*, *99*(3), Article 3.
https://doi.org/10.1038/sj.hdy.6801009

Dufay, M., Vaudey, V., De Cauwer, I., Touzet, P., Cuguen, J., & Arnaud, J. -F. (2008).
Variation in pollen production and pollen viability in natural populations of
gynodioecious Beta vulgaris ssp. maritima : Evidence for a cost of restoration of
male function? *Journal of Evolutionary Biology*, *21*(1), 202-212.
https://doi.org/10.1111/j.1420-9101.2007.01454.x

Ehlers, B. K., Maurice, S., & Bataillon, T. (2005). Sex inheritance in gynodioecious
species : A polygenic view. *Proceedings of the Royal Society B: Biological*
Sciences, *272*(1574), 1795-1802. https://doi.org/10.1098/rspb.2005.3168

Frank, S. A. (1989). The Evolutionary Dynamics of Cytoplasmic Male Sterility. *The*
American Naturalist, *133*(3), 345-376. https://doi.org/10.1086/284923

Gouyon, P.-H., Vichot, F., & Damme, J. M. M. V. (1991). Nuclear-Cytoplasmic Male
Sterility : Single-Point Equilibria Versus Limit Cycles. *The American Naturalist*,
137(4), 498-514. http://www.jstor.org/stable/2462377

Huang, W., Hu, J., Yu, C., Huang, Q., Wan, L., Wang, L., Qin, X., Ji, Y., Zhu, R., Li, S., &
Zhu, Y. (2012). Two non-allelic nuclear genes restore fertility in a gametophytic
pattern and enhance abiotic stress tolerance in the hybrid rice plant. *Theoretical and*
Applied Genetics, *124*(5), 799-807. https://doi.org/10.1007/s00122-011-1755-9

Huang, W., Yu, C., Hu, J., Wang, L., Dan, Z., Zhou, W., He, C., Zeng, Y., Yao, G., Qi, J.,

Zhang, Z., Zhu, R., Chen, X., & Zhu, Y. (2015). Pentatricopeptide-repeat family

protein RF6 functions with hexokinase 6 to rescue rice cytoplasmic male sterility.

Proceedings of the National Academy of Sciences, *112*(48), 14984-14989.

https://doi.org/10.1073/pnas.1511748112

Kim, Y.-J., & Zhang, D. (2018). Molecular Control of Male Fertility for Crop Hybrid

Breeding. *Trends in Plant Science*, *23*(1), 53-65.

https://doi.org/10.1016/j.tplants.2017.10.001

Koelewijn, H. P. (2003). Variation in restorer genes and primary sexual investment in

gynodioecious Plantago coronopus : The trade-off between male and female

function. *Proceedings of the Royal Society of London. Series B: Biological*

Sciences, *270*(1527), 1939-1945. https://doi.org/10.1098/rspb.2003.2462

Koelewijn, H. P., & Van Damme, J. M. (1995). Genetics of male sterility in gynodioecious

Plantago coronopus. II. Nuclear genetic variation. *Genetics*, *139*(4), 1759-1775.

https://doi.org/10.1093/genetics/139.4.1759

Laporte, V., Viard, F., Bena, G., Valero, M., & Cuguen, J. (2001). The Spatial Structure of

Sexual and Cytonuclear Polymorphism in the Gynodioecious Beta vulgaris ssp.

maritima : I/ at a Local Scale. *Genetics*, *157*(4), 1699-1710.

https://doi.org/10.1093/genetics/157.4.1699

Laser, K. D., & Lersten, N. R. (1972). Anatomy and cytology of microsporogenesis in

cytoplasmic male sterile angiosperms. *The Botanical Review*, *38*(3), 425-454.

https://doi.org/10.1007/BF02860010

Laugier, F., Saclier, N., Béthune, K., Braun, A., Konecny, L., Lefébure, T., Luquet, E.,

Plénet, S., Romiguier, J., & David, P. (2024). Both nuclear and cytoplasmic

polymorphisms are involved in genetic conflicts over male fertility in the

gynodioecious snail Physa acuta. *Evolution*, qpae053.

https://doi.org/10.1093/evolut/qpae053

Lewis, D. (1941). Male Sterility in Natural Populations of Hermaphrodite Plants the

Equilibrium Between Females and Hermaphrodites to Be Expected with Different

Types of Inheritance. *New Phytologist*, *40*(1), 56-63.

https://doi.org/10.1111/j.1469-8137.1941.tb07028.x

Manicacci, D., Atlan, A., & Couvet, D. (1997). Spatial structure of nuclear factors

involved in sex determination in the gynodioecious Thymus vulgaris L. *Journal of*

Evolutionary Biology, *10*(6), 889-907. https://doi.org/10.1111/j.1420-

9101.1997.tb00004.x

Matsuhira, H., Kagami, H., Kurata, M., Kitazaki, K., Matsunaga, M., Hamaguchi, Y.,

Hagihara, E., Ueda, M., Harada, M., Muramatsu, A., Yui-Kurino, R., Taguchi, K.,

Tamagake, H., Mikami, T., & Kubo, T. (2012). Unusual and Typical Features of a

Novel Restorer-of-Fertility Gene of Sugar Beet (Beta vulgaris L.). *Genetics*,

192(4), 1347-1358. https://doi.org/10.1534/genetics.112.145409

Mccauley, D. E., & Olson, M. S. (2003). Associations among cytoplasmic molecular

markers, gender, and components of fitness in Silene vulgaris, a gynodioecious

plant. *Molecular Ecology*, *12*(3), 777-787. https://doi.org/10.1046/j.1365-

294X.2003.01764.x

Noël, E., Chemtob, Y., Janicke, T., Sarda, V., Pélissié, B., Jarne, P., & David, P. (2016).

Reduced mate availability leads to evolution of self-fertilization and purging of

inbreeding depression in a hermaphrodite. *Evolution*, *70*(3), 625-640.

https://doi.org/10.1111/evo.12886

Ross, M. D., & Gregorius, H. R. (1985). SELECTION WITH GENE-CYTOPLASM

INTERACTIONS. II. MAINTENANCE OF GYNODIOECY. *Genetics*, *109*(2),

427-439. https://doi.org/10.1093/genetics/109.2.427

Saumitou-Laprade, P., Cuguen, J., & Vernet, P. (1994). Cytoplasmic male sterility in

plants : Molecular evidence and the nucleocytoplasmic conflict. *Trends in Ecology*

& Evolution, *9*(11), 431-435. https://doi.org/10.1016/0169-5347(94)90126-0

Sloan, D. B., Alverson, A. J., Wu, M., Palmer, J. D., & Taylor, D. R. (2012). Recent

Acceleration of Plastid Sequence and Structural Evolution Coincides with Extreme

Mitochondrial Divergence in the Angiosperm Genus Silene. *Genome Biology and*

Evolution, *4*(3), 294-306. https://doi.org/10.1093/gbe/evs006

Touzet, P., Hueber, N., Bürkholz, A., Barnes, S., & Cuguen, J. (2004). Genetic analysis of

male fertility restoration in wild cytoplasmic male sterility G of beet. *Theoretical*

and Applied Genetics, *109*(1), 240-247. https://doi.org/10.1007/s00122-004-1627-7

van Damme, E. (1984). A relation between perfect equilibria in extensive form games and

proper equilibria in normal form games. *International Journal of Game Theory*,

13(1), 1-13. https://doi.org/10.1007/BF01769861

van Damme, J. M. M., Hundscheid, M. P. J., Ivanovic, S., & Koelewijn, H. P. (2004).

Multiple CMS–restorer gene polymorphism in gynodioecious Plantago coronopus.

Heredity, *93*(2), 175-181. https://doi.org/10.1038/sj.hdy.6800490

van Damme, J. M. M., & van Delden, W. (1982). Gynodioecy in Plantago lanceolata L I.

polymorphism for plasmon type. *Heredity*, *49*(3), 303-318.

https://doi.org/10.1038/hdy.1982.104

Vellnow, N., Vizoso, D. B., Viktorin, G., & Schärer, L. (2017). No evidence for strong

cytonuclear conflict over sex allocation in a simultaneously hermaphroditic

flatworm. *BMC Evolutionary Biology*, *17*(1), 103. https://doi.org/10.1186/s12862-

017-0952-9

Werren, J. H., & Beukeboom, L. W. (1998). SEX DETERMINATION, SEX RATIOS,

AND GENETIC CONFLICT. *Annual Review of Ecology, Evolution and*

Systematics, *29*(Volume 29, 1998), 233-261.

https://doi.org/10.1146/annurev.ecolsys.29.1.233

Yamamoto, M. P., Shinada, H., Onodera, Y., Komaki, C., Mikami, T., & Kubo, T. (2008).

A male sterility-associated mitochondrial protein in wild beets causes pollen

disruption in transgenic plants. *The Plant Journal*, *54*(6), 1027-1036.

https://doi.org/10.1111/j.1365-313X.2008.03473.x