Development of an immunoassay for the detection and diagnosis of microbially-influenced corrosion caused by methanogenic Archaea

Sven Lahme and Jorge Mantilla Aguas

ExxonMobil Technology and Engineering Company, Spring, Texas, USA

Abstract

Microbially-influenced corrosion (MIC) is a costly problem across several industries. The steadily rising use of advanced molecular biological methods to investigate MIC allowed ever deeper insights in the underlying microbial community structure and function. However, currently available technologies do not allow accurate detection and diagnosis of MIC in the field. Recent discovery of a special hydrogenase in certain corrosive methanogenic Archaea allowed for the development of a first ever MIC biomarker termed here *micH*. The *micH* gene encodes the large subunit of a special [NiFe] hydrogenase involved in MIC. Here we describe the development of a recombinant antibody that enable the specific and sensitive detection of the MicH protein in western blot immunoassays. Using a recombinant MicH protein we determined the lower limit of detection per assay to be around 0.3 fg MicH. The immunoassay was able to detect a strong signal for the MicH protein in *micH*-positive pure cultures of *Methanobacterium*-like. strain IM1 that was cultivated on iron granules, and the signal was over 500 times lower in a *micH*-negative *Methanococcus maripaludis* S2 culture. To further evaluate the ability to differentiate corrosive from non-corrosive microbial communities, we tested ten oil field enrichment cultures that showed a wide range of corrosion rates (0.02 – 0.48 mm/yr). We detected the MicH protein in planktonic (36.5 – 1473.5 pg/mL) and carbon steel biofilm samples (41.0 – 7971.3 pg/cm^2) from corrosive methanogenic enrichments (0.17 – 0.48 mm/yr) but did not detect MicH in any of the non-

corrosive tests (<0.08 mm/yr) despite methanogenic activity. The results indicate that corrosion

was likely caused by methanogenic Archaea expressing a corrosive [NiFe] hydrogenase and that

the newly developed MicH-specific immunoassay can detect and monitor their activity. The

development of a specific and sensitive immunoassays to detect a MIC biomarker allow corrosion

scientists and field practitioners to detect and monitor the activity of corrosive methanogenic

Archaea.

Introduction

Microbially-influenced corrosion (MIC) negatively affects steel infrastructures and causes

significant costs across multiple industries ever year. In anaerobic systems, such as pipelines used

for the production and transportation of mixtures of oil, water and gas, MIC is often linked to the

presences and activity of sulfate-reducing bacteria (SRB) and methanogenic Archaea[1–5].

The monitoring of corrosive biofilms in industrial settings often involves the quantification of cell

numbers by culture-based or molecular biological methods such as the quantification of adenosine

triphosphate (ATP) or universal and functional genes by quantitative PCR (qPCR)[6–9]. Despite the

introduction of the latter into industrial operations, reliable detection of MIC during operation is

still challenged. Firstly, not all microorganisms are causing industrially relevant corrosion and

secondly, even members of the same metabolic groups (e.g. SRB or methanogenic Archaea) have

been shown to have vastly different corrosion rates[1,10]. This suggest likely genetic differences that

enable some microorganisms to cause higher rates of corrosion than others.

A recent study identified a specific gene islands in genomes of corrosive methanogenic Archaea

that are absent in non-corrosive close relative strains. The gene islands encode among other genes

a unique and phylogenetically highly conserved bidirectional [NiFe] hydrogenase (termed 'MIC

hydrogenase'), a Tat system for protein translocation as well as enzymes with potential

involvement in protein glycosylation[5,11–13]. The 'MIC hydrogenase' was only detected in corrosive

pure cultures and corrosive oil field enrichment cultures but not in the ones showing low corrosion,

explaining the corrosive character of some methanogenic Archaea.

We recently proposed to use the large subunit of the 'MIC hydrogenase' as biomarker (termed

micH) to detect and monitor certain corrosive methanogenic Archaea in the field[5]. By utilizing

specific qPCR assays, the MIC biomarker showed excellent correlation with MIC in both

laboratory tests and field applications[5,14]. Here, we discuss the development of a MicH-specific

recombinant antibody to detect and monitor corrosive methanogenic Archaea in laboratory tests

and in the field. In addition, protein detection may reduce potential false positive that could be

associated with qPCR-based detection. A recombinant antibody able to detect MIC protein

biomarker would also allow the development of a field-friendly rapid test assays (e.g. lateral flow

tests) to support MIC monitoring and detection in the field.

Materials and Methods

Growth of pure cultures and enrichments

Actively growing cultures of *Methanococcus maripaludis* strain S2 (= DSM 14266), *Desulfovibrio*

ferrophilus strain IS5 (= DSM 15579) and *Oleidesulfovibrio alaskensis* strain G20 (= DSM 17464)

were acquired from the German strain collection DSMZ. Culture aliquots of *Methanobacterium-*

like strain IM1 were donated by Dr Florin Musat (Helmholtz Centre for Environmental Research,

Leipzig, Germany). Pure cultures and enrichments in this study were routinely grown in the

presence of 1 − 2 mm iron granules (40 µg/ml medium; Thermo Fisher Scientific, Waltham,

Massachusetts, US), 0.5 mM sodium acetate, 0.5 mM sodium sulfide and 10 mM sodium sulfate

in anaerobic synthetic brine that was modelled after an oil field produced water as previously

described[5].

For establishing field enrichment cultures 100 ml Duran® glass bottles were filled with 2 g iron

granules and dry heat sterilized at 180° C for 1 h. After cooling down the bottle headspace was

exchanged with N_2/CO_2 (79:21) and sealed with sterile butyl rubber stoppers and screw cups. The

sterile anoxic glass bottles were inoculated with 50 ml of produced water on site. Production fluids

(oil, gas and water) were collected in sterile polypropylene bottles and oil and water allowed to

separate by gravity. Once separated, the water was collected using sterile syringes with attached

sterilized polypropylene tubes. After collection the tube was carefully replace by a sterile needle

and the anoxic glass bottles inoculated. After arrival in the destination lab, the cultures were

incubated at 32° C for at least two weeks before used as inoculum for corrosion bottle tests.

Enrichment cultures were established this way from eight pipelines of two separate West African

oil field operations. In addition, two additional enrichment were established using pipeline solids

('pigging debris') collected as described previously[5] from a pipeline from an U.S. West Coast

operation and a U.S. onshore oil field operation during maintenance cleaning. Upon arrival in the

lab approximately 3 g solids were transfer into sterile bottle containing 50 ml of the previously

mentioned synthetic produced water[5] and closed with sterile butyl rubber stoppers under N_2/CO_2

(79:21) atmosphere.

Corrosion bottle tests

Corrosion tests were performed in one-liter Duran® glass bottles that were filled with 600 ml

anoxic synthetic brine to which three pre-cleaned and sterilized API 5L X52 steel coupons were

added in polyether ether ketone coupon holder as described earlier[5,15]. The bottles were sealed with

butyl rubber stoppers using sterile Hungate techniques and a N_2/CO_2 (79:21) headspace. Test were

inoculated with 1% of respective cultures and incubated on a rotary shaker at 75 rpm and 32°C for

eight weeks. Planktonic samples were collected for protein and DNA analysis using sterile syringes

and 0.2 μm polyethersulfone filter membranes or sterile swabs for biofilm collection as mentioned

previously[5]. Samples were immediately frozen at −80°C until further analysis. Removal of

corrosion products and weigh loss corrosion analysis of corrosion coupons was performed at the

end of the test as outlined earlier[5].

Recombinant antibody development

The recombinant antibody development workflow was conducted by Genscript Biotech

(Piscataway, NJ) and will be briefly outline below. For the initial development of polyclonal

antibodies, two peptide sequences were chosen from the protein sequence of the target gene *micH*

of the corrosive methanogen *M. maripaludis* strain OS7 (see supplementary material for details)

and conjugated to the metalloprotein keyhole limpet hemocyanin (KLH). Per peptide four rabbits

were immunized with three additional consecutive boosters. After four months, blood was

harvested from the rabbits and polyclonal antibodies partially purified for evaluation. Tests were

performed by indirect enzyme-linked immunosorbent assay (ELISA) using either of the two

peptides as target and via Western blotting as outlined below using protein extracts from

Methanobacterium-like strain IM1 as positive control[16] and protein extracts from *M. maripaludis*

strain S2 was used as MicH-negative control (Fig. S1)[11]. Animals with best performing polyclonal

antibodies were selected for cell fusion of spleen B cells with myeloma cell line. Screening and

selection of clones were performed via immunoassays as described above. The best performing

monoclonal antibody 'A11-1' was chosen for DNA sequencing (see supplementary material) and

recombinant antibody production by Genscript Biotech.

Protein extraction and detection via Western blot

Proteins were extracted from filters or swabs by placing them in sterile 2 ml screw cup

microcentrifuge tubes. 900 μl lysis buffer (10 mM HEPES, 1% Triton X-100, 5 mM EDTA, 100

mM DTT and 100mM Na_2CO_3, 1% SDS, 12% sucrose) were added to the filters or swabs and

placed in a heating block (105°C) for exactly 2 min. The tubes were removed and cooled to room

temperature and then vigorously mixed and centrifuged for 5 min at 21,100 g. 200 µl of the

supernatant was further processed to remove detergents using Pierce™ HiPPR Detergent Removal

Spin Column Kit (Thermo Fisher Scientific, Waltham, MA) according to manufacturer's protocol.

The detergent-free protein samples were further processed using Pierce™ Protein Concentrators

with a 10 kDa molecular weight cut-off polyether sulfone membrane (Thermo Fisher Scientific,

Waltham, MA). Protein samples were concentrated to an approximate volume of 25 µl and either

directly used for Wester blotting or stored at −80°C.

Western blotting was performed on a Protein Simple JESS Western blot platform (Protein Simple,

San Jose, CA) using the 12 − 230 kDa Separation Module and the Jess Anti-Rabbit Detection

Module. Protein analysis was performed according to the manufacturer protocol using a 1:20

antibody dilution (approx. 25 µg/ml) of A11-1 and 3 µl of protein sample. Protein Simple's Total

Protein module was used for normalization. For quantification a recombinant MicH protein was

purchased from Genscript Biotech (Piscataway, NJ) developed following their large-scale *E. coli*

expression protocol and His-tag purification.

DNA extraction and qPCR analysis

For DNA extraction cells were collected via filtration and filters transferred to 5 ml cryo vials

containing 1 ml DNA/RNA shield (Zymo Research, Irvine, CA) as preservative. Filters were

stored at −80°C until further analysis. DNA extraction and *micH* qPCR analysis were performed

by Microbial Insights Inc. (Knoxville, TN) as described previously[5].

Adenosine triphosphate analysis

ATP was quantified using the 2nd Generation ATP® test kit and PhotoMaster™ luminometer

(LuminUltra Technologies Ltd., Fredericton, Canada). For analysis of planktonic cells 20 ml of

medium was collected via filtration using the provided filters and syringes following the

manufacturers protocol. Biofilms were collected using a minimum of two sterile swabs per coupon

and the swab tips used for ATP quantification as outlined in the kit's manual.

Gas chromatography

Gas chromatographic analysis of methane was performed on an Agilent 490 Micro gas

chromatograph using a thermal conductivity detector. Separation was achieved on a heated Agilent

PoraPlot U column (50°C, 22.1 lb/in^2, and 10 m) in argon gas.

Results

Recombinant Anti-MicH antibody showed excellent affinity for recombinant MicH protein.

In order to optimize the signal to noise ratio (S/N) we performed first a dilution series with the

Anti-MicH antibody and a fixed concentration of 1 ng/ml of MicH$_{rec}$. A signal was detected

between approximately 50 – 60 kDa molecular weight that is in good agreement with the expected

molecular weight of the MicH protein containing the His-tag used for purification (54.5 kDa; Fig.

1A). Test performed with different antibody dilutions of Anti-MicH showed best results for S/N

ratio and low baseline with dilutions 1:50, 1:20 and 1:10 (Fig. 1B). For subsequent tests an

antibody dilution of 1:20 was chosen to minimize potential background noise while achieving an

overall strong signal.

In addition, a calibration curve of recombinant MicH protein showed excellent linear response

with concentrations between 20 pg/ml and 10 ng/ml (Fig. 2 and Fig. S2). A clearly detectable signal

with acceptable S/N ration (>30) was still detected at 20 pg/ml MicH$_{rec}$ therefore the assay was

able to reliably detect as little as 0.3 fg of recombinant MicH corresponding to approximately $3 \cdot 10^3$

protein molecules.

MicH-specific immunoassay showed excellent selectivity for corrosive methanogenic cultures

Unlike qPCR assays, immunoassays are less selective and may show potential background signal

due to cross-reactivity with non-target proteins or peptides in absence or low concentration of their

specific target[17]. To test for assay specificity and the ability to distinguish between corrosive and

non-corrosive methanogenic Archaea as well as non-methanogenic cultures, we tested the MicH-

specific immunoassay on protein extracts from methanogenic and sulfate-reducing pure cultures

(Fig. 3).

Protein extract of *Methanobacterium*-like strain IM1 obtained from cultures, that inflicted

significant corrosion of 0.3 mm/yr, showed a strong signal in the Western blot (Fig. 3 and Fig.

S3A). Furthermore, the *micH* gene was also detected by qPCR in DNA extracted from cultures of

strain IM1, confirming both the presence and expression of the 'MIC hydrogenase' during

corrosion. In contrast, the MicH-specific assay showed a more than 500x lower signals for protein

extracts from *M. maripaludis* strain S2 known to lack the 'MIC hydrogenase'[11]. In agreement,

strain S2 caused only negligible corrosion of 0.04 mm/yr (Fig. 3). Similar low signals were

observed in protein extracts from *Desulfovibrio ferrophilus* IS5 and *Desulfovibrio alaskensis* G20.

DNA analysis via qPCR confirmed absence of the *micH* gene in those cultures (Fig. 3). The results

confirm that the MicH-specific immunoassay selectively detects the MicH protein while showing

low cross-reactivity with proteins extracted from non-corrosive methanogenic strain S2 and SRB

pure cultures (Fig. 3 and Fig. S3A).

MicH-specific immunoassay allowed detection of MicH in corrosive methanogenic field cultures

Next, we wanted to better understand the performance of the MicH-specific immunoassay and

selectivity of the Anti-MicH antibody in more complex context of oil field produced water

enrichment cultures. Therefore, ten field enrichment cultures that showed varying degree of MIC

activity were compared in terms of corrosion as well as presence of the *micH* gene and expression

of the MicH protein (Fig. 4). Five of the ten cultures were classified as '+MIC' showing averaged

corrosion rates between $0.48 - 0.17$ mm/yr, which according to the Association for Materials

Protection and Performance (AMPP) is considered high to severe (AMPP-SP0775-2013). The

other five cultures were classified as '-MIC' showing rates between $0.02 - 0.08$ mm/yr, considered

as low to moderate corrosion according to AMPP (Fig. 4A).

Of the ten cultures eight showed detectable methanogenic activity having formed between 0.1 and

1.0 mmol of methane at the end of the test (Fig. 4A). ATP results indicated that biofilms have

formed on steel surfaces to a varying degree with generally higher numbers detected on corroded

steel surfaces (+MIC; $278 - 71299$ pg/cm^2) compared to less corroded ones (-MIC; $6 - 1762$

pg/cm^2; Fig. 4B). The *micH* gene was detected in planktonic cells as well as all five biofilms

collected from corroded steel coupons (>0.17 mm/yr; Fig. 4C). The MicH protein was similarely

detected in planktonic ($36.5 - 1473.5$ pg/mL) as well as biofilm samples ($41.0 - 7971.3$ pg /cm^2)

collected from '+MIC' enrichments, indicating active expression of the 'MIC hydrogenase' in

those cultures (Fig. 4D and Fig. S3B). Interestingly, we detected a small amount of the *micH* gene

in one of the enrichments showing low corrosion (0.04 mm/yr). However, the MicH protein has

not been detected, suggesting that the *micH* gene may not have been expressed and no corrosive

biofilm formed (Fig. 4).

Discussion

Multiple studies have highlighted the importance of understanding the underlying MIC

mechanisms to better detect and diagnose MIC[5,7,18,19]. Here we presented the development of a

MicH-specific immunoassay that showed strong affinity for the large subunit of a the 'MIC

hydrogenase' shown to be involved in corrosion by certain methanogenic Archaea[5,11]. The results

obtained with pure and field enrichment cultures suggest that the Anti-MicH antibody is highly

selective for the MicH protein found in certain corrosive methanogenic Archaea while showing

little to no cross-reactivity with proteins from SRB or non-corrosive methanogenic Archaea. This

also included oil field enrichment cultures showing methanogenic activity. In addition, the assay

confirmed that corrosive methanogenic cultures indeed actively expressed the 'MIC hydrogenase'

thereby accelerating the iron oxidation of carbon steel as previously shown in pure cultures and

genetically modified *Methanococcus maripaludis* strains[11,12].

Interestingly, we were able to detect the *micH* gene as well as the MicH protein in cultures of the

corrosive *Methanobacterium*-like strain IM1[16]. To our knowledge this is the first dataset that

allowed a deeper insight into the underlying molecular mechanisms of this corrosive methanogenic

archaeon. Previous potential step measurements have suggested a biologically catalyzed H_2

production pathway on graphite electrodes[20]. The identification of the *micH* gene and protein in

strain IM1 further supports an enzymatic catalyzed pathway for proton reduction on steel surfaces.

The detecting of the MicH protein not only in biofilms but also planktonic cells suggest that the

assay may be applicable to both MIC monitoring in biofilms as well as water samples. The latter

is often readily available in the field, while it is often challenging to obtain relevant biofilm

samples in the field to monitor pipeline corrosion. Biofilms act as planktonic cell factories

continuously shedding cells into fluids streams[21,22] that can then be collected at pipeline outlets

and can be analyzed for presents and expression of the MIC biomarker. We could previously show

the feasibility of using qPCR assays to detect MIC biomarkers in produced water samples as well

as pipeline solids from various global operations [5,14] and it would be interesting to see whether

similar can be accomplished using the novel immunoassay.

Overall, the development of a MicH-specific immunoassay offers an additional tool for corrosion

scientists and field practitioners to detect, monitor and understand corrosive methanogenic

biofilms. Furthermore, the detection of the MicH protein in multiple field enrichment cultures from

different locations together with previous qPCR results[5] highlight the importance of this MIC

mechanism in anaerobic systems.

Conclusion

Here, we built on previous research and findings to develop specific immunoassay for the detection

of an industrially important MIC mechanism. The assay offers corrosion scientists and field

practitioners a novel way to detect, monitor and understand corrosive methanogenic biofilms.

Furthermore, the detection of the MicH protein in multiple field enrichment cultures from different

locations together with previous qPCR results5 highlight the importance of this MIC mechanism

in anaerobic systems on a global scale.

Establishment and optimization of a novel Western blot-based immunoassay to detect an

important MIC biomarker ('MicH') found in certain highly corrosive methanogenic Archaea.

Analysis of recombinant MicH protein and pure cultures of corrosive and non-corrosive

methanogens and SRB showed high sensitivity and selectivity of the immunoassay

Test with corrosive and non-corrosive oil field enrichment cultures showed applicability

for Mic detection in the field.

Acknowledgement

The authors like to thank Manuel Castelan for assistance with gas analysis and Jaspreet Mand for providing protein extracts for the study as well as Dennis Enning for the fruitful discussions. In addition, we are grateful to Florin Musat for providing a culture of strain IM1 for our study.

References

1. Uchiyama, T., K. Ito, K. Mori, H. Tsurumaru, and S. Harayama, *Appl. Environ. Microbiol.* 76 (2010): pp. 1783–1788.

2. Okoro, C., S. Smith, L. Chiejina, R. Lumactud, D. An, H.S. Park, J. Voordouw, B.P. Lomans, and G. Voordouw, *J Ind Microbiol Biotechnol* 41 (2014): pp. 665–678.

3. Enning, D., and J. Garrelfs, *Appl. Environ. Microbiol.* 80 (2014): pp. 1226–1236.

4. Mand, J., H.S. Park, C. Okoro, B.P. Lomans, S. Smith, L. Chiejina, and G. Voordouw, *Front. Microbiol.* 6 (2016): fmicb.2015.01538

5. Lahme, S., J. Mand, J. Longwell, R. Smith, and D. Enning, *Applied and Environmental Microbiology* 87 (2021): pp. e01819-20.

6. Lomans, B.P., R. de Paula, B. Geissler, C.A. Kuijvenhoven, and N. Tsesmetzis, Paper presented at the SPE International Oilfield Corrosion Conference and Exhibition, Aberdeen, Scotland, UK (2016): SPE-179919-MS

7. Eckert, R.B., and T.L. Skovhus, *International Biodeterioration & Biodegradation* 126 (2018): pp. 169–176.

8. TM0194-2014-SG, Field Monitoring of Bacterial Growth in Oil and Gas Systems, AMPP

9. TM0212-2018-SG, Detection, Testing, and Evaluation of Microbiologically Influenced Corrosion on Internal Surfaces of Pipelines, AMPP

10. Enning, D., H. Venzlaff, J. Garrelfs, H.T. Dinh, V. Meyer, K. Mayrhofer, A.W. Hassel, M. Stratmann, and F. Widdel, *Environmental Microbiology* 14 (2012): pp. 1772–1787.

11. Tsurumaru, H., N. Ito, K. Mori, S. Wakai, T. Uchiyama, T. Iino, A. Hosoyama, H. Ataku, K.

Nishijima, M. Mise, A. Shimizu, T. Harada, H. Horikawa, N. Ichikawa, T. Sekigawa, K.

Jinno, S. Tanikawa, J. Yamazaki, K. Sasaki, S. Yamazaki, N. Fujita, and S. Harayama,

Scientific Reports 8 (2018): p. 15149.

12. Holten, M.P., D.R. Fonseca, and K.C. Costa, *Applied and Environmental Microbiology* 87

(2021): pp. e00995-21.

13. Kawaichi, S., R. Kotoky, J. Fiutowski, and A.-E. Rotaru, *BioRxiv* (2024): p.

2024.03.06.583737.

14. Lahme, S., J. Mand, K. Oparaodu, and D. Enning, Paper presented at the ADIPEC, Abu

Dhabi, UAE (2022): SPE-211238-MS

15. Lahme, S., D. Enning, C.M. Callbeck, D.M. Vega, T.P. Curtis, I.M. Head, and C.R.J. Hubert,

Appl. Environ. Microbiol. 85 (2019): pp. e01891-18.

16. Dinh, H.T., J. Kuever, M. Mußmann, A.W. Hassel, M. Stratmann, and F. Widdel, *Nature* 427

(2004): pp. 829–832.

17. Kurien, B.T., and R.H. Scofield, *Methods* 38 (2006): pp. 283–293.

18. Knisz, J., R. Eckert, L.M. Gieg, A. Koerdt, J.S. Lee, E.R. Silva, T.L. Skovhus, B.A.

An Stepec, and S.A. Wade, *FEMS Microbiology Reviews* 47 (2023): p. fuad041.

19. Little, B.J., D.J. Blackwood, J. Hinks, F.M. Lauro, E. Marsili, A. Okamoto, S.A. Rice, S.A.

Wade, and H.-C. Flemming, *Corrosion Science* 170 (2020): p. 108641.

20. Beese-Vasbender, P.F., J.-P. Grote, J. Garrelfs, M. Stratmann, and K.J.J. Mayrhofer,

Bioelectrochemistry 102 (2015): pp. 50–55.

21. Bester, E., G.M. Wolfaardt, N.B. Aznaveh, and J. Greener, *International Journal of

Molecular Sciences* 14 (2013): pp. 21965–21982.

22. Klopper, K.B., E. Bester, M. van Schalkwyk, and G.M. Wolfaardt, *Environmental*

Microbiology Reports 16 (2024): p. e70010.

Figure 1 Performance of recombinant antibody Anti-MicH using a recombinant MicH protein as

target. A) Chemiluminescence measured as arbitrary units (A.U.) showing signal at the expected

molecular weight of $MicH_{rec}$. Test was performed with a 1:20 dilution of the antibody. B) Results

of tests performed with varying dilutions of Anti-MicH and 1 ng/ml $MicH_{rec}$.

Figure 2 Western Blots performed with dilution series of recombinant MicH protein and Anti-

MicH antibody dilutions of 1:20. The error bar represent the standard deviation of technical

triplicates.

Figure 3 Analysis of protein extracts with MicH-specific Western blot immunoassay and DNA

extracts with micH qPCR assay. A) Recombinant MicH Protein (10 ng/mL), B) *Methanobacterium*

sp. strain IM1, C) *Methanococcus maripaludis* strain S2, D) *Desulfovibrio ferrophilus* IS5, E)

Desulfovibrio alaskensis G20. Average corrosion rates are shown below the bar plot. The error

bars where seen indicate the standard deviations of triplicate cultures or technical triplicates in

case of recombinant MicH.

Figure 4 A) Weight loss corrosion rates and methane formation in oil field produced water enrichment cultures classified as corrosive ['+MIC'] and non-corrosive ['-MIC'] enrichments. B) MicH protein numbers detected in corrosion coupon biofilms (open diamonds) and planktonic cells (closed diamonds). Enrichments 1, 2, 4, 6, 7, 9, 10 – First West African oil field. Enrichment 3 – Second West African oil field. Enrichment 5 – U.S. offshore oil field. Enrichment 8 – U.S. onshore oil field. Error bars in A) indicate standard deviation of weight loss analysis from three corrosion coupons.