**A genome-wide association study identifies markers and candidate genes**

**affecting tolerance to the wheat pathogen *Zymoseptoria tritici***

Alexey Mikaberidze[1*], Bruce A. McDonald[2], Lukas Kronenberg[3]

[1]*School of Agriculture, Policy and Development; University of Reading, Reading, United*

*Kingdom*

[2]*Plant Pathology, Institute of Integrative Biology, ETH Zurich, Zurich, Switzerland.*

[3]*Crop Genetics, The John Innes Centre, Norwich, United Kingdom.*

[*]Corresponding author: A. Mikaberidze; Email: a.mikaberidze@reading.ac.uk

Keywords: leaf tolerance, trade-off, GWAS, disease resistance, marker-trait association,

candidate genes, *Zymoseptoria tritici*, septoria tritici blotch, plant defense

**Abstract**

Plants defend themselves against pathogens using either resistance, measured as the

host's ability to limit pathogen multiplication, or tolerance, measured as the host's ability

to reduce the negative effects of infection. Tolerance is a promising trait for crop

breeding, but its genetic basis has rarely been studied and remains poorly understood.

Here, we reveal the genetic basis of leaf tolerance to the fungal pathogen *Zymoseptoria*

*tritici* that causes the globally important septoria tritici blotch disease on wheat. Leaf

tolerance to *Z. tritici* is a quantitative trait that was recently discovered in wheat by using

automated image analyses that quantified the symptomatic leaf area and counted the

number of pycnidia found on the same leaf. A genome-wide association study including

both tolerance and resistance to STB found a strong negative genetic correlation

between these traits, indicative of a trade-off. We identified four chromosome intervals

associated with tolerance and a separate chromosome interval associated with

resistance. Within these intervals, we identified candidate genes, including wall-

associated kinases similar to *Stb6*, the first cloned STB resistance gene. A trade-off

between tolerance and resistance would hinder breeding simultaneously for both traits,

but our findings suggest a way forward using marker-assisted breeding. We expect that

the methods described here can be used to characterize tolerance to other fungal

diseases that produce visible fruiting bodies, such as speckled leaf blotch on barley,

potentially unveiling conserved tolerance mechanisms shared among plant species.

**Introduction**

Plants defend themselves against pathogens using either resistance, measured as the host's ability to limit pathogen multiplication, or tolerance, measured as the host's ability to reduce the negative effects of infection (Pagan and Garcia-Arenal 2020). A fundamental difference between these two strategies is that resistance reduces the multiplication rate of the pathogen, whereas tolerance does not. Tolerance was first recognized by plant pathologists in 1894 (Cobb 1894), and it is thought to be a host defense strategy as common and important as resistance (Pagan and Garcia-Arenal 2020). Yet, our knowledge of the mechanisms and genes controlling tolerance pales in comparison to our knowledge of the mechanisms and genes underlying resistance. This stems in part from the difficulty of measuring tolerance in plants and also from a lack of agreement in how tolerance should be defined. The excellent, comprehensive review by Pagan and Garcia-Arenal (2020) presents well-reasoned definitions that should resolve the latter difficulty.

The genetic basis of tolerance in plants has rarely been studied and remains poorly understood. The Pagan and Garcia-Arenal (2020) review identified ten examples where tolerance was inferred and its genetic basis was analyzed. In most of these cases, tolerance appeared to be a quantitative trait, involving from one to 70 quantitative trait loci (QTL) or candidate genes, but in some of these cases, it remains unclear whether the measured trait was truly tolerance or a form of resistance that was treated as

tolerance (Ayala et al., 2002; Williams et al., 2003; Han et al., 2008). For example, Han

et al. (2008) identified eight QTLs associated with tolerance to *Phytophthora sojae* in

soybeans, using the proportion of surviving plants as a proxy for tolerance. But it is

possible that the plants surviving an exposure to *P. sojae* were displaying partial

resistance instead of tolerance to this pathogen. The most extensive work on tolerance

has been conducted with plant viruses, including a study where tolerance to the barley

yellow dwarf virus in wheat was ascribed to 22 QTLs of minor effect (Ayala et al. 2002).

In tomatoes, a MAP kinase was found to enhance tolerance to the tomato yellow leaf

curl virus (TYLCV) by regulating salicylic acid and jasmonic acid signaling (Li et al.

2017). More recently, the overexpression of a cellulose synthase-like gene was shown

to boost tolerance to TYLCV (Choe et al., 2021). Functional alleles of flowering

repressor genes in *Arabidopsis thaliana* were found to contribute to plant tolerance to

cucumber mosaic virus (Shukla et al., 2021). Tamisier et al. (2022) identified candidate

genes for potato virus Y tolerance in peppers (*Capsicum annuum*), including a cluster of

NBS-LRR genes. But we are not aware of any studies that have identified candidate

genes specifically associated with tolerance to fungal plant pathogens until now.

Septoria tritici blotch (STB) is the most damaging disease of wheat in Europe

(Jorgensen et al. 2014) and among the most important diseases of wheat globally

(Savary et al. 2019). STB is caused by the fungus *Zymoseptoria tritici*, a pathogen that

has co-evolved with wheat for more than 10,000 years (Stukenbrock et al. 2007) and

has a high evolutionary potential (McDonald et al. 2022). The most common strategies

for controlling STB are deployments of fungicides and STB-resistant wheat cultivars.

Exposed *Z. tritici* populations typically evolve resistance to fungicides and virulence

against resistant cultivars within a few years of deployment (McDonald and Mundt 2016;

McDonald et al. 2019; Kildea et al. 2020). The genetic basis of fungicide resistance and

virulence has been explored in several populations of *Z. tritici*, leading to the discovery,

cloning and functional validation of several of the underlying genes for both traits (Meile

et al. 2018; Zhong et al. 2017; Amezrou et al. 2023; Garnault et al. 2019). Several STB

resistance genes in wheat have also been cloned and functionally validated (Saintenac

et al. 2021; Saintenac et al. 2018; Hafeez et al. 2023). The cloned pathogen avirulence

genes and cloned wheat resistance genes have been shown to largely conform to the -

gene-for-gene concept of plant-pathogen interactions, though resistance to STB does

not appear to involve a hypersensitive response (Saintenac et al. 2018).

Leaf-level tolerance to *Z. tritici* is a quantitative trait that was recently discovered in

wheat (Mikaberidze and McDonald 2020) by using automated image analyses that could

accurately quantify the leaf area affected by STB and count the number of fungal fruiting

bodies (called pycnidia) found on the same leaf (Stewart et al. 2016; Karisto et al. 2018).

These measures were then used to quantify degrees of STB resistance and STB

tolerance in wheat, the latter using a novel measure called kappa, in 335 elite winter

wheat cultivars growing in the same field. Resistance was quantified as the average

number of pycnidia on a leaf, $N_p$, adjusted for the overall leaf area. More resistant plant

genotypes suppressed pathogen reproduction, and consequently carried fewer pycnidia

on their leaves. Kappa is an exponential slope that characterizes the negative

relationship between green leaf area and the number of pycnidia on a leaf. Lower kappa

values correspond to higher tolerance: when inhabited by pathogen populations of the

same size (i.e., when leaves carry the same numbers of pycnidia), more tolerant plant

genotypes retain larger green leaf areas than less tolerant genotypes (Figure 1).

Mikaberidze & McDonald (2020) showed that there was a wide, continuous variation in

both resistance and leaf tolerance across the 335 wheat cultivars. They also found a

negative relationship between tolerance and resistance, indicative of a trade-off

between these traits.

Before the discovery of leaf tolerance, we used these data to conduct a genome-wide

association study (GWAS) to analyze the genetic architecture of STB resistance in the

335 wheat cultivars and identified several chromosome regions that contained

interesting candidate STB resistance genes (Yates et al., 2019). Here we use the leaf

tolerance trait kappa to conduct a GWAS aiming to elucidate the genetic architecture of

tolerance and identify candidate genes that may be associated with leaf tolerance. We

also sought to determine if the trade-off between tolerance and resistance to STB has a

genetic basis. We discovered that the genetic associations for tolerance were

independent from the previously described genetic associations for resistance to this

pathogen. We identified four chromosome intervals associated with leaf-level tolerance

and a separate chromosome interval associated with resistance. A bivariate GWAS that

jointly considered both tolerance and resistance did not provide significant marker

associations, even though there was a significant negative genetic correlation between

these traits. Within each of the significant chromosome intervals, we identified candidate

genes associated with tolerance that were overrepresented relative to the rest of the

genome.

**Results**

**Figure 1.** Illustration of leaf tolerance. STB symptoms on wheat leaves can be seen as characteristic necrotic lesions with pycnidia (seen as small, black, round structures within lesions) on cultivar Intact (A, tolerant) vs cultivar Lynx (B, non-tolerant). The number of pycnidia is similar in the two images, but the non-tolerant leaf has a larger disease-induced necrotic area (and a smaller green leaf area), and therefore suffers

more damage from disease compared to the tolerant leaf. Black rectangles are 4x

magnified in the insets, where two characteristic pycnidia are indicated by arrows. To

quantify leaf tolerance it is not sufficient to compare two leaves, instead, we analyzed

large numbers of diseased leaves from each cultivar and estimated tolerance as the

negative exponential slope of how green leaf areas decrease versus the number of

pycnidia on a leaf (see Figure 4 and Mikaberidze & McDonald 2020).

Leaf tolerance (quantified as kappa) and resistance to STB (quantified as the number of

pycnidia per leaf, $N_p$) were measured based on the analysis of 11,152 individual images

of naturally infected leaves coming from 335 elite winter wheat cultivars growing in a

replicated field experiment (Karisto et al. 2018). On average, each leaf was infected by a

different strain of *Z. tritici* (Lorrain et al. 2024; McDonald et al. 2022), hence the

measures of leaf tolerance and resistance calculated for each cultivar represent average

values across a very large number of pathogen strains. Despite the large variance

associated with infections involving thousands of pathogen strains, the heritabilities

were high for both leaf tolerance (0.44 for kappa) and resistance (0.88 for $N_p$). There

were strong phenotypic ($r_p$ = -0.40, *p*<0.001) and genetic ($r_g$ = -0.67, *p*<0.001; Figure 2)

correlations between leaf tolerance and resistance.

$r_g = -0.67, p = 4.4\text{e-}06$

**Figure 2.** Genetic correlation between tolerance (quantified as kappa) and resistance

(quantified as $N_p$). Purple dots represent the empirical best linear unbiased predictors

extracted from the bivariate model (Eq. 3) and gray lines represent the standard

deviations.

The leaf tolerance and resistance phenotypes were used to conduct a GWAS that

included, after filtering, 9,125 single nucleotide polymorphisms (SNPs) in 330 of the

wheat cultivars. The GWAS identified three marker-trait associations (MTAs) for leaf

tolerance that exceeded the Bonferroni significance threshold, located on chromosomes

2D, 6B and 7D (Figure 3). An additional leaf tolerance MTA on chromosome 7A fell just

below the Bonferroni threshold (LOD = 5.19, Figure 3), but was included in further

analyses. For resistance, $N_p$, one significant MTA was detected on chromosome 5A.

The bivariate GWAS that jointly considered leaf tolerance and resistance did not detect

any significant associations, though the MTAs observed on chromosomes 5A, 6B and

7D in the univariate analyses remained distinct relative to other SNPs (Figure 3, panel

C). Though it was not significant at the Bonferroni threshold, a distinctive peak visible on

chromosome 7A (LOD = 4.01, Figure 3, panel C) was included in further analyses.

**Figure 3:** Manhattan and QQ-plots depicting the GWAS results for tolerance (kappa)

(A), resistance (B) and the combined, bivariate GWAS (C). Green, red and purple dots

in the Manhattan plots represent the SNPs significantly associated with the respective

traits: kappa, resistance and the bivariate model combining kappa and resistance.

Crossed circles represent marker-trait associations that were not significant at the

Bonferonni threshold, but were nevertheless considered in additional analyses. The

legends in the QQ-plots indicate the applied GWAS model and the genome-wide

inflation coefficient λ.

The SNPs and chromosome positions for each of the six identified MTAs are shown in

Table 1. For each MTA, a chromosome interval was defined by including the DNA

sequences positioned 2.5 million base pairs in each direction from the significant SNP,

i.e., by placing the most significant SNP at the center of a 5 Mb interval on the IWGSC

reference sequence v1.0 (IWGSC, 2018). The positions of these chromosome intervals

were then compared to the positions of chromosome intervals containing STB

resistance QTL that were reported in recent publications (Yates et al. 2019; Zakieh et al.

2023; Mekonnen et al. 2021; Alemu et al. 2021; Mahboubi et al. 2022) to determine if

there were any overlaps (Supplementary Table S1). The leaf tolerance MTA on

chromosome 2D was found to be 0.9 Mbp upstream of an STB resistance MTA that was

previously detected in a greenhouse experiment where seedlings of 316 Nordic

breeding lines were inoculated with two Nordic strains of *Z. tritici* (Zakieh et al. 2023).

The resistance MTA on chromosome 5A was found to be within the same interval as a

different STB resistance trait, PLACL (percentage of leaf area covered by lesions), that

was identified in our earlier analyses of the same dataset (Yates et al. 2019). The

tolerance MTA on chromosome 7D was found to be 1.6 Mbp downstream of the STB

resistance QTL detected in a greenhouse experiment where seedlings of 185 wheat

genotypes of globally diverse origin were inoculated with ten different *Z. tritici* strains of

global origin (Mahboubi et al., 2022). Using the consensus map by Wang et al. (2014),

we also compared our results to earlier publications reviewed in Brown et al, (2015).

There, we found that the resistance MTA on chromosome 5A and the tolerance MTAs

on chromosomes 6B, 7A and 7D were within the reported intervals for MQTL19, 21, 24

and 27, respectively. However, these intervals are too large (45, 22, 55 and 13 cM,

respectively) to be confident that these overlaps are biologically meaningful.  Therefore,

while our tolerance MTA on chromosomes 2D, and 7D fall within previously reported

resistance intervals, our tolerance MTA on chromosomes 6B and 7A are likely to

constitute new loci or refined loci for tolerance within meta-QTL previously associated

with resistance.

**Table 1:** Marker-trait associations (MTAs) for tolerance (kappa), resistance and the

combined, bivariate GWAS. For each MTA, we report the physical position, minor allele

frequency (MAF), LOD score, the SNP effect estimates, and the percentage of

phenotypic variance explained (PVE). Non-significant MTAs are indicated with "n.s."

| Interval | SNP | Position | MAF | LOD | Effect | PVE |
|---|---|---|---|---|---|---|
| kappa 2D | IAAV8779 | 647,278,479 | 0.288 | 5.46 | 0.178 | 4.47 |

| | | | | | | |
|---|---|---|---|---|---|---|
| resistance 5A | Ra_c7322_2294 | 20,816,495 | 0.297 | 8.12 | -0.286 | 12.72 |
| kappa 6B | wsnp_Ex_c702_1383612 | 21,415,561 | 0.053 | 8.04 | 0.446 | 33.61 |
| kappa 7D | wsnp_Ex_c12102_19361467 | 557,843,433 | 0.198 | 5.39 | 0.195 | 8.02 |
| kappa 7A (n.s.) | BS00099804_51 | 89,835,842 | 0.139 | 5.19 | -0.226 | 7.55 |
| (kappa, resistance) 7A (n.s.) | CAP7_c7296_88 | 642,736,799 | 0.212 | 4.01 | (-0.233; 0.025) | - |

In order to identify candidate genes, we searched the IWGSC refseq1.0 annotation

(IWGSC, 2018) within 5 Mb windows around each of the six identified MTAs. We

selected candidate genes within these regions based on their functional description

(Supplementary Tables S2-S7) and classified them into 16 motif groups. Next, we

compared the occurrence of these groups within each interval to their occurrence in

10,000 5 Mbp intervals randomly chosen across the genome (see Materials and

Methods). Twelve of these motif groups showed a significant (p<0.05)

overrepresentation compared to random genome intervals and contained a total of 50

candidate genes (Table 2). For details regarding the motif groups and individual genes,

see Supplementary Tables S8, S9 and S10.

Around the resistance MTA on chromosome 5A, we found three occurrences of "leucine

rich repeat" (LRR) motifs, whereas in 95% of the 10'000 random intervals, no more than

two LRR motifs were present. Thus, LRR motifs are considered significantly

overrepresented around the resistance MTA on chromosome 5A (Table 2, observed = 3,

$p < 0.05$). Likewise, around the MTAs for leaf tolerance, we found a significant

overrepresentation of gene motifs associated with programmed cell death (cysteine

proteinase inhibitors, cysteine protease, RPP13, DCD domain), disease resistance

motifs (GCN1, NBS-LRRs, RPM1, G/L Lectin RLK, Aspartate--tRNA ligase, wall-

associated kinase), cell wall associated Ran-binding and wound responsive motifs. The

bivariate MTA on chromosome 7A also showed a significant overrepresentation of cell

wall associated Ran-binding and wound-responsive motifs. Among these candidate

genes, we consider the wall associated kinase near the kappa MTA on chromosome 6B

(TraesCS6B01G035300, $p < 0.05$) to be especially interesting because *STB6,* the first

cloned resistance gene for STB, is also a wall associated serine/threonine kinase with a

galacturonan-binding domain (Saintenac et al. 2018).

**Table 2: Candidate gene motifs within 5 Mbp intervals around marker-trait associations**. The

significance of motif overrepresentation was determined based on the probability distribution of their

occurrence across random samples. Column 4 shows the total occurrence of motifs across the entire

genome. Columns 5-8 show the expected occurrence of motifs in 90%, 95%, 99% and 99.9% of the

random samples, respectively. Column 9 shows the observed occurrence of motifs in the intervals around

MTAs. '**' denotes significant overrepresentation for observed > q99 (i.e. $p<0.01$), '*' denotes observed >

q95 (i.e. $p<0.05$), '.' denotes observed > q90 (i.e. $p<0.1$) and 'n.s.' denotes no significant

overrepresentation of a motif within a given associated interval.

| interval | genes in interval | gene motif group | occurrence genome | expected <= probability quantile | | | | observed | |
|---|---|---|---|---|---|---|---|---|---|
| | | | | q90 | q95 | q99 | q99.9 | | |
| kappa 2D | 97 | Cysteine proteinase inhibitor | 101 | 0 | 0 | 1 | 5 | 4 | ** |
| | | GCN1 | 6 | 0 | 0 | 0 | 1 | 1 | ** |
| | | RPM13 | 46 | 0 | 0 | 1 | 3 | 2 | ** |
| | | Cysteine protease | 225 | 0 | 1 | 3 | 11 | 2 | * |
| | | NBS-LRR | 1908 | 4 | 7 | 17 | 26 | 8 | * |
| | | RPM1 | 376 | 1 | 2 | 5 | 16 | 5 | * |
| | | Disease resistance protein | 144 | 0 | 1 | 2 | 5 | 1 | . |
| | | Expansin | 262 | 0 | 1 | 3 | 7 | 1 | . |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | | Wat1 | 194 | 0 | 1 | 3 | 6 | 1 | . |
| | | LRR | 746 | 1 | 2 | 8 | 20 | 1 | n.s. |
| kappa 6B | 62 | NBS-LRR | 1908 | 4 | 7 | 17 | 26 | 10 | * |
| | | RPM1 | 376 | 1 | 2 | 5 | 16 | 5 | * |
| | | Wall-associated kinase | 95 | 0 | 0 | 1 | 3 | 1 | * |
| | | LRR | 746 | 1 | 2 | 8 | 20 | 2 | . |
| kappa 7A (n.s.) | 43 | DCD domain | 34 | 0 | 0 | 1 | 2 | 1 | * |
| | | Lectin RLK | 209 | 0 | 1 | 2 | 4 | 1 | . |
| kappa 7D | 30 | Aspartate--tRNA ligase | 16 | 0 | 0 | 0 | 1 | 1 | ** |
| | | Ran-binding | 12 | 0 | 0 | 0 | 2 | 1 | ** |
| | | Wound-responsive | 69 | 0 | 0 | 1 | 14 | 2 | ** |
| resistance 5A | 50 | LRR | 746 | 1 | 2 | 8 | 20 | 3 | * |
| | | Cysteine protease | 225 | 0 | 1 | 3 | 11 | 1 | . |
| | | Disease resistance protein | 144 | 0 | 1 | 2 | 5 | 1 | . |
| | | NBS-LRR | 1908 | 4 | 7 | 17 | 26 | 4 | n.s. |
| (kappa, resistance) | 61 | Ran-binding | 12 | 0 | 0 | 0 | 2 | 1 | ** |
| | | Wound-responsive | 69 | 0 | 0 | 1 | 14 | 3 | ** |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| 7A (n.s.) | | LRR | 746 | 1 | 2 | 8 | 20 | 1 | n.s. |

**Discussion**

Though several studies have analyzed the genetics of disease tolerance in plants

(Pagan & Garcia Arenal 2020), we believe this is the first study to reveal the genetic

basis of tolerance to a fungal plant pathogen and identify candidate genes that may

confer the leaf tolerance phenotype. Among these candidates are wall-associated

kinases with galacturonan-binding and serine/threonine kinase domains similar to *Stb6*,

the first cloned STB resistance gene (Saintenac et al. 2018). Our automated analyses of

more than 11,000 scanned leaves revealed that wheat's response to an STB infection

can be dissected into resistance and tolerance components that display high and

moderate heritabilities, respectively. In the analyzed population of 330 elite winter wheat

cultivars, the two traits showed a strongly negative genetic correlation (Figure 2),

supporting our previous report of a substantial negative phenotypic correlation indicative

of a trade-off between tolerance and resistance (Mikaberidze & McDonald 2020).

While screening for germplasm that performs well during an STB epidemic, wheat

breeders cannot distinguish between the leaf tolerance and resistance traits using

traditional visual scoring. A more detailed phenotyping based on leaf image analysis is

required to distinguish between them. As a result, breeding based on visual

assessments may select germplasm that has higher resistance or higher tolerance, both

of which could produce an improved yield response to STB infections compared to STB

susceptible germplasm. This may explain why the STB tolerance QTL that we identified

on chromosome 2A overlapped with an STB resistance QTL identified in Nordic

breeding lines based on traditional visual scoring of STB under greenhouse conditions

(Supplementary Table S1; Zakieh et al., 2023). Furthermore, because breeders have

been selecting for better performance in elite European winter wheat over several

decades using visual scoring, we would expect these efforts to result in cultivars with

high values for both tolerance and resistance. But it was rare to find elite winter wheat

cultivars that show high values for both traits (Figure 2), suggesting that there is indeed

a trade-off between the two traits. These interpretations are consistent with our previous

analysis, where we found a positive correlation between cultivar release year and

degree of tolerance and an absence of such a relationship for resistance, which

suggested that European wheat breeders may have been selecting for tolerance instead

of resistance to STB during recent decades (Mikaberidze & McDonald 2020).

Should wheat breeding programs seek to limit STB damage by combining STB

resistance and tolerance into the same cultivar? This would require that the two traits

are encoded by a shared pathway or by separate sets of genes that can be recombined

into the same lineage, as well as the absence of epistasis. Our analyses reveal a

nuanced picture of genetic connections between STB resistance and tolerance. The

significant and strongly negative genetic correlation between the two traits suggests that

the traits are not independent of each other. Yet the univariate GWAS analyses

identified different chromosome intervals associated with tolerance and resistance, while

the bivariate GWAS revealed only one association that was visually striking, but not

statistically significant. A similar pattern was found in studies of resistance and tolerance

of pepper plants to potato virus Y: the two traits exhibited significant negative phenotypic

and genetic correlations, the GWAS identified markers that were significantly associated

with either tolerance or resistance, but none of the markers were shared between the

two traits (Tamisier et al., 2020; Tamisier et al., 2022). A possible explanation of these

outcomes is that there could be several molecular pathways contributing to tolerance.

Some of the pathways exhibit a negative genetic correlation with resistance and are

underpinned by a large number of genes of small effect. Here, a negative correlation

between tolerance and resistance may result from these genes exhibiting negative

pleiotropy, whereby the same gene contributes to an increase in tolerance, but a

decrease in resistance (or vice versa). Alternatively, a negative correlation between the

two traits can be caused by linked monotropic genes: among several linked genes,

some contribute to an increase in tolerance, while others contribute to a decrease in

resistance (Gardner and Latta, 2007). Under either of these scenarios, we would not be

able to capture these genes via MTAs in the GWAS. Other tolerance pathways could be

independent of resistance, and conferred by fewer genes with larger effects. These

would be identified as significant MTAs for tolerance in the GWAS, none of which were

significantly associated with resistance. Hence, purely phenotypic selection for tolerance

may inadvertently select against resistance and vice versa. However, marker-assisted

selection for components of tolerance that are independent of resistance could avoid

this pitfall and select for tolerance without compromising resistance. Additional

experiments will be needed to further validate the tolerance MTAs and the associated

candidate genes to enable this approach.

We identified tolerance candidate genes and discovered a significant overrepresentation

of genes encoding programmed cell death, leucine-rich repeats, responses to wounding,

and wall-associated kinases. Many of these candidate genes encode functions that are

typically associated with disease resistance, such as NBS-LRRs, lectin receptor-like

kinases, and wall associated kinases, including genes that are similar to *Stb6*, the first

cloned STB resistance gene (Saintenac et al. 2018). Similarly, Tamisier et al (2022)

found a cluster of candidate NBS-LRR genes to be associated with tolerance of pepper

plants to potato virus Y. This suggests that there may be a common genetic architecture

underlying plant resistance and tolerance. In particular the same pathogen sensing,

recognition, and signaling processes may be involved at the beginning of the tolerance

and resistance response pathways. Under this scenario, the same gene may confer

tolerance to one pathogen, but resistance to a different pathogen. For example, a

candidate NBS-LRR gene for potato virus Y tolerance in pepper (*Capsicum annuum*)

shares 87.1% nucleotide identity with the *Bs2* gene in a different pepper species

(*Capsicum chacoense*) known to confer resistance to bacterial spot disease (Tamisier et

al., 2022). For tolerance, we found a greater overrepresentation of gene motifs

associated with wound responses and programmed cell death (PCD) compared to the

resistance trait (Table 2). We speculate that inhibition or appropriate regulation of wound

response or PCD pathways may lead to increased leaf tolerance via reduction of

excessive necrosis of leaf tissue.

While this is the first work to identify candidate genes affecting tolerance to fungal

pathogens, we expect that the methods described in this paper can be applied to many

other fungal diseases that produce visible fruiting bodies, such as speckled leaf blotch

on barley, septoria leaf blotch on oats, or septoria leaf spot of tomatoes. As tolerance is

analyzed in other plant pathosystems, it will become possible to compare candidate

genes identified across different systems to determine if tolerance, like resistance, is

encoded by conserved mechanisms shared among many plant species.

**Materials and Methods**

Naturally infected penultimate leaves from 335 elite winter wheat cultivars were sampled

from replicated plots during the 2016 field season as described in earlier publications

(Karisto et al., 2018; Yates et al., 2019; Mikaberidze and McDonald, 2020). We estimate

that at least half a million *Z. tritici* genotypes were present in the sampled plots

(McDonald et al. 2022; Lorrain et al. 2024), thus our results are relevant for epidemics

caused by highly diverse, natural pathogen populations. On average, 16 infected leaves

from each plot were imaged using a flatbed scanner (Canon CanoScan LiDe 220) at

1200 dpi resolution to obtain the percentage of leaf area covered by lesions (PLACL, a

damage function that reduces the green leaf area, a measure of disease-induced

reduction of plant fitness for each leaf) and numbers of pycnidia ($N_p$, associated with

pathogen reproduction, a measure of the pathogen burden in each leaf) (Stewart et al.,

2016; Karisto et al., 2018). In order to control for the effect of total leaf area on the

number of pycnidia per leaf, we performed the adjustment $N_{p,i} \rightarrow (A_{tot}/A_{tot,i})\, N_{p,i}$, where $N_{p,i}$

and $A_{tot,i}$ are the number of pycnidia and the total area of an individual leaf $i$, and $A_{tot}$ is

the mean total leaf area averaged over the entire dataset (Mikaberidze and McDonald,

2020).

These measures were then used to quantify degrees of STB resistance and tolerance,

the latter using a novel measure called kappa. Kappa is an exponential slope that

characterizes the negative relationship between green leaf area and the number of

pycnidia on each leaf (Figure 4; Mikaberidze & McDonald, 2020). Small values of kappa

indicate a high level of leaf tolerance, while large values of kappa indicate a low level of

leaf tolerance. The kappa values were calculated for each cultivar based on

measurements from approximately 32 leaves (each cultivar was replicated twice in the

experiment), with each leaf typically infected by a different pathogen strain (McDonald et

al. 2022; Lorrain et al. 2024). Hence the tolerance measures represented average

values across a wide range of pathogen genotypes (≈32 *Z. tritici* strains) for each wheat

cultivar and the range of kappa values encompassed a wide range of host genotypes

(~330 wheat cultivars). Resistance was quantified as the average number of pycnidia

found on each leaf ($N_p$; adjusted for the total leaf area). More susceptible cultivars allow

higher numbers of pycnidia per leaf that translate to more pathogen reproduction, while

more resistant cultivars limit the numbers of pycnidia per leaf and therefore reduce

pathogen reproduction. Additional details on how tolerance and resistance were

calculated for each cultivar can be found in Mikaberidze & McDonald 2020. The

workflow for data acquisition and analysis is illustrated in Figure 4. The raw data

stemming from the image analysis of each individual diseased leaf reported by Karisto

et al. (2018) are available via Dryad Digital Repository:

https://doi.org/10.5061/dryad.171q4. The processed phenotypic and genomic datasets

underlying the outcomes of this study are available via (xxx; to be specified upon the

acceptance of the manuscript); the code that can be used the reproduce the analyses is

available via (xxx; to be specified upon the acceptance of the manuscript).

**Figure 4.** Workflow for the acquisition of phenotypic data, estimation of leaf tolerance

and GWAS [modified from Figure 1 in (Yates et al., 2019) and Figure 3 in (Mikaberidze

& McDonald, 2020].

*Statistical Analysis*

An initial data inspection revealed strong skews in the distributions of both tolerance and

resistance, which resulted in violations of the assumption of independence and

normality of the residual distribution in the subsequent linear modeling (Supplementary

Figure S1). For this reason, the raw data from individual plots was subjected to a rank-

based inverse normal transformation (RINT) using the R-package RNOmni (v1.0.1.2;

McCaw, 2023) before conducting further analyses. This brought the distributions of

tolerance and resistance close to the normal distribution and resulted in independent,

normally distributed residuals with a mean of zero.

To obtain best linear unbiased estimates (BLUEs) across replications while accounting

for spatial variability, a spatial model using two-dimensional p-splines was fitted in the R-

package SpATS (v1.0.18; Rodríguez-Álvarez *et al.*, 2017). The two complete blocks

were allocated diagonally in a virtual grid (see Kronenberg et al. 2021, Pérez-Valencia

et al. 2022 for details), with rows and columns corresponding to the relative plot

positions within each replicate of the experiment. The fitted model was:

$Y_{ijk} = \mu + G_i + f(r,c) + r_j + c_k + e_{ijk}$                             (1)

where $Y_{ijk}$ is the plot value of the respective trait: tolerance quantified as kappa or

resistance quantified as $N_p$. $\mu$ is the global intercept and $G_i$ is the response of the

genotype *i*. To account for spatial variability, $r_j$ and $c_k$ represent the effects of the row *j*

and column *k*, respectively, while *f(r,c)* is a smoothed bivariate surface across rows and

columns within the virtual grid, thus fitting an independent spatial trend to each of the

two replicates. From this model, the BLUEs were extracted to be used in the GWAS

whereas spatially corrected plot values, comprising the BLUEs and residual errors but

omitting spatial trends and other unwanted design factors, were used to calculate

heritability and genetic correlations.

To calculate heritability, the model:

$\widehat{Y}_i = \mu + G_i + e_i$                                                                              (2)

was fitted using the R-package asreml-R (v4.2.0.302, VSNi Team 2023), where $\widehat{Y}_i$ is the

spatially corrected plot value from Eq. (1) for the respective trait, $G_i$ is the random

genotype response with known variance-covariance structure based on the genome-

wide identity by state (IBS) relationship matrix calculated from single nucleotide

polymorphism (SNP) data using the R-package SNPRelate (v1.30.1; Zheng *et al.*, 2012)

and $e_i$ is the residual error. Heritability was estimated on a genotype difference basis:

$H^2_{\Delta BLUP}$ according to Eq. (24) in (Schmidt et al., 2019).

Phenotypic correlations were calculated as Pearson's *r* based on the adjusted genotype

means extracted from Eq. (1). To calculate the genetic correlation between kappa and

resistance, Eq. (2) was expanded to the bivariate model:

$\begin{pmatrix} \widehat{Y}_i^1 \\ \widehat{Y}_i^2 \end{pmatrix} = \begin{pmatrix} \mu^1 \\ \mu^2 \end{pmatrix} + \begin{pmatrix} G_i^1 \\ G_i^2 \end{pmatrix} + \begin{pmatrix} e_i^1 \\ e_i^2 \end{pmatrix}$                                           (3)

in the R-package asreml-R, where the superscripts 1 and 2 denote the two traits: kappa

and $N_p$, respectively. $G_i^1$ and $G_i^2$ were again set as random with known variance-

covariance structure based on IBS.

The genetic correlation was then calculated from the estimated variance and covariance

components from Eq. (3) following Holland et al. (2001):

$$Corr\left(G^1, G^2\right) = \frac{Cov\left(G^1, G^2\right)}{\sqrt{Var\left(G^1\right) Var\left(G^2\right)}}$$     (4)

*Genome-wide association studies (GWAS)*

Genome-wide association studies for the traits kappa, $N_p$ and the combined, bivariate

response were conducted following the same workflow as described by Roth &

Kronenberg et al. (2024). Marker data were supplied by the GABI wheat consortium

(Gogna *et al.*, 2022) for the GABI genotypes and by the Agroscope wheat breeding

program for the Swiss genotypes (Fossati and Brabant, 2003). Markers were mapped to

the *Triticum aestivum* reference sequence v.1.0 (IWGSC et al., 2018) using ncbi-blast+

(v2.9.0-2). Equivocally mapped markers were excluded, and the remaining markers

were filtered for a missing rate <0.05 and a minor allele frequency >0.05, resulting in

9125 SNPs for 330 wheat cultivars. The remaining missing markers were imputed using

*fastPHASE* (v.1.4; Scheet and Stephens, 2006) implemented in

https://github.com/mwylerCH/HapMap_Imputation.

The univariate GWAS analyses for kappa and $N_p$ were conducted using the Blink model

(Huang *et al.*, 2019) implemented in the R-package GAPIT3 (v.3.1.0) (Wang and Zhang,

2021). The bivariate GWAS was conducted using the software GEMMA (v0.98.1, Zhou

and Stephens, 2014) using the first three principal components among SNP genotypes

and the genome-wide IBS matrix to correct for population structure and relatedness,

respectively.

Taking into account advances in GWAS methodologies in wheat since the publication of

(Yates et al., 2019), we adapted our analyses in four ways. We used (i) an adjustment

for spatial variability using the R-package SpATS (Rodríguez-Álvarez et al., 2017); (ii) a

different transformation [RINT-transformation (McCaw et al., 2020) instead of log-

transformation] that better satisfies the assumption of normality in the residuals of the

applied linear models; (iii) a more comprehensive marker panel that includes Swiss

cultivars, thus adding 11 genotypes previously excluded due to missing marker data;

improved imputation of missing genotype data based on similarity of haplotype clusters

around the missing genotype; and (iv) a different GWA model [single marker-based

BLINK (Huang et al., 2019) instead of haplotype based PLINK (Purcell et al., 2007)] that

better manages the systematic inflation of p-values in genetic association tests. To

ensure reproducibility and better understand the effects of these four modifications, we

reanalyzed the data of (Yates et al., 2019), and compared those results with the

outcomes of our new GWAS pipeline applied to the same phenotypes. We illustrate the

comparison of the two analyses in Supplemental Figure S2.

*Identification of candidate genes*

To identify candidate genes, we searched the IWGSC refseq 1.0 annotation (IWGSC,

2018) for 2.5 Mb in each direction from the genome position of the associated SNP

markers identified by GWAS. In a first step, likely candidates for either leaf tolerance or

resistance were identified based on their functional description. Genes were categorized

into motif groups based on their description (see Table 2; Supplementary Tables S8 &

S9). Then the likelihood of occurrence for each motif was quantified using a

bootstrapping approach. We examined 10,000 random 5 Mb intervals across the entire

genome, counted the occurrence of the selected candidate motifs identified in the

intervals around the MTAs and calculated quantile distributions for the occurrence of the

respective motifs across the random samples. A motif was considered significantly

overrepresented if the occurrence in the identified interval was larger than the

occurrence in 95% of the random intervals.

**Acknowledgements**

We thank the group of Crop Science at ETH Zurich for allowing us to sample STB in

their field experiment and in particular Andreas Hund for designing the field experiment

and Hansueli Zellweger for crop husbandry. We are grateful to Petteri Karisto for leading

the collection and processing of leaf samples, for help with preparing Figure 4, and for

fruitful discussions, and to Steven Yates for conducting initial analyses. This work was

supported by the Swiss National Foundation (SNF) in the framework of the project

DisPhenHiT (grant no. 206826). AM gratefully acknowledges financial support from the

SNF through the Ambizione grant PZ00P3_161453.

**506   Literature Cited**

Alemu, A., Brazauskas, G., Gaikpa, D.S., Henriksson, T., Islamov, B., Jørgensen, L.N.,

Koppel, M., Koppel, R., Liatukas, Ž., Svensson, J.T. and Chawade, A. 2021. Genome-

wide association analysis and genomic prediction for adult-plant resistance to Septoria

tritici blotch and powdery mildew in winter wheat. Frontiers in Genetics. 12:661742.

Amezrou, R., Audéon, C., Compain, J., Gélisse, S., Ducasse, A., Saintenac, C., Lapalu,

N., Louet, C., Orford, S., Croll, D. and Amselem, J. 2023. A secreted protease-like

protein in Zymoseptoria tritici is responsible for avirulence on Stb9 resistance gene in

wheat. PLoS Pathogens, 19:e1011376.

Ayala, L., Henry, M., van Ginkel, M., Singh, R., Keller, B., Khairallah, M. 2002.

Identification of QTLs for BYDV tolerance in bread wheat. Euphytica. 128:249–59.

Brown, J.K., Chartrain, L., Lasserre-Zuber, P. and Saintenac, C. 2015. Genetics of

resistance to Zymoseptoria tritici and applications to wheat breeding. Fungal Genetics

and Biology, 79:33-41.

Choe, S., Choi, B., Kang, J.H., Seo, J.K. 2021. Tolerance to tomato yellow leaf curl virus

in transgenic tomato overexpressing a cellulose synthase-like gene. Plant Biotechnol J.,

19:657-659.

Cobb, N. 1894. Contributions to an economic knowledge of Australian rusts

(Uredineae). Agric. Gaz. N.S.W. 5:239–50.

Covarrubias-Pazaran, G. 2016. Genome-Assisted Prediction of Quantitative Traits

Using the R Package sommer. PLOS ONE 11:e0156744.

Fossati, D., Brabant, C. 2003. Die Weizenzüchtung in der Schweiz. AGRARForschung

10:447–458.

Garnault, M., Duplaix, C., Leroux, P., Couleaud, G., Carpentier, F., David, O. and

Walker, A.S. 2019. Spatiotemporal dynamics of fungicide resistance in the wheat

pathogen Zymoseptoria tritici in France. Pest Management Science 75:1794-1807.

Gogna, A., Schulthess, A.W., Röder, M.S., Ganal, M.W., Reif, J.C. 2022. Gabi wheat a

panel of European elite lines as central stock for wheat genetic research. Scientific Data

9:538.

Hafeez, A.N., Chartrain, L., Cong, F., Cambon, F., Clarke, M., Griffiths, S., Hayta, S.,

Jiang, M., Keller, B., Kirby, R. and Kolodziej, M.C., 2023. Septoria tritici blotch

resistance gene Stb15 encodes a lectin receptor-like kinase. bioRxiv

2023.09.11.557217; doi: https://doi.org/10.1101/2023.09.11.557217.

Han Y, Teng W, Yu K, Poysa V, Anderson T, et al. 2008. Mapping QTL tolerance to

Phytophthora root rot in soybean using microsatellite and RAPD/SCAR derived markers.

Euphytica 162:231–39.

Holland, J.B., Frey, K.J., Hammond, E.G. 2001. Correlated responses of fatty acid

composition, grain quality, and agronomic traits to nine cycles of recurrent selection for

increased oil content in oat. Euphytica 122:69–79.

Huang, M., Liu, X., Zhou, Y., Summers, R.M., Zhang, Z. 2019. BLINK: a package for the

next level of genome-wide association studies with both individuals and markers in the

millions. GigaScience 8:giy154; https://doi.org/10.1093/gigascience/giy154.

The International Wheat Genome Sequencing Consortium (IWGSC), Appels R. et al.,

2018. Shifting the limits in wheat research and breeding using a fully annotated

reference genome. Science 361:eaar7191.

Jørgensen, L.N., Hovmøller, M. S., Hansen, J. G., Lassen, P., Clark, B., Bayles, R.,

Rodemann, B., Flath, K., Jahn, M., Goral, T., Czembor, J.J., Cheyron, P., Maumene, C.,

De Pope, C., Ban, R., Nielsen, G.C., Berg, G. 2014. IPM strategies and their dilemmas

including an introduction to www.eurowheat.org. Journal of Integrative Agriculture

13:265–281.

Karisto, P., Walter, A., Hund, A., Yu, K., Anderegg, J., Mascher, F., McDonald, B.A.,

Mikaberidze, A. 2018. Ranking quantitative resistance to Septoria tritici blotch in elite

wheat cultivars using automated image analysis. Phytopathology 108:568-581.

Kildea, S., Byrne, J.J., Cucak, M. and Hutton, F. 2020. First report of virulence to the

septoria tritici blotch resistance gene Stb16q in the Irish *Zymoseptoria tritici* population.

New Disease Reports 41:13.

Kronenberg, L., Yates, S., Boer, M. P., Kirchgessner, N., Walter, A., & Hund, A. 2021.

Temperature response of wheat affects final height and the timing of stem elongation

under field conditions. Journal of Experimental Botany, 72:700–717.

https://doi.org/10.1093/jxb/eraa47

Li, Y., Qin, L., Zhao, J., Muhammad, T., Cao, H., Li, H., Zhang, Y. and Liang, Y. 2017.

SlMAPK3 enhances tolerance to Tomato yellow leaf curl virus (TYLCV) by regulating

salicylic acid and jasmonic acid signaling in tomato (Solanum lycopersicum). PLOS

ONE 12:e0172466

Lorrain, C., Feurtey, A., Alassimone, J. and McDonald, B. 2024. A novel genome-wide

association approach reveals wheat pathogen genes involved in host specialization.

Research Square, in review. https://doi.org/10.21203/rs.3.rs-4486034/v1

Mahboubi, M., Talebi, R., Mehrabi, R., Mohammad Naji, A., Maccaferri, M. and Kema,

G.H. 2022. Genetic analysis of novel resistance sources and genome-wide association

mapping identified novel QTLs for resistance to Zymoseptoria tritici, the causal agent of

septoria tritici blotch in wheat. Journal of Applied Genetics 63:429-445.

McCaw, Z., Lane, J. M., Saxena, R., Redline, S., Lin, X. 2020. Operating Characteristics

of the Rank-Based Inverse Normal Transformation for Quantitative Trait Analysis in

Genome-Wide Association Studies. Biometrics 76:1262–1272.

McCaw, Z.R. 2023. RNOmni: Rank Normal Transformation Omnibus Test.

https://doi.org/10.32614/CRAN.package.RNOmni

McDonald, B.A., Mundt, C.C. 2016. How knowledge of pathogen population biology

informs management of Septoria tritici blotch. Phytopathology 106:948-955.

McDonald, B.A., Suffert, F., Bernasconi, A. and Mikaberidze, A., 2022. How large and

diverse are field populations of fungal plant pathogens? The case of *Zymoseptoria tritici*.

Evolutionary Applications, 15:1360-1373.

McDonald, M.C., Renkin, M., Spackman, M., Orchard, B., Croll, D., Solomon, P.S. and

Milgate, A., 2019. Rapid parallel evolution of azole fungicide resistance in Australian

populations of the wheat pathogen *Zymoseptoria tritici*. Applied and Environmental

Microbiology, 85:e01908-18.

Meile, L., Croll, D., Brunner, P., Plissonneau, C., Hartmann, F., McDonald, B.A.,

Sánchez-Vallet, A. 2018. A fungal avirulence factor encoded in a highly plastic genomic

region triggers partial resistance to septoria tritici blotch. New Phytologist 219:1048–

1061.

Mekonnen, T., Sneller, C.H., Haileselassie, T., Ziyomo, C., Abeyo, B.G., Goodwin, S.B.,

Lule, D. and Tesfaye, K. 2021. Genome-wide association study reveals novel genetic

loci for quantitative resistance to septoria tritici blotch in wheat (Triticum aestivum L.).

Frontiers in Plant Science 12:671323.

Mikaberidze, A. and McDonald B.A. 2020. A tradeoff between tolerance and resistance

to a major fungal pathogen in elite wheat cultivars. New Phytologist 226:879-890.

Pérez-Valencia, D. M., Rodríguez-Álvarez, M. X., Boer, M. P., Kronenberg, L., Hund, A.,

Cabrera-Bosquet, L., Millet, E. J., and van Eeuwijk, F. A. 2022. A two-stage approach

for the spatio-temporal analysis of high-throughput phenotyping data. Scientific Reports

12:3177. https://doi.org/10.1038/s41598-022-06935-9

Pagán, I. and García-Arenal, F. 2020. Tolerance of plants to pathogens: a unifying view.

Annual Review of Phytopathology, 58:77-96.

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D., Maller,

J., Sklar, P., de Bakker, P.I., Daly, M.J., Sham, P.C. 2007. PLINK: a tool set for whole-

genome association and population-based linkage analyses. Am. J. Hum. Genet.

81:559-575.

Rodríguez-Álvarez MX, Boer MP, Eeuwijk FA van, Eilers PHC. 2017. Correcting for

spatial heterogeneity in plant breeding experiments with P-splines. Spatial Statistics

23:52–71.

Roth, L., Kronenberg, L., Aasen, H., Walter, A., Hartung, J., van Eeuwijk, F., Piepho, H.-

P., and Hund, A. 2024. High-throughput field phenotyping reveals that selection in

breeding has affected the phenology and temperature response of wheat in the stem

elongation phase. Journal of Experimental Botany 75:2084–2099

https://doi.org/10.1093/jxb/erad481

Saintenac, C., Cambon, F., Aouini, L., Verstappen, E., Ghaffary, S.M.T., Poucet, T.,

Marande, W., Berges, H., Xu, S., Jaouannet, M. and Favery, B. 2021. A wheat cysteine-

rich receptor-like kinase confers broad-spectrum resistance against Septoria tritici

blotch. Nature Communications, 12:433.

Saintenac, C., Lee, W.S., Cambon, F., Rudd, J.J., King, R.C., Marande, W., Powers,

S.J., Bergès, H., Phillips, A.L., Uauy, C. and Hammond-Kosack, K.E., 2018. Wheat

receptor-kinase-like protein Stb6 controls gene-for-gene resistance to fungal pathogen

*Zymoseptoria tritici*. Nature Genetics 50:368-374.

Savary, S., Willocquet, L., Pethybridge, S. J., Esker, P., McRoberts, N., and Nelson, A.

2019. The global burden of pathogens and pests on major food crops. Nature Ecology &

Evolution 3:430–439.

Scheet, P., and Stephens, M. 2006. A fast and flexible statistical model for large-scale

population genotype data: applications to inferring missing genotypes and haplotypic

phase. The American Journal of Human Genetics 78:629-644.

Schmidt, P., Hartung, J, Bennewitz, J, Piepho, H.-P. 2019. Heritability in Plant Breeding

on a Genotype-Difference Basis, Genetics 212:991–1008.

Shukla, A., Pagán, I., Crevillén, P., Alonso-Blanco, C. and García-Arenal, F. 2022. A

role of flowering genes in the tolerance of *Arabidopsis thaliana* to cucumber mosaic

virus. Molecular Plant Pathology 23:175–187.

Stewart, E.L., Hagerty, C.H., Mikaberidze, A., Mundt, C. C., Zhong, Z., McDonald, B.A.

2016. An improved method for measuring quantitative resistance to the wheat pathogen

Zymoseptoria tritici using high throughput automated image analysis. Phytopathology

106:782-788.

Stukenbrock, E. H., Banke, S., Javan-Nikkhah, M., and McDonald, B. A. 2007. Origin

and domestication of the fungal wheat pathogen *Mycosphaerella graminicola* via

sympatric speciation. Mol. Biol. Evol. 24:398-411.

Tamisier, L., Szadkowski, M., Girardot, G., Djian-Caporalino, C., Palloix, A., Hirsch, J.,

Moury, B. 2022. Concurrent evolution of resistance and tolerance to potato virus Y in

*Capsicum annuum* revealed by genome-wide association. Molecular Plant Pathology

23:254–264. https://doi.org/10.1111/mpp.13157

Tamisier, L., Szadkowski, M., Nemouchi, G., Lefebvre, V., Szadkowski, E., Duboscq, R.,

Santoni, S., Sarah, G., Sauvage, C., Palloix, A., Moury, B. 2020. Genome-wide

association mapping of QTLs implied in potato virus Y population sizes in pepper:

evidence for widespread resistance QTL pyramiding. Mol. Plant Pathol. 21:3-16.

The VSNi Team. 2023. asreml: Fits Linear Mixed Models using REML. R package

version 4.2.0.302, www.vsni.co.uk.

Wang, J. and Zhang, Z. 2021. GAPIT Version 3: Boosting Power and Accuracy for

Genomic Association and Prediction. Genomics, Proteomics & Bioinformatics 19:629-

640. https://doi.org/10.1016/j.gpb.2021.08.005

Williams, K.J., Lewis, J.G., Bogacki, P., Pallotta, M.A.,Willsmore, K.L., et al. 2003.

Mapping of a QTL contributing to cereal cyst nematode tolerance and resistance in

wheat. Aust. J. Agric. Res. 54:731–37.

Yates, S., Mikaberidze, A., Krattinger, S.G., Abrouk, M., Hund, A., Yu, K., Studer, B.,

Fouche, S., Meile, L., Pereira, D., Karisto, P. and McDonald, B.A. 2019. Precision

phenotyping reveals novel loci for quantitative resistance to septoria tritici blotch. Plant

Phenomics 3285904.

Zakieh, M., Alemu, A., Henriksson, T. et al. 2023. Exploring GWAS and genomic

prediction to improve Septoria tritici blotch resistance in wheat. Sci Rep 13:15651.

https://doi.org/10.1038/s41598-023-42856-x

Zheng, X., Levine, D., Shen, J., Gogarten, S.M., Laurie, C., Weir, B.S. 2012. A high-

performance computing toolset for relatedness and principal component analysis of

SNP data. Bioinformatics 28:3326–3328.

Zhong, Z., Marcel, T., Hartmann, F.E., Ma, X., Plissonneau,C., Zala, M., Ducasse, A.,

Compain, J., Lapalu, N., Amselem, J., McDonald, B.A., Croll, D. and Palma-Guerrero, J.

2017. A small secreted protein in *Zymoseptoria tritici* is responsible for avirulence on

wheat cultivars carrying the *Stb6* resistance gene. New Phytologist 214:619-631.

Zhou, X. and Stephens, M. 2014. Efficient multivariate linear mixed model algorithms for

genome-wide association studies. Nature Methods 11:407–409.

**Supplementary Material**

Supplementary Figure S1:  Residual vs. fitted values of the SpATS model, before and

after rank based inverse normal transformation.

Supplementary Figure S2: Reproduction of Yates et al. (2019) GWAS outcomes using

different models and marker sets.

Supplementary Table S1: Comparison of detected MTA to previously reported QTL for

*Z. tritici* resistance.

Supplementary Table S2: Gene list for the associated interval for kappa on

chromosome 2D.

Supplementary Table S3: Gene list for the associated interval for kappa on

chromosome 6B.

Supplementary Table S4: Gene list for the associated interval for kappa on

chromosome 7A.

Supplementary Table S5: Gene list for the associated interval for kappa on

chromosome 7D.

Supplementary Table S6: Gene list for the associated interval for resistance on

chromosome 5A.

Supplementary Table S7: Gene list for the associated interval for the bivariate GWAS

on chromosome 7A.

Supplementary Table S8: Gene motif group details including functional descriptions

comprised in the motif groups, total occurrence of motifs across the genome, and gene

IDs comprised in the motif groups.

Supplementary Table S9:  Identified candidate gene motifs within 5 Mbp intervals

around the respective marker-trait associations including candidate gene IDs per motif

group.

Supplementary Table S10:  Details for the identified candidate genes including motif

group, gene ID, start, end, functional description, Pfam description, interpro description,

GO description and GO IDs from the IWGSC ref. seq. v1.0.