Title:

Dynamic Wt1 expression in the gastrulation-stage mouse embryo specifies vascular and visceral smooth muscle cell fate independently from mesothelial fate.

Authors: Suad H. Alsukari[1,3], Huei Teng Ng[1], Lilly Lang[1], Sharna Lunn[2], Shanthi Beglinger[1], Lauren Carr[1], Michael Boyes[1], David A. Turner[2] and Bettina Wilm[1]*

Affiliations:

[1]Department of Women's and Children's Health, Institute of Life Course and Medical Sciences, Faculty of Health and Life Sciences, University of Liverpool, Liverpool, UK

[2]Department of Musculoskeletal and Ageing Science, Institute of Life Course and Medical Sciences, Faculty of Health and Life Sciences, University of Liverpool, Liverpool, UK

[3]current address: Department of Biology, Albaha University, Albaha, KSA

*To whom correspondence is to be addressed (bwilm@liverpool.ac.uk)

Acknowledgements:

We acknowledge funding by the BBSRC Alert13, grant number BB/L014947/1, for the Zeiss Z.1 Light sheet microscope. We are also indebted to the University of Liverpool's (UoL) Centre for Cell Imaging (CCI) facility for provision of state-of-the-art imaging equipment funded through grants awarded to the University of Liverpool's CCI by the MRC (MR/K015931/1) and the BBSRC (BB/M012441/1, BB/R01390X/1), and for excellent technical support, assistance, and training. We express our thanks to the Biomedical Services Unit at the University of Liverpool, for expert support in mouse maintenance, breeding and timed mating. We acknowledge Sumaya Dauleh and Thomas P. Wilm for technical assistance.

Author Contributions: BW conceived and designed the study. SHA, HTN, LL, SL, SB, LC, MB, DAT and BW performed experiments and analysed the data. HTN analysed the scRNAseq datasets and interpreted it with DAT and BW. DAT and BW wrote the manuscript

with input from HTN and SHA. All authors discussed the results and commented on the
manuscript.

Funding:

BW was supported by MRC project grant MR/M012751/1 and the BBSRC Doctoral Training
Partnership scheme (BB/M011186/1 for MB; BB/T008695/1 for HTN), and acknowledges
generous funding from the Albaha University, Saudi Arabia (PhD Studentship to SHA). SB
was funded by an NHS Student Bursary. LL and LC were MRes students self-funding their
studies. DAT was supported in this work by a BBSRC New Investigator Grant
(BB/X000907/1; SL) and a David Sainsbury NC3Rs Fellowship (NC/P001467/1).

1. Abstract

The Wilms' Tumour protein (Wt1) had previously been shown to specify both the mesothelial lineage and vascular smooth muscle cell (vSMC) lineage in the mouse intestine, with evidence suggesting that intestinal vSMCs arise from the mesothelium. Here, we dissect the temporal relationship between these two mesodermal lineages, reporting that unexpectedly, Wt1 expression already in gastrulation stage embryos specifies a population of SMC precursor cells in the lateral plate mesoderm (LPM). Tamoxifen-induced genetic lineage tracing of Wt1-expressing cells revealed that vSMC and visceral smooth muscle cells (visSMCs) of the foetal mid-gut are labelled when Tamoxifen was administered at E7.5 or E8.5. By contrast, intestinal mesothelial cells were labelled after Tamoxifen dosing from E9.5 and later. Analysis of scRNAseq data of gastrulation stage mouse embryos and confocal microscopy demonstrated for the first time Wt1 expression in epiblast and primitive streak, emerging mesoderm and, from E7.5 onwards, in the LPM. In E8.5 mouse embryos, the co-expression of key smooth muscle fate markers *Tagln*, *Acta2*, *MrtfA/B* and *Pdgfrb* in *Wt1*-expressing cells revealed that vSMC and visSMC fate is specified independently from visceral mesothelium formation. Together, our study provides insight into the developmental lineage of smooth muscle specified by Wt1 expression at gastrulation stages.

2. Introduction

The embryonic origin of vascular smooth muscle cells (vSMCs) is relatively diverse as different regions of the vasculature arise from different progenitors including neural crest cells (NCCs), lateral plate mesoderm (LPM) and somatic mesoderm [1]. Differentiation of vSMCs and tissue formation in vertebrates are driven by a series of canonical cell signalling events and mechanical forces which regulate progenitor cell recruitment, differentiation, and maturation [2-4]. Platelet derived growth factor receptor beta (Pdgfrb) signalling is a key regulatory pathway in vSMC myogenesis since disruption of Pdgfrb signalling reduces vSMC proliferation and recruitment [4,5]. Specification of vSMC is marked by expression of smooth muscle cell markers such as transgelin (Tagln), alpha-smooth muscle actin (Acta2), smooth muscle myosin heavy chain (Myh11), desmin and others [6,7]. However, the timepoint when these genes start to be expressed in embryonic development, and their roles during the early stages of vSMC specification and differentiation remain unclear.

Wilms' tumour protein 1 (Wt1) is a transcription factor involved in the development of several tissues and organs in normal embryogenesis. The earliest Wt1 expression was identified in the intermediate mesoderm (IM) and LPM in E9.5 mouse embryos [8]. Its most prominent roles are in the development of the kidneys, arising from the IM, and of the heart, where the Wt1-expressing epicardium contributes to coronary vessel formation [9,10]. The epicardium is the mesothelial layer of the heart, which is continuous with the pericardial mesothelium of the pericardial cavity. Wt1 is also expressed in the pleural and peritoneal mesothelium, lining the cavities and covering the organs housed within, from about E9.5 onwards and throughout adulthood [8,11-15].

Previously, studies showed that the epicardium can give rise to vSMC of the developing coronary vasculature [16,17]. Genetic lineage tracing experiments using a transgenic mouse line expressing Cre recombinase under control of regulatory elements of the human WT1 gene (*Tg(WT1-cre)AG11Dbdr*, in short: *Wt1-Cre*) in combination with the *Rosa^LacZ/LacZ* reporter mouse revealed that a population Wt1-expressing serosa and pleural mesothelium cells give rise to vSMC in the peritoneal and lung cavity [13,15]. These results suggested that the mesothelium and vasculature of the intestine and lungs have a developmental relationship similar to that of the epicardium and the coronary vasculature. Similar studies using *mWt1/IRES/GFP-Cre* (*Wt1cre*) transgenic mice crossed with the *Rosa26R^EYFP* reporter mouse also found a contribution of Wt1-expressing mesothelial cells to the vasculature of the developing intestine [18].

In order to determine when mesothelial cells become committed to give rise to intestinal
vascular smooth muscle, we conducted pulse-chase studies of postnatal and adult mice
using a tamoxifen-inducible temporally-controlled and *Wt1*-driven Cre system
(*Wt1^tm2(cre/ERT2)Wtp*, *Wt1^CreERT2*)[17]. Our results showed that Wt1-expressing peritoneal cells
maintained the mesothelium over the intestine and peritoneal wall (and contributed to
visceral fat, [19]), but failed to give rise to intestinal vasculature components. However, Wt1-
expressing epicardium was observed to contribute towards formation of coronary vessels
postnatally and in adult mice [14]. These findings suggested that the capacity of Wt1-
expressing mesothelial cells to differentiate into vSMCs must be restricted to developmental
stages.

We propose that there is an embryonic contribution of Wt1-expressing cells towards the
vSMC fate in the peritoneum and intestine, of which the temporo-spatial resolution and
molecular mechanisms have remained unresolved so far. Here, to address this, we used
the tamoxifen-inducible *Wt1^CreERT2* mouse line in combination with Rosa-based reporter mice
for lineage tracing analysis to determine the contribution of Wt1-expressing cells towards
intestinal vasculature in mouse embryo. Our results indicate that during embryonic
development, two distinct populations of Wt1-expressing cells emerge, one which defines
mesothelial cells from approximately E9.5 onwards, and one which defines SMC fate
predominantly at stages around E7.5 to E8.5. Lineage-traced cells were found not only as
vSMCs but also as visceral SMCs (visSMCs) in the intestinal wall. We provide, for the first
time, a detailed analysis of Wt1 expression in E6.5-E8.5 mouse embryos. *In silico* analysis
of single cell RNAseq data revealed that *Wt1* is co-expressed with the emerging mesoderm
marker *Brachyury* (*Bra*) and later with SMC markers in E6.5-E8.5 mouse embryos. We then
describe expression of Wt1 in the E7.5 and E8.5 mouse embryo in the primitive streak and
the emerging neuro-mesodermal progenitor cells using cutting edge confocal microscopy
techniques. Lastly, we show that Wt1-derived cells within the mesentery of the developing
intestine aligned with the developing endothelial vascular network and co-expressed Pdgfrb.
Together, these data reveal a novel developmental differentiation pathway of visceral and
vascular smooth muscle cells influenced by Wt1 expression and independent of
mesothelium formation.

3. Results

Contribution of *Wt1*-expressing cells to SMCs in the peritoneal cavity.

To determine when during embryogenesis Wt1-expressing cells became fated to develop into vSMCs, we combined the *Wt1$^{CreERT2/+}$* lineage driver with either the *Rosa26R* (lacZ) or the *Rosa26^{mTmG}* lineage reporter in mice, gave tamoxifen (Tam) once at different embryonic stages (pulse), and performed endpoint analysis at stages around E18.5 (chase; see **Fig. 1A, Supplementary Fig. 1A** and **Supplementary Table 1**). We obtained similar results with both reporter systems: Tam administration at E9.5 or up to E13.5, followed by endpoint whole mount analysis, revealed almost exclusive labelling of mesothelial cells in the mesentery and intestine at mid-intestinal level (**Fig. 1B-E**, **Supplementary Fig. 1C-F**, **Supplementary Table 1**). When Tam was given to pregnant dams at E7.5 or E8.5 followed by endpoint analysis, we observed instead labelled cells surrounding the developing vessels of the mesentery, and in the visceral smooth muscle of the intestine at mid-intestinal level, but barely in the mesothelium (**Fig. 1F-J**; **Supplementary Fig. 1G-K**). Littermates carrying no *Wt1^{CreERT2}* allele showed no labelled cells (**Supplementary Fig. 1L, M**).

We also analysed the stomach and colon of foetuses under this experimental design, and observed that Tam dosing at E7.5 resulted in labelling of visceral smooth muscle cells but not mesothelial cells. By contrast, foetuses after Tam dosing at E8.5 showed a mixed population of labelled cells in the stomach, mostly consisting of visceral smooth muscle; in the colon, there were only labelled visceral smooth muscle cells. In foetuses which had received Tam at E9.5 up to E14.5, cell labelling occurred predominantly in mesothelial cells (**Supplementary Fig. 2**, **Supplementary Table 1**). However, after Tam dosing at E9.5 and up to E11.5 we observed in a few foetuses occasional visceral smooth muscle cells that were LacZ-positive, in the stomach and colon.

Interestingly, the spleen was not labelled when Tam was administered at E7.5, indicating that Wt1 was not expressed yet at this stage in the cells that later will become splenic precursors forming the putative splenic mesenchyme [20].

Figure 1: Embryonic contribution of cells of the Wt1-lineage to visceral mesothelium or visceral and vascular smooth muscle is dependent on timing of tamoxifen dosing. A. Schematic of the study design: CD1 dams were time mated with $Wt1^{CreERT2/+};Rosa26^{LacZ/LacZ}$ males and received tamoxifen (Tam) at the respective embryonic stages (embryonic day, E); embryos were dissected between E17.5 and E19.5, and XGal stained. **B.** When Tam was given at E13.5 and embryos analysed at E17.5, the mesentery (M) and intestine (I) were well covered by lacZ+ cells. **C, D.** When Tam had been given at E12.5, the coverage of LacZ-positive cells in the mesentery was less complete at E19.5. This loss of coverage with LacZ-positive cells was still more pronounced E17.5 when Tam had been given at E11.5 (D). **E.** Similarly, there were only patches of LacZ-positive cells in the mesentery and over the intestine at E17.5 after Tam administration at E9.5. **F, G.** When Tam had been given at E8.5, LacZ-positive cells were observed in small patches in the mesentery (arrow, F), or in cell chains near the vasculature (arrowhead, F), or in distinct rings (open arrowheads, G) surrounding the mesenteric veins (filled arrowhead, G) in the second arcade of the mesenteric vascular tree. The corresponding artery is depicted with an arrow (G). H-J. Tam dosing at E7.5 resulted at E18.5 in LacZ-positive cells scattered within the mesentery (H), in the circular musculature of the small intestine (white arrowheads, I), and surrounding a segment of mesenteric artery of the second arcade (black arrowheads, J). Scale bars, 500 µm (B, C, D, F, H), 200 µm (E, G, I, J).

Immunofluorescence staining on sections through intestine and mesentery of $Wt1^{CreERT2/+}$;

$Rosa26^{mTmG/+}$ foetuses confirmed that after inducing Tam-based lineage tracing at E8.5,

visSMCs and vSMCs co-expressed Acta2 and GFP and were found close to the CD31-

positive vascular endothelial cells, while hardly any mesothelial cells (pan-cytokeratin, Wt1)

were GFP-positive. By contrast, after Tam dosing at E11.5, mesothelial cells were GFP-

labelled but no visSMCs or vSMCs (**Fig. 2**).

These results suggest that during embryonic development, two distinct populations of Wt1-

expressing cells emerge, one which defines mesothelial cells from approximately E9.5

onwards, and one which defines visceral and vascular SMC fate predominantly at stages

around E7.5 to E8.5.

An additional observation in the mice after Tam dosing at E7.5 and E8.5, was that labelled

visSMCs and vSMCs could only be found in a short fragment of the mid-intestinal region

including mesentery (**Supplementary Fig. 1G**). This suggests a short pulse-like or transient

period of Wt1 expression during which Tam exposure allowed the recombination and

labelling of future visSMCs and vSMCs cells in the midgut region. However, the labelled

visSMC in the stomach and colon additionally indicate contribution along fore- and hindgut

regions not limited to initiation at E7.5-E8.5.

Figure 2: Immunofluorescence analysis of embryonic Wt1-lineage cells in intestine and mesentery. Frozen sections of intestine and mesentery of embryos after tamoxifen (Tam) dosing at E11.5 or E8.5, and dissection at E19.5 or E17.5, respectively) were analysed by immunofluorescence for co-expression of GFP with cytokeratin (A, A', E, E'), Wt1 (B, B', F, F'), CD31 (C, C', G, G') and Acta2 (D, D' H, H', I, I'). **A, A', B, B'.** After Tam at E11.5, the mesothelial markers cytokeratin and Wt1 were detected in the mesothelium covering the intestine and co-expression with GFP was observed in the visceral mesothelium (arrowheads). **E, E', F, F'.** After Tam at E8.5, co-expression of GFP with mesothelial markers was sparse (filled arrowhead). **C, C', D, D'.** After Tam at E11.5, expression analysis for the vascular markers CD31 and Acta2 showed no co-expression with GFP within the mesentery. **G, G'.** After Tam at E8.5, GFP-positive cells were found in close vicinity to CD31 expressed in the mesenteric vessels (arrowhead). **H, H', I, I'.** At this stage, GFP was co-expressed with Acta2 in the mesenteric vasculature (arrowhead, H, H'), and the visceral smooth muscle (arrowhead, I, I'). Scale bars, 50 μm (all).

Wt1 transcript is expressed from E6.5 in mouse embryos and co-expressed with markers for smooth muscle lineage

Next, we analysed expression of *Wt1* transcripts during mouse gastrulation stages E6.5 to E8.5 using a published single cell RNAseq data set [21]. We found that *Wt1* was expressed in mouse embryos throughout this developmental period in 0.65% to 1.66% of total embryonic cells (**Supplementary Fig. 3.1A**). Mean expression for *Wt1* was between 0.955 and 1.13 with cells considered expressing *Wt1* if levels were above 0 (**Supplementary Fig. 3.1B**). Most of the *Wt1*-expressing cells in the E7.5 embryo were found in extraembryonic ectoderm, the rostral neuroectoderm, mesenchyme, epiblast and primitive streak (**Supplementary Fig. 3.1C**). In E8.5 embryos, most *Wt1*-expressing cells were in the mesenchyme, surface ectoderm, pharyngeal mesoderm and cardiomyocytes (**Supplementary Fig. 3.1D**). Since visSMCs and vSMCs have their origin in the lateral plate mesoderm (LPM) but the dataset annotation lacked a clear indication which of the mesoderm tissues were LPM, we characterised in a candidate approach somatic, paraxial, nascent, mixed and intermediate mesoderm together with primitive streak, notochord, epiblast and mesenchyme for classical LPM and other mesodermal markers, confirming that 'mesenchyme' was representing the LPM population (**Supplementary Fig. 3.1E, F**).

Next, we selected *Wt1*-expressing cells from the epiblast, primitive streak, and mesodermal tissues at stages that mapped to gastrulation (E6.5-E8.5; **Fig. 3A**). We observed a shift from *Wt1*-expression mainly in the epiblast and primitive streak (E6.5 to E7.25) to expression predominantly in the 'mesenchyme'/LPM from E7.5 onwards (**Fig. 3A**). When we probed for *Wt1* and *Bra* co-expression (*Wt1+/Bra+*) in this cell population, we found that between E6.5 and E7.5, these cells were mostly found in the primitive streak, epiblast and nascent mesoderm. By contrast, from E7.5 onwards, there was a shift of *Wt1+/Bra+* co-expressing cells to the intermediate mesoderm, LPM, somitic mesoderm and neuro-mesodermal progenitor pool (NMP; **Fig. 3B**). Mean expression levels in *Wt1+ Bra+* co-expressing cells were higher for *Bra* (range 1.2 to 1.8) compared to *Wt1* (0.95-1.05; **Fig. 3C**). These data show that *Wt1* expression shifted from the emerging mesoderm to the LPM as gastrulation progressed. Furthermore, *Wt1* was co-expressed with gastrulation marker *Bra* in a few cells of key tissues of the forming mesoderm.

We then further interrogated the *Wt1*-expressing population (epiblast, primitive streak, mesodermal) for co-expression of LPM and smooth muscle markers, including *Hand1*, *Transgelin* (*Tagln*), *Acta2* and *Pdgfrb*. Our analysis found that *Wt1* and *Hand1* were co-

expressed already in the epiblast, primitive streak, nascent mesoderm and LPM at E7.0,

and from E7.5 onwards predominantly in the LPM (**Fig. 3D**). Mean expression levels of

Hand1 rose over the course of development to 3.5 at E7.5, while levels of *Wt1* were steady

at about 1 (**Fig. 3H**). Cells co-expressing *Wt1* and *Tagln* were already present in epiblast

cells at E6.5, before being found in primitive streak, nascent mesoderm and at later stages

predominantly in the LPM (**Fig. 3E**, **Supplementary Fig. 3.2A**). Mean expression levels of

Tagln rose over the course of development to 3 at E8.5, while levels of *Wt1* increased to

just above 1 (**Fig. 3I**). Similar patterns were observed for cells co-expressing *Wt1* and *Acta2*

(**Fig. 3F, J**) or *Wt1* and *Pdgfrb* (**Fig. 3G, K**), respectively, starting only to be co-expressed

from E7.5 onwards, but again predominantly in the LPM. Mean expression levels for *Wt1*

and *Acta2* in these cells followed a similar trajectory as for *Wt1* and *Tagln*, where *Acta2*

reached a level of 4, while *Wt1* rose to just above 1 (**Fig. 3J**). By contrast, the mean

expression levels of *Pdgfrb* and *Wt1* in co-expressing cells dropped and then increased

again from E7.5 to E8.5 (**Fig. 3K**). Mesodermal cells that co-expressed *Wt1*, *Tagln* and

Hand1 were found at E7.0 in the primitive streak, nascent mesoderm and LPM (**Fig. 3L**).

Cells that co-expressed *Wt1*, *Tagln* and *Acta2*, were identified almost exclusively in the LPM

from E7.5 to E8.5 (**Fig. 3M**). Furthermore, the trajectories of mean expression levels of these

markers and cell numbers of the triple-expressing cells were similar to those observed for

dual co-expressing cells (**Fig. 3L, M, N, O**), indicating that in the population of *Wt1*-

expressing cells at stages between E7.5 to E8.5 a considerable proportion consisted of

Hand1-, *Tagln*- and *Acta2*-co-expressing cells. This is confirmed in a Venn diagram showing

that on average 25 mesodermal cells (23.1%) in E8.5 embryos expressing *Wt1*, co-

expressed *Hand1*, *Acta2*, *Tagln* and *Pdgfrb* (**Fig. 3T**), indicating their commitment to a SMC

fate. Of all mesodermal cells that expressed *Hand1* and/or *Acta2* and/or *Tagln* and/or *Pdgfrb*

and <u>not </u>*Wt1*, only 9.2% expressed all 4 genes (**Fig. 3U**). It is important to note that by

selecting for *Wt1*-expression in the mesodermal population of *Hand1*/*Acta2*/*Tagln*/*Pdgfrb*-

expressing cells, cell numbers are smaller than in the mesodermal population of

Hand1/*Acta2*/*Tagln*/*Pdgfrb*-expressing cells that did not express *Wt1*. This suggests that

Wt1-expression is observed in a subpopulation of this group of cells, but that these cells

have a clear commitment along the SMC lineage.

These data suggest that a consistent population of *Wt1*-expressing cells arises from the

epiblast, via the primitive streak to the emerging mesoderm and the LPM which co-

expressed *Hand1* as LPM marker, and further key markers and regulators of smooth muscle

fate and differentiation. Importantly, levels of *Acta2* and *Tagln* expression rose during these

developmental stages also in cells that did not co-express *Wt1* (**Supplementary Fig. 3.2B**).

We further probed the relationship of *Wt1* expression in cells undergoing differentiation
towards smooth muscle by assessing the co-expression of *Wt1* with myocardin-related
factors *MrtfA* and *MrtfB* (**Fig. 3P, Q, R, S**). Both smooth muscle regulating genes were co-
expressed with *Wt1* in cells of the epiblast and primitive streak, shifting from E7.5
predominantly towards LPM cells. Mean expression levels of *Wt1* and *MrtfA* or *Wt1* and
MrtfB were very similar at levels between 0.5 and 1 (*Wt1*/*MrtfA*) and 0.7 and just above 1
(*Wt1*/*MrtfB*). These results indicate that *Wt1* is co-expressed with markers and regulators of
smooth muscle differentiation during stages before, during and immediately after mesoderm
formation and specification.

Figure 3: Distribution of *Wt1*-expressing cells in the mesoderm and presumptive smooth muscle cell lineage in gastrulation stage embryos. A-C. Distribution of *Wt1*-expressing cells from E6.5 to E8.5 together with early mesoderm marker, *Bra*, in mesodermal cells. *Wt1* transcripts were originally detected in epiblast and primitive streak cells and gradually (E7.5 to E7.75) shifted into cells of mesoderm lineage, predominantly lateral plate mesoderm and its derivatives. **D-K.** Distribution of *Wt1*-expressing cells in mesodermal tissues that co-expressed LPM and early SMC markers *Hand1*, *Tagln*, *Acta2* or *Pdgfrb*. *Wt1* transcripts were identified in epiblast and primitive streak cells that co-expressed *Hand1* at E7.0 (D, H), or *Tagln* at E6.5 (E, I), but from E7.5 onwards, these cells were mainly found in the LPM. Similar dynamics were observed for *Wt1*-expressing cells that co-expressed *Acta2* (F, J) or *Pdgfrb* (G, K). A small number of cells co-expressing *Wt1* and any of the SMC markers also contributed to intermediate mesoderm cells. **L-O.** Triple co-expression of *Wt1* with *Tagln* and *Hand1* or *Wt1* with *Tagln* and *Acta2* was predominantly detected in cells of the LPM, from E7.5 onwards. **P-S.** Co-expression of *Wt1* with the myocardin-related transcription factors (Mrtf) *MrtfA* and *MrtfB* in mesodermal tissues was found in small number of cells at E6.5, where expression was found in epiblast and primitive streak cells. Co-expressing cells were later detected in the lateral plate mesoderm, intermediate mesoderm and paraxial mesoderm. **T, U.** Venn diagram of mesodermal cells that expressed *Wt1* and co-expressed *Hand1*, *Acta2*, *Tagln* and/or *Pdgfrb* (T). Note that these numbers encompass the cells of the four E8.5 embryos that had been sequenced in (Pijuan-Sala et al. 2019). (U) shows the Venn diagram for mesodermal cells expressing *Hand1*, *Acta2*, *Tagln* and/or *Pdgfrb* when *Wt1*-expressing cells are not considered.

Our *in vivo* lineage tracing results indicated that Wt1 expression at between around E7.5 to

E8.5 primed a group of emerging mesoderm cells towards SM fate, but not mesothelial fate.

To determine whether mesothelial markers were co-expressed with *Wt1* during gastrulation

stages, we interrogated the scRNAseq data for co-expression of *Wt1* with mesothelial

progenitor marker *Hand2* (see also paralogous gene *Hand1* in **Fig. 3**), mesothelial-specific

markers *Podoplanin* (*Pdpn*), *Epicardin* (*Tcf21*), *Uroplakin 3b* (*Upk3b)* and *Tbx18*. Our results

showed that cells co-expressing *Wt1* and *Hand2* were mostly found in the LPM from E7.0

onwards, and expression levels of *Hand2* reached 1.4 (**Supplementary Fig. 3.2G, K**).

Onset of mesothelial gene expression in *Wt1*-expressing cells varied between E6.5 (*Pdpn*,

Supplementary Fig. 3.2D, H, L) and E8.25 (*Tbx18*, **Supplementary Fig. 3.2D, O, P**). *Tcf21*

and *Upk3b* started to be co-expressed in *Wt1*-expressing cells at E7.0 and E7.5,

respectively (**Supplementary Fig. 3.2 D, I, J, M, N**). In all cases, the majority of co-

expressing cells were found in the LPM. However, there was little overlap in the co-

expression between these mesothelial markers with *Wt1* in the E8.5 mouse embryo

(**Supplementary Fig. 3.2Q, R**). This suggests that mesothelial cell identity or fate was not

robustly established yet in *Wt1*-expressing mesodermal cells in the LPM in the E8.5 embryo.

Wt1 protein is expressed in the primitive streak and emerging mesoderm in E7.5 and

E8.5 mouse embryos

Next, we determined Wt1 protein expression in E7.5 and E8.5 mouse embryos using

confocal immunofluorescence microscopy of whole mount embryos, and found that Wt1 was

expressed in the tail bud of E8.5 embryos, as well as extra-embryonic tissues including the

yolk sac and allantois (**Fig. 4 A-C**). Co-immunostaining with Bra and confocal imaging

clearly identified Wt1-expressing cells in the Bra-domains of primitive streak and IM/LPM
region (**Fig. 4A, C, D-G**; compare **Supplementary Fig. 4** based on the EMAP eMouse Atlas
project [22]). Co-expression analysis of individual focal planes revealed that in the most
posterior region around the primitive streak, Wt1 expression was found in some Bra-positive
cells emerging from the node (**Fig. 4E**), the mesenchyme including neuromesodermal
progenitor cells and LPM/IM, the neural plate, primitive blood cells and allantois (**Fig. 4D-G;**
and insets). Quantification of co-expression of Wt1 and Bra in the insets of **Fig. 4E, F**
revealed that cells with strong Wt1 expression, also were positive for strong Bra expression
(**Fig. 4H**), confirming that Wt1 was co-expressed in Bra-positive cells emerging from the
mesoderm. Of note, the scRNAseq dataset suggested that very few cells contained
transcripts of both *Wt1* and *Bra*, while protein for both genes seemed to be detectable in
more cells.

At E7.5, the co-expression domain between Wt1 and Bra was more restricted, with Wt1
mostly expressed in the epiblast, where it was co-expressed with Bra, and in the primitive
streak (**Supplementary Fig. 5B**).

Further confocal image analysis of triple whole mount immunofluorescence for Wt1, Bra and
Tagln in E8.5 mouse tailbuds demonstrated protein co-expression of Wt1 with Tagln in the
mesoderm/paraxial mesoderm and LPM (**Fig. 4I-N** and **Supplementary Fig. 5A**). These
data support the observation that co-expression of Wt1 and Tagln in the emerging LPM
define the smooth muscle population.

Taken together, these data show that Wt1 is dynamically expressed in the gastrulating
embryo: in cells of the primitive streak and emerging mesoderm including the
neuromesodermal progenitor cells and LPM, where it is co-expressed with SMC markers.

Figure 4: Wt1 is expressed in the primitive streak, emerging mesoderm, NMP and LPM in E8.5 mouse embryos. A-D. Whole mount staining for Wt1 and Bra in the E8.5 mouse tail bud. A and B are MIP images generated from 150 z-sections, while C and D are focal images at sagittal level. Wt1 was expressed in the tail bud, in the allantois and yolk sac, as well as primitive streak, NMP, LPM/IM, where there was co-expression with Bra. In D (based on E), the different tissue domains are outlined. **E-G.** Images at two focal planes through the tail bud including node, primitive streak, notochord, NMP, LPM/IM and neural plate, in sagittal (slice 109, E), and near-sagittal (slice 268, F) position. Boxes in E and F indicate position of higher magnified inserts. Lines in E indicate position of focal views in G1-4. Arrows in G-14 indicate position of focal planes for E and F. **H.** Quantification of co-expression levels of both Wt1 and Bra at the two focal slices indicated shown in E and F and indicated in G1-4. AFU: arbitrary fluorescence units. **I.** Whole mount staining for Wt1, Bra and Tagln in the E8.5 mouse tail bud. Image of oblique focal plane; stipled line indicates focal plane for images J-N. **J.** LPM and notochordal plate (NCP) are outlined in transverse focal plane image. **K-M**. Expression of Wt1 (K), Bra (L) and Tagln (M). Wt1 and Tagln were co-expressed in the surface cells covering the paraxial mesoderm and in the LPM. Scale bars, 100 μm (A-E), 50 μm (I-N).

Cells arising from *Wt1*-lineage tracing after Tam dosing at E7.5 or E8.5 localise to three main tissues at E9.5: pro-epicardium, urogenital ridge and neural tube.

To obtain a better temporal resolution of the cells that had expressed Wt1 at gastrulation stages for lineage tracing of SMCs, we decided to repeat the E7.5 and E8.5 Tam dosing and observe the range and localisation of GFP-positive cells at the earlier E9.5 time-point (**Fig. 5**).

When we administered Tam at E8.5 and analysed at E9.5, abundant GFP-positive cells where localised to the pro-epicardium at the caudal side of the forming heart (**Supplementary Fig. 6A-D**), and to the urogenital ridge (**Fig 5**, **Supplementary Fig. 6A, B, C'**). We observed GFP-positive cells consistently towards the ventral midline, including the mesentery, in sections, and by using light sheet microscopy of the trunk region of the E9.5 embryo (**Fig. 5**, **Supplementary Fig. 7**, **Supplementary Movie, Supplementary Table 2**). By contrast, Tam dosing at E7.5 resulted in less GFP-positive cells, which were mostly found in the pro-epicardium (**Supplementary Fig. 6E-G**). In three litters analysed, we found that all 15 embryos had between 2 and 20 GFP-positive cells in the pro-epicardium, while GFP-positive cells (range 1-12) in the posterior trunk region were observed only in 9 out of 15 embryos (**Supplementary Table 3**, **Supplementary Fig. 8**). Importantly, we also detected GFP-positive cells in the neural tube of embryos after Tam at E7.5 (**Supplementary Fig. 6H-K**) and E8.5 (not shown), in line with our observations in **Figs. 3** and **4** that Wt1 was also expressed in the emerging neuroectoderm.

Together, these results demonstrate that the $Wt1^{CreERT2/+}$ lineage tracing system detected cells that expressed Wt1 at around E7.5 and E8.5 within a similar spectrum of cell progeny, but GFP-positive cell numbers where smaller when Tam was given at E7.5. This was in agreement with the scRNAseq data analysis (**Fig. 3**) and distribution of Wt1-expressing cells by whole mount immunofluorescence (**Fig. 4**). Moreover, we could identify GFP-positive cells that appeared in the region of the mesentery, as potential precursors for visceral and vascular smooth muscle of the developing intestinal tract.

Figure 5: Localisation of Wt1-lineage cells in the urogenital ridge and close to the mesentery near endothelial cells. A, B. Immunofluorescence staining for GFP and Wt1 (D) and GFP and CD31 (E) on sections through E9.5 embryos after Tam dosing at E8.5. GFP-positive cells were detected in the region between dorsal aorta (da) and mesentery, besides presence in the urogenital ridge (ur). **C.** Representative overview image of embryo at E10.5 after Tam dosing at E8.5. GFP-positive cells were most prominent in the heart (ht), and urogenital ridge (ur). Arrowheads point to individual cells dorsally away from the main ur expression domain. **D, D'.** Immunofluorescence staining for GFP and SMA on sections through the region just below the forelimb bud. GFP-positive cells were localized in the urogenital ridge and close to the dorsal aorta (da)(arrowheads). **E, E'.** Immunofluorescence staining for GFP and CD31 on section through similar embryonic level as in D. GFP-positive cells were also found ventral to the dorsal aorta and dorsal to the mesentery, closely associated with cells of the endothelial plexus (filled arrowhead: GFP+, open arrowhead: CD31+). da, dorsal aorta, e, endoderm, flb, forelimb bud, hlb, hindlimb bud, ur, urogenital ridge. Scale bars, 1000 μm (A), 500 μm (B), 250 μm (B'), 100 μm (C, C', D), 50 μm (E).

Cells of the *Wt1*-lineage contribute to the development of the vascular and visceral smooth muscle between E10.5 and E15.5

To elucidate the developmental path and fate of the vSMCs and visSMCs using the emergence of the GFP-positive cells, we analysed embryos at E10.5 after Tam dosing at E8.5. At this stage, GFP-positive cells were predominantly seen in the region of the urogenital ridge and the heart (**Fig. 5C**). In tissue sections, GFP-positive cells localised in the region between the dorsal aorta and the ventral mesentery, and within the urogenital ridge (**Fig. 5D, D'**). Some of the cells within the region between dorsal aorta and ventral mesentery appeared to be situated close to endothelial cells (**Fig. 5E, E'**).

In embryos at E12.5, we found GFP-positive cells in a spotty appearance within the mesentery of the developing small intestinal segment of the intestinal tract (**Fig. 6B, B'**). Immunofluorescence analysis on sections through this region revealed that GFP-positive cells were localised within the mesentery and in the developing intestinal wall near the forming vascular plexus (**Fig. 6C, D**). GFP-expressing cells were distinct from endothelial cells but closely associated (**Fig. 6D**). Furthermore, our analysis showed that some of the GFP-labelled cells within the mesentery and the intestinal wall co-expressed Pdgfrb, suggesting that they had started the smooth muscle differentiation programme (**Fig. 6E, F-F''**). Occasionally, we found a few GFP-positive cells in the mesothelial covering of the mesentery (**Fig. 6E, F**). Together, these results indicate that *Wt1*-lineage traced cells labelled at around E8.5, appeared in the vicinity of the developing vasculature and co-expressed Pdgfra, a key marker of vascular smooth muscle progenitor cells during stages of intestinal patterning and angiogenesis. When endpoint analysis was performed at E15.5, we detected GFP-positive cells within the visceral smooth muscle and surrounding mesenteric vessels (**Fig. 6G-H**), similar to findings from the endpoint analysis at around E18.5 (**Fig. 1I**).

Figure 6. Contribution of *Wt1*-lineage cells to the mesentery near developing vasculature and to the intestinal wall. A. Schematic of the study design: analysis of embryos at E12.5 or E15.5 after time mating of *Wt1$^{CreERT2/+}$;Rosa26$^{mTmG/mTmG}$* males with CD1 females and Tam dosing at E8.5. **B, B'.** View of the intestinal loop and mesentery. GFP-positive cells (chevrons) were found within the mesentery of the developing intestinal tract (arrow; star depicts caecum). Note the abundant GFP fluorescence in the liver (Li). **C-D.** Immunofluorescence of CD31 and GFP on sections through this region showed GFP-positive cells within the mesentery and in the intestinal wall, near the developing microvascular plexus (C). Higher magnification (D) reveals that GFP-positive cells (chevrons) are often arranged close to neighbouring endothelial cells (arrows) of the vascular plexus in the intestinal wall and mesentery. **E-F''.** Immunofluorescence for Pdgfrb and GFP on neighbouring sections showed that GFP-positive cells within the intestinal wall mesenchyme and mesentery co-expressed Pdgfrb (chevrons), while a few mesothelial GFP-positive cells were Pdgfrb-negative (F-F'', arrow). **G, H.** Representative images from two different embryos at E15.5 after tamoxifen dosing at E8.5. GFP-positive cells (arrowheads) were found within a segment of the small intestine in a circular arrangement. da, dorsal aorta, e, endoderm, m, mesentery, ma, mesenteric artery, mv, mesenteric vein. Scale bars, 500 μm (G, H), 300 μm (B, B'), 100 μm (C, E, F-F''), 50 μm (D).

Discussion

In this study, we have for the first time determined the temporal and spatial relationship between the emerging intestinal vascular and visceral smooth muscle cells and the expression of the transcriptional regulator Wt1 during gastrulation and post-gastrulation stages of mouse embryonic development. Our analysis revealed that Wt1 is expressed in a progenitor cell pool in the emerging mesoderm of the gastrulating embryo. Some of these cells differentiate into smooth muscle cells in the intestine and mesentery. Our data indicate that this process is temporally independent of the formation of the mesothelium covering these tissues, which also expresses Wt1. Thus, our data shed new light onto smooth muscle development independent from mesothelium development, and the temporo-spatial involvement of Wt1 in this process (**Supplementary Fig. 9**).

Here, using Tam-dependent, inducible genetic lineage tracing, we analysed the embryonic contribution of Wt1-expressing cells to visSMCs and vSMCs of the intestine. In previous studies, we had shown that Wt1-expesing cells give rise to vSMCs in the mesentery and intestine using a non-inducible Wt1-based lineage tracing system [13]. Similarly, by using the same or an alternative Wt1-driven genetic reporter system, it had been demonstrated that Wt1-expressing progenitor cells give rise to visSMCs and vSMCs not only in the intestine but also in the lungs [15,18,23], however, the temporal involvement and spatial significance of Wt1 expression during smooth muscle specification and differentiation had not been determined.

In a follow-on study, we determined that Wt1-expressing cells in postnatal stages fail to give rise to intestinal visSMCs and vSMCs, by making use of the same Tam-dependent temporal activation process for lineage analysis as utilised here (Wilm et al., 2021). We concluded that the source of intestinal and mesenteric SMCs had to be cells expressing Wt1 during embryonic development; however, it was not clear whether these cells arise from the overlying mesothelium which also expresses Wt1 during embryonic development, as had been our previous hypothesis [13].

To address this question, we administered Tam between E7.5 and E13.5 using the *Wt1^CreERT2/+* driver in combination with either the *Rosa26^LacZ* or *Rosa26^mTmG* reporter systems, and analysed the presence of LacZ- and GFP-labelled cells in foetuses just before birth (E17.5, E18.5 or E19.5). Our analysis showed that Tam dosing at E7.5 or E8.5 resulted in labelling of vSMC and visSMCs which had originated from Wt1-expressing cells. When

Tam was given at E9.5 or later stages, the labelled cells were confined to the mesothelial covering of the mesentery and intestine. This finding demonstrates a clearly defined separate mechanism of the emergence of the mesothelial lineage, from the vascular/visceral smooth muscle lineage, indicating a role for Wt1 in distinct mechanisms.

Other studies have previously used the Tam-inducible $Wt1^{CreERT2/+}$ driver in combination with reporter systems to elucidate the lineage and origin of smooth muscle cells in the lungs. Contribution to airway smooth muscle, vascular smooth muscle and parenchyma was observed in late-stage mouse foetuses when Tam had been given between E9.5 and E11.5 [24,25]. Time lapse experiments of *ex vivo* cultured lungs from lineage-traced E12.5 embryos after Tam at E10.5 revealed a direct contribution of mesothelial cells towards the deeper lung parenchyma [24]. These results suggest that the smooth muscle of the airways and lung vasculature may have a contribution via the mesothelium at later developmental stages, in a different mechanism to the one described here for the intestinal tract.

In the heart, Wt1-based lineage tracing had demonstrated contribution of epicardial cells to coronary vessel formation [13,17,26,27]. Based on *in situ* hybridisation and immunofluorescence on sections of E8.5 embryos, Wt1 was not expressed in the developing heart at this stage. By contrast, non-inducible $Wt1^{CreEGFP}$;$Rosa26^{mTmG}$ embryos at E8.5 had a number of GFP-expressing cells in the forming heart, indicating an early contribution of Wt1-expressing cells [26]. However, Wt1 is also expressed in non-epicardial cells in the heart, including myocardium, which makes interpretation of lineage experiments more challenging [26]. In addition, our own post-natal lineage tracing experiments using the Tam-inducible $Wt1^{CreERT2/+}$ driver had shown that after birth Wt1-expessing cells contribute to coronary SMCs [14], suggesting a different mechanism of coronary angiogenesis and maintenance of coronary vasculature than in the intestinal tract and mesentery.

The Tam-inducible lineage tracing system is dependent on the Tam-driven recombination activity which then irreversibly switches on the reporter gene expression. Several studies have analysed the temporal delay between Tam administration and reporter expression, indicating that depending on route of administration (oral gavage or IP injection), a temporal delay of between 12-18 hours could be observed in other genetic systems [28,29]. This temporal delay needs to be factored in when interpreting the results of the current study, suggesting that about 0.5 embryonic days may have to be added to the stages indicated for the recombination initiation.

One of the surprising findings of this study emerged from our analysis of scRNAseq data [21]

showing that from E6.5 onwards, *Wt1* is expressed in the epiblast, primitive streak and

emerging mesoderm, and in the LPM where it is co-expressed with key markers of SMC

development. We could confirm co-expression of Wt1 protein with Bra in the primitive

streak/emerging mesoderm, and with the key SM marker Tagln in the LPM.

Analysis of Wt1 expression in early stage mouse embryos had been previously performed

by in situ hybridisation showing *Wt1* transcripts in the intermediate mesoderm/LPM in E9.0

mouse embryos [8]. Because further insights into Wt1 expression or function at gastrulation

stages have not been published yet, our results are the first to shed light onto Wt1

gastrulation-stage expression or engagement. We have identified Wt1 expression in epiblast

and primitive streak cells in embryos up to about E7.25, and then a shift to specifically

identified mesodermal cells in E8.5 embryos where they are predominantly present in the

LPM. Our results reveal critical expression of Wt1 in a progenitor pool of SMCs, which may

represent the precursors of the visSMCs and vSMCs labelled in the lineage tracing studies.

With these results we have demonstrated that Wt1-expressing cells unexpectedly emerge

already in gastrulation stage embryos as the precursors for intestinal visSMC and vSMC,

providing a direct link for Wt1 with SMC fate.

The peritoneal mesothelium emerges due to separation of the LPM into somatic and

splanchnic mesoderm, which surround the peritoneal cavity; the splanchnic mesoderm

forms the mesodermal component of the intestinal tract [30]. Detailed analysis of the

emergence of the mesothelium has been provided in chick and quail embryo models [31,32],

demonstrating that the bilateral splanchnopleure consisting of splanchnic mesoderm and

endoderm, turns into a trilaminar tissue with a mesenchymal layer emerging between an

outer mesodermal epithelium and the endoderm. While in the chick embryo, the outer

epithelium is situated on a basal lamina already at HH13 (~19 somites, 48 hours),

cytokeratin expression as indicator for epithelium, starts only at HH19 (~40 somites, 71

hours; roughly corresponding to mouse embryos at Theiler stage 17 or E10.5), and Wt1 is

not expressed at this stage yet [32]. The outer epithelium is defined as mesothelium only from

HH29 (6 days) onwards (chick and quail), when it has organised into a simple squamous

epithelium [31,32].

In zebrafish, *hand2* expression has been presented as a marker for mesothelial progenitor

cells emerging from the lateral most region of the LPM at tailbud stage (i.e. at late

gastrulation stages) [33]. Co-expression with Tagln in the LPM-emerging cells, and a more

limited expression of *wt1a/b*, that was restricted to the cells of the anterior visceral

peritoneum, and absent posteriorly and in the parietal mesothelium, suggested that *hand2*

was a more encompassing marker for arising mesothelium. Furthermore, *hand2* mutant

zebrafish had disorganised LPM-derived cells, in particular a lack of parietal mesothelium,

indicating an upstream regulatory role of *hand2* in mesothelium formation [33].

In the mouse embryo, the peritoneal mesothelium has been shown to arise between E10.5

and E11.5 since at E11.5, a continuous layer of cells expressing cytokeratin and Wt1 is

present [13]. Further information on mesothelial precursor populations and lineage in mouse

embryonic development is limited: non-inducible Hand2 or Hand1-based lineage tracing

showed labelling of mesothelium over the liver in E14.5 or E15.5 mouse embryos but no

temporal analysis [33]. Non-inducible Hand1-lineage has also been shown to label the visceral

smooth muscle in mouse embryos at E14.5 [34]. Therefore, it is not clear whether in the

mouse, Hand1 or Hand2 can be considered markers of the mesothelial precursor

population.

Visceral smooth muscle arises from around E13 in the mouse embryo based on Acta2

immunofluorescence [13,35], while the vascular smooth muscle in the intestine and mesentery

can be detected from E12.5 during angiogenic remodelling, with mature vessels covered by

mural cells, detectable from E15.5 onwards [13,36]. A complex signalling network including the

Myocardin/serum response factor (SRF) interaction controls SMC differentiation by

regulating expression of cytoskeletal SM proteins including Acta2, Transgelin and smooth

muscle myosin heavy chain. Furthermore, Pdgfrb signalling regulates vSMC proliferation,

migration and differentiation [1]. In this study, we have shown that some of the key regulators

and markers of SMC fate and differentiation are co-expressed with Wt1 already from E7.5

onwards in the LPM. In addition, at E12.5, Wt1-lineage traced cells in the developing

intestinal /mesenteric vascular network co-expressed Pdgfrb after activation of the reporter

at around E8.5. These results strongly suggest a role for Wt1 in smooth muscle

development, originating in cells of the emerging mesoderm. Furthermore, our analysis of

Wt1-lineage tracing at earlier stages (E9.5, E10.5) of development, suggest that the Wt1-

lineage traced cells originating at E8.5, may migrate into the mesentery and towards the

developing vascular network.

Taken together, our data clearly demonstrate that the visSMCs and vSMCs that arose from

Wt1-expressing cells at around E7.5 or E8.5, are independent in their specification and

differentiation from the mesothelial cells of the peritoneum. These can only be tagged after

initiation of Wt1-controlled recombination from around E9.5 onwards. These results indicate

an early emergence of visSMC and vSMC lineage in the gastrulating embryo, and an

uncoupling from the mesothelial lineage. The role of Wt1 in both processes remains to be

elucidated.

Mice:

The following compound mutants were used in this study and bred in-house: $Wt1^{CreERT2/+}$; $Rosa26^{LacZ/LacZ}$ ($Wt1^{tm2(cre/ERT2)Wtp}$; $Gt(ROSA)26Sor/J)^{17,37}$, $Wt1^{CreERT2/+}$; $Rosa26^{mTmG/+}$ ($Wt1^{tm2(cre/ERT2)Wtp}$; $Gt(ROSA)26Sor^{tm4(ACTB-tdTomato,-EGFP)Luo}/J)^{38}$. CD1 female mice (at least 10 weeks old, Charles River, Harlow, UK) were time-mated with $Wt1^{CreERT2/+}$;$Rosa26^{LacZ/LacZ}$ or $Wt1^{CreERT2/+}$;$Rosa26^{mTmG/+}$ males, and when a vaginal plug was detected, noon of the day was considered as embryonic day 0.5 (E0.5). CD1 female mice time-mated by Charles River were ordered in. Mice were housed in individually ventilated cages under a 12-hour light/dark cycle, with *ad libitum* access to standard food and water. All animal experiments were performed under a Home Office licence granted under the UK Animals (Scientific Procedures) Act 1986 and were approved by the University of Liverpool AWERB committee. Experiments are reported in line with the ARRIVE guidelines.

Tamoxifen dosing

For Wt1 lineage tracing in embryos, pregnant CD1 mice after mating with $Wt1^{CreERT2/+}$; $Rosa26^{LacZ/LacZ}$ or $Wt1^{CreERT2/+}$; $Rosa26^{mTmG/+}$ male mice were dosed with 100 μg/g body weight of Tam (T5648, SigmaAldrich; 40 mg/ml, in corn oil (C8267, SigmaAldrich)) via oral gavage once on various days and analysed between E9.5 and E19.5. Pregnant dams were monitored for their well-being including daily weight checks.

XGal staining

Dissected embryos or tissues were fixed in 2% paraformaldehyde (PFA)/0.25% glutaraldehyde in phosphate buffered saline (PBS) for between 1 and 1.5 hours, whole-mount XGal staining performed overnight according to standard protocols, followed by overnight post-fixation in 4% PFA (PBS) at 4°C.

Immunofluorescence

The following primary antibodies were used: anti-Wt1 rabbit polyclonal (1:200 to 1:500, clone C-19, sc-192, Santa Cruz), anti-Wt1 mouse monoclonal (1:50, clone 6F-H2, M3561, Dako), anti-SMA mouse monoclonal (1:100 to 1:200, clone 1A4, A2547, Sigma), anti-Pecam/CD31 rat monoclonal (1:50, 550274, Pharmingen), anti-Brachyury goat polyclonal (1:200, PA5-46984, Invitrogen/ThermoScientific), anti-Tagln/Sm22 mouse monoclonal (1:100, 60213-1-

Ig, Proteintech) anti-GFP rabbit or goat polyclonal (1:5000, ab6556 or ab6673, Abcam). The
anti-SMA and anti-Tagln/Sm22 antibodies were directly labelled using Zenon direct labelling
kit (Invitrogen/ThermoScientific) according to manufacturer's instructions. Secondary
antibodies were Alexa fluorophore-coupled (Invitrogen/ThermoScientific) and were used at
a dilution of 1:1000. DAPI (D9542, SigmaAldrich) was used at 1:250-1000 for nuclear
counterstaining.

Frozen sections: Tissues were fixed in 4% PFA for between 30 to 90 mins, protected in 30%
sucrose overnight, placed in Cryomatrix (Thermo Scientific) and snap frozen. Frozen
sections were generated at 8-10 µm on a Thermo Scientific HM525 NX Cryostat.
Immunofluorescence (IF) staining was performed following standard protocols [13]. Two
bleaching steps of 20 min each in 3% H_2O_2/MeOH were included for embryos or their tissues
carrying the Rosa26$^{mTmG/+}$ mutation in order to remove the tdTomato fluorescence [39].
Subsequently to completed IF, sections were coverslipped with Fluoro-Gel (with Tris buffer;
Electron Microscopy Sciences, USA), sealed and stored in the dark at 4°C.

Whole tissues and embryos: Whole tissues/embryos were fixed in 4% PFA for 60 minutes
on ice and stored in 0.0025% Triton X-100 (Sigma Aldrich) in PBS at 4°C until used.
Tissues/embryos were permeabilised using 0.25% Triton X-100 (PBS) for 60 minutes on
ice. IF staining was performed using previously published protocols [40-42]. E7.5 or E8.5 IF-
stained embryos were mounted on coverslips within SecureSeal imaging spacers (Grace
Bio-Labs) and embedded in RapiClear 1.52 solution (SUNJin Lab). Samples were stored in
the dark at 4°C before imaging.

Microscopy

Epifluorescence microscopy: IF sections were imaged on a Leica DM 2500 upright
microscope with a Leica DFC350 FX digital camera and a Leica EL6000 fluorescence light
source supported by the Leica Application Suite software package (LAS, version 3 or 4;
Leica Microsystems, Germany/Switzerland).

Whole mount microscopy: XGal-stained and GFP-positive embryos and tissues were
imaged using a Leica MZ 16F dissecting microscope equipped with a Leica DFC420 C
digital camera and a Leica EL6000 fluorescence light source supported by the LAS.

Confocal microscopy: Imaging of whole mount embryos was performed using an Andor
Dragonfly spinning disc confocal microscope (Andor, Oxford Instruments) mounted on an
inverted Leica DMi8 base using either a 10x (0.45NA) or a 25x water-dipping objective

(NA0.95). The z-steps for the 10x and 25x objectives were 1.32 μm and 0.35 μm,
respectively. DAPI, Alexa- 488, −568 and −647 were sequentially excited using 405nm 561,
637nm laser diodes, respectively, and emitted light reflected through 450/50 nm, 525/50nm,
600/50nm and 700/25nm bandpass filters, respectively. An iXon 888 EM-CCD camera was
used to collect emitted light, and data were captured using Fusion version 5.5 (Bitplane) with
subsequent image analysis performed using either Fiji [43] or Imaris (Bitplane). Images were
manually quantified by circling visible nuclei within a single z-plane to generate regions of
interest (ROI) using the ROI manager tool in FiJi. The average grey values were then
measured for the selected regions for each fluorescent channel.

Light sheet microscopy: The unfixed trunk area of E9.5 *Wt1$^{CreERT2/+}$; Rosa26$^{mTmG/+}$* embryos
dosed with tamoxifen at E8.5, was mounted in 1% low melting point agarose (ROTI Garose,
Carl Roth, Karlsruhe, Germany), transferred into a capillary (Size 4, Brand GmbH,
Wertheim, Germany) and placed in the imaging chamber of a Zeiss Light Sheet Z.1
microscope with one 5x detection objective and two 5x illumination objectives supported
with the Zen software (Carl Zeiss, Jena, Germany). Imaging was performed using the 488
nm channel to capture GFP-positive cells, and the 561 nm channel to image membrane-
bound dTomato to provide tissue context. Images were adjusted to colours using the Imaris
software and the movie edited to highlight the shape of the selected area by using
the surface tool, and by labelling GFP-positive cells with coloured spots to indicate their
different locations.

Analysis of single-cell mouse gastrulation atlas.

Data analysis from a previously published dataset [21] was performed using R (version 4.3.2).
Data were accessed through Bioconductor 'MouseGastrulationData' package (version
1.16.0), and only processed data were extracted for downstream analysis. Extracted data
were normalised using *logNormCounts* function from 'scuttle' package (version 1.12.0).
Normalisation of cell count per sample was carried out by dividing total number of cells by
sample size and rounding up to the nearest integer. All plots were created using 'ggplot2'
package (version 3.4.3).

4. References

Donadon, M. & Santoro, M. M. The origin and mechanisms of smooth muscle cell development in vertebrates. *Development* **148**, doi:10.1242/dev.197384 (2021).

Monte, G. d. *et al.* Differential Notch Signaling in the Epicardium Is Required for Cardiac Inflow Development and Coronary Vessel Morphogenesis. *Circulation Research* **108**, 824-836, doi:doi:10.1161/CIRCRESAHA.110.229062 (2011).

Qi, L., Kong, W., Fu, Y., Fu, X. & Xu, T. Vascular Smooth Muscle Cell Development and Cardiovascular Malformations. *Cardiology Discovery* **01**, 259-268, doi:doi:10.1097/CD9.0000000000000035 (2021).

Stratman, A. N. *et al.* Chemokine mediated signalling within arteries promotes vascular smooth muscle cell recruitment. *Communications Biology* **3**, 734, doi:10.1038/s42003-020-01462-7 (2020).

Hellstrom, M., Kalen, M., Lindahl, P., Abramsson, A. & Betsholtz, C. Role of PDGF-B and PDGFR-beta in recruitment of vascular smooth muscle cells and pericytes during embryonic blood vessel formation in the mouse. *Development* **126**, 3047-3055 (1999).

Smyth, L. C. D. *et al.* Markers for human brain pericytes and smooth muscle cells. *Journal of Chemical Neuroanatomy* **92**, 48-60, doi:https://doi.org/10.1016/j.jchemneu.2018.06.001 (2018).

Assinder, S. J., Stanton, J. A. & Prasad, P. D. Transgelin: an actin-binding protein and tumour suppressor. *Int J Biochem Cell Biol* **41**, 482-486, doi:10.1016/j.biocel.2008.02.011 (2009).

Armstrong, J. F., Pritchard-Jones, K., Bickmore, W. A., Hastie, N. D. & Bard, J. B. The expression of the Wilms' tumour gene, WT1, in the developing mammalian embryo. *Mech Dev* **40**, 85-97 (1993).

Kreidberg, J. A. *et al.* WT-1 is required for early kidney development. *Cell* **74**, 679-691, doi:https://doi.org/10.1016/0092-8674(93)90515-R (1993).

Rackley, R. R. *et al.* Expression of the Wilms' tumor suppressor gene WT1 during mouse embryogenesis. *Cell Growth Differ* **4**, 1023-1031 (1993).

Moore, A. W. *et al.* YAC transgenic analysis reveals Wilms' tumour 1 gene activity in the proliferating coelomic epithelium, developing diaphragm and limb. *Mech Dev* **79**, 169-184 (1998).

Moore, A. W., McInnes, L., Kreidberg, J., Hastie, N. D. & Schedl, A. YAC complementation shows a requirement for Wt1 in the development of epicardium, adrenal gland and throughout nephrogenesis. *Development* **126**, 1845-1857, doi:10.1242/dev.126.9.1845 (1999).

Wilm, B., Ipenberg, A., Hastie, N. D., Burch, J. B. E. & Bader, D. M. The serosal mesothelium is a major source of smooth muscle cells of the gut vasculature. *Development* **132**, 5317-5328, doi:10.1242/dev.02141 (2005).

Wilm, T. P. *et al.* Restricted differentiative capacity of Wt1-expressing peritoneal mesothelium in postnatal and adult mice. *Scientific reports* **11**, 15940, doi:10.1038/s41598-021-95380-1 (2021).

Que, J. *et al.* Mesothelium contributes to vascular smooth muscle and mesenchyme during lung development. *Proc Natl Acad Sci U S A* **105**, 16626-16630, doi:10.1073/pnas.0808649105 (2008).

Guadix, J. A., Carmona, R., Munoz-Chapuli, R. & Perez-Pomares, J. M. In vivo and in vitro analysis of the vasculogenic potential of avian proepicardial and epicardial cells. *Dev Dyn* **235**, 1014-1026, doi:10.1002/dvdy.20685 (2006).

Zhou, B. *et al.* Epicardial progenitors contribute to the cardiomyocyte lineage in the developing heart. *Nature* **454**, 109-113, doi:10.1038/nature07060 (2008).

18   Carmona, R., Cano, E., Mattiotti, A., Gaztambide, J. & Munoz-Chapuli, R. Cells
derived from the coelomic epithelium contribute to multiple gastrointestinal tissues in
mouse embryos. *PloS one* **8**, e55890, doi:10.1371/journal.pone.0055890 (2013).
19   Chau, Y. Y. *et al.* Acute multiple organ failure in adult mice deleted for the
developmental   regulator   Wt1.   *PLoS   Genet*   **7**,   e1002404,
doi:10.1371/journal.pgen.1002404 (2011).
20   Hecksher-Sorensen, J. *et al.* The splanchnic mesodermal plate directs spleen and
pancreatic laterality, and is regulated by Bapx1/Nkx3.2. *Development* **131**, 4665-
4675, doi:10.1242/dev.01364 (2004).
21   Pijuan-Sala, B. *et al.* A single-cell molecular map of mouse gastrulation and early
organogenesis. *Nature* **566**, 490-495, doi:10.1038/s41586-019-0933-9 (2019).
22   Richardson, L. *et al.* EMAGE mouse embryo spatial gene expression database: 2014
update. *Nucleic Acids Res* **42**, D835-844, doi:10.1093/nar/gkt1155 (2014).
23   Cano, E., Carmona, R. & Munoz-Chapuli, R. Wt1-expressing progenitors contribute
to multiple tissues in the developing lung. *Am J Physiol Lung Cell Mol Physiol* **305**,
L322-332, doi:10.1152/ajplung.00424.2012 (2013).
24   Dixit, R., Ai, X. & Fine, A. Derivation of lung mesenchymal lineages from the fetal
mesothelium requires hedgehog signaling for mesothelial cell entry. *Development*
140, 4398-4406, doi:10.1242/dev.098079 (2013).
25   Moiseenko, A. *et al.* Origin and characterization of alpha smooth muscle actin-
positive cells during murine lung development. *Stem cells (Dayton, Ohio)* **35**, 1566-
1578, doi:10.1002/stem.2615 (2017).
26   Rudat, C. & Kispert, A. Wt1 and epicardial fate mapping. *Circulation research* **111**,
165-169, doi:10.1161/CIRCRESAHA.112.273946 (2012).
27   Quijada, P., Trembley, M. A. & Small, E. M. The Role of the Epicardium During Heart
Development   and   Repair.   *Circulation   research*   **126**,   377-394,
doi:10.1161/CIRCRESAHA.119.315857 (2020).
28   Nguyen, M. T., Zhu, J., Nakamura, E., Bao, X. & Mackem, S. Tamoxifen-dependent,
inducible Hoxb6CreERT recombinase function in lateral plate and limb mesoderm,
CNS isthmic organizer, posterior trunk neural crest, hindgut, and tailbud. *Dev Dyn*
238, 467-474, doi:10.1002/dvdy.21846 (2009).
29   Zhu, J. *et al.* Uncoupling Sonic hedgehog control of pattern and expansion of the
developing   limb   bud.   *Developmental   cell*   **14**,   624-632,
doi:10.1016/j.devcel.2008.01.008 (2008).
30   Funayama, N., Sato, Y., Matsumoto, K., Ogura, T. & Takahashi, Y. Coelom formation:
binary decision of the lateral plate mesoderm is controlled by the ectoderm.
Development **126**, 4129-4138, doi:10.1242/dev.126.18.4129 (1999).
31   Thomason, R. T., Bader, D. M. & Winters, N. I. Comprehensive timeline of
mesodermal development in the quail small intestine. *Dev Dyn* **241**, 1678-1694,
doi:10.1002/dvdy.23855 (2012).
32   Winters, N. I., Thomason, R. T. & Bader, D. M. Identification of a novel developmental
mechanism in the generation of mesothelia. *Development* **139**, 2926-2934,
doi:10.1242/dev.082396 (2012).
33   Prummel, K. D. *et al.* Hand2 delineates mesothelium progenitors and is reactivated
in mesothelioma. *Nat Commun* **13**, 1677, doi:10.1038/s41467-022-29311-7 (2022).
34   Barnes, R. M., Firulli, B. A., Conway, S. J., Vincentz, J. W. & Firulli, A. B. Analysis of
the Hand1 cell lineage reveals novel contributions to cardiovascular, neural crest,
extra-embryonic, and lateral mesoderm derivatives. *Dev Dyn* **239**, 3086-3097,
doi:10.1002/dvdy.22428 (2010).
35   Walton, K. D. *et al.* Villification in the mouse: Bmp signals control intestinal villus
patterning. *Development* **143**, 427-436, doi:10.1242/dev.130112 (2016).

36   Hatch, J. & Mukouyama, Y. S. Spatiotemporal mapping of vascularization and
innervation in the fetal murine intestine. *Dev Dyn* **244**, 56-68, doi:10.1002/dvdy.24178
(2015).
37   Soriano, P. Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat*
Genet **21**, 70-71 (1999).
38   Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L. & Luo, L. A global double-fluorescent
Cre reporter mouse. *Genesis* **45**, 593-605, doi:10.1002/dvg.20335 (2007).
39   Lua, I., Li, Y., Pappoe, L. S. & Asahina, K. Myofibroblastic Conversion and
Regeneration of Mesothelial Cells in Peritoneal and Liver Fibrosis. *The American*
journal of pathology **185**, 3258-3273, doi:10.1016/j.ajpath.2015.08.009 (2015).
40   Baillie-Johnson, P., van den Brink, S. C., Balayo, T., Turner, D. A. & Martinez Arias,
A. Generation of Aggregates of Mouse Embryonic Stem Cells that Show Symmetry
Breaking, Polarization and Emergent Collective Behaviour In Vitro. *J Vis Exp*,
doi:10.3791/53252 (2015).
41   Beccari, L. *et al.* Multi-axial self-organization properties of mouse embryonic stem
cells into gastruloids. *Nature* **562**, 272-276, doi:10.1038/s41586-018-0578-0 (2018).
42   Turner, D. A. *et al.* Anteroposterior polarity and elongation in the absence of extra-
embryonic tissues and of spatially localised signalling in gastruloids: mammalian
embryonic organoids. *Development* **144**, 3894-3906, doi:10.1242/dev.150391
(2017).
43   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676-682, doi:10.1038/nmeth.2019 (2012).
716
717