Title: Rapid GIT transit time in volant vertebrates, with implications for convergence in

microbiome composition

Target Journals: The Journal of Physiological Sciences (Short Communication)

Authors: Emily Cornelius Ruhs[1,2φ], Katherine McFerrin[1φ], Devin N. Jones[3], Natalia Cortes-

Delgado[1], Ny Anjara Fifi Ravelomanantsoa[4], Carl J. Yeoman[3,5], Raina K. Plowright[3,6], Cara E.

Brook[1]

φCo-first author

Correspondence: ecruhs@uchicago.edu
Affiliations:
1. University of Chicago, Department of Ecology and Evolution, Chicago, IL 60637
2. Field Museum of Natural History, Grainger Bioinformatics Center, Chicago, IL, 60605
3. Montana State University, Department of Microbiology & Cell Biology, 109 Lewis Hall,
Bozeman, MT 59717
4. University of Antananarivo, Department of Zoology and Animal Biodiversity, Antananarivo,
Madagascar
5. Department of Animal and Range Sciences, Montana State University, Bozeman, MT 59717
6. College of Veterinary Medicine, Department of Public & Ecosystem Health, Cornell
University, Ithaca, NY, 14850
23
ORCID:
ECR: https://orcid.org/0000-0002-0546-6593
DNJ: https://orcid.org/0000-0001-9215-2930
NAFR: https://orcid.org/0000-0002-0977-8383
CYJ: https://orcid.org/0000-0002-8775-7879
RKP: https://orcid.org/0000-0002-3338-6590
CEB: https://orcid.org/0000-0003-4276-073X

ABSTRACT:

Flying birds and bats have simplified gastrointestinal tracts (GITs) and low intestinal mass to

facilitate flight. Previous work showed reduced GIT transit times in birds relative to other

vertebrates—but GIT transit has never been collectively quantified for bats. Unique among

mammals, bat GIT microbiomes are dominated by Pseudomonadota bacteria (previously

Proteobacteria), which also dominate the microbiomes of flying birds – we hypothesized this

convergence to result from rapid GIT transit times for both volant taxa. We conducted a meta-

analysis of vertebrate GIT transit times which showed that bats and flying birds have

significantly faster transit times relative to nonvolant vertebrates. Additionally, within the bat

order (Chiroptera), we demonstrated decreasing transit times associated with increasing body

mass, a pattern contrasting other vertebrates (including volant birds) and possibly influencing

GIT microbiome composition. This inverted mass-transit association is likely driven by diet as

fruit- and nectar-consuming Pteropodids are the largest of all bats.

BACKGROUND

Flying vertebrates (e.g., birds and bats) have unique gastrointestinal physiologies, likely evolved to facilitate powered flight [1]. As one such adaptation, flying birds are known to have rapid gastrointestinal tract (GIT) transit times [2,3], i.e., the time taken for food to pass through the stomach, small and large intestine, and colon prior to excretion [4]. Rapid GIT transit is hypothesized to help volant animals maintain low body masses for flight [2] while maximizing energy intake [5]. Faster GIT transit times also enable an increase in consumption rate during foraging [6]. GIT transit time influences the composition and abundance of the GIT microbiome; specifically, in humans, GIT transit is directly, positively correlated with total bacterial count [7,8] and positively associated with the presence of slow-growing gut microbes, such as methanogens [9,10]. In humans, the small intestine (where transit time ranges from 3-5 hours) has lower microbial diversity and abundance relative to the colon (where transit time can exceed 30 hours) [11]. Most mammalian GIT microbiomes that have been surveyed to date (chiefly those of Artiodactyla, Carnivora, Primates, and Xenarthra) are dominated by the phyla Firmicutes and Bacteroidota (formerly Bacteroidetes)[12]. By contrast, both bird and bat GIT microbiomes tend to be dominated by the phylum Pseudomonadota (previously called Proteobacteria) [13]. It is possible, therefore, that the convergence in composition of flying bird and bat GIT microbiomes could be explained by similarly rapid GIT transit. Though rapid GIT transit times are well-described for flying birds, GIT transit has never been collectively summarized for bats.

The passage of digesta through the GIT is affected by many factors, several of which converge in flying birds and bats. Anatomically, longer and more complex GITs (e.g., multiple stomach chambers such as those found in herbivorous ungulates) increase digesta passage time

[14], but shorter, simpler GITs (such as those found in small birds and bats) decrease transit time

[15,16]. As previously reviewed [15], relative to non-flying mammals, both small birds and bats

have short, simple digestive tracts, with reduced surface area and intestinal tissue [16,17]. Most

bat GITs also lack a cecum and appendix, organs with specialized digestive properties in other

taxa [17–20]. To compensate for the loss of intestinal absorptive surface area important for

nutrient uptake in the GIT, both flying birds and bats have expanded villi in the GIT epithelium

and exhibit greater reliance on paracellular absorption, the passive uptake of water soluble

nutrients in the GIT, for meeting energetic demands [16].

In addition to anatomy, diet composition plays a key role in modulating GIT transit times.

The water and fat content, consistency, and hardness of ingested food all impact passage time

[21]. For instance, insectivorous bats have longer food retention times than

frugivorous/nectivorous bats, likely due to a diet of harder, higher consistency food [5,22,23].

Evidence from pollen-eating bats indicates that the degree of dietary specialization is related to

the efficiency of digestion, as well: for example, flower-specialist bats digest pollen content into

protein resources more rapidly than generalist species [24,25]. Similarly, among insectivorous

bats, the shortest GIT transit times have been recorded for trophic specialists [22]. It is

hypothesized that the microbiome may play a larger role in nutrient extraction for diet specialist

species and may explain some of the differences in the efficiency of digestion and

gastrointestinal design [26].

Finally, digesta passage rate is also modulated by interactions between the host diet and

GIT microbial community [15]. For example, increased fiber intake can decrease passage time as

its breakdown by microbes produces short chain fatty acids (SCFAs), which stimulate intestinal

motility to reduce transit time. High SCFA concentrations are also typically associated with

lower GIT pH, which is, in turn, associated with reduced transit time [27]. Nonetheless, the

interaction among SCFA-producing bacteria, SCFAs, and pH is complex: lower pH

environments are associated with reductions in *Bacteroides* spp., typically one of the most

abundant SCFA-producing genera in the mammalian GIT, and decreased production of the two

most abundant SCFAs (Acetate and Propionate). By contrast, low pH environments favor some

bacteria (e.g. *Eubacterium spp., Roseburia spp., and Faecalibacterium spp.*) that produce greater

amounts of another SCFA, Butyrate [28] [29,30]. In humans, the more rapidly transiting small

intestine is additionally dominated by microbial 'pioneer species' in the phylum

Pseudomonadota, which have high growth rates to facilitate rapid colonization of new substrates

[31,32]. For the same reason, Pseudomonadota are also some of the earliest microbes to establish

in human neonate GIT microbiomes [33] and manifest at higher proportions in the GIT

microbiomes of human hosts experimentally induced to rapidly transit food [34].

Pseudomonadota are highly diverse and, as a result, are thought to be responsible for

considerable functional variation in the tasks performed by the human GIT microbiome, despite

being lower in abundance than Firmicutes and Bacteroidetes [35,36].

We compiled GIT transit times from the literature for a variety of food types across a

wide range of volant (bat and bird) and non-volant (carnivores, ungulates, rodents, primates,

reptiles) vertebrate taxa. We aimed to compare digesta passage times to elucidate their possible

role in structuring differences in the GIT microbial community across these taxa. We

demonstrate that flying birds and bats do indeed have significantly faster gut transit times than

non-volant vertebrates, offering one possible explanation for their uniquely congruent GIT

microbiomes.

METHODS

Literature survey and statistical approach

We collected GIT transit times from the primary literature across a wide range of

vertebrate taxa to assess the impact of flight on the duration of time required for food passage in

the GIT. To compile our database, we searched Web of Science for publications which cited the

following search criteria in the title or abstract: "gut OR gastrointestinal OR food" AND "transit

OR passage" AND "rate OR time." We restricted our search using the "NOT" function to

exclude studies that cited the following off-target terms: human[s], [wo]men, obesity, diabetes,

adult, child[ren], patient[s], volunteer[s], disease, mice/mouse, rat[s]. This exclusion thus

focused our study on GIT transit in non-model animal systems.

This search returned 1169 results, which excluded human and laboratory mouse and/or

rat studies. We then individually reviewed each of the 1169 studies, including any study that

reported "minimum gut transit" (MGT), "mean retention time" (MRT), "transit time", or "gut

passage time" for any vertebrate taxon, in our final database. We examined each study to

confirm that the value reported corresponded to the duration of time passed from consumption to

first passage in feces to ensure these values were comparable. Though mean retention time

(MRT), corresponding to the average period of time food is retained in the gut, offers an

arguably more accurate representation of the gut passage rates relevant to the establishment of

GIT microbial communities [37], we found MGT to be more widely reported in the literature

across multiple vertebrate taxa, offering a larger pool for comparison. As a result, we report

results from MGT records in the main analysis and results for MRT in the supplemental analysis

(appendix 1). Most of the studies returned from the search were irrelevant to our research

agenda; we identified 107 unique publications that reported MGTs or equivalent for 112 unique

species (MRT for only 36 unique species) for inclusion in our final database

(https://github.com/brooklabteam/git-transit-time/tree/main). Some studies reported MGT or

MRT for multiple taxa, leading to multiple entries in the database, which we summarized prior to

statistical analyses.

From each study, we extracted the following information for each animal surveyed and

included it in the final database (Supplementary Dataset 1): (1)

Order/Family/Genus/Species/Common Name, (2) diet classification (classed as

frugivore/nectarivore, insectivore, omnivore, folivore, carnivore), (3) mass (grams), (4) food

included in feeding trial (broad description), (5) food type in feeding trial (specific classification

among four possible categories: fruit/nectar/pollen, protein, fiber/foliage, or a label-solution

typical to these studies, such as barium sulfate), (6) mean or median MGT (or MRT) and, when

available, minimum/maximum, and standard error/deviation of reported MGT, and finally (7)

number of individuals and number of trials included in survey. In cases where some of the

species-level information (i.e., mass) was missing from the original study, we searched the

literature to find averages of these values for the taxon in question. We recorded the source (if

not the original study) and the method of data summarization for each entry in our database,

adjacent to the metric in question.

We next compared GIT transit times across the different taxonomic groups surveyed.

First, we used the unique transit time value for each entry in the database, preferentially selecting

the median reported value when available, then the mean if median was not reported. We then

calculated an average transit time value for each combination of unique species and diet type in

our database, weighting database entries by the number of individuals surveyed and the number

of trials conducted in each original study. In cases where this original sample size information

was not included in the study (n=9), we assumed a sample size and trial size of only one

individual in the weighting. Once the dataset was collapsed to a single transit time metric for

each combination of species and diet type (a total of 142 entries representing 109 unique

species), we then compared gut transit across disparate taxa.

Due to small sample sizes, we elected to categorize all non-mammalian vertebrates

taxonomically by Class (specifically, Birds, Fish, Amphibians, Reptiles) and all mammals by

Order. We separated birds into "Flying" and "Non-Flying" categories, though only two non-

flying birds (*Spheniscus demersus*, the African penguin, and *Strutheo camelus*, the common

ostrich) were represented in the database. After assigning these groupings, we subset the

database to compare only those taxonomic groupings where more than five entries were reported,

thus reducing our final database for comparison down to 140 entries across 103 unique species

(47 bats, 14 carnivores, 16 even-toed ungulates [order Artiodactyla], 16 flying birds [order

Aves], 22 primates, 8 rodents, 17 reptiles; supplemental table 1).

Statistical Comparison of GIT Transit Times by Taxa

Using our most stringent data subset, we first asked the question: *What is the relationship*

between taxonomic grouping and digesta passage time? To address this statistically, we fit a

linear mixed effect model in the Gaussian family from the lme4 package in R (version 2023.9.1)

[38], assuming a response variable of the $\log_{10}$ transit time (in hours), a single predictor variable

of taxonomic grouping (modeled as a categorical factor) and a random effect of food type (Table

1). We used the "Rodent" grouping (which demonstrated the lowest median GIT transit outside

the two volant taxa) as the reference level for comparison and computed p-values using

Satterthwaite's degrees of freedom method in the "lmerTest" package [39].

To elaborate on this interaction, we next asked the question: *What is the relationship*

between the interaction of host body mass, taxonomic grouping, and the rapidity of GIT transit

time? To address this question statistically, we fit another linear mixed-effects model in the lme4

package in R, this time querying the response variable of the $\log_{10}$ transit time (in hours) against

the predicter variable of the interaction of the $\log_{10}$ of the host's body mass (in kg) with its

taxonomic grouping and included a random effect of taxonomic grouping in this model (Table

2).

190
RESULTS

Relationship between taxonomic grouping and the rapidity of GIT transit time

We found strong evidence that each taxonomic grouping maintained a significantly

different GIT transit time (as reported by MGT) compared to the reference level ("Rodent";

Table 1, Figure 1a). Carnivores, primates, ungulates, and reptiles had significantly greater GIT

transit times relative to rodents, whereas birds and bats had shorter GIT transit times relative to

rodents (Figure 2a).

Random effect of food type on the correlation between taxonomic group and GIT transit time

In the first mixed model (without inclusion of a covariate predictor of body mass), we

found a significant random effect of food type on the relationship between taxonomic grouping

and GIT transit time, with fruit/nectar/pollen diets associated with the most rapid gut transits

(**Figure 2b**). There was no evidence to suggest that fiber/foliage or protein diets had an effect on

MGT but labeled solutions (e.g., dye or barium-sulfate solution to monitor gastrointestinal transit

time) were found to be associated with longer gut transit times (increased MGT).

Relationship between the interaction of host body mass, taxonomic grouping, and the rapidity of

GIT transit time

Four taxonomic groups demonstrated significant positive interaction effects with $\log_{10}$

body mass to modulate GIT transit time (as reported by MGT; Table 2): larger body size was

associated with longer transit times in the case of all taxonomic groups, except for ungulates and

bats (Figure 2c). By contrast, bats had marginally negative effects with $\log_{10}$ body mass to

modulate GIT transit time – showing that larger body size was associated with shorter gut transit

times.

Relationship between the interaction of host body mass, sub-grouping (e.g., family-level), and

the rapidity of GIT transit time in the order Chiroptera

Given this reversal of the GIT transit time association with mass identified in the bat data,

as well as the significant random effect of diet type in model 1 (Table 1), we next explored

whether family-level groupings (associated with individual diet types) could further explain this

relationship between host body mass and GIT transit time (as reported by MGT). We only

considered data from bat families with more than one-entry (Phyllostomidae, Pteropodidae, and

Vespertilionidae). While none of the family-level groupings had significantly different slopes

from one-another, Pteropodidae – with a strictly fruit/nectar/pollen diet and the largest bodied bat

family - was the only family with a non-negative slope (Figure 1C; Table 3).

Discussion

Understanding the factors influencing gastrointestinal transit (GIT) times across

vertebrate taxa is crucial for unraveling the intricate relationship between digestive physiology,

diet, and evolutionary adaptations. In this study, we aimed to compare GIT transit times among

volant (bats and birds) and non-volant vertebrates while considering the effects of body mass and

diet. Our findings shed light on the unique digestive adaptations of flying vertebrates and their

implications for GIT microbial communities. Consistent with previous research, our study

reveals significant differences in GIT transit times among taxonomic groups. The taxonomic

groupings of carnivora, primates, ungulates, and reptiles exhibited longer GIT transit times

compared to rodents, whereas flying birds and bats demonstrated shorter transit times. This

pattern of accelerated transit time was consistent for bats regardless of whether using the metric

MGT or MRT as a proxy for GIT transit time. This highlights the diverse strategies employed by

vertebrates to process and assimilate nutrients, allowing them to adapt to unique life histories and

dietary preferences shaped by their evolutionary histories.

In our second analysis, host body mass emerged as a key determinant of GIT transit times

across taxonomic groups. Generally, larger-bodied species exhibited longer transit times,

consistent with expectations based on the scaling of gastrointestinal tract length [40]. Indeed,

flying vertebrates have shorter intestines than non-volant mammals and reptiles to facilitate flight

through reductions in body mass [41]. However, these anatomical changes in flying birds and

bats have necessitated physiological adaptions for rapid nutrient absorption and water regulation

which could, in turn, influence food passage times [16,42,43]. For example, we observed a

reversal of this trend in Pteropodid bats, where larger body size was associated with shorter

transit times (though this association was not statistically significant) [3]. Bats in the family

Pteropodidae (Figure 1c), are on average, the largest of all bat species and maintain a fruit and

nectar-based diet, which we showed to be associated with shorter GIT transit time. Nonetheless,

while fruit and nectar-based diets were associated with more rapid gut transit than other diets in

our analysis (Figure 2b), taxonomic grouping remained a significant predictor of transit time

even after accounting for diet—suggesting a unique digestive physiology in fruit bats

independent of dietary differences. By contrast, label solutions were associated with longer

transit times in our analysis, independent of vertebrate taxa, likely because these standard

markers have foreign compositions that vertebrate guts are not optimized to digest.

The observed differences in GIT transit times among vertebrate taxa have significant

implications for GIT microbial communities. Accelerated transit times in flying birds and bats

may facilitate the rapid turnover of gut contents, shaping microbial diversity and composition;

supported by the dominance of Pseudomonadota in volant species [13]. For many taxa,

distinctive GIT microbial communities colonize different regions of the digestive tract and

perform specific digestive functions [44]. By contrast, microbiome composition and abundance

may be relatively consistent throughout regions of the bat GI tract (except in some frugivores,

which demonstrate moderate shifts in relative abundance of different microbial taxa across the

length of the GIT [45]). In general, consumption of difficult-to-digest, highly fibrous foliage is

rare in both birds and bats [3]. Frugivorous bats disgorge most of the fibrous material (i.e., fruit

pulp) in their diets and only consume the nectar and juice [46]. The increased importance of

paracellular nutrient absorption in volant vertebrates may then negate the need for microbiota to

enhance these processes in both birds and bats [15]. The convergence of GIT microbial

communities between birds and bats, despite their phylogenetic distance, underscores the

importance of GIT physiology in shaping microbial ecology.

Future directions

Our study provides valuable insights into the factors which underpin variation in GIT

transit times across vertebrate taxa. By elucidating the interplay between diet, body size, and

evolutionary history, we contribute to a deeper understanding of digestive adaptations in flying

vertebrates and their implications for GIT microbial communities. Future research should

explore the functional significance of these adaptations and their role in vertebrate ecology and

evolution.

DECLARATIONS

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

The datasets generated and the code used for analysis for the present study are available at:

https://github.com/brooklabteam/git-transit-time.git

Competing interests

The authors declare no competing interests.

Funding

The authors acknowledge funding support from a Branco Weiss 'Society in Science' fellowship

to CEB, an NIH-NIAD DP2 award to CEB (1DP2AI171120), a pilot grant from the Microbiome

Center at the University of Chicago to CEB, the DARPA PREEMPT Program Cooperative

Agreement #D18AC00031 to RPK (with a subgrant to CEB), and a U.S. National Science

Foundation grant (DEB-1716698, EF-2133763/EF-2231624) to RKP. CJY was funded through

the Montana Agricultural Experiment Station (MONB00113). ECR was funded by a Grainger

Bioinformatics Fellowship through the Field Museum of Natural History (Chicago, IL), and

NCD was funded by a Chicago Fellows scholarship through the Biological Sciences Division of

the University of Chicago. The content of the information does not necessarily reflect the

position or the policy of the U.S. government, and no official endorsement should be inferred.

Author's contributions

ECR conducted the final data analysis and wrote the manuscript. KM collected data from the

literature and helped with the data analysis. DNJ conceptualized the project, helped with the data

collection, and wrote the manuscript. NAFR helped with the data analysis. NCD reviewed the

data and wrote the manuscript. CJY helped with the data analysis. RKP conceptualized the

project and provided funding. CEB conceptualized the project, collected the original data, helped

with data analysis, funded the project, and wrote the manuscript. All authors reviewed and

provided edits to the final manuscript.

Acknowledgements

The authors thank the Brook lab at the University of Chicago for helpful comments on the

manuscript.

323 **Table 1.** Relationship between taxonomic grouping and the rapidity of GIT transit time.

Taxonomic Grouping	Slope	Std. error	Degrees of freedom	t-value	p-value
(Intercept)	0.5473	0.1353	40.5218	4.045	0.000228 ***
Flying Birds	-0.587	0.1492	139.7	-3.936	0.000130 ***
Bats	-0.427	0.1387	135.9	-3.080	0.002505 **
Carnivores	0.495	0.1635	133.1	3.028	0.002955 **
Primates	0.839	0.1435	13.85	5.849	3.34e-08 ***
Ungulates	1.017	0.1487	139.57	6.840	2.29e-10 ***
Reptiles	1.462	0.1488	139.97	9.827	< 2e-16 ***
random effect of food type: p-value = 0.02377*					
* at p <.05; ** at p < .01; ***at p < .001					

324 Results of the mixed effects model of GIT transit times, as predicted by taxonomic grouping with

325 a random effect of diet classification. *Model: $\log_{10}(transit\ time) \sim Class + (1|\ Diet),\ REML=F$*.

327   **Table 2**. Relationship between the interaction of host body mass, taxonomic grouping, and the

328   rapidity of GIT transit time.

log10(mass): taxonomic grouping	Slope	Std. error	Degrees of freedom	t-value	p-value
(Intercept)	0.89115	0.25511	7.13648	3.493	0.00978 **
$\log_{10}$(mass):Flying Birds	0.50171	0.08364	136.43675	5.998	1.69e-08 ***
$\log_{10}$(mass):Bats	-0.16267	0.08661	139.23271	-1.878	0.06244 .
$\log_{10}$(mass):Rodents	0.27406	0.11004	133.81241	2.491	0.01397 *
$\log_{10}$(mass):Carnivores	0.48411	0.19952	139.91864	2.426	0.01652 *
$\log_{10}$(mass):Primates	0.29365	0.13470	137.56043	2.180	0.03095 *
$\log_{10}$(mass):Ungulates	0.03099	0.09638	135.84425	0.322	0.74826
$\log_{10}$(mass):Reptiles	0.12151	0.06620	132.74398	1.836	0.06865 .
*random effect of class: p-value = < 2.2e-16 ****					
* at p <.05; ** at p < .01; ***at p < .001					

329   Results of the mixed effects model of GIT transit times, as predicted by the interaction of body

330   mass and taxonomic grouping, with a random effect of taxonomic grouping. *Model: $\log_{10}$(transit*

331   *time) ~ $\log_{10}$(mass):Class + (1|Class), REML=F.*

338   **Table 3**. Relationship between the interaction of host body mass, sub-grouping (e.g. family-

339   level), and the rapidity of GIT transit time in the order Chiroptera.

log10(mass): taxonomic grouping	Slope	Std. error	Degrees of freedom	t-value	p-value
(Intercept)	-0.114	0.148	45.000	-0.771	0.445
$\log_{10}$(mass):Phyllostomidae	-0.143	0.099	45.000	-1.447	0.155
$\log_{10}$(mass):Pteropodidae	0.002	0.137	45.000	0.017	0.987
$\log_{10}$(mass):Vespertilionidae	-0.133	0.085	45.000	-1.574	0.122
random effect of family: p-value = 1					
* at p <.05; ** at p < .01; ***at p < .001					

340   Results of the mixed effects model of GIT transit times, as predicted by the interaction of body

341   mass and family-level grouping, with a random effect of family-level grouping. *Model:*

342   *$\log_{10}$(transit time) ~ $\log_{10}$(mass):Family + (1|Family), REML=F.*

Figure 1: *Minimum GIT transit times (as reported by MGT) from experimental studies across*

volant and nonvolant vertebrates. **(A)** Boxplots show the interquartile range of minimum GIT

transit times (on a $\log_{10}$ scale) recovered from studies across diverse vertebrate taxa (x-axis).

Shapes correspond to food type used in each study, according to the legend. Flying birds and

bats have significantly faster GIT transit times relative to the reference taxon (rodents), and

consequently, all other non-flying vertebrates (Table 1; linear mixed effects model; *=p-

value<0.01, **=p-value<0.001, and ***=p-value< 0.0001). **(B)** Minimum GIT transit time (y-

axis, $\log_{10}$ scale) for each taxon relative to their body mass (in kg; x-axis), with taxa

distinguished via color and food type indicated by shape as highlighted in (A). Trend lines (Table

2; Linear mixed effects model) indicate that MGT increases with body mass in all taxa except

bats, which show the opposite trend (p-value =0.06). Polygons encircle the full range of species

surveyed in each taxa. **(C)** Bat-only data, showing the reversed direction of the relationship

between mass and GIT transit for bats (from positive for other taxa to negative in bats). The

largest-bodied Pteropodidae fruit bats show some of the most rapid transit times and appear to

lack a relationship with body mass (Table 3).

Figure 2: *Fixed and random partial effects on GIT transit time (as reported by MGT).* Fixed **(a)** and random **(b)** effects sizes from a linear mixed effects model with taxonomic class as a fixed effect and diet category as a random effect. Bats and flying birds show significant negative slopes, indicating more rapid GIT transit times than rodents, while all other taxa demonstrate significant positive slopes, indicating slower GIT transit times than rodents. Fruit/nectar/pollen diets show the most negative interaction with GIT transit, whereas label-solution (a medical solution marker, such as barium sulfate) shows a positive interaction. Fixed **(c)** and random **(d)** effects from a linear mixed effects model with log10(mass): taxonomic class as a fixed effect interaction and taxonomic class as a random effect. Bats are the only taxonomic group to show that mass has a marginally inverse effect on GIT transit.

REFERENCES

1. Maina, J.N. (2000) What it Takes to Fly: The Structural and Functional Respiratory Refinements in Birds and Bats. *Journal of Experimental Biology* 203, 3045–3064
2. McWhorter, T.J. *et al.* (2009) The integration of digestion and osmoregulation in the avian gut. *Biol Rev Camb Philos Soc* 84, 533–565
3. Price, E.R. *et al.* (2015) Digestive Adaptations of Aerial Lifestyles. *Physiology* 30, 69–78
4. Müller, M. *et al.* (2018) Gastrointestinal Transit Time, Glucose Homeostasis and Metabolic Health: Modulation by Dietary Fibers. *Nutrients* 10, 275
5. Laska, M. (1990) Food transit times and carbohydrate use in three Phyllostomid bat species. *Z. Säugetierkd* 55, 49–54
6. Staliński, J. (1994) Digestion, defecation and food passage rate in the insectivorous bat Myotis myotis. *Acta Theriologica* 39, 1–11
7. Flint, H.J. *et al.* (2012) The role of the gut microbiota in nutrition and health. *Nat Rev Gastroenterol Hepatol* 9, 577–589
8. Stephen, A.M. *et al.* (1987) Effect of changing transit time on colonic microbial metabolism in man. *Gut* 28, 601–609
9. El Oufir, L. *et al.* (1996) Relations between transit time, fermentation products, and hydrogen consuming flora in healthy humans. *Gut* 38, 870–877
10. Jeffery, I.B. and O'Toole, P.W. (2013) Diet-Microbiota Interactions and Their Implications for Healthy Living. *Nutrients* 5, 234–252
11. Martinez-Guryn, K. *et al.* (2019) Regional Diversity of the Gastrointestinal Microbiome. *Cell Host & Microbe* 26, 314–324
12. Nishida, A.H. and Ochman, H. (2018) Rates of gut microbiome divergence in mammals. *Mol Ecol* 27, 1884–1897
13. Song, S.J. *et al.* (2020) Comparative Analyses of Vertebrate Gut Microbiomes Reveal Convergence between Birds and Bats. *mBio* 11, e02901-19, /mbio/11/1/mBio.02901-19.atom
14. Clemens, E.T. and Stevens, C.E. (1980) A comparison of gastrointestinal transit time in ten species of mammal. *The Journal of Agricultural Science* 94, 735–737
15. Jones, D.N. *et al.* (2022) Do gastrointestinal microbiomes play a role in bats' unique viral hosting capacity? *Trends in Microbiology* 30, 632–642
16. Caviedes-Vidal, E. *et al.* (2007) The digestive adaptation of flying vertebrates: High intestinal paracellular absorption compensates for smaller guts. *Proceedings of the National Academy of Sciences* 104, 19132–19137
17. Klite, P.D. (1965) Intestinal bacterial flora and transit time of three neotropical bat species. *J Bacteriol* 90, 375–379
18. Popov, I.V. *et al.* (2021) Gut microbiota of bats: pro-mutagenic properties and possible frontiers in preventing emerging disease. *Sci Rep* 11, 21075
19. Gadelha-Alves, R. *et al.* (2008) Comparative Intestinal Histomorphology of Five Species of Phyllostomid Bats (Phyllostomidae, Microchiroptera): Ecomorphological Relations with Alimentary Habits. *International Journal of Morphology* 26, 591–602
20. Guinane, C.M. *et al.* (2013) Microbial Composition of Human Appendices from Patients following Appendectomy. *mBio* 4, e00366-12
21. Traveset, A. (1998) Effect of seed passage through vertebrate frugivores' guts on germination: a review. *Perspectives in Plant Ecology, Evolution and Systematics* 1/2, 151–190

22. Roswag, A. *et al.* (2012) Inter- and intraspecific comparisons of retention time in
insectivorous bat species (Vespertilionidae). *Journal of Zoology* 288, 85–92
23. Buchler, E.R. (1975) Food Transit Time in Myotis lucifugus Chiroptera: Vespertilionidae).
Journal of Mammalogy 56, 252–255
24. Herrera M., L.G. and Martínez Del Río, C. (1998) Pollen Digestion by New World Bats:
Effects of Processing Time and Feeding Habits. *Ecology* 79, 2828–2838
25. Mancina, C.A. *et al.* (2005) Pollen digestion by nectarivorous and frugivorous Antillean bats.
Mammalian Biology 70, 282–290
26. Ingala, M.R. *et al.* (2021) You are more than what you eat: potentially adaptive enrichment
of microbiome functions across bat dietary niches. *Animal Microbiome* 3, 82
27. Procházková, N. *et al.* (2023) Advancing human gut microbiota research by considering gut
transit time. *Gut* 72, 180–191
28. Duncan, S.H. *et al.* (2009) The role of pH in determining the species composition of the
human colonic microbiota. *Environmental Microbiology* 11, 2112–2122
29. Firrman, J. *et al.* (2022) The impact of environmental pH on the gut microbiota community
structure and short chain fatty acid production. *FEMS Microbiology Ecology* 98, fiac038
30. Walker, A.W. *et al.* (2005) pH and Peptide Supply Can Radically Alter Bacterial Populations
and Short-Chain Fatty Acid Ratios within Microbial Communities from the Human Colon.
Applied and Environmental Microbiology 71, 3692–3700
31. Zoetendal, E.G. *et al.* (2012) The human small intestinal microbiota is driven by rapid uptake
and conversion of simple carbohydrates. *ISME J* 6, 1415–1426
32. Le Bouguénec, C. and Schouler, C. (2011) Sugar metabolism, an additional virulence factor
in enterobacteria. *International Journal of Medical Microbiology* 301, 1–6
33. Fusco, W. *et al.* (2023) Short-Chain Fatty-Acid-Producing Bacteria: Key Components of the
Human Gut Microbiota. *Nutrients* 15, 2211
34. Minnebo, Y. *et al.* (2023) Gut microbiota response to in vitro transit time variation is
mediated by microbial growth rates, nutrient use efficiency and adaptation to in vivo transit
time. *Microbiome* 11, 240
35. Bradley, P.H. and Pollard, K.S. (2017) Proteobacteria explain significant functional
variability in the human gut microbiome. *Microbiome* 5, 36
36. Moon, C.D. *et al.* (2018) Metagenomic insights into the roles of Proteobacteria in the
gastrointestinal microbiomes of healthy dogs and cats. *Microbiologyopen* 7, e00677
37. De Cuyper, A. *et al.* (2018) How does dietary particle size affect carnivore gastrointestinal
transit: A dog model. *J Anim Physiol Anim Nutr (Berl)* 102, e615–e622
38. Bates, D. *et al.* (2015) Fitting linear mixed-effects models using lme4. *Journal of Statistical*
Software
39. Kuznetsova, A. *et al.* (2017) lmerTest Package: Tests in Linear Mixed Effects Models.
Journal of Statistical Software 82, 1–26
40. Calder, W.A. (1983) Ecological scaling: mammals and birds. *Annual Review of Ecology and*
Systematics 14, 213–230
41. Duque-Correa, M.J. *et al.* (2022) Diet, habitat and flight characteristics correlate with
intestine length in birds. *Proceedings of the Royal Society B: Biological Sciences* 289,
20220675
42. Schondube, J.E. *et al.* (2001) Diet and the evolution of digestion and renal function in
phyllostomid bats. *Zoology (Jena)* 104, 59–73

43. Levey, D.J. and Rio, C.M. del (2001) It Takes Guts (and More) to Eat Fruit: Lessons From
Avian Nutritional Ecology. *The Auk* 118, 819–831
44. Savage, D.C. (1977) Microbial Ecology of the Gastrointestinal Tract. *Annu. Rev. Microbiol.*
31, 107–133
45. Carrillo-Araujo, M. *et al.* (2015) Phyllostomid bat microbiome composition is associated to
host phylogeny and feeding strategies. *Front. Microbiol.* 6
46. Kunz, T.H. *et al.* (2011) Ecosystem services provided by bats: Ecosystem services provided
by bats. *Annals of the New York Academy of Sciences* 1223, 1–38
479