Title: Chickpea NCR13 disulfide cross-linking variants exhibit profound differences in

antifungal activity and modes of action

Authors: James Godwin[a,1], Arnaud Thierry Djami-Tchatchou[a,1], Siva L. S. Velivelli[a], Meenakshi

Tetorya[a], Ravi Kalunke[a], Ambika Pokhrel[a], Mowei Zhou[c,2], Garry W. Buchko[c,d], Kirk J.

Czymmek[a,b], and Dilip M. Shah[a,3]

Author Affiliations:

[a]Donald Danforth Plant Science Center, St Louis, MO. 63132, USA

[b]Advanced Bioimaging Laboratory, Donald Danforth Plant Science Center, St Louis, MO. 63132,

USA

[c]Earth and Biological Sciences Directorate, Pacific Northwest National Laboratory, Richland, WA

99354, USA

[d]School of Molecular Biosciences, Washington State University, Pullman, WA 99164, USA

1. These authors contributed equally to the manuscript.

2. Current address: Department of Chemistry, Zhejiang University, 866 Yuhangtang Road,

Hangzhou, Zhejiang 310058, China

3.   Corresponding Author:

Correspondence and requests for materials should be addressed to D.M.S. (Email:

dshah@danforthcenter.org)

Abstract

Small cysteine-rich antifungal peptides with multi-site modes of action (MoA) have potential for

development as biofungicides. In particular, legumes of the inverted-repeat lacking clade express

a large family of nodule-specific cysteine-rich (NCR) peptides that orchestrate differentiation of

nitrogen-fixing bacteria into bacteroids. These NCRs can form 2 or 3 intramolecular disulfide

bonds and a subset of these peptides with high cationicity exhibits antifungal activity. However,

the importance of intramolecular disulfide pairing and MoA against fungal pathogens for most of

these plant peptides remains to be elucidated. Our study focused on a highly cationic chickpea

NCR13, which has a net charge of +9 and contains six cysteines capable of forming three disulfide

bonds. NCR13 expression in *Pichia pastoris* resulted in formation of two peptide folding variants,

NCR13_PFV1 and NCR13_PFV2, that differed in the pairing of two out of three disulfide bonds

despite having an identical amino acid sequence. The NMR structure of each PFV revealed a

unique three-dimensional fold with the PFV1 structure being more compact but less dynamic.

Surprisingly, PFV1 and PFV2 differed profoundly in the potency of antifungal activity against

several fungal plant pathogens and their multi-faceted MoA. PFV1 showed significantly faster

fungal cell-permeabilizing and cell entry capabilities as well as greater stability once inside the

fungal cells. Additionally, PFV1 was more effective in binding fungal ribosomal RNA and

inhibiting protein translation *in vitro*. Furthermore, when sprayed on pepper and tomato plants,

PFV1 was more effective in reducing disease symptoms caused by *Botrytis cinerea*, causal agent

of gray mold disease in fruits, vegetables and flowers. In conclusion, our work highlights the

significant impact of disulfide pairing on the antifungal activity and MoA of NCR13 and provides

structural framework for design of novel, potent antifungal peptides for agricultural use.

Author Summary

Fungal pathogens cause significant pre-harvest and post-harvest losses of crop yield, making them a serious biological threat to global food security. Chemical fungicides are effective in controlling fungal diseases across various crops. However, rapid evolution of fungal pathogen resistance to single-site chemical fungicides in agriculture has created an urgent need for development of safe, sustainable, and cost-effective multi-site fungicides. Nodule-specific cysteine-rich (NCR) peptides expressed in the inverted-repeat lacking clade legume plants exhibit potent antifungal activity; however, their modes of action (MoA) are poorly understood. Particularly, the specific contribution of disulfide pairing to the potency and spectrum of antifungal activity against fungal plant pathogens and MoA of these peptides remains to be identified. Chickpea NCR13 expressed in *P. pastoris* generates two peptide variants that differ in their disulfide cross-linking pattern. These variants exhibit striking differences in their three-dimensional structures and potency of antifungal activity against multiple fungal pathogens and MoA. They also differ in their ability to confer resistance to gray mold in pepper and tomato plants. Our study highlights the major impact a specific pattern of disulfide pairing can have on the *in vitro* and *in planta* antifungal activity of an NCR peptide.

Introduction

The management of fungal diseases using chemical fungicides is one of the mainstays of modern agriculture. Unfortunately, modern fungicides that have a single site mode of action (MoA) are losing their effectiveness owing to an increase in development of fungal resistance (1). A major limitation in fungicide development is finding molecules with novel structures and multi-site MoA to mitigate evolution of resistance.

Plants express many antimicrobial peptides that are vital components of the innate immune system and act as a first line of defense against bacterial and fungal pathogens (2, 3). Moreover, plants express a multitude of cationic antifungal peptides differing in structure and sequence (4). They are versatile, have large sequence space, and can be tuned for broad-spectrum antifungal activity against plant fungal pathogens. Thus, cationic antimicrobial peptides represent promising templates for a new generation of fungicides with broad action against fungal pathogens (5).

Several antifungal peptides expressed in plants are cysteine-rich with a characteristic number of cysteine-residues at conserved positions and exhibit high degree of sequence diversity (4, 6). Defensins and defensin-like peptides are the best characterized examples of cysteine-rich peptides employed by plants to combat fungal pathogens (7-9). Legumes of the inverted repeat-lacking clade (IRLC) express many nodule-specific cysteine-rich (NCR) defensin-like peptides that cause differentiation of nitrogen-fixing bacteria to bacteroids. In *Medicago truncatula* and chickpea, terminal differentiation process is orchestrated by 639 and 63 nodule-specific cysteine-rich (NCR) peptides respectively, with conserved cysteine motifs (10). NCRs cause bacterial genome endoreduplication, changes in transcription and translation, and induction of morphological changes leading to the formation of bacteroids with unique morphology (8, 11-13). Each NCR is synthesized as a precursor protein containing an N-terminal signal peptide and a C-terminal mature

peptide. Several NCRs in each legume are cationic with a net charge ranging from +3 to +11 and

rich in hydrophobic residues, dual hallmarks of defensin-like antimicrobial peptides (10, 14).

Large landscape of natural NCRs from IRLC legumes offer an unprecedented opportunity for

discovery of the novel antimicrobial functions in plants. Several NCR peptides exhibiting

antibacterial activity against Gram-negative and Gram-positive bacteria have so far been identified

(15-18). Another report detailed the evaluation of 19 cationic *M. truncatula* NCRs for their ability

to inhibit the growth of a human fungal pathogen *Candida albicans*. Of these, nine with a net

charge above +9 inhibited the growth of this pathogen at concentrations ranging from 1.5 to 10.5

µM (14). Previously, we reported *M. truncatula* NCR044 having broad-spectrum antifungal

activity and multi-faceted MoA. In addition, when sprayed on leaf surfaces of tomato and

Nicotiana benthamiana plants, NCR044 conferred resistance to the gray mold disease caused by

the fungal pathogen *B. cinerea* (19).

NCRs exhibit diverse primary sequences and net charges, with a notable conserved feature

being the presence of four or six conserved cysteine residues. These cysteines form intramolecular

disulfide bonds which enhance the stability of the peptides (10, 20). The influence of cysteines and

disulfide cross-linking on the structure and function of *M. truncatula* NCR247 and NCR169, each

containing four cysteines, has been previously investigated. Four conserved cysteines and disulfide

cross-linking had major influence on the symbiotic activities of NCR247 tested against

Sinorhizobium meliloti (21, 22). NCR169 produced two structurally distinct disulfide variants that

exhibited differences in phospholipid binding and antibacterial activity (23). However, the

significance of disulfide cross-linking for the structures, antifungal activity, and MoA of majority

of NCRs is poorly understood.

In this study, we show that a highly cationic chickpea NCR13 peptide expressed in *Pichia pastoris* forms two structurally unique disulfide variants despite having an identical amino acid sequence. Interestingly, these two variants exhibited distinct differences in their antifungal potency against economically important plant fungal pathogens. NCR13 disulfide variants displayed notable differences in fungal cell entry, cell permeabilization, stability and MoA against the fungal pathogen *B. cinerea*. While both disulfide variants reduced gray mold disease symptoms by spray application on the leaves of pepper and tomato plants, the NCR13 variant with potent *in vitro* antifungal activity was more effective in controlling the disease *in planta*. Our study reveals the remarkable impact of disulfide cross-linking on the structure-activity relationships and MoA of an antifungal defensin-like NCR peptide.

Results

Chickpea NCR13 is expressed in *P. pastoris* as two peptide folding variants differing in disulfide pairing

In chickpea, NCR13 is one of 63 predicted NCR peptides expressed only in nodules undergoing differentiation during symbiosis (10, 24). NCR13 is a 32-residue peptide with six cysteines predicted to form three disulfide bonds (Fig 1A). Two other homologs of this peptide, NCR07 and NCR15, are also expressed in chickpea nodules with highly conserved cysteine residues (Fig 1A). No other IRLC legumes express NCRs with significant homology to NCR13. NCR13 has 66.6% and 48.4% sequence identity with NCR15 and NCR07, respectively, carrying a net charge of +8 and a hydrophobic amino acid content of 28%, hallmark features of a potential antifungal peptide (Fig 1B).

To investigate the antifungal properties of the NCR13 peptide, it was expressed in a

heterologous *Pichia pastoris* system. The peptide was secreted into the growth medium and

purified using the cation exchange chromatography. When the NCR13 peptide was subjected to

C18 reverse-phase high-performance liquid chromatography (HPLC), we consistently observed

two peaks (Fig 1C). In contrast, chemically synthesized NCR13 produced only one peak (Fig 1D).

These observations led us to suspect that oxidized NCR13 was expressed in our *P. pastoris*

expression system as two species differing in length or disulfide cross-linking pattern. Mass

spectrometry analysis of peptides purified from both HPLC peaks revealed identical molecular

weights (S1 Fig) as expected for NCR13 with three disulfide bonds, narrowing our suspicion to

different disulfide bonding patterns. Across various batches of purified NCR13, peak 1 was

consistently observed to contain less peptide than peak 2 (S1 Table). Peak 1 and peak 2 peptides

are hereafter referred to as NCR13_peptide folding variant 1 (NCR13_PFV1) and NCR13_PFV2

respectively.

A

```
CaNCR13  -TKPCQSDKDCKKFACRKPKVPKCINGFCKCVR-  32
CaNCR15  TKQPCKSRKHCKTYRCPTPKVPNCVNGFCKCVR-  33
CaNCR07  KKMPCKRRRDCKTYPCPHPKVRDCVKGYCKCVVR  34
```

B

Peptide	Net Charge	% Hydrophobic amino acids	Predicted MW (kDa)	Isoelectric point	% identity with NCR13
CaNCR13	+8	28.1	3.66	9.59	100
CaNCR15	+9	24.2	3.81	9.82	66.6
CaNCR07	+10	26.5	4.08	9.85	48.4

Fig 1. Primary amino acid sequence and properties of NCR13. (A) Primary amino acid sequences for Chickpea (*C. arientinum*) NCR13 and its homologs NCR15 and NCR07. Conserved cysteine residues are highlighted in red. The number of amino acids is indicated at the end of each sequence. (B) Peptide characteristics of CaNCR13, CaNCR15 and CaNCR07. (C) High-performance liquid chromatography (HPLC) analyses of NCR13 produced recombinantly in *Pichia pastoris*, monitored at 260 nm, reveals the presence of two major products. mAU represents milli-Absorbing Units. (D) Chemically synthesized NCR13 elutes as one HPLC peak.

To further characterize both NCR13_PFVs, we performed NMR structural analysis of ^{15}N-labeled NCR13_PFV1 and NCR13_PFV2. Fig 2A and B show the assigned ^{1}H-^{15}N HSQC spectra for NCR13_PFV1 and NCR13_PFV2, respectively. The broad dispersion of amide chemical shifts in both spectral dimensions is a standard feature of a structured peptide (25). As illustrated in Fig 2, assignment of both spectra was consistent with the primary amino acid sequence of NCR13.

However, the pattern of the fingerprint ^{1}H-^{15}N HSQC spectrum for both peptides was clearly

different, suggesting their structures were unique. Given that the primary amino acid sequence of

NCR13 contains six cysteine residues (C4, C10, C15, C23, C28, and C30) that can form fifteen

different patterns of intramolecular disulfide bonds when all six are oxidized, our attention

immediately focused on a different disulfide bond pattern for the two peptides. Treatment of both

peptides with the reducing agents dithiothreitol (DTT) or tris(2-carboxyethyl) phosphine resulted

in the identical ^{1}H-^{15}N HSQC spectrum shown in Fig 2C, illustrating that the parent primary amino

sequence was identical for both peptides. Note that when the NCR13 cysteines are all in the

reduced (dithiol) state, dispersion of amide chemical shifts in both spectral dimensions were

distinctively narrow, a standard feature of an unstructured protein (25, 26). As observed for *M.*

truncatula NCR044 (19), disulfide bond formation is essential to stabilize the elements of

secondary structure observed in the oxidized state because in the reduced state the peptide is

largely disordered.

Disulfide bond formation between all six cysteine residues in NCR13_PFV1 and

NCR13_PFV2 was unambiguously verified from the NMR chemical shifts of the cysteine side

chain β-carbon. In general, the cysteine ^{13}C$^{\beta}$ chemical shift is greater than 35 ppm in the oxidized

state and under 32 ppm in the reduced state. As summarized in S2 Table, the ^{13}C$^{\beta}$ chemical shift

for all six cysteines in both isoforms of oxidized NCR13 are between 37 and 48 ppm, indicating

they are all oxidized in NCR13_PFV1 and NCR13_PFV2. Due to the number of potential disulfide

bond patterns that the six cysteine residues can adopt, it was not possible to use NMR structures

in early calculations to unambiguously deduce disulfide bond pairs from ^{1}H-^{1}H NOEs that arise

due to proximal γ-sulphur atoms in the disulfide state. Instead, as previously applied to NCR044,

the disulfide bond pattern was inferred from the analysis of mass spectral data for trypsin-digested,

unlabeled protein in the oxidized and reduced states (19). For NCR13_PFV2, this approach

showed that the disulfide bonds were C4-C10, C15-C30, and C23-C28 (S2 Fig). For

NCR13_PFV1, despite extensive modification of the experimental procedures, this approach

failed due to the hyperstability of this isoform towards trypsinization. Consequently, the disulfide

bond pattern predicted by AlphaFold (27) C4-C23, C10-C28, and C15-C30 was used to calculate

the ensemble of NMR structures for NCR13_PFV1. The two NCR13 isoforms share one identical

disulfide bond, C15-C30, and S2 Table shows that these cysteine $^{13}C^{\beta}$ chemical shifts differ less

than 0.5 ppm, strongly suggesting this disulfide bond is present in both isoforms. As will be

discussed below, this disulfide bond pattern best supports the experimental NOE data.

Fig 2. Fingerprint ^{1}H-^{15}N HSQC spectra for oxidized and reduced NCR13. Assigned ^{1}H-^{15}N
HSQC spectra for oxidized (disulfide) NCR13_PFV1 (A) and NCR13_PFV2 (B). (C) The ^{1}H-^{15}N
HSQC spectrum of fully reduced (dithiol) NCR13_PFV1 and NCR13_PFV2 were identical (the
spectrum for NCR13_PFV2 is illustrated). All spectra collected on ~ 1 mM samples at 20°C in 20
mM sodium acetate, 50 mM NaCl, pH 5.3 at a ^{1}H resonance frequency of 600 MHz. Amide side
chain resonance pairs are connected by a red dashed line. Not within the displayed spectral
boundaries is the G26 amide resonance for NCR13_PFV1 and NCR13_PFV2 at 8.66/103.1 ppm
and 7.99/101.7 ppm, respectively, and the I24 amide resonance for NCR13_PFV1 at 8.72/132.0
ppm.

Three-dimensional NMR solution structures of NCR13_PFV1 and NCR13_PFV2

To obtain structural insights into NCR13_PFVs, we performed NMR analyses of the oxidized forms of these two peptides. Fig 3 illustrates a superposition of cartoon representations of the final ensemble of structures calculated for the two NCR13 PFVs (left) with the main chain of a single structure shown to the right of the ensemble with the disulfide bonds highlighted in different colors. The elements of secondary structure are essentially identical in both isoforms, one anti-parallel β-sheet (V20-I24, F27-V31) and a short a-helix (Q5 or S6–C10). However, the disulfide bond pattern markedly alters the relative orientation of these elements of secondary structure. Both peptides contain one identical disulfide bond, C15-C30, that ties the N-terminal end of the peptide to approximately the middle of the linker region between the two elements of secondary structure. The consequence is that in NCR13_PFV1 the α-helix sits over the top of the face of the β-sheet while in NCR13_PFV2 the α-helix sits on the side of the β-sheet. This difference is better illustrated in S3 Fig, a cartoon illustration of two representative structures superimposed on the β-strand. This results in a smaller accessible surface area for NCR13_PFV1 (3188 ± 63 Å^2) than for NCR13_PFV2 (3357 ± 129 Å^2). The more compact structure for NCR13_PFV1, with the disulfide bonds formed between C4, C10, C23, and C28 directed into the interior of the peptide, may protect these disulfide bonds from reduction and contribute to the hyperstability of NCR13_PFV1 towards reduction. Disulfide bonds may also be protected from reduction because this NCR13 isomer is less dynamic. This suggestion is indirectly made by comparing the ensemble of calculated solution structures illustrated in Fig 3. The ensemble of structures calculated for NCR13_PFV1 converge to a greater extent than for NCR13_PFV2 with the backbone ($N-C^a-C=O$) RMSD to the mean for the ordered residues almost twice as large for NCR13_PFV2 (0.41 ± 13 Å versus 0.76 ± 20 Å, respectively; S3 Table). While this difference reflects the greater number of NOEs observed for

NCR13_PFV1 (316) than for NCR13_PFV2 (119) (S3 Table) due to a more compact structure, it

also indirectly suggests the NCR13_PFV1 explores less conformational space than NCR13_PFV2.

To further verify that the AlphaFold predicted disulfide bond pattern was correct for

NCR13_PFV1, an NMR structure was calculated using only the NOE and TALOS+ (dihedral

angle) data (no hydrogen bonds) with the other possible disulfide bond pattern (C4-C28, C10-C23,

and C15-C30). In S4A Fig, two representative cartoon structures that result from the two possible

disulfide bond patterns are superimposed on the β-strands. In the reversed, C4-C28/C10-C23

disulfide pattern (yellow), the direction of α1 is also reversed relative to the β-strand and this

prevents the N-terminal region (T1-P3) from making close contact with the β-strand. While most

of the NOEs were satisfied in both structure calculations, long range experimental NOEs between

the N-terminal region of NCR13_PFV1 (T1-P3) and the β-strand were only satisfied with the C4-

C23/C10-C28 disulfide pattern used in the structure deposited to the PDB (8UTM), suggesting

this pattern was correct. Furthermore, as shown in S4B Fig, the predicted AlphaFold structure for

NCR13_PFV1 was like the experimental structure (8UTM). In both structures, the β-sheet is

almost identical with the major difference a slightly shorter α-helix displaced two residues (D7-

K11 versus Q5-C10 experimentally) that packs against the β-strand a bit differently in the

AlphaFold structure (green). Most relatively, the N-terminal region (T1-P3) folds towards the β-

strand in the AlphaFold predicted structure, as verified experimentally with the long-range NOEs

between these regions.

Fig 3. Three-Dimensional NMR solution structure of NCR13_PFV1 and NCR13_PFV2.
Cartoon representation of the backbone superposition of the ordered regions in the ensemble of 20 structures calculated for oxidized NCR13_PFV1 (8ULM) and NCR13_PFV2 (7TH8). β-strands are colored blue and labeled sequentially starting from the N-terminal and the lone α-helix is colored wheat. Next to each ensemble is a backbone stick representation of a single structure with the six oxidized cysteine side chains highlighted in yellow, red, or cyan. Above the structures is a schematic summary of the elements of secondary structure observed for the two NMR structures, color-coded similarly. Also shown in the schematic are the disulfide bond connections in both isomers. One disulfide bond, between C15 and C30 (red), is identical in both isomers with C10 and C28 swapping disulfide bond partners.

NCR13_PFV1 exhibits much greater antifungal activity against fungal pathogens than NCR13_PFV2

Having found striking differences in their NMR structures, we investigated the *in vitro* antifungal activity of NCR13_PFV1 and NCR13_PFV2 against three necrotrophic plant fungal pathogens (19). NCR13_PFV1 inhibited the growth of *B. cinerea* at the Minimal Inhibitory Concentration (MIC) of 0.09 µM. Conversely, NCR13_PFV2 inhibited *B. cinerea* at the MIC of 1.5 µM (Figs 4A-C), seventeen-fold higher than that of NCR13_PFV1. Chemically synthesized NCR13 had a MIC of 3 µM, two-fold higher than NCR13_PFV2 (Fig 4B), suggesting that this fully reduced synthesized product was yet another isoform of NCR13 and illustrating the importance of disulfide bonds for antifungal activity. Similarly, NCR13_PFV1 showed several fold greater antifungal activity against *Fusarium virguliforme,* and *Sclerotinia sclerotiorum* compared to NCR13_PFV2 (Fig 4A). Overall, these results reveal that NCR13_PFV1 is a much more potent antifungal peptide compared to NCR13_PFV2.

A

Pathogen	NCR13_PFV1	NCR13_PFV2
	MIC (µM)	
Botrytis cinerea	0.09	1.5
Fusarium virguliforme	0.09	1.5
Sclerotinia sclerotiorum	6.0	24

Fig 4. NCR13_PFV1 exhibits potent antifungal activity compared to NCR13_PFV2. (A) Minimal Inhibitory Concentration (MIC) values of NCR13_PFVs for different fungal pathogens. (B) Antifungal activity of NCR13_PFV1, NCR13_PFV2, and chemically synthesized NCR13 (NCR13_CS) against *B. cinerea*. Fungal cell viability assay using resazurin dye. Change from blue

to pink/colorless signals resazurin reduction and indicates metabolically active *B. cinerea*
germlings after 60 h. NCR13_PFVs are used at concentrations of 0.09 - 6 µM. NCR13_PFV1 at a
concentration 0.045 µM had no antifungal activity (data not shown). N = 3, N indicates biological
replicates. (C) Representative microscopic images showing the inhibition of *B. cinerea* growth 24
h after treatment with 0.09 µM NCR13_PFVs (Right). *B. cinerea* without peptide added served as
a negative control (Left). Scale bar = 100 µm.
296

Disulfide bonds in NCR13_PFV1 contribute to the antifungal activity and formation of

peptide folding variants

Fully oxidized NCR13_PFV1 contains three disulfide bonds: C4-C23, C10-C28, and C15-C30.

The potent antifungal activity of NCR13_PFV1 against *B. cinerea* at nanomolar concentrations

prompted us to investigate the contribution of each disulfide bond towards antifungal activity and

the formation of folded variants when expressed in *P. pastoris*. We carried out functional analysis

of these disulfide bonds by creating three dual Cys to Ser variants to systematically remove one

disulfide bond: NCR13_C4S-C23S, NCR13_C10S-C28S, and NCR13_C15S-C30S variants (Fig

5A). These knockout variants were expressed in *P. pastoris* and purified. HPLC profiles of

NCR13_C4S-C23S, and NCR13_C15S-C30S revealed a single peak whereas NCR13_C10S-

C28S revealed two peaks (Fig 5B), the latter observation suggesting NCR13_C10S-C28S folded

with at least two different disulfide bonding patterns. Mass spectrometry analysis confirmed the

expected molecular mass of each peptide (S4 Table), consistent with them being in an oxidized

state. Therefore, our results indicate that the disulfide pairing of C4-C23 and C15-C30 is essential

for the formation of NCR13_PFVs.

　　All three single disulfide bond knockouts had lower antifungal activity against *B. cinerea*

compared to NCR13_PFV1, as summarized in Fig 5. NCR13_C4S-C23S and NCR13_C15S-

C30S both inhibited *B. cinerea* with a MIC value of 1.5 and 0.75 µM respectively, while the

products in the two NCR13_C10S-C28S HPLC peaks had a MIC of 3 µM (Fig 5C and D). While

we were unable to confirm the disulfide bonding pattern in each of these knockout variants, the

MIC data indicates that each disulfide bond contributes to the antifungal activity of NCR13_PFV1

(Fig 5C and D). Indeed, a NCR13 construct with all six Cys residues replaced with Ser (NSR13)

was inactive against *B. cinerea* even at the highest tested concentration, 24 µM (Fig 5C and D, S4

Fig). It is likely that NSR13 is unfolded, as suggested by the NMR data for reduced NCR13 (Fig

2) and its late elution time on a reverse phase column. Therefore, it may be concluded that each

disulfide bond contributes to the antifungal activity of the NCR13.

Fig 5. Removing disulfide bonds in NCR13_PFV1 strongly reduces antifungal activity
against *B. cinerea in vitro*. (A) Four NCR13 constructs are used to assay the importance of
disulfide bond formation for antifungal activity by substituting pairs of Cys residues with Ser
(highlighted in red). The numbers above the sequence indicate the position of the Cys substitution.

(B) HPLC analyses of NCR13_C4S-C23S, NCR13_C10S-C28S, and NCR13_C15S-C30S
recombinantly produced in *P. pastoris* monitored at 260 nm (mAU = milli-Absorbing Units). (C)
Fungal cell viability assay of disulfide knockout variants. A color change from blue to pink/clear
indicative of metabolically active (live) *B. cinerea* germlings after 60 h. (D) Summary of the MIC
values for NCR13_PFV1 and NCR13 disulfide knockout variants.

Identification of the sequence motif governing the antifungal activity of NCR13

To identify the sequence motif governing the antifungal activity of NCR13 against *B. cinerea*, we

purchased chemically synthesized alanine scanning variants NCR13_AlaV1 through

NCR13_Ala5 (Fig 6A). These variants contained a string of three to five consecutive Alanine (Ala)

residues substituted throughout the NCR13 sequence. We then compared their antifungal activity

relative to a chemically synthesized NCR13. Notably, our findings revealed that NCR13_AlaV2,

NCR13_AlaV3, and NCR13_AlaV4 had no antifungal activity against *B. cinerea* at the highest

tested concentration, 12 µM (Fig 6A, S6A Fig), whereas NCR13 inhibited this pathogen at an MIC

of 3 µM. This data suggested that a 13-residue, Lys-rich motif, KKFA**C**RKPKVPK**C,** located in

the random coil and β-strand region, was a major determinant of NCR13 antifungal activity. We

further explored the importance of this motif by testing antifungal activity of NCR13_AlaV3

expressed in *P. pastoris* to allow formation of disulfide bonds and found that it also lacked

antifungal activity against *B. cinerea* like the chemically synthesized NCR13_AlaV3 (S6B Fig). It

is noteworthy that NCR13_AlaV1 and NCR13_AlaV5 also displayed 2- and 4-fold reduction,

respectively, in the antifungal activity against *B. cinerea* indicating that the N-terminal QSDKD

and C-terminal INGF motifs are relatively minor contributors to the antifungal activity of this

peptide. (Fig 6A).

A

Peptide	Sequence	MIC (μM)
NCR13_CS	TKPCQSDKDCKKFACRKPKVPKCINGFCKCVR	3
NCR13_AlaV1	TKPC**AAAAA**CKKFACRKPKVPKCINGFCKCVR	6
NCR13_AlaV2	TKPCQSDKDC**AAAA**CRKPKVPKCINGFCKCVR	Inactive (>12)
NCR13_AlaV3	TKPCQSDKDCKKFAC**AAAA**VPKCINGFCKCVR	Inactive (>12)
NCR13_AlaV4	TKPCQSDKDCKKFACRKPK**AAA**CINGFCKCVR	Inactive (>12)
NCR13_AlaV5	TKPCQSDKDCKKFACRKPKVPKC**AAAA**CKCVR	12

B

Fig 6. Active motif of NCR13 required for antifungal activity. (A) Antifungal activity of NCR13
synthetic peptide and NCR13 alanine variants against *B. cinerea*. Alanine scanning mutagenesis
was done by substituting alanine within the NCR13 core sequence, leading to the creation of the
NCR13_AlaV1-V5. MIC values were determined against *B. cinerea*. The amino acid substitutions
are shown in red. Representative pictures are shown in S5 Fig. (B) Active motif of NCR13
(highlighted with a bold red font) required for its antifungal activity against *B. cinerea*. The
secondary structure elements are shown above the amino acid sequence. The gray line and green
arrow indicate α-helix (α1) and β-strand (β1) respectively.

NCR13_PFV1 disrupts the plasma membrane of *B. cinerea* faster than NCR13_PFV2

Since NCR13_PFV1 exhibits ~17-fold more potent antifungal activity against *B. cinerea* than

NCR13_PFV2, we hypothesized that the difference in antifungal activity lies in differential

membrane permeabilization ability. To test this hypothesis, we performed SYTOX^TM Green (SG)

assays to examine the ability of NCR13_PFV1 and NCR13_PFV2 at 0.09 μM to permeabilize the

plasma membrane of *B. cinerea* germlings. SG, a membrane-impermeant dye, binds to nucleic

acids and is an indicator of membrane permeability. Average fluorescence intensities were

measured along different time points and NCR13_PFV1 showed much higher fluorescence

intensity within exposed cells than NCR13_PFV2 (Fig 7A and B). After 10 min, SG uptake was

observed in germlings challenged with NCR13_PFV1 but not in germlings challenged with

NCR13_PFV2. After 20 min, a very strong SG uptake signal was observed in NCR13_PFV1

treated germlings, but only a modest signal in NCR13_PFV2 treated germlings (Fig 7A and B).

Thus, NCR13_PFV1 inhibited fungal growth to a greater degree than NCR13_PFV2 at nanomolar

concentration in part by permeabilizing the plasma membrane of *B. cinerea*.

NCR13_PFV1 and NCR13_PFV2 reveal quantitative differences in their binding to multiple

membrane phospholipids

Antifungal plant defensins gain entry into fungal cells and bind to a variety of bioactive

phospholipids resident on the plasma membrane (9). We speculated that NCR13_PFV1 and

NCR13_PFV2 differ in their phospholipid binding specificity in the context of membrane bilayer.

Therefore, we used phospholipid strip assays to determine if NCR13_PFVs bind to different

phospholipids. Our findings revealed that both NCR13_PFV1 and NCR13_PFV2 interacted with

several phospholipids, including phosphatidylinositol 4-phosphate [PI(4)P] and PI(4,5)P_2 (Fig

7C). NCR13_PFV2 displayed significantly stronger binding affinity to phosphoinositides such as

PI(3)P, PI(4)P, PI(5)P, PI(4,5)P_2, and PI(3,4,5)P_3 than NCR13_PFV1 (Fig 7D). The interaction of

PI(4)P and PI(4,5)P_2 with NCR13_PFV1 and NCR13_PFV2 were further validated using a

polyPIPosome binding assay (Fig 7E). Nevertheless, the significance of the phospholipid binding

differences between the two peptides for their antifungal activity requires further investigation.

Since several antifungal plant defensins bind to membrane phospholipids to oligomerize and

induce membrane permeabilization (28), we tested the ability of NCR13_PFV1 and NCR13_PFV2

to oligomerize by incubating the peptides with the biochemical cross-linker

bis(sulfosuccinimidyl)suberate (BS₃) in the presence and absence of PI(4,5)P₂. NCR13_PFV1

formed oligomers both in the presence and absence of PI(4,5)P2. On the other hand, NCR13_PFV2

formed higher order oligomers, but, only at higher concentrations (115 µM and 330 µM) of

PI(4,5)P₂ (S7 Fig). Therefore, further investigation is required to determine whether phospholipid

binding, including PI(4,5)P2, and the oligomerization of NCR13_PFVs are critical for antifungal

activity.

Fig 7. NCR13 PFV1 permeabilizes the cell membrane of *B. cinerea* rapidly. (A) Confocal

microscopy images of SYTOX Green (SG) uptake in *B. cinerea* germlings at different time points.

Germlings were treated with 0.09 µM NCR13_PFVs and SG dye simultaneously for 20 min. No peptide is used as a control. Scale bars = 10 µm. The experiment was repeated thrice with similar results. (B) Quantitative measurements of average fluorescence intensity over time to assess cell membrane permeability in *B. cinerea* germlings treated with 0.09 µM NCR13 PFVs. (C) PIP strip showing binding of NCR13 PFVs to multiple phospholipids, including phosphatidylinositol 4,5-bisphosphate (PI(4,5)P_2), phosphatidylinositol (3,4,5)-trisphosphate PI(3,4,5)P_3 and phosphatidylinositol monophosphates PI(3)P, PI(4)P, and PI(5)P, phosphatidylinositol bisphosphate PI(3,4)P_2. (D) Integrated density measurement of PIP strip probed with NCR13_PFVs. Datapoints are means ± SE of two independent biological replicates ($N = 2$). (E) Binding of NCR13_PFVs to PI(4)P and PI(4,5)P2 (PIP$_2$) containing liposomes. NCR13_PFV's were incubated without liposomes (None) and liposomes bearing no phosphoinositides (control), S- supernatant fraction and P- pellet fraction.

NCR13 PFV1 is internalized faster in germling cells of *B. cinerea* than NCR13_PFV2

Due to its much greater antifungal activity *in vitro* against *B. cinerea*, we hypothesized that NCR13_PFV1 exhibited faster entry into fungal cells at its MIC than NCR13_PFV2. To test this hypothesis, we exposed *B. cinerea* germlings to 0.09 µM DyLight550-labeled NCR13_PFV1 and NCR13_PFV2 and monitored their entry into germlings of *B. cinerea* using confocal microscopy. Within ~30 min, NCR13_PFV1 penetrated the germlings; however, NCR13_PFV2 did not enter the germlings (Fig 8A and B). At a concentration of 1.5 µM, NCR13_PFV1 started entering germlings in ~4 min, while NCR13_PFV2, after briefly gathering on the cell surface (Fig 8C, white arrowheads), started entering germlings at ~8 min. Unlike NCR13_PFV2's lack of entry at 0.09 µM, delayed entry of NCR13_PFV2 was evident at 1.5 µM (Fig 8C and D).

Following entry into *B. cinerea* germling cells, both NCR13_PFV1 and NCR13_PFV2 accumulated in distinct subcellular foci (Fig 8E, white arrows). Further localization studies using the nuclear staining dye Hoechst 33258 revealed that both peptides were localized in the nuclei.

To determine if each peptide was sub-localized within the nucleolus, we used the nucleolus-

specific RNA staining dye Nucleolus Bright Green. As shown in Fig 8F, each peptide was localized

within the nucleolus of germling cells suggesting that each peptide likely binds to ribosomal RNA

(rRNA) or proteins and interferes with protein synthesis.

To gain further insight into the MoA of NCR13_PFV1 and NCR13_PFV2, we assessed the

stability of each peptide once inside *B. cinerea* cells. We challenged *B. cinerea* germlings with

each peptide separately for two different time points, 2 h and 4 h, and tested the stability of each

peptide in fungal cells through western blot analysis. An NCR13 polyclonal antibody capable of

binding to each peptide with equal affinity was used for this analysis (S8 Fig). NCR13_PFV1

accumulated in germling cells within 2 h and accumulates further after 4 h. In contrast, much less

NCR13_PFV2 accumulated in germling cells within 2 h and was strongly decreased by 4 h (Fig

8E). Therefore, we highlight NCR13_PFV1 more rapidly enters fungal cells than NCR13_PFV2,

and once inside NCR13_PFV1 is likely more stable than NCR13_PFV2.

Fig 8. Internalization and localization of NCR13 PFVs in *B. cinerea* germlings. (A) Confocal microscopy images of *B. cinerea* germlings at different points treated with 0.09 μM DyLight550-labeled NCR13_PFV1 (left) and NCR_PFV2 (right) for 30 minutes. Scale bars = 10 μm. Data represents representative results of three independent experiments. (B) Quantification of peptide internalization was conducted by analyzing the average fluorescence intensities over time (in minutes) for NCR13_PFVs. The upper threshold for background was set at 23 for Fiji analysis. (C) Confocal microscopy images of *B. cinerea* germlings treated with 1.5 μM DyLight550-labeled NCR13_PFVs for 30 minutes. White arrowheads indicate NCR13_PFV2 on the cell surface. Scale bars = 10 μm. (D) Quantitative measurements of peptide internalization in *B. cinerea* germlings treated with 1.5 μM NCR13_PFV1 (left) and NCR13_PFV2 (right). (E) Internalized DyLight550-labeled NCR13_PFV1 (left) and NCR13_PFV2 (right) concentrate in specific foci (white arrowheads) within *B. cinerea* germlings. Scale bars = 10 μm. (F) NCR13_PFV1 and NCR13_PFV2 localize to the nucleolus. DyLight550-labeled NCR13_PFV1 (upper panel) and PFV2 (lower panel) colocalize with rRNA-specific Nucleolus Bright Green stain. Hoechst 33258 is used for staining nucleus. White arrowheads indicate nucleolar localization. Scale bars = 5 μm. (G) Semi-quantitative analysis of NCR13_PFVs within *B. cinerea* germlings. Germlings were treated with 1.5 μM NCR13_PFV1 and NCR13_PFV2 for 0, 2, and 4 hours. Total proteins were extracted from fungal cells and immunoblotted using anti-NCR13 antibody. Equal loading of proteins was verified using Ponceau staining. The bar graph represents integrated density measurements. Statistical significance was tested using paired student's t-test. Data are shown as mean ± SD (N = 3, where N refers to biological replicates).

NCR13_PFV1 binds to rRNA and is a more potent inhibitor of protein translation *in vitro* than NCR13_PFV2

Since NCR13_PFVs are localized in the nucleoli of *B. cinerea*, we decided to test the binding of each peptide to rRNA. This was accomplished with an Electrophoretic Mobility Shift Assay (EMSA) (29) by incubating each peptide with total rRNA isolated from *B. cinerea* germlings. Our results showed that NCR13_PFV1 retarded the mobility of the rRNA at all concentrations greater than 0.18 μM, while NCR13_PFV2 showed no binding to rRNA until 3 μM (Fig 9A, S8 Fig). Therefore, it is evident that NCR13_PFV1 exhibits stronger binding affinity to fungal rRNA than NCR13_PFV2.

Next, we tested the ability of the NCR13_PFVs to inhibit protein translation *in vitro* because we

hypothesized that NCR13_PFV1 may be a more potent inhibitor of protein translation than

NCR13_PFV2. This was accomplished using a wheat germ *in vitro* translation system based on a

luciferase activity (30). Protein translation inhibitory activity of each peptide was concentration

dependent (Fig 9B). NCR13_PFV1 inhibited translation more effectively than NCR13_PFV2 at

concentrations more than 0.75 µM. At 24 µM, NCR13_PFV1 almost completely halted translation

compared to the ~50% inhibition by NCR13_PFV2 (Fig 9B). Based on these results, the difference

in the antifungal potency of NCR13_PFVs may be partly due to the inhibition of protein translation

in vivo with NCR13_PFV1 being a more potent protein translation inhibitor in *B. cinerea*.

Fig 9. NCR13_PFV1 exhibits rRNA binding and higher translation inhibition activity. (A)
rRNA binding by NCR13_PFVs tested using electrophoretic mobility shift assay (EMSA). *Botrytis*
cinerea rRNA was used to assess the binding activity of the peptides NCR13_PFV1 and

NCR13_PFV2. The binding reactions were subjected to electrophoresis, and the mobility shifts
were visualized on an agarose gel (1%). The concentration gradient of peptides is indicated above
the lanes. The first lane on the left contains a molecular weight marker. (B) Relative *in vitro*
translational inhibition (%) in the presence of NCR13_PFVs at different concentrations. Sterile
water was used as negative control and cycloheximide (96 µM) was used as positive control. N=
3, 'N' indicates biological replicates. Statistical significance is determined using One-way ANOVA
with Tukey multiple comparisons test (**** indicates p value < 0.0001).
502

NCR13_PFV1 is more potent than NCR13_PFV2 semi-*in planta*

Next, we compared the semi-*in planta* antifungal activity of NCR13_PFV1 and NCR13_PFV2

using detached leaves of pepper as previously described (31). To eliminate leaf-to-leaf variation,

both peptides were added instantaneously with freshly prepared *B. cinerea* spores to the same leaf,

along with a no-peptide control, at different concentrations (3, 1.5, and 0.09 µM) (S9A Fig). Leaves

were evaluated for reduction of gray mold lesions through measurement of photosynthetic

efficiency Fv/Fm (variable fluorescence over saturation level of fluorescence) using CropReporter

images. At 3 µM and 1.5 µM NCR13_PFVs, no disease symptoms were visualized (S9B Fig) and

no significant difference in photosynthetic efficiency (S9C Fig) was observed, whereas the no-

peptide control contained visible lesions and reduced photosynthetic efficiency. At 0.09 µM, only

NCR13_PFV1 effectively suppressed the formation of visible lesions, while NCR13_PFV2

resulted in lesion sizes similar to the no-peptide control (S9B Fig). Moreover, a decrease in

photosynthetic efficiency observed for NCR13_PFV1 was milder than observed for

NCR13_PFV2. Together, this data further illustrates that NCR13_PFV1 is a more potent antifungal

agent than NCR13_PFV2 in terms of preventing and reducing gray mold lesions on peppers (S9B

and C Fig). These observations correlate well with the *in vitro* antifungal activity of each peptide.

Based on this comparative study of their antifungal activity using a detached leaf assay,

NCR13_PFV1 was judged to be a more promising peptide-based fungicide *in planta* (see below).

Spray-applied NCR13_PFV1 is more effective in reducing gray mold disease symptoms in pepper and tomato plants

To determine if the *in vitro* and semi-*in planta* antifungal activity of these peptides translated to antifungal activity *in planta*, we tested the curative protection afforded by NCR13_PFV1 and NCR13_PFV2 against *B. cinerea* by spray application on pepper plants. Four-week-old pepper plants were pre-inoculated with freshly prepared *B. cinerea* spores. At 8 h post-inoculation, peptide at 1.5 and 3 µM concentration was sprayed onto pepper leaves. At 72 h post-inoculation. disease symptoms were evaluated by comparing photosynthetic efficiency (Fv/Fm) of the pathogen-challenged peptide-sprayed plants with the no-peptide control. Green areas indicate a high photosynthetic rate and healthy tissue, while yellow to red areas indicate reduced photosynthetic rate and diseased tissue. As evidenced by their higher Fv/Fm values, leaves of pepper plants sprayed with NCR13_PFV1 at 1.5 and 3 µM had greater reduction in gray mold lesions than the leaves of pepper plants sprayed with NCR13_PFV2 at the same concentrations (Fig 10A and B). Pepper plants sprayed with 1.5 µM NCR13_PFV2 showed no significant control ($p > 0.05$) of gray mold when compared to the no-peptide control plants suggesting NCR13_PFV2 is unable to reduce gray mold disease symptoms at lower concentration (Fig 10A-C). Correspondingly, the data aligns with the observation that NCR13_PFV1 markedly decreased the percentage of disease severity relative to NCR13_PFV2 at both treatment concentrations (Fig 10C). However, NCR13_PFV2 sprayed at 3 µM was able to provide significant curative protection compared to control plants (Fig 10A-C), but it was not as effective as NCR13_PFV1. A similar observation was made when four-week-old tomato plants were also pre-inoculated with *B. cinerea* spores followed by spraying 3 µM of each peptide (S10A-C Fig). The NCR13_PFVs significantly decreased the disease severity, with NCR13_PFV1 demonstrating the most pronounced effect (S10B Fig).

544 NCR13_PFV1 and NCR13_PFV2 inhibited gray mold disease 90% and 40%, respectively (S10C

545 Fig). In summary, these data indicate the higher curative antifungal potential of NCR13_PFV1 for

546 managing gray mold disease in pepper and tomato plants.

Fig 10. NCR13 PFV1 exhibits enhanced curative antifungal activity against gray mold in
pepper (*Capsicum annuum*) plants. (A) Curative antifungal activity of NCR13_PFV1 and PFV2.
Four-week-old pepper plants were first spray-inoculated using 1 mL of 5×10^4 *B. cinerea* spores.

At 8 hpi (hours post inoculation), the plants were sprayed with 2 mL of peptide (NCR13_PFV1 or PFV2) at a concentration of 1.5 or 3 µM or control (*B. cinerea* alone). Mock-treated plants were only sprayed with 1X SFM (no fungal spore suspension). (B) Calculated photosynthetic quantum yield (Fv/Fm) for individual pepper plants. (C) Disease severity (%) was calculated as described in materials and methods. (B-C) Each data point represents the average of 4 leaves per plant for 4 plants for each treatment. Horizontal lines represent the median and boxes indicate the 25th and 75th percentiles. The Kruskal-Wallis test with Dunn's multiple comparisons test is used to determine statistically significant differences between control and peptide-treated inoculated plants. Two independent experiments were performed with similar results.

Discussion

NCR13 is one of the highly cationic NCR peptides among the 63 found in chickpea nodules (10). In this study, we discovered that NCR13 expressed in *P. pastoris* forms two isomers, NCR13_PFV1 and NCR13_PFV2, which have identical amino acid sequence and molecular mass but differ in their disulfide pairing patterns. We report a comparative analysis of the structure, *in vitro* and semi-*in planta* antifungal activity and MoA of NCR13_PFVs. Our results support the proposal that disulfide cross-linking provides an additional source of structural and functional diversity in NCR peptides (21).

Four- or six-cysteine NCRs are highly diverse in their primary sequences and are predicted to be structurally different. Previously, two solution NMR structures of NCR044 and NCR169, each containing two disulfide bonds, were solved (19, 23). The NCR044 structure was largely disordered, dynamic and contained a single α-helix and a β-sheet (19). Structural analysis revealed that the two NCR169 isomers contained only one short, two-stranded β-sheet, with one isomer being more extended and flexible while the other was more compact and constrained (23). To date, a three-dimensional structure for a six-cysteine NCR peptide has not been reported. Here, we report the first solution structures of two NCR13 disulfide isomers each containing an α-helix and β-sheet. Nevertheless, unique pairing of two out of the three disulfide bonds significantly alters how

these secondary structure elements fold upon each other rendering NCR13_PFV1 more compact

and resistant to digestion by trypsin or reduction by dithiothreitol. It is striking that *P. pastoris*

generates two disulfide variants that are structurally different and raises the question of whether

these variants are also expressed in the differentiating nodules of chickpea to perform different

biological activities. In an earlier study, three disulfide variants of NCR247 containing four

cysteines were generated using chemical synthesis and *in vitro* folding. NCR247 variants were

found to have distinct patterns of biological activities such as bacterial gene expression, inhibition

of bacterial cell division, protein translation and antibacterial activity (21, 22). These observations

suggest that NCR peptides can form disulfide variants *in vitro* or in an appropriate expression host,

leading to diverse biological activity. Whether disulfide variants of NCR13, NCR247, or NCR169

are formed *in vivo* in nodules to control symbiosis in IRLC legumes remains an open question. In

contrast to NCR peptides, plant defensins with eight cysteines expressed in *P. pastoris* are only

observed as one species with an invariant disulfide pairing pattern (32).

To our knowledge, this study is the first detailed investigation of a chickpea NCR peptide and

its antifungal action. Our data demonstrates that disulfide folding variants of NCR13 produced in

P. pastoris exhibit profound differences in antifungal activity against multiple fungal pathogens of

economic importance. NCR13_PFV1 possesses highly potent antifungal activity against *B. cinerea*

and other pathogens, showing a four- to sixteen-fold greater activity compared to NCR13_PFV2.

Remarkably, we found that the antifungal activity of NCR13_PFV1 is greater than other NCR

peptides reported to date in literature (33). It is noteworthy that enhanced antifungal activity of

both NCR13_PFVs produced in *P. pastoris* compared to chemically synthesized NCR13 without

disulfide bond, underscores the critical role of disulfide bonds for antifungal activity. Furthermore,

the formation of all three disulfide bonds is essential for highly potent antifungal activity of

NCR13_PFV1 as removal of each disulfide bond resulted in a significant loss of its antifungal

activity against *B. cinerea*. In addition, Cys residues are essential for the antifungal activity of

NCR13 as their replacement with Ser leads to complete loss of antifungal activity. Acting as

molecular anchors, disulfide bonds provide stability to the three-dimensional structure of proteins

and peptides. They set the constraints for distance and angles between connected cysteine residues,

thus favoring the stability of the protein's folded state over its unfolded form (34). Furthermore,

the insertion of cysteine residues into *de novo* designed peptides has produced hyperstable peptides

with unique tertiary structures that form the basis for new peptide-based drugs (35, 36). The impact

on antifungal activity of knocking out two instead of just one disulfide bond is yet to be

investigated. Further, evaluating the three-dimensional structures of these double disulfide bond

knockout variants will provide deeper insight into structure-activity relationships of this unique

NCR peptide. *P. pastoris* has been the eukaryotic organism of choice for producing cysteine-rich

peptides to ensure formation of the correct disulfide bond patterns (37). It is intriguing that NCR13

expression results in the formation of two disulfide folding variants in *P. pastoris*. In this context,

it is critical to observe that knocking out either the C4-C23 or C15-C30 disulfide bond in

NCR13_PFV1 results in the formation of one peak on HPLC, whereas knocking out the C10-C28

bond still results in the formation of two distinct peaks. Mass spectrometry analysis shows that

peptides from two HPLC peaks of the C10-C28 knockout and one HPLC peak in the C4-C23 and

C15-C30 knockouts are in the oxidized state. Therefore, it is likely they are in a folded/oxidized

form. Finally, understanding the molecular mechanisms governing the generation of disulfide

variants of specific antifungal NCR peptides could pave the way for production of targeted

disulfide structural variants in *P. pastoris* with enhanced antifungal activity.

Antifungal peptides with potent fungicidal activity and multiple MoA are attractive candidates

for development as spray-on fungicides. NCR13 has a promise as one such candidate. For instance,

ability to permeabilize the plasma membrane of fungal pathogens is a common MoA of plant

cysteine-rich antifungal peptides including NCRs (14). NCR13_PFV1 can permeabilize the

plasma membrane of *B. cinerea* rapidly at very low concentrations. The membrane disrupting

ability of NCR13_PFV1 is far greater than that of NCR13_PFV2 and NCR044 (19). Both

NCR_PFVs bind to the same phospholipids on lipid strips with different binding intensity.

However, it is not clear what role, if any, phospholipid binding plays in conferring increased

membrane disrupting ability of NCR13_PFV1 and this aspect remains to be elucidated.   In

addition, NCR13_PFV1 gains rapid entry into the interior of fungal cells compared to

NCR13_PFV2. It is plausible that more compact folding of NCR13_PFV1 allows it to more

quickly penetrate the cell wall and pass through the outer and inner leaflets of the plasma

membrane. This is supported by the fact that *B. cinerea* germlings challenged with 1.5 µM of each

peptide showed NCR13_PFV1 enters fungal cells in much greater levels after 2 h than

NCR13_PFV2. Moreover, once inside the cells, NCR13_PFV1 accumulates stably for at least 4 h

whereas NCR13_PFV2 tends to degrade in this time. Greater stability of NCR13_PFV1 to

proteolysis might be a major contributing factor to its much greater antifungal activity against this

pathogen, an observation that warrants further investigation.

Subcellular localization studies showed that NCR_PFVs are targeted to nucleoli in the

germlings of *B. cinerea* suggesting that these peptides are targeting ribosomal biogenesis or protein

translation machinery as part of their MoA as proposed earlier for the MoA of NCR044 (19). In

support of this hypothesis, we found that addition of each NCR13_PFVs to the *in vitro* translation

system inhibited protein synthesis in a concentration dependent manner. NCR13_PFV1 proved to

be more effective in inhibiting protein translation *in vitro*. NCR13_PFV1 also binds to *B. cinerea*

rRNA at a lower concentration (0.37 µM) than NCR13_PFV2 (6 µM). Further research is needed

to determine if differences in rRNA binding affinity and protein translation inhibition *in vitro*

directly translate to profound differences in their antifungal activity *in vivo*. NCR peptides appear

to target protein translation as part of their antimicrobial mechanism as evidenced by *M. truncatula*

NCR247 inhibiting protein translation *in vitro* by binding to ribosomal proteins (21, 38). Plant

defensin MtDef4 also inhibits protein translation *in vitro* by binding to ribosomal proteins, but not

to ribosomal RNA (39). Besides that, it is likely that NCR13_PFV1 and NCR13_PFV2 have

different, yet unidentified, intracellular targets which will be the subject of future investigation.

Fungicides currently used commercially for management of fungal diseases target ergosterol

biosynthesis, succinate dehydrogenase and cytochrome bc_1 complex. These targets are prone to

rapid development of fungal resistance (40). Small NCR peptides with multi-faceted MoA have

emerged as potential candidates for development as bioinspired fungicides. We have previously

demonstrated efficacy of NCR044 as a spray-on fungicide for management of gray mold in tomato

(19). Here, we report NCR13 is also a multi-target antifungal peptide that reduces gray mold

disease symptoms curatively when sprayed on pepper and tomato plants. Consistent with its *in*

vitro and semi-in planta antifungal activity, NCR13_PFV1 affords greater protection against gray

mold disease than NCR13_PFV2 in both pepper and tomato plants. NCR13 is amenable to large

scale recombinant production in a bio-fermenter using *P. pastoris* expression system. However,

further characterization of NCR13 is needed, including demonstrating its efficacy for control of

gray mold and other diseases in large-scale greenhouse and field tests, assessment of off-target

activity against mammalian cells and phytotoxicity. In summary, this study demonstrated that

disulfide bond pairing pattern has a profound impact on the *in vitro* antifungal activity, MoA, and

in planta antifungal activity of chickpea NCR13. As illustrated in Fig 11, the graphical abstract

illustrates the key findings of our study. Overall, our work underscores the significance of disulfide

cross-linking in enhancing the efficacy of antifungal peptides for agricultural use.

Fig 11. Graphical abstract illustrating the biosynthesis of disulfide linkage variants of NCR13

675 (PFV1 and PFV2) in *P. pastoris*, their structures, antifungal activity and modes of action in *B.*

676 *cinerea* germlings. The figure depicts the differences in cell penetration, localization, rRNA

677 binding and translation inhibition, membrane permeabilization, phospholipid binding, and overall

678 fungal cell killing potency. Created using BioRender.com.

Materials and Methods

Fungal Cultures and Spore Suspension

B. cinerea strain T4 was cultured in V8 medium for 3 weeks at 25°C. Fungal spores were collected by flooding the fungal growth plates with water and scraped using sterile spreader. The spore suspension was filtered through two layers of Miracloth, followed by centrifugation at 12,000 g for 2 minutes (min) and subsequently washed twice with sterile water. Lastly, fungal spores were resuspended in Synthetic Fungal Media (SFM) (41). The spore suspension was adjusted to the desired spore density of about 10^5 spores/mL using a hemocytometer. *Fusarium virguliforme (F. virguliforme)* was grown and maintained regularly in potato dextrose agar (PDA) (Difco Laboratories) medium. Two weeks old *F. virguliforme* plates were used to collect spores as described above for *B. cinerea* followed by centrifugation and washing step. Spores were counted using a hemocytometer and adjusted to the final concentration of 10^5 spores/mL in 2X SFM media. *Sclerotinia sclerotiorum* obtained from the field of pennycress in Illinois was grown on PDA medium at room temperature.

Expression and purification of NCR13

For expression in *P. pastoris*, the published amino acid sequence of the mature NCR13 peptide (24) was used for synthesis of the codon-optimized gene (GenScript). The NCR13 gene was expressed in *P. pastoris* X33 using either a SacI linearized pPICZα-A integration vector (Invitrogen) or at XhoI and XbaI restriction sites. NCR13 was purified using a CM Sephadex C-25 cation-exchange chromatography as described (19). After induction, cell-free supernatant was collected after centrifugation (10,000×g) at 4 °C for 20 min, and the pH was adjusted to 6.0. The supernatant was passed through cation-exchange resin, CM Sephadex C-25 (Sigma-Aldrich, Cat No: C25120) previously equilibrated with binding buffer (25 mM Sodium Acetate Anhydrous, pH

6.0) using an AKTA FPLC. Peptides bound to the resin were eluted using the elution buffer (1M

NaCl, 50 mM Tris, pH 7.6). The FPLC fraction containing NCR13 was concentrated and desalted

using an Amicon Stirred Cell with a 3-kD cutoff membrane (Millipore, Cat No: PLBC04310) and

further purified by reverse phase C18-HPLC. HPLC fractions collected separately for

NCR13_PFV1 and NCR13_PFV2 were lyophilized and re-suspended in nuclease-free water.

Peptide concentrations were determined using the Pierce™ BCA Protein Assay Kit (Thermo

Scientific). NCR13 and alanine variants of NCR13 were chemically synthesized (Biomatik) and

further purified by in-house HPLC.

Molecular weight determination

Purified peptides were analyzed by LC-MS for accurate mass determination. Firstly, peptide

samples were cleaned using Thermo Scientific HyperSep C18 microscale SPE extraction tips and,

after dry-down, resuspended in 0.1% formic acid for LC-MS analysis. LC-MS was performed with

a Dionex RSLCnano HPLC coupled to an OrbiTrap Fusion Lumos (Thermo Scientific) mass

spectrometer using 80 min gradient (1-90% acetonitrile). Sample was resolved using a 75 μm x 15

cm PepMap C18 column (Thermo Scientific). MS spectra of protein ions of different charge-states

were acquired in positive ion mode with a resolution setting of 120,000 (at m/z 200). Mass

deconvolution was performed using ProteinProspector, Xcalibur (Thermo Scientific), and ESIprot

Online.

Disulfide bond pattern identification for NCR13_PFV2 using mass spectrometry

An aliquot of unlabeled, oxidzed NCR13_PFV2 was mixed with 20-fold less (in mass) trypsin,

incubated at room temperature for 1 h, and diluted in 50% acetonitrile, 0.1% formic acid to ~0.1

μM. The diluted samples were infused at 2 μL/min into a Thermo Q-Exactive Exploris 480

Orbitrap mass spectrometer with positive mode electrospray at 3.4 kV. The ion transfer tube was

at 320 °C, sheath gas at 2 psi, and auxiliary gas at 5 psi. Data-dependent acquisition was used

(MS1 resolution 60k, MS2 resolution 30k) over 8 min. Fragmentation spectra were collected with

higher-energy collisional dissociation (HCD) at 30% collision energy. Data were analyzed and

deconvoluted in FreeStyle (Thermo Scientific). Crosslinked peptides were identified using Byonic

(ProteinMetrics). Under the same conditions, NCR13_PFV1 could not be digested by trypsin.

^{15}N isotopic labeling of NCR13 for NMR structural analysis

The ^{15}N-labeled NCR13_PFVs were prepared according to the protocol as described previously

(32) with minor modifications. NCR13-expressing *P. pastoris* X33 was grown at 28 ± 2°C for two

days on a YPD plate containing 150 μg/mL Zeocin. A single colony was then inoculated into 5 mL

of YPD broth and grown overnight at 28 ± 2 °C on a rotary shaker at 225 ± 25 rpm. After overnight

growth, a 5 mL culture was transferred aseptically to 100 mL of FM23 medium (0.8% YNB

without amino acids and ammonium sulfate, 2 mg/L biotin, 1.2% $(NH_4)_2SO_4$, 0.3% K_2HPO_4,

0.28% KH_2PO_4, 2 ml/L PTM1 salts) containing 3% glucose. After 24 h, an additional 2% glucose

was added to the culture medium. After a growth phase of ~30 h and just prior to methanol

induction, the culture medium was adjusted to 0.02% ^{15}N-ammonium chloride (^{15}NH$_4$Cl) and 0.1%

glucose. After 40 h, cells were harvested by centrifugation at 5,000 rpm at room temperature for

10 min, briefly washed with 0.2% glycerol, pelleted, and re-suspended in 100 mL FM23 medium

containing 1.2% ^{15}NH$_4$Cl and grown for 4-5 days at 25 °C at 225 ± 25 rpm. In the induction phase,

a low concentration of 0.2% methanol was initially added to the culture medium to induce protein

expression and gradually increased by 0.4% from 72 to 120 h (1.4% final). The labeled NCR13

peaks were purified from the *P. pastoris* growth medium following the same protocol used for the

unlabeled peptide.

NMR spectroscopy of NCR13

All NMR data were collected at 20˚C on ^{15}N-labeled samples of oxidized NCR13_PFV1 and NCR13_PFV2 (~ 1 mM, 20 mM sodium acetate, 50 mM NaCl, pH 5.3) using Varian (Inova-600) or Bruker (Advent-750) spectrometers equipped with an HCN-probe and pulse field gradients. Except for the carbonyl resonances (which could not be assigned because making ^{13}C-labeled samples in *P. pastoris* was too expensive), the ^{1}H, ^{13}C, and ^{15}N chemical shifts of the backbone and side chain resonances were assigned from the analysis of two-dimensional ^{1}H-^{15}N HSQC, ^{1}H-^{13}C HSQC, ^{1}H-^{1}H TOCSY, ^{1}H-^{1}H NOESY, and ^{1}H-^{1}H COSY spectra and three-dimensional ^{15}N-edited NOESY-HSQC (mixing time = 90 ms) and ^{15}N-edited TOCSY spectra. To obtain additional chemical shift assignments, inter- and intra-residue proton side chain restraints for proton near the residual water resonance, most of the two-dimensional data sets were also collected by lyophilizing and resuspending the sample in 99.8% D_2O. In preparing the latter sample, a ^{1}H-^{15}N HSQC spectrum was collected ~10 min after the addition of D_2O to identify slowly exchanging amides that could be used as hydrogen bond restraints in the structure calculations. To confirm oxidized NCR13_PFV1 and NCR13_PFV2 arose from different disulfide bond patterns of a common precursor primary amino acid sequence, both oxidized samples were reduced by making the NMR samples 5 mM in Dithiothreitol (DTT). While NCR13_PFV2 was reduced rapidly, it took more than 5 – 7 days to fully reduce NCR13_PFV1. All NMR data were processed using Felix 2007 (MSI, Molecular Simulations) software and analyzed with the program Poky (v3.115) (42). The assigned ^{1}H, ^{13}C, and ^{15}N chemical shifts were referenced using indirect methods (DSS = 0 ppm) and deposited into the BioMagResBank database (www.bmrb.wisc.edu) under the accession numbers BMRB-31111 (oxidized NCR13_PFV1 8ULM) and BMRB-30979 (oxidized NCR13_PFV2 7TH8).

NMR solution structure calculations for oxidized NCR13_PFV1 and NCR13_PFV2

The protocols used to obtain solution structures for both peptides followed a similar approach starting with the ^{1}H, and ^{15}N chemical shift assignments and peak-picked NOESY data serving as initial experimental inputs in iterative structure calculations with the program CYANA (v 2.1) (43). The ^{1}H, ^{13}C, and ^{15}N chemical shift assignments were also the primary basis for the introduction of 24 (NCR13_PFV1) and 25 (NCR13_PFV2) dihedral Psi (ψ) and Phi (φ) angle restraints identified with the program TALOS+ using the online webserver (https://spin.niddk.nih.gov/bax/nmrserver/talos/) (44). Nine restraints between the side chain sulfur atoms of C4-C23, C15-C30, and C10-C28 (NCR13_PFV1) and C4-C10, C15-C30, and C23-C28 (NCR13_PFV1) (2.0-2.1 Å, 3.0-3.1 Å, and 3.0-3.1 Å for the Sγ-Sγ, Sγ-Cβ, and Cβ-Sγ distances, respectively) (19, 45) were introduced on the basis of AlphaFold predictions for NCR13_PFV1 and high resolution mass spectrometry analysis for NCR13_PFV2. Exhaustive attempts to identify the disulfide bond pattern for NCR13_PFV1 using various mass spectrometric methods failed likely due to its hyperstability to reduction by DTT or tris(2-carboxyethyl) phosphine. Backbone hydrogen bond restraints (1.8 – 2.0 Å and 2.7 – 3.0 Å for the NH–O and N-O distances, respectively) were added to the calculations for proximate backbone amide-carbonyl groups present in early structure calculations that were accompanied with slowly exchanging amides in the deuterium exchange experiment. The final ensemble of 20 CYANA derived structures was then refined in explicit water with CNS (version 1.1) using the PARAM19 force field and force constants of 500, 500, and 1000 kcal for the NOE, hydrogen bond, and dihedral restraints, respectively. For these water refinement calculations, 10 and 0 % were added to the upper boundary of the CYANA distance restraints for NCR13_PFV1 and NCR13_PFV2, respectively, and the lower boundary set to the van der Waals limit for both. Structural quality was

assessed using the online Protein Structure Validation Suite (PSVS, v1.5) (46). Complete structural

statistics are summarized in S3 Table. The atomic coordinates for the final ensemble of 20

structures have been deposited in the Research Collaboratory for Structural Bioinformatics

(RSCB) under PDB codes 8ULM (NCR13_ PFV1) and 7TH8 (NCR13_ PFV2).

Peptide uptake and stability assessment within *B. cinerea*

B. cinerea ($\sim 10^5$ spores/mL) was inoculated in 10 mL of Potato Dextrose Broth (PDB) and

incubated in a shaker at 28ºC for 18 h. *B. cinerea* germlings were treated with 1.5 µM

NCR13_PFV1 and NCR13_PFV2 for 0, 2 and 4 h. Total proteins were extracted from the fungal

cells using 300 µL of HEPES buffer (100 mM HEPES, 5mM EDTA, 0.1% Triton-100, ProBlock™

Gold Yeast/Fungi Protease Inhibitor, pH 7.5) and centrifuged at 14000 rpm to remove cell debris.

The concentration of protein present in the supernatant was measured using Pierce™ BCA Protein

Assay Kit (Thermo Scientific). Total proteins (10 µg) were separated on a 4–20% SDS-PAGE gel

(Bio-Rad, TGX Stain-Free™ precast gels) and transferred to a nitrocellulose membrane. Equal

loading of proteins was checked using Ponceau S staining (Thermo Scientific). Immunoblotting

was performed using the custom synthesized anti-NCR13 polyclonal antibody (Biomatik) at a

concentration of 0.1 µg/mL followed by an incubation in the secondary anti-rabbit antibody

(Cytiva, Cat No: RPN4301) with a dilution of 1:20,000.

***In vitro* antifungal assay**

The antifungal activity of NCR13 against fungal pathogens was determined using the resazurin

cell viability assay as described previously (47). Briefly, 45 µL of each peptide dilution (0, 0.09,

0.1875, 0.375, 0.75, 1.5, 3 and 6 µM) was added to each well of the microtiter plate containing 45

µL ($\sim 10^5$ spores/mL) of spore suspension. Three biological replicates were used for each treatment

and the experiments were repeated three times. Resazurin at 0.1% (w/v) was added to each well

after incubation for 48 h. Images were taken at ~24 h. The Minimum Inhibitory Concentration

(MIC) for each peptide denotes the lowest concentration at which the peptide halts the growth of

fungus evident through a lack of color change from blue to colorless. The *in vitro* antifungal

activity of NCR13_PFVs against *S. sclerotiorum* was determined using a 24-well plate assay as

described (48). Mycelial plugs (1 mm in diameter) cut from the growing edge of the two-day-old

actively growing colony were transferred from PDA media to a series of wells containing 250 µL

of SFM.

Internalization and permeabilization assays using confocal microscopy

NCR13_PFV1 and PFV2 were labeled with DyLight550 amine reactive fluorescent dye following

the manufacturer's instructions (Thermo Scientific). *B. cinerea* germlings (50 µL, ~5x10^4

spores/mL) were treated with DyLight 550 labeled NCR13_PFV1 and NCR13_PFV2 (50 µL, 0.09

µM and 1.5 µM), and the fluorescence measured with excitation and emission wavelengths of 550

and 560-600 nm, respectively. Time-lapse confocal microscopy data with a z-stack at each time

point were acquired using a 63X water immersion objective lens (HC PL Apochromat CS2) at 30-

second intervals. For the membrane permeabilization assay, fluorescent dye SYTOX^{TM} Green

(Thermo Scientific) at a concentration of 0.25 µM was used to determine the membrane

permeabilization of *B. cinerea* germlings as described previously (31). Time lapse imaging was

performed to visualize the SYTOX^{TM} Green entry upon treatment with 50 µL of 0.09 and 1.5 µM

NCR13_PFVs. Imaging was performed on a Leica SP8 confocal microscope with excitation at 488

nm and emission at 508–544 nm. Average fluorescence intensity measurements were obtained

using Fiji (49).

Phospholipid binding assays

Phospholipid binding of NCR13_PFVs were performed with PIP Strips (Echelon Biosciences, Cat No: P-6001) that are spotted with 100 pmol of various biologically active phospholipids. Peptide binding to the membrane strips was detected using monoclonal NCR13 antibody (0.1 µg/mL) followed by goat anti-rabbit IgG HRP (Cytiva RPN4301) at 1:20,000 dilution. The binding signals were detected using the SuperSignal West Pico Chemiluminescent Substrate Kit (Thermo Scientific) following the manufacturer's protocol. Quantitative analysis of the blot intensity was performed using Fiji (49).

Oligomerization assay

To determine the oligomerization status of the NCR13_PFVs, the peptides were prepared in PBS buffer at 1.5 mg/mL (5 µL/reaction) and incubated with specified concentrations of PI(4,5)P2. The cross-linking reaction was initiated by addition of 12.5 mM bis(sulfosuccinimidyl)suberate BS3 (Thermo Scientific) and then incubated for 30 min at room temperature. After incubation, tricine sample buffer was added to each sample and heat denatured at 95 ºC for 5 min. Samples were loaded on a 16.5% Tris-Tricine precast gel (Bio-Rad Laboratories). For SDS-PAGE, 5 µL of Precision Plus Dual Xtra Protein Standards (2–250 kDa) were used (Bio-Rad Laboratories). Oligomerization patterns were visualized using silver staining.

PolyPIPosome binding assay

The validation of the binding of NCR13_PFV1 and NCR13_PFV2 to phospholipids was further performed using a liposome binding assay. In this assay, polyPIPosome binding was conducted in 200 µL of binding buffer (20 mM HEPES pH 7.4, 120 mM NaCl, 1 mM EGTA, 1 mM $MgCl_2$, 1 mg/mL BSA, 0.2 mM $CaCl_2$, 5 mM KCl) plus 10 µL each of control PolyPIPosome, PI4P PolyPIPosomes, and PIP_2 PolyPIPosome (Echelon Biosciences). Subsequently, 2 µg of each

peptide was added, and the mixture incubated for 1 h at room temperature. Following incubation,

liposomes were collected by centrifugation at 16,000 × g for 20 min, and the supernatant was

collected. The pellet was subjected to three washes in 1 mL of binding buffer. After the washes,

the pellet was resuspended in 40 µL of 2X Laemmli sample buffer. Both samples were reduced

with 100 mM DTT and separated on a 4–20% TGX Stain-Free™ precast gels (Bio-Rad

Laboratories). Proteins were then transferred to a nitrocellulose membrane and subjected to

blotting using the NCR13 antibody.

Electrophoretic mobility shift assay

Total RNA was extracted and purified from *B. cinerea* grown in Potato Dextrose Broth (Fisher

Scientific) using the NucleoSpin RNA Plus Mini kit (Takara Bio) according to manufacturer

instructions. The gel shift experiments were performed by mixing 200 ng of the total RNA with

different concentrations of NCR13 PFV1 and PFV2 (0-3 µM) in 20 µL of binding buffer (10 mM

Tris-HCl, pH 8.0, 1 mM EDTA). The reaction mixtures were incubated for 1 h at room temperature

and mixed with 4 µL of 6X gel loading dye (New England Biolabs). Following electrophoresis in

1.2% agarose gel containing SYBR safe DNA gel stain (Thermo Scientific) with TAE buffer at

120 V for 45 min, the gels were visualized using a Bio-Rad ChemiDoc XRS+ system.

In vitro **translation assay**

In vitro translation was performed using the RiboMAX™ Large Scale RNA Production system

and Wheat Germ Extract *in vitro* translation system (Promega). First, the luciferase DNA template

(Promega) was transcribed to mRNA and translated to luciferase following the manufacturer's

protocol. The luciferase enzyme activity was detected by the luciferase kit (Promega) using a

luminometer (Tecan). Different concentrations of NCR13_PFV1 and NCR13_PFV2 were used to

determine their translation inhibition ability. Sterile water was added to the *in vitro* translation

system as a negative control and a fungal translation inhibitor cycloheximide (48 µM) was used as

a positive control. Translation inhibition (TI) percentage is calculated using this formula TI % =

(1− a/b) x 100, where 'a' equals the translation level with peptide and 'b' equals the translation

level without peptide.

892 *In planta* and Semi-*In planta* antifungal assays

Pepper (*Capsicum annuum*, California Wonder 300 TMR) plants used in this study were grown in

a greenhouse for 4 weeks under 14 h light/10 h dark cycle. Tomato (*Solanum lycopersicum* cv.

Mountain Spring) plants were grown in a controlled environment growth chamber for 4 weeks,

under a 16 h light/ 8 h dark cycle. Semi-*in planta* antifungal activity of each peptide against *B.*

cinerea was determined using detached leaves of pepper. NCR13_PFV1 and NCR13_PFV2 were

tested at 6, 1.5, and 0.09 µM. Following incubation of 10 µL peptide with 10 µL of *B. cinerea*

spores (~10^5 spores/mL) under a humid environment for 48 h, the leaves were photographed in

white light. High-resolution fluorescence images were also taken using CropReporter

(PhenoVation). These images depicted the calculated Fv/Fm (maximum quantum yield of

photosystem II) values of the diseased area affected by *B. cinerea* infection. Colors in the images

show five different classes ranging from class I to class V (0.000 to 0.700) depicting varying

degrees of tissue damage.

To test the curative antifungal activity of peptides against *B. cinerea*, 4-week-old pepper and

tomato plants were used. Plants were first sprayed with 1 mL of *B. cinerea* spores (about 5×10^4)

suspended in 1× SFM. After 8 h of *B. cinerea* challenge, 2 mL of NCR13_PFV1 or NCR13_PFV2

(3.0 and 1.5 µM) were sprayed per plant. Control plants were sprayed with 2 mL of water. All

plants were placed in a high humid environment for 72 h. Plants were imaged at 72 h after infection

in white light and high-resolution fluorescence images were taken using CropReporter

(PhenoVation). The quantification of plant health was carried out using the calculated Fv/Fm

(maximum quantum yield of photosystem II) images of the CropReporter showing the efficiency

of photosynthesis in false colors.

For the curative assay using pepper and tomato plants, the picture of the whole plants and detached

leaves were taken as described above. The disease symptoms on individual leaves of each plant,

were visually assessed in % of disease portion (disease severity) and on a rating scale of 0-5 where

each rank would represent the following proportion of diseased tissue per leaf : Scale 0: No obvious

signs/symptoms of disease, Scale 1: $0 > x \le 20$ % diseased portion of leaf, Scale 2: $20 > x \le 40$ %

diseased portion of leaf, Scale 3: $40 > x \le 60$ % diseased portion of leaf, Scale 4: $60 > x \le 80$ %

diseased portion of leaf, Scale 5: $80 > x \le 100$ % diseased portion of leaf. Four plants were used

per treatment and for each plant the disease was assessed per leaf. Besides disease severity (DS),

the inhibition of disease by treatment was also evaluated. The following formula was used to assess

the inhibition of disease severity by each treatment: % Disease Inhibition= (Average DS for

Control -Treatment DS)/(Average DS for Control) x 100%. Where treatment DS is the disease

severity for each leaf except those of the control. Plots were generated using the 'ggplot2' package

in R version 4.3.2.

References

1.   McLaughlin MS, Roy M, Abbasi PA, Carisse O, Yurgel SN, Ali S. Why Do We Need Alternative
Methods for Fungal Disease Management in Plants? Plants (Basel). 2023;12(22).
2.   Goyal RK, Mattoo AK. Multitasking antimicrobial peptides in plant development and host
defense against biotic/abiotic stress. Plant Sci. 2014;228:135-49.
3.   Dos Santos-Silva CA, Zupin L, Oliveira-Lima M, Vilela LMB, Bezerra-Neto JP, Ferreira-Neto
JR, et al. Plant Antimicrobial Peptides: State of the Art, In Silico Prediction and Perspectives in the
Omics Era. Bioinform Biol Insights. 2020;14:1177932220952739.
4.   van der Weerden NL, Bleackley MR, Anderson MA. Properties and mechanisms of action of
naturally occurring antifungal peptides. Cellular and molecular life sciences : CMLS.
2013;70(19):3545-70.
5.   Lobo F, Boto A. Host-defense peptides as new generation phytosanitaries: Low toxicity and
low induction of antimicrobial resistance. Agronomy. 2022;12:1614.
6.   Tavormina P, De Coninck B, Nikonorova N, De Smet I, Cammue BP. The Plant Peptidome: An
Expanding Repertoire of Structural Features and Biological Functions. Plant Cell. 2015;27(8):2095-
118.
7.   Cools TL, Struyfs C, Cammue BP, Thevissen K. Antifungal plant defensins: increased insight
in their mode of action as a basis for their use to combat fungal infections. Future Microbiol.
2017;12:441-54.
8.   Maroti G, Downie JA, Kondorosi E. Plant cysteine-rich peptides that inhibit pathogen growth
and control rhizobial differentiation in legume nodules. Current opinion in plant biology.
2015;26:57-63.
9.   Parisi K, Shafee TMA, Quimbar P, van der Weerden NL, Bleackley MR, Anderson MA. The
evolution, function and mechanisms of action for plant defensins. Semin Cell Dev Biol.
2019;88:107-18.
10.   Montiel J, Downie JA, Farkas A, Bihari P, Herczeg R, Balint B, et al. Morphotype of bacteroids
in different legumes correlates with the number and type of symbiotic NCR peptides. Proc Natl
Acad Sci U S A. 2017;114(19):5041-6.
11.   Kondorosi E, Mergaert P, Kereszt A. A paradigm for endosymbiotic life: cell differentiation of
Rhizobium bacteria provoked by host plant factors. Annu Rev Microbiol. 2013;67:611-28.
12.   Kereszt A, Mergaert P, Montiel J, Endre G, Kondorosi E. Impact of Plant Peptides on
Symbiotic Nodule Development and Functioning. Front Plant Sci. 2018;9:1026.
13.   Mergaert P. Role of antimicrobial peptides in controlling symbiotic bacterial populations.
Nat Prod Rep. 2018;35(4):336-56.
14.   Ordogh L, Voros A, Nagy I, Kondorosi E, Kereszt A. Symbiotic plant peptides eliminate
Candida albicans both in vitro and in an epithelial infection model and inhibit the proliferation of
immortalized human cells. Biomed Res Int. 2014;2014:320796.
15.   Farkas A, Maroti G, Kereszt A, Kondorosi E. Comparative Analysis of the Bacterial
Membrane Disruption Effect of Two Natural Plant Antimicrobial Peptides. Frontiers in microbiology.
2017;8:51.
16.   Tiricz H, Szűcs A, Farkas A, Pap B, Lima RM, Maróti G, et al. Antimicrobial Nodule-Specific
Cysteine-Rich Peptides Induce Membrane Depolarization-Associated Changes in the
Transcriptome. Applied and Environmental Microbiology. 2013;79(21):6737.
17.   Balogh EP, Mosolygo T, Tiricz H, Szabo AM, Karai A, Kerekes F, et al. Anti-chlamydial effect of
plant peptides. Acta Microbiol Immunol Hung. 2014;61(2):229-39.

18.   Alhhazmi AA, Alluhibi SS, Alhujaily R, Alenazi ME, Aljohani TL, Al-Jazzar AA, et al. Novel
antimicrobial peptides identified in legume plant, Medicago truncatula. Microbiol Spectr.
2024;12(2):e0182723.
19.   Velivelli SLS, Czymmek KJ, Li H, Shaw JB, Buchko GW, Shah DM. Antifungal symbiotic
peptide NCR044 exhibits unique structure and multifaceted mechanisms of action that confer
plant protection. Proc Natl Acad Sci U S A. 2020;117(27):16043-54.
20.   Bulleid NJ, Ellgaard L. Multiple ways to make disulfides. Trends Biochem Sci.
2011;36(9):485-92.
21.   Shabab M, Arnold MF, Penterman J, Wommack AJ, Bocker HT, Price PA, et al. Disulfide
cross-linking influences symbiotic activities of nodule peptide NCR247. Proc Natl Acad Sci U S A.
2016;113(36):10157-62.
22.   Haag AF, Kerscher B, Dall'Angelo S, Sani M, Longhi R, Baloban M, et al. Role of cysteine
residues and disulfide bonds in the activity of a legume root nodule-specific, cysteine-rich peptide.
J Biol Chem. 2012;287(14):10791-8.
23.   Isozumi N, Masubuchi Y, Imamura T, Mori M, Koga H, Ohki S. Structure and antimicrobial
activity of NCR169, a nodule-specific cysteine-rich peptide of Medicago truncatula. Sci Rep.
2021;11(1):9923.
24.   Montiel J, Szucs A, Boboescu IZ, Gherman VD, Kondorosi E, Kereszt A. Terminal Bacteroid
Differentiation Is Associated With Variable Morphological Changes in Legume Species Belonging to
the Inverted Repeat-Lacking Clade. Mol Plant Microbe Interact. 2016;29(3):210-9.
25.   Yee A, Chang X, Pineda-Lucena A, Wu B, Semesi A, Le B, et al. An NMR approach to
structural proteomics. Proc Natl Acad Sci U S A. 2002;99(4):1825-30.
26.   Shaw WJ, Tarasevich BJ, Buchko GW, Arachchige RMJ, Burton SD. Controls of nature:
Secondary, tertiary, and quaternary structure of the enamel protein amelogenin in solution and on
hydroxyapatite. J Struct Biol. 2020;212(3):107630.
27.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate
protein structure prediction with AlphaFold. Nature. 2021;596(7873):583-9.
28.   Poon I, Baxter AA, Lay FT, Mills GD, Adda CG, Payne JA, et al. Phosphoinositide-mediated
oligomerization of a defensin induces cell lysis. eLife. 2014;3:e01808.
29.   Hellman LM, Fried MG. Electrophoretic mobility shift assay (EMSA) for detecting protein-
nucleic acid interactions. Nature protocols. 2007;2(8):1849-61.
30.   Olliver CL, Grobler-Rabie A, Boyd CD. In vitro translation of messenger RNA in a wheat germ
extract cell-free system. Methods Mol Biol. 1985;2:137-44.
31.   Li H, Velivelli SL, Shah DM. Antifungal potency and modes of action of a novel olive tree
defensin against closely related ascomycete fungal pathogens. . Mol Plant Microbe Interact.
2019;32:164901664.
32.   Kovaleva V, Bukhteeva I, Kit OY, Nesmelova IV. Plant Defensins from a Structural
Perspective. Int J Mol Sci. 2020;21(15).
33.   Lima RM, Kylarova S, Mergaert P, Kondorosi E. Unexplored Arsenals of Legume Peptides
With Potential for Their Applications in Medicine and Agriculture. Front Microbiol. 2020;11:1307.
34.   Wedemeyer WJ, Welker E, Narayan M, Scheraga HA. Disulfide bonds and protein folding.
Biochemistry. 2000;39(23):7032.
35.   Bhardwaj G, Mulligan VK, Bahl CD, Gilmore JM, Harvey PJ, Cheneval O, et al. Accurate de
novo design of hyperstable constrained peptides. Nature. 2016;538(7625):329-35.
36.   Buchko GW, Pulavarti S, Ovchinnikov V, Shaw EA, Rettie SA, Myler PJ, et al. Cytosolic
expression, solution structures, and molecular dynamics simulation of genetically encodable
disulfide-rich de novo designed peptides. Protein Sci. 2018;27(9):1611-23.

37.   Karbalaei M, Rezaee SA, Farsiani H. Pichia pastoris: A highly successful expression system
for optimal synthesis of heterologous proteins. J Cell Physiol. 2020;235(9):5867-81.
38.   Farkas A, Maroti G, Durgo H, Gyorgypal Z, Lima RM, Medzihradszky KF, et al. Medicago
truncatula symbiotic peptide NCR247 contributes to bacteroid differentiation through multiple
mechanisms. Proc Natl Acad Sci U S A. 2014;111(14):5183-8.
39.   Li H, Kalunke R, Tetorya M, Czymmek KJ, Shah DM. Modes of action and potential as a
peptide-based biofungicide of a plant defensin MtDef4. Mol Plant Pathol. 2024;25, e134538.
https://doi.org/10.1111/mpp.134538.
40.   Hawkins NJ, Fraaije BA. Fitness Penalties in the Evolution of Fungicide Resistance. Annu Rev
Phytopathol. 2018;56:339-60.
41.   Liang J, Shah DM, Wu Y, Rosenberger CA, Hakimi SM, inventors; Monsanto Company,
assignee. Antifungal polypeptide from alfalfa and methods for controlling plant pathogenic fungi.
USA patent US 6316407. 2001 November 13, 2001.
42.   Lee W, Rahimi M, Lee Y, Chiu A. POKY: a software suite for multidimensional NMR and 3D
structure calculation of biomolecules. Bioinformatics. 2021;37(18):3041-2.
43.   Guntert P. Automated NMR structure calculation with CYANA. Methods Mol Biol.
2004;278:353-78.
44.   Shen Y, Delaglio F, Cornilescu G, Bax A. TALOS+: a hybrid method for predicting protein
backbone torsion angles from NMR chemical shifts. J Biomol NMR. 2009;44(4):213-23.
45.   Buchko GW, Hewitt SN, Van Voorhis WC, Myler PJ. Solution NMR structures of oxidized and
reduced Ehrlichia chaffeensis thioredoxin: NMR-invisible structure owing to backbone dynamics.
Acta Crystallogr F Struct Biol Commun. 2018;74(Pt 1):46-56.
46.   Bhattacharya A, Tejero R, Montelione GT. Evaluating protein structures determined by
structural genomics consortia. Proteins. 2007;66(4):778-95.
47.   Chadha S, Kale SP. Simple fluorescence-based high throughput cell viability assay for
filamentous fungi. Lett Appl Microbiol. 2015;61(3):238-44.
48.   Djami-Tchatchou AT, Tetorya M, Godwin J, Codjoe JM, Li H, Shah DM. Small Cationic
Cysteine-Rich Defensin-Derived Antifungal Peptide Controls White Mold in Soybean. J Fungi
(Basel). 2023;9(9).
49.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.

1057

Acknowledgements

The work presented here was supported by the National Science Foundation grant IOS-2037981

and the National Institute of Food and Agriculture GRANT13515963 to D.S and K.C. The

structures for NCR13 were solved using resources at the W.R. Wiley Environmental Molecular

Sciences Laboratory, a national scientific user facility sponsored by U.S. Department of Energy's

Office of Biological and Environmental Research (BER) program located at Pacific Northwest

National Laboratory (PNNL). Battelle operates PNNL for the U.S. Department of Energy. We also

acknowledge imaging support from the Advanced Bioimaging Laboratory (RRID:SCR_018951) at

the Donald Danforth Plant Science Center (DDPSC) and usage of the Leica SPX-8 acquired

through an NSF Major Research Instrumentation grant (DBI-1337680). We acknowledge the

DDPSC Proteomics & Mass Spectrometry Facility and supported by the National Science

Foundation under Grant No. DBI-1827534 for acquisition of the Orbitrap Fusion Lumos LC-

MS/MS. The greenhouse and growth chamber space provided by the DDPSC Plant Growth

Facility is acknowledged. We also thank Dr. Ruby Tiwari for providing her comments on this

manuscript.

1073

1074 **Author Contributions**

1075 K.C. and D.S. conceived the research. J.G., A.D-C., S.V., M.T., K.C., M.Z., G.B. and D.S. designed

1076 the research. J.G., A.D-C., S.V., M.T., A.P., R.K, M.Z. and G.B. conducted the experiments. J.G.,

1077 A.D-C., S.V., M.T., K.C., M.Z., G.B. and D.S. analyzed and interpreted the results. J.G., G.B.,

1078 K.C. and D.S. wrote the manuscript.

1079

1080 **Competing interests**. The authors declare no competing interests.

1081

1082 **Data deposition**: The coordinates of the NCR13 NMR structure have been deposited in the Protein

1083 Data Bank, www.rcsb.org 8ULM (NCR13_ PFV1) and 7TH8 (NCR13_ PFV2) and the NMR

1084 chemical shifts have been deposited in the BioMagResBank, www.bmrb.wisc.edu BMRB-31111

1085 (oxidized NCR13_PFV1) and BMRB-30979 (oxidized NCR13_PFV2).