**Title: Chickpea NCR13 disulfide cross-linking variants exhibit profound differences in**

**antifungal activity and modes of action**

**Authors:** James Godwin[a,1]**,** Arnaud Thierry Djami-Tchatchou[a,1]**,** Siva L. S. Velivelli[a], Meenakshi

Tetorya[a], Raviraj Kalunke[a], Ambika Pokhrel[a], Mowei Zhou[c,2], Garry W. Buchko[c,d], Kirk J.

Czymmek[a,b], and Dilip M. Shah[a,3]

**Author Affiliations:**

[a]Donald Danforth Plant Science Center, St Louis, MO. 63132, USA

[b]Advanced Bioimaging Laboratory, Donald Danforth Plant Science Center, St Louis, MO. 63132,

USA

[c]Earth and Biological Sciences Directorate, Pacific Northwest National Laboratory, Richland, WA

99354, USA

[d]School of Molecular Biosciences, Washington State University, Pullman, WA 99164, USA

1. These authors contributed equally to the manuscript.

2. Current address: Department of Chemistry, Zhejiang University, 866 Yuhangtang Road,

Hangzhou, Zhejiang 310058, China

**3.    Corresponding Author:** (dshah@danforthcenter.org)

**Abstract**

Cationic nodule-specific cysteine-rich (NCR) peptides of nitrogen-fixing legume plants exhibit antifungal activity and can be repurposed for development as biofungicides. Chickpea NCR13 is a highly cationic peptide with six cysteines forming three disulfide bonds. Expression of NCR13 in *Pichia pastoris* resulted in formation of two peptide folding variants, NCR13_PFV1 and NCR13_PFV2, that differed in the pairing of two out of three disulfide bonds despite having an identical amino acid sequence. The NMR structure of each PFV revealed a unique three-dimensional fold with the PFV1 structure being more compact but less dynamic. PFV1 and PFV2 differed profoundly in the potency of antifungal activity and their multi-faceted modes of action (MoA). PFV1 showed significantly faster fungal cell-permeabilizing and cell entry capabilities and greater stability once inside the fungal cells. PFV1 was more effective in binding to fungal ribosomal RNA and inhibiting protein translation *in vitro*. When sprayed on pepper and tomato plants, PFV1 was more effective in controlling the gray mold disease caused by *Botrytis cinerea*. Our work highlights the significant impact of disulfide pairing on the antifungal activity and MoA of NCR13 and provides structural framework for design of novel, potent antifungal peptides for agricultural use.

**Introduction**

The management of fungal diseases using chemical fungicides is one of the mainstays of modern agriculture. Unfortunately, modern fungicides that have a single site mode of action (MoA) are losing their effectiveness owing to an increase in development of fungal resistance (McLaughlin et al., 2023). A major limitation in fungicide development is finding molecules with novel structures and multi-site MoA to mitigate evolution of resistance.

Plants express many antimicrobial peptides that are vital components of the innate immune system and act as a first line of defense against bacterial and fungal pathogens (Dos Santos-Silva et al., 2020; Goyal & Mattoo, 2014). Moreover, plants express a multitude of cationic antifungal peptides differing in structure and sequence (van der Weerden et al., 2013). They are versatile, have large sequence space, and can be tuned for broad-spectrum antifungal activity against plant fungal pathogens. Thus, cationic antimicrobial peptides represent promising templates for a new generation of fungicides with broad action against fungal pathogens (Lobo & Boto, 2022).

Several antifungal peptides expressed in plants are cysteine-rich with a characteristic number of cysteine-residues at conserved positions and exhibit high degree of sequence diversity (Tavormina et al., 2015; van der Weerden et al., 2013). Defensins and defensin-like peptides are the best characterized examples of cysteine-rich peptides employed by plants to combat fungal pathogens (Cools et al., 2017; Maroti et al., 2015; Parisi et al., 2019). Legumes of the inverted repeat-lacking clade (IRLC) express many nodule-specific cysteine-rich (NCR) defensin-like peptides that cause differentiation of nitrogen-fixing bacteria to bacteroids. In *Medicago truncatula* and chickpea, terminal differentiation process is orchestrated by 639 and 63 nodule-specific cysteine-rich (NCR) peptides, respectively, with conserved cysteine motifs (Montiel et al., 2017). NCRs cause bacterial genome endoreduplication, changes in transcription and translation,

and induction of morphological changes leading to the formation of bacteroids with unique

morphology (Kereszt et al., 2018; Kondorosi et al., 2013; Maroti et al., 2015; Mergaert, 2018).

Each NCR is synthesized as a precursor protein containing an N-terminal signal peptide and a C-

terminal mature peptide. Several NCRs in each legume are cationic with a net charge ranging from

+3 to +11 and rich in hydrophobic residues, dual hallmarks of defensin-like antimicrobial peptides

(Montiel et al., 2017; Ordogh et al., 2014). Large landscape of natural NCRs from IRLC legumes

offers an unprecedented opportunity for discovery of the novel antimicrobial functions in plants.

Several NCR peptides exhibiting antibacterial activity against Gram-negative and Gram-positive

bacteria have so far been identified (Alhhazmi et al., 2024; Balogh et al., 2014; Farkas et al., 2017;

Tiricz et al., 2013). Another report detailed the evaluation of 19 cationic *M. truncatula* NCRs for

their ability to inhibit the growth of a human fungal pathogen *Candida albicans*. Of these, nine

with a net charge above +9 inhibited the growth of this pathogen at concentrations ranging from

1.5 to 10.5 μM (Ordogh et al., 2014). Previously, we reported *M. truncatula* NCR044 having

broad-spectrum antifungal activity and multi-faceted MoA. In addition, when sprayed on leaf

surfaces of tomato and *Nicotiana benthamiana* plants, NCR044 conferred resistance to the gray

mold disease caused by the fungal pathogen *B. cinerea* (Velivelli et al., 2020).

NCRs exhibit diverse primary sequences and net charges, with a notable conserved feature

being the presence of four or six conserved cysteine residues. These cysteines form intramolecular

disulfide bonds which enhance the stability of the peptides (Bulleid & Ellgaard, 2011; Montiel et

al., 2017). The influence of cysteines and disulfide cross-linking on the structure and function of

*M. truncatula* NCR247 and NCR169, each containing four cysteines, has been previously

investigated. Four conserved cysteines and disulfide cross-linking had major influence on the

symbiotic activities of NCR247 tested against *Sinorhizobium meliloti* (Haag et al., 2012; Shabab

et al., 2016). NCR169 produced two structurally distinct disulfide variants that exhibited differences in phospholipid binding and antibacterial activity (Isozumi et al., 2021). However, the significance of disulfide cross-linking for the structures, antifungal activity, and MoA of many NCRs remains poorly understood.

In this study, we show that chickpea NCR13, a highly cationic 32 residue peptide, was expressed in *Pichia pastoris* as two structurally unique disulfide variants with an identical amino acid sequence. Interestingly, these two variants exhibited distinct differences in their antifungal potency against economically important plant fungal pathogens. NCR13 disulfide variants displayed notable differences in fungal cell entry, cell permeabilization, stability and MoA against the fungal pathogen *B. cinerea*. While both disulfide variants reduced gray mold disease symptoms by spray application on the leaves of pepper and tomato plants, the NCR13 variant with highly potent *in vitro* antifungal activity was more effective in controlling the disease *in planta*. Our study reveals the remarkable impact of disulfide cross-linking on the structure-activity relationships and MoA of an antifungal defensin-like NCR peptide.

**Results**

**Chickpea NCR13 is expressed in *P. pastoris* as two peptide folding variants differing in disulfide pairing**

In chickpea, NCR13 is one of 63 predicted NCR peptides expressed only in nodules undergoing differentiation during symbiosis (Montiel et al., 2017; Montiel et al., 2016). NCR13 is a 32-residue peptide with six cysteines predicted to form three disulfide bonds (Figure 1A). Two other homologs of this peptide, NCR07 and NCR15, are also expressed in chickpea nodules with highly conserved cysteine residues (Figure 1A). No other IRLC legumes express NCRs with

significant homology to NCR13. NCR13 has 66.6% and 48.4% sequence identity with NCR15

and NCR07, respectively, carrying a net charge of +8 and a hydrophobic amino acid content of

28%, hallmark features of a potential antifungal peptide (Figure 1B).

To investigate the antifungal properties of the NCR13 peptide, it was expressed in a

heterologous *Pichia pastoris* system. The peptide was secreted into the growth medium and

purified using the cation exchange chromatography.  When the NCR13 peptide was subjected to

C18 reverse-phase high-performance liquid chromatography (HPLC), we consistently observed

two peaks (Figure 1C). In contrast, chemically synthesized NCR13 produced only one peak

(Figure 1D). These observations led us to suspect that oxidized NCR13 was expressed in our *P.*

*pastoris* expression system as two species differing in length or disulfide cross-linking pattern.

Mass spectrometry analysis of peptides purified from both HPLC peaks revealed identical

molecular weights (Supplementary Figure S1) as expected for NCR13 with three disulfide bonds,

narrowing our suspicion to different disulfide bonding patterns. Across various batches of purified

NCR13, peak 1 was consistently observed to contain less peptide than peak 2 (Supplementary

Table S1). Peak 1 and peak 2 peptides are hereafter referred to as NCR13_peptide folding variant

1 (NCR13_PFV1) and NCR13_PFV2 respectively.

To further characterize both NCR13_PFVs, we performed NMR structural analysis of $^{15}$N-

labeled NCR13_PFV1 and NCR13_PFV2.  Figure 2A and B show the assigned $^{1}$H-$^{15}$N HSQC

spectra for NCR13_PFV1 and NCR13_PFV2, respectively. The broad dispersion of amide

chemical shifts in both spectral dimensions is a standard feature of a structured peptide (Yee et al.,

2002). As illustrated in Figure 2, assignment of both spectra was consistent with the primary amino

acid sequence of NCR13. However, the pattern of the fingerprint $^{1}$H-$^{15}$N HSQC spectrum for both

peptides was clearly different, suggesting their structures were unique. Given that the primary

amino acid sequence of NCR13 contains six cysteine residues (C4, C10, C15, C23, C28, and C30)

that can form fifteen different patterns of intramolecular disulfide bonds when all six are oxidized,

our attention immediately focused on a different disulfide bond pattern for the two peptides.

Treatment of both peptides with the reducing agents dithiothreitol (DTT) or tris(2-carboxyethyl)

phosphine resulted in the identical $^1$H-$^{15}$N HSQC spectrum shown in Figure 2C, illustrating that

the parent primary amino sequence was identical for both peptides. Note that when the NCR13

cysteines are all in the reduced (dithiol) state, dispersion of amide chemical shifts in both spectral

dimensions were distinctively narrow, a standard feature of an unstructured protein (Shaw et al.,

2020; Yee et al., 2002). As observed for *M. truncatula* NCR044 (Velivelli et al., 2020), disulfide

bond formation is essential to stabilize the elements of secondary structure observed in the

oxidized state because in the reduced state the peptide is largely disordered.

Disulfide bond formation between all six cysteine residues in NCR13_PFV1 and

NCR13_PFV2 was unambiguously verified from the NMR chemical shifts of the cysteine side

chain β-carbon. In general, the cysteine $^{13}$C$^β$ chemical shift is greater than 35 ppm in the oxidized

state and under 32 ppm in the reduced state. As summarized in Supplementary Table S2, the $^{13}$C$^β$

chemical shift for all six cysteines in both isoforms of oxidized NCR13 are between 37 and 48

ppm, indicating they are all oxidized in NCR13_PFV1 and NCR13_PFV2. Due to the number of

potential disulfide bond patterns that the six cysteine residues can adopt, it was not possible to use

NMR structures in early calculations to unambiguously deduce disulfide bond pairs from $^1$H-$^1$H

NOEs that arise due to proximal γ-sulphur atoms in the disulfide state. Instead, as previously

applied to NCR044, the disulfide bond pattern was inferred from the analysis of mass spectral data

for trypsin-digested, unlabeled protein in the oxidized and reduced states (Velivelli et al., 2020).

For NCR13_PFV2, this approach showed that the disulfide bonds were C4-C10, C15-C30, and

C23-C28 (Supplementary Figure S2). For NCR13_PFV1, despite extensive modification of the

experimental procedures, this approach failed due to the hyperstability of this isoform towards

trypsinization. Consequently, the disulfide bond pattern predicted by AlphaFold (Jumper et al.,

2021) C4-C23, C10-C28, and C15-C30 was used to calculate the ensemble of NMR structures for

NCR13_PFV1. The two NCR13 isoforms share one identical disulfide bond, C15-C30, and

Supplementary Table S2 shows that these cysteine $^{13}C^{\beta}$ chemical shifts differ less than 0.5 ppm,

strongly suggesting this disulfide bond is present in both isoforms. As will be discussed below, this

disulfide bond pattern best supports the experimental NOE data.

**Three-dimensional NMR solution structures of NCR13_PFV1 and NCR13_PFV2**

To obtain structural insights into NCR13_PFVs, we performed NMR analyses of the oxidized

forms of these two peptides. Figure 3 illustrates a superposition of cartoon representations of the

final ensemble of structures calculated for the two NCR13 PFVs (left) with the main chain of a

single structure shown to the right of the ensemble with the disulfide bonds highlighted in different

colors. The elements of secondary structure are essentially identical in both isoforms, one anti-

parallel β-sheet (V20-I24, F27-V31) and a short a-helix (Q5 or S6–C10). However, the disulfide

bond pattern markedly alters the relative orientation of these elements of secondary structure. Both

peptides contain one identical disulfide bond, C15-C30, that ties the N-terminal end of the peptide

to approximately the middle of the linker region between the two elements of secondary structure.

The consequence is that in NCR13_PFV1 the α-helix sits over the top of the face of the β-sheet

while in NCR13_PFV2 the α-helix sits on the side of the β-sheet. This difference is better

illustrated in Supplementary Figure S3, a cartoon illustration of two representative structures

superimposed on the β-strand. This results in a smaller accessible surface area for NCR13_PFV1

(3188 $\pm$ 63 Å$^2$) than for NCR13_PFV2 (3357 $\pm$ 129 Å$^2$). The more compact structure for

NCR13_PFV1, with the disulfide bonds formed between C4, C10, C23, and C28 directed into the

interior of the peptide, may protect these disulfide bonds from reduction and contribute to the

hyperstability of NCR13_PFV1 towards reduction. Disulfide bonds may also be protected from

reduction because this NCR13 isomer is less dynamic. This suggestion is indirectly made by

comparing the ensemble of calculated solution structures illustrated in Figure 3. The ensemble of

structures calculated for NCR13_PFV1 converge to a greater extent than for NCR13_PFV2 with

the backbone (N-C$^a$-C=O) RMSD to the mean for the ordered residues almost twice as large for

NCR13_PFV2 (0.41 $\pm$ 13 Å versus 0.76 $\pm$ 20 Å, respectively; Supplementary Table S3). While

this difference reflects the greater number of NOEs observed for NCR13_PFV1 (316) than for

NCR13_PFV2 (119) (Supplementary Table S3) due to a more compact structure, it also indirectly

suggests the NCR13_PFV1 explores less conformational space than NCR13_PFV2.

To further verify that the AlphaFold predicted disulfide bond pattern was correct for

NCR13_PFV1, an NMR structure was calculated using only the NOE and TALOS+ (dihedral

angle) data (no hydrogen bonds) with the other possible disulfide bond pattern (C4-C28, C10-C23,

and C15-C30). In Supplementary Figure S4A, two representative cartoon structures that result

from the two possible disulfide bond patterns are superimposed on the β-strands. In the reversed,

C4-C28/C10-C23 disulfide pattern (yellow), the direction of α1 is also reversed relative to the β-

strand and this prevents the N-terminal region (T1-P3) from making close contact with the β-

strand. While most of the NOEs were satisfied in both structure calculations, long range

experimental NOEs between the N-terminal region of NCR13_PFV1 (T1-P3) and the β-strand

were only satisfied with the C4-C23/C10-C28 disulfide pattern used in the structure deposited to

the PDB (8UTM), suggesting this pattern was correct. Furthermore, as shown in Supplementary

Figure S4B, the predicted AlphaFold structure for NCR13_PFV1 was like the experimental

structure (8UTM). In both structures, the β-sheet is almost identical with the major difference a

slightly shorter α-helix displaced two residues (D7-K11 versus Q5-C10 experimentally) that packs

against the β-strand a bit differently in the AlphaFold structure (green). Most relatively, the N-

terminal region (T1-P3) folds towards the β-strand in the AlphaFold predicted structure, as verified

experimentally with the long-range NOEs between these regions.

**219 NCR13_PFV1 exhibits much greater antifungal activity against fungal pathogens than**
**220 NCR13_PFV2**

Having found striking differences in their NMR structures, we investigated the *in vitro*

antifungal activity of NCR13_PFV1 and NCR13_PFV2 against three necrotrophic plant fungal

pathogens (Velivelli et al., 2020). NCR13_PFV1 inhibited the growth of *B. cinerea* at the Minimal

Inhibitory Concentration (MIC) of 0.09 µM. Conversely, NCR13_PFV2 inhibited *B. cinerea* at the

MIC of 1.5 µM (Figure 4A-C), seventeen-fold higher than that of NCR13_PFV1. Chemically

synthesized NCR13 had a MIC of 3 µM, two-fold higher than NCR13_PFV2 (Figure 4B),

suggesting that this fully reduced synthesized product was yet another isoform of NCR13 and

illustrating the importance of disulfide bonds for antifungal activity. Similarly, NCR13_PFV1

showed several fold greater antifungal activity against *Fusarium virguliforme,* and *Sclerotinia*

*sclerotiorum* compared to NCR13_PFV2 (Figure 4A). Overall, these results reveal that

NCR13_PFV1 is a much more potent antifungal peptide compared to NCR13_PFV2.

**233 Disulfide bonds in NCR13_PFV1 contribute to the antifungal activity and formation of**
**234 peptide folding variants**

Fully oxidized NCR13_PFV1 contains three disulfide bonds:  C4-C23, C10-C28, and C15-

C30. The potent antifungal activity of NCR13_PFV1 against *B. cinerea* at nanomolar

concentrations prompted us to investigate the contribution of each disulfide bond towards

antifungal activity and the formation of folded variants when expressed in *P. pastoris*. We carried

out functional analysis of these disulfide bonds by creating three dual Cys to Ser variants to

systematically remove one disulfide bond:  NCR13_C4S-C23S, NCR13_C10S-C28S, and

NCR13_C15S-C30S variants (Figure 5A). These knockout variants were expressed in *P. pastoris*

and purified. HPLC profiles of NCR13_C4S-C23S, and NCR13_C15S-C30S revealed a single

peak whereas NCR13_C10S-C28S revealed two peaks (Figure 5B), the latter observation

suggesting NCR13_C10S-C28S folded with at least two different disulfide bonding patterns. Mass

spectrometry analysis confirmed the expected molecular mass of each peptide (Supplementary

Table S4), consistent with them being in an oxidized state. Therefore, our results indicate that the

disulfide pairing of C4-C23 and C15-C30 is essential for the formation of NCR13_PFVs.

All three single disulfide bond knockouts had lower antifungal activity against *B. cinerea*

compared to NCR13_PFV1, as summarized in Figure 5.  NCR13_C4S-C23S and NCR13_C15S-

C30S both inhibited *B. cinerea* with a MIC value of 1.5 and 0.75 µM respectively, while the

products in the two NCR13_C10S-C28S HPLC peaks had a MIC of 3 µM (Figure 5C and D).

While we were unable to confirm the disulfide bonding pattern in each of these knockout variants,

the MIC data indicates that each disulfide bond contributes to the antifungal activity of

NCR13_PFV1 (Figure 5C and D).  Indeed, a NCR13 construct with all six Cys residues replaced

with Ser (NSR13) was inactive against *B. cinerea* even at the highest tested concentration, 24 µM

(Figure 5C and D; Supplementary Figure S4).  It is likely that NSR13 is unfolded, as suggested by

the NMR data for reduced NCR13 (Figure 2) and its late elution time on a reverse phase column.

Therefore, it can be concluded that each disulfide bond contributes to the antifungal activity of the

NCR13.

**Identification of the sequence motif governing the antifungal activity of NCR13**

To identify the sequence motif governing the antifungal activity of NCR13 against *B.*

*cinerea*, we purchased chemically synthesized alanine scanning variants NCR13_AlaV1 through

NCR13_Ala5 (Figure 6A). These variants contained a string of three to five consecutive Ala

residues substituted throughout the NCR13 sequence. We then compared their antifungal activity

relative to a chemically synthesized NCR13. Notably, our findings revealed that NCR13_AlaV2,

NCR13_AlaV3, and NCR13_AlaV4 had no antifungal activity against *B. cinerea* at the highest

tested concentration, 12 µM (Figure 6A, Supplementary Figure S6A), whereas NCR13 inhibited

this pathogen at an MIC of 3 µM. This data suggested that a 13-residue, Lys-rich motif,

KKFA**C**RKPKVPK**C,** located in the random coil and β-strand region, was a major determinant of

NCR13 antifungal activity. We further explored the importance of this motif by testing antifungal

activity of NCR13_AlaV3 expressed in *P. pastoris* to allow formation of disulfide bonds and found

that it also lacked antifungal activity against *B. cinerea* like the chemically synthesized

NCR13_AlaV3 (Supplementary Figure S6B). It is noteworthy that NCR13_AlaV1 and

NCR13_AlaV5 also displayed 2- and 4-fold reduction, respectively, in the antifungal activity

against *B. cinerea* indicating that the N-terminal QSDKD and C-terminal INGF motifs are

relatively minor contributors to the antifungal activity of this peptide. (Figure 6A).

**NCR13_PFV1 disrupts the plasma membrane of *B. cinerea* faster than NCR13_PFV2**

Since NCR13_PFV1 exhibits ~17-fold more potent antifungal activity against *B. cinerea*

than NCR13_PFV2, we hypothesized that the difference in antifungal activity lies in differential

membrane permeabilization ability. To test this hypothesis, we performed SYTOX$^{TM}$ Green (SG)

assays to examine the ability of NCR13_PFV1 and NCR13_PFV2 at 0.09 μM to permeabilize the

plasma membrane of *B. cinerea* germlings. SG, a membrane-impermeant dye, binds to nucleic

acids and is an indicator of membrane permeability. Average fluorescence intensities were

measured along different time points and NCR13_PFV1 showed much higher fluorescence

intensity within exposed cells than NCR13_PFV2 (Figure 7A and B). After 10 min, SG uptake

was observed in germlings challenged with NCR13_PFV1 but not in germlings challenged with

NCR13_PFV2. After 20 min, a very strong SG uptake signal was observed in NCR13_PFV1

treated germlings, but only a modest signal in NCR13_PFV2 treated germlings (Figure 7A and B).

Thus, NCR13_PFV1 inhibited fungal growth to a greater degree than NCR13_PFV2 at nanomolar

concentration in part by permeabilizing the plasma membrane of *B. cinerea*.

**NCR13_PFV1 and NCR13_PFV2 reveal quantitative differences in their binding to multiple**

**membrane phospholipids**

Antifungal plant defensins gain entry into fungal cells and bind to a variety of bioactive

phospholipids resident on the plasma membrane (Parisi et al., 2019). We speculated that

NCR13_PFV1 and NCR13_PFV2  differ in their phospholipid binding specificity in the context

of membrane bilayer. Therefore, we used  phospholipid strip assays to determine if NCR13_PFVs

bind to different phospholipids. Our findings revealed that both NCR13_PFV1 and NCR13_PFV2

interacted with several phospholipids, including phosphatidylinositol 4-phosphate [PI(4)P] and

PI(4,5)P$_2$ (Figure 7C). NCR13_PFV2 displayed significantly stronger binding affinity to

phosphoinositides such as PI(3)P, PI(4)P, PI(5)P, PI(4,5)P$_2$, and PI(3,4,5)P$_3$ than NCR13_PFV1

(Figure 7D). The interaction of PI(4)P and PI(4,5)P$_2$ with NCR13_PFV1 and NCR13_PFV2 were

further validated using a polyPIPosome binding assay (Figure 7E). Nevertheless, the significance

of the phospholipid binding differences between the two peptides for their antifungal activity

requires further investigation. Since several antifungal plant defensins bind to membrane

phospholipids to oligomerize and induce membrane permeabilization (Poon et al., 2014), we tested

the ability of NCR13_PFV1 and NCR13_PFV2 to oligomerize by incubating the peptides with the

biochemical cross-linker bis(sulfosuccinimidyl)suberate (BS$_3$) in the presence and absence of

PI(4,5)P$_2$. NCR13_PFV1 formed oligomers both in the presence and absence of PI(4,5)P2. On the

other hand, NCR13_PFV2 formed higher order oligomers, but, only at higher concentrations (115

μM and 330 μM) of PI(4,5)P$_2$ (Supplementary Figure S7). Therefore, further investigation is

required to determine whether phospholipid binding, including PI(4,5)P2, and the oligomerization

of NCR13_PFVs are critical for fungal cell killing.

**NCR13 PFV1 is internalized faster in germling cells of *B. cinerea* than NCR13_PFV2**

Due to its much greater antifungal activity *in vitro* against *B. cinerea*, we hypothesized that

NCR13_PFV1 exhibited faster entry into fungal cells at its MIC than NCR13_PFV2. To test this

hypothesis, we exposed *B. cinerea* germlings to 0.09 μM DyLight550-labeled NCR13_PFV1 and

NCR13_PFV2 and monitored their entry into germlings of *B. cinerea* using confocal microscopy.

Within ~30 min, NCR13_PFV1 penetrated the germlings; however, NCR13_PFV2 did not enter

the germlings (Figure 8A and B). At a concentration of 1.5 μM, NCR13_PFV1 started entering

germlings in ~4 min, while NCR13_PFV2, after briefly gathering on the cell surface (Figure 8C,

white arrowheads), started entering germlings at ~8 min. Unlike NCR13_PFV2's lack of entry at

0.09 µM, delayed entry of NCR13_PFV2 was evident at 1.5 µM (Figure 8C and D).

Following entry into *B. cinerea* germling cells, both NCR13_PFV1 and NCR13_PFV2 accumulated in distinct subcellular foci (Figure 8E, white arrows). Further localization studies using the nuclear staining dye Hoechst 33258 revealed that both peptides were localized in the nuclei. To determine if each peptide was sub-localized within the nucleolus, we used the nucleolus-specific RNA staining dye Nucleolus Bright Green. As shown in Figure 8F, each peptide was localized within the nucleolus of germling cells suggesting that each peptide likely binds to ribosomal RNA (rRNA) or proteins and interferes with protein synthesis.

To gain further insight into the MoA of NCR13_PFV1 and NCR13_PFV2, we assessed the stability of each peptide once inside *B. cinerea* cells. We challenged *B. cinerea* germlings with each peptide separately for two different time points, 2 h and 4 h, and tested the stability of each peptide in fungal cells through western blot analysis. An NCR13 polyclonal antibody capable of binding to each peptide with equal affinity was used for this analysis (Supplementary Figure S8). NCR13_PFV1 accumulated in germling cells within 2 h and accumulates further after 4 h. In contrast, much less NCR13_PFV2 accumulated in germling cells within 2 h and was strongly decreased by 4 h (Supplementary Figure 8E). Therefore, we highlight NCR13_PFV1 more rapidly enters fungal cells than NCR13_PFV2, and once inside NCR13_PFV1 is more stable than NCR13_PFV2.

**NCR13_PFV1 binds to rRNA and is a more potent inhibitor of protein translation *in vitro* than NCR13_PFV2**

Since NCR13_PFVs are localized in the nucleoli of *B. cinerea*, we decided to test the binding of each peptide to rRNA. This was accomplished with an Electrophoretic Mobility Shift Assay (EMSA) (Hellman & Fried, 2007) by incubating each peptide with total rRNA isolated from *B. cinerea* germlings. Our results showed that NCR13_PFV1 retarded the mobility of the rRNA at all concentrations greater than 0.18 µM, while NCR13_PFV2 showed no binding to rRNA until 3 µM (Figure 9A; Supplementary Figure S8). Therefore, it is evident that NCR13_PFV1 exhibits stronger binding affinity to rRNA than NCR13_PFV2.

Next, we tested the ability of the NCR13_PFVs to inhibit protein translation *in vitro* because we hypothesized that NCR13_PFV1 may be a more potent inhibitor of protein translation than NCR13_PFV2. This was accomplished using a wheat germ *in vitro* translation system based on a luciferase activity (Olliver et al., 1985). Protein translation inhibitory activity of each peptide was concentration dependent (Figure 9B). NCR13_PFV1 inhibited translation more effectively than NCR13_PFV2 at concentrations more than 0.75 µM. At 24 µM, NCR13_PFV1 almost completely halted translation compared to the ~50% inhibition by NCR13_PFV2 (Figure 9B). Based on these results, the difference in the antifungal potency of NCR13_PFVs may be partly due to the inhibition of protein translation *in vivo* with NCR13_PFV1 being a more potent protein translation inhibitor in *B. cinerea*.

**NCR13_PFV1 is more potent than NCR13_PFV2 semi-*in planta***

Next, we compared the semi-*in planta* antifungal activity of NCR13_PFV1 and NCR13_PFV2 using detached leaves of pepper as previously described (Li et al., 2019). To eliminate leaf-to-leaf variation, both peptides were added instantaneously with freshly prepared *B. cinerea* spores to the same leaf, along with a no-peptide control, at different concentrations (3, 1.5,

and 0.09 μM) (Supplementary Figure S9A). Leaves were evaluated for reduction of gray mold

lesions through measurement of photosynthetic efficiency Fv/Fm (variable fluorescence over

saturation level of fluorescence) using CropReporter images. At 3 μM and 1.5 μM NCR13_PFVs,

no disease symptoms were visualized (Supplementary Figure S9B) and no significant difference

in photosynthetic efficiency (Supplementary Figure S9C) was observed, whereas the no-peptide

control contained visible lesions and reduced photosynthetic efficiency. At 0.09 μM, only

NCR13_PFV1 effectively suppressed the formation of visible lesions, while NCR13_PFV2

resulted in lesion sizes similar to the no-peptide control (Supplementary Figure S9B). Moreover,

a decrease in photosynthetic efficiency observed for NCR13_PFV1 was milder than observed for

NCR13_PFV2. Together, this data further illustrates that NCR13_PFV1 is a more potent antifungal

agent than NCR13_PFV2 in terms of preventing and reducing gray mold lesions on peppers

(Supplementary Figure S9B and C). These observations correlate well with the *in vitro* antifungal

activity of each peptide. Based on this comparative study of their antifungal activity using a

detached leaf assay, NCR13_PFV1 was judged to be a more promising peptide-based fungicide *in*

*planta* (see below).

**Spray-applied NCR13_PFV1 is more effective in reducing gray mold disease symptoms in**

**pepper and tomato plants than NCR13_PFV2**

To determine if the *in vitro* and semi-*in planta* antifungal activity of these peptides

translated to antifungal activity *in planta*, we tested the curative protection afforded by

NCR13_PFV1 and NCR13_PFV2 against *B. cinerea* by spray application on pepper plants. Four-

week-old pepper plants were pre-inoculated with freshly prepared *B. cinerea* spores. At 8 h post-

inoculation, peptide at 1.5 and 3 μM concentration was sprayed onto pepper leaves. At 72 h post-

inoculation. disease symptoms were evaluated by comparing photosynthetic efficiency (Fv/Fm) of

the pathogen-challenged peptide-sprayed plants with the no-peptide control. Green areas indicate

a high photosynthetic rate and healthy tissue, while yellow to red areas indicate reduced

photosynthetic rate and diseased tissue. As evidenced by their higher Fv/Fm values, leaves of

pepper plants sprayed with NCR13_PFV1 at 1.5 and 3 μM had greater reduction in gray mold

lesions than the leaves of pepper plants sprayed with NCR13_PFV2 at the same concentrations

(Figure 10A and B). Pepper plants sprayed with 1.5 μM NCR13_PFV2 showed no significant

control (p>0.05) of gray mold when compared to the no-peptide control plants suggesting

NCR13_PFV2 is unable to reduce gray mold disease symptoms at lower concentration (Figure

10A-C). Correspondingly, the data aligns with the observation that NCR13_PFV1 markedly

decreased the percentage of disease severity relative to NCR13_PFV2 at both treatment

concentrations (Figure 10C). However, NCR13_PFV2 sprayed at 3 μM was able to provide

significant curative protection compared to control plants (Figure 10A-C), but it was not as

effective as NCR13_PFV1. A similar observation was made when four-week-old tomato plants

were also pre-inoculated with *B. cinerea* spores followed by spraying 3 μM of each peptide

(Supplementary Figure S10A-C). The NCR13_PFVs significantly decreased the disease severity,

with NCR13_PFV1 demonstrating the most pronounced effect (Figure S10B). NCR13_PFV1 and

NCR13_PFV2 inhibited gray mold disease 90% and 40%, respectively (Supplementary Figure

S10C). In summary, these data indicate higher potential of NCR13_PFV1 for management of the

gray mold disease in pepper and tomato plants.

**Discussion**

NCR13 is a highly cationic peptide among the 63 predicted NCR peptides expressed in chickpea

nodules (Montiel et al., 2017). In this study, we discovered that NCR13 expressed in *P. pastoris*

forms two isomers, NCR13_PFV1 and NCR13_PFV2, which have identical amino acid sequence

and molecular mass but differ in their disulfide pairing patterns. We report a comparative analysis

of the structure, *in vitro* and semi-*in planta* antifungal activity and MoA of NCR13_PFVs. Our

results support the proposal that disulfide cross-linking provides an additional source of structural

and functional diversity in NCR peptides (Shabab et al., 2016).

Four- or six-cysteine NCRs are highly diverse in their primary sequences and are predicted to

be structurally different. Previously, two solution NMR structures of NCR044 and NCR169, each

containing two disulfide bonds, were solved (Isozumi et al., 2021; Velivelli et al., 2020). The

NCR044 structure was largely disordered, dynamic and contained a single α-helix and a β-sheet

(Velivelli et al., 2020). Structural analysis revealed that the two NCR169 isomers contained only

one short, two-stranded β-sheet, with one isomer being more extended and flexible while the other

was more compact and constrained (Isozumi et al., 2021). To date, a three-dimensional structure

for a six-cysteine NCR peptide has not been reported. Here, we report the first solution structures

of two NCR13 disulfide isomers each containing an α-helix and β-sheet. Nevertheless, unique

pairing of two out of the three disulfide bonds significantly alters how these secondary structure

elements fold upon each other rendering NCR13_PFV1 more compact and resistant to digestion

by trypsin or reduction by dithiothreitol. It is striking that *P. pastoris* generates two disulfide

variants that are structurally different and raises the question of whether these variants are also

expressed in the differentiating nodules of chickpea to perform different biological activities. In

an earlier study, three disulfide variants of NCR247 containing four cysteines were generated using

chemical synthesis and *in vitro* folding. NCR247 variants were found to have distinct patterns of

biological activities such as bacterial gene expression, inhibition of bacterial cell division, protein

translation and antibacterial activity (Haag et al., 2012; Shabab et al., 2016). These observations

suggest that NCR peptides can form disulfide variants *in vitro* or in an appropriate expression host,

leading to diverse biological activity. Whether disulfide variants of NCR13, NCR247, or NCR169

are formed *in vivo* in nodules to control symbiosis in IRLC legumes remains an open question. In

contrast to NCR peptides, plant defensins with eight cysteines expressed in *P. pastoris* are only

observed as one species with an invariant disulfide pairing pattern (Kovaleva et al., 2020).

To our knowledge, this study is the first detailed investigation of a chickpea NCR peptide and

its antifungal action. Our data demonstrates that disulfide folding variants of NCR13 produced in

*P. pastoris* exhibit profound differences in antifungal activity against multiple fungal pathogens of

economic importance. NCR13_PFV1 possesses highly potent antifungal activity against *B. cinerea*

and other pathogens, showing a four- to sixteen-fold greater activity compared to NCR13_PFV2.

Remarkably, we found that the antifungal activity of NCR13_PFV1 is greater than other NCR

peptides reported to date in literature (Lima et al., 2020). It is noteworthy that enhanced antifungal

activity of both NCR13_PFVs produced in *P. pastoris* compared to chemically synthesized NCR13

without disulfide bond, underscores the critical role of disulfide bonds for antifungal activity.

Furthermore, the formation of all three disulfide bonds is essential for highly potent antifungal

activity of NCR13_PFV1 as removal of each disulfide bond resulted in a significant loss of its

antifungal activity against *B. cinerea*. In addition, Cys residues are essential for the antifungal

activity of NCR13 as their replacement with Ser leads to complete loss of antifungal activity.

Acting as molecular anchors, disulfide bonds provide stability to the three-dimensional structure

of proteins and peptides. They set the constraints for distance and angles between connected

cysteine residues, thus favoring the stability of the protein's folded state over its unfolded form

(Wedemeyer et al., 2000). Furthermore, the insertion of cysteine residues into *de novo* designed

peptides has produced hyper-stable peptides with unique tertiary structures that form the basis for

new peptide-based drugs (Bhardwaj et al., 2016; Buchko, Pulavarti, et al., 2018). The impact on

antifungal activity of knocking out two instead of just one disulfide bond is yet to be investigated.

Further, evaluating the three-dimensional structures of these double disulfide bond knockout

variants will provide deeper insight into structure-activity relationships of this unique NCR

peptide. *P. pastoris* has been the eukaryotic organism of choice for producing cysteine-rich

peptides to ensure formation of the correct disulfide bond patterns (Karbalaei et al., 2020). It is

intriguing that NCR13 expression results in the formation of two disulfide folding variants in *P.*

*pastoris*. In this context, it is critical to observe that knocking out either the C4-C23 or C15-C30

disulfide bond in NCR13_PFV1 results in the formation of one peak on HPLC, whereas knocking

out the C10-C28 bond still results in the formation of two distinct peaks. Mass spectrometry

analysis shows that peptides from two HPLC peaks of the C10-C28 knockout and one HPLC peak

in the C4-C23 and C15-C30 knockouts are in the oxidized state. Therefore, it is likely they are in

a folded/oxidized form. Finally, understanding the molecular mechanisms governing the

generation of disulfide variants of specific antifungal NCR peptides could pave the way for

production of targeted disulfide structural variants in *P. pastoris* with enhanced antifungal activity.

Antifungal peptides with potent fungicidal activity and multiple MoA are attractive candidates

for development as spray-on fungicides. NCR13 has a promise as one such candidate. For instance,

ability to permeabilize the plasma membrane of fungal pathogens is a common MoA of plant

cysteine-rich antifungal peptides including NCRs (Ordogh et al., 2014). NCR13_PFV1 can

permeabilize the plasma membrane of *B. cinerea* rapidly at very low concentrations. The

membrane disrupting ability of NCR13_PFV1 is far greater than that of NCR13_PFV2 and

NCR044 (Velivelli et al., 2020). Both NCR_PFVs bind to the same phospholipids on lipid strips

with different binding intensity. However, it is not clear what role, if any, phospholipid binding

plays in conferring increased membrane disrupting ability of NCR13_PFV1 and this aspect

remains to be elucidated. In addition, NCR13_PFV1 gains rapid entry into the interior of fungal

cells compared to NCR13_PFV2. It is plausible that more compact folding of NCR13_PFV1

allows it to more quickly penetrate the cell wall and pass through the outer and inner leaflets of

the plasma membrane. This is supported by the fact that *B. cinerea* germlings challenged with 1.5

µM of each peptide showed NCR13_PFV1 enters fungal cells in much greater levels after 2 h than

NCR13_PFV2. Moreover, once inside the cells, NCR13_PFV1 accumulates stably for at least 4 h

whereas NCR13_PFV2 tends to degrade in this time. Greater stability of NCR13_PFV1 to

proteolysis might be a major contributing factor to its much greater antifungal activity against this

pathogen, an observation that warrants further investigation.

Subcellular localization studies showed that NCR_PFVs are targeted to nucleoli in the

germlings of *B. cinerea* suggesting that these peptides are targeting ribosomal biogenesis or protein

translation machinery as part of their MoA as proposed earlier for the MoA of NCR044 (Velivelli

et al., 2020). In support of this hypothesis, we found that addition of each NCR13_PFVs to the *in*

*vitro* translation system inhibited protein synthesis in a concentration dependent manner.

NCR13_PFV1 proved to be more effective in inhibiting protein translation *in vitro*. NCR13_PFV1

also binds to *B. cinerea* rRNA at a lower concentration (0.37 µM) than NCR13_PFV2 (6 µM).

Further research is needed to determine if differences in rRNA binding affinity and protein

translation inhibition *in vitro* translate to profound differences in their antifungal activity *in vivo*.

NCR peptides appear to target protein translation as part of their antimicrobial mechanism as

evidenced by *M. truncatula* NCR247 inhibiting protein translation *in vitro* by binding to ribosomal

proteins (Farkas et al., 2014; Shabab et al., 2016). Plant defensin MtDef4 also inhibits protein

translation *in vitro* by binding to ribosomal proteins, but not to ribosomal RNA (Li et al., 2024).

Besides that, it is likely that NCR13_PFV1 and NCR13_PFV2 have different, yet unidentified,

intracellular targets which will be the subject of future investigation.

Fungicides currently used commercially for management of fungal diseases target ergosterol

biosynthesis, succinate dehydrogenase and cytochrome $bc_1$ complex. These targets are prone to

rapid development of fungal resistance (Hawkins & Fraaije, 2018). Small NCR peptides with

multi-faceted MoA have emerged as potential candidates for development as bioinspired

fungicides. We have previously demonstrated efficacy of NCR044 as a spray-on fungicide for

management of gray mold in tomato (Velivelli et al., 2020). Here, we report NCR13 is also a multi-

target antifungal peptide that reduces gray mold disease symptoms curatively when sprayed on

pepper and tomato plants. Consistent with its *in vitro* and semi-in planta antifungal activity,

NCR13_PFV1 affords greater protection against gray mold disease than NCR13_PFV2 in both

pepper and tomato plants. NCR13 is amenable to large scale recombinant production in a bio-

fermenter using *P. pastoris* expression system. However, further characterization of NCR13 is

needed, including demonstrating its efficacy for control of gray mold and other diseases in large-

scale greenhouse and field tests, assessment of off-target activity against mammalian cells and

phytotoxicity. In summary, this study demonstrated that disulfide bond pairing pattern has a

profound impact on the *in vitro* antifungal activity, MoA, and *in planta* antifungal activity of

chickpea NCR13. As illustrated in Figure 11, the graphical abstract illustrates the key findings of

our study.  Overall, our work underscores the significance of disulfide cross-linking in enhancing

the efficacy of antifungal peptides for agricultural use.

## Materials and Methods

### Fungal Cultures and Spore Suspension

*B. cinerea* strain T4 was cultured in V8 medium for 3 weeks at 25°C. Fungal spores were collected by flooding the fungal growth plates with water and scraped using sterile spreader. The spore suspension was filtered through two layers of Miracloth, followed by centrifugation at 12,000 g for 2 minutes (min) and subsequently washed twice with sterile water. Lastly, fungal spores were resuspended in Synthetic Fungal Media (SFM) (Liang et al., 2001). The spore suspension was adjusted to the desired spore density of about $10^5$ spores/mL using a hemocytometer. *Fusarium virguliforme (F. virguliforme)* was grown and maintained regularly in potato dextrose agar (PDA) (Difco Laboratories) medium. Two weeks old *F. virguliforme* plates were used to collect spores as described above for *B. cinerea* followed by centrifugation and washing step. Spores were counted using a hemocytometer and adjusted to the final concentration of $10^5$ spores/mL in 2X SFM media. *Sclerotinia sclerotiorum* obtained from the field of pennycress in Illinois was grown on PDA medium at room temperature.

### Expression and purification of NCR13

For expression in *P. pastoris*, the published amino acid sequence of the mature NCR13 peptide (Montiel et al., 2016) was used for synthesis of the codon-optimized gene (GenScript). The NCR13 gene was expressed in *P. pastoris* X33 using either a SacI linearized pPICZα-A integration vector (Invitrogen) or at XhoI and XbaI restriction sites. NCR13 was purified using a CM Sephadex C-25 cation-exchange chromatography as described (Velivelli et al., 2020). After induction, cell-free supernatant was collected after centrifugation (10,000×g) at 4 °C for 20 min, and the pH was adjusted to 6.0. The supernatant was passed through cation-exchange resin, CM Sephadex C-25 (Sigma-Aldrich, Cat No: C25120) previously equilibrated with binding buffer (25 mM Sodium

Acetate Anhydrous, pH 6.0) using an AKTA FPLC. Peptides bound to the resin were eluted using

the elution buffer (1M NaCl, 50 mM Tris, pH 7.6). The FPLC fraction containing NCR13 was

concentrated and desalted using an Amicon Stirred Cell with a 3-kD cutoff membrane (Millipore,

Cat No: PLBC04310) and further purified by reverse phase C18-HPLC. HPLC fractions collected

separately for NCR13_PFV1 and NCR13_PFV2 were lyophilized and re-suspended in nuclease-

free water. Peptide concentrations were determined using the Pierce™ BCA Protein Assay Kit

(Thermo Scientific). NCR13 and alanine variants of NCR13 were chemically synthesized

(Biomatik) and further purified by in-house HPLC.

**Molecular weight determination**

Purified peptides were analyzed by LC-MS for accurate mass determination. Firstly, peptide

samples were cleaned using Thermo Scientific HyperSep C18 microscale SPE extraction tips and,

after dry-down, resuspended in 0.1% formic acid for LC-MS analysis. LC-MS was performed with

a Dionex RSLCnano HPLC coupled to an OrbiTrap Fusion Lumos (Thermo Scientific) mass

spectrometer using 80 min gradient (1-90% acetonitrile). Sample was resolved using a 75 μm x 15

cm PepMap C18 column (Thermo Scientific). MS spectra of protein ions of different charge-states

were acquired in positive ion mode with a resolution setting of 120,000 (at m/z 200). Mass

deconvolution was performed using ProteinProspector, Xcalibur (Thermo Scientific), and ESIprot

Online.

**Disulfide bond pattern identification for NCR13_PFV2 using mass spectrometry**

An aliquot of unlabelled, oxidzed NCR13_PFV2 was mixed with 20-fold less (in mass) trypsin,

incubated at room temperature for 1 h, and diluted in 50% acetonitrile, 0.1% formic acid to ~0.1

μM. The diluted samples were infused at 2 μL/min into a Thermo Q-Exactive Exploris 480

Orbitrap mass spectrometer with positive mode electrospray at 3.4 kV. The ion transfer tube was

at 320 °C, sheath gas at 2 psi, and auxiliary gas at 5 psi. Data-dependent acquisition was used (MS1 resolution 60k, MS2 resolution 30k) over 8 min. Fragmentation spectra were collected with higher-energy collisional dissociation (HCD) at 30% collision energy. Data were analyzed and deconvoluted in FreeStyle (Thermo Scientific). Crosslinked peptides were identified using Byonic (ProteinMetrics). Under the same conditions, NCR13_PFV1 could not be digested by trypsin (data not shown).

**$^{15}$N isotopic labeling of NCR13 for NMR structural analysis**

The $^{15}$N-labeled NCR13_PFVs were prepared according to the protocol as described previously (32) with minor modifications. NCR13-expressing *P. pastoris* X33 was grown at $28 \pm 2$°C for two days on a YPD plate containing 150 µg/mL Zeocin. A single colony was then inoculated into 5 mL of YPD broth and grown overnight at $28 \pm 2$ °C on a rotary shaker at $225 \pm 25$ rpm. After overnight growth, a 5 mL culture was transferred aseptically to 100 mL of FM23 medium (0.8% YNB without amino acids and ammonium sulfate, 2 mg/L biotin, 1.2% $(NH_4)_2SO_4$, 0.3% $K_2HPO_4$, 0.28% $KH_2PO_4$, 2 ml/L PTM1 salts) containing 3% glucose. After 24 h, an additional 2% glucose was added to the culture medium. After a growth phase of ~30 h and just prior to methanol induction, the culture medium was adjusted to 0.02% $^{15}$N-ammonium chloride ($^{15}NH_4Cl$) and 0.1% glucose. After 40 h, cells were harvested by centrifugation at 5,000 rpm at room temperature for 10 min, briefly washed with 0.2% glycerol, pelleted, and re-suspended in 100 mL FM23 medium containing 1.2% $^{15}NH_4Cl$ and grown for 4-5 days at 25 °C at $225 \pm 25$ rpm. In the induction phase, a low concentration of 0.2% methanol was initially added to the culture medium to induce protein expression and gradually increased by 0.4% from 72 to 120 h (1.4% final). The labeled NCR13 peaks were purified from the *P. pastoris* growth medium following the same protocol used for the unlabeled peptide.

**NMR spectroscopy of NCR13**

All NMR data were collected at 20˚C on [15]N-labeled samples of oxidized NCR13_PFV1 and NCR13_PFV2 (~ 1 mM, 20 mM sodium acetate, 50 mM NaCl, pH 5.3) using Varian (Inova-600) or Bruker (Advent-750) spectrometers equipped with an HCN-probe and pulse field gradients. Except for the carbonyl resonances (which could not be assigned because making [13]C-labeled samples in *P. pastoris* was too expensive), the [1]H, [13]C, and [15]N chemical shifts of the backbone and side chain resonances were assigned from the analysis of two-dimensional [1]H-[15]N HSQC, [1]H-[13]C HSQC, [1]H-[1]H TOCSY, [1]H-[1]H NOESY, and [1]H-[1]H COSY spectra and three-dimensional [15]N-edited NOESY-HSQC (mixing time = 90 ms) and [15]N-edited TOCSY spectra. To obtain additional chemical shift assignments, inter- and intra-residue proton side chain restraints for proton near the residual water resonance, most of the two-dimensional data sets were also collected by lyophilizing and resuspending the sample in 99.8% $D_2O$. In preparing the latter sample, a [1]H-[15]N HSQC spectrum was collected ~10 min after the addition of $D_2O$ to identify slowly exchanging amides that could be used as hydrogen bond restraints in the structure calculations. To confirm oxidized NCR13_PFV1 and NCR13_PFV2 arose from different disulfide bond patterns of a common precursor primary amino acid sequence, both oxidized samples were reduced by making the NMR samples 5 mM in Dithiothreitol (DTT). While NCR13_PFV2 was reduced rapidly, it took more than 5 – 7 days to fully reduce NCR13_PFV1. All NMR data were processed using Felix 2007 (MSI, Molecular Simulations) software and analyzed with the program Poky (v3.115) (Lee et al., 2021). The assigned [1]H, [13]C, and [15]N chemical shifts were referenced using indirect methods (DSS = 0 ppm) and deposited into the BioMagResBank database (www.bmrb.wisc.edu) under the accession numbers BMRB-31111 (oxidized NCR13_PFV1 8ULM) and BMRB-30979 (oxidized NCR13_PFV2 7TH8).

**NMR solution structure calculations for oxidized NCR13_PFV1 and NCR13_PFV2**

The protocols used to obtain solution structures for both peptides followed a similar approach starting with the $^1$H, and $^{15}$N chemical shift assignments and peak-picked NOESY data serving as initial experimental inputs in iterative structure calculations with the program CYANA (v 2.1) (Guntert, 2004). The $^1$H, $^{13}$C, and $^{15}$N chemical shift assignments were also the primary basis for the introduction of 24 (NCR13_PFV1) and 25 (NCR13_PFV2) dihedral Psi ($\psi$) and Phi ($\varphi$) angle restraints identified with the program TALOS+ using the online webserver (https://spin.niddk.nih.gov/bax/nmrserver/talos/) (Shen et al., 2009). Nine restraints between the side chain sulfur atoms of C4-C23, C15-C30, and C10-C28 (NCR13_PFV1) and C4-C10, C15-C30, and C23-C28 (NCR13_PFV1) (2.0-2.1 Å, 3.0-3.1 Å, and 3.0-3.1 Å for the S$\gamma$-S$\gamma$, S$\gamma$-C$\beta$, and C$\beta$-S$\gamma$ distances, respectively) (Buchko, Hewitt, et al., 2018; Velivelli et al., 2020) were introduced on the basis of AlphaFold predictions for NCR13_PFV1 and high resolution mass spectrometry analysis for NCR13_PFV2. Exhaustive attempts to identify the disulfide bond pattern for NCR13_PFV1 using various mass spectrometric methods failed likely due to its hyperstability to reduction by DTT or tris(2-carboxyethyl) phosphine. Backbone hydrogen bond restraints (1.8 – 2.0 Å and 2.7 – 3.0 Å for the NH–O and N-O distances, respectively) were added to the calculations for proximate backbone amide-carbonyl groups present in early structure calculations that were accompanied with slowly exchanging amides in the deuterium exchange experiment. The final ensemble of 20 CYANA derived structures was then refined in explicit water with CNS (version 1.1) using the PARAM19 force field and force constants of 500, 500, and 1000 kcal for the NOE, hydrogen bond, and dihedral restraints, respectively. For these water refinement calculations, 10 and 0 % were added to the upper boundary of the CYANA distance restraints for NCR13_PFV1 and NCR13_PFV2, respectively, and the lower boundary set to the van der Waals limit for both.

Structural quality was assessed using the online Protein Structure Validation Suite (PSVS, v1.5)

(Bhattacharya et al., 2007). Complete structural statistics are summarized in Supplementary Table

S3. The atomic coordinates for the final ensemble of 20 structures have been deposited in the

Research Collaboratory for Structural Bioinformatics (RSCB) under PDB codes 8ULM (NCR13_

PFV1) and 7TH8 (NCR13_ PFV2).

**Peptide uptake and stability assessment within *B. cinerea***

*B. cinerea* (~$10^5$ spores/mL) was inoculated in 10 mL of Potato Dextrose Broth (PDB) and

incubated in a shaker at 28ºC for 18 h. *B. cinerea* germlings were treated with 1.5 µM

NCR13_PFV1 and NCR13_PFV2 for 0, 2 and 4 h. Total proteins were extracted from the fungal

cells using 300 µL of HEPES buffer (100 mM HEPES, 5mM EDTA, 0.1% Triton-100, ProBlock™

Gold Yeast/Fungi Protease Inhibitor, pH 7.5) and centrifuged at 14000 rpm to remove cell debris.

The concentration of protein present in the supernatant was measured using Pierce™ BCA Protein

Assay Kit (Thermo Scientific). Total proteins (10 µg) were separated on a 4–20% SDS-PAGE gel

(Bio-Rad, TGX Stain-Free™ precast gels) and transferred to a nitrocellulose membrane. Equal

loading of proteins was checked using Ponceau S staining (Thermo Scientific). Immunoblotting

was performed using the custom synthesized anti-NCR13 polyclonal antibody (Biomatik) at a

concentration of 0.1 µg/mL followed by an incubation in the secondary anti-rabbit antibody

(Cytiva, Cat No: RPN4301) with a dilution of 1:20,000. ***In vitro* antifungal assay**

The antifungal activity of NCR13 against fungal pathogens was determined using the resazurin

cell viability assay as described previously (Chadha & Kale, 2015). Briefly, 45 µL of each peptide

dilution (0, 0.09, 0.1875, 0.375, 0.75, 1.5, 3 and 6 µM) was added to each well of the microtiter

plate containing 45 µL (~$10^5$ spores/mL) of spore suspension. Three biological replicates were

used for each treatment and the experiments were repeated three times. Resazurin at 0.1% (w/v)

was added to each well after incubation for 48 h. Images were taken at ~24 h. The Minimum

Inhibitory Concentration (MIC) for each peptide denotes the lowest concentration at which the

peptide halts the growth of fungus evident through a lack of color change from blue to colorless.

The *in vitro* antifungal activity of NCR13_PFVs against *S. sclerotiorum* was determined using a

24-well plate assay as described (Djami-Tchatchou et al., 2023). Mycelial plugs (1 mm in

diameter) cut from the growing edge of the two-day-old actively growing colony were transferred

from PDA media to a series of wells containing 250 µL of SFM.

**Internalization and permeabilization assays using confocal microscopy**

NCR13_PFV1 and PFV2 were labeled with DyLight550 amine reactive fluorescent dye following

the manufacturer's instructions (Thermo Scientific). *B. cinerea* germlings (50 µL, ~5x10$^4$

spores/mL) were treated with DyLight 550 labeled NCR13_PFV1 and NCR13_PFV2 (50 µL, 0.09

µM and 1.5 µM), and the fluorescence measured with excitation and emission wavelengths of 550

and 560-600 nm, respectively. Time-lapse confocal microscopy data with a z-stack at each time

point were acquired using a 63X water immersion objective lens (HC PL Apochromat CS2) at 30-

second intervals. For the membrane permeabilization assay, fluorescent dye SYTOX$^{TM}$ Green

(Thermo Scientific) at a concentration of 0.25 µM was used to determine the membrane

permeabilization of *B. cinerea* germlings as described previously (Li et al., 2019). Time lapse

imaging was performed to visualize the SYTOX$^{TM}$ Green entry upon treatment with 50 µL of 0.09

and 1.5 µM NCR13_PFVs. Imaging was performed on a Leica SP8 confocal microscope with

excitation at 488 nm and emission at 508–544 nm. Average fluorescence intensity measurements

were obtained using Fiji (Schindelin et al., 2012).

**Phospholipid binding assays**

Phospholipid binding of NCR13_PFVs were performed with PIP Strips (Echelon Biosciences) that

are spotted with 100 pmol of various biologically active phospholipids. Peptide binding to the

membrane strips was detected using monoclonal NCR13 antibody (0.1 µg/mL) followed by goat

anti-rabbit IgG HRP (Cytiva RPN4301) at 1:20,000 dilution. The binding signals were detected

using the SuperSignal West Pico Chemiluminescent Substrate Kit (Thermo Scientific) following

the manufacturer's protocol. Quantitative analysis of the blot intensity was performed using Fiji

(Schindelin et al., 2012).

**Oligomerization assay**

To determine the oligomerization status of the NCR13_PFVs, the peptides were prepared in PBS

buffer at 1.5 mg/mL (5 µL/reaction) and incubated with specified concentrations of PI(4,5)P2. The

cross-linking reaction was initiated by addition of 12.5 mM bis(sulfosuccinimidyl)suberate BS3

(Thermo Scientific) and then incubated for 30 min at room temperature. After incubation, tricine

sample buffer was added to each sample and heat denatured at 95 ºC for 5 min. Samples were

loaded on a 16.5% Tris-Tricine precast gel (Bio-Rad Laboratories). For SDS-PAGE, 5 µL of

Precision Plus Dual Xtra Protein Standards (2–250 kDa) were used (Bio-Rad Laboratories).

Oligomerization patterns were visualized using silver staining.

**PolyPIPosome binding assay**

The validation of the binding of NCR13_PFV1 and NCR13_PFV2 to phospholipids was further

performed using a liposome binding assay. In this assay, polyPIPosome binding was conducted in

200 µL of binding buffer (20 mM HEPES pH 7.4, 120 mM NaCl, 1 mM EGTA, 1 mM MgCl$_2$, 1

mg/mL BSA, 0.2 mM CaCl$_2$, 5 mM KCl) plus 10 µL each of control PolyPIPosome, PI4P

PolyPIPosomes, and PIP$_2$ PolyPIPosome (Echelon Biosciences). Subsequently, 2 µg of each

peptide was added, and the mixture incubated for 1 h at room temperature. Following incubation,

liposomes were collected by centrifugation at $16,000 \times g$ for 20 min, and the supernatant was

collected. The pellet was subjected to three washes in 1 mL of binding buffer. After the washes,

the pellet was resuspended in 40 µL of  2X Laemmli sample buffer. Both samples were reduced

with 100 mM DTT and separated on a 4–20% TGX Stain-Free™ precast gels (Bio-Rad

Laboratories). Proteins were then transferred to a nitrocellulose membrane and subjected to

blotting using the NCR13 antibody.

**Electrophoretic mobility shift assay**

Total RNA was extracted and purified from *B. cinerea* grown in Potato Dextrose Broth (Fisher

Scientific) using the NucleoSpin RNA Plus Mini kit (Takara Bio) according to manufacturer

instructions. The gel shift experiments were performed by mixing 200 ng of the total RNA with

different concentrations of NCR13 PFV1 and PFV2 (0-3 µM) in 20 µL of binding buffer (10 mM

Tris-HCl, pH 8.0, 1 mM EDTA). The reaction mixtures were incubated for 1 h at room temperature

and mixed with 4 µL of 6X gel loading dye (New England Biolabs). Following electrophoresis in

1.2% agarose gel containing SYBR safe DNA gel stain (Thermo Scientific) with TAE buffer at

120 V for 45 min, the gels were visualized using a Bio-Rad ChemiDoc XRS+ system.

*In vitro* **translation assay**

*In vitro* translation was performed using the RiboMAX™ Large Scale RNA Production system

and Wheat Germ Extract *in vitro* translation system (Promega). First, the luciferase DNA template

(Promega) was transcribed to mRNA and translated to luciferase following the manufacturer's

protocol. The luciferase enzyme activity was detected by the luciferase kit (Promega) using a

luminometer (Tecan). Different concentrations of NCR13_PFV1 and NCR13_PFV2 were used to

determine their translation inhibition ability. Sterile water was added to the *in vitro* translation

system as a negative control and a fungal translation inhibitor cycloheximide (48 µM) was used as

a positive control. Translation inhibition (TI) percentage is calculated using this formula TI % =

(1− a/b) x 100, where 'a' equals the translation level with peptide and 'b' equals the translation

level without peptide.

***In planta* and Semi-*In planta* antifungal assays**

Pepper (*Capsicum annuum*, California Wonder 300 TMR) plants used in this study were grown in

a greenhouse for 4 weeks under 14 h light/10 h dark cycle. Tomato (*Solanum lycopersicum* cv.

Mountain Spring) plants were grown in a controlled environment growth chamber for 4 weeks,

under a 16 h light/ 8 h dark cycle. Semi-*in planta* antifungal activity of each peptide against *B.*

*cinerea* was determined using detached leaves of pepper. NCR13_PFV1 and NCR13_PFV2 were

tested at 6, 1.5, and 0.09 μM. Following incubation of 10 μL peptide with 10 μL of *B. cinerea*

spores (~$10^5$ spores/mL) under a humid environment for 48 h, the leaves were photographed in

white light. High-resolution fluorescence images were also taken using CropReporter

(PhenoVation). These images depicted the calculated Fv/Fm (maximum quantum yield of

photosystem II ) values of the diseased area affected by *B. cinerea* infection. Colors in the images

show five different classes ranging from class I to class V (0.000 to 0.700) depicting varying

degrees of tissue damage.

To test the curative antifungal activity of peptides against *B. cinerea*, 4-week-old pepper and

tomato plants were used. Plants were first sprayed with 1 mL of *B. cinerea* spores (about $5 \times 10^4$)

suspended in 1× SFM. After 8 h of *B. cinerea* challenge, 2 mL of NCR13_PFV1 or NCR13_PFV2

(3.0 and 1.5 μM) were sprayed per plant. Control plants were sprayed with 2 mL of water. All

plants were placed in a high humid environment for 72 h. Plants were imaged at 72 h after infection

in white light and high-resolution fluorescence images were taken using CropReporter

(PhenoVation). The quantification of plant health was carried out using the calculated Fv/Fm

(maximum quantum yield of photosystem II) images of the CropReporter showing the efficiency

of photosynthesis in false colors.

For the curative assay using pepper and tomato plants, the picture of the whole plants and detached

leaves were taken as described above. The disease symptoms on individual leaves of each plant,

were visually assessed in % of disease portion (disease severity) and on a rating scale of 0-5 where

each rank would represent the following proportion of diseased tissue per leaf : Scale 0: No obvious

signs/symptoms of disease, Scale 1: $0 > x \leq 20$ % diseased portion of leaf, Scale 2: $20 > x \leq 40$ %

diseased portion of leaf, Scale 3: $40 > x \leq 60$ % diseased portion of leaf, Scale 4: $60 > x \leq 80$ %

diseased portion of leaf, Scale 5: $80 > x \leq 100$ % diseased portion of leaf. Four plants were used

per treatment and for each plant the disease was assessed per leaf. Besides disease severity (DS),

the inhibition of disease by treatment was also evaluated. The following formula was used to assess

the inhibition of disease severity by each treatment: % Disease Inhibition= (Average DS for

Control -Treatment DS)/(Average DS for Control ) x 100%. Where treatment DS is the disease

severity for each leaf except those of the control. Plots were generated using the 'ggplot2' package

in R version 4.3.2.

**Acknowledgements**

The work presented here was supported by the National Science Foundation grant IOS-2037981

and the National Institute of Food and Agriculture GRANT13515963 to D.S and K.C. The

structures for NCR13 were solved using resources at the W.R. Wiley Environmental Molecular

Sciences Laboratory, a national scientific user facility sponsored by U.S. Department of Energy's

Office of Biological and Environmental Research (BER) program located at Pacific Northwest

National Laboratory (PNNL). Battelle operates PNNL for the U.S. Department of Energy. We also

acknowledge imaging support from the Advanced Bioimaging Laboratory (RRID:SCR_018951) at

the Donald Danforth Plant Science Center (DDPSC) and usage of the Leica SPX-8 acquired

through an NSF Major Research Instrumentation grant (DBI-1337680). The Washington

University undergraduate summer intern Bram Osterhout contributed to the early work on NCR13

peptide. We also thank Dr. Ruby Tiwari for providing her comments on this manuscript. We

acknowledge the DDPSC Proteomics & Mass Spectrometry Facility and supported by the National

Science Foundation under Grant No. DBI-1827534 for acquisition of the Orbitrap Fusion Lumos

LC-MS/MS. The greenhouse and growth chamber space provided by the DDPSC Plant Growth

Facility is acknowledged. We also thank Dr. Ruby Tiwari for providing her comments on this

manuscript.

**Author Contributions**

K.C., G.B. and D.S. conceived the research. J.G., A.D-C., S.V., M.T., K.C., M.Z., G.B. and D.S.

designed the research. J.G., A.D-C., S.V., M.T., A.P., R.K, M.Z. and G.B. conducted the

experiments. J.G., A.D-C., S.V., M.T., K.C., M.Z., G.B. and D.S. analyzed and interpreted the

results. J.G., G.B., K.C. and D.S. wrote the manuscript.

**Competing interests**. The authors declare no competing interests.

**Data deposition**: The coordinates of the NCR13 NMR structure have been deposited in the Protein

Data Bank, www.rcsb.org 8ULM (NCR13_ PFV1) and 7TH8 (NCR13_ PFV2) and the NMR

chemical shifts have been deposited in the BioMagResBank, www.bmrb.wisc.edu BMRB-31111

(oxidized NCR13_PFV1) and BMRB-30979 (oxidized NCR13_PFV2).

A

```
CaNCR13  -TKPCQSDKDCKKFACRKPKVPKCINGFCKCVR-  32
CaNCR15  TKQPCKSRKHCKTYRCPTPKVPNCVNGFCKCVR-  33
CaNCR07  KKMPCKRRRDCKTYPCPHPKVRDCVKGYCKCVVR  34
```

B

| Peptide | Net Charge | % Hydrophobic amino acids | Predicted MW (kDa) | Isoelectric point | % identity with NCR13 |
|---------|-----------|---------------------------|--------------------|--------------------|-----------------------|
| CaNCR13 | +8 | 28.1 | 3.66 | 9.59 | 100 |
| CaNCR15 | +9 | 24.2 | 3.81 | 9.82 | 66.6 |
| CaNCR07 | +10 | 26.5 | 4.08 | 9.85 | 48.4 |

**Figure 1. Primary amino acid sequence and properties of NCR13.** (A) Primary amino acid sequences for Chickpea (*C. arientinum*) NCR13 and its homologs NCR15 and NCR07. Conserved

818 cysteine residues are highlighted in red. The number of amino acids is indicated at the end of each
819 sequence. (B) Peptide characteristics of CaNCR13, CaNCR15 and CaNCR07. (C) High-
820 performance liquid chromatography (HPLC) analyses of NCR13 produced recombinantly in
821 *Pichia pastoris*, monitored at 260 nm, reveals the presence of two major products. mAU represents
822 milli-Absorbing Units. (D) Chemically synthesized NCR13 elutes as one HPLC peak.

**Figure 2. Fingerprint $^1$H-$^{15}$N HSQC spectra for oxidized and reduced NCR13.** Assigned $^1$H-
$^{15}$N HSQC spectra for oxidized (disulfide) NCR13_PFV1 (A) and NCR13_PFV2 (B). (C) The $^1$H-
$^{15}$N HSQC spectrum of fully reduced (dithiol) NCR13_PFV1 and NCR13_PFV2 were identical
(the spectrum for NCR13_PFV2 is illustrated). All spectra collected on ~ 1 mM samples at 20°C
in 20 mM sodium acetate, 50 mM NaCl, pH 5.3 at a $^1$H resonance frequency of 600 MHz. Amide
side chain resonance pairs are connected by a red dashed line. Not within the displayed spectral
boundaries is the G26 amide resonance for NCR13_PFV1 and NCR13_PFV2 at 8.66/103.1 ppm
and 7.99/101.7 ppm, respectively, and the I24 amide resonance for NCR13_PFV1 at 8.72/132.0
ppm.

**Figure 3. Three-Dimensional NMR solution structure of NCR13_PFV1 and NCR13_PFV2.**
Cartoon representation of the backbone superposition of the ordered regions in the ensemble of 20 structures calculated for oxidized NCR13_PFV1 (8ULM) and NCR13_PFV2 (7TH8). β-strands are colored blue and labeled sequentially starting from the N-terminal and the lone α-helix is colored wheat. Next to each ensemble is a backbone stick representation of a single structure with the six oxidized cysteine side chains highlighted in yellow, red, or cyan. Above the structures is a schematic summary of the elements of secondary structure observed for the two NMR structures, color-coded similarly. Also shown in the schematic are the disulfide bond connections in both isomers. One disulfide bond, between C15 and C30 (red), is identical in both isomers with C10 and C28 swapping disulfide bond partners.

A

|  | 4 10 15 23 28 30 |
|---|---|
| NCR13 | TKP**C**QSDKD**C**KKFA**C**RKPKVPK**C**INGF**C**K**C**VR |
| NCR13_C4S-C23S | TKP**S**QSDKDCKKFACRKPKVPK**S**INGFCKCVR |
| NCR13_C10S-C28S | TKPCQSDKD**S**KKFACRKPKVPKCINGF**S**KCVR |
| NCR13_C15S-C30S | TKPCQSDKDCKKFA**S**RKPKVPKCINGFCK**S**VR |
| NSR13 | TKP**S**QSDKD**S**KKFA**S**RKPKVPK**S**INGF**S**K**S**VR |

B

C

D

| Peptide Name | MIC (µM) |
|---|---|
| NCR13_PFV1 | 0.09 |
| NCR13_C4S-C23S | 1.5 |
| NCR13_C10S-C28S P1 | 3 |
| NCR13_C10S-C28S P2 | 3 |
| NCR13_C15S-C30S | 0.75 |
| NSR13 | Inactive |

**Figure 5. Removing disulfide bonds in NCR13_PFV1 reduces antifungal activity against *B. cinerea in vitro*.** (A) Four NCR13 constructs are used to assay the importance of disulfide bond formation for antifungal activity by substituting pairs of Cys residues with Ser (highlighted in red). The numbers above the sequence indicate the position of the Cys substitution. (B) HPLC analyses of NCR13_C4S-C23S, NCR13_C10S-C28S, and NCR13_C15S-C30S recombinantly produced in *P. pastoris* monitored at 260 nm (mAU = milli-Absorbing Units). (C) Fungal cell viability assay of disulfide knockout variants. A color change from blue to pink/clear indicative of metabolically active (live) *B. cinerea* germlings after 60 h. (D) Summary of the MIC values for NCR13_PFV1 and NCR13 disulfide knockout variants.

A

| Peptide | Sequence | MIC (μM) |
|---|---|---|
| NCR13_CS | TKPCQSDKDCKKFACRKPKVPKCINGFCKCVR | 3 |
| NCR13_AlaV1 | TKPC**AAAAA**CKKFACRKPKVPKCINGFCKCVR | 6 |
| NCR13_AlaV2 | TKPCQSDKDC**AAAA**CRKPKVPKCINGFCKCVR | Inactive (>12) |
| NCR13_AlaV3 | TKPCQSDKDCKKFAC**AAAA**VPKCINGFCKCVR | Inactive (>12) |
| NCR13_AlaV4 | TKPCQSDKDCKKFACRKPK**AAA**CINGFCKCVR | Inactive (>12) |
| NCR13_AlaV5 | TKPCQSDKDCKKFACRKPKVPKC**AAAA**CKCVR | 12 |

B

**Figure 6. Active motif of NCR13 required for antifungal activity.** (A) Antifungal activity of
NCR13 synthetic peptide and NCR13 alanine variants against *B. cinerea*. Alanine scanning
mutagenesis was done by substituting alanine within the NCR13 core sequence, leading to the
creation of the NCR13_AlaV1-V5. MIC values were determined against *B. cinerea*. The amino
acid substitutions are shown in red. Representative pictures are shown in Fig. *S5*. (B) Active motif
of NCR13 (highlighted with a bold red font) required for its antifungal activity against *B. cinerea*.
The secondary structure elements are shown above the amino acid sequence. The gray line and
green arrow indicate α-helix (α1) and β-strand (β1) respectively.

**Figure 7. NCR13 PFV1 permeabilizes the cell membrane of *B. cinerea* rapidly.** (A) Confocal microscopy images of SYTOX Green (SG) uptake in *B. cinerea* germlings at different time points. Germlings were treated with 0.09 μM NCR13_PFVs and SG dye simultaneously for 20 min. No peptide is used as a control. Scale bars = 10 μm. The experiment was repeated thrice with similar results. (B) Quantitative measurements of average fluorescence intensity over time to assess cell membrane permeability in *B. cinerea* germlings treated with 0.09 μM NCR13 PFVs. (C) PIP strip showing binding of NCR13 PFVs to multiple phospholipids, including phosphatidylinositol 4,5-bisphosphate (PI(4,5)P$_2$), phosphatidylinositol (3,4,5)-trisphosphate PI(3,4,5)P$_3$ and phosphatidylinositol monophosphates PI(3)P, PI(4)P, and PI(5)P, phosphatidylinositol

bisphosphate PI(3,4)P$_2$. (D) Integrated density measurement of PIP strip probed with NCR13_PFVs. Datapoints are means ± SE of two independent biological replicates ($N = 2$). (E) Binding of NCR13_PFVs to PI(4)P and PI(4,5)P2 (PIP$_2$) containing liposomes. NCR13_PFV's were incubated without liposomes (None) and liposomes bearing no phosphoinositides (control), S- supernatant fraction and P- pellet fraction.

**Figure 8. Internalization and localization of NCR13 PFVs in *B. cinerea* germlings.** (A)
Confocal microscopy images of *B. cinerea* germlings at different points treated with 0.09 μM
DyLight550-labeled NCR13_PFV1 (left) and NCR_PFV2 (right) for 30 minutes. Scale bars = 10
μm. Data represents representative results of three independent experiments. (B) Quantification of
peptide internalization was conducted by analyzing the average fluorescence intensities over time
(in minutes) for NCR13_PFVs. The upper threshold for background was set at 23 for Fiji analysis.
(C) Confocal microscopy images of *B. cinerea* germlings treated with 1.5 μM DyLight550-labeled
NCR13_PFVs for 30 minutes. White arrowheads indicate NCR13_PFV2 on the cell surface. Scale
bars = 10 μm. (D) Quantitative measurements of peptide internalization in *B. cinerea* germlings
treated with 1.5 μM NCR13_PFV1 (left) and NCR13_PFV2 (right). (E) Internalized DyLight550-
labeled NCR13_PFV1 (left) and NCR13_PFV2 (right) concentrate in specific foci (white
arrowheads) within *B. cinerea* germlings. Scale bars = 10 μm. (F) NCR13_PFV1 and
NCR13_PFV2 localize to the nucleolus. DyLight550-labeled NCR13_PFV1 (upper panel) and
PFV2 (lower panel) colocalize with rRNA-specific Nucleolus Bright Green stain. Hoechst 33258
is used for staining nucleus. White arrowheads indicate nucleolar localization. Scale bars = 5 μm.
(G) Semi-quantitative analysis of NCR13_PFVs within *B. cinerea* germlings. Germlings were
treated with 1.5 μM NCR13_PFV1 and NCR13_PFV2 for 0, 2, and 4 hours. Total proteins were
extracted from fungal cells and immunoblotted using anti-NCR13 antibody. Equal loading of
proteins was verified using Ponceau staining. The bar graph represents integrated density
measurements. Statistical significance was tested using paired student's t-test. Data are shown as
mean ± SD (N = 3, where N refers to biological replicates).

**Figure 9. NCR13_PFV1 exhibits rRNA binding and higher translation inhibition activity.** (A) rRNA binding by NCR13_PFVs tested using electrophoretic mobility shift assay (EMSA). *Botrytis cinerea* rRNA was used to assess the binding activity of the peptides NCR13_PFV1 and NCR13_PFV2. The binding reactions were subjected to electrophoresis, and the mobility shifts were visualized on an agarose gel (1%). The concentration gradient of peptides is indicated above the lanes. The first lane on the left contains a molecular weight marker. (B) Relative *in vitro* translational inhibition (%) in the presence of NCR13_PFVs at different concentrations. Sterile water was used as negative control and cycloheximide (96 µM) was used as positive control. N= 3, 'N' indicates biological replicates. Statistical significance is determined using One-way ANOVA with Tukey multiple comparisons test (**** indicates p value < 0.0001).

**Figure 10. NCR13 PFV1 exhibits enhanced curative antifungal activity against gray mold in pepper (*Capsicum annuum*) plants.** (A) Curative antifungal activity of NCR13_PFV1 and PFV2. Four-week-old pepper plants were first spray-inoculated using 1 mL of $5 \times 10^4$ *B. cinerea* spores. At 8 hpi (hours post inoculation), the plants were sprayed with 2 mL of peptide (NCR13_PFV1 or PFV2) at a concentration of 1.5 or 3 μM or control (*B. cinerea* alone). Mock-treated plants were only sprayed with 1X SFM (no fungal spore suspension). (B) Calculated photosynthetic quantum yield (Fv/Fm) for individual pepper plants. (C) Disease severity (%) was calculated as described

in materials and methods. (B-C) Each data point represents the average of 4 leaves per plant for 4 plants for each treatment. Horizontal lines represent the median and boxes indicate the 25th and 75th percentiles. The Kruskal-Wallis test with Dunn's multiple comparisons test is used to determine statistically significant differences between control and peptide-treated inoculated plants. Two independent experiments were performed with similar results.

1009

**Figure 11.** Graphical abstract illustrating the biosynthesis of disulfide linkage variants of NCR13_PFV1 and NCR13_PFV2 in *P. pastoris*, their structures and their effects on antifungal activity and modes of action in *B. cinerea* germlings. The figure depicts the differences in cell penetration, localization, rRNA binding and translation inhibition, membrane permeabilization, phospholipid binding, and overall fungal cell killing potency. Created using BioRender.com.

1015

1016

1017

1018

1019

1020

1021

## References

1022

Alhhazmi, A. A., Alluhibi, S. S., Alhujaily, R., Alenazi, M. E., Aljohani, T. L., Al-Jazzar, A. A., Aljabri, A.
D., Albaqami, R., Almutairi, D., Alhelali, L. K., Albasri, H. M., Almutawif, Y. A., Alturkostani,
M. A., & Almutairi, A. Z. (2024). Novel antimicrobial peptides identified in legume plant,
Medicago truncatula. *Microbiol Spectr*, *12*(2), e0182723.
https://doi.org/10.1128/spectrum.01827-23
Balogh, E. P., Mosolygo, T., Tiricz, H., Szabo, A. M., Karai, A., Kerekes, F., Virok, D. P., Kondorosi, E., &
Burian, K. (2014). Anti-chlamydial effect of plant peptides. *Acta Microbiol Immunol Hung*,
*61*(2), 229-239. https://doi.org/10.1556/AMicr.61.2014.2.12
Bhardwaj, G., Mulligan, V. K., Bahl, C. D., Gilmore, J. M., Harvey, P. J., Cheneval, O., Buchko, G. W.,
Pulavarti, S. V., Kaas, Q., Eletsky, A., Huang, P. S., Johnsen, W. A., Greisen, P. J., Rocklin, G.
J., Song, Y., Linsky, T. W., Watkins, A., Rettie, S. A., Xu, X.,...Baker, D. (2016). Accurate de
novo design of hyperstable constrained peptides. *Nature*, *538*(7625), 329-335.
https://doi.org/10.1038/nature19791
Bhattacharya, A., Tejero, R., & Montelione, G. T. (2007). Evaluating protein structures determined by
structural genomics consortia. *Proteins*, *66*(4), 778-795. https://doi.org/10.1002/prot.21165
Buchko, G. W., Hewitt, S. N., Van Voorhis, W. C., & Myler, P. J. (2018). Solution NMR structures of
oxidized and reduced Ehrlichia chaffeensis thioredoxin: NMR-invisible structure owing to
backbone dynamics. *Acta Crystallogr F Struct Biol Commun*, *74*(Pt 1), 46-56.
https://doi.org/10.1107/S2053230X1701799X
Buchko, G. W., Pulavarti, S., Ovchinnikov, V., Shaw, E. A., Rettie, S. A., Myler, P. J., Karplus, M.,
Szyperski, T., Baker, D., & Bahl, C. D. (2018). Cytosolic expression, solution structures, and
molecular dynamics simulation of genetically encodable disulfide-rich de novo designed
peptides. *Protein Sci*, *27*(9), 1611-1623. https://doi.org/10.1002/pro.3453
Bulleid, N. J., & Ellgaard, L. (2011). Multiple ways to make disulfides. *Trends Biochem Sci*, *36*(9),
485-492. https://doi.org/10.1016/j.tibs.2011.05.004
Chadha, S., & Kale, S. P. (2015). Simple fluorescence-based high throughput cell viability assay for
filamentous fungi. *Lett Appl Microbiol*, *61*(3), 238-244. https://doi.org/10.1111/lam.12460
Cools, T. L., Struyfs, C., Cammue, B. P., & Thevissen, K. (2017). Antifungal plant defensins:
increased insight in their mode of action as a basis for their use to combat fungal infections.
*Future Microbiol*, *12*, 441-454. https://doi.org/10.2217/fmb-2016-0181
Djami-Tchatchou, A. T., Tetorya, M., Godwin, J., Codjoe, J. M., Li, H., & Shah, D. M. (2023). Small
Cationic Cysteine-Rich Defensin-Derived Antifungal Peptide Controls White Mold in
Soybean. *J Fungi (Basel)*, *9*(9). https://doi.org/10.3390/jof9090873
Dos Santos-Silva, C. A., Zupin, L., Oliveira-Lima, M., Vilela, L. M. B., Bezerra-Neto, J. P., Ferreira-
Neto, J. R., Ferreira, J. D. C., de Oliveira-Silva, R. L., Pires, C. J., Aburjaile, F. F., de Oliveira, M.
F., Kido, E. A., Crovella, S., & Benko-Iseppon, A. M. (2020). Plant Antimicrobial Peptides:
State of the Art, In Silico Prediction and Perspectives in the Omics Era. *Bioinform Biol
Insights*, *14*, 1177932220952739. https://doi.org/10.1177/1177932220952739
Farkas, A., Maroti, G., Durgo, H., Gyorgypal, Z., Lima, R. M., Medzihradszky, K. F., Kereszt, A.,
Mergaert, P., & Kondorosi, E. (2014). Medicago truncatula symbiotic peptide NCR247
contributes to bacteroid differentiation through multiple mechanisms. *Proc Natl Acad Sci U
S A*, *111*(14), 5183-5188. https://doi.org/10.1073/pnas.1404169111
Farkas, A., Maroti, G., Kereszt, A., & Kondorosi, E. (2017). Comparative Analysis of the Bacterial
Membrane Disruption Effect of Two Natural Plant Antimicrobial Peptides. *Front Microbiol*, *8*,
51. https://doi.org/10.3389/fmicb.2017.00051

Goyal, R. K., & Mattoo, A. K. (2014). Multitasking antimicrobial peptides in plant development and
host defense against biotic/abiotic stress. *Plant Sci*, *228*, 135-149.
https://doi.org/10.1016/j.plantsci.2014.05.012
Guntert, P. (2004). Automated NMR structure calculation with CYANA. *Methods Mol Biol*, *278*, 353-
378. https://doi.org/10.1385/1-59259-809-9:353
Haag, A. F., Kerscher, B., Dall'Angelo, S., Sani, M., Longhi, R., Baloban, M., Wilson, H. M., Mergaert,
P., Zanda, M., & Ferguson, G. P. (2012). Role of cysteine residues and disulfide bonds in the
activity of a legume root nodule-specific, cysteine-rich peptide. *J Biol Chem*, *287*(14),
10791-10798. https://doi.org/10.1074/jbc.M111.311316
Hawkins, N. J., & Fraaije, B. A. (2018). Fitness Penalties in the Evolution of Fungicide Resistance.
*Annu Rev Phytopathol*, *56*, 339-360. https://doi.org/10.1146/annurev-phyto-080417-050012
Hellman, L. M., & Fried, M. G. (2007). Electrophoretic mobility shift assay (EMSA) for detecting
protein-nucleic acid interactions. *Nat Protoc*, *2*(8), 1849-1861.
https://doi.org/10.1038/nprot.2007.249
Isozumi, N., Masubuchi, Y., Imamura, T., Mori, M., Koga, H., & Ohki, S. (2021). Structure and
antimicrobial activity of NCR169, a nodule-specific cysteine-rich peptide of Medicago
truncatula. *Sci Rep*, *11*(1), 9923. https://doi.org/10.1038/s41598-021-89485-w
Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
Bates, R., Zidek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J.,
Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J.,...Hassabis, D. (2021). Highly
accurate protein structure prediction with AlphaFold. *Nature*, *596*(7873), 583-589.
https://doi.org/10.1038/s41586-021-03819-2
Karbalaei, M., Rezaee, S. A., & Farsiani, H. (2020). Pichia pastoris: A highly successful expression
system for optimal synthesis of heterologous proteins. *J Cell Physiol*, *235*(9), 5867-5881.
https://doi.org/10.1002/jcp.29583
Kereszt, A., Mergaert, P., Montiel, J., Endre, G., & Kondorosi, E. (2018). Impact of Plant Peptides on
Symbiotic Nodule Development and Functioning. *Front Plant Sci*, *9*, 1026.
https://doi.org/10.3389/fpls.2018.01026
Kondorosi, E., Mergaert, P., & Kereszt, A. (2013). A paradigm for endosymbiotic life: cell
differentiation of Rhizobium bacteria provoked by host plant factors. *Annu Rev Microbiol*,
*67*, 611-628. https://doi.org/10.1146/annurev-micro-092412-155630
Kovaleva, V., Bukhteeva, I., Kit, O. Y., & Nesmelova, I. V. (2020). Plant Defensins from a Structural
Perspective. *Int J Mol Sci*, *21*(15). https://doi.org/10.3390/ijms21155307
Lee, W., Rahimi, M., Lee, Y., & Chiu, A. (2021). POKY: a software suite for multidimensional NMR and
3D structure calculation of biomolecules. *Bioinformatics*, *37*(18), 3041-3042.
https://doi.org/10.1093/bioinformatics/btab180
Li, H., Kalunke, R., Tetorya, M., Czymmek, K. J., & Shah, D. M. (2024). Modes of action and potential
as a peptide-based biofungicide of a plant defensin MtDef4. *Mol Plant Pathol*, *25, e134538*.
*https://doi.org/10.1111/mpp.134538*.
Li, H., Velivelli, S. L., & Shah, D. M. (2019). Antifungal potency and modes of action of a novel olive
tree defensin against closely related ascomycete fungal pathogens. . *Mol Plant Microbe*
*Interact*, *32*, 164901664. https://doi.org/10.1094/MPMI-08-19-0224-R
Liang, J., Shah, D. M., Wu, Y., Rosenberger, C. A., & Hakimi, S. M. (2001). *Antifungal polypeptide*
*from alfalfa and methods for controlling plant pathogenic fungi* (USA Patent No. US
6316407).
Lima, R. M., Kylarova, S., Mergaert, P., & Kondorosi, E. (2020). Unexplored Arsenals of Legume
Peptides With Potential for Their Applications in Medicine and Agriculture. *Front Microbiol*,
*11*, 1307. https://doi.org/10.3389/fmicb.2020.01307

Lobo, F., & Boto, A. (2022). Host-defense peptides as new generation phytosanitaries: Low toxicity
and low induction of antimicrobial resistance. *Agronomy*, *12*, 1614.
https://doi.org/https://doi.org/10.3390/agronomy12071614
Maroti, G., Downie, J. A., & Kondorosi, E. (2015). Plant cysteine-rich peptides that inhibit pathogen
growth and control rhizobial differentiation in legume nodules. *Curr Opin Plant Biol*, *26*, 57-
63. https://doi.org/10.1016/j.pbi.2015.05.031
McLaughlin, M. S., Roy, M., Abbasi, P. A., Carisse, O., Yurgel, S. N., & Ali, S. (2023). Why Do We Need
Alternative Methods for Fungal Disease Management in Plants? *Plants (Basel)*, *12*(22).
https://doi.org/10.3390/plants12223822
Mergaert, P. (2018). Role of antimicrobial peptides in controlling symbiotic bacterial populations.
*Nat Prod Rep*, *35*(4), 336-356. https://doi.org/10.1039/c7np00056a
Montiel, J., Downie, J. A., Farkas, A., Bihari, P., Herczeg, R., Balint, B., Mergaert, P., Kereszt, A., &
Kondorosi, E. (2017). Morphotype of bacteroids in different legumes correlates with the
number and type of symbiotic NCR peptides. *Proc Natl Acad Sci U S A, 114*(19), 5041-5046.
https://doi.org/10.1073/pnas.1704217114
Montiel, J., Szucs, A., Boboescu, I. Z., Gherman, V. D., Kondorosi, E., & Kereszt, A. (2016). Terminal
Bacteroid Differentiation Is Associated With Variable Morphological Changes in Legume
Species Belonging to the Inverted Repeat-Lacking Clade. *Mol Plant Microbe Interact*, *29*(3),
210-219. https://doi.org/10.1094/MPMI-09-15-0213-R
Olliver, C. L., Grobler-Rabie, A., & Boyd, C. D. (1985). In vitro translation of messenger RNA in a
wheat germ extract cell-free system. *Methods Mol Biol*, *2*, 137-144.
https://doi.org/10.1385/0-89603-064-4:137
Ordogh, L., Voros, A., Nagy, I., Kondorosi, E., & Kereszt, A. (2014). Symbiotic plant peptides
eliminate Candida albicans both in vitro and in an epithelial infection model and inhibit the
proliferation of immortalized human cells. *Biomed Res Int*, *2014*, 320796.
https://doi.org/10.1155/2014/320796
Parisi, K., Shafee, T. M. A., Quimbar, P., van der Weerden, N. L., Bleackley, M. R., & Anderson, M. A.
(2019). The evolution, function and mechanisms of action for plant defensins. *Semin Cell
Dev Biol*, *88*, 107-118. https://doi.org/10.1016/j.semcdb.2018.02.004
Poon, I., Baxter, A. A., Lay, F. T., Mills, G. D., Adda, C. G., Payne, J. A., Phan, T. K., Ryan, G. F., White, J.
A., Veneer, P. K., van der Weerden, N. L., Anderson, M. A., Kvansakul, M., & Hulett, M. D.
(2014). Phosphoinositide-mediated oligomerization of a defensin induces cell lysis. *Elife*, *3*,
e01808. https://doi.org/10.7554/eLife.01808
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K.,
Tomancak, P., & Cardona, A. (2012). Fiji: an open-source platform for biological-image
analysis. *Nat Methods*, *9*(7), 676-682. https://doi.org/10.1038/nmeth.2019
Shabab, M., Arnold, M. F., Penterman, J., Wommack, A. J., Bocker, H. T., Price, P. A., Griffitts, J. S.,
Nolan, E. M., & Walker, G. C. (2016). Disulfide cross-linking influences symbiotic activities
of nodule peptide NCR247. *Proc Natl Acad Sci U S A, 113*(36), 10157-10162.
https://doi.org/10.1073/pnas.1610724113
Shaw, W. J., Tarasevich, B. J., Buchko, G. W., Arachchige, R. M. J., & Burton, S. D. (2020). Controls of
nature: Secondary, tertiary, and quaternary structure of the enamel protein amelogenin in
solution and on hydroxyapatite. *J Struct Biol*, *212*(3), 107630.
https://doi.org/10.1016/j.jsb.2020.107630
Shen, Y., Delaglio, F., Cornilescu, G., & Bax, A. (2009). TALOS+: a hybrid method for predicting
protein backbone torsion angles from NMR chemical shifts. *J Biomol NMR*, *44*(4), 213-223.
https://doi.org/10.1007/s10858-009-9333-z

Tavormina, P., De Coninck, B., Nikonorova, N., De Smet, I., & Cammue, B. P. (2015). The Plant
Peptidome: An Expanding Repertoire of Structural Features and Biological Functions. *Plant
Cell*, *27*(8), 2095-2118. https://doi.org/10.1105/tpc.15.00440

Tiricz, H., Szűcs, A., Farkas, A., Pap, B., Lima, R. M., Maróti, G., Kondorosi, É., & Kereszt, A. (2013).
Antimicrobial Nodule-Specific Cysteine-Rich Peptides Induce Membrane Depolarization-
Associated Changes in the Transcriptome. *Applied and Environmental Microbiology*, *79*(21),
6737. https://doi.org/10.1128/AEM.01791-13

van der Weerden, N. L., Bleackley, M. R., & Anderson, M. A. (2013). Properties and mechanisms of
action of naturally occurring antifungal peptides. *Cell Mol Life Sci*, *70*(19), 3545-3570.
https://doi.org/10.1007/s00018-013-1260-1

Velivelli, S. L. S., Czymmek, K. J., Li, H., Shaw, J. B., Buchko, G. W., & Shah, D. M. (2020). Antifungal
symbiotic peptide NCR044 exhibits unique structure and multifaceted mechanisms of
action that confer plant protection. *Proc Natl Acad Sci U S A*, *117*(27), 16043-16054.
https://doi.org/10.1073/pnas.2003526117

Wedemeyer, W. J., Welker, E., Narayan, M., & Scheraga, H. A. (2000). Disulfide bonds and protein
folding. *Biochemistry*, *39*(23), 7032. https://doi.org/10.1021/bi005111p

Yee, A., Chang, X., Pineda-Lucena, A., Wu, B., Semesi, A., Le, B., Ramelot, T., Lee, G. M.,
Bhattacharyya, S., Gutierrez, P., Denisov, A., Lee, C. H., Cort, J. R., Kozlov, G., Liao, J., Finak,
G., Chen, L., Wishart, D., Lee, W.,...Arrowsmith, C. H. (2002). An NMR approach to
structural proteomics. *Proc Natl Acad Sci U S A*, *99*(4), 1825-1830.
https://doi.org/10.1073/pnas.042684599

1185