Infectious subgenomic amplicon strategies for Japanese encephalitis and West Nile viruses

Prince Pal Singh[1,2], Nguyen Phuong Khanh Le[1], Uladzimir Karniychuk[1,*]

[1]Department of Veterinary Biosciences, College of Veterinary Medicine, The Ohio State

University, Columbus, OH, USA

[2]School of Public Health, University of Saskatchewan, Saskatoon, Canada

*Correspondence: Uladzimir Karniychuk

Department of Veterinary Biosciences, College of Veterinary Medicine, The Ohio State

University, Columbus, OH, USA

karniychuk.1@osu.edu

ABSTRACT

Classical methods for constructing infectious cDNA clones of flaviviruses are often hindered by instability and toxicity. The Infectious-Subgenomic-Amplicons (ISA) method is an advancement which utilizes overlapping DNA fragments representing viral genomic sequence and in-cell recombination to bypass bacterial plasmid assembly. However, the ISA method has limitations due to the toxicity of some ISA DNA fragments in bacteria during synthetic production. We validated modified ISA strategies for producing toxic ISA Japanese encephalitis virus (JEV) and West Nile virus (WNV) DNA fragments. Three approaches were explored including subdividing toxic DNA fragments into two sub-fragments for synthetic clonal production, using a low-copy bacterial plasmid, and subdividing the toxic DNA fragments into four short overlapping sub-fragments, each up to 1.8 kb. The latter novel approach in ISA applications enabled the synthesis of entirely bacteria-free ISA DNA fragments. Our results demonstrate that subdividing toxic fragments into sub-fragments smaller than 1.8 kb for synthesis is the efficient strategy, circumventing the need for bacterial plasmids and ensuring rapid production of synthetic flaviviruses. This method also shortens the production timeline. We also compared the efficacy of JEV and WNV ISA in zinc finger antiviral protein 1 (ZAP) wild-type and knockout cells and found that knockout cells may be more effective for ISA rescue of flaviviruses, including attenuated strains for live attenuated vaccines. The validated modified ISA strategies provide an efficient approach for producing synthetic JEV and WNV. This will enable rapid research during outbreaks of emerging flaviviruses by facilitating the quick generation of new virus variants.

Flavivirus, Japanese encephalitis virus, West Nile virus, dengue virus, infectious subgenomic amplicons, ZAP, reverse genetics, infectious clones

INTRODUCTION

Flaviviruses are single-stranded RNA viruses transmitted mainly by arthropods, as well as sexually and transplacentally. The genus Flavivirus (family *Flaviviridae*) comprises viruses which cause human disease, including dengue virus (DENV), Japanese encephalitis virus (JEV), and West Nile virus (WNV). These flaviviruses infect millions of people each year, causing symptoms from mild fevers to fatal hemorrhagic and neurological diseases [1]. The rising number of global flavivirus infections, frequent outbreaks, and expanding geographical range pose significant public health concerns [1]. Flaviviruses also pose constant threats to livestock industries [2]. There are no approved human vaccines for certain flaviviruses, such as Zika virus and WNV. Moreover, no licensed antivirals exist against flaviviruses.

There are four DENV serotypes which cause illnesses ranging from mild dengue fever to severe life-threatening dengue. Dengue virus is the major cause of hospitalization and death in children in Southeast Asian countries. Annually, DENV results in approximately 390 million infections, 100 million clinically apparent cases, and 500,000 severe cases, with at least 2.5 billion people at risk [1,3]. Over the past 70 years, infection rates have steadily increased, making DENV the most prevalent arthropod-borne viral disease worldwide. By comparison, only sporadic DENV epidemics were documented before the Second World War [1,4]. Between 2010 and 2022, more than 33,000 locally acquired dengue cases have been reported in the USA. Since 2014, there have been fewer than 1,000 cases per year, but 2022 was the first time in a decade that cases reached four digits. In 2023, Florida officials issued a warning about dengue as the cases continued to increase.

Japanese encephalitis virus causes encephalitis in humans in the Asia-Pacific region [5], with

68,000 annual cases and 15,000 deaths [6]. JEV epidemics were originally described in Japan in

the 19th century, and the virus was first recovered in 1935 from an infected human in Tokyo [1].

The enzootic cycle of JEV is between water birds and *Culex* mosquitoes, with pigs also serving

as an amplifying host [1]. Humans are considered incidental dead-end hosts and generally do not

produce viremia sufficient to infect mosquitoes. Though many human infections are

asymptomatic, around 20-30% of clinical infections are fatal, and 30-50% of the survivors

develop life-long neurological sequelae [5]. Human transplacental and fetal JEV infections are

also reported [7]. The geographic range of emerging JEV keeps expanding from the South of

Russia to Australia, Japan, Eastern China, India, and South-East Asia. Also, severe travel-related

cases are reported around the globe [8]. There is a concern that JEV can be introduced into North

America given a large population of susceptible avian species and amplifying hosts—pigs, wild

boars; susceptible *Culex* mosquitoes are also ubiquitous [9-11]. In addition to morbidity and

mortality in humans, JEV causes economically significant clinical disease in pigs, mostly

abortions and stillbirth [11,12]. Japanese encephalitis virus keeps expanding—the 2022 JEV

outbreak in Australia with infections in swine herds, zoonotic transmission, and human deaths

was caused by genotype IV which was not associated with outbreaks before [13].

West Nile virus (WNV) is classified by the National Institute of Allergy and Infectious Diseases

as a "Category B Priority" pathogen, indicating its high threat level. West Nile virus cycles

between *Culex* mosquitoes and birds, but it can also infect humans, horses, and other mammals.

The virus is widespread, found in Africa, the Middle East, West Asia, Australia, the Americas,

and Europe. The first major European outbreak occurred in Romania in 1996, with the largest in

2018 affecting 12 countries. In the U.S., WNV emerged in 1999, rapidly spreading and causing

significant morbidity and annual outbreaks, notably in 2002, 2003, and 2012 [1,14].

West Nile virus is neuroinvasive, leading to central nervous system damage, including

meningitis and encephalitis, which can be fatal or cause long-term disabilities such as a

poliomyelitis-like syndrome. The virus's spread has increased due to expanding mosquito

habitats and changing bird migration patterns caused by climate change. Currently, it is the most

common insect-borne zoonotic virus in the U.S., with no approved human vaccines or antivirals.

From 1999 to 2010, blood supply screening estimated 2 to 4 million total infections in the U.S.

[15]. In 2022, Centers for Disease Control and Prevention reported 1,126 cases and 93 deaths in

the U.S., and 2,406 cases and 1,599 neuroinvasive diseases in 2023, underscoring growing cases

and inadequate environmental controls.

Reverse genetics, defined as the generation of viruses entirely from complementary DNA

(cDNA) [16], has revolutionized virology research, including research of flaviviruses. This

powerful technique enables the experimental editing of viral genomes to investigate effects of

point mutations, host-pathogen interactions, virulence factors, mechanisms of immune evasion,

and the development of reporter viruses, vaccines, and antiviral testing [17,18]. The classical

method involves constructing an infectious DNA clone. This clone is typically a bacterial DNA

plasmid or two plasmids containing the entire cDNA of the viral genome and a suitable

promoter, such as the T7 promoter [17,19]. In this approach, viral RNA is obtained from

biological specimens, and a double-stranded cDNA copy of the RNA virus genome is generated

and incorporated into plasmids, which are then amplified in bacteria. After plasmid purification,

infectious viruses are produced either by direct transfection of permissive cells or by transfection

of genomic RNA obtained through *in vitro* transcription [17,19].

Despite advances in molecular techniques, classical methods for constructing infectious cDNA

clones are time-consuming, unpredictable, and labor-intensive and frequently associated with

undesirable mutations or unstable and toxic clones in bacteria [17,19]. Flavivirus infectious DNA

clones are particularly challenging due to the instability and toxicity of some viral sequences in

bacterial hosts [17-23]. To mitigate these difficulties, bacteria-free approaches have been

developed, including long PCR, circular polymerase extension reaction, Gibson assembly, and

Infectious-Subgenomic-Amplicons (ISA) [17].

The ISA method represents a significant advancement in reverse genetics for flaviviruses [24]. It

allows to bypass the problematic step of assembling and amplifying the full-length flavivirus

genome in bacterial plasmids and bacterial cells. Instead, the ISA process relies on in-cell

recombination, which simplifies and shortens the procedure where overlapping PCR-amplified

ISA DNA fragments, flanked with a pCMV promoter and HDR/SV40pA sequences, are directly

transfected into mammalian cells, insect cells, or *in vivo* in mice [17,25]. For example, in the

most recent study using the classical ISA the authors rescued DENV; in contrast cloning the full-

length DENV genome in bacteria failed due to toxicity [19].

While ISA is termed a "bacteria-free" approach, it is not completely free of bacteria and

associated toxicity issues. Specifically, to our knowledge all sixteen published studies utilizing

ISA for flavivirus rescue have used PCR amplification of fragments from pre-existing full-length

flavivirus DNA infectious clones, infected biological specimens, or synthetic overlapping DNA

ISA fragments. These later synthetic DNA fragments, provided by commercial suppliers, are

assembled, inserted into bacterial plasmids, and amplified in bacteria [19,24-38]. Although this

approach avoids constructing unstable and toxic plasmids containing sequences of the entire viral

genome, some synthetic ISA DNA fragments still may pose bacterial toxicity, slowing or halting

their commercial production. Indeed, in several of these papers, along with commercially

generated ISA DNA fragments, the authors PCR-amplified some ISA DNA fragments directly

from historically preexisting entire-genome-length flavivirus infectious DNA clones or from

infected biological specimens to rescue yellow fever virus, DENV, and JEV [24,34]. While the

reason for applying this mixed, not entirely synthetic reverse genetics approach was not

explicitly explained in these publications, it is likely due to the toxicity of ISA DNA fragments

preventing commercial generation of fragments for fully synthetic flaviviruses.

In our studies, we encountered similar technical difficulties with the toxicity of flavivirus ISA

DNA fragments, which impeded commercial suppliers' ability to provide the required fragments.

Given the importance of generating entirely synthetic flavivirus stocks for certain experimental

objectives, in this study we applied and validated modified ISA strategies. These approaches

allowed us to generate JEV and WNV using a completely synthetic method for toxic ISA

flavivirus fragments.

MATERIALS AND METHODS

Wild-type and CCCH-type zinc finger antiviral protein (ZAP) knockout cells, *in silico*

permutation of the viral genome, ISA, DENV, JEV, and WNV stock generation and titration,

comparative ISA, mouse experiments, RT-qPCR, and statistics are described in supplemental

materials and methods.

RESULTS

Classical Infectious-Subgenomic-Amplicons method for dengue virus

As described before [19], transfection of three overlapping ISA DNA fragments generated via

synthetic clonal genes with medium copy number bacterial plasmids (**Table S1**) and

amplification in bacteria (**Fig 1**) consistently rescued infectious DENV in all five well replicates.

The virus induced a cytopathic effect (CPE) in all transfected wells at passages 0 and 1 (**Fig 2A**),

as well as in flasks during subsequent passaging and virus stock generation. Sanger sequencing

confirmed correct sequences in the DENV stock. The DENV stock produced in VERO-ZAP-KO

cells exhibited high infectious titers when titrated on both VERO-ZAP-WT and VERO-ZAP-KO

cells (**Fig 2B**), along with high viral RNA loads (**Fig 2C**) and E protein expression (**Fig 2D**).

Altogether, as in the previous study [19], we confirmed that classical ISA (**Fig 1**) with three

overlapping DNA fragments generated via *synthetic clonal genes* with bacterial plasmids is a

highly efficient and robust technique for type 2 DENV reverse genetics, enabling the rescue of

synthetic recombinant viruses. The properties of ISA-derived DENV have been extensively

characterized in prior research [19]; thus, we further focused on the more challenging ISA

strategies for JEV and WNV.

Infectious-Subgenomic-Amplicons method to overcome bacterial toxicity caused by

Japanese encephalitis virus sequences

In contrast to DENV, classical ISA failed to rescue the JEV SA14-14-2 wild-type strain (JEV-WT). The JEV Fragment I exhibited bacterial toxicity across three different medium and low copy bacterial plasmids (**Table S1, Fig 3**). To overcome this, we divided the toxic Fragment I sequence (4,550 bp) into two sub-fragments, Fragment-I-A and Fragment-I-B, each 2,315-2,316 bp. These sub-fragments were produced via *synthetic clonal genes* using medium copy number bacterial plasmids and amplified in bacteria (**Table S1, Fig 3, File S1**). Transfection of four overlapping ISA DNA fragments generated via *synthetic clonal genes* (**Fig 3**) consistently rescued infectious JEV-WT in all five replicates. The virus induced CPE in all transfected wells at passages 0 and 1 (**Fig 4A**), as well as in flasks during subsequent passaging and virus stock generation. Sanger sequencing confirmed correct sequences in the JEV-WT stock. The JEV-WT stock produced in VERO-ZAP-KO cells exhibited high infectious titers when titrated on both VERO-ZAP-WT and VERO-ZAP-KO cells (**Fig 4B**), along with high viral RNA loads (**Fig 4B**) and E protein expression (**Fig 4C**).

Enrichment of CpG dinucleotides in viral genomes is an emerging approach for live attenuated vaccines and oncolytics [26-29,39]. We generated CpG-enriched JEV (JEV-CpG) (**Table S1**, **Table S2**, **File S2**) using the same ISA strategy as for JEV-WT (**Fig 3**). The Fragment I sequence was subdivided into two sub-fragments, Fragment-I-A and Fragment-I-B, produced using *synthetic clonal genes* in medium copy number bacterial plasmids. Transfection of four overlapping ISA DNA fragments successfully rescued infectious JEV-CpG in all five replicates,

with the virus inducing CPE in all transfected wells at passages 0 and 1 (**Fig 4D**). Sanger

sequencing confirmed the correct sequences in the JEV-CpG stock. The JEV-CpG stock

produced in VERO-ZAP-KO cells showed high infectious titers on both VERO-ZAP-WT and

VERO-ZAP-KO cells (**Fig 4E**), along with high viral RNA loads (**Fig 4E**) and E protein

expression (**Fig 4G**). Consistent with previous enrichment studies in Zika virus [27,28], the

CpG-enriched JEV variant exhibited an attenuated phenotype in VERO-ZAP-WT cells (**Fig 4F**).

Next, we conducted a mouse study to identify whether JEV-WT and JEV-CpG obtained with the

synthetic ISA approach causes typical infection phenotypes *in vivo*. The SA14-14-2 JEV strain

does not cause infection in conventional mice; thus, we used *Ifnar1*[-/-] mice deficient for the type I

IFN receptor function. The wild SA14-14-2 JEV strain causes clinical signs and lethality in

Ifnar1[-/-] mice [40]. Five-week-old mice, six mice per group, were injected subcutaneously with

virus-free media (MOCK) or ISA-derived JEV-WT or JEV-CpG. Mice in the MOCK group did

not show clinical signs (**Fig 4H**); one MOCK mouse lost weight after manipulation and injection

but recovered. Mice in the JEV-WT group showed a high 50% mortality due to paralysis and loss

of 20% baseline body weight (**Fig 4H**). Another 50% of mice in the JEV-WT group also showed

clinical signs—ataxia, depression and hunching but survived. Consistent with previous

enrichment studies in Zika virus [27,28], the JEV-CpG variant showed an attenuated phenotype

with no mortality or disease in mice (**Fig 4H**).

We tested spleen and brain from MOCK and JEV groups with the virus-specific RT-qPCR assay.

As expected, JEV RNA was not detected in the MOCK group (**Fig 4I**). All samples from the

JEV-WT group showed high viral RNA loads—$10^{5.58-7.52}$ RNA copies per g of the brain, and

$10^{5.44-7.60}$ RNA copies per g of the spleen (**Fig 4I**). In contrast, all brain samples from the JEV-

CpG group were negative in RT-qPCR (**Fig 4I**). JEV-CpG RNA loads in the spleen were

considerably lower ($P < 0.0001$, t-test) than JEV-WT loads (**Fig 4I**).

Infection phenotypes in our study with the ISA-derived JEV-WT were comparable to the

infection phenotypes in the previous study where *Ifnar1*$^{-/-}$ mice injected with the wild (not

synthetic) SA14-14-2 JEV strain also showed mortality, clinical signs, and virus in the spleen

[40]. In contrast to our study however, the authors did not find wild SA14-14-2 JEV in the brain.

The discrepancy can be explained by the different sampling time—at 12 days after injection in

our study, and 4 days in the previous study [40]. Also, in the previous study, mice injected with

$10^{6.7}$ plaque forming units (PFU) showed 100% mortality, while mice in our study

(10^7 TCID$_{50}$/mouse) showed 50% mortality and 100% morbidity. The difference may be due to

different methods to quantify viral loads for injection (PFU versus TCID$_{50}$ assay), the route of

injection—intraperitoneal previously and subcutaneous in our study, and potentially reduced

viral population complexity in synthetically rescued JEV compared to wild viruses.

Altogether, dividing the toxic JEV fragment into two sub-fragments provides a highly efficient

and robust ISA approach for reverse genetics. This method overcomes the bacterial toxicity

caused by JEV sequences and enables the successful rescue of synthetic recombinant JEV.

Infectious-Subgenomic-Amplicons methods to overcome bacterial toxicity caused by West Nile virus sequences

During the rescue of the wild-type WNV NY99 strain (WNV-WT), Fragment I exhibited bacterial toxicity when using a medium copy bacterial plasmid (**Fig 5**). To address this, a low copy plasmid was utilized (**Fig 5**), which enabled the successful rescue of Fragment I after several attempts over a three-month period.

As observed with JEV (**Fig 3**), classical ISA failed to rescue the WNV NY99 strain which was permutated (WNV-Per) to generate the permutated control virus (see materials and methods; **File S2**, **Table S2**) across three different medium and low copy bacterial plasmids (**Table S1, Fig 6**). To overcome this issue, the toxic Fragment I sequence (4,358 bp) was divided into three sub-fragments: Fragment-I-A (1,198 bp), Fragment-I-B (1,126 bp), and Fragment-I-c (2,179 bp) (**Table S1, File S1**). The first two fragments were made shorter to facilitate bacteria-free *synthesis of DNA* (available only for fragments shorter than 1.8 kb) based on the assumption that the 5' flanking CMV promoter sequence contributed to the toxicity in bacteria. The longer Fragment-I-c was used for *synthetic clonal genes* with medium copy plasmid and bacterial amplification (**Table S1, File S1**). However, due to continued toxicity and failure, Fragment-I-c was further subdivided into overlapping Fragment-I-C (1,162 bp) and Fragment-I-D (1,089 bp), both small enough for bacteria-free *synthesis of DNA*. This troubleshooting led to the six overlapping ISA fragments for WNV-Per (**Fig 6, Table S1, File S1**).

Transfection of the three overlapping ISA DNA fragments for WNV-WT and six overlapping

ISA DNA fragments for WNV-Per consistently rescued the corresponding infectious WNV

variants in all five replicates. Sanger sequencing confirmed correct sequences in WNV stocks.

Viruses induced CPE in all transfected wells at passages 0 and 1 (**Fig 7A, B**), as well as in flasks

during subsequent passaging and virus stock generation. Both WNV-WT and WNV-Per stocks

produced in VERO-ZAP-KO cells had high infectious titers when titrated on both VERO-ZAP-

WT and VERO-ZAP-KO cells (**Fig 7C, D**), along with high viral RNA loads (**Fig 7C, D**) and E

protein expression (**Fig 7E, F**).

Next, we conducted a mouse study to identify whether WNV-WT and WNV-Per variants

obtained with the synthetic ISA approach cause typical infection phenotypes *in vivo*. We used a

well-established C57BL/6J mouse model for severe WNV infection [41-43]. Six-week-old

C57BL/6J mice (14 mice per group) were MOCK infected or injected intraperitoneally (IP) with

WNV-WT or WNV-Per variants. Mice in the MOCK group did not show clinical signs, weight

loss, or mortality (**Fig 7G**). Mice in the WNV-WT and WNV-Per groups showed high

mortality—93% and 79% (**Fig 7G**), ataxia, depression, paralysis, loss of 20% body weight, and

death.

We tested spleen and brain from MOCK and WNV groups with the virus-specific RT-qPCR

assay. As expected, WNV RNA was not detected in the MOCK group (**Fig 7H**). In contrast, all

samples from the WNV-WT and WNV-Per groups showed high WNV RNA loads—$10^{4.98-6.97}$

RNA copies per g of the spleen, and $10^{5.78-8.72}$ RNA copies per g of the brain (**Fig 7H**). The

difference in viral RNA loads between WNV-WT and WNV-Per groups was not statistically

significant ($P < 0.05$), which is expected because permutation in the WNV-Per variant does not

affect encoded proteins and does not considerably affect genome coding parameters (**Table S2**).

Infection phenotypes in our study with the ISA-derived WNV variants were comparable to the

infection phenotypes in the previous study where C57BL/6J mice injected with the wild (not

synthetic) WNV NY99 strain also showed mortality, clinical signs, and virus in the spleen and

brain [41-43].

Comparative Infectious-Subgenomic-Amplicons in ZAP wild-type and ZAP knockout cells

The zinc finger antiviral protein 1 (ZAP) exhibits broad-spectrum antiviral activity. Recent

studies, including our own, have demonstrated that flaviviruses such as Zika virus, JEV, and

WNV are susceptible to ZAP antiviral effects in VERO cells [44-46]. Here, we compared the

efficiency of JEV and WNV ISA in wild-type (VERO-ZAP-WT) and ZAP knockout (VERO-

ZAP-KO) cells. We selected JEV-WT, JEV-CpG, and WNV-Per variants, using four and six ISA

fragments (**Figs 3, 6**) for comparative ISA. Infectious-Subgenomic-Amplicons for each variant

were performed concurrently, as described in the supplemental materials and methods, with DNA

amplicons for transfection into VERO-ZAP-WT or VERO-ZAP-KO cells derived from the same

PCR reactions.

JEV-WT induced CPE in all five replicates of both ZAP-WT and ZAP-KO cells during passage

0, with more pronounced CPE in ZAP-KO cells (**Fig 8A**). JEV-CpG also induced CPE in all five

replicates of both ZAP-WT and ZAP-KO cells during passage 0, with more pronounced CPE in

ZAP-KO cells (**Fig 8A**). WNV-Per caused equally strong CPE in both ZAP-WT and ZAP-KO

cells (**Fig 8A**). During passage 1, JEV-WT, JEV-CpG, and WNV-Per infections in ZAP-KO cells

were associated with higher E protein expression compared to ZAP-WT cells (***$P < 0.0001$;

Figs 8B, D, F). Consequently, JEV-CpG and WNV-Per $TCID_{50}$ infectious titers in supernatants

were significantly higher in ZAP-KO cells than in ZAP-WT cells ($P = 0.014$-0.017) (**Figs 8C, E,**

J).

Overall, both wild-type and ZAP knockout cells demonstrated 100% ISA efficiency, successfully

rescuing JEV and WNV in all five replicates. However, ISA in ZAP-KO cells resulted in the

considerably higher number of virus-positive cells and elevated infectious virus loads.

DISCUSSION

The goal of this study was to validate modified ISA strategies for producing toxic ISA flavivirus

DNA fragments synthetically, overcoming bacterial toxicity and eliminating the need to amplify

toxic fragments from biological specimens or other preexisting sources. These ISA strategies

enable the generation of entirely synthetic JEV and WNV stocks. The strategies involved: (i)

Subdividing the problematic toxic DNA fragment into two sub-fragments for subsequent

synthetic clonal gene production with bacterial plasmids and bacteria. (ii) Using a low copy

bacterial plasmid. (iii) Subdividing the toxic DNA fragment into four short overlapping sub-

fragments, each up to 1.8 kb, for subsequent bacteria-free *synthesis of DNA* production. To our

knowledge, the latter strategy had never been applied for ISA.

As expected, and previously demonstrated for the DENV 2 serotype [19], the classical ISA

approach of subdividing the flavivirus genome into three overlapping fragments of

approximately 4 kb each works well for DENV. These fragments were synthesized quickly and

without technical difficulties by a commercial supplier using a medium copy bacterial plasmid

(**Table S1**).

In contrast, this classical three-fragment ISA approach failed when we attempted to rescue JEV-

WT Fragment I using three different medium and low copy bacterial plasmids (**Table S1**). The

solution was to divide the toxic fragment into two sub-fragments. We encountered similar

technical difficulties producing WNV-WT Fragment I. Here, replacing the medium copy plasmid

with a low copy plasmid resolved the issue, allowing the rescue of the entirely synthetic WNV-

WT variant, but this took several attempts and over three months.

More challenging difficulties arose when we attempted to rescue the permuted WNV-Per variant.

Similar to JEV ISA, we failed to produce Fragment I using three different medium and low copy

bacterial plasmids. Subdividing the toxic Fragment I into three sub-fragments (A, B, and c) did

not work because the sub-fragment Fragment I-c was also toxic to bacteria (**Table S1**).

Subsequently, we divided the highly toxic WNV-Per Fragment I into four sub-fragments, each

smaller than 1.8 kb. This allowed the *synthesis of DNA* commercial production process that is

bacteria-free (**Table S1**).

While various approaches have been employed to overcome technical difficulties in constructing

cDNA molecules representing the full-length flavivirus genomes in bacterial vectors, this process

remains challenging, laborious, and time-consuming. As demonstrated by our difficulties during

the rescue of synthetic JEV and WNV (**Table S1**), these challenges are also present in the

classical ISA approach. We propose that subdividing toxic fragments into sub-fragments smaller

than 1.8 kb suitable for the *synthesis of DNA* process is currently the most rapid and robust

strategy to overcome the need for bacterial plasmids and bacteria and ensure the fast rescue of

entirely synthetic flaviviruses.

This strategy significantly shortens the time required to produce recombinant flaviviruses. For

example, the *synthesis of DNA* process by a commercial supplier (**Table S1**) has turnaround time

2-4 days. Additionally, the synthesis of DNA is cost-effective with current 0.07 USD per bp; for

example, 4.3 kb DNA fragment for WNV ISA ordered via *synthetic clonal genes* process with

bacterial plasmids and amplification in bacteria costs around 840 USD with 3-4 weeks

turnaround time, while the same fragment divided into four sub-fragments and ordered via

bacteria-free *synthesis of DNA* costs 360 USD with 2-4 days turnaround time. During production,

the provider flanks DNA fragments with short 22 bp adapters (**File S1**), facilitating efficient PCR

amplification of ordered ISA fragments from the provided construct. The provider also applies

NGS to verify correct sequences, reducing both time and costs for virus rescue. We anticipate

that ongoing improvements in DNA synthesis technology and associated cost reductions will

provide even faster and more cost-effective bacteria-free fragments for ISA, enabling the even

more rapid production of entirely synthetic flaviviruses.

The zinc finger CCCH-type antiviral protein 1, also known as ZAP, ZC3HAV1, or PARP13, is a

cellular protein with broad antiviral activity. ZAP binds viral RNA and evokes antiviral activity

by mediating viral RNA degradation and translational inhibition [47]. It also interacts with

various cellular proteins that may act as co-factors, enhancing its antiviral effects [48].

Furthermore, ZAP can augment other antiviral systems [44,49]. Recent studies, including our

own, have demonstrated that flaviviruses such as Zika virus, JEV, and WNV are susceptible to

ZAP antiviral effects in VERO cells [44-46]. Thus, we also compared the efficacy of JEV-WT.

JEV-CpG, and WNV-Per ISA in ZAP-WT and ZAP-KO cells. Both cell types showed 100%

efficiency, with viruses rescued in all five replicates. However, the increased number of virus-

positive cells and infectious titers in ZAP-KO cells suggests that these cells may be more

effective for rescuing modified flaviviruses, such as attenuated variants for live attenuated

vaccines, including CpG-enriched vaccines. This is significant because CpG dinucleotide

enrichment is a novel vaccine approach. Increasing CpG dinucleotides in viral RNA, while

preserving the original amino acid composition, may weaken the virus, preventing disease but

potentially inducing strong antiviral immunity. However, CpG-enriched vaccine candidates are

attenuated both *in vitro* and *in vivo*, which can complicate their rescue and the generation of

high-titer vaccine stocks [27,28]. Using VERO-ZAP-KO cells could aid in developing CpG-

enriched vaccine candidates, and it is advantageous that VERO cells are already used as a GMP

line for producing many vaccines.

In conclusion, the ISA method is a robust and straightforward approach to reverse genetics for

JEV and WNV. In this study, we validated a modified ISA strategy that involves subdividing the

toxic DNA fragment into four short overlapping sub-fragments, each up to 1.8 kb, for subsequent

bacteria-free *DNA synthesis* production. To our knowledge, this strategy had never been applied

to ISA before, enabling the rapid generation of flaviviruses using a completely synthetic method

for toxic ISA flavivirus fragments. The strategy can be particularly beneficial for research on

emerging flaviviruses where a rapid response is needed to rescue new virus variants during

outbreaks.

ETHICS APPROVAL STATEMENT

The animal studies were approved by The Ohio State University Institutional Biosafety

Committee (e-Protocol # 2022R00000111) and The Institutional Animal Care and Use

Committee (e-Protocol #2023A00000095).

DISCLOSURE STATEMENT

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

Conceptualization: PPS, NPKL, UK. Investigation: PPS, NPKL, UK. Data analysis: PPS, NPKL,

UK. Funding: UK. Writing—original draft preparation: PPS, NPKL, UK. Writing—review and

editing: PPS, NPKL, UK.

DATA AVAILABILITY

All data are available in the manuscript or in the supplemental data.

FUNDING

REFERENCES

1. Pierson TC, Diamond MS. The continued threat of emerging flaviviruses. Nat Microbiol. 2020 Jun;5(6):796-812.
2. Blackburn NK, Swanepoel R. An investigation of flavivirus infections of cattle in Zimbabwe Rhodesia with particular reference to Wesselsbron virus. J Hyg (Lond). 1980 Aug;85(1):1-33.
3. Bhatt S, Gething PW, Brady OJ, et al. The global distribution and burden of dengue. Nature. 2013 Apr 25;496(7446):504-7.
4. Gubler DJ. Dengue/dengue haemorrhagic fever: history and current status. Novartis Found Symp. 2006;277:3-16; discussion 16-22, 71-3, 251-3.
5. Amicizia D, Zangrillo F, Lai PL, et al. Overview of Japanese encephalitis disease and its prevention. Focus on IC51 vaccine (IXIARO®). Journal of Preventive Medicine and Hygiene2018. p. E99-E107.
6. Tarantola A, Goutard F, Newton P, et al. Estimating the burden of Japanese encephalitis virus and other encephalitides in countries of the mekong region. In: Gray DJ, editor. PLoS neglected tropical diseases2014. p. e2533.
7. Chaturvedi UC, Mathur A, Chandra A, et al. Transplacental Infection with Japanese Encephalitis Virus. Journal of Infectious Diseases: Oxford University Press; 1980. p. 712-715.
8. Lindquist L. Recent and historical trends in the epidemiology of Japanese encephalitis and its implication for risk assessment in travellers. Journal of Travel Medicine2018. p. S3-S9.
9. Oliveira ARS, Piaggio J, Cohnstaedt LW, et al. Introduction of the Japanese encephalitis virus (JEV) in the United States – A qualitative risk assessment. Transboundary and Emerging Diseases2019. p. 1558-1574.
10. Huang YJS, Harbin JN, Hettenbach SM, et al. Susceptibility of a North American Culex quinquefasciatus to Japanese Encephalitis Virus. Vector borne and zoonotic diseases (Larchmont, N.Y.): Vector Borne Zoonotic Dis; 2015. p. 709-711.
11. Faizah AN, Kobayashi D, Matsumura R, et al. Blood meal source identification and RNA virome determination in Japanese encephalitis virus vectors collected in Ishikawa Prefecture, Japan, show distinct avian/mammalian host preference. J Med Entomol. 2023 Apr 7.
12. Shimizu T, Kawakami Y, Fukuhara S, et al. Experimental stillbirth in pregnant swine infected with Japanese encephalitis virus. The Japanese journal of experimental medicine1954. p. 363-375.
13. Howard-Jones AR, Pham D, Jeoffreys N, et al. Emerging Genotype IV Japanese Encephalitis Virus Outbreak in New South Wales, Australia. Viruses. 2022 Aug 24;14(9).
14. Barrett ADT. West Nile in Europe: an increasing public health problem. J Travel Med. 2018 Jan 1;25(1).
15. Petersen LR, Carson PJ, Biggerstaff BJ, et al. Estimated cumulative incidence of West Nile virus infection in US adults, 1999-2010. Epidemiol Infect. 2013 Mar;141(3):591-5.
16. Neumann G, Kawaoka Y. Reverse genetics systems for the generation of segmented negative-sense RNA viruses entirely from cloned cDNA. Curr Top Microbiol Immunol. 2004;283:43-60.
17. Aubry F, Nougairede A, Gould EA, et al. Flavivirus reverse genetic systems, construction techniques and applications: a historical perspective. Antiviral Res. 2015 Feb;114:67-85.
18. Ruggli N, Rice CM. Functional cDNA clones of the Flaviviridae: strategies and applications. Adv Virus Res. 1999;53:183-207.
19. Park CJ, Lee YA, Yoo SM, et al. Efficient reverse genetics approach involving infectious subgenomic amplicon for engineering dengue virus. J Med Virol. 2023 Aug;95(8):e28978.
20. Gritsun TS, Gould EA. Infectious transcripts of tick-borne encephalitis virus, generated in days by RT-PCR. Virology. 1995 Dec 20;214(2):611-8.
21. Rice CM, Grakoui A, Galler R, et al. Transcription of infectious yellow fever RNA from full-length cDNA templates produced by in vitro ligation. New Biol. 1989 Dec;1(3):285-96.

22. Kapoor M, Zhang L, Mohan PM, et al. Synthesis and characterization of an infectious dengue
virus type-2 RNA genome (New Guinea C strain). Gene. 1995 Sep 11;162(2):175-80.
23. Sumiyoshi H, Hoke CH, Trent DW. Infectious Japanese encephalitis virus RNA can be synthesized
from in vitro-ligated cDNA templates. J Virol. 1992 Sep;66(9):5425-31.
24. Aubry F, Nougairède A, de Fabritus L, et al. Single-stranded positive-sense RNA viruses generated
in days using infectious subgenomic amplicons. The Journal of general virology2014. p. 2462-7.
25. Cochin M, Driouich JS, Moureau G, et al. In vivo rescue of arboviruses directly from subgenomic
DNA fragments. Emerg Microbes Infect. 2024 Dec;13(1):2356140.
26. Trus I, Berube N, Jiang P, et al. Zika Virus with Increased CpG Dinucleotide Frequencies Shows
Oncolytic Activity in Glioblastoma Stem Cells. Viruses: Multidisciplinary Digital Publishing
Institute; 2020. p. 579.
27. Trus I, Udenze D, Berube N, et al. CpG-Recoding in Zika Virus Genome Causes Host-Age-
Dependent Attenuation of Infection With Protection Against Lethal Heterologous Challenge in
Mice. Front Immunol. 2019;10:3077.
28. Trus I, Udenze D, Karniychuk U. Generation of CpG-Recoded Zika Virus Vaccine Candidates.
Methods in molecular biology (Clifton, N.J.): Methods Mol Biol; 2022. p. 289-302.
29. Udenze D, Trus I, Berube N, et al. CpG content in the Zika virus genome affects infection
phenotypes in the adult brain and fetal lymph nodes. Front Immunol. 2022;13:943481.
30. Hu F, Zhu T, Guo X, et al. Generation of duck Tembusu virus using a simple reverse genetic system
in duck embryo fibroblast cells. J Virol Methods. 2022 Feb;300:114385.
31. Hoornweg TE, Godeke GJ, Hoogerwerf MN, et al. Rescue and in vitro characterization of a
divergent TBEV-Eu strain from the Netherlands. Sci Rep. 2023 Feb 18;13(1):2872.
32. Atieh T, Nougairède A, Klitting R, et al. New reverse genetics and transfection methods to rescue
arboviruses in mosquito cells. Scientific Reports: Nature Publishing Group; 2017. p. 13983.
33. Driouich JS, Moureau G, de Lamballerie X, et al. Reverse Genetics of RNA Viruses: ISA-Based
Approach to Control Viral Population Diversity without Modifying Virus Phenotype. Viruses. 2019
Jul 20;11(7).
34. Atieh T, Baronti C, de Lamballerie X, et al. Simple reverse genetics systems for Asian and African
Zika viruses. Scientific reports: Nature Publishing Group; 2016. p. 39384.
35. Touret F, Gilles M, Klitting R, et al. Live Zika virus chimeric vaccine candidate based on a yellow
fever 17-D attenuated backbone. Emerg Microbes Infect. 2018 Sep 26;7(1):161.
36. Klitting R, Roth L, Rey FA, et al. Molecular determinants of Yellow Fever Virus pathogenicity in
Syrian Golden Hamsters: one mutation away from virulence. Emerg Microbes Infect. 2018 Mar
29;7(1):51.
37. Fiacre L, Nougairede A, Migne C, et al. Different viral genes modulate virulence in model
mammal hosts and Culex pipiens vector competence in Mediterranean basin lineage 1 West Nile
virus strains. Front Microbiol. 2023;14:1324069.
38. Aubry F, Nougairède A, de Fabritus L, et al. "ISA-Lation" of Single-Stranded Positive-Sense RNA
Viruses from Non-Infectious Clinical/Animal Samples. In: Ng LFP, editor. PLOS ONE: Public Library
of Science; 2015. p. e0138703.
39. Takata MA, Goncalves-Carneiro D, Zang TM, et al. CG dinucleotide suppression enables antiviral
defence targeting non-self RNA. Nature. 2017 Oct 5;550(7674):124-127.
40. Liu J, Jing W, Fang Y, et al. The Infection of the Japanese Encephalitis Virus SA14-14-2 Strain
Induces Lethal Peripheral Inflammatory Responses in IFNAR Deficiency Mice. Frontiers in
microbiology: Front Microbiol; 2022.
41. Zhang YN, Li N, Zhang QY, et al. Rational design of West Nile virus vaccine through large
replacement of 3' UTR with internal poly(A). EMBO Mol Med. 2021 Sep 7;13(9):e14108.

42.   Daffis S, Samuel MA, Keller BC, et al. Cell-specific IRF-3 responses protect against West Nile virus
524   infection by interferon-dependent and -independent mechanisms. PLoS Pathog. 2007 Jul
525   27;3(7):e106.
526   43.   Daffis S, Samuel MA, Suthar MS, et al. Interferon regulatory factor IRF-7 induces the antiviral
527   alpha interferon response and protects against lethal West Nile virus infection. J Virol. 2008
528   Sep;82(17):8465-75.
529   44.   Nguyen Phuong Khanh Le PPS, Ahmad Jawad Sabir, Ivan Trus, Uladzimir Karniychuk. Endogenous
530   ZAP is associated with altered global cellular gene expression during Zika virus infection. bioRxiv.
531   2024.
532   45.   Chiu HP, Chiu H, Yang CF, et al. Inhibition of Japanese encephalitis virus infection by the host zinc-
533   finger antiviral protein. PLoS Pathogens: Public Library of Science; 2018.
534   46.   Sabir AJ, Le NPK, Singh PP, et al. Endogenous ZAP affects Zika virus RNA interactome. RNA Biol.
535   2024 Jan;21(1):1-10.
536   47.   Li MMH, Aguilar EG, Michailidis E, et al. Characterization of Novel Splice Variants of Zinc Finger
537   Antiviral Protein (ZAP). J Virol. 2019 Sep 15;93(18).
538   48.   Goodier JL, Pereira GC, Cheung LE, et al. The Broad-Spectrum Antiviral Protein ZAP Restricts
539   Human Retrotransposition. PLOS Genetics: Public Library of Science; 2015. p. e1005252.
540   49.   Nguyen LP, Aldana KS, Yang E, et al. Alphavirus Evasion of Zinc Finger Antiviral Protein (ZAP)
541   Correlates with CpG Suppression in a Specific Viral nsP2 Gene Sequence. Viruses. 2023 Mar
542   24;15(4).

FIGURE LEGENDS

Figure 1. Classical ISA method for dengue virus.

Figure 2. Dengue virus ISA and stock generation. (**A**) Cytopathic effect in DENV ISA-transfected ZAP-KO BHK+VERO cells, passage 0 day 7 after transfection. All five replicate wells showed CPE at passages 0 and 1; the representative images are shown. (**B**) Infectious $TCID_{50}$ titers of the DENV stock produced in ZAP-KO cells. The stock was titrated in VERO-ZAP-WT or VERO-ZAP-KO cells. (**C**) DENV RNA loads of the stock determined by virus-specific RT-qPCR. (**D**) Images of cells positive for the DENV E protein (red staining) at 7 days after infection with MOI 0.01. Magnification x200. The experiment was done in 4 replicates. Figures represent the general patterns in all replicates.

Figure 3. The adapted ISA method for Japanese encephalitis virus. To overcome toxicity, we divided the toxic Fragment I into two sub-fragments (**Table S1**, **File S1**).

Figure 4. JEV ISA and stock generation. Cytopathic effect in JEV-WT (**A**) and JEV-CpG (**D**) ISA-transfected ZAP-KO BHK+VERO cells, passage 0, day 6 after transfection. All five replicates showed CPE; the representative images are shown. Infectious $TCID_{50}$ titers of the JEV-WT (**B**) and JEV-CpG (**E**) stocks produced in ZAP-KO cells. The stock was titrated in VERO-ZAP-WT or VERO-ZAP-KO cells. JEV RNA loads of the stock determined by virus-specific RT-qPCR. Images of cells positive for the JEV-WT (**C**) and JEV-CpG (**G**) E protein (red staining) at 3 days after infection, MOI 0.01. Magnification x200. The experiment was done in 3 biological replicates. Figures represent the general staining patterns in replicates. (**F**) The fold increase of JEV-WT or JEV-CpG RNA loads in cell supernatants collected at 1h and 72h after inoculation. VERO-WT and VERO-ZAP-KO cells were inoculated with MOI 0.01. Virus inoculums were removed and replaced with media. Supernatants were collected at 1h (to normalize for leftover inoculum RNA) and at 72h. Three biological x 3 technical replicates. ***P=0.0017; unpaired t-test. (**H**) Mean body weight and survival in MOCK *Ifnar1*[-/-] mice and *Ifnar1*[-/-] mice injected with the ISA-derived JEV-WT or JWV-CpG. The following clinical scoring was used: 0—no visible abnormalities; 1—mild ataxia or tremors; 2—obvious ataxia or tremors; 3—depression, hunching, reluctance to walk, and falling to the side when walking; 4—close to moribund but still somewhat responsive; 5—paralysis; 6—loos of 20% baseline body weight; 7—found dead. Scores 4, 5, or 6 were used for the endpoint and mouse euthanasia. In body weight a standard error of the mean (SEM) is shown. (**I**) JEV loads in mouse spleen and brain sampled at euthanasia due to severe clinical signs or at the end of the study—at 12 days after injection.

Figure 5. The adapted ISA method for the wild-type WNV variant (WNV-WT). To overcome toxicity, we generated Fragment I in the low copy pCC1 bacterial plasmid (**Table S1**, **File S1**).

Figure 6. The adapted ISA method for the permuted WNV variant (WNV-Per). To overcome toxicity, we subdivide the toxic Fragment I into four sub-fragments for bacteria-free *synthesis of DNA* production (**Table S1**, **File S1**).

Figure 7. WNV ISA and stock generation. Cytopathic effect in WNV-WT (**A**) and WNV-Per (**B**) ISA-transfected ZAP-KO BHK+VERO cells, passage 0, day 6 after transfection. All five replicates for each WNV variant showed CPE; representative images are shown. Infectious $TCID_{50}$ titers and RT-qPCR RNA loads of the WNV-WT (**C**) and WNV-Per (**D**) stocks produced in ZAP-KO cells. For $TCID_{50}$ virus stocks were titrated in VERO-ZAP-WT or VERO-ZAP-KO cells. Images of cells positive for the WNV-WT (**E**) and WNV-Per (**F**) E protein (red staining) at 3 days after infection, MOI 0.01. Magnification x200. The experiment was done in 3 biological replicates. Figures represent the general patterns in all replicates. (**G**) Mean body weight and survival in MOCK C57BL/6J mice and C57BL/6J mice injected with the ISA-derived WNV-WT and WNV-Per. The following clinical scoring was used: 0—no visible abnormalities; 1—mild ataxia or tremors; 2—obvious ataxia or tremors; 3—depression, hunching, reluctance to walk, and falling to the side when walking; 4—close to moribund but still somewhat responsive; 5—paralysis; 6—loos of 20% baseline body weight; 7—found dead. Scores 4, 5, or 6 were used for the endpoint and mouse euthanasia. In body weight a standard error of the mean (SEM) is shown. (**H**) WNV loads in mouse spleen and brain sampled at euthanasia due to severe clinical signs or at the end of the study.

Figure 8. Comparative ISA in ZAP-WT and ZAP-KO cells. (**A**) Cytopathic effect in JEV-WT, JEV-CpG, and WNV-Per ISA-transfected ZAP-WT and ZAP-KO BHK+VERO cells, passage 0, day 5 (JEV-WT and JEV-CpG) and day 4 (WNV-Per) after transfection. All five replicates for JEV-WT, JEV-CpG, and WNV-Per showed CPE; the representative images are shown. The digital quantification of JEV-WT (**B**), JEV-CpG (**D**), and WNV-Per (**F**) E protein-positive cells in all 5-well replicates at day 4 (JEV) and day 2 (WNV) during passage 1 of comparative ISA. For technical replicates, three random images were obtained from each well. Examples of JEV-WT, JEV-CpG, and WNV-Per E protein IHC are in **Figures 4C, G,** and **7F**. ***P<0.0001, unpaired t-test. JEV-WT (**C**), JEV-CpG (**E**), and WNV-Per (**J**) infectious $TCID_{50}$ titers in supernatants collected at day 4 (JEV) and day 2 (WNV-Per) during passage 1 of comparative ISA. Data from 4-well replicates are shown. *P<0.014-0.017, unpaired t-test.

SUPPORTING INFORMATION CAPTIONS

Supplemental materials and methods.

Table S1 ISA fragments and primers.

Table S2 JEV and WNV genomic parameters.

File S1 ISA fragments.

File S2 JEV-CpG and WNV-Permuted reference sequences.

651
652
653

Fig. 1

Fig. 2

JEV-WT and JEV-CpG whole genome

Fig. 3

679
680
681

Fig. 4

Fig. 5

WNV-Permuted whole genome

Fig. 6

Fig. 7

714
715
716

Fig. 8

SUPPLEMENTAL MATERIALS AND METHODS

Infectious subgenomic amplicon strategies for Japanese encephalitis and West Nile viruses

Prince Pal Singh[1,2], Nguyen Phuong Khanh Le[1], Uladzimir Karniychuk[1,*]

[1]Department of Veterinary Biosciences, College of Veterinary Medicine, The Ohio State University, Columbus, OH, USA

[2]School of Public Health, University of Saskatchewan, Saskatoon, Canada

*Correspondence: Uladzimir Karniychuk

Department of Veterinary Biosciences, College of Veterinary Medicine, The Ohio State University, Columbus, OH, USA

karniychuk.1@osu.edu

SUPPLEMENTAL MATERIALS AND METHODS

Cells

We maintained wild-type VERO E6 (VERO-ZAP-WT) and BHK-21 (BHK-ZAP-WT) cells,

along with CCCH-type zinc finger antiviral protein (ZAP) knockout (ZAP-KO) derivatives

(VERO-ZAP-KO and BHK-ZAP-KO), in DMEM (Fisher, MA, USA; #11-965-118)

supplemented with 10% heat-inactivated fetal bovine serum (FBS, Fisher, MA, USA;

#A5256801), 1x Penicillin-Streptomycin (Fisher, MA, USA; #15140122), and 2.67 mM sodium

bicarbonate (Fisher, MA, USA; # 25080094) at +37°C in a 5% CO_2 humidified incubator.

To generate the VERO-ZAP-KO cell line, we previously used the guide RNA (gRNA)

GTCTCTGGCAGTACTTGCGA targeting the first exon of the *Chlorocebus sabaeus* (Gene ID:

103226990) ZC3HAV1 gene, necessary for expressing all ZAP isoforms. A non-targeting gRNA

(ACGGAGGCTAAGCGTCGCAA) served as the control [1,2]. These gRNAs were transiently

transfected into cells using GenCrispr NLS-Cas9-NLS Nuclease (GenScript). Transfected cells

were then seeded in 96-well plates through limiting dilution to generate isogenic single clones.

After expansion, the clones were genotyped via Sanger sequencing using primers

ATCGCTGGGCTGGACTAACG and GCAGAGAAGGGAGTGGCTGAA to identify location

of indels on *Chlorocebus sabaeus* genome: -2 bp deletion, NW_023666072.1: 4677550-

4677551. The knockout subclone confirmed by genotyping was further validated by western blot

for ZAP expression in our previous studies [1,2]. Negative control cells were not isolated into

single clones but were validated by Sanger sequencing to confirm the absence of indels.

The same strategy was used to generate BHK-ZAP-KO cells with gRNA

GGAGATCGTAGGCCAGACCGGGG targeting the first exon of the *Mesocricetus auratus*

(Gene ID: 110343172) Zc3hav1 gene and the same non-targeting control gRNA. After

expansion, the clones were genotyped via Sanger sequencing with primers

GCTGCTTCATCACCAAGATCC and GGTCGGGAAGACCCTTTCAC to identify location of

indels on *Mesocricetus auratus* genome: -29 bp deletion, NW_024429200.1: 3564916-3564944.

We confirmed the absence of mycoplasma contamination in all cells using the LookOut

Mycoplasma PCR Detection Kit (Millipore Sigma, MA, USA; #MP0035).

***In silico* CpG enrichment and permutation in viral genomes**

For CpG dinucleotide enrichment in the JEV genome, specifically in the region encoding the E

protein (**File S2**), we used the SSE 1.3 software package [3], as we previously described [4,5].

The genomic parameters of the wild-type and CpG-enriched JEV variants are listed in **Table S2**.

Certain experiments involving viral genomic mutations need permuted or scrambled virus

control variants. Permutation can potentially impact the efficiency of reverse genetics methods.

To simulate this scenario for ISA, we generated a permuted control for WNV. For permutation

of WNV sequences we selected genomic regions encoding E and NS1 (**File S2**) because these

proteins are important for flavivirus pathogenesis, modulation of the infection cycle, viral RNA

replication, and host immune evasion. Moreover, we previously permuted these genomic regions

in the Zika virus genome; this permutation did not affect Zika infection phenotypes *in vitro* and

in mice [4]. We used an SSE 1.3 software package [3], specifically the CDLR randomizing in the

SSE 1.3 [3,6-8]. During randomization we sought to avoid areas of the genome containing RNA

elements that are required for replication or translation of the virus genome, such as cis-

replicating elements, gene start or gene-end signals; we also avoided regions with prominent

secondary structures. The CDLR randomizes the order of codons within the sequence while

maintaining encoded proteins, dinucleotide frequencies, and coding parameters (**Table S2**).

Infectious-Subgenomic-Amplicons (ISA) and virus stock generation

The following reference sequences were used for *de novo* synthesis of flaviviruses: DENV type 2

strain 16681 [GenBank: U87411.1], JEV strain SA14-14-2 [GenBank: MK585066.1], and WNV

strain NY99 [GenBank: DQ211652.1]. To rescue DENV, JEV, and WNV, initially, we used

three overlapping (70-80 nt overlap) ISA DNA fragments (3,371-4,550 bp length) covering the

entire viral genome (**Table S1**; **File S1**); three overlapping DNA fragments is common ISA

strategy for flaviviruses [9,10]; thus, we consider it as a classical approach. The first and last

fragments for each virus are flanked with pCMV promoter and HDR/SV40pA sequences (**File**

S1).

Initially all DNA fragments for ISA were ordered through GeneScript (NJ, USA) or Twist

Biosciences (CA, USA) *synthetic clonal genes* services where DNA fragments are synthesized,

inserted into the bacterial plasmids (**Table S1**), amplified in bacteria, extracted, sequenced, and

plasmids with inserted fragments are delivered as lyophilized DNA with known concentration.

Some flavivirus ISA DNA fragments were toxic to bacteria, in both medium copy and low copy

plasmids (**Table S1**), and after multiple attempts *synthetic clonal genes* approach failed (see

results for details). To overcome this flavivirus DNA sequence toxicity to bacteria, we split toxic

clonal genes into smaller overlapping fragments below 1.8 kb length. These smaller size

fragments allowed *synthesis of DNA* services from Twist Biosciences were DNA fragments,

flanked with 22 bp adapters (**File S1**), are rapidly (2-4 days service) synthesized without using

bacterial plasmids and bacteria. Twist Biosciences and GeneScript confirmed the correct

sequences of DNA fragments by next-generation-sequencing (NGS).

Overlapping clonal genes or synthetic DNA ISA fragments were amplified with high fidelity

Invitrogen Platinum PCR SuperMix High Fidelity (Fisher, MA, USA; #12-532-016) (primers are

in **Table S1**), mixed in equimolar concentration to obtain the final 1µg of DNA for transfection,

and transfected into BHK-ZAP-KO cell monolayers (6-well plates, five transfection replicates,

one mock-transfection control well) with Lipofectamine 3000 (Fisher, MA, USA; #L3000015) in

640 µl of OptiMEM (Fisher, MA, USA; #11-058-021) for 12 h at +37°C, 5% CO_2. Afterward,

OptiMEM media was removed and replaced with 3 ml of DMEM with 10% of FBS containing

$4x10^5$ VERO-ZAP-KO cells on the top of BHK-ZAP-KO cells and incubated plates for

additional 6-7 days (passage 0). For passage 1, 400 µl of media from each well of passage 0 was

transferred to wells with VERO-ZAP-KO cells and incubated for 4-7 days. During passages 0

and 1, cells were monitored for cytopathic effect (CPE). Generated flaviviruses were additionally

passaged 2-3 times on VERO-ZAP-KO cells to generate virus stocks; flasks were monitored for

CPE. Cell culture media from the final stocks were centrifuged (12,000 g, 20 min, +4°C),

aliquoted, and frozen (-80°C). Viral titers were quantified in duplicate in VERO-ZAP-WT and

VERO-ZAP-KO cells with endpoint dilution assay described below. The absence of mycoplasma

contamination in all virus stocks and cell cultures was confirmed using PCR Detection Kit

(Millipore Sigma, MA, USA; #MP0035).

To validate virus stock sequences, RNA was extracted from virus stock culture supernatants as

described below. PCR was performed using SuperScript™ IV One-Step RT-PCR System (Fisher,

MA, USA; #12-594-025) as per manufacturer's instructions with the primers targeting sequences

encoding E protein: DENV (Forward 5- GACAGCTGTCACTCCTTCAAT GAC-3 and Reverse

5- CATGCTCTTCCCCTGAGTGAG G-3); JEV-WT (Forward 5-

GTTTTAATTGTCTGGGAATGGGC-3 and Reverse 5-GCCAACTTTGTATTTGATGTTTTC-

3); JEV-CpG (Forward 5-CAGTTTTAATTGTCTGGGAATGGG-3 and Reverse 5-

GTCAGCATGCACATTGGTCG-3); WNV-WT (Forward 5-

CTATTGCTTTTGGTGGCCCCAG-3 and Reverse 5-GTAGTTGGTCCATGGACAAA-3);

WNV-Per (Forward 5-CTATTGCTTTTGGTGGCCCCAG-3 and Reverse 5-

GTAGTTGGTCCATGGACAAA-3). PCR products were run on a 1% agarose gel, and DNA was

purified and concentrated using DNA Clean & Concentrator-5 (Zymo Research, CA, USA; #

D4013) as per manufacturer's instructions. DNA was sequenced by Sanger sequencing and

sequences were aligned with corresponding ISA sequences (**File S1**). Sanger sequencing

confirmed correct sequences in DENV, JEV, and WNV stocks.

Comparative ISA

To compare ISA efficiency on ZAP-WT and ZAP-KO cells, JEV and WNV fragments were

transfected into ZAP-WT or ZAP-KO cells as above. These experiments were conducted

simultaneously; also, the DNA amplicons for transfection in ZAP-WT or ZAP-KO cells were

obtained from the same PCR reactions. During passage 0, ZAP-WT and ZAP-KO cells were

observed for 5 (JEV variants) and 4 (WNV variants) days to compare CPE; the time was shorter

for WNV because it causes more aggressive infection and CPE. During the 1st passage, CPE was

observed, supernatants were collected on days 2-4 (2 for WNV variants; 4 for JEV variants) from

each well, and infectious virus titers were quantified with the end-point dilution assay as

described below. The shorter passage duration in the comparative ISA than in the initial ISA

used for virus rescue were applied to preserve cell monolayers for quantitative

immunohistochemistry (IHC) described below. The quantitative difference between ZAP-WT

and ZAP-KO conditions was represented by $TCID_{50}$ infectious titers in cell culture supernatants

from four replicates. Additionally, cellular monolayers from each transfected and mock-

transfected well (five replicates) were fixed and stained for viral E protein via IHC as described

in the infectious virus titration below. To quantify and compare IHC staining between ZAP-WT

and ZAP-KO conditions, three images (magnification x200) were randomly acquired in all five

well replicates with the same microscopic settings. Afterward, each image was processed in

ImageJ (NIH) (https://imagej.net/imaging/particle-analysis) with Automatic Particle counting

[2].

Mouse experiments

The animal studies were approved by The Ohio State University Institutional Biosafety

Committee (e-Protocol # 2022R00000111) and IACUC (e-Protocol #2023A00000095). Mice

were euthanized by deeply anesthetizing them with isoflurane, followed by cervical dislocation.

We conducted mouse experiments to identify whether ISA-rescued synthetic JEV and WNV

cause expected infection phenotypes which were previously described in mice infected with wild

or field isolates. The wild SA14-14-2 JEV strain does not cause clinical signs in conventional

mice; thus, we used *Ifnar1*$^{-/-}$ mice (The Jackson Laboratory, USA; Strain #:028288). The SA14-

14-2 JEV strain causes clinical signs or lethality in infected *Ifnar1*$^{-/-}$ mice [11]. After one week of

acclimatization, five-week-old mice (6 mice per group) were injected subcutaneously with the

virus-free media (MOCK) or ISA-derived JEV-WT or JEV-CpG variants (**Table S1**). We

standardized mouse injection doses by normalization of viral RNA loads (determined by RT-

qPCR, see below) in the JEV-WT and JEV-CpG inoculums which resulted in 10^{10} RNA

copies/mouse for both variants (JEV-WT – 10^7 TCID$_{50}$/mouse; JEV-CpG – $10^{6.3}$ TCID$_{50}$/mouse;

titers quantified on VERO-ZAP-WT cells, see below). Afterward, mice were monitored for

clinical signs and body weight. Mice were sacrificed when exhibiting greater than 20% weight

loss or severe clinical signs. We sampled spleen and brain from deceased mice, or from survived

mice at the end of the study to quantify JEV loads.

The wild NY99 WNV strain causes severe infection in conventional C57BL/6J mice; thus, we

used the well-characterized C57BL/6J mouse lethal model [12-14]. Five-week-old C57BL/6J

mice (Strain #:000664; 14 mice per group) were ordered from The Jackson Laboratory,

acclimatized for a week, and injected intraperitoneally (IP) with the virus-free media (MOCK),

ISA-derived WNV wild-type (WNV-WT), or ISA-derived WNV permuted (WNV-Per) variant

(**Table S1**). We standardized mouse injection doses by normalization of viral RNA loads

(determined by RT-qPCR, see below) in the WNV-WT and WNV-Per inoculums which resulted

in 10^7 RNA copies/mouse for both variants (WNV-WT - 10^5 TCID$_{50}$/mouse; WNV-Permuted -

$10^{5.6}$ TCID$_{50}$/mouse; titers quantified on VERO-ZAP-WT cells, see below). Afterward, mice

were monitored for clinical signs and body weight changes. Mice were sacrificed when

exhibiting 20% or greater weight loss or severe clinical signs, or at the end of the study to sample

spleen and brain from mice.

Infectious virus titration

To quantify infectious loads in DENV, JEV and WNV stocks, the endpoint dilution assay was used

to determine the 50% tissue culture infectious dose ($TCID_{50}$), with two replicates, as we had

described before [4,15-19]. Viral stocks were serially diluted in four replicates in media and 50 µl

of each dilution was added to confluent VERO-ZAP-WT or VERO-ZAP-KO cells cultured in 96-

well plates. Dilutions were made in DMEM media with 1% FBS. After 2 h at 37°C, 150 µl of

media was added. The cells were incubated for 5-7 days (5 days for WNV; 7 days for DENV and

JEV) at 37°C, 5% CO_2. Afterward, plates were washed, dried, formalin fixed, and stained with

specific antibodies (Ab) [4,15-19]. For DENV and WNV IHC, we used primary anti-pan flavivirus

envelop (E) protein monoclonal Ab D1-4G2-4-15 (ATCC, VA, USA; #HB-112) followed by Goat

anti-Mouse IgG HRP Ab (Abcam, MA, USA; #ab97023). For JEV, we used JEV E polyclonal Ab

(GeneTex, CA, USA; #GTX125867;) followed by Goat Anti-Rabbit IgG H&L (HRP) Ab (Abcam,

MA, USA; #ab97051) [4,15-19]. Infectious titers were calculated by the Spearman-Kärber formula

and expressed in $\log_{10}$ $TCID_{50}$ per ml. Media from mock-inoculated cells were used as negative

controls.

RNA extraction and virus-specific reverse transcriptase quantitative polymerase chain

reaction assays (RT-qPCR)

The RNA from cell culture supernatants was extracted with QIAamp Viral RNA Mini Kit

(QIAGEN) according to the manufacturer's instructions. The RNA from mouse tissues was

extracted with PureLink RNA Mini Kit (Invitrogen) according to the manufacturer's instructions.

All RT-qPCR reactions were conducted on the QuantStudio 3 real-time PCR system (Applied

Biosystems) and analyzed using QuantStudio Design & Analysis Software v1.5.2.

For DENV (F: 5′-AAGGACTAGAGGTTAKAGGAGACCC-3′; R: 5′-

GGCGYTCTGTGCCTGGAWTGATG-3′; Probe: 56-JOEN/AA CAG CAT A/ZEN/T TGA CGC

TGG GAA AGA CC-3IABkFQ), JEV-WT and JEV-CpG (F: 5′-GCCACCCAGGAGGTCCTT-

3′; R: 5′-CCCCAAAACCGCAGGAAT-3′; Probe: 56-FAM-CAAGAGGTG/ZEN/GACGGCC-

3IABkFQ), and WNV-WT and WNV-Per (F: 5′-AGTAGTTCGCCTGTGTGAGC-3′; R: 5′-

GCCCTCCTGGTTTCTTAGA-3′; Probe: FAM-

AATCCTCACAAACACTACTAAGTTTGTCA-TAMRA) we used the previously validated

probe-based one-step RT-qPCR assays [20-22]. The Luna Universal Probe One-Step RT-qPCR

Kit (NEB) reaction mixture (20 μL) consisted of 10 μL Luna Universal Probe One-Step Reaction

Mix, 1 μL Luna WarmStart RT Enzyme Mix, 1 μL (10 μM) of forward and reverse primers, 0.5

μL of the probe, 2.5 μL nuclease-free water and 4 μL of RNA template. The reverse transcription

and enzyme activation steps of 10 min at 55˚C and 1 min at 95˚C were followed by 40

amplification cycles (10 s at 95˚C and 30 s at 60˚C). A standard curve was used to quantify viral

RNA loads as we described [15].

PCR values were corrected for fluid volumes or tissue weights and upon logarithmical

transformation expressed as virus RNA genome copies per ml or g. Strict precautions were taken

to prevent PCR contamination. All master mix preparations were done in the dedicated PCR

cabinet. Aerosol-resistant filter pipette tips and disposable gloves were used. Kit reagent controls

were included in every RNA extraction and PCR run.

Statistics

We used GraphPad 9 PRISM for statistics. For the comparison of the viral RNA loads, infectious

titers, and quantitative IHC we used an unpaired t-test. *P*-value < 0.05 was considered statistically

significant.

REFERENCES

1. Nguyen Phuong Khanh Le, Prince Pal Singh, Ahmad Jawad Sabir, Ivan Trus, Uladzimir Karniychuk. Endogenous ZAP is associated with altered global cellular gene expression during Zika virus infection. bioRxiv. 2024.
2. Sabir AJ, Le NPK, Singh PP, et al. Endogenous ZAP affects Zika virus RNA interactome. RNA Biol. 2024 Jan;21(1):1-10.
3. Simmonds P. SSE: a nucleotide and amino acid sequence analysis platform. BMC research notes: BioMed Central; 2012. p. 50.
4. Trus I, Udenze D, Berube N, et al. CpG-Recoding in Zika Virus Genome Causes Host-Age-Dependent Attenuation of Infection With Protection Against Lethal Heterologous Challenge in Mice. Front Immunol. 2019;10:3077.
5. Trus I, Udenze D, Karniychuk U. Generation of CpG-Recoded Zika Virus Vaccine Candidates. Methods in molecular biology (Clifton, N.J.): Methods Mol Biol; 2022. p. 289-302.
6. Gaunt E, Wise HM, Zhang H, et al. Elevation of CpG frequencies in influenza A genome attenuates pathogenicity but enhances host response to infection. eLife2016. p. e12735.
7. Atkinson NJ, Witteveldt J, Evans DJ, et al. The influence of CpG and UpA dinucleotide frequencies on RNA virus replication and characterization of the innate cellular pathways underlying virus attenuation and enhanced replication. Nucleic acids research2014. p. 4527-45.
8. Tulloch F, Atkinson NJ, Evans DJ, et al. RNA virus attenuation by codon pair deoptimisation is an artefact of increases in CpG/UpA dinucleotide frequencies. eLife2014. p. 1-15.
9. Park CJ, Lee YA, Yoo SM, et al. Efficient reverse genetics approach involving infectious subgenomic amplicon for engineering dengue virus. J Med Virol. 2023 Aug;95(8):e28978.
10. Aubry F, Nougairede A, Gould EA, et al. Flavivirus reverse genetic systems, construction techniques and applications: a historical perspective. Antiviral Res. 2015 Feb;114:67-85.
11. Liu J, Jing W, Fang Y, et al. The Infection of the Japanese Encephalitis Virus SA14-14-2 Strain Induces Lethal Peripheral Inflammatory Responses in IFNAR Deficiency Mice. Frontiers in microbiology: Front Microbiol; 2022.
12. Zhang YN, Li N, Zhang QY, et al. Rational design of West Nile virus vaccine through large replacement of 3' UTR with internal poly(A). EMBO Mol Med. 2021 Sep 7;13(9):e14108.
13. Daffis S, Samuel MA, Keller BC, et al. Cell-specific IRF-3 responses protect against West Nile virus infection by interferon-dependent and -independent mechanisms. PLoS Pathog. 2007 Jul 27;3(7):e106.
14. Daffis S, Samuel MA, Suthar MS, et al. Interferon regulatory factor IRF-7 induces the antiviral alpha interferon response and protects against lethal West Nile virus infection. J Virol. 2008 Sep;82(17):8465-75.

15. Chapagain S, Pal Singh P, Le K, et al. Japanese encephalitis virus persists in the human
reproductive epithelium and porcine reproductive tissues. PLoS Negl Trop Dis. 2022
Jul;16(7):e0010656.
16. Sabir AJ, Singh PP, Trus I, et al. Asian Zika virus can acquire generic African-lineage mutations
during in utero infection. Emerg Microbes Infect. 2023 Dec;12(2):2263592.
17. Darbellay J, Lai K, Babiuk S, et al. Neonatal pigs are susceptible to experimental Zika virus
infection. Emerg Microbes Infect. 2017 Feb 15;6(2):e6.
18. Darbellay J, Cox B, Lai K, et al. Zika Virus Causes Persistent Infection in Porcine Conceptuses and
may Impair Health in Offspring. EBioMedicine. 25:73-86.2017.
19. Trus I, Udenze D, Cox B, et al. Subclinical in utero Zika virus infection is associated with interferon
alpha sequelae and sex-specific molecular brain pathology in asymptomatic porcine offspring. In:
Pierson TC, editor. PLoS Pathogens2019. p. e1008038.
20. Eiden M, Vina-Rodriguez A, Hoffmann B, et al. Two new real-time quantitative reverse
transcription polymerase chain reaction assays with unique target sites for the specific and
sensitive detection of lineages 1 and 2 West Nile virus strains. J Vet Diagn Invest. 2010
Sep;22(5):748-53.
21. Shao N, Li F, Nie K, et al. TaqMan Real-time RT-PCR Assay for Detecting and Differentiating
Japanese Encephalitis Virus. Biomedical and environmental sciences : BES: Biomed Environ Sci;
2018. p. 208-214.
22. Callahan JD, Wu SJ, Dion-Schultz A, et al. Development and evaluation of serotype- and group-
specific fluorogenic reverse transcriptase PCR (TaqMan) assays for dengue virus. J Clin Microbiol.
2001 Nov;39(11):4119-24.