Probing intracellular determinants of PARP inhibitor selectivity and pharmacology with CeTEAM

Maria J. Pires[1], Alen Lovric[2], Emanuele Fabbrizi[3], Dante Rotili[3], Mikael Altun[1,*], Nicholas C.K. Valerie[1,*]

[1]Science for Life Laboratory, Division of Clinical Physiology, Department of Laboratory Medicine, Karolinska Institutet, Karolinska University Hospital; Huddinge, SE-141 52, Sweden.
[2]Division of Clinical Physiology, Department of Laboratory Medicine, Karolinska Institutet, Karolinska University Hospital; Huddinge, SE-141 52, Sweden.
[3]Department of Chemistry and Technology of Drugs, Sapienza University of Rome; 00185 Roma RM, Italy.

*Corresponding author: nicholas.valerie@ki.se, mikael.altun@ki.se

Keywords

PARP inhibitors, CeTEAM, HPF1, cellular target engagement, drug profiling, pharmacology, biosensor, fluorescence, protein engineering, drug discovery

Abstract

PARP inhibitors (PARPi) predominantly targeting PARP1 and PARP2 have revolutionized cancer therapy by exploiting synthetic lethality and selectively killing cancer cells with defective DNA repair. However, achieving PARP1 or PARP2-selective inhibitors is difficult due to their close structural homology. Selectivity profiling is typically done with purified proteins, but these lack the complexity of intracellular environments and could therefore be inaccurate. The cellular target engagement by accumulation of mutant (CeTEAM) method provides insights into drug binding *in cellulo* by means of conditionally stabilized biosensors, thus offering a dynamic view of pharmacological events in living cells. Here, we duplex PARP1 L713F-GFP and PARP2 L269A-mCherry biosensors to systematically characterize potential PARPi binding and cell cycle alterations at the single cell level. Our results reveal that most PARPi are generally equipotent for both PARPs or have slight biases only towards PARP1, not PARP2. AZD5305, a reported PARP1-selective inhibitor, was the exception and appears ~1600-fold more potent towards PARP1. Surprisingly, niraparib was >10-fold more selective for PARP1, despite reported equipotent biochemical activity. Meanwhile, the next generation PARPi, senaparib, was a potent PARP1/2 binder and DNA trapper. We also assessed the effect of the PARP1/2 active site component, HPF1, on intracellular PARPi binding and see that HPF1 depletion elicits slight deviations in apparent binding potency, while contributing additively to PARP-DNA trapping. These results highlight that multiplexing CeTEAM biosensors and layered genetic perturbations can systematically profile determinants of intracellular drug selectivity. Furthermore, the PARP1/2 CeTEAM platform should facilitate the discovery of selective PARPi for better targeted therapies.

Introduction

PARP inhibitors (PARPi) have revolutionized treatment for cancers characterized by deficiencies in DNA repair mechanisms, particularly those involving mutations to the homologous recombination repair proteins, BRCA1 and BRCA2 (1). These drugs preferentially block the ability of PARP1 and PARP2 to initiate repair of DNA damage and selectively kill cancer cells that are unable to repair double strand breaks (DSBs) based on the concept of synthetic lethality (2, 3). This effect is further driven by the ability of some PARPi to trap PARP1 and PARP2 onto DNA, thereby creating more irreparable DNA damage (1, 4)

Most PARPi indiscriminately target both PARP1 and PARP2 and, to a lesser extent, other PARP paralogs (5). PARP1 and PARP2 have substantially different N-terminal DNA binding domains but share homologous catalytic domains and allosteric regulatory mechanisms (6, 7). Therefore, specific targeting of their enzymatic activities has been challenging to date, as the structural variation in current PARPi conveys similar inhibition of both paralogs but very different allosteric outcomes (6, 8). Nonetheless, each has unique biological functions warranting preferential targeting (9), which could convey clinical ramifications but also create therapeutic opportunities. For example, the hematological toxicity of PARPi has been attributed to an on-target, adverse effect of PARP2 targeting due to its essential role in erythroid progenitor differentiation (10, 11). Thus, despite an abundance of potent, clinical-grade inhibitors, the need for truly selective, next generation PARPi remains high.

Small molecule selectivity is traditionally assessed with purified proteins *in vitro*, as more exact calculations of the inhibitory potency and characterizations of the binding modality can be performed. However, selectivity profiling in cellular contexts, as opposed to with purified proteins, has unveiled disparities in inhibitor performance, suggesting that the physiological, intracellular environment can significantly influence the behavior of inhibitors (12). For example, the interaction of PARP1 and PARP2 with co-factors like Histone PARylation Factor 1 (HPF1) can further complicate the pharmacodynamics of PARPi (13). HPF1 modulates the PARylation activity of PARP1 and PARP2 by effectively completing their enzymatic pockets, and, in turn, tuning their ability to interact with and modify histones during chromatin remodeling and DNA repair (13, 14). Thus, it is reasonable to expect HPF1 to influence PARPi binding and activity. Indeed, early indications suggest that HPF1 can modulate PARPi binding potency *in vitro* (15, 16) and PARPi sensitivity *in cellulo* (17), but the effect on PARPi binding in cells has not been elucidated.

Here, we utilize the recently developed cellular target engagement by accumulation of mutants (CeTEAM) assay (18) to comprehensively profile intracellular selectivity of PARPi towards PARP1 and PARP2. CeTEAM leverages stability-dependent biosensors that can be multiplexed with other readouts of downstream pharmacology to follow drug-target interactions *in cellulo*. By means of a duplexed, dual fluorescence CeTEAM platform, we compare the binding dynamics of reported PARPi for PARP1 and PARP2 drug biosensors and integrate ensuing cell cycle changes to generate a compendium of PARP1/2 binding and trapping profiles in live cells. While most PARPi equipotently engage PARP1/2, some possess significant selectivity for PARP1 (>10-fold), including AZD5305, but none favor PARP2. In general, binding potency correlates positively with PARP trapping phenotypes, except for veliparib. By incorporating conditional depletion of HPF1 to understand its influence on PARP1/2 and trapping by PARPi, we find that HPF1 knockdown modestly improves binding to PARP2 and additively promotes PARP trapping phenotypes. Our findings provide an avenue for integrating genetic screens into cell-based drug binding and selectivity assays by means of CeTEAM, as well as a valuable tool for evaluating the PARP1/2 binding and trapping selectivity of next-generation PARP inhibitors.

Results

Validation of a duplexed PARP1/2 CeTEAM fluorescent biosensor system

Despite almost all clinical PARPi targeting both PARP1 and PARP2, there is a paucity of data regarding intracellular PARP2 drug-target engagement, especially regarding relative potency (19–21). In our hands, we were unable to detect discernible shifts in PARP2 stability by the cellular thermal shift assay (CETSA), despite seeing expected increases in soluble PARP1 upon olaparib addition (22, 23), which may partially explain the limited data on PARP2 engagement in cells (**Supplementary Fig. 1A-C**). We previously demonstrated the CeTEAM amenability of GFP-tagged PARP1 L713F and the analogous mutant in PARP2 (L269A) individually with multiple PARPi by comprehensive dose-response profiling (18). To build a truly multiplexed platform for PARPi selectivity profiling, we utilized spectrally distinct fluorescent tags for simultaneous monitoring of PARP1 L713F (GFP) and PARP2 L269A (mCherry) within the same cellular context (**Fig. 1**). We validated this system by western blot analysis after a 24-hour PARPi treatment using talazoparib, olaparib, niraparib, veliparib, and 3-aminobenzamide (3-AB), with iniparib as a negative control (**Fig. 1A and B**). All PARPi showed dose-dependent stabilization of PARP1 and PARP2 except for iniparib. As before (18), 3-AB required higher concentrations for similar engagement. Unexpectedly, niraparib stabilized PARP2 L269A significantly worse than PARP1 L713F at 1 µM.

Live-cell fluorescence microscopy confirmed these results, demonstrating essentially equipotent stabilization EC_{50} values with both biosensors, which ranged from ~2 nM (talazoparib) to ~300-500 µM (3-AB). These values were generally in close agreement with reported biochemical IC_{50} and K_i values (**Fig. 1C-F**). Notably, however, niraparib binding to PARP2 L269A-mCherry was worse by >10-fold (~11 nM *vs* ~133 nM) (24). Findings with L269A-mCherry were consistent with earlier stabilization EC_{50} values from PARP2 L269A-GFP (18), suggesting that the choice of fluorescent protein tag did not specifically contribute to observed stabilization changes (**Fig. 1F**). Thus, our PARP1/2 duplexed CeTEAM biosensor system is a capable platform for detailed pharmacological assessments of PARP inhibitors.

Re-screening of PARP1 L713F-nLuc stabilizers with the PARP1/2 CeTEAM assay

We previously screened a ~1200 compound drug-like library for PARP1 L713F biophysical perturbagens and found >90% of potent PARP1i, as well as several non-PARPi that also stabilized the nLuc biosensor (18). To determine if these molecules also affected PARP2 L269A stability, we then re-screened the original hits with the duplexed PARP1/2 CeTEAM assay (**Fig. 2A**). This also afforded the opportunity to validate PARP1 L713F stabilization by an orthogonal readout and integrate cell cycle measurements in response to drug treatment, which could further inform on differential pharmacology. In total, the re-screen contained the annotated PARPi and 14 non-PARPi tested previously, as well as two additional PARPi reported to be PARP1- (MC2050) and PARP2-selective (MC3474; Compound 1), respectively (25). The compounds were tested at 10 µM and biosensor abundance was measured after 16 hours. Hits were defined as having statistically significant ($p < 0.05$) abundance changes from the DMSO (negative) control.

All PARPi significantly increased PARP1 L713F-GFP abundance, but only a few non-PARPi (PF-3758309, bisindolylmaleimide I, AZ960, Go 6983, and ACY-1083) also met this threshold, essentially mirroring the L713F-nLuc biosensor reported earlier (18). Notably, the DNMT trappers, decitabine and azacitidine, did not meaningfully increase L713F-GFP, in contrast to the nLuc reporter reported earlier (18). Almost all PARPi also significantly increased PARP2 L269A-mCherry abundance, except for MC2050 and MC3474 (**Fig. 2C**). AZD5305, a compound previously identified as a highly specific PARP1 inhibitor (26, 27), exhibited comparable potency against PARP2 at this high dose (**Fig. 2B and C**). The same non-PARPi that increased L713F-GFP also affected PARP2 L269A-mCherry, except for the two protein kinase C inhibitors (PKCi; bisindolylmaleimide I and Go 6983), which, in

contrast to PARP1 L713F, significantly decreased PARP2 L269A abundance. Another non-PARPi, AT9283, showed a modest but significant effect on PARP2 that was not seen with PARP1.

We then performed a cross-comparison of hits for PARP1 and PARP2 to help identify any clear trends (**Fig. 2D**). Of the compounds tested, 32 significantly stabilized both PARP1 and PARP2, while 8 did not appreciably affect either. All non-responsive molecules were annotated non-PARPi. Therefore, apart from select molecules, effects on PARP2 L269A abundance generally aligned with PARP1 L713F at the 10 µM screening concentration.

Bisindolylmaleimide I and Go 6983 are autofluorescence artifacts

We then performed a follow-up dose-response with tested non-PARPi to determine if any trends developed with increasing compound concentrations up to 10 µM (**Supplementary Fig. 2A-C**). Most of the hits from the initial screen with marginal effects were not recapitulated by dose-response profiling. The only compounds to elicit a dose-responsive increase in biosensor abundance were the PKC inhibitors, bisindolylmaleimide I and Go 6983, which were specific for PARP1 L713F-GFP (**Supplementary Fig. 2B and C and Supplementary Fig. 3**). Significant increases in PARP1 L713F-nLuc abundance with these compounds was also observed previously (18).

Intrigued by their preferential effect on PARP1 L713F-GFP abundance, we took a closer look at the raw images. Interestingly, both PKC inhibitors appeared to induce the translocation of PARP1 L713F-GFP from the nucleus to the perinuclear space, suggesting an alteration in PARP1 subcellular distribution (**Supplementary Fig. 3A**). A similar effect was also observed with WT PARP1-GFP, indicating it was not specific to the mutant protein (**Supplementary Fig. 3B**). However, when we switched the fusion tags (to eliminate their potential contributions), we saw the inverse: only PARP2-GFP was affected by bisindolylmaleimide I, not PARP1-mCherry (**Supplementary Fig. 3C**). In cells only expressing PARP1-mCherry, bisindolylmaleimide I still elicited perinuclear staining in the CELLCYTE X green channel (emission$_\lambda$: 502-561 nm), suggesting the compound itself was responsible for the signal. Indeed, both bisindolylmaleimide I and Go 6983 have near identical structures and previous studies suggested that both exhibit autofluorescence in the green-yellow visible spectrum (28) (**Supplementary Fig. 3D**). These results were validated by immunofluorescence (IF) confocal microscopy for endogenous PARP1 with Alexa Fluor 488 or 568 secondary antibodies, a set-up which afforded better spectral separation. Bisindolylmaleimide I treatment alone gave the same perinuclear staining in the 568 channel (spectral free band (VF) 590-620nm) but was filtered out in the 488 channel (spectral free band (VF) 490-530nm). Meanwhile, localization of endogenous PARP1 by Alexa Fluor 488 was unchanged from control cells (**Supplementary Fig. 3E**). Thus, bisindolylmaleimide I and Go 6983 are autofluorescence artifacts apparently unrelated to PARP1 or PARP2 stability.

Intracellular selectivity profiling of PARPi with CeTEAM

In parallel with our analysis of non-PARPi, we conducted dose-response experiments with the expanded PARPi library using the duplexed PARP1/2 biosensors to establish a compendium of PARP1 and PARP2 binding in live cells (**Fig. 3A**). Comparative analysis of the dose-responses for PARP1 L713F-GFP (**Fig. 3B**) and PARP2 L269A-mCherry (**Fig. 3C**), alongside literature-derived IC$_{50}$ values, facilitated a robust evaluation of each inhibitor's cellular efficacy, as summarized in **Table 1** and depicted in **Supplementary Figure 4**. As before, talazoparib, olaparib, veliparib, and 3-AB were equipotent for PARP1 and PARP2, while iniparib was inactive and niraparib was ~14-fold more potent towards PARP1.

Unsurprisingly, clinical-grade PARP1/2i were typically the most potent for both PARPs (low nanomolar), while early generation PARPi (*e.g.*, PJ34) or those not intended to directly target PARP1/2 (*e.g.*, Tankyrase-IN-2, ME0328) were significantly worse. As suggested by the name, Tankyrase-IN-2 predominantly targets TNKS1/2 but has low level activity towards PARP1(29), while ME0328 preferentially targets PARP3 (30). Furthermore, the next generation PARPi, senaparib (IMP4297) (31)

equipotently engaged PARP1/2 biosensors in the low nanomolar range like other potent PARPi, such as A-966492 (**Fig. 3D and Table 1**). Senaparib reportedly has biochemical IC_{50} values of 0.48 nM and 1.6 nM for PARP1 and PARP2, respectively (31, 32) , which align well with our intracellular binding data.

Most of the expanded PARPi library were equipotent binders of both PARPs, but those that demonstrated meaningful selectivity were only biased towards PARP1, not PARP2. To facilitate comparisons of PARPi selectivity, we performed a pairwise analysis of PARP1 and PARP2 biosensor saturation for each PARPi and compared their apparent EC_{50} values (**Fig. 3D**). While some inhibitors had slight selectivity towards PARP1 (*e.g.*, rucaparib, MC2050, and mefuparib), AZD5305, INO-1001, niraparib, and Tankyrase-IN-2 had >10-fold preference for PARP1 binding, although Tankyrase-IN-2 was significantly less potent overall. At 10 μM, we initially saw that AZD5305 exhibited comparable stabilization of both biosensors (**Fig. 2B and C**). However, it displayed markedly higher specificity towards PARP1 over PARP2 by extended dose-response, as evidenced by a substantial difference in EC_{50} values – 3.9 nM *vs* 6.2 μM (~1600-fold potency difference; **Table 1, Fig. 3**). Our measures of apparent PARP1/2 binding for AZD5305 align well with reported biochemical and intracellular IC_{50} values (26, 27). We could also confirm the slight specificity of MC2050 (25) for PARP1 in our cellular assays (PARP1: 625 nM, PARP2: 2482 nM), albeit with significantly worse efficacy than AZD5305 despite their similar biochemical potencies. Interestingly, the early generation PARPi, INO-1001, was quite robust in our assay and yielded a surprising ~30-fold preference for PARP1 over PARP2 (apparent PARP1 EC_{50}: ~7.9 nM, PARP2 EC_{50}: ~217.2 nM), although we did not find reports with conclusive data describing its PARP1/2 inhibitory potency.

Some PARPi also bound PARP1 and PARP2 much worse than anticipated. For example, PJ34 was >10-fold worse towards both PARPs in our assay as compared to reported IC_{50} values (**Fig. 3D and Table 1**). Additionally, despite published specificity for PARP2 (25), MC3474 was one of the least potent PARPi in our assay, which may relate to its relatively high biochemical IC_{50} values (**Supplementary Fig. 5 and Table 1**). These demonstrated a similar phenomenon to other early PARPi (*e.g.*, 3-AB), where intracellular potency is significantly compromised.

In addition to live-cell fluorescence microscopy, we validated these findings for many PARPi by western blot analysis – confirming that detected fluorescence changes were proportional to protein abundance changes (**Supplementary Fig. 5A and B**). Collectively, these results suggest that a duplexed PARP1/2 CeTEAM platform can effectively profile PARPi selectivity in a scalable assay with live, single cells.

PARP trapping phenotypes strongly correlate with PARP1/2 binding potency

In addition to the synthetic lethal DNA damage elicited in homologous recombination-defective cancer cells, many clinical PARPi can trap PARP1 and PARP2 on DNA as part of their anticancer mechanism of action (1, 4). It has previously been shown that measuring DNA content (cell cycle) is a suitable readout of the DNA replication stress/S-phase delay and subsequent G2 arrest resulting from PARP trapping (33). Similarly, we have also utilized the cell permeable stain, Hoechst 33342, as a proxy to profile trapping dynamics in live cells with the PARP1 L713F-GFP CeTEAM assay (18). We defined PARP trapping as PARPi that both engaged the PARP1 biosensor and elicited an S/G2 shift in the cell cycle. To profile how this expanded library of PARPi influenced cell cycle dynamics in relation to PARP1 and PARP2 binding, our PARPi library dose-response experiments in PARP1/2 biosensor cells were complemented with Hoechst 33342 (**Fig. 3E-G and Supplementary Fig. 6**). As before, this method enabled us to generate crude estimations of the cell cycle and estimated DNA trapping EC_{50}. In this case, a lower EC_{50} indicates more effective PARPi trapping, whereas a higher EC_{50} suggests reduced trapping. Further, juxtaposing this data with PARP1/2 biosensor binding can give insights to potential on- *vs* off-target activity of PARPi.

Overall, inhibitors that were the most potent binders of both PARP1 and PARP2 generally yielded the most potent trapping phenotypes, whereas the less potent molecules were generally poor at generating these phenotypes (**Fig. 3G**). Among these, the next generation PARPi, senaparib, was among the most potent elicitors of PARP trapping phenotypes (0.40 nM trapping EC_{50}), along with talazoparib and AZ9482 (0.28 and 0.35 nM, respectively). There was one exception to this trend – veliparib was a potent binder to both PARP1 and PARP2 but was exceptionally worse at trapping PARP to DNA (528 nM trapping EC_{50}), in line with previous data (4, 18). On the other hand, PJ34, Tankyrase-IN-2, and 3-AB, which were among the worst PARP1/2 binders, had minimal (PJ34 – 3.75 µM EC_{50}) or no observable trapping activity (Tankyrase-IN-2, 3-AB).

Interestingly, the PARP1-selective inhibitor, AZD5305 also demonstrated a trapping phenotype but was comparatively inferior to other potent PARP1/2 dual inhibitors, such as senaparib (60.7 nM *vs* 0.4 nM trapping EC_{50}, respectively; **Fig. 3F-G and Table 2**). Thus, while intracellular PARP1 abundance is predominant over PARP2, there is likely a significant contribution of PARP2 DNA trapping to the overall effect.

HPF1 depletion modestly affects PARPi binding potency but additively promotes PARP trapping

Histone PARylation Factor 1 (HPF1) is known for its essential role in regulating PARP activity *via* redirection of PARylation activity towards histone serine residues in response to DNA damage (14). As part of this mechanism, HPF1 contributes a key glutamine residue (Glu284) to the PARP1 and PARP2 active sites that facilitates rapid PAR initiation (13, 34). Despite its low cellular abundance relative to PARP1, HPF1 effectively influences PAR synthesis by a "hit-and-run", catalytic-like mechanism (34). Earlier *in vitro* studies have suggested that the presence of HPF1 slightly improves affinity of PARPi for PARP1 but not PARP2 (15), however its impact on PARPi binding in cells has not been comprehensively studied. To determine how HPF1 influences PARPi binding to PARP1/2, we complemented the duplexed PARP1/2 CeTEAM biosensor cells with doxycycline-regulable shRNA towards HPF1 to conditionally deplete HPF1 (**Fig. 4**). First, we generated three shRNAs targeting HPF1 and validated their knockdown efficiency at both the mRNA level by RT-qPCR (each >~90%) and protein level by Western blot analysis (each >~80%), demonstrating significant reductions in HPF1 expression (**Fig. 4A-C**). We then selected one hairpin (shHPF1 #1) for functional characterization.

We then assessed how HPF1 knockdown affected the binding dynamics of talazoparib, olaparib, niraparib, veliparib, 3-AB, and iniparib (**Fig. 4D-G**). PARPi binding curves for a control hairpin (shNT) were similar to earlier profiles (**Fig. 1E and F**). As expected, both biosensors were unaffected by iniparib irrespective of HPF1 levels. Like data with purified proteins, HPF1 had subtle effects on observed binding of other PARPi to PARP1 and PARP2. Observed stabilization of the PARP1 biosensor was generally unchanged with clinical grade inhibitors while PARP2 binding was generally enhanced by ~2-3-fold following HPF1 depletion (**Fig. 4G and Table 3**). Notably, talazoparib binding was improved by ~2-fold by HPF1 depletion for both PARP1 and PARP2. Thus, HPF1 can influence PARPi binding to PARP1 and PARP2 in cells, which has important implications for intracellular PARPi selectivity. Additionally, our data demonstrates that genetic components can be layered onto CeTEAM-based assays to investigate the contribution of other proteins to observed binding potencies of a given target.

In addition to assessing differences in target engagement following HPF1 knockdown, we sought to evaluate its impact on PARP inhibitor trapping effects. As HPF1 binds PARP1/2 following HD domain destabilization (13), HPF1 depletion should theoretically increase PARP trapping on DNA by increasing flexibility between the HD and catalytic domains. Indeed, as seen previously (17), HPF1 depletion induced an S/G2-phase delay consistent with PARP trapping-related replication stress (**Fig. 4H-J and Table 4**). HPF1 depletion-related S/G2 character appeared to additively compound the effects of all tested trapping PARPi, whereas profiles for iniparib and 3-AB only exhibited the basal increase from HPF1 knockdown (**Fig. 4H-J and Supplementary Fig. 7**). Overall, the data suggests that HPF1

knowdown has an additive effect on PARPi-induced trapping – presumably by already retaining a fraction of the PARP pool onto DNA.

Discussion

Poly-ADP-ribose polymerases (PARPs) are crucial players in cellular DNA repair and attractive drug targets. However, PARP1 and PARP2 have highly homologous catalytic domains, so truly selective PARPi have historically been difficult to identify. Traditional selectivity profiling is done with purified proteins, but this approach is highly dependent on the experimental conditions and omits several factors that can influence ligand binding, such as post-translational modifications (PTMs), biomolecular interaction partners, and general space constraints within the intracellular environment. Thus, cell-based selectivity assays can afford novel insights to drug interactomes and overcome some limitations of biochemical approaches (12). Here, we evaluated PARPi binding to PARP1 and PARP2 using a multiplexed cellular target engagement by accumulation of mutant (CeTEAM) approach (18) to generate live-cell selectivity and pharmacology profiles.

Overall, we found most PARPi are roughly equipotent for both PARP1 and PARP2 in cells. Among these was the next generation PARPi, senaparib, which has recently shown efficacy as a first-line option for advanced ovarian cancer (31, 35). Despite its promise in the clinic, relatively little is known about its potency and selectivity. By duplexed CeTEAM assay, we saw robust binding to both PARP1 and PARP2 (10.7 and 5.3 nM, respectively), similar to other potent, clinical-grade PARPi, thereby supporting its efficacious clinical activity.

There were also PARPi with divergent selectivity (arbitrary selectivity ratio >10-fold), but only with preference towards PARP1 – including niraparib, INO-1001, Tankyrase-IN-2, and AZD5305. Tankyrase-IN-2 is a potent tankyrase 1/2 inhibitor (PARP5A/B) with residual activity towards PARP1 (10 and 7 nM IC_{50} vs 710 nM, respectively). We could confirm similar binding activity towards PARP1 intracellularly by CeTEAM and residual activity towards PARP2 (789 nM vs 13.2 µM, respectively). Niraparib is an established and potent PARP1/2 inhibitor with high bioavailability (PARP1/2 biochemical IC_{50}: 3.8 and 2.1 nM, respectively (36)), but we surprisingly saw >10-fold better binding to PARP1 over PARP2 by CeTEAM (10.9 vs 133 nM stabilization EC_{50}, respectively).

The discrepancy in niraparib selectivity was unexpected given the abundance of *in vitro* data supporting its equipotency. A possible explanation for this is that structural deviations present in PARP2 L269A alter the catalytic site and affect PARPi binding. However, the majority of PARPi screened aligned well with reported selectivity and potency profiles. Alternatively, it was reported that niraparib can sterically clash with the N-terminus of αF in the PARP2 HD domain (24). Yet, other PARPi interacting similarly to niraparib (talazoparib, rucaparib) did not show such selectivity disagreements in our study. Furthermore, the L269A mutation increases flexibility within the HD, which might further invalidate this explanation. However, *in vitro* data of niraparib binding using fluorescence polarization, has also indicated an ~50-fold preference for PARP1 over PARP2 (27), suggesting that it could be more selective of a binder than originally thought.

AZD5305 has been touted as one of the first truly PARP1-selective inhibitors to date (26, 27), and our datasets with live cell CeTEAM selectivity profiling confirm its high selectivity for PARP1 over PARP2 by a remarkable ~1600-fold (3.9 nM vs 6.2 µM, respectively). Of all inhibitors tested, it was the only one that had such a high selectivity. It was recently reported that AZD5305 can still allosterically trap PARP2 onto DNA at higher concentrations (>1 µM), suggesting that it binds PARP2 above 1 µM and could influence intracellular selectivity and toxicity profiles (24). Indeed, binding and trapping of PARP2 has been implicated as the driver of hematological toxicities in patients treated with current PARPi (10, 11). However, based on our findings, there is a significant intracellular selectivity window (at least 1000-fold), implying that this can be largely avoided in a clinical setting. Thus, AZD5305 should

be a promising molecule for selective PARP1 targeting and validates the use of a duplexed PARP1/2 CeTEAM assay to screen for novel, highly selective PARPi.

In addition to allosteric modulation of the HD domains within PARP1 and PARP2, trapping is believed to be primarily driven by inhibitory potency – as inhibition also blocks PARP1/2 autoPARylation, thereby promoting their retention on DNA (6, 16, 37, 38). In agreement with this notion, we generally saw that trapping correlates highly with binding potency and particularly when targeting PARP1/2 equipotently. Veliparib was the clear exception to this trend, as it was the only potent PARPi that poorly traps (trapping EC_{50}: ~529 nM). This unique feature is likely driven by its considerably smaller size, which is unable to directly contact the HD domains and further promote PARP retention on DNA (24, 38). Interestingly, significant trapping phenotypes by veliparib arise near the binding saturation point of the PARP1/2 biosensors, suggesting that its trapping effect is primarily driven by inhibition – an insight afforded by the unique capabilities of the CeTEAM platform. In agreement with earlier reports (26), we also see that AZD5305 induces PARP trapping but to a lesser extent than most equipotent, clinical-grade PARPi (trapping EC_{50}: ~61 nM). This is conceivably due to the lack of extensive PARP2 trapping, which occurs more readily with inhibitors equipotently targeting PARP1 and PARP2. Still, the high trapping potency of AZD5305 underscores the significant contribution of higher intracellular PARP1 abundance to overall PARP trapping phenotypes.

Another surprise was that the early generation PARPi, INO-1001 (Inotek/Genentech) – which was halted after Phase 1 clinical trials (39), was a potent PARP1/2 binder with impressive selectivity for PARP1 over PARP2 (~30-fold; 7.9 vs 217.2 nM, respectively). Notably, the INO-1001 selectivity profile was like that of niraparib, perhaps due to structural similarity, although insights to its PARP binding modality are unavailable for direct comparisons. Like AZD5305, INO-1001 also induced potent PARP trapping phenotypes (~60 nM), but to a lesser extent than other clinical-grade inhibitors. Interestingly, dose-limiting toxicities of INO-1001 were hematological, which may, in retrospect, relate to inhibition or trapping of PARP2. It is unclear why clinical development was ceased, as no serious adverse effects were reported from the clinical trials and pharmacokinetics data was promising, although concerns about liver toxicity may have played a role (40, 41). Nevertheless, our data suggests it might be worth considering INO-1001 as a tool for studies differentiating PARP1 and PARP2 biology.

Besides multiplexing drug biosensors, we added to the capabilities of CeTEAM by layering on a genetic perturbation. Here, we sought to understand how HPF1, a critical component of the PARP1 and PARP2 catalytic sites, could influence binding of PARPi in cells. HPF1 was previously found to increase binding affinity of some PARPi for PARP1 but not PARP2 (6, 16). In our experiments with PARP1 L713F and PARP2 L269A biosensors, we instead saw generally improved PARPi binding when HPF1 was absent, particularly for PARP2. These discrepancies could be due to the nature of our PARP variants, as HPF1 preferentially binds PARP1/2 when the HD is in the open conformation and our mutants are constitutively flexible in the HD (13). Thus, HPF1 and ligand binding could have a compound effect on PARP biosensor stability. In line with this, as before (17), we saw that knockdown of HPF1 induced a PARP trapping-like phenotype, which could relate to how HPF1 affects PARP1/2 allostery and release from DNA. The HPF1 knockdown yielded an additive effect on PARP trapping following addition of PARPi, again suggesting that both factors contribute to HD conformational dynamics and, thus, dictate PARP1/2 retention.

In summary, we expand the capabilities of CeTEAM by both multiplexing PARP1 and PARP2 drug biosensors, while also incorporating knockdown of HPF1 to discern effects on PARPi binding and PARP trapping. The expansion into multiplexed CeTEAM biosensors opens the door for higher throughput intracellular selectivity profiling assays, yielding potentially more accurate selectivity data overall. Meanwhile, the integration of genetic perturbations into CeTEAM platforms can facilitate the discovery of novel regulators of drug-target interactions and expand the scope of drug interactomes. In

the context of PARP inhibitor development, our duplexed PARP1/2 CeTEAM platform should be a useful tool to aid the discovery of next generation, PARP1- or PARP2-selective inhibitors.

Experimental procedures

Cell lines and culturing conditions

U-2 OS, osteosarcoma cells, HEK293T, embryonic kidney epithelial cells, and CCRF-CEM, human T lymphoblasts, were obtained from the American Type Culture Collection (ATCC, Manassass, VA, USA). U-2 OS and HEK293T cells were cultured in DMEM high glucose GlutaMAX medium (Thermo Fisher Scientific) supplemented with 1% Penicillin-Streptomycin (Thermo Fisher Scientific) and 10% fetal bovine serum – FBS (Thermo Fisher Scientific), where CCRF-CEM cells were cultured in RPMI 1640 Medium, GlutaMAX™ Supplement (Thermo Fisher Scientific) supplemented with 1% Penicillin-Streptomycin (Thermo Fisher Scientific) and 10% fetal bovine serum. For *in vitro* fluorescent microscopy read-outs, DMEM, high glucose, no glutamine, no phenol red media (Thermo Fisher Scientific) supplemented with 1x GlutaMAX, 1% Penicillin-Streptomycin (Thermo Fisher Scientific) and 10% fetal bovine serum – FBS (Thermo Fisher Scientific) was used. Cell cultures were maintained at 37°C with 5% CO_2 in a humidified incubator.

Antibodies and chemicals

anti-mCherry (rabbit polyclonal, cat. No. PA5-34974), donkey anti-rabbit Alexa Fluor 568 (cat. No. A10042) and goat anti-rabbit Alexa Fluor 488 (cat. No. A11008) were obtained from Thermo Fisher, anti-PARP2 (rabbit polyclonal, cat. No. 55149-1-AP) and anti-PARP1 recombinant antibody (rabbit recombinant, cat. No 80174-1-RR-20) were acquired from Proteintech, anti-GAPDH (polyclonal rabbit, cat. No. ab9485) was purchased by Abcam, anti β-actin (mouse monoclonal, A5441) was obtained from Sigma, anti-HPF1 (rabbit monoclonal, 90876), was acquired from Cell Signaling, anti-PARP1 (mouse monoclonal, sc8007) and anti-GFP (mouse monoclonal, sc-9996) were purchased from Santa Cruz Biotechnologies. Doxycycline hydrochloride (Sigma-Aldrich) was dissolved in MilliQ water (2 mg/mL) and used at 0.75 μg/mL. Talazoparib, niraparib, olaparib, veliparib, AZ9482, senaparib, tankyrase-IN2 and INO-1001 (MedChemExpress) were dissolved in DMSO to a stock of 10 mM. Iniparib (MedChemExpress) was dissolved in DMSO to a stock of 20 mM. 3-aminobenzamide (3-AB; Sigma-Aldrich) was dissolved in DMSO to a stock of 100 mM. A-966492, pamiparib, PJ34, MC2050, MC3474, mefuparib, AZD5305, and rucaparib were kindly provided by the Rotili Lab and dissolved in DMSO at 10 mM. Bisbenzimide H 33342 trihydrochloride (Hoechst 33342, Thermo Fisher) was dissolved in MilliQ water to a stock of 20 mg/mL.

Recombinant DNA cloning

PARP2 L269A was subcloned (**Table S1**) from pENTR1a-PARP2 L269A-GFP (7, 18) into pENTR1a-C-mCherry by flanking SalI/NotI restriction sites and subsequently transferred to pLenti CMV Hygromycin DEST (Addgene plasmid #17454) or Ef1a-Tta3G-P2A-Blast (42). pENTR1a-PARP1 L713F-C-GFP was transferred to pLenti CMV Blast DEST (Addgene plasmid #17451).

Short hairpin RNA (shRNA) HPF1#1, #2 and #3, and NT (non-targeting) were oligo annealed into a prSITEP-TetR-Neomycin-akaLuc. The shRNA system backbone of the pRSITEP-U6Tet-(sh)-EF1-TetRep-2A-Puro-P2A-RFP670 plasmid (43) was modified to include a Tet-Repressor for controlled shRNA expression via doxycycline, a Neomycin resistance gene for antibiotic selection, and a luminescent reporter, akaLuc. The entire sequence is transferable using XbaI/SalI restriction sites. All subcloning into entry vectors was validated by automated sequencing; while shuttling into destination vectors was performed with Gateway LR Clonase II (ThermoFisher Scientific) and positive clones were confirmed by colony PCR. All plasmids, synthetic genes, and primers (Eurofins Genomics) are listed in **Table S1**.

Lentivirus production and transduction

Lentiviral production was performed following transfection of third generation lentiviral packing vectors by calcium phosphate precipitation. prSITEP Neo, pLenti CMV Hygro, pLenti CMV Blast, or pLenti CMV Puro lentiviral constructs were co-transfected with lentiviral packaging vectors (Gag-Pol, Rev, and VSV-G envelope) into subconfluent HEK293T cells. Viral particles were harvested at 48- and 72-hours post-transfection, and target cells were transduced at 1:1 dilution of lentivirus and fresh, complete medium in the presence of polybrene (8 μg/mL). Forty-eight hours post-transduction, target cells were re-plated at low density in the presence of G418/neomycin (Sigma-Aldrich, 450 μg/mL for six days; pINDUCER20 – Addgene plasmid #44012), puromycin (Sigma-Aldrich, 1 μg/mL for three days; pLenti CMV Puro – Addgene plasmid #17452), blasticidin (Sigma-Aldrich, 5 μg/mL for four days; pLenti CMV Blast – Addgene plasmid #17451), or hygromycin (Sigma-Aldrich, 100 μg/mL for five days; pLenti CMV Hygro – Addgene plasmid #17454) that was replenished at three-day intervals.

Reverse transcription quantitative PCR (RT-qPCR)

U-2 OS cells were plated at 60,000 cells/well in 6-well plates in the absence or presence of 0.75 μg/mL DOX. After 72 hours, the cells were harvested with TRIzol (Thermo Fisher Scientific). RNA was purified with the Direct-zol RNA MiniPrep kit (Zymo Research) according to the manufacturer's instructions and quantified on a De Novix DS-11 FX Spectophotometer/Fluorometer. cDNA was then generated with the iScript cDNA Synthesis Kit (Bio-Rad) according to the manufacturer's instructions. qPCR was performed with 2.5 ng cDNA per sample and iTaq Universal SYBR Green Supermix (Bio-Rad) using a Bio-Rad CFX96 Real-Time PCR Detection System. Relative quantity of target genes was calculated using the ΔΔCt method *via* normalization to *β-actin*. All qPCR primers are listed in **Table S1** and were ordered from Eurofins.

Western blotting

Cells were collected by trypsinization and lysed in RIPA buffer (20 minutes on ice with occasional mixing), and the clarified lysate was supplemented with Laemmli buffer (1x final). Following heating for 5 minutes at 95°C, the samples were either directly loaded for electrophoresis or frozen at -20°C for later use. Protein samples were separated on 4-20% gradient Mini-PROTEAN gels (Bio-Rad) or Midi-PROTEAN gels (Bio-Rad) prior to transferring onto 0.2 μm nitrocellulose with a Trans-Blot Turbo Transfer System (Bio-Rad). After blocking with ROTI®Block (Carl Roth) for 1 hour at room temperature primary antibodies were applied in 1x ROTI®Stock TBST (Carl Roth) at specified concentrations and left to incubate on a shaker at 4°C overnight. The primary antibodies used included anti-GFP probe (mouse monoclonal, 1:500), anti-mCherry (rabbit polyclonal, 1:2000), anti-PARP1 (mouse monoclonal, 1:500), anti-PARP2 (rabbit polyclonal, 1:2000), anti-GAPDH (rabbit, 1:2500), anti-β-actin (mouse monoclonal, 1:5000), and anti-HPF1 (rabbit, 1:1000). LI-COR secondary antibodies were diluted in 1x ROTI®Stock TBST (Carl Roth) at 1:10,000 and incubated at room temperature for one hour. The blots were then imaged using a LI-COR Odyssey Fc system and analysed with Image Studio Software (LI-COR).

Live cell fluorescence microscopy

For PARP L713F and PARP2 L269A experiments in live U-2 OS cells, 44,000 cells were plated in transparent, clear bottom 96-well plates (BD Falcon) on day 0 in complete medium. The following day media was changed to complete DMEM phenol-free medium, and inhibitors were added to their indicated final concentrations in complete DMEM phenol-free medium (final DMSO 0.1% or 1% [3-AB] v/v). After 16 or 24 hours, cell-permeable Hoechst 33342 was added to a final concentration of 1μg/mL for 30 minutes prior to imaging.

For shHPF1 experiments in live U-2 OS cells, Dox 0.75ug/mL was added 72h prior the start of the experiment. After this period, cells were trypsined and 44,000 cells were plated in transparent, clear bottom 96-well plates (BD Falcon) on day 0 in DMEM phenol-free medium. The next day, inhibitors were added to their indicated final concentrations in complete DMEM phenol-free medium (final DMSO 0.1% or 1% [3-AB] v/v). After 24 hours, cell-permeable Hoechst 33342 was added to a final concentration of 1μg/mL for 30 minutes prior to imaging.

Imaging was performed using the CellCyteX microscope (Cytena) at 10x magnification. The microscope was maintained at 37°C with 5% CO2 in a humidified incubator. Image analysis was then performed with CellProfiler software (Broad Institute) where the mean nuclei intensity upper quartile intensity mask of both GFP and mCherry were used to understand the fold changes and the integrated Hoechst intensity was used for cell cycle analysis.

Small molecule screen details
Composition, storage, and plating of screened compounds

Hit molecules from an earlier PARP1 L713F stability screen (18) were handled and plated as before but into clear plates 96-well plates cell culture plates (Starstedt) using an Echo 550™ acoustic liquid handler (LabCyte). An overview of the small molecules screened is detailed in **Table S2**.

Screen execution

U-2 OS PARP1 L713F-GFP/PARP2 L269A-mCherry cells were plated at 10,000 cells per well into drug-containing assay plates (final concentration 10 µM) using a Multidrop Combi liquid dispenser (Thermo Scientific). Cells were then incubated with drugs for 16 hours at 37 °C with 5% CO_2 in a humidified incubator. To minimize edge effects, the plates were placed in self-made humidity chambers that limited evaporation in the outer ring of wells. Imaging was performed using the CellCyteX microscope (Cytena) at 10x magnification. The microscope was maintained at 37°C with 5% CO2 in a humidified incubator. Image analysis was then performed with CellProfiler software (Broad Institute) where the mean nuclei intensity upper quartile intensity mask of both GFP and mCherry were used to understand the fold changes. The same strategy was applied to measure the dose-curves of all the non-PARPi (DC-05, A-395, ACY-1083, droxinostat, KDM5-C70, bisindolylmaleimide I (GF109203X), AZ960, Go 6983, linifanib (ABT-869), paclitaxel, PF-3758309, AT9283, azacitidine and decitabine).

Cellular thermal shift assay (CETSA)

CCRF-CEM cells ($1x10^6$ per treatment) were harvested by centrifugation at 500 x g for 5 minutes and washed twice with phosphate-buffered saline (PBS). Post-washing, cells were resuspended in 4mL Tris-buffered saline (TBS) containing a cocktail of proteasome inhibitors (EDTA-free, Roche). Cell lysis was achieved by alternately exposing cells to dry ice and absolute ethanol (VWR) for 3 minutes and to a temperature of 37°C for 3 minutes, repeating this cycle three times. Subsequently, lysates were centrifuged at 20,000 x g for 20 minutes at 4°C. The supernatant was collected and aliquoted into 100 µL fractions. Each aliquot was treated with either DMSO or 10 µM olaparib and incubated at room temperature for 20 minutes. Thermal denaturation was conducted at a range of temperatures: 37°C, 40°C, 43°C, 46°C, 49°C, 52°C, 55°C, 58°C, 61°C, 64°C, 67°C, and 70°C, each for 3 minutes. Samples were then cooled down at room temperature for 3 minutes before being centrifuged at 20,000 x g for 20 minutes at 4°C. Samples were prepared by adding Laemmli 4X sample buffer and heated to 95°C for complete denaturation. Proteins were then resolved by SDS-PAGE and transferred onto membranes for Western blot analysis, as above.

Immunofluorescence (IF) microscopy

U-2 OS cells were seeded at a density of 4.44×10^4 cells per well in a PhenoPlate 96-well plate (Perkin Elmer). The next day, DMSO (0.1% v/v, final) or 10 µM bisindolylmaleimide I (GF109203X) was added to each well. After 24 hours the cells were fixed with 4% paraformaldehyde (PFA - Histolab) for 15 minutes and then permeabilized with 0.1% Triton X-100 (Sigma) in PBS for 15 minutes. Afterwards, the cells were blocked with 2% bovine serum albumin (BSA - Sigma) in PBS for one hour at room temperature. The cells were then incubated with a primary anti-PARP1 recombinant antibody (rabbit recombinant), diluted 1:200 in 0.1% BSA in PBS, for two hours at room temperature. Alexa Fluor 568 and Alexa Fluor 488-conjugated secondary antibodies (anti-rabbit) were diluted 1:500 in 0.1% BSA in PBS and applied to the cells at room temperature in the dark. After incubation with secondary antibodies. Cells were stained with Hoechst at a 1:2000 dilution for 5 minutes at room temperature. Cells were imaged using a A1R+ Nikon confocal microscope system at 20x magnification. The 568-channel spectral free band (VF) ranged between 590-620 nm and the 488-channel spectral free band (VF) ranged between 490-530 nm.

Statistical analysis

All graphing and statistical analyses were performed using GraphPad Prism V10. Saturation curve fitting was performed using the [agonist] vs response four parameter variable slope model in GraphPad Prism. Specific post-hoc tests, variations, and statistical significances for relevant experiments are described within individual figure legends. For **Tables 3 and 4** the $logEC_{50}$ p values were measured using unpaired, two-tailed t test assuming both populations have the same SD (standard deviation).

Supporting information: This article contains supporting information.

(1) Valerie, N. C. K.; Sanjiv, K.; Mortusewicz, O.; Zhang, S. M.; Alam, S.; Pires, M. J.; Stigsdotter, H.; Rasti, A.; Langelier, M.-F.; Rehling, D.; Throup, A.; Desroses, M.; Onireti, J.; Wakchaure, P.; Almlöf, I.; Boström, J.; Bevc, L.; Benzi, G.; Stenmark, P.; Pascal, J. M.; Helleday, T.; Page, B. D. G.; Altun, M. Coupling Cellular Drug-Target Engagement to Downstream Pharmacology with CeTEAM. *bioRxiv* **2024**, 2022.09.19.505646. https://doi.org/10.1101/2022.09.19.505646.
(2) Langelier, M.-F.; Riccio, A. A.; Pascal, J. M. NAR Breakthrough Article PARP-2 and PARP-3 Are Selectively Activated by 5 Phosphorylated DNA Breaks through an Allosteric Regulatory Mechanism Shared with PARP-1. *Nucleic Acids Res.* **2014**, *42* (12), 7762–7775. https://doi.org/10.1093/nar/gku474.

Acknowledgments: We would like to express our gratitude to Seher Alam for the insightful discussions about the project, and to the SciLifeLab Compound Center, part of the Chemical Biology Consortium Sweden (CBCS), for assistance with compound management and spotting.

Funding: This research was supported by the Swedish Childhood Cancer Society (TJ2019-0036 – NCKV), Cancer Research KI (Karolinska Institutet) Blue Sky Grant (NCKV), Felix Mindus Contribution to Leukemia Research (2019-01992 – NCKV), Loo and Hans Osterman Foundation (2020-01208 – NCKV), Karolinska Institutet Research Foundation (2020-01685, 2022-01749 – NCKV), Swedish Cancer Society (21 0352 PT – NV), Hållsten Foundation (MA), SciLifeLab Technology Development Project Grant (MA), Novo Nordisk Pioneer Innovator Grant 1 (NNF22OC0076798 – MA, NV), and MJP was supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 859860.

Author contributions: Conceptualization (NCKV, MA), Methodology (MJP, NCKV, MA), Investigation (MJP, NCKV), Resources (NCKV, MA), Writing – original draft (MJP), Writing – review & editing: (MJP, AL, EF, DR, MA, NCKV), Visualization: (MJP, AL, NCKV, MA), Supervision (NCKV, MA, DR), Project administration (NCKV, MA), Funding acquisition (NCKV, MA, DR)

Conflict of interests: MA and NCKV are inventors on a patent application describing CeTEAM and its uses (PCT/EP2019/073769). The remaining authors declare no competing interests.

References

1. Lord, C. J., and Ashworth, A. (2017) PARP inhibitors: Synthetic lethality in the clinic. *Science*. **355**, 1152–1158

2. Bryant, H. E., Schultz, N., Thomas, H. D., Parker, K. M., Flower, D., Lopez, E., Kyle, S., Meuth, M., Curtin, N. J., and Helleday, T. (2005) Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. *Nature*. **434**, 913–917

3. Farmer, H., McCabe, H., Lord, C. J., Tutt, A. H. J., Johnson, D. A., Richardson, T. B., Santarosa, M., Dillon, K. J., Hickson, I., Knights, C., Martin, N. M. B., Jackson, S. P., Smith, G. C. M., and Ashworth, A. (2005) Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature*. **434**, 917–921

4. Murai, J., Huang, S. Y. N., Das, B. B., Renaud, A., Zhang, Y., Doroshow, J. H., Ji, J., Takeda, S., and Pommier, Y. (2012) Trapping of PARP1 and PARP2 by Clinical PARP Inhibitors. *Cancer Res*. **72**, 5588–5599

5. Thorsell, A. G., Ekblad, T., Karlberg, T., Löw, M., Pinto, A. F., Trésaugues, L., Moche, M., Cohen, M. S., and Schüler, H. (2017) Structural Basis for Potency and Promiscuity in Poly(ADP-ribose) Polymerase (PARP) and Tankyrase Inhibitors. *J Med Chem*. **60**, 1262–1271

6. Rudolph, J., Jung, K., and Luger, K. (2022) Inhibitors of PARP: Number crunching and structure gazing. *Proc Natl Acad Sci U S A*. 10.1073/PNAS.2121979119/-/DCSUPPLEMENTAL

7. Langelier, M. F., Riccio, A. A., and Pascal, J. M. (2014) PARP-2 and PARP-3 are selectively activated by 5' phosphorylated DNA breaks through an allosteric regulatory mechanism shared with PARP-1. *Nucleic Acids Res*. **42**, 7762–7775

8. Rouleau-Turcotte, É., and Pascal, J. M. (2023) ADP-ribose contributions to genome stability and PARP enzyme trapping on sites of DNA damage; paradigm shifts for a coming-of-age modification. *J Biol Chem*. 10.1016/J.JBC.2023.105397

9. Bai, P. (2015) Biology of Poly(ADP-Ribose) Polymerases: The Factotums of Cell Maintenance. *Mol Cell*. **58**, 947–958

10. LaFargue, C. J., Dal Molin, G. Z., Sood, A. K., and Coleman, R. L. (2019) Exploring and comparing adverse events between PARP inhibitors. *Lancet Oncol*. **20**, e15–e28

11. Farrés, J., Llacuna, L., Martin-Caballero, J., Martínez, C., Lozano, J. J., Ampurdanés, C., López-Contreras, A. J., Florensa, L., Navarro, J., Ottina, E., Dantzer, F., Schreiber, V., Villunger, A., Fernández-Capetillo, O., and Yélamos, J. (2015) PARP-2 sustains erythropoiesis in mice by limiting replicative stress in erythroid progenitors. *Cell Death Differ*. **22**, 1144–1157

12. Vasta, J. D., Corona, C. R., Wilkinson, J., Zimprich, C. A., Hartnett, J. R., Ingold, M. R., Zimmerman, K., Machleidt, T., Kirkland, T. A., Huwiler, K. G., Ohana, R. F., Slater, M., Otto, P., Cong, M., Wells, C. I., Berger, B. T., Hanke, T., Glas, C., Ding, K., Drewry, D. H., Huber, K. V. M., Willson, T. M., Knapp, S., Müller, S., Meisenheimer, P. L., Fan, F., Wood, K. V., and Robers, M. B. (2018) Quantitative, Wide-Spectrum Kinase Profiling in Live Cells for Assessing the Effect of Cellular ATP on Target Engagement. *Cell Chem Biol*. **25**, 206-214.e11

13. Suskiewicz, M. J., Zobel, F., Ogden, T. E. H., Fontana, P., Ariza, A., Yang, J. C., Zhu, K., Bracken, L., Hawthorne, W. J., Ahel, D., Neuhaus, D., and Ahel, I. (2020) HPF1 completes the PARP active site for DNA damage-induced ADP-ribosylation. *Nature*. **579**, 598–602

14. Gibbs-Seymour, I., Fontana, P., Rack, J. G. M., and Ahel, I. (2016) HPF1/C4orf27 Is a PARP-1-Interacting Protein that Regulates PARP-1 ADP-Ribosylation Activity. *Mol Cell*. **62**, 432–442

15. Rudolph, J., Roberts, G., and Luger, K. (2021) Histone Parylation factor 1 contributes to the inhibition of PARP1 by cancer drugs. *Nat Commun*. 10.1038/S41467-021-20998-8

16. Stojanovic, P., Luger, K., and Rudolph, J. (2023) Slow Dissociation from the PARP1-HPF1 Complex Drives Inhibitor Potency. *Biochemistry*. **62**, 2382–2390

17. Prokhorova, E., Zobel, F., Smith, R., Zentout, S., Gibbs-Seymour, I., Schützenhofer, K., Peters, A., Groslambert, J., Zorzini, V., Agnew, T., Brognard, J., Nielsen, M. L., Ahel, D., Huet, S., Suskiewicz, M. J., and Ahel, I. (2021) Serine-linked PARP1 auto-modification controls PARP inhibitor response. *Nat Commun*. 10.1038/S41467-021-24361-9

18. Valerie, N. C. K., Sanjiv, K., Mortusewicz, O., Zhang, S. M., Alam, S., Pires, M. J., Stigsdotter, H., Rasti, A., Langelier, M.-F., Rehling, D., Throup, A., Desroses, M., Onireti, J., Wakchaure, P., Almlöf, I., Boström, J., Bevc, L., Benzi, G., Stenmark, P., Pascal, J. M., Helleday, T., Page, B. D. G., and Altun, M. (2024) Coupling cellular drug-target engagement to downstream pharmacology with CeTEAM. *bioRxiv*. 10.1101/2022.09.19.505646

19. Burtscher, M. L., Gade, S., Garrido-Rodriguez, M., Rutkowska, A., Werner, T., Eberl, H. C., Petretich, M., Knopf, N., Zirngibl, K., Grandi, P., Bergamini, G., Bantscheff, M., Fälth-Savitski, M., and Saez-Rodriguez, J. (2024) Network integration of thermal proteome profiling with multi-omics data decodes PARP inhibition. *Mol Syst Biol*. **20**, 458–474

20. Knezevic, C. E., Wright, G., Remsing Rix, L. L., Kim, W., Kuenzi, B. M., Luo, Y., Watters, J. M., Koomen, J. M., Haura, E. B., Monteiro, A. N., Radu, C., Lawrence, H. R., and Rix, U. (2016) Proteome-wide Profiling of Clinical PARP Inhibitors Reveals Compound-Specific Secondary Targets. *Cell Chem Biol*. **23**, 1490–1503

21. Howard, R. T., Hemsley, P., Petteruti, P., Saunders, C. N., Molina Bermejo, J. A., Scott, J. S., Johannes, J. W., and Tate, E. W. (2020) Structure-Guided Design and In-Cell Target Profiling of a Cell-Active Target Engagement Probe for PARP Inhibitors. *ACS Chem Biol*. **15**, 325–333

22. Shaw, J., Dale, I., Hemsley, P., Leach, L., Dekki, N., Orme, J. P., Talbot, V., Narvaez, A. J., Bista, M., Martinez Molina, D., Dabrowski, M., Main, M. J., and Gianni, D. (2019) Positioning High-Throughput CETSA in Early Drug Discovery through Screening against B-Raf and PARP1. *SLAS Discov*. **24**, 121–132

23. Molina, D. M., Jafari, R., Ignatushchenko, M., Seki, T., Larsson, E. A., Dan, C., Sreekumar, L., Cao, Y., and Nordlund, P. (2013) Monitoring drug target engagement in cells and tissues using the cellular thermal shift assay. *Science*. **341**, 84–87

24. Langelier, M. F., Lin, X., Zha, S., and Pascal, J. M. (2023) Clinical PARP inhibitors allosterically induce PARP2 retention on DNA. *Sci Adv*. 10.1126/SCIADV.ADF7175

25. Tomassi, S., Pfahler, J., Mautone, N., Rovere, A., Esposito, C., Passeri, D., Pellicciari, R., Novellino, E., Pannek, M., Steegborn, C., Paiardini, A., Mai, A., and Rotili, D. (2020) From PARP1 to TNKS2 Inhibition: A Structure-Based Approach. *ACS Med Chem Lett*. **11**, 862–868

26. Illuzzi, G., Staniszewska, A. D., Gill, S. J., Pike, A., McWilliams, L., Critchlow, S. E., Cronin, A., Fawell, S., Hawthorne, G., Jamal, K., Johannes, J., Leonard, E., Macdonald, R., Maglennon, G., Nikkilä, J., O'Connor, M. J., Smith, A., Southgate, H., Wilson, J., Yates, J., Cosulich, S., and Leo, E. (2022) Preclinical Characterization of AZD5305, A Next-Generation, Highly Selective PARP1 Inhibitor and Trapper. *Clin Cancer Res*. **28**, 4724–4736

27. Johannes, J. W., Balazs, A., Barratt, D., Bista, M., Chuba, M. D., Cosulich, S., Critchlow, S. E., Degorce, S. L., Di Fruscia, P., Edmondson, S. D., Embrey, K., Fawell, S., Ghosh, A., Gill, S. J.,

Gunnarsson, A., Hande, S. M., Heightman, T. D., Hemsley, P., Illuzzi, G., Lane, J., Larner, C., Leo, E., Liu, L., Madin, A., Martin, S., McWilliams, L., O'Connor, M. J., Orme, J. P., Pachl, F., Packer, M. J., Pei, X., Pike, A., Schimpl, M., She, H., Staniszewska, A. D., Talbot, V., Underwood, E., Varnes, J. G., Xue, L., Yao, T., Zhang, K., Zhang, A. X., and Zheng, X. (2021) Discovery of 5-{4-[(7-Ethyl-6-oxo-5,6-dihydro-1,5-naphthyridin-3-yl)methyl]piperazin-1-yl}-N-methylpyridine-2-carboxamide (AZD5305): A PARP1-DNA Trapper with High Selectivity for PARP1 over PARP2 and Other PARPs. *J Med Chem*. **64**, 14498–14512

28. Tamir, T. Y., Drewry, D. H., Wells, C., Major, M. Ben, and Axtman, A. D. (2020) PKIS deep dive yields a chemical starting point for dark kinases and a cell active BRSK2 inhibitor. *Sci Rep*. 10.1038/S41598-020-72869-9

29. Buchstaller, H. P., Anlauf, U., Dorsch, D., Kuhn, D., Lehmann, M., Leuthner, B., Musil, D., Radtki, D., Ritzert, C., Rohdich, F., Schneider, R., and Esdar, C. (2019) Discovery and Optimization of 2-Arylquinazolin-4-ones into a Potent and Selective Tankyrase Inhibitor Modulating Wnt Pathway Activity. *J Med Chem*. **62**, 7897–7909

30. Sharif-Askari, B., Amrein, L., Aloyz, R., and Panasci, L. (2018) PARP3 inhibitors ME0328 and olaparib potentiate vinorelbine sensitization in breast cancer cell lines. *Breast Cancer Res Treat*. **172**, 23–32

31. Cai, S. X., Ma, N., Wang, X., Guo, M., Jiang, Y., and Tian, Y. E. (2024) The discovery of a potent PARP1 inhibitor Senaparib. *Mol Cancer Ther*. 10.1158/1535-7163.MCT-23-0625

32. Cai, S. X., Xu, Q., Hu, X., Jiang, Y., Wang, G., and Tian, E. (2016) Abstract LB-057: 1-Benzylquinazoline-2,4(1H,3H)-diones as potent PARP inhibitors. SAR of the quinazoline group and in vivo efficacy of clinical candidate IMP4297. in *Cancer Research*, p. LB-057, American Association for Cancer Research, **76**, LB-057

33. Michelena, J., Lezaja, A., Teloni, F., Schmid, T., Imhof, R., and Altmeyer, M. (2018) Analysis of PARP inhibitor toxicity by multidimensional fluorescence microscopy reveals mechanisms of sensitivity and resistance. *Nat Commun*. 10.1038/S41467-018-05031-9

34. Langelier, M. F., Billur, R., Sverzhinsky, A., Black, B. E., and Pascal, J. M. (2021) HPF1 dynamically controls the PARP1/2 balance between initiating and elongating ADP-ribose modifications. *Nat Commun*. 10.1038/S41467-021-27043-8

35. Wu, X., Liu, J., Wang, J., Wang, L., Lin, Z., Wang, X., Zhu, J., Kong, B., Fei, J., Tang, Y., Xia, B., Liang, Z., Wang, K., Huang, Y., Zheng, H., Lin, A., Jiang, K., Wang, W., Wang, X., Lou, G., Pan, H., Yao, S., Li, G., Hao, M., Cai, Y., Chen, X., Yang, Z., Chen, Y., Wen, H., Qu, P., Xu, C., Hsieh, C.-Y., Cui, M., Chen, L., Cheng, Y., Zhao, W., Pan, M., Tang, Y., Zhang, Y., Lin, X., Abulizi, G., Duan, W., and Zeng, L. (2024) Senaparib as first-line maintenance therapy in advanced ovarian cancer: a randomized phase 3 trial. *Nat Med*. 10.1038/S41591-024-03003-9

36. Jones, P., Altamura, S., Boueres, J., Ferrigno, F., Fonsi, M., Giomini, C., Lamartina, S., Monteagudo, E., Ontoria, J. M., Orsale, M. V., Palumbi, M. C., Pesci, S., Roscilli, G., Scarpelli, R., Schultz-Fademrecht, C., Toniatti, C., and Rowley, M. (2009) Discovery of 2-{4-[(3S)-piperidin-3-yl]phenyl}-2H-indazole-7-carboxamide (MK-4827): a novel oral poly(ADP-ribose)polymerase (PARP) inhibitor efficacious in BRCA-1 and -2 mutant tumors. *J Med Chem*. **52**, 7170–7185

37. Xue, H., Bhardwaj, A., Yin, Y., Fijen, C., Ephstein, A., Zhang, L., Ding, X., Pascal, J. M., VanArsdale, T. L., and Rothenberg, E. (2022) A two-step mechanism governing PARP1-DNA retention by PARP inhibitors. *Sci Adv*. 10.1126/SCIADV.ABQ0414

38. Zandarashvili, L., Langelier, M. F., Velagapudi, U. K., Hancock, M. A., Steffen, J. D., Billur, R., Hannan, Z. M., Wicks, A. J., Krastev, D. B., Pettitt, S. J., Lord, C. J., Talele, T. T., Pascal, J. M., and Black, B. E. (2020) Structural basis for allosteric PARP-1 retention on DNA breaks. *Science*. 10.1126/SCIENCE.AAX6367

39.   Yuan, Y., Liao, Y. M., Hsueh, C. T., and Mirshahidi, H. R. (2011) Novel targeted therapeutics: inhibitors of MDM2, ALK and PARP. *J Hematol Oncol*. 10.1186/1756-8722-4-16

40.   Bedikian, A. Y., Papadopoulos, N. E., Kim, K. B., Hwu, W. J., Homsi, J., Glass, M. R., Cain, S., Rudewicz, P., Vernillet, L., and Hwu, P. (2009) A phase IB trial of intravenous INO-1001 plus oral temozolomide in subjects with unresectable stage-III or IV melanoma. *Cancer Invest*. **27**, 756–763

41.   Morrow, D. A., Brickman, C. M., Murphy, S. A., Baran, K., Krakover, R., Dauerman, H., Kumar, S., Slomowitz, N., Grip, L., McCabe, C. H., and Salzman, A. L. (2009) A randomized, placebo-controlled trial to evaluate the tolerability, safety, pharmacokinetics, and pharmacodynamics of a potent inhibitor of poly(ADP-ribose) polymerase (INO-1001) in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention: results of the TIMI 37 trial. *J Thromb Thrombolysis*. **27**, 359–364

42.   Kanellis, D. C., Espinoza, J. A., Zisi, A., Sakkas, E., Bartkova, J., Katsori, A. M., Boström, J., Dyrskjøt, L., Broholm, H., Altun, M., Elsässer, S. J., Lindström, M. S., and Bartek, J. (2021) The exon-junction complex helicase eIF4A3 controls cell fate via coordinated regulation of ribosome biogenesis and translational output. *Sci Adv*. 10.1126/SCIADV.ABF7561

43.   Eshtad, S., Mavajian, Z., Rudd, S. G., Visnes, T., Boström, J., Altun, M., and Helleday, T. (2016) hMYH and hMTH1 cooperate for survival in mismatch repair defective T-cell acute lymphoblastic leukemia. *Oncogenesis*. 10.1038/ONCSIS.2016.72

Figures

Figure 1. Establishment of a duplexed PARP1/2 CeTEAM drug biosensor platform. A. A representative western blot of PARP1 L713F-GFP and PARP2 L269A-mCherry abundance after 24-hour treatment with PARP inhibitors (PARPi) at indicated concentrations. **B.** Densitometric quantification of PARP1 L713F-GFP and PARP2 L269A-mCherry relative to GAPDH and normalized to the highest value observed for each reporter. Means shown ± SD (n=3). Statistical significance indicated as not significant (ns), $* - p < 0.05$, $** - p < 0.01$, $*** - p < 0.001$, and $**** - p < 0.0001$ as determined by one-way ANOVA with Dunnett's post-test comparing each condition to the DMSO vehicle control. **C.** Representative live-cell fluorescence micrographs of L713F-GFP (green) and L269A-mCherry (red) treated with DMSO or 10 µM talazoparib for 24 hours. Cell nuclei identified by Hoechst staining (blue) and scale bar = 100µm. **D.** Live-cell L713F-GFP fold change by fluorescence microscopy following a concentration gradient with the indicated PARPi for 24 hours and normalization to DMSO controls. Means ± SD (n=3) with lines-of-best-fit shown. **E.** Live-cell L269A-mCherry fold change by fluorescence microscopy, as in **D**. Means ± SD (n=3) with lines-of-best-fit shown. **F.** Observed stabilization EC50 values for PARP1 L713F-GFP and PARP2 L269A-mCherry with indicated PARPi compared with PARP2 L269A-GFP data from *Valerie et al.* Reported biochemical IC50 values given for reference. ND – not determined.

Figure 2. Rescreening of potential PARP1/2 biophysical modulators with duplexed PARPi biosensors. A. Overview of rescreening procedure for PARP1 (GFP)/PARP2 (mCherry) target engagement by live-cell fluorescent microscopy. Nuclei are identified by Hoechst 33342 staining. **B.** PARP1 L713F-GFP intensity following 10 μM PARPi (orange, 31 total) and non-PARPi (yellow, 14 total). Negative (DMSO, grey) and positive controls (10 μM veliparib, blue) are shown with intensity reference lines. Means shown for three wells (four images per well) ± SD. Statistical significance denoted as not significant (ns), $* - p < 0.05$, $** - p < 0.01$, $*** - p < 0.001$, and $**** - p < 0.0001$ as determined by one-way ANOVA with Dunnett's post-test comparing each condition to the DMSO vehicle control. **C.** PARP2 L269A-mCherry intensities for the rescreened compounds, as in **B**. **D.** Venn diagram representation of statistically significant hits for PARP1 and PARP2. Compounds affecting neither target are also shown, while those stabilizing only one target are named. RFU – relative fluorescence units.

Figure 3. Intracellular PARP inhibitor selectivity and trapping profiling with CeTEAM.
A. Experimental procedure for dose-dependent PARP1 L713F-GFP and PARP2 L269A-mCherry stabilization and cell cycle perturbations (Hoechst 33342) by PARPi with live-cell

fluorescent microscopy. **B.** Relative PARP1 L713F-GFP intensity expressed as fold change DMSO control after 24 hours of a PARPi gradient. Means shown from n=3. **C.** Relative PARP2 L269A-mCherry intensity expressed as fold change DMSO control after 24 hours of a PARPi gradient. Means shown from n=3. **D.** Two-dimensional comparison of PARPi stabilization EC_{50} values towards PARP1 L713F-GFP and PARP2 L269A-mCherry. Green – ≥10-fold PARP selectivity (PARP1 selective); red – ≥10-fold PARP2 selectivity (PARP2 selective); grey – <10-fold selectivity difference (equipotent). Values derived from n=3 replicates from **B** and **C**. **E.** Derivation of PARP trapping EC_{50} from the median Hoechst intensity (DNA content). **F.** Representative cell cycle profile dynamics for a PARP1-selective (AZD5305), an equipotent, trapping (talazoparib), and an equipotent, poorly-trapping PARPi (PJ34), as shown by Ridgeline plot. Red circles – approximate stabilization EC_{50} values for PARP1 (closed) or PARP2 (open) biosensors. **G.** Three-dimensional comparison of PARP1 (x-axis) and PARP2 (y-axis) engagement EC_{50} values and PARP trapping EC_{50} values (circle size) for tested PARPi. Circle size – 0.20 nM (smallest) to ≥3000 nM (largest), or open circle – EC_{50} not determined (ND). A representative experiment is used for trapping EC_{50} overlay.

Figure 4. An integrated CeTEAM-genetics approach to assess the influence of HPF1 on intracellular PARPi binding and trapping. A. RT-qPCR analysis of gene silencing efficiency by shRNAs targeting *HPF1* (shHPF1#1, shHPF1#2, or shHPF1#3) compared to non-targeting control (shNT). Gene expression levels were normalized to *β-actin* and expressed as percentage of shNT control. Means ± SD from n=2 replicates. **** – $p < 0.0001$ as determined by one-way ANOVA with Dunnett's post-test. **B.** Representative HPF1 protein knockdown after 72 hours DOX by Western blot. **C.** Quantification of HPF1 protein levels post-knockdown (related to **B**). Means ± SD from n=3 replicates. ** – $p < 0.01$ and *** – $p < 0.001$ as determined by one-way ANOVA and Dunnett's post-test. **D.** Experimental set-up for live cell tracking PARP1 (GFP) and PARP2 (mCherry) target engagement and PARPi-induced replication stress (cell cycle, Hoechst) by fluorescent microscopy following HPF1 knockdown. **E.** PARP1 L713F-GFP intensity fold change in control (solid line) or HPF1 knockdown (dashed line) cells after talazoparib (blue), olaparib (red), niraparib (green), veliparib (purple), iniparib (orange), or 3-AB (black) gradients for 24 hours. Data normalized to DMSO control. Lines-of-best-fit from n=3 experiments. **F.** PARP2 L269A-mCherry intensity fold change, as in **E**. **G.** shHPF1/shNT EC₅₀ value ratios (fold change) of talazoparib, olaparib, niraparib, veliparib, iniparib, and 3-AB

stabilization towards PARP1 (L713F-GFP) or PARP2 (L269A-mCherry). **H.** Representative cell cycle profile dynamics of control (shNT) or shHPF1 cells after a talazoparib, veliparib (**I**), or 3-AB gradient (**J**) for 24 hours, represented as a Ridgeline plot (related to **E** and **F**). Red circles – approximate stabilization EC_{50} values for PARP1 (closed) or PARP2 (open) biosensors.

Table 1. Comparison of CeTEAM stabilization EC$_{50}$ values and reported biochemical IC$_{50}$ values of tested PARPi towards PARP1/2 (M)

PARPi	PARP1 IC$_{50}$	PARP1 L713F-GFP EC$_{50}$	PARP2 IC$_{50}$	PARP2 L269A-mCherry EC$_{50}$
Talazoparib	1.20e-009[a]	1.854e-009	8.70e-010[a]	1.875e-009
Pamiparib	9.00e-010[a]	3.839e-009	5.00e-010[a]	1.467e-009
Niraparib	3.80e-009[a]	1.093e-008	2.10e-009[a]	1.325e-007
Veliparib	5.20e-009[a]	2.274e-008	2.90e-009[a]	1.062e-008
Rucaparib	1.40e-009[a]	1.110e-009	1.40e-009[a]	7.013e-009
A-966492	1.00e-009[b]	5.081e-009	1.50e-009[b]	1.201e-008
Olaparib	5.00e-009[a]	6.520e-008	1.00e-009[a]	2.689e-008
PJ34	1.10e-07[a]	1.913e-006	8.60e-008[a]	1.471e-006
Mefuparib	3.20e-009[a]	9.477e-009	1.90e-009[a]	9.156e-008
MC3474	2.04e-006[a]	ND	3.4e-007[a]	ND
MC2050	9.30e-009[a]	6.246e-007	5.49e-007[a]	2.496e-006
AZD5305	1.50e-009[b]	3.847e-009	6.53e-007[b]	6.172e-006
Tankyrase-IN-2	7.1e-007[a]	7.885e-007	-	1.315e-005
AZ9482	1.00e-009[b]	2.701e-008	1.00e-009[b]	1.849e-008
Senaparib	4.80e-010[a]	1.074e-008	1.60e-009[a]	5.257e-009
INO-1001	<5.0e-008[b]	7.198e-009	-	2.172e-007
ME0328	6.30e-006[a]	ND	1.08e-005[a]	ND
3-AB	5.00e-008[b]	5.338e-004	-	2.953e-004
Iniparib	-	ND	-	ND

[a] Biochemical assay; [b] Cell-based assay; ND – not determined

Table 2. DNA trapping EC$_{50}$ values of tested PARPi from median Hoechst intensity changes

PARPi	DNA trapping EC$_{50}$ (M)
Talazoparib	2.78e-010
Pamiparib	2.96e-009
Niraparib	2.85e-009
Veliparib	5.28e-007
Rucaparib	2.37e-008
A-966492	4.16e-008
Olaparib	2.57e-008
PJ34	3.75e-006
Mefuparib	2.36e-009
MC3474	ND
MC2050	9.25e-008
AZD5305	6.07e-008
Tankyrase-IN-2	ND
AZ9482	3.51e-010
Senaparib	3.98e-010
INO-1001	5.97e-008
ME0328	ND
3-AB	ND
Iniparib	ND

ND – not determined

Table 3. Comparison of PARPi EC_{50} values in shNT or shHPF1 cells

PARP1 L713F-GFP EC_{50}

PARPi	shNT (nM)	shHPF1 (nM)	logEC$_{50}$ p value*	shHPF1/shNT EC_{50} ratio
Talazoparib	1.949	0.963	0.0209	2.0
Olaparib	63.80	30.96	0.4754	0.7
Niraparib	28.49	27.60	0.9528	1.0
Veliparib	63.47	81.47	0.7858	0.8
Iniparib	NA	NA	NA	NA
3-AB	147,300	187,000	0.3837	0.8

PARP2 L269A-mCherry EC_{50}

PARPi	shNT (nM)	shHPF1 (nM)	logEC$_{50}$ p value*	shHPF1/shNT EC_{50} ratio
Talazoparib	1.819	0.7461	0.0180	2.4
Olaparib	16.01	5.993	0.0155	2.7
Niraparib	140.2	78.68	0.6835	1.8
Veliparib	18.63	9.159	0.3355	2.0
Iniparib	NA	NA	NA	NA
3-AB	145,100	206,900	0.6325	0.7

* – two-tailed, unpaired t test; others are not significant

Table 4. Comparison of shNT and shHPF1 trapping EC_{50} values by PARPi

PARPi	shNT (nM)	shHPF1 (nM)	$logEC_{50}$ p value*	shHPF1/shNT EC_{50} ratio
Talazoparib	0.29	0.09	0.1001	3.2
Olaparib	14.4	1.89	0.1382	7.6
Niraparib	7.85	1.74	0.1295	4.5
Veliparib	709.1	55.0	0.0745	12.9
Iniparib	NA	NA	NA	NA
3-AB	NA	NA	NA	NA

* – two-tailed, unpaired t test; others are not significant