Genetic predisposition towards multicellularity in *Chlamydomonas reinhardtii*

I-Chen Kimberly Chen[1], Shania Khatri[1], Matthew D. Herron[1], Frank Rosenzweig[1]

[1]School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA, USA

Keywords: multicellularity, genetic predisposition, experimental evolution, *C. reinhardtii*

Authors for correspondence:

I-Chen Kimberly Chen

kichen.evol@gmail.com

Frank Rosenzweig

frank.rosenzweig@biology.gatech.edu

Abstract

The evolution from unicellular to multicellular organisms facilitates further phenotypic

innovations, notably cellular differentiation. Multiple research groups have shown that in the

laboratory simple, obligate multicellularity can evolve from a unicellular ancestor under

appropriate selection. However, little is known about the extent to which deterministic factors

like ancestral genotype and environmental context influence the likelihood of this evolutionary

transition. To test whether certain genotypes are predisposed to evolve multicellularity in

different environments, we carried out a set of 24 evolution experiments each founded by a

population consisting of 10 different strains of the unicellular green alga *Chlamydomonas*

reinhardtii, all in equal proportions. Twelve of the initially identical replicate populations were

subjected to predation by the protist *Parmecium tetraurelia* while the other 12 were subjected to

settling selection by slow centrifugation. Population subsamples were transferred to fresh media

on a weekly basis for a total of 40 transfers (~600 generations). Heritable multicellular structures

arose in four of 12 predation-selected populations (6 multicellular isolates in total), but never in

the settling selection populations. By comparing whole genome sequences of the founder and

evolved strains, we discovered that every multicellular isolate arose from one of two founders.

Cell cluster size varied not only among evolved strains derived from different ancestors but

among strains derived from the same ancestor. These findings show that both deterministic and

stochastic factors influence whether initially unicellular populations can evolve simple

multicellular structures.

Introduction

A longstanding question in population biology centers around the relative roles that stochastic processes (e.g., mutation, recombination and genetic drift) versus deterministic process (e.g., natural selection) play in generating evolutionary trajectories and outcomes. In many instances of convergent evolution, the deterministic process of natural selection seems to prevail (1-3). For example, on each of the Caribbean's four largest islands (Puerto Rico, Jamaica, Cuba and Hispaniola), lizards of the genus *Anolis* exhibit four basic morphologies: slender grass-dwelling forms with long tails, ground-dwelling forms with long legs, short-legged forms that creep on twigs, and canopy-dwelling forms with large toe pads. These four body types independently adapted to similar habitats on each island, making it one of the best-studied examples of how evolutionary outcomes can be predictable (2). On the other hand, stochastic processes, such as a lineage's ancestral history and the levels of variation maintained by mutation and drift, can make evolutionary outcomes largely unpredictable (4). For instance, in *Escherichia coli* and *Salmonella typhimurium* resistance to the antibiotic streptomycin caused lower rate of protein synthesis, thereby reducing the fitness of cells when the drug was not present (5, 6). This fitness cost can be diminished by compensatory mutations at other sites in the bacteria's genomes (7, 8). However, when many of those compensatory mutations were introduced into wild-type, streptomycin-sensitive backgrounds, they were found to be detrimental, indicating that the fitness effect of those compensatory mutations depended on genetic backgrounds (5, 6, 9). Had those compensatory mutations occurred without the prior streptomycin-resistant mutations, their evolutionary trajectories and fitness outcomes could have been very different.

Laboratory microbial evolution has proved to be a useful tool for investigating under what

conditions stochastic versus deterministic factors exert the stronger influence on adaptation

evolution. Well-controlled, highly replicated experiments can be carried out for hundreds or even

thousands of generations under different types and/or intensities of selection. Such studies have

repeatedly shown that similar environments tend to select for similar phenotypes, suggesting that

evolution can be deterministic and predictable. For example, all 12 lines in Richard Lenski's

long-term evolution experiment (LTEE) in *E. coli* evolved faster growth and larger cell size than

the ancestor (10), as well as similar overall patterns of gene expression (11). In asexual haploid

Saccharomyces cerevisiae populations, sterility and whole genome duplication repeatedly

evolves (12, 13). Of course, stochastic processes like mutations can produce these predictable

results, provided that the genetic architecture of a selectively advantageous phenotype is

sufficiently complex that it can be approached via different mutational trajectories (14). But

sometimes, stochastic events like mutations can produce wholly unexpected outcomes. For

example, in one of the twelve LTEE *E. coli* populations that had evolved for >31,000 generations

in glucose-limiting, citrate-buffered medium, a weakly beneficial mutation followed by a

refining mutation together conferred the ability to exploit citrate (Cit^+) as a growth substrate.

These mutations were strongly selected, resulting in a Cit^+ dominated, Cit^+/Cit^- population (15-

17). A diagnostic feature of wild-type *E. coli* is its inability to ferment citrate under oxic

conditions. Thus, this two mutation combo, a tandem duplication that captured an aerobically

expressed promoter for the expression of a previously silent citrate:succinate antiporter gene,

citT, and a promoter mutation that activated expression of C_4-dicarboxylate:H^+ symporter *dctA,*

brought about a key evolutionary innovation (17). This innovation, which was contingent on an

earlier mutation in the citrate synthase gene *gltA,* benefited from even earlier mutations that

increased acetate production after the optimization of glucose metabolism, fundamentally altered

the ecological and evolutionary trajectory of the affected population (15, 18).

While the roles that stochastic versus deterministic factors play in producing

microevolutionary outcomes have been assessed in the lab, it is less clear how such factors have

impacted macroevolutionary outcomes, including major evolutionary transitions in the history of

life (19). Among these transitions is the evolution of multicellular life-forms from unicellular

ancestors, a transition that has occurred repeatedly across the Tree of Life (20-23), and that has

served as a gateway to further innovations, such as division of labor among cells, including

germ-soma differentiation (19). Historically, our understanding of the mechanisms that have

driven this transition has derived chiefly from paleontology (24-26), comparative morphology

(27-29), and genomics (30-37), all of which are retrospective in nature. However, in recent years

a number of investigators have adopted a prospective approach, and shown that simple, heritable,

obligate multicellularity can evolve under selection in the laboratory (38-42). Here, using

laboratory evolution we investigate the extent to which two different deterministic processes,

selection mode and genetic ancestry, enable a unicellular eukaryote, the green alga

Chlamydomonas reinhardtii, to make the evolutionary transition from single cells to

multicellularity.

C. reinhardtii is a member of the volvocine algae (Division Chlorophyta), a group that

encompasses species exhibiting different levels of complexity that range from single-celled *C.*

reinhardtii, to colonial *Gonium pectorale*, to fully differentiated *Volvox carteri*. Two prior

evolution experiments generated heritable multicellularity in *C. reinhardtii* via selective

regimens that favor increased size. Using settling selection, Ratcliff, Herron (41) evolved

amorphous multicellular clusters of $>10^2$ cells bound to one another by a transparent

extracellular matrix. Using predation selection by the protist *Paramecium tetraurelia*, Herron,

Borin (42) evolved a variety of stereotypical multicellular structures, consisting of 2^2 - 2^5 cells

surrounded by a common cell wall. These structures bear a striking resemblance to morphologies

exhibited by some of the more 'primitive' multicellular members of the volvocine algae, e.g.,

Pandorina and *Yamagishiella*.

Unfortunately, instead of starting from the same ancestral strain, the ancestors used in these

two selection experiments consisted of different outbred populations, making it impossible to

establish whether it was ancestral genotype or the selective regimen that determined which form

of multicellularity evolved. To test whether certain genotypes are predisposed to evolve

multicellularity in certain forms, or if particular environments select for specific multicellular

phenotypes, we carried out two sets of evolution experiments each founded by a population

consisting of 10 different strains of *C. reinhardtii*, all in equal proportions. Twelve initially

identical replicate populations were subjected to predation by *P. tetraurelia* and 12 replicate

populations were subjected to settling selection. Subsamples of these populations were

transferred to fresh media every week for a total of 40 transfers (~600 generations). Because the

medium used in this experiment contains abundant nitrate, we expect all reproduction to be

asexual (the entry into the sexual cycle is controlled by nitrogen deprivation in *C. reinhardtii*,

(43)). Founder strains included two lab strains and eight field isolates, each of which was

phenotypically (e.g., cell sizes and chlorophyll content) and genetically different from the others

(44). In particular, each of our founder strains contained unique stable polymorphisms that serve

as markers for identifying the probable ancestor(s) of evolved multicellular strains.

In our experiments, after ~600 generations, heritable, obligate multicellular structures arose

in four of 12 predation-selected populations, but never in the settling rate selected populations.

We found that only two of the original ten strains were 'parents' for these multicellular

descendants, suggesting that certain genetic backgrounds may predispose those strains to evolve

multicellularity under this selective regimen. However, the size and structure of the multicellular

types were different even in isolates originating from the same 'parent' strain, indicating that

stochastic processes like mutation accumulation can lead to strikingly different multicellular

forms in populations evolved under the same condition. Together these findings show that both

deterministic and stochastic factors can play important roles in shaping the path by which

populations evolve simple multicellularity.

Materials and Methods

Experimental evolution

A list of the *C. reinhardtii* founder strains used in this study is provided in **Table 1**. Strains

were selected on the basis of their geographic isolation from one another, their phenotypic and

phylogenetic diversity, and the fact that the genome of each strain has been sequenced (44). In

this study, each founder strain was grown to stationary phase (~6-8 x 10^6 cells/mL) in TAP

liquid medium (45); equal volumes of those cultures were mixed and used as the master stock to

start the selection experiment. Twelve replicate populations were subjected to predation selection

by *Paramecium tetraurelia* (Carolina Biological Catalogue #131560) as described by Herron,

Borin (42) and 12 replicate populations were subjected to settling rate selection (41) (**Figure 1**).

Each population was cultured on a 14:10-hour light:dark cycle at 22.5 °C in 15 mL of COMBO

medium. COMBO is a fresh water medium suitable for co-culturing algae and zooplankton (46).

We used COMBO for both predation and settling rate selection experiments so that the two

treatments could be compared. Subsamples of each population were transferred to fresh COMBO

medium every week for a total of 40 transfers, with approximately 15 generations elapsing

between transfers. For experiments under predation selection, 1.5 mL of co-cultured *C.*

reinhardtii and *P. tetraurelia* were transferred to fresh medium every week. For the settling rate

selection, each population was transferred to a 15 mL Falcon tube and centrifuged at 100 x *g* for

10 s and only the bottom 0.25 mL was transferred to fresh medium. Under both types of

selection, the bottleneck size at each transfer was approximately 1-2 x 10^6 cells.

Every two months (i.e., every eight transfers or 120 generations), 0.1 mL samples of each

experimental population were diluted 100-fold and 30 μL was plated on 0.5% TAP agar to

screen for multicellular isolates. Unicellular isolates are normally motile and form large colonies

on 0.5% agar plates, whereas multicellular isolates are normally non-motile and form small

colonies (preliminary data not shown). Three small colonies per population (presumably

multicellular isolates labelled "A", "B", and "C") were picked up using sterile toothpicks,

inoculated into 2 mL of TAP medium in 24-well plates, incubated overnight and screened the

next day for multicellular phenotypes by visual examination under light microscopy. Isolates that

demonstrated multicellular phenotypes were diluted 10-fold, then 30 μL was replated on solid

medium. The same procedure was repeated three times sequentially to ensure that each isolate

was clonal and that multicellular phenotypes of isolates from the predation treatment were not

transiently induced by the presence of the *P. tetraurelia* predator. All founder strains used to start

the evolution experiment and evolved multicellular isolates were cryopreserved at -80°C using

the GeneArt Cryopreservation Kit for Algae (Invitrogen, Thermal Fisher Scientific).

Cells per cluster in the multicellular isolates

At the end of our evolution experiments we obtained pure culture isolates of several

multicellular strains. To understand how cluster size distribution patterns varied among strains

over a 6-day incubation period as well as the maximum number of cells per cluster, we subjected

each strain to epifluorescence microscopy. Cells of evolved strains and their ancestors were

scraped from TAP agar plates (3-4 inoculation loops, 0.09 cm in diameter), inoculated into

15 mL of TAP medium then cultured for five days to high density (approximately 6-8 x 10^6

cells/mL). Next, we transferred cultures of each strain into 15 mL of TAP medium at a starting

density of 10^4 cells/mL, cultured on a 14:10-hour light:dark cycle at 22.5 °C, and sampled every

24 hours over the course of 6-days. We used TAP medium here because it is a standard medium

commonly used in *Chlamydomonas* research (45); multicellular phenotypes were stable in TAP.

Subsets of cell cultures were removed from well-mixed test tubes and transferred into 1.5 mL

Eppendorf tubes (from day 1 to day 6, the amounts of cell cultures transferred were 1200, 900,

400, 300, 200, 100 μL). Samples were centrifuged at 14,000 × g for 1 minute and the supernatant

was replaced by a 50% v/v ethanol-water solution to fix cells and render cell membranes

permeable to the fluorescent nucleotide stain DAPI (4′,6-diamidino-2-phenylindole). After a 10-

minute fixation period, samples were centrifuged again at 14000 × g for 1 minute and the

supernatant replaced with distilled water containing 1 μg/mL DAPI. Cells stained with DAPI

were kept in darkness for one hour at 25°C then moved to a 4°C refrigerator without light to

prevent photobleaching until being imaged on an inverted epifluorescence microscope (the

Eclipse Ti series, Nikon).

Prior to imaging, samples were removed from the refrigerator and gently pipetted. A 10 μL

aliquot of each sample was mounted on glass microscope slides, then multiple images were

recorded at 20x magnification using bright field and the DAPI filter set (Excitation/Emission:

358/461 nm) overlapping on top of each other. Images were taken of at least 20 clusters or cells

per sample, providing a robust dataset for computing the distribution of cluster sizes within each

population at a given time point. Next, images were manually screened to demarcate cluster

boundaries and record cell number within each boundary using Cell Counter in ImageJ

(https://imagej.nih.gov/ij/). Cell number was estimated by counting DAPI stained nuclei

confined within cluster boundaries (e.g., **Figure 2**). This procedure was repeated four times for

each of the evolved strains and twice for the ancestral strains.

Modes of reproduction in multicellular isolates

Multicellularity can arise either by many cells aggregating (47, 48) or by one cell

reproducing clonally and its daughter cells remaining intimately associated (49, 50). In order to

distinguish between these two forms of multicellularity, ancestral and evolved strains were

imaged using time-lapse transmission microscopy over 72 hours of growth. As before, cells were

scraped from TAP agar plates (3-4 inoculation loops, 0.09 cm in diameter), inoculated into liquid

TAP medium and cultured for five days to high density (approximately 6-8 x 10^6 cells/mL).

Then, each culture was vortexed, transferred 1:100 into fresh TAP medium, and allowed to grow

for two more days. Exponentially growing cultures were first mixed by vortexing, then diluted

1:10 in TAP medium, whereafter 100 μL from each culture was inoculated into the wells of a 96-

well tissue culture plate containing100 μL of TAP medium per well. Well positions for each

strain were determined by a random number generator. Wells not inoculated with algae were

filled instead with 100 μL of TAP cell-free medium. For time-lapse microscopy, the 96-well

plate was imaged at 200x magnification using the Nikon Eclipse Ti inverted microscope, where

the field of view was positioned on cells near the center and at the bottom of wells. The time-

lapse was run for 72 hours, capturing images of each well every 30 minutes. For each strain, five

technical replicates were performed on the 96-well plate.

Identifying the ancestors of evolved multicellular isolates

MATING TYPE ASSAY: To ascertain whether certain ancestral strains were more likely to

evolve multicellularity than others, we identified alleles that differed among founder strains

(such that each strain has a specific combination), and used those alleles as markers to determine

the probable ancestor(s) of evolved multicellular strains. We first established the alleles at the

mating type locus in evolved isolates, as the mating types of the founder strains had been

previously identified. Each evolved isolate was subjected to PCR-amplification using either the

mid or *fus1* locus (the *mid* gene is unique to mating type minus (mt-) cells, while the *fus1* is

unique to mating type plus (mt+) cells). The *mid* locus was amplified using the primer pair

MTM1F (CTG CTG GTA CAA AGG TGT GGC ACG) and MTM2R (CAT GCA GTC TCT

CTC ACC CAT TCG GC), which were expected to produce a 750bp piece of the *mid* gene; the

fus1 locus was amplified using the primer pair MTP2F (GCT GGC ATT CCT GTA TCC TTG

ACG C) and MTP2R (GCG GCG TAA CAT AAA GGA GGG TCG), which were expected to

produce a 434bp piece of the *fus1* gene (51). After identifying the mating type of evolved

isolates, we next compared how their whole genome sequences differed from the founder strains,

giving us deeper insight into which ancestors produced multicellular isolates.

WHOLE GENOME SEQUENCING: Genomic DNA from evolved isolates and the ancestral strain

CC-1690 (a wild-type strain in our lab) was extracted, using the *Chlamydomonas* DNA

extraction protocol recommended by the Joint Genome Institute (https://www.pacb.com/wp-

content/uploads/2015/09/Experimental-Protocol-DNA-extraction-of-Chlamydomonas-using-

CTAB-JGI.pdf). Genomic DNA libraries were prepared, and samples were sequenced at 2 x

151bp length using the Illumina NextSeq platform at Microbial Genome Sequencing Center

(MiGS, https://www.migscenter.com/). For the founder strains other than CC-1690, their

sequenced reads were downloaded from NCBI. The quality of sequenced reads of evolved strains

and founders was examined by FastQC and trimmed using Trimmomatic-0.36 (*options*:

HEADCROP:16 LEADING:30 TRAILING:28 SLIDINGWINDOW:4:15 MINLEN:54). Paired

reads were mapped to the Joint Genome Institute v5.5 assembly of the *C. reinhardtii* CC503 mt+

genome (52) using BWA-MEM default setting (53). Genomic variants (small polymorphisms

such as SNPs (single nucleotide polymorphisms), indels (insertions and deletions), MNPs (multi-

nucleotide polymorphisms) and complex events (composite insertion and substitution events))

were detected using FreeBayes v1.0.2-29 (*options*: -p 1 -m 3 -q 1) (54). Variants with quality

scores greater than 20, read depth greater than 5X, and allele frequencies greater than 0.85, were

analyzed. According to a previous study (44), each founder strain contains a genome fingerprint

that distinguish them from each other (e.g., see Flowers et al. 2015 **Figure 2B**). Our approach

was to generate a filtered genetic variant data set that consisted of nonreference variants

observed in all strains together, and identify ancestral identities of multicellular isolates using

strain-specific variants along with mating types in founder strains. Sequencing data were

deposited at the NCBI Sequence Read Archive with project number PRJNA820479 and

accession numbers SAMN27009621-4.

Results

Multicellular structures evolved under predation-selection but not under settling selection.

A founder population of *C. reinhardtii* was created by mixing in equal proportion each the 10

strains described in **Table 1**; subsamples of this founder population were subjected to 12-fold

replicated experimental evolution according to the scheme depicted in **Figure 1**. Hereafter, we

will refer to Predation Selection lines as PS01-PS12 and Settling Rate Selection Lines as SRS01-

SRS12. Under the predator selective regime, after 16 weekly transfers we observed multicellular

structures in two of the 12 replicate populations, PS05 and PS11. However, evolved isolates

from those two populations did not maintain their multicellular phenotypes after serial plate

transfers in the laboratory; neither did multicellular morphologies persist in these two

populations in the following screen after eight weeks. After 32 weekly transfers, multicellular

structures appeared in three populations, PS09, PS10 and PS11; one evolved strain was isolated

from each population (PS09-32-A, PS10-32-B and PS11-32-B respectively). Other than PS10-

32-B from PS10, the multicellular strains that evolved in populations PS09 and PS11 after 32

weekly transfers maintained multicellular morphologies after being plated on solid medium three

times in succession, as well as in the experiments described below (**Figure 2: Panels A-B**). After

an additional 8 weekly transfers (40 transfers altogether), multicellular structures were detected

in two other predation-selected populations, PS02 and PS12; three evolved strains were isolated

from PS02 (PS02-40-A, PS02-40-B, PS02-40-C) and one strain was isolated from PS12 (PS12-

40-C). Each of the evolved strains isolated from PS02 and PS12 after 40 weekly transfers

maintained their multicellular structures after being plated on solid medium three times in

succession, as well as in the experiments described below (**Figure 2: Panels C-F**). Together

these observations indicate that the evolved multicellular phenotypes under predation selection

are stably heritable. Under the settling rate selective regime, we observed multicellular structures

in two of the 12 replicate populations, SRS05 and SRS 07, after 16 weekly transfers, and again in

populations SRS03, SRS08 and SRS11 after 32 weekly transfers. However, isolates from those

populations did not maintain their multicellular phenotypes after serial plate transfers in the

laboratory. Thus, heritably stable multicellular structures were never observed in any of the

settling rate selected populations over the course of 40 weekly transfers, or ~600 generations.

In the unicellular ancestors, most cells exist as single cells in their respective populations

(**Figure 2: Panels G-H**). By contrast, within each multicellular structure of the evolved strains,

cell walls are visible surrounding individual two, four or eight cells, and these are held in place

by transparent extracellular matrix (**Figure 2: Panels A-F**). Most evolved strains form clusters

approximately 20-100 μm in size, except PS09-32-A, which can form clusters whose diameters

range up to several hundred micrometers (**Figure 2: Panel A**). Development in the evolved

multicellular strains is strictly clonal. Time-lapse microscopy over 72 hours shows that,

compared to the ancestors where dividing cells only transiently form clusters before daughter

cells are released (e.g., CC-2936, **Supplemental Video 1**), multicellular isolates develop by

parental cells dividing into daughter cells within clusters over multiple rounds of cell division

(**Supplemental Videos 2-7**). Because cells within each multicellular cluster are genetically

identical, clusters likely function as units of selection (55, 56) under predation selection.

Cells per cluster in the multicellular isolates

In order to determine cluster size distributions for evolved strains, we sampled, stained, and

imaged growing cultures every 24 hours over a 6-day period, which approximated the serial

transfer interval of our evolution experiments. Four biological replicates were performed for

each of the evolved strains, and two biological replicates were performed for the ancestral

strains. As distributions tended to be right-skewed, data were represented as boxplots that depict

clusters' central tendencies. Large outlier clusters were occasionally present in each population.

These outliers, defined as data points that exceeded a distance of 1.5 times the interquartile range

(IQR) above the 3rd quartile, were excluded from further analyses.

We observed that multicellular cluster size varied among evolved strains over the course of

6-days. The median cluster size of multicellular strain PS09-32-A was >80 cells on day-1 and

then maintained at >100 cells throughout the growth cycle, with a maximum of 156 cells on day-

6 (**Figure 3A**). By contrast, median cluster sizes of PS11-32-B, PS02-40-A, PS02-40-B, and

PS12-40-C ranged between 2 and 10 cells, with maximum sizes being reached on day-1 (6, 10, 7

and 7 cells, respectively) then gradually decreasing thereafter (**Figure 3B, C, D, F**). The median

cluster size of strain PS02-40-C remained at 4 cells between day-1 and day-5 before declining to

2 cells on day-6 (**Figure 3E**). Median cluster sizes of the ancestral strains (CC-2936 and CC-

1690, see the following paragraph for identification details) remained one cell per cluster over

growth cycle, with multicellular clusters occurring occasionally during reproduction (**Figure 3G-**

H).

We found that there was a significant difference among evolved strains and ancestral strains

in the median cluster sizes (Kruskal-Wallis Test, H = 242.22, df = 7, p << 0.01). To ensure our

comparisons were of similar developmental stages, the median for each strain was measured at

the time point at which it reached its maximum value, as in Herron, Borin (42). Particularly, the

median cluster size of PS09-32-A (156 cells) was significantly larger than those of other

multicellular strains (6, 10, 7, 4 and 7 cells from PS11-32-B, PS02-40-A, PS02-40-B, PS02-40-

C, PS12-40-C, respectively) and those of their ancestors (1 and 1 cell from CC-2936 and CC-

1690) (**Table 2**). There were no significant differences in median cluster sizes among all the

multicellular strains other than PS09-32-A (**Table 2**).

Multicellular isolates evolved from the founder strains CC-2936 and CC-1690

PCR analysis of mating types in the evolved multicellular isolates revealed that all were mt+,

suggesting they originated from at least one of the mt+ founder strains (**Supplemental Figure**

1). Therefore, for the following genomic analysis, we only compared evolved isolates with mt+

founders to identify which ancestor genotype(s) they were most likely to have originated from.

We performed whole genome sequencing of PS09-32-A, PS11-32-B, PS02-40-A in PS02 and

PS12-40-C to an aligned sequencing depth of ~15X, 15X, 14X and 18X using paired end (2 x

151bp) Illumina sequencing. One of the mt+ founder strains, CC-1690, was re-sequenced for a

separate study (not published) to an aligned depth of 44X using the same Illumina sequencing

method. We did not sequence PS2-40-B and PS2-40-C in PS02 owing to the fact that they are

phenotypically similar to PS02-40-A, and are likely the same genotype as PS2-40-A, or only

differ by a few mutations, as they were isolated from the same population. For the remaining

four mt+ founder strains, their sequenced reads were downloaded from NCBI using accession

numbers SAMN03272879, SAMN03272875, SAMN03272877 and SAMN03272872 for strains

CC-2937, CC-2936, CC-2344 and CC2343, respectively (44). Based on the sequence alignments

to ~112 Mb of the *C. reinhardtii* CC-503 mt+ reference genome (52), we generated a filtered

genetic variant data set that consisted of 4,714,560 nonreference variants in all strains together.

Each strain contains a genome fingerprint based on the location of their genetic variants across

17 *C. reinhardtii* chromosomes. By visually comparing the genome-wide distributions of those

variants and through direct comparisons, we can infer the most likely ancestors of evolved

multicellular isolates in the populations PS02, PS09, PS11 and PS12.

Among mt+ founder strains, lab strain CC-1690 is diverged from reference strain CC-503 by

90,242 variants in our filtered data set, while the field isolates CC-2937, CC-2936, CC-2344 and

CC-2343 differ from CC-503, respectively, by 1,651,964, 1,194,250, 2,102,758, 2,187,600

variants in 111,098,438 sites across the reference genome (**Table 3**). Among evolved strains,

PS09-32-A, PS11-32-B and PS02-40-A contain 1,169,953, 1,110,266 and 1,144,326

nonreference variants, respectively, while PS12-40-C only contains 77,708 variants (**Table 3**).

When comparing the genome-wide distributions of nonreference variants between mt+ founders

and evolved strains, we found that the distributions in PS09-32-A, PS11-32-B and PS02-40-A

visually match to that in CC-2936 while the distribution in PS12-40-C visually match to that in

CC-1690 (**Figure 4**). When directly comparing the variants between mt+ founders and evolved

strains, we found that > 99% of variants identified in the evolved strains PS09-32-A, PS11-32-B

and PS02-40-A matched to the founder strain CC-2936, while 100% of the variants identified in

PS12-40-C matched to the founder strain CC-1690 (**Table 3**), suggesting that PS09-32-A, PS11-

32-B, PS02-40-A evolved from CC-2936 and that PS12-40-C evolved from CC-1690. The < 1%

of variants in PS09-32-A, PS11-32-B, PS02-40-A that did not match to CC-2936 is likely due

either to different sequencing methods (e.g., CC-2936 was previously sequenced at 2 x 51bp

length using the Illumina HiSeq 2000 sequencer (44) while the evolved strains here were

sequenced at 2 x 151bp length using the Illumina NextSeq 2000 sequencer), or to mutations that

accumulated in these lines since they were sequenced by Flowers, Hazzouri (44).

To further illustrate that specific allele combinations in the founder strains CC-2936 and CC-

1690 are also present in their respective evolved isolates, **Table 4** lists 25 loci across different

chromosomal locations that differ among five mt+ founders. The nucleotides in those 25

positions in PS09-32-A, PS11-32-B and PS02-40-A all match to the ones in CC-2936, while the

nucleotides in those positions in PS12-40-C all match to CC-1690 (**Table 4**), indicating that CC-

2936 is the ancestor of PS09-32-A, PS11-32-B and PS02-40-A and that CC-1690 is the ancestor

of PS12-40-C.

Discussion

The emergence of multicellular organisms and the emergence of eukaryotes are widely regarded as major evolutionary transitions in the history of life (57). However, unlike eukaryogenesis, which seems to have been a singular event (58, 59), multicellularity has arisen at least 25 times across widely divergent lineages (20). Within the volvocine algae, multicellularity arose in some lineages as recently as the Jurassic and Cretaceous periods (126-49 million years ago) (37, 60-62). Still, even though multicellularity has arisen repeatedly and in diverse lineages, little is known about the relative roles that stochastic versus deterministic factors play in driving this transition. Previous work has shown that in the lab simple, obligate multicellularity can evolve in the unicellular green alga *C. reinhardtii* under either predation or settling selection (41, 42). Here, we used experimental evolution to assess the influences that genetic ancestry and selective environment have on whether single cells become stably multicellular. We established a founder population consisting of 10 different strains of *C. reinhardtii*, all in equal proportion. This founder population was used to generate 24 replicate populations: 12 were subjected to predation by the protist *P. tetraurelia*, 12 were subjected to settling selection that favors large cell clusters that settle rapidly. After ~600 generations, heritable, obligate multicellular structures arose in four of 12 predation-selected populations, but never in the settling rate selected populations. Furthermore, we found that multicellular isolates evolved from only two founders CC-1690 and CC-2936, and that the size and structure of some multicellular isolates from the founder CC-2936 differed from each other. Whether the obligate multicellularity we observed was at a stable point or at some point along a continuum of evolving multicellularity, our results

indicate that both deterministic and stochastic factors play significant roles on the trajectories by

which populations evolve multicellularity.

If the evolutionary transition to multicellularity was purely deterministic, we would expect

that similar selective environments would result in similar phenotypes regardless of ancestral

genotypes. However, the fact that we observed different types of multicellularity evolve under

predation selection, even from the same ancestor (CC-2936), suggests that stochastic processes

like the accumulation of random mutations can lead to largely different outcomes. That being

said, the large clusters formed by PS09-32-A (> 100 cells most of time) can be costly, as the

growth of a colony may be limited because interior cells may not have the same access to

nutrients or environmental signals (e.g. oxygen or lights) compared to exterior cells (63, 64). If

this is true, then it is possible that PS09-32-A might have evolved a smaller cluster size similar to

what we observed in other multicellular isolates had we allowed the experiment to run longer.

Such a response to selection would demonstrate that evolutionary process is still deterministic at

the phenotypic level.

We also found that the multicellular isolates evolved in three of 12 predation-selected

populations came from founder CC-2936, while the multicellular isolate evolved in the other

predation-selected population descended from founder CC-1690. We did not observe

multicellular isolates evolved from any other founders. These results suggest that particular

suites of genetic variants had accumulated in those strains prior to our evolution experiments,

and that these facilitated the transition to multicellularity in our selective regimen. An inverse

question to ask is: why are some founders more resistant than others to selection pressures

expected to favor evolution of multicellularity? Were the populations quickly purged of certain

founders just because they grew less quickly or to a lower stationary phase density? In other

words, were the results biased at the onset because certain strains have poor competitive ability,

relative to others? To test these ideas, future work should be directed toward competition

experiments among founders, and/or similar evolutions using the same founders where

populations are sequenced at different time points to monitor changes in the frequency of allelic

markers by which founders can be differentiated.

Multiple reasons could explain why multicellularity did not evolve under our settling

selection regimen. First, in our experiments settling rate selection was episodic, occurring only

once a week, whereas predation selection was continuous. Second, our settling rate selection

(centrifugation at 100 g for 10 s, with pelleted cells transferred to fresh medium once a week)

differed slightly from that of Ratcliff, Herron (41) (100 g for 5 s, with pelleted cells transferred

every 3 days). Third, so that the two treatments could be compared it was essential that the

settling selection and the predation experiments be carried out in same medium. We elected to

use COMBO, a defined freshwater medium for co-culturing algae and zooplankton (46) that had

been used successfully in our earlier predation experiment (42). Significantly, COMBO does not

contain an organic carbon source, therefore, it only supports photoautotrophic growth of *C.*

reinhardtii (65). In this key respect, COMBO differs from the TAP medium more commonly

used in *Chlamydomonas* research (including the Ratcliff, Herron (41) settling selection

experiment). TAP contains acetate, which supports heterotrophic growth by *C. reinhardtii*.

COMBO necessarily brings about a different physiological state than TAP, and this difference

may influence *C. reinhardtii*'s latent tendency to form clonal assemblages and/or extracellular

matrices that cause bodies to settle rapidly in aqueous media. Fourth, by luck of the draw we

might not have initiated our experiment with the right set of ancestral genotypes, or with

sufficient standing genetic variation. Ratcliff, Herron (41) initiated their evolutions using an

outbred population that had a high level of standing genetic variation; moreover, all populations

went through one round of sexual reproduction (within-population mating) during the evolutions.

After 73 transfers (~315 generations) with settling rate selection imposed every three days,

multicellular *C. reinhardtii* evolved in only one of 10 replicate populations. The difference

between our results and those of Ratcliff, Herron (41) suggests that the amount of standing

genetic variation might play a role in the tempo and mode by which multicellularity emerges.

Finally, perhaps multicellularity did not evolve under the settling selection here simply because

the outcome is stochastic. After all, in Ratcliff, Herron (41) only one of 10 replicate populations

evolved multicellularity under settling selection. If we use this as an estimate of the probability

of multicellularity evolving under settling selection, the binomial probability of zero of twelve

replicates evolving multicellularity would be 0.28 ($p(x=0)$ given 12 trials, each with a probability

of success of 0.1), far too likely to rule out.

Regarding the genetic basis of multicellularity in our experiments, we only sequenced

evolved isolates exhibiting multicellularity at low coverages (14-18X), as our objective was to

determine ancestry. Evolved unicellular isolates, in particular ones sharing a common ancestor

with the evolved multicellular isolates, were not sequenced due to budget constraints. Therefore,

comparisons between unicellular and multicellular evolved isolates, or between founder strains

and their evolved multicellular isolates, are not possible for the investigation on the mutational

trajectories to multicellularity here. Future directions include repeated backcrossing multicellular

traits into the founder or wild-type strains to generate near isogenic lines, or directly sequencing

the candidate genes known to cause multicellular phenotypes to identify the mutation(s)

responsible for the transitions to multicellularity. For example, previous work has shown that the

retinoblastoma (*RB*) cell cycle regulatory pathway is involved in cluster formation in the alga

Gonium, an undifferentiated colonial relative of *Chlamydomonas* (33). Indeed, expression of the

Gonium RB gene in a *Chlamydomonas* strain lacking its *RB* gene caused this mutant to form

colonial assemblies containing 2-16 cells (33). It is plausible that mutations involved in the cell

cycle pathway such as *RB* may underlie the transition from unicellular to multicellular

phenotypes in our evolution experiments.

Taken together, our findings show that past genetic changes in some strains of

Chlamydomonas can shift the probabilities of alternative evolutionary paths towards

multicellularity. Further, subsequent mutations can play a significant role in shaping divergent

multicellular forms even in populations evolved under the same condition.

Acknowledgements

This work was supported by NASA Astrobiology Institute Award NNA17BB05A (Rosenzweig

PI, Herron co-I) and by NSF Rules of Life (Rosenzweig co-PI). The authors thank Ozan Bozdag

and Alireza Zamani-Dahaj in the Ratcliff lab for useful discussion and for their assistance with

epifluorescence microscopy. We also thank Peter Conlin, Jim Elser, Ross Lindsey, and Emily

Cook for helpful comments on earlier drafts of this manuscript. This material is based upon work

while MDH was serving at the National Science Foundation.

Figure 1. Experimental evolution diagram. A founder population consisting of 10 different strains of *C. reinhardtii*, all in equal proportion, was used to generate 24 replicate populations: 12 were subjected to settling selection that favors large cell clusters that settle rapidly; 12 were subjected to predation by the protist *P. tetraurelia*. Subsamples of each population were transferred to fresh medium every week, with approximately 15 generations elapsing between transfers, for a total of 40 transfers. For the settling rate selection, each population was transferred to a 15 mL Falcon tube and centrifuged at 100 x *g* for 10 s and only the bottom 0.25 mL was transferred to fresh medium. For populations under predation selection, 1.5 mL of co-cultured *C. reinhardtii* and *P. tetraurelia* were transferred to fresh medium every week.

Figure 2. Microscopic images of each evolved strain and their unicellular ancestors stained by DAPI. (A) PS09-32-A; (B) PS11-32-B; (C) PS02-40-A; (D) PS02-40-B; (E) PS02-40-C; (F) PS12-40-C; (G) CC-2936; (H) CC-1690. Black arrows indicate examples of extracellular matrix.

Genomic data indicate that strain PS12-40-C evolved from CC-1690 while the rest of

multicellular strains evolved from CC-2936 (see **Table 3**).

Figure 3. Box plots of cells per cluster over a 6-day period. Cluster sizes were measured by sampling cultures over six days of growth, staining nuclei with DAPI, and imaging using fluorescent microscopy. (A-F) Cluster sizes in the evolved strains; (G-H) Cluster sizes in the ancestors. An x or cross represents the mean of cells per cluster; the crossbar indicates the median of cells per cluster. The grey horizontal bar in each figure represents one cell per cluster. Note the y-axis scales vary among strains.

Figure 4. Genome-wide distributions of nonreference variants in evolved and mt+ founder strains on 17 *Chlamydomonas* chromosomes.

Table 1. *C. reinhardtii* founder strains used in this experiment.

Strain	Mating type*	Origin	Location	Reference(s)
CC-1690	mt+	Lab	-	Sager (1955)
CC-5080	mt-	Lab	-	Gallaher et al (2015)
CC-2937	mt+	Field	Quebec, Canada	Sack et al (1994)
CC-2936	mt+	Field	Quebec, Canada	Sack et al (1994)
CC-2935	mt-	Field	Quebec, Canada	Sack et al (1994)
CC-2344	mt+	Field	Ralston, PA	Spanier et al (1992)
CC-2342	mt-	Field	Pittsburgh, PA	Spanier et al (1992)
CC-2343	mt+	Field	Melbourne, FL	Spanier et al (1992)
CC-2931	mt-	Field	Durham, NC	Pröschold et al (2005)
CC-2290	mt-	Field	Minnesota	Gross et al (1988)

*There are two mating types, identical in appearance and known as mating type *plus* and *minus*, in *C. reinhardtii*. The + or - sign in each parenthesis denotes the mating type of each strain. The entry into the sexual cycle is controlled by nitrogen deprivation in *C. reinhardtii*. The media used in the experiment sustain normal cell growth and we expect all reproduction to be asexual.

Table 2. P-values for Dunn's Test for all pairwise comparisons between the maximum cluster sizes in different strains. Single asterisks indicate p-values of <0.05. Double asterisks indicate p-values of < 0.01.

	Maximum cluster size (median)							
	PS09-32-A (156 cells)	PS11-32-B (6 cells)	PS02-40-A (10 cells)	PS02-40-B (7 cells)	PS02-40-C (4 cells)	PS12-40-C (7 cells)	CC-2936 (1 cell)	CC-1690 (1 cell)
PS09-32-A		**	*	**	**	**	**	**
PS11-32-B			0.56	0.81	0.40	0.48	**	**
PS02-40-A				0.72	0.28	0.31	**	**
PS02-40-B					0.38	0.43	**	**
PS02-40-C						0.98	**	**
PS12-40-C							**	**
CC-2936								0.24
CC-1690								

Table 3. Comparisons between genomic variants in evolved and mt+ founder strains. The numbers in parenthesis after strain names indicate the number of nonreference variants identified in each strain (see Methods). Heat maps are color-coded (light grey/low percentage match < grey/middle percentage match < dark grey/high percentage match).

	CC-1690 (90,242)	CC-2937 (1,651,964)	CC-2936 (1,194,250)	CC-2344 (2,102,758)	CC-2343 (2,187,600)
PS09-32-A (1,169,953)	59.56%	35.46%	99.93%	22.50%	18.41%
PS11-32-B (1,110,266)	59.15%	35.11%	99.90%	22.28%	18.18%
PS02-40-A (1,144,326)	59.59%	35.40%	99.93%	22.42%	18.32%
PS12-40-C (77,708)	100%	2.02%	3.58%	1.33%	1.09%

Table 4. Examples of genomic variants that differ among mt+ founder strains and their presence and absence in the evolved strains.

Strain Position	CC-1690	CC-2937	CC-2936	CC-2344	CC-2343	PS09-32-A	PS11-32-B	PS02-40-A	PS12-40-C
Chrom_1 1782815	A	T	T	T	T	T	T	T	A
Chrom_1 1796517	A	G	A	A	A	A	A	A	A
Chrom_2 531304	G	G	A	G	G	A	A	A	G
Chrom_2 906332	T	T	T	C	T	T	T	T	T
Chrom_2 1216720	C	C	C	C	T	C	C	C	C
Chrom_3 666196	C	A	T	A	A	T	T	T	C
Chrom_3 983155	AA	AG	GG	GG	AG	GG	GG	GG	AA
Chrom_3 1128242	C	C	T	C	C	T	T	T	C
Chrom_4 14191	TGA	TGA	TGA	TA	TGA	TGA	TGA	TGA	TGA
Chrom_4 236504	G	G	G	G	A	G	G	G	G
Chrom_5 763807	G	A	A	A	A	A	A	A	G
Chrom_5 888052	C	T	C	C	C	C	C	C	C
Chrom_5 1624406	G	G	A	G	G	A	A	A	G
Chrom_6 122423	ACG	ACA	ACA	ATG	ACG	ACA	ACA	ACA	ACG
Chrom_6 204099	T	T	T	T	C	T	T	T	T
Chrom_7 1392564	CCCGC	CC	CC	CC	CC	CC	CC	CC	CCCGC
Chrom_8 301955	TG	AA	TG	TG	TG	TG	TG	TG	TG
Chrom_8 533535	A	A	G	A	A	G	G	G	A
Chrom_9 133015	A	A	A	C	A	A	A	A	A
Chrom_10 69703	C	C	C	C	T	C	C	C	C
Chrom_11 448221	A	C	C	C	C	C	C	C	A
Chrom_13 679947	ATCTAT	AAC	ATCTAT	AAC	AAC	ATCTAT	ATCTAT	ATCTAT	ATCTAT
Chrom_14 153285	G	G	A	G	G	A	A	A	G
Chrom_15 135507	C	C	C	T	C	C	C	C	C
Chrom_16 312391	C	C	C	C	G	C	C	C	C

References

1. Conway Morris S. 2003 *Life's Solution: Inevitable Humans in a Lonely Universe.* Cambridge Univ. Press.
2. Losos JB. 2010 Adaptive Radiation, Ecological Opportunity, and Evolutionary Determinism. American Society of Naturalists E. O. Wilson Award Address. Am. Nat. **175**(6), 623-39. (doi:10.1086/652433)
3. McGhee G. 2011 *Convergent Evolution: Limited Forms Most Beautiful*. MIT Press.
4. Gould SJ. 1989 *Wonderful Life: The Burgess Shale and the Nature of History*. W. W. Norton & Co..
5. Schrag SJ, Perrot V, Levin BR. 1997 Adaptation to the fitness costs of antibiotic resistance in *Escherichia coli*. Proc. Biol. Sci. **264**(1386), 1287-91. (doi:10.1098/rspb.1997.0178)
6. Maisnier-Patin S, Berg OG, Liljas L, Andersson DI. 2002 Compensatory adaptation to the deleterious effect of antibiotic resistance in *Salmonella typhimurium*. Mol. Microbiol. **46**(2), 355-66. (doi:10.1046/j.1365-2958.2002.03173.x)
7. Schrag SJ, Perrot V. 1996 Reducing antibiotic resistance. Nature **381**(6578), 120-1. (doi:10.1038/381120b0)
8. Bjorkman J, Samuelsson P, Andersson DI, Hughes D. 1999 Novel ribosomal mutations affecting translational accuracy, antibiotic resistance and virulence of *Salmonella typhimurium*. Mol. Microbiol. **31**(1), 53-8. (doi:10.1046/j.1365-2958.1999.01142.x)
9. Weinreich DM, Watson RA, Chao L. 2005 Perspective: Sign epistasis and genetic constraint on evolutionary trajectories. Evolution **59**(6), 1165-74. (doi:10.1111/j.0014-3820.2005.tb01768.x)
10. Lenski RE, Travisano M. 1994 Dynamics of adaptation and diversification: a 10,000-generation experiment with bacterial populations. Proc. Natl Acad. Sci. USA **91**(15), 6808-14. (doi:10.1073/pnas.91.15.6808)
11. Cooper TF, Rozen DE, Lenski RE. 2003 Parallel changes in gene expression after 20,000 generations of evolution in *Escherichia coli*. Proc. Natl Acad. Sci. USA **100**(3), 1072-7. (doi:10.1073/pnas.0334340100)
12. Fisher KJ, Buskirk SW, Vignogna RC, Marad DA, Lang GI. 2018 Adaptive genome duplication affects patterns of molecular evolution in *Saccharomyces cerevisiae*. PLoS Genet. **14**(5), e1007396. (doi:10.1371/journal.pgen.1007396)
13. Lang GI, Botstein D, Desai MM. 2011 Genetic variation and the fate of beneficial mutations in asexual populations. Genetics **188**(3), 647-61. (doi:10.1534/genetics.111.128942)
14. Kvitek DJ, Sherlock G. 2013 Whole genome, whole population sequencing reveals that loss of signaling networks is the major adaptive strategy in a constant environment. PLoS Genet **9**(11), e1003972. (doi:10.1371/journal.pgen.1003972)
15. Blount ZD, Borland CZ, Lenski RE. 2008 Historical contingency and the evolution of a key innovation in an experimental population of *Escherichia coli*. Proc. Natl Acad. Sci. USA **105**(23), 7899-906. (doi:10.1073/pnas.0803151105)
16. Quandt EM, Deatherage DE, Ellington AD, Georgiou G, Barrick JE. 2014 Recursive genomewide recombination and sequencing reveals a key refinement step in the evolution

of a metabolic innovation in *Escherichia coli*. Proc. Natl Acad. Sci. USA **111**(6), 2217-22. (doi:10.1073/pnas.1314561111)

17. Blount ZD, Barrick JE, Davidson CJ, Lenski RE. 2012 Genomic analysis of a key innovation in an experimental *Escherichia coli* population. Nature **489**(7417), 513-8. (doi:10.1038/nature11514)

18. Quandt EM, Gollihar J, Blount ZD, Ellington AD, Georgiou G, Barrick JE. 2015 Fine-tuning citrate synthase flux potentiates and refines metabolic innovation in the Lenski evolution experiment. eLife **4**, e09696. (doi:10.7554/eLife.09696)

19. Maynard Smith J, Szathmáry E. 1995 *The Major Transitions in Evolution*. Oxford University Press.

20. Grosberg RK, Strathmann RR. 2007 The Evolution of Multicellularity: A Minor Major Transition? Annu. Rev. Ecol. Evol. Syst. **38**, 621-54. (doi:10.1146/annurev.ecolsys.36.102403.114735)

21. Knoll AH. 2011 The Multiple Origins of Complex Multicellularity. Annu. Rev. Earth Planet. Sci. **39**, 217-39. (doi:10.1146/annurev.earth.031208.100209)

22. Sebe-Pedros A, Degnan BM, Ruiz-Trillo I. 2017 The origin of Metazoa: a unicellular perspective. Nat. Rev. Genet. **18**(8), 498-512. (doi:10.1038/nrg.2017.21)

23. King N. 2004 The unicellular ancestry of animal development. Dev. Cell **7**(3), 313-25. (doi:10.1016/j.devcel.2004.08.010)

24. El Albani A, Mangano MG, Buatois LA, Bengtson S, Riboulleau A, Bekker A, et al. 2019 Organism motility in an oxygenated shallow-marine environment 2.1 billion years ago. Proc. Natl Acad. Sci. **116**(9), 3431-6. (doi:10.1073/pnas.1815721116).

25. Gibson TM, Shih PM, Cumming VM, Fischer WW, Crockford PW, Hodgskiss MSW, et al. 2018 Precise age of *Bangiomorpha pubescens* dates the origin of eukaryotic photosynthesis. Geology **46**(2), 135-8. (doi:10.1130/G39829.1)

26. Tang Q, Pang K, Yuan X, Xiao S. 2020 A one-billion-year-old multicellular chlorophyte. Nat. Ecol. Evol. **4**(4), 543-9. (doi:10.1038/s41559-020-1122-9)

27. Tikhonenkov DV, Hehenberger E, Esaulov AS, Belyakova OI, Mazei YA, Mylnikov AP, et al. 2020 Insights into the origin of metazoan multicellularity from predatory unicellular relatives of animals. BMC Biol. **18**(1): 39. (doi:10.1186/s12915-020-0762-1)

28. Brunet T, King N. 2017 The Origin of Animal Multicellularity and Cell Differentiation. Dev. Cell **43**(2), 124-40. (doi:10.1016/j.devcel.2017.09.016)

29. Umen J, Herron MD. 2021 Green Algal Models for Multicellularity. Annu. Rev. Genet. **55**, 603-32. (doi:10.1146/annurev-genet-032321-091533)

30. King N, Westbrook MJ, Young SL, Kuo A, Abedin M, Chapman J, et al. 2008 The genome of the choanoflagellate *Monosiga brevicollis* and the origin of metazoans. Nature **451**(7180), 783-8. (doi:10.1038/nature06617)

31. Prochnik SE, Umen J, Nedelcu AM, Hallmann A, Miller SM, Nishii I, et al. 2010 Genomic analysis of organismal complexity in the multicellular green alga *Volvox carteri*. Science **329**(5988), 223-6. (doi:10.1126/science.1188800)

32. Suga H, Chen Z, de Mendoza A, Sebe-Pedros A, Brown MW, Kramer E, et al. 2013 The *Capsaspora* genome reveals a complex unicellular prehistory of animals. Nat. Commun. **4**, 2325. (doi:10.1038/ncomms3325)

33. Hanschen ER, Marriage TN, Ferris PJ, Hamaji T, Toyoda A, Fujiyama A, et al. 2016 The *Gonium pectorale* genome demonstrates co-option of cell cycle regulation during the evolution of multicellularity. Nat. Commun. **7**, 11370. (doi:10.1038/ncomms11370)

34. Paps J, Holland PWH. 2018 Reconstruction of the ancestral metazoan genome reveals an increase in genomic novelty. Nat. Commun. **9**(1), 1730. (doi:10.1038/s41467-018-04136-5)

35. Sebe-Pedros A, Ballare C, Parra-Acero H, Chiva C, Tena JJ, Sabido E, et al. 2016 The Dynamic Regulatory Genome of *Capsaspora* and the Origin of Animal Multicellularity. Cell **165**(5), 1224-37. (doi:10.1016/j.cell.2016.03.034)

36. Featherston J, Arakaki Y, Hanschen ER, Ferris PJ, Michod RE, Olson B, et al. 2018 The 4-Celled *Tetrabaena socialis* Nuclear Genome Reveals the Essential Components for Genetic Control of Cell Number at the Origin of Multicellularity in the Volvocine Lineage. Mol. Biol. Evol. **35**(4), 855-70. (doi:10.1093/molbev/msx332)

37. Lindsey CR, Knoll AH, Herron MD, Rosenzweig F. 2024 Fossil-calibrated molecular clock data enable reconstruction of steps leading to differentiated multicellularity and anisogamy in the Volvocine algae. BMC Biol. **22**, 79. (doi:10.1186/s12915-024-01878-1)

38. Boraas ME, Seale DB, Boxhorn JE. 1998 Phagotrophy by a flagellate selects for colonial prey: A possible origin of multicellularity. Evo. Ecol. **12**, 153-64. (doi:10.1023/A:1006527528063)

39. Becks L, Ellner SP, Jones LE, Hairston NG, Jr. 2010 Reduction of adaptive genetic diversity radically alters eco-evolutionary community dynamics. Ecol. Lett. **13**(8), 989-97. (doi:10.1111/j.1461-0248.2010.01490.x)

40. Ratcliff WC, Denison RF, Borrello M, Travisano M. 2012 Experimental evolution of multicellularity. Proc. Natl Acad. Sci. USA **109**(5), 1595-600. (doi:10.1073/pnas.1115323109)

41. Ratcliff WC, Herron MD, Howell K, Pentz JT, Rosenzweig F, Travisano M. 2013 Experimental evolution of an alternating uni- and multicellular life cycle in *Chlamydomonas reinhardtii*. Nat. Commun. **4**, 2742. (doi:10.1038/ncomms3742)

42. Herron MD, Borin JM, Boswell JC, Walker J, Chen ICK, Knox CA, et al. 2019 *De novo* origins of multicellularity in response to predation. Sci. Rep. **9**(1), 2328. (doi:10.1038/s41598-019-39558-8)

43. Sager R, Granick S. 1954 Nutritional control of sexuality in *Chlamydomonas reinhardi*. J. Gen. Physiol. **37**(6), 729-42. (doi:10.1085/jgp.37.6.729)

44. Flowers JM, Hazzouri KM, Pham GM, Rosas U, Bahmani T, Khraiwesh B, et al. 2015 Whole-Genome Resequencing Reveals Extensive Natural Variation in the Model Green Alga *Chlamydomonas reinhardtii*. Plant Cell **27**(9), 2353-69. (doi:10.1105/tpc.15.00492)

45. Gorman DS, Levine RP. 1965 Cytochrome f and plastocyanin: their sequence in the photosynthetic electron transport chain of *Chlamydomonas reinhardi*. Proc. Natl Acad. Sci. USA **54**(6), 1665-9. (doi:10.1073/pnas.54.6.1665)

46. Kilham SS, Kreeger DA, Lynn SG, Goulden CE, Herrera L. 1998 COMBO: a defined freshwater culture medium for algae and zooplankton. Hydrobiologia **377**, 147-59. (doi:10.1023/A:1003231628)

47. Du Q, Kawabe Y, Schilde C, Chen ZH, Schaap P. 2015 The Evolution of Aggregative Multicellularity and Cell-Cell Communication in the *Dictyostelia*. J. Mol. Biol. **427**(23), 3722-33. (doi:10.1016/j.jmb.2015.08.008)

48. Bonner JT. 2000 *First Signals: The Evolution of Multicellular Development*. Princeton University Press.

49. Tarnita CE, Taubes CH, Nowak MA. 2013 Evolutionary construction by staying together and coming together. J. Theor. Biol. **320**, 10-22. (doi:10.1016/j.jtbi.2012.11.022)

50. Bonner JT. 1998 The origins of multicellularity. Integrative Biology Issues News and Reviews. **1**(1): 27-36. (doi:10.1002/(SICI)1520-6602(1998)1:1<27::AID-INBI4>3.0.CO;2-6)
51. Zamora I, Feldman JL, Marshall WF. 2004 PCR-based assay for mating type and diploidy in *Chlamydomonas*. Biotechniques **37**(4), 534-6. (doi:10.2144/04374BM01)
52. Merchant SS, Prochnik SE, Vallon O, Harris EH, Karpowicz SJ, Witman GB, et al. 2007 The *Chlamydomonas* genome reveals the evolution of key animal and plant functions. Science **318**(5848), 245-50. (doi:10.1126/science.1143609)
53. Li H. 2013 Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv 1303.3997. (doi:10.48550/arXiv.1303.3997)
54. Garrison E, Marth G. 2012 Haplotype-based variant detection from short-read sequencing. arXiv 1207.3907. (doi:10.48550/arXiv.1207.3907)
55. Michod RE, Roze D. 2001 Cooperation and conflict in the evolution of multicellularity. Heredity **86**, 1-7. (doi:10.1046/j.1365-2540.2001.00808.x)
56. Lewontin R. 1970 The Units of Selection. Annu. Rev. Ecol. Syst **1**, 1-18.
57. Herron MD. 2021 What are the major transitions? Biol. Philos. 36, 2. (doi:10.1007/s10539-020-09773-z)
58. Andersson SG, Zomorodipour A, Andersson JO, Sicheritz-Ponten T, Alsmark UC, Podowski RM, et al. 1998 The genome sequence of *Rickettsia prowazekii* and the origin of mitochondria. Nature **396**(6707):133-40. (doi:10.1038/24094)
59. Wang Z, Wu M. 2015 An integrated phylogenomic approach toward pinpointing the origin of mitochondria. Sci. Rep. **5**, 7949. (doi:10.1038/srep0794)
60. Herron MD, Hackett JD, Aylward FO, Michod RE. 2009 Triassic origin and early radiation of multicellular volvocine algae. Proc. Natl Acad. Sci. USA **106**(9), 3254-8. (doi:10.1073/pnas.0811205106)
61. Ma X, Shi X, Wang Q, Zhao M, Zhang Z, Zhong B. 2023 A Reinvestigation of Multiple Independent Evolution and Triassic-Jurassic Origins of Multicellular Volvocine Algae. Genome Biol. Evol. **15**(8), evad142. (doi:10.1093/gbe/evad142)
62. Lindsey CR, Rosenzweig F, Herron MD. 2021 Phylotranscriptomics points to multiple independent origins of multicellularity and cellular differentiation in the volvocine algae. BMC Biol. **19**(1),182. (doi:10.1186/s12915-021-01087-0)
63. Knoll AH, Hewitt D. 2011 Phylogenetic, Functional, and Geological Perspectives on Complex Multicellularity. In The Major Transitions in Evolution Revisited, eds. Calcott B, Sterelny K, p. 251-70. (doi:10.7551/mitpress/9780262015240.003.0013)
64. Bonner JT. 2011 *Why size matters: from bacteria to blue whales*. Princeton University Press.
65. Harris EH. 2009 *The Chlamydomonas Sourcebook, Second Edition*. Academic Press.