Running title: Early HPI axis adaptation

Early life challenge enhances cortisol regulation in zebrafish larvae

Luis A. Castillo-Ramírez[1], Ulrich Herget[1,2], Soojin Ryu[1,3], Rodrigo J. De Marco[1,4]§

[1]Max Planck Institute for Medical Research, Jahnstr. 29, 69120 Heidelberg, Germany

[2]Division of Biology and Biological Engineering, California Institute of Technology, 1200

East California Boulevard, Pasadena, California 91125, USA

[3]Living Systems Institute, College of Medicine and Health, University of Exeter, Stocker

Road EX4 4QD Exeter, UK

[4]School of Biological and Environmental Sciences, Faculty of Science, Liverpool John

Moores University, Byrom Street, L3 3AF Liverpool, UK

ORCID

Luis A. Castillo-Ramírez https://orcid.org/0009-0005-4896-5586

Ulrich Herget https://orcid.org/0000-0002-4419-1414

Soojin Ryu https://orcid.org/0000-0002-7059-0160

Rodrigo J. De Marco https://orcid.org/0000-0002-2082-4264

§**Author for correspondence**:

De Marco, Rodrigo <R.J.DeMarco@ljmu.ac.uk>

ABSTRACT

The hypothalamic-pituitary-adrenal (HPA) axis in mammals and the hypothalamic-pituitary-interrenal (HPI) axis in fish are open systems that adapt to the environment during development. Little is known about how this adaptation begins and regulates early stress responses. We used larval zebrafish to examine the impact of prolonged forced swimming at 5 days post-fertilization (dpf), termed early-life challenge (ELC), on cortisol responses, neuropeptide expression in the nucleus preopticus (NPO), and gene transcript levels. At 6 dpf, ELC-exposed larvae showed normal baseline cortisol but reduced reactivity to an initial stressor. Conversely, they showed increased reactivity to a second stressor within the 30-minute refractory period, when cortisol responses are typically suppressed. ELC larvae had fewer corticotropin-releasing hormone (*crh*), arginine vasopressin (*avp*), and oxytocin (*oxt*)-positive cells in the NPO, with reduced crh and avp co-expression. Gene expression analysis revealed upregulation of genes related to cortisol metabolism (*hsd11b2*, *cyp11c1*), steroidogenesis (*star*), and stress modulation (*crh*, *avp*, *oxt*). These results suggest that early environmental challenge initiates adaptive plasticity in the HPI axis, tuning cortisol regulation to balance responsiveness and protection during repeated stress. Future studies should explore the broader physiological effects of prolonged forced swimming and its long-term impact on cortisol regulation and stress-related circuits.

Keywords: Zebrafish larvae, HPI axis, Cortisol regulation, Early life challenge, Stress response, Developmental programming

Summary: This study explores how early-life challenges in zebrafish affect stress responses and hormone regulation, offering insights into developmental adaptability and stress management mechanisms.

INTRODUCTION

The HPA/I axis functions as an adaptive system closely linked to bodily responses to environmental challenges that disrupt homeostasis and trigger protective adaptations (Wendelaar Bonga, 1997; Charmandari et al., 2005; Chrousos, 2009). As an open system, it continually adjusts to interactions between the organism and its environment throughout development, enhancing the ability to cope with stress. However, excessive or poorly timed stress can lead to high allostatic load, accelerating maladaptive processes (de Kloet et al., 2005; Danese and McEwen, 2012; Chen and Baram, 2016). While the effects of stress in adulthood are often transient, early-life stress can alter brain development, resulting in long-lasting behavioural changes (Danese and McEwen, 2012; Bick and Nelson, 2016; Teicher et al., 2016; Agorastos et al., 2019). These behavioural symptoms may manifest in childhood, adolescence, or later in adulthood, when the delayed effects of stress-induced alterations in brain development become apparent (Lupien et al., 2009). Additionally, human infants exposed to high cortisol levels *in utero* show elevated baseline cortisol and reduced cortisol responses to separation stress (O'Connor et al., 2013). Similarly, rodent studies show that early postnatal dexamethasone treatment reduces corticosterone responses to stress later in life (Felszeghy et al., 2000), while prolonged exposure to low-dose corticosterone further suppresses reactivity (Kinlein et al., 2015). These findings illustrate how both pre- and post-natal environments shape stress regulation and long-term HPA axis function, although the exact mechanisms remain unclear.

Larval zebrafish (*Danio rerio*) are a valuable model for exploring HPA/I axis adaptation during early development due to their genetic accessibility and external development, which facilitate the analysis of early-life stress (ELS) (Alsop and Vijayan, 2008; Alsop and Vijayan, 2009; Egan et al., 2009; Champagne and Richardson, 2013; Biran et al., 2015; Eachus et al., 2021; Tan et al., 2022). As a non-mammalian vertebrate model

(Grunwald and Eisen, 2002; Holtzman et al., 2016), zebrafish have become prominent in

ELS research, offering practical advantages such as small size, high reproductive capacity,

and low maintenance costs. Their external development and lack of parental care allow for

controlled experiments without the confounding effects of prenatal stress. The first stage of

stress processing in zebrafish occurs in the nucleus preopticus (NPO), which is homologous

to the mammalian paraventricular nucleus (PVN) (Herget et al., 2014). The process involves

key neuropeptides like corticotropin-releasing hormone (CRH), which stimulates

adrenocorticotropic hormone (ACTH) production in the pituitary, leading to glucocorticoid

(GC) release from the interrenal gland, functionally similar to the adrenal gland in mammals

(Alderman and Bernier, 2009). The increased secretion of GCs like cortisol after the onset of

stress, known as glucocorticoid reactivity (GC_R), is crucial for the organism's response to

challenge. Beyond their role in stress, GCs regulate a wide range of bodily functions,

including metabolism, maintenance of water and electrolyte balance, immunity, growth,

cardiovascular function, mood, cognition, reproduction, and development. Primarily

produced in the adrenal cortex, along with aldosterone and dehydro-epi-androsterone, all

derived from cholesterol, GCs are also synthesized at extra-adrenal sites such as the thymus,

brain, and epithelial barriers. These localized productions contribute to spatial specificity in

steroid actions, operating independently of systemic and stress-induced fluctuations (Munck

et al., 1984; Chrousos, 1998; de Kloet et al., 1998; McEwen, 2007; Timmermans et al.,

2019; de Kloet and Joëls, 2023). Together, these features make zebrafish a powerful handle

for investigating the molecular mechanisms by which early environments influence brain

development and behaviour, as well as for exploring potential therapeutic interventions for

stress-related disorders (Veenstra-VanderWeele and Warren, 2015; Bick and Nelson, 2016).

By 4-6 dpf, when the HPI axis becomes functional (Alsop and Vijayan, 2008;

Alderman and Bernier, 2009; De Marco et al., 2016), zebrafish larvae in laboratory settings

primarily encounter visual cues from conspecifics and mechanosensory inputs from other

larvae, unless raised in isolation, along with self-generated motion. Temperature,

illumination, and food availability can be tightly controlled, creating an environment with

limited variability, ideal for analyzing early HPI axis calibration to external stimuli. For

example, their response to water currents, where they adjust swimming to match flow

strength, allows for environments with varying physical challenges. Zebrafish larvae adapt

to water vortex flows through rheotaxis, orienting themselves against the current, which

imposes high energy demands and activates the HPI axis. Controlled vortex flows at specific

revolutions per minute (rpm) induce a rapid and significant rise in cortisol levels, unlike

conditions without flow, establishing vortex flow as an effective stressor (Castillo-Ramírez

et al., 2019). While the metabolic or cardiorespiratory effects of vortex exposure have not

yet been measured, the cortisol increase indicates significant energy demands and

physiological stress. Additionally, vortex flows offer several advantages as a stressor,

minimizing confounding variables such as salt concentration or pH changes and utilizing an

innate, reproducible behaviour, reducing experimental variability. Consequently, vortices

and less predictable water motions have been used to study stress responses in zebrafish

larvae (De Marco et al., 2014; vom Berg-Maurer et al., 2016; Ryu and De Marco, 2017;

Castillo-Ramírez et al., 2019; Langebeck-Jensen et al., 2019; Herget et al., 2023; Castillo-

Ramírez et al., 2024). Notably, as vortex strength (rpm) increases, cortisol levels rise

proportionally, as does larval swimming effort to counteract the flow (Castillo-Ramírez et

al., 2019). This quantifiable relationship between vortex intensity and GC_R allows precise

categorization of stress levels, making vortex flows ideal for repeated stress assays. Using

these highly controlled water vortices, we established a high-throughput forced swim test for

larval zebrafish. This test revealed that early-life challenge (ELC), in the form of sustained

involuntary swimming, triggers prolonged HPI axis activation and alters the cortisol

response to a single instance of either homotypic or heterotypic stress 1-4 days post-ELC.

ELC also enhances spontaneous activity, reduces startle reactivity, and improves energy

efficiency during rheotaxis (Castillo-Ramírez et al., 2019). The mechanisms underlying

these effects remain unknown.

To address this gap, we set out to characterize features of the HPI axis and cortisol

response in zebrafish larvae following ELC induced by vortex flows. First, we re-examined

how ELC in the form of sustained involuntary swimming at 5 dpf affects cortisol response

dynamics following a 3-minute vortex at 6 dpf. The rationale for selecting the 5-6 dpf time

window is based on a key observation: the machinery for GC synthesis and signalling

becomes fully functional around hatching, as evidenced by significant increases in gene

expression (Alsop and Vijayan, 2008; Alderman and Bernier, 2009). This corresponds with

the gradual rise in whole-body cortisol levels between 2 and 8 dpf. Importantly, cortisol

responsiveness to vortex-induced stress peaks at 6 dpf (Castillo-Ramírez et al., 2019),

indicating increased developmental sensitivity of the HPI axis at 5–6 dpf. We then examined

how ELC impacts the expression and co-expression of *crh*, *avp* (arginine vasopressin), and

oxt (oxytocin) in the larval NPO. Focusing on these peptides is justified by their established

roles in mammals and their conserved functions in stress response and homeostasis in

zebrafish. CRH initiates the body's response to stress and influences autonomic and

behavioural responses; AVP plays a crucial role in stress response, social behaviour, and

cardiovascular functions, including osmoregulation and blood pressure regulation; and OXT

is known for its role in modulating stress and anxiety, social bonding, and reproductive

behaviour (Swanson and Sawchenko, 1983; Herman and Cullinan, 1997; Eaton et al., 2008).

Assuming that the expression and coexpression profile of NPO cells can undergo stress-

derived plasticity, as occurs with PVN cells (Kiss, 1988; Harbuz and Lightman, 1989;

Swanson, 1991; Kurrasch et al., 2009), we predicted that ELC may reduce the degree of

expression and coexpression of key NPO neuropeptides involved in HPI axis activation.

Next, we explored the effect of ELC at 5 dpf on the transcript levels of genes

involved in HPI axis activation, cortisol synthesis and inactivation, and GR signalling, both

under baseline conditions and after vortex exposure at 6 dpf. These genes included *hsd11b2* (11-hydroxysteroid dehydrogenase type 2), essential for cortisol inactivation (Krozowski, 1999; Alderman and Vijayan, 2012; Theodoridi et al., 2021); *cyp11c1* (11β-hydroxylase), critical for the synthesis of 11-ketotestosterone and cortisol (Nelson et al., 2013; Tokarz et al., 2015); and *star* (steroidogenic acute regulatory protein), involved in cholesterol transport (Stocco, 2000). We also analysed *crh*, *avp*, *oxt*, *crhr1* (corticotropin-releasing hormone receptor 1), *pomca* (pro-opiomelanocortin, the ACTH precursor), *mc2r* (adrenocorticotropic hormone receptor), *nr3c2* (mineralocorticoid receptor, MR), *nr3c1* (glucocorticoid receptor, GR), and *fkbp5* (FK506 binding protein 5), a stress-responsive regulator of GR (Sinars et al., 2003; Zannas et al., 2016; Hartmann et al., 2021). To explore potential ELC-derived patterns in transcript abundance, we employed a whole-body RNA approach without isolating changes specific to individual HPI axis components (NPO, pituitary, interrenal gland). This exploratory approach is inherently limited, as some target genes are ubiquitously expressed while others are more restricted, with expression often extending beyond the HPI axis. Despite this limitation, the approach allows us to test whether the overall transcript profile reflects increased cortisol inactivation under both baseline and stress conditions, and whether it indicates an enhanced capacity for cortisol synthesis and stress modulation in larvae previously exposed to ELC. Such patterns would be consistent with an early challenge that imposed prolonged HPI axis activity and elevated cortisol levels.

Lastly, we examined how ELC at 5 dpf affects the cortisol response of 6 dpf larvae to a second vortex applied within the 30-minute refractory period following an initial exposure (Castillo-Ramírez et al., 2024). The refractory period reflects the action of the glucocorticoid feedback loop, during which cortisol levels typically return to baseline and are unresponsive to additional stress. This period has only recently been identified in zebrafish larvae and represents a critical window to assess altered cortisol regulation. In previous work (Castillo-Ramírez et al., 2024), we showed that cortisol levels rise after a 3-minute vortex and return

to baseline within 30–40 minutes, depending on the intensity of the vortex. A second vortex

administered within 30 minutes of the first fails to elevate cortisol, but blocking GR disrupts

this refractory period, indicating GR-dependent feedback. We hypothesized that if ELC at 5

dpf enhances cortisol regulation, larvae exposed to ELC may show an altered response to

repeated stress during the refractory period, potentially amplifying the cortisol response to a

second vortex. The results are discussed in the context of how ELC influences the initial

adjustments of cortisol responsiveness in zebrafish larvae when they encounter

environmental challenges for the first time after being in a largely constant environment.

This early-life impact on HPI axis activity further supports the suitability of larval zebrafish

as a model for studying the developmental programming of HPA/I axis function.

RESULTS

Forced swimming at 5 dpf reduces a larva's cortisol response to a subsequent exposure

to the same type of stress

We first exposed 5 dpf larvae to either vortex flows for 9 hours (ELC larvae) or equal

handling without vortex exposure (control larvae) (see Materials and Methods for details).

At 6 dpf, we compared the basal whole-body cortisol levels (Fig. 1A) and the cortisol time

course following a 3-minute vortex (Fig. 1B) between ELC and control larvae. Consistent

with previous data (Castillo-Ramírez et al., 2019), mean baseline whole-body cortisol levels

at 6 dpf were similar between control and ELC larvae (Fig. 1C, Mann-Whitney test: U=17,

p=0.90; N=6 per group), although cortisol levels in ELC larvae were less variable

(coefficient of variation: Control, 46.8%, ELC, 18.1%). The 3-minute vortex transiently

increased whole-cortisol in both groups, yet ELC larvae showed reduced GC_R to the vortex

(Fig. 1D, two-way ANOVA on square root-transformed data: group: $F_{(1,50)}$=35.8,

p<0.0001; time: $F_{(4,50)}$=202.9, p<0.0001; group x time: $F_{(4,50)}$=15.4, p<0.0001;

Bonferroni's tests: 10': $p<0.0001$, 20': $p<0.0001$, 30': $p=0.0032$, 40': $p=0.125$, 60': $p=0.99$;

N=6 per group). These results confirm that prolonged forced swimming at 5 dpf reduces a

larva's cortisol response to subsequent homotypic stress, likely due to adjustments in the

mechanisms regulating HPI axis reactivity. This raises questions about how these changes

relate to the expression and co-expression levels of NPO neuropeptides, which may

influence downstream HPI axis activity.

NPO imaging shows expression and coexpression of key neuropeptides linked to early life challenge

To examine whether the altered cortisol response to homotypic stress observed in ELC

larvae at 6 dpf (Fig. 1) is associated with changes in the expression of *crh*, *avp*, and *oxt* in

the larval NPO, we employed multicolor fluorescent *in situ* hybridization at 6 dpf following

the ELC protocol implemented at 5 dpf (see Materials and Methods). We compared the

numbers of *crh*-, *avp*-, or *oxt*-positive cells in the NPO of unstressed subjects from both

groups, ELC and control larvae. Quantification of cell numbers revealed that, compared to

control larvae, ELC larvae had fewer *crh*-positive cells within the NPO, as well as fewer

oxt-positive and fewer *avp*-positive cells (Fig. 2A, Mann-Whitney tests: *oxt*: U=167, $p=0.03$;

avp: U=112, $p=0.0004$; *crh*: U=85, $p<0.0001$; Group sizes: ELC, N=20, Control, N=27), in

line with the idea that ELC can alter the expression of NPO neuropeptides. Additionally, we

examined the degree of coexpression between sets of two neuropeptides in the same cell in

ELC and control larvae using systematic cell by cell comparisons of pairwise combinatorial

ISH staining of *avp*, *crh*, and *oxt* in the larval NPO (Fig. 2B). In both groups, ELC and

control larvae, coexpression was consistently high for the combination of *avp* and *crh*, low

or moderate for the *crh* and *oxt* combination, and absent or rare for the *oxt* and *avp*

combination, in line with previous data (Herget and Ryu, 2015). Also, coexpression was

consistently lower in ELC larvae for the combination *avp* and *crh* staining compared to

control larvae (Fig. 2C, Mann-Whitney tests: *crh + avp*: U=787.5, *p*=0.02; *avp + crh*:

U=686, *p*=0.002; Group sizes: ELC, N=20, Control, N=27). These results indicate that the

changes in GC_R observed at 6 dpf in ELC larvae correlate with altered levels of both the

expression and coexpression of *avp*, *crh*, and *oxt* in the larval NPO.

Early life challenge correlates with changes in the transcript abundance of genes

involved in cortisol metabolism, steroidogenesis, and stress modulation

Next, we explored whether ELC at 5 dpf is associated with changes in the transcript levels

of *hsd11b2*, *cyp11c1*, *star*, *crh*, *avp*, *oxt*, *pomca*, *mc2r*, *crhr1*, *nr3c2*, *nr3c1*, and *fkbp5* at 6

dpf, both at baseline and following a 3-minute vortex. These genes play roles in cortisol

synthesis, HPI axis activation, cortisol inactivation, and GR signalling. We employed RT-

qPCR to quantify total RNAs from both ELC and control larvae, measuring the fold change

in transcript abundance of the 12 genes (Table 1) relative to reference samples from control

larvae. Samples were taken at baseline and at 30, 60, and 120 minutes after exposure to the

3-minute vortex (see Materials and Methods). The results indicated that ELC at 5 dpf was

associated with altered expression of several genes by 6 dpf (Fig. 3). Compared to controls,

ELC larvae showed consistently higher expression of *hsd11b2* under both baseline and

stress conditions (Fig. 3A, two-way ANOVA on log-transformed data: group: $F_{(1,32)}=53.4$,

p<0.0001; time: $F_{(3,32)}=0.06$, *p*=0.98; group x time: $F_{(3,32)}=0.25$, *p*<0.86; Bonferroni's

tests: basal: *p*=0.007, 30': *p*=0.003, 60': *p*=0.0006, 120': *p*=0.013; N=5 per group). Post-

stressor, ELC larvae showed elevated expression of *cyp11c1* and *star* (Fig. 3B,C, two-way

ANOVAs; *cyp11c1*: group: $F_{(1,32)}=20.8$, *p*<0.0001; time: $F_{(3,32)}=0.99$, *p*=0.41; group x

time: $F_{(3,32)}=1.15$, *p*=0.34; Bonferroni's tests: 60': *p*=0.007; *star* log-transformed: group:

$F_{(1,32)}=26.2$, *p*<0.0001; time: $F_{(3,32)}=2.02$, *p*=0.13; group x time: $F_{(3,32)}=1.41$, *p*=0.26;

Bonferroni's tests: 60': p=0.014, 120': p=0.008). Increases in *pomca*, *crh*, and *oxt*

expression were also observed post-vortex (Fig. 3D-F; *pomca*: group: $F_{(1,32)}=4.99$,

$p=0.033$; time: $F_{(3,32)}=0.45$, $p=0.72$; group x time: $F_{(3,32)}=2.26$, $p=0.10$; Bonferroni's

tests: 120': $p=0.038$; *crh* log-transformed: group: $F_{(1,32)}=6.41$, $p=0.017$; time:

$F_{(3,32)}=1.05$, $p=0.38$; group x time: $F_{(3,32)}=0.76$, $p=0.52$; *oxt* sqrt-transformed: group:

$F_{(1,32)}=7.24$, $p=0.011$; time: $F_{(3,32)}=2.65$, $p=0.07$; group x time: $F_{(3,32)}=1.23$, $p=0.32$).

Expression of *avp* and *nr3c2* was influenced by both group and time factors (Fig. 3G,H; *avp*:

group: $F_{(1,32)}=6.9$, $p=0.013$; time: $F_{(3,32)}=4.98$, $p=0.006$; group x time: $F_{(3,32)}=0.23$,

$p=0.88$; *nr3c2*: group: $F_{(1,32)}=4.5$, $p=0.042$; time: $F_{(3,32)}=9.42$, $p=0.0001$; group x time:

$F_{(3,32)}=2.77$, $p=0.06$). Although not statistically significant, *nr3c1* levels tended to increase

in ELC larvae post-stressor (Fig. 3I, group: $F_{(1,32)}=2.76$, $p=0.11$; time: $F_{(3,32)}=2.2$,

$p=0.11$; group x time: $F_{(3,32)}=1.2$, $p=0.32$). For *crhr1* and *mc2r*, only time influenced

expression (Fig. 3J,K; *crhr1*: group: $F_{(1,32)}=0.45$, $p=0.51$; time: $F_{(3,32)}=3.9$, $p=0.02$; group

x time: $F_{(3,32)}=1.1$, $p=0.37$; *mc2r*: time: group: $F_{(1,32)}=0.06$, $p=0.81$; time: $F_{(3,32)}=6.2$,

$p=0.002$; group x time: $F_{(3,32)}=0.31$, $p=0.82$). Lastly, *fkbp5* transcript levels rose initially

and declined later in both ELC and control larvae (Fig. 3L, log-transformed: group:

$F_{(1,32)}=1.7$, $p=0.21$; time: $F_{(3,32)}=45.7$, $p<0.0001$; group $\times$ time: $F_{(3,32)}=4.6$, $p=0.009$). In

summary, we observed differences in transcript levels between control and ELC larvae at 6

dpf. Expression of *hsd11b2* was elevated in both baseline and stress conditions, while

cyp11c1 and *star* increased specifically after vortex exposure. Stressed ELC larvae also

showed higher levels of *pomca*, *crh*, and *oxt*. Expression of *avp* and *nr3c2* was influenced

by both group and time factors, while trends in *nr3c1*, *crhr1*, and *mc2r* were primarily

driven by time following stressor exposure. Finally, *fkbp5* levels initially increased and then

declined in both groups (Fig. 3M).

Early life challenge modifies glucocorticoid reactivity to a second vortex during the

refractory period

A reduced capacity of the NPO to influence HPI axis reactivity, along with enhanced

cortisol inactivation following ELC, may partly account for the diminished GC_R observed

after the initial post-ELC vortex. Furthermore, increased cortisol synthesis capacity and the

availability of stress modulators could influence GC_R dynamics during repeated homotypic

stress, potentially affecting the response to a second vortex. To investigate this possibility,

we compared whole-body cortisol levels between ELC and control larvae at 6 dpf after two

consecutive 3-minute vortices, applied within the 30-minute refractory period following the

initial exposure (Castillo-Ramírez et al., 2024) (Fig. 4A). ELC larvae showed increased

whole-body cortisol in response to the second vortex applied during this refractory period,

whereas control larvae did not (Fig. 4B, Unpaired two-tailed *t*-test: $t(10)=13.9$, $p<0.0001$;

N=6 per group). Both ELC and control larvae had similar baseline cortisol levels (Fig. 4C,

One-sample *t*-test against fold change of '1': $t(5)=0.9$, $p=0.40$; N=6 per group; data from

Fig. 1C). However, ELC larvae showed a reduced cortisol response to the first vortex as

well as an enhanced response to the second vortex (Fig. 4D, two-way ANOVA: group:

$F(1,20)=9.8$, $p=0.0052$; vortex: $F(1,20)=208.1$, $p<0.0001$; group $\times$ vortex: $F(1,20)=149.5$,

$p<0.0001$; Bonferroni's tests: vortex #1: $p<0.0001$, vortex #2: $p<0.0001$; N=6 per group;

data from Fig. 1D and Fig. 4B). As a result, control and ELC larvae showed distinct cortisol

reactivity to the second vortex. While ELC larvae showed a GC_R to the second vortex that

was 91% of their response to the first vortex, control larvae showed only 18% (Fig. 4E,

Unpaired two-tailed *t*-test; $t(10)=20.7$, $p<0.0001$; N=6 per group; data from Fig. 4D). These

findings indicate that prolonged forced swimming at 5 dpf improves the ability of larvae at 6

dpf to regulate cortisol levels and adjust HPI axis activity in response to stress, enhancing

their capacity to cope with repeated exposure to the same stressor.

DISCUSSION

Here we provide strong evidence for extended effects of early forced swimming on cortisol

regulation and stress responsiveness in zebrafish larvae. We observed that ELC at 5 dpf

resulted in significant alterations in cortisol reactivity to subsequent stress events.

Specifically, ELC larvae had similar baseline cortisol levels compared to controls but

showed reduced GC_R to a single vortex at 6 dpf. This finding is consistent with previous

data (Castillo-Ramírez et al., 2019) and reveals adaptive adjustments in stress axis function

following prolonged early-life stressor exposure. Notably, while ELC larvae showed

reduced GC_R to the initial stressor, they had increased cortisol responsiveness to a second

stress event occurring within the 30-minute refractory period following the first vortex

(Castillo-Ramírez et al., 2024). This dual response pattern suggests that ELC larvae have

enhanced stress responsiveness during homotypic stress, potentially due to adaptive changes

in HPI axis signalling and cortisol metabolism.

Our findings are consistent with research in fish and other vertebrates showing that

early life stress and GC manipulation can have lasting effects on HPA/I axis function and

stress responsiveness. In larval zebrafish, optogenetic techniques enable non-invasive

control of endogenous GC levels (De Marco et al., 2013; 2016). Transgenic larvae

expressing Beggiatoa photoactivated adenylyl cyclase (bPAC) (Ryu et al., 2010; Stierl et al.,

2011) under a StAR promoter can induce light-dependent activation of interrenal gland

steroidogenic cells, resulting in elevated cortisol levels (Gutierrez-Triana et al., 2015).

Persistent activation of these cells during early development leads to chronically high

cortisol levels and a diminished cortisol response to later stressors (Nagpal et al., 2024).

Additionally, optogenetically elevated GC levels early in life disrupt hypothalamic

neurogenesis, causing precocious development, failed maturation, and impaired feeding and

growth (Eachus et al., 2024). Chronic stress induced by random shocks in developing

zebrafish also causes sustained high cortisol levels and increased expression of

glucocorticoid and mineralocorticoid receptors, which correlate with increased anxiety-like

behaviour and elevated cortisol levels later in life (Chin et al., 2022). Zebrafish exposed to

air-exposure stress at different developmental stages showed increased cortisol levels,

altered stress responses, and changes in ion concentrations (Hare et al., 2021). Similar trends

are observed in other species. In rainbow trout, chronic cortisol intake and daily stress

exposure reduce GC_R (Barton et al., 1987). Rodent studies show that neonatal rats treated

with dexamethasone show reduced GC_R in response to cold and restraint stress later in life

(Felszeghy et al., 2000), while mice subjected to prolonged low-dose corticosterone

exposure show suppressed GC_R, adrenal atrophy, and decreased CRH mRNA levels in the

PVN (Kinlein et al., 2015). In humans, infants exposed to high cortisol levels *in utero* show

elevated baseline cortisol and reduced GC_R in response to separation stress (O'Connor et al.,

2013). Collectively, these observations underscore the complex interplay between ELC, GC

regulation, and stress responsiveness early in life. However, the mechanisms through which

ELC affects HPA/I axis function and leads to long-term changes in stress response patterns

are still largely unclear, particularly regarding the initial phase of this interaction.

The HPA axis regulates its stress-induced activation through negative feedback

mechanisms (Dallman and Yates, 1969; Dallman et al., 1994; Dallman, 2005; Shipston,

2022). In mammals, this feedback acts on the PVN, where GCs suppress the expression of

stress modulators like *crh* (Malkoski et al., 1999). Given the structural and

chemoarchitectural similarities between the mammalian PVN and the zebrafish NPO

(Herget et al., 2014), it is likely that the NPO is subject to similar GC-mediated regulatory

feedback (Castillo-Ramírez et al., 2024). Supporting this, zebrafish with disrupted GC

signalling show increased numbers of *crh*-expressing cells in the NPO (Ziv et al., 2013;

Facchinello et al., 2017). In mammals, regulation of the PVN involves not only the

expression of individual modulators like *crh* but also the co-expression of multiple

neuropeptides within the same cells (Sawchenko et al., 1984; Whitnall et al., 1985; Simmons

and Swanson, 2009; Biag et al., 2012). However, in the zebrafish NPO, neuropeptides tend

to be produced by distinct, tightly packed cells, with limited co-expression, although certain combinations show occasional to moderate co-expression (Herget and Ryu, 2015). Given that stress can induce plastic changes in the expression and co-expression of neuropeptides in the PVN (Kiss, 1988; Harbuz and Lightman, 1989; Swanson, 1991; Kurrasch et al., 2009), we hypothesized that ELC might similarly alter the expression and co-expression profiles of key neuropeptides in the NPO. Our findings show that ELC larvae had fewer *crh*-, *avp*-, and *oxt*-positive NPO cells than controls, with a marked reduction in *crh* and *avp* co-expression. These results suggest that, analogous to the mammalian PVN (Ziv et al., 2013; Herget et al., 2014; Facchinello et al., 2017), the zebrafish NPO serves as a critical site for GC feedback regulation. Moreover, they indicate that ELC induces plasticity in neuropeptide expression within the developing NPO, potentially affecting downstream HPI axis activity. The reduced number of *crh*- and *avp*-positive cells in the NPO may contribute to the reduced GC_R observed following the initial stress exposure in ELC larvae.

While the expression profile of steroidogenic enzymes involved in GC biosynthesis has been characterized in developing zebrafish (Weger et al., 2018), our data revealed group differences between ELC and control larvae. Specifically, ELC larvae showed upregulation of *hsd11b2* under both baseline and stress conditions, as well as increased expression of *cyp11c1*, *star*, *pomca*, *crh*, *oxt*, *avp*, and *nr3c2* following a 3-minute vortex. Additionally, transcripts for *avp*, *nr3c2*, *crhr1*, *mc2r*, and *fkbp5* showed time-dependent changes post-stressor. Interestingly, previous studies in fish have linked whole-body cortisol levels with *hsd11b2* and *cyp11c1* mRNA expression (Tsalafouta et al., 2014), while prolonged cortisol release has been observed in zebrafish lacking functional Hsd11b2 (Theodoridi et al., 2021). Notably, although ELC larvae showed fewer *crh*-positive cells in the NPO, their whole-body *crh* levels increased after acute stress. This suggests that ELC may influence the maturation of stress-related circuits involving NPO *crh*-positive cells (Bolton et al., 2022). While whole-body qPCR cannot isolate transcript changes specific to the HPI axis, given that some

genes are ubiquitously expressed while others have restricted but broader distribution, the

overall transcript pattern in 6 dpf ELC larvae points to enhanced cortisol inactivation under

both baseline and stress conditions. Additionally, the upregulation of steroidogenic enzymes

and stress modulators indicates an increased capacity for cortisol synthesis and regulation

during stress. These transcriptional changes, alongside altered *avp*, *crh*, and *oxt* expression

and co-expression patterns in the NPO, suggest that ELC enhances cortisol regulation,

preventing excessive cortisol release and promoting an appropriate response to homotypic

stress (Fig. 5).

Our findings contribute to a better understanding of how early-life environments

shape GC_R, NPO plasticity, and GC pathways during the early stages of HPI axis

development. They also highlight the value of larval zebrafish as a model organism for

studying the developmental programming of HPI axis function (Weinstock, 2008; Glover et

al., 2010; Moisiadis and Matthews, 2014; de Abreu et al., 2021; Eachus et al., 2021;

Swaminathan et al., 2023). While this study focused primarily on cortisol dynamics, future

research should compare forced swimming in vortex conditions to a larva's routine

swimming patterns, investigating the broader physiological effects of vortex-induced stress,

which offers an excellent tool for high-throughput testing. Additional studies could evaluate

measures such as whole-body lactate and cholesterol levels, cardiac activity, and anaerobic

metabolism to provide a more comprehensive understanding of the physiological demands

imposed by vortex exposure. Moreover, examining how prolonged early-life forced

swimming affects cortisol production in interrenal cells (Wilson et al., 2016), the maturation

of stress-related circuits in the NPO, and stress responses in later life stages could reveal

long-lasting impacts on cortisol regulation and stress-related pathways. Such research would

clarify how vortex exposure as an early-life challenge influences not only immediate stress

responses but also long-term physiological and behavioural outcomes, offering valuable

insights into stress-related developmental programming.

MATERIALS AND METHODS

Zebrafish husbandry, handling, and experimental unit

Zebrafish breeding and maintenance were conducted under standard conditions (Westerfield,

2000). Groups of thirty wild-type eggs (cross of AB and TL strains, AB/TL) were collected

in the morning, and the embryos were raised on a 12:12 light/dark cycle at 28°C in 35 mm

Petri dishes with 5 ml of Embryo-medium2 (E2). The E2 medium (0.5x E2, 1 L) consisted

of 5 mM NaCl, 0.25 mM KCl, 0.5 mM $MgSO_4 \times 7 H_2O$, 0.15 mM KH_2PO_4, 0.05 mM

Na_2HPO_4, 0.5 mM $CaCl_2$, and 0.71 mM $NaHCO_3$. At 3 dpf, the E2 medium was renewed,

and chorions and debris were removed from the dishes. Experiments were performed with 5-

6 dpf larvae. In all experiments, each experimental unit (replicate) comprised a group of

thirty larvae, maintained in a 35 mm Petri dish. All dishes were kept under identical

conditions in the incubator on top of the stirrer plate (see below) to ensure no perturbation.

Zebrafish experimental procedures were conducted according to the guidelines of the

German animal welfare law and were approved by the local government

(Regierungspräsidium Karlsruhe; G-29/12).

Water vortex flows

We used water vortices in a high-throughput manner to induce rheotaxis and cortisol

elevation (Castillo-Ramírez et al., 2019). Groups of thirty larvae (either 5 or 6 dpf,

depending on the experiment) were placed in 35 mm Petri dishes containing 5 ml of E2

medium (experimental units) and exposed to controlled vortices generated by the spinning

movements of small encapsulated (coated) magnetic stir bars (6 x 3 mm, Fisherbrand,

#11888882, Fisher Scientific, Leicestershire, UK) inside the dishes. Each Petri dish, either

containing a stir bar or not (depending on the experimental group), was always positioned on

a magnetic stirrer plate (Variomag, Poly 15; Thermo Fisher Scientific, Leicestershire, UK)

and maintained at 28°C in an incubator (RuMed 3101, Rubarth Apparate GmbH, Laatzen,

Germany). Vortex flows were induced by the movements of the stir bars, driven by

magnetic field inversions of the stirrer plates set at 330 revolutions per minute (rpm). The

magnetic field inversions alone did not affect whole-body cortisol levels in larval zebrafish,

and we limited the speed to 330 rpm to avoid potential ceiling effects in vortex-dependent

cortisol increases, as previously reported (Castillo-Ramírez et al., 2019). At 5 dpf, larvae

were either exposed to 9 hours of continuous vortices (ELC larvae) or to magnetic field

inversions without vortices (Control larvae), as no stir bars were present in the control

group's Petri dishes. The 9-hour exposure period was designed to maximize the duration of

vortex exposure while adhering to the light/dark cycle. Larvae at 5 dpf exposed to vortex

flows for 9 hours show elevated whole-body cortisol levels, peaking shortly after vortex

initiation and remaining high for up to four hours compared to controls, i.e., unexposed

larvae that are handled similarly but without vortex flows in their medium. By six hours

after vortex onset, cortisol levels in both exposed and control larvae are comparable.

Additionally, exposed larvae consistently engage in positive rheotaxis throughout the vortex

exposure, as indicated by their body angle relative to the incoming current, even 8.5 hours

after the vortex began (Castillo-Ramírez et al., 2019). At 6 dpf, larvae previously exposed

(ELC) or not exposed (Control) to the 9-hour forced swimming period at 5 dpf were used for

the following analyses: 1) measurement of baseline whole-body cortisol levels (Fig. 1C),

extraction of total RNA for quantification of baseline transcript abundance of 12 genes, or

whole-mount fluorescent *in situ* hybridization and immunohistochemistry to assess cell

number and coexpression (Fig. 2); 2) measurement of whole-body cortisol at 10, 20, 30, 40,

and 60 minutes after a single 3-minute vortex (330 rpm) (Fig. 1D), or total RNA extraction

to quantify transcript abundance of the 12 genes post-vortex (Fig. 3); and 3) measurement of

whole-body cortisol after a second 3-minute vortex administered 30 minutes after the first

(Fig. 4). The selected 10, 20, 30, 40, and 60-minute intervals (Fig. 1D) effectively captured

cortisol dynamics following the initial stressor, providing a broad range to account for

variability in stress response timing. This design ensured experimental rigor while adhering

to ethical guidelines for subject use. At 6 dpf, larvae were immobilized in ice water before

cortisol detection or total RNA extraction (see below).

Whole-body cortisol

At 6 dpf, groups of 30 ELC or Control larvae (replicates), either unexposed to the 3-minute

vortex (baseline measurement) or exposed to one or two vortices applied within 30 minutes,

were immobilized in ice water (<2°C) to minimize stress. Excess water was removed, and

larvae were immediately frozen in an ethanol/dry-ice bath to ensure rapid euthanasia and

preservation. Samples were then stored at −20°C for subsequent cortisol extraction, which

took place between 10:30 and 11:30 hours. The procedures for cortisol measurements and

the homemade ELISA were as previously described (Yeh et al., 2013). The homemade

ELISA used for cortisol detection was validated through assessments of intra-assay and

inter-assay precision, recovery rates, and cross-reactivity. Results across batches and

sessions were consistent, with no significant differences in baseline cortisol levels, further

confirming the reliability of the assay. Full details of the validation process, including

comparisons with a commercial kit, can be found in Castillo-Ramírez et al. (2024).

Quantitative Real-Time PCR (qPCR)

Total RNA was extracted from groups of thirty larvae, as previously described for cortisol

extraction. RNA isolation was performed using the RNeasy Micro Kit (Qiagen, Hilden,

Germany), according to the manufacturer's protocol. The purity and concentration of RNA

were determined using a NanoDrop spectrophotometer (Thermo Fisher Scientific), and the

integrity of the RNA was confirmed using an RNA 6000 Pico Kit chip (Agilent

Technologies) on the Agilent 2100 Bioanalyzer. Only samples with RNA Integrity Numbers

(RIN) above 7.0 were used in subsequent qPCR analysis. qPCR reactions were carried out

using the Power SYBR Green RNA-to-Ct 1-Step Kit (Thermo Fisher Scientific,

Leicestershire, UK) on a 7500 Real-Time PCR system (Applied Biosystems). RNA was

reverse transcribed and amplified in a single-step reaction. Primers for the target genes

(listed in Table 1) were either designed in-house using Primer3 software or obtained from

published sources. Primer efficiency was confirmed using a standard curve approach,

ensuring that only primers with an efficiency of 90% to 110% were used in further

experiments. The gene Elongation Factor 1-alpha (EF1α) (F:

CTGGAGGCCAGCTCAAACGT   R: ATCAAGAAGAGTAGTACCGCTA) was selected

as the reference gene based on its known stability in zebrafish across various tissues and

treatments (McCurley and Callard, 2008). To ensure that EF1α maintained stable expression

in our experimental setup, the cycle threshold (Ct) values of EF1α were examined across all

experimental groups. Consistency in Ct values across conditions indicated stable expression

of EF1α, validating its use as a reliable internal control for normalizing the expression of

target genes. Minimal variation in Ct values across all treatments and replicates

demonstrated that the gene was unaffected by experimental conditions. Each 20 µL qPCR

reaction contained 10 µL of SYBR Green Master Mix, 0.5 µL of forward and reverse

primers (10 µM), 0.4 µL of ROX reference dye, and 100 ng of total RNA. The thermal

cycling program included an initial reverse transcription step at 48°C for 30 min, followed

by denaturation at 95°C for 10 min, and 40 cycles of 95°C for 15 s and 60°C for 1 min.

Relative gene expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method (Livak and

Schmittgen, 2001), with EF1α serving as the reference gene for normalization. The fold

change in gene expression was calculated by comparing the treated samples to the control

group, and statistical analyses were performed on the ΔCt values to determine significance.

Whole-mount fluorescent *in situ* hybridization, immunohistochemistry, and imaging

Whole-mount fluorescent *in situ* hybridization and immunohistochemistry were performed

as described elsewhere (Kastenhuber et al., 2010; Lauter et al. 2011), using riboprobes for

avp (Eaton et al., 2008), *oxt* (Unger and Glasgow, 2003), and *crh* (Löhr et al., 2009), and a

primary chicken antibody labeling GFP (1:500, Abcam), with a secondary anti-chicken

Alexa 488 antibody (1:1000, invitrogen). For imaging, specimens were cleared in 80%

glycerol (Gerbu Biotechnik GmbH, Heidelberg, Germany) in PBS for 1 h. Confocal stacks

were recorded using a Leica SP5 confocal microscope (Leica Microsystems GmbH,

Wetzlar, Germany) with a Nikon 20x glycerol objective (Nikon, Tokyo, Japan). Each

channel was recorded sequentially to reduce interfering signals from overlapping emission

spectra. Zoom, dimensions, gain, offset, average, and speed were adjusted for each stack to

obtain the optimal image quality of the desired volume. Stacks were evaluated using Amira

5.4 (Thermo Fisher Scientific, Leicestershire, UK) to create maximum intensity projections,

which were spatially restricted to the volume of interest, excluding signals from planes in

front or behind. Brightness and contrast were adjusted for each channel.

General design and statistical analysis

Cortisol and RT-qPCR measurements were conducted on distinct groups of thirty larvae

(replicates), with larval density per well kept constant. For each measurement, all thirty

larvae in a well were used. Each replicate was fully independent of the others. For cell

counting measurements, each replicate consisted of a single larva. In all experiments,

treatments were randomly assigned to replicates, and blinding was implemented. An initial

experimenter conducted the treatments, collected, and labelled the samples. A second

experimenter then performed the measurements on the labelled samples, assigning new

labels. The first experimenter subsequently quantified the results using these newly encoded

samples. Our sample sizes are consistent with those commonly used in the field and align

with previous publications (Castillo-Ramírez et al., 2019; De Marco et al., 2013; Yeh et al.,

2013; De Marco et al., 2016; vom Berg-Maurer et al., 2016; Herget et al., 2023;). They are

based on prior work that established acceptable coefficients of variation for measurements

while minimizing the use of subjects, in accordance with ethical guidelines. Data were tested

for normality and homoscedasticity using the Shapiro-Wilk and KS normality tests and the

Brown–Forsythe and Bartlett's tests, respectively. We employed Mann-Whitney tests for

pairwise comparisons involving non-normally distributed data, two-way ANOVAs followed

by Bonferroni's post-hoc tests for multiple comparisons. In cases where the data did not

meet the assumption of homoscedasticity, we applied log transformations to normalize

variance. Unpaired two-tailed t-tests were used for comparing independent groups, and one-

sample t-tests were performed when comparing data to a hypothetical mean (i.e., a fold

change of 1). No data points or samples were excluded from the analysis. Statistical analyses

were performed using MS-Excel (Microsoft Corp; Redmond, WA, USA) and Prism 10.2.0

(Graphpad Software Inc, San Diego, CA, USA).

Acknowledgments

We would like to thank L. Flores-García for assistance with the experiments, Karl J.

Iremonger and Chrysanthi Fergani for providing useful comments on a previous draft, R.

Singer and A. Schoell for expert fish care, and R. Rödel for technical support.

Competing Interests

561   The authors declare that the research was conducted in the absence of any commercial or

562   financial relationships that could be construed as a potential conflict of interest.

Author Contributions

L.A.C-R., Conceptualization, Data curation, Formal analysis, Investigation, Validation,

Methodology, Writing - original draft; U.H., Data curation, Formal analysis, Methodology,

Writing – original draft; S.R., Conceptualization, Resources, Supervision, Funding

acquisition, Methodology, Project administration, Writing – review; R.J.DM.,

Conceptualization, Resources, Supervision, Funding acquisition, Investigation, Validation,

Methodology, Data curation, Formal analysis, Visualization, Project administration, Writing

- original draft, Writing - review and editing.

Funding

This work was supported by the Max Planck Society and Liverpool John Moores University.

Data availability

The experimental datasets presented in this study are available upon request from the

authors.

References

Agorastos, A., Pervanidou, P., Chrousos, G. P., and Baker, D.G. (2019). Developmental

trajectories of early life stress and trauma: a narrative review on neurobiological aspects

beyond stress system dysregulation. *Front. Psychiatry* **10,**118. doi:

10.3389/fpsyt.2019.00118.

Alderman, S.L. and Bernier, N.J. (2009). Ontogeny of the corticotropin-releasing factor

system in zebrafish. *Gen. Comp. Endocrinol.* **164**(1),61-9. doi:

10.1016/j.ygcen.2009.04.007.

Alderman, S.L. and Vijayan, M.M. (2012). 11β-Hydroxysteroid dehydrogenase type 2 in

zebrafish brain: a functional role in hypothalamus-pituitary-interrenal axis regulation. *J.*

Endocrinol. **215**(3),393-402. doi: 10.1530/JOE-12-0379.

Alsop, D. and Vijayan, M.M. (2008). Development of the corticosteroid stress axis and

receptor expression in zebrafish. *Am. J. Physiol. Regul. Integr. Comp. Physiol.*

294(3),R711-9. doi.org/10.1152/ajpregu.00671.2007

Alsop, D. and Vijayan, M.M. (2009). Molecular programming of the corticosteroid stress

axis during zebrafish development. *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.*

153(1),49-54. doi.org/10.1016/j.cbpa.2008.12.008

Barton, B.A., Schreck, C.B. and Barton, L.D. (1987). Effects of chronic cortisol

administration and daily acute stress on growth, physiological conditions, and stress

responses in juvenile rainbow trout. *Dis. Aquat. Org.* **2**,173–185.

doi:10.3354/DAO002173.

Biag, J., Huang, Y., Gou, L., Hintiryan, H., Askarinam, A., Hahn, J.D., Toga, A.W.,

and Dong, H.W. (2012). Cyto- and chemoarchitecture of the hypothalamic

paraventricular nucleus in the C57BL/6J male mouse: a study of immunostaining and

multiple fluorescent tract tracing. *J. Comp. Neurol.* **520**(1),6-33. doi: 10.1002/cne.22698.

Bick, J., and Nelson, C.A. (2016). Early adverse experiences and the developing brain.

Neuropsychopharmacology **41**,177–196. doi: 10.1038/npp.2015.252.

Biran, J., Tahor, M., Wircer, E. and Levkowitz, G. (2015). Role of developmental factors

in hypothalamic function. *Front. Neuroanat.* **9**,47. doi: 10.3389/fnana.2015.00047.

Bolton, J.L., Short, A.K., Othy, S., Kooiker, C.L., Shao, M., Gunn, B.G., Beck, J., Bai,

X., Law, S.M., Savage, J.C., et al. (2022). Early stress-induced impaired microglial

pruning of excitatory synapses on immature CRH-expressing neurons provokes aberrant

adult stress responses. *Cell Rep.* **38**(13),110600. doi: 10.1016/j.celrep.2022.110600.

Castillo-Ramírez, L.A., Ryu, S., and De Marco, R.J. (2019). Active behaviour during early development shapes glucocorticoid reactivity. *Sci. Rep.* **9**(1),12796. doi: 10.1038/s41598-019-49388-3.

Castillo-Ramírez, L.A., Ryu, S., and De Marco, R.J. (2024). Cortisol dynamics and GR-dependent feedback regulation in zebrafish larvae exposed to repeated stress. bioRxiv doi: https://doi.org/10.1101/2024.08.07.606983

Champagne, D.L., and Richardson, M.K. (2013). Patterns of cortisol reactivity to stress and the development of coping style in zebrafish *Danio rerio*. *J. Neuroendocrinol.* **25**(9),750-61. doi: 10.1111/jne.12053.

Charmandari, E., Tsigos, C., and Chrousos, G. (2005). Endocrinology of the stress response. *Annu. Rev. Physiol.* **67**,259-84. doi: 10.1146/annurev.physiol.67.040403.120816.

Chen, Y., and Baram, T. Z. (2016). Toward understanding how early-life stress reprograms cognitive and emotional brain networks. *Neuropsychopharmacology* **41**,197–206. doi: 10.1038/npp.2015.181.

Chin, J.S.R., Phan, T.N., Albert, L.T., Keene, A.C., and Duboué, E.R. (2022). Long lasting anxiety following early life stress is dependent on glucocorticoid signaling in zebrafish. *Sci. Rep.* **12**(1),12826. doi: 10.1038/s41598-022-16257-5.

Chrousos, G.P. (1998). Stressors, stress, and neuroendocrine integration of the adaptive response. The 1997 Hans Selye Memorial Lecture. *Ann. N. Y. Acad. Sci.* **851**,311-35. doi: 10.1111/j.1749-6632.1998.tb09006.x.

Chrousos, G.P. (2009). Stress and disorders of the stress system. *Nat. Rev. Endocrinol.* **5**, 374–381. doi: 10.1038/nrendo.2009.106.

Dallman, M.F. (2005). Fast glucocorticoid actions on brain: Back to the future. *Front. Neuroendocrinol.* **26**(3-4),103-8. doi: 10.1016/j.yfrne.2005.08.001.

Dallman, M.F., and Yates, F.E. (1969). Dynamic asymmetries in the corticosteroid

feedback path and distribution-metabolism-binding elements of the adrenocortical

system. *Ann. N. Y. Acad. Sci*. **156**(2),696-721. doi: 10.1111/j.1749-6632.1969.tb14008.x.

Dallman, M.F., Akana, S.F., Levin, N., Walker, C.D., Bradbury, M.J., Suemaru, S.,

and Scribner, K.S. (1994). Corticosteroids and the control of function in the

hypothalamo-pituitary-adrenal (HPA) axis. *Ann. N. Y. Acad. Sci.* **746**,22-31. doi:

10.1111/j.1749-6632.1994.tb39206.x.

Danese, A., and McEwen, B.S. (2012). Adverse childhood experiences, allostasis, allostatic

load, and age-related disease. *Physiol. Behav.* **106**,29–39. doi:

10.1016/j.physbeh.2011.08.019.

de Abreu, M.S., Demin, K.A., Giacomini, A.C.V.V., Amstislavskaya, T.G., Strekalova,

T., Maslov, G.O., Kositsin, Y., Petersen, E.V., and Kalueff, A.V. (2021).

Understanding how stress responses and stress-related behaviors have evolved in

zebrafish and mammals. *Neurobiol. Stress.* **15**,100405. doi: 10.1016/j.ynstr.2021.100405.

de Kloet, E.R., and Joëls, M. (2023). The cortisol switch between vulnerability and

resilience. *Mol. Psychiatry* **29**(1),20-34. doi: 10.1038/s41380-022-01934-8.

de Kloet, E.R., Joëls, M., and Holsboer, F. (2005). Stress and the brain: from adaptation to

disease. *Nat. Rev. Neurosci.* **6**,463–475. doi: 10.1038/nrn1683.

de Kloet, E.R., Vreugdenhil, E., Oitzl, M.S., and Joëls, M. (1998). Brain corticosteroid

receptor balance in health and disease. *Endocr. Rev.* **19**(3),269-301. doi:

10.1210/edrv.19.3.0331.

De Marco, R.J., Groneberg, A.H., Yeh, C.M., Castillo-Ramírez, L.A., and Ryu, S.

(2013). Optogenetic elevation of endogenous glucocorticoid level in larval zebrafish.

Front. Neural Circuits **7**,82. doi: 10.3389/fncir.2013.00082.

De Marco, R.J., Groneberg, A.H., Yeh, C.M., Treviño, M., and Ryu, S. (2014). The

behavior of larval zebrafish reveals stressor-mediated anorexia during early vertebrate

development. *Front. Behav. Neurosci.* **8**,367. doi: 10.3389/fnbeh.2014.00367.

De Marco, R.J., Thiemann, T., Groneberg, A.H., Herget, U., and Ryu, S. (2016).

Optogenetically enhanced pituitary corticotroph cell activity post-stress onset causes

rapid organizing effects on behaviour. *Nat. Commun.* **7**,12620. doi:

10.1038/ncomms12620

Eachus, H., Choi, M.K., and Ryu, S. (2021). The Effects of Early Life Stress on the Brain

and Behaviour: Insights From Zebrafish Models. *Front. Cell Dev. Biol.* **9**,657591. doi:

10.3389/fcell.2021.657591.

Eachus, H., Choi, M.K., Tochwin, A., Kaspareit, J., Ho, M., and Ryu, S. (2024).

Elevated glucocorticoid alters the developmental dynamics of hypothalamic neurogenesis

in zebrafish. *Commun. Biol.* **7**(1),416. doi: 10.1038/s42003-024-06060-5.

Eaton, J.L., Holmqvist, B., and Glasgow, E. (2008). Ontogeny of vasotocin-expressing

cells in zebrafish: selective requirement for the transcriptional regulators orthopedia and

single-minded 1 in the preoptic area. *Dev. Dyn.* **237**(4),995-1005. doi:

10.1002/dvdy.21503.

Egan, R.J., Bergner, C.L., Hart, P.C., Cachat, J.M., Canavello, P.R., Elegante, M.F.,

Elkhayat, S.I., Bartels, B.K., Tien, A.K., Tien, D.H., et al. (2009). Understanding

behavioral and physiological phenotypes of stress and anxiety in zebrafish. *Behav. Brain*

Res. **205**(1),38-44. doi: 10.1016/j.bbr.2009.06.022.

Facchinello, N., Skobo, T., Meneghetti, G., Colletti, E., Dinarello A, Tiso N, Costa R,

Gioacchini G, Carnevali O, Argenton F, et al. (2017). nr3c1 null mutant zebrafish are

viable and reveal DNA-binding-independent activities of the glucocorticoid receptor. *Sci.*

Rep. **7**(1),4371. doi: 10.1038/s41598-017-04535-6.

Felszeghy, K., Bagdy, G., and Nyakas, C. (2000). Blunted pituitary-adrenocortical stress

response in adult rats following neonatal dexamethasone treatment. *J. Neuroendocrinol.*

12(10),1014-21. doi: 10.1046/j.1365-2826.2000.00551.x.

Glover, V., O'Connor, T.G., and O'Donnell, K. (2010). Prenatal stress and the

programming of the HPA axis. *Neurosci. Biobehav. Rev.* **35**(1),17-22. doi:

10.1016/j.neubiorev.2009.11.008.

Grunwald, D.J., and Eisen, J.S. (2002). Headwaters of the zebrafish — emergence of a

new model vertebrate. *Nat. Rev. Genet.* **3**,717–724. doi: 10.1038/nrg891.

Gutierrez-Triana, J.A., Herget, U., Castillo-Ramírez, L.A., Lutz, M., Yeh, C.M., De

Marco, R.J., and Ryu. S. (2015). Manipulation of Interrenal Cell Function in

Developing Zebrafish Using Genetically Targeted Ablation and an Optogenetic Tool.

Endocrinology **156**(9),3394-401. doi: 10.1210/EN.2015-1021.

Hare, A.J., Zimmer, A.M., LePabic, R., Morgan, A.L., and Gilmour, K.M. (2021).

Early-life stress influences ion balance in developing zebrafish (*Danio rerio*). *J. Comp.*

Physiol. B. **191**(1),69-84. doi: 10.1007/s00360-020-01319-9.

Harbuz, M.S., and Lightman, S.L. (1989). Responses of hypothalamic and pituitary

mRNA to physical and psychological stress in the rat. *J. Endocrinol.* **122**(3),705-11. doi:

10.1677/joe.0.1220705.

Hartmann, J., Bajaj, T., Klengel, C., Chatzinakos, C., Ebert, T., Dedic, N.,

McCullough, K.M., Lardenoije, R., Joëls, M., Meijer, O.C., et al. (2021).

Mineralocorticoid receptors dampen glucocorticoid receptor sensitivity to stress via

regulation of FKBP5. *Cell Rep.* **35**(9),109185. doi: 10.1016/j.celrep.2021.109185.

Herget, U., and Ryu, S. (2015). Coexpression analysis of nine neuropeptides in the

neurosecretory preoptic area of larval zebrafish. *Front. Neuroanat.* **9**,2. doi:

10.3389/fnana.2015.00002.

Herget, U., Wolf, A., Wullimann, M.F., and Ryu, S. (2014). Molecular neuroanatomy and

chemoarchitecture of the neurosecretory preoptic-hypothalamic area in zebrafish larvae.

J. Comp. Neurol. **522**(7),1542-1564. doi: 10.1002/cne.23480.

Herget, U., Ryu, S., and De Marco, R.J. (2023) Altered glucocorticoid reactivity and

behavioral phenotype in rx3-/- larval zebrafish. *Front. Endocrinol.* **6**(14),1187327. doi:

10.3389/fendo.2023.1187327.

Herman, J.P., and Cullinan, W.E. (1997). Neurocircuitry of stress: central control of the

hypothalamo-pituitary-adrenocortical axis. *Trends Neurosci.* **20**(2),78-84. doi:

10.1016/s0166-2236(96)10069-2.

Holtzman, N.G., Kathryn Iovine, M., Liang, J.O., and Morris, J. (2016). Learning to fish

with genetics: a primer on the vertebrate model *Danio rerio*. *Genetics* **203**,1069–1089.

doi: 10.1534/genetics.116.190843.

Kastenhuber, E., Kratochwil, C.F., Ryu, S., Schweitzer, J., and Driever, W. (2010).

Genetic dissection of dopaminergic and noradrenergic contributions to catecholaminergic

tracts in early larval zebrafish. *J. Comp. Neurol.* **518**(4),439-58. doi: 10.1002/cne.22214.

Kinlein, S.A., Wilson, C.D., and Karatsoreos, I.N. (2015). Dysregulated hypothalamic-

pituitary-adrenal axis function contributes to altered endocrine and neurobehavioral

responses to acute stress. *Front. Psychiatry.* **6**,31. doi: 10.3389/fpsyt.2015.00031.

Kiss, J.Z. (1988). Dynamism of chemoarchitecture in the hypothalamic paraventricular

nucleus. *Brain Res. Bull.* **20**(6),699-708. doi: 10.1016/0361-9230(88)90080-9.

Krozowski, Z. (1999). The 11beta-hydroxysteroid dehydrogenases: functions and

physiological effects. *Mol. Cell. Endocrinol.* **151**(1-2),121-7. doi: 10.1016/s0303-

7207(98)00256-1.

Kurrasch, D.M., Nevin, L.M., Wong, J.S., Baier, H., and Ingraham, H.A. (2009).

Neuroendocrine transcriptional programs adapt dynamically to the supply and demand

for neuropeptides as revealed in NSF mutant zebrafish. *Neural Dev.* **4**,22. doi:

10.1186/1749-8104-4-22.

Langebeck-Jensen, K., Shahar, O.D., Schuman, E.M., Langer, J.D., and Ryu, S. (2019).

Larval Zebrafish Proteome Regulation in Response to an Environmental Challenge.

Proteomics. **19**(14),e1900028. doi: 10.1002/pmic.201900028.

Lauter, G., Söll, I., and Hauptmann, G. (2011). Multicolor fluorescent in situ

hybridization to define abutting and overlapping gene expression in the embryonic

zebrafish brain. *Neural Dev.* **6**,10. doi: 10.1186/1749-8104-6-10.

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using

real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**(4),402-8.

doi: 10.1006/meth.2001.1262.

Löhr, H., Ryu, S., and Driever, W. (2009). Zebrafish diencephalic A11-related

dopaminergic neurons share a conserved transcriptional network with neuroendocrine cell

lineages. *Development* **136**(6),1007-17. doi: 10.1242/dev.033878.

Lupien, S.J., McEwen, B.S., Gunnar, M.R., and Heim, C. (2009). Effects of stress

throughout the lifespan on the brain, behaviour and cognition. *Nat. Rev. Neurosci.*

10,434–445. doi: 10.1038/nrn2639.

Malkoski, S.P., and Dorin, R.I. (1999). Composite glucocorticoid regulation at a

functionally defined negative glucocorticoid response element of the human

corticotropin-releasing hormone gene. *Mol. Endocrinol.* **13**(10),1629-44. doi:

10.1210/mend.13.10.0351.

McCurley, A.T., and Callard, G.V. (2008). Characterization of housekeeping genes in

zebrafish: male-female differences and effects of tissue type, developmental stage and

chemical treatment. *BMC Molecular Biol.* **9**,102. https://doi.org/10.1186/1471-2199-9-

102.

McEwen, B.S. (2007). Physiology and neurobiology of stress and adaptation: central role of

the brain. *Physiol. Rev.* **87**(3),873-904. doi: 10.1152/physrev.00041.2006.

Moisiadis, V.G., and Matthews, S.G. (2014). Glucocorticoids and fetal programming part

1: Outcomes. *Nat. Rev. Endocrinol.* **10**(7),391-402. doi: 10.1038/nrendo.2014.73.

Munck, A., Guyre, P.M., and Holbrook, N.J. (1984). Physiological functions of

glucocorticoids in stress and their relation to pharmacological actions. *Endocr. Rev.*

5(1),25-44. doi: 10.1210/edrv-5-1-25.

Nagpal, J., Eachus, H., Lityagina, O., and Ryu, S. (2024). Optogenetic induction of

chronic glucocorticoid exposure in early-life leads to blunted stress-response in larval

zebrafish. *Eur. J. Neurosci.* **59**(11),3134-3146. doi: 10.1111/ejn.16301.

Nelson, D.R., Goldstone, J.V., and Stegeman, J.J. (2013). The cytochrome P450 genesis

locus: the origin and evolution of animal cytochrome P450s. *Philos. Trans. R. Soc. Lond.*

B Biol. Sci. **368**(1612),20120474. doi: 10.1098/rstb.2012.0474.

O'Connor, T.G., Bergman, K., Sarkar, P., and Glover, V. (2013). Prenatal cortisol

exposure predicts infant cortisol response to acute stress. *Dev. Psychobiol.* **55**(2),145-55.

doi: 10.1002/dev.21007.

Ryu, S., and De Marco, R.J. (2017). Performance on innate behaviour during early

development as a function of stress level. *Sci. Rep.* **7**,7840.

https://doi.org/10.1038/s41598-017-08400-4.

Ryu, M.H., Moskvin, O.V., Siltberg-Liberles, J., and Gomelsky, M. (2010). Natural and

engineered photoactivated nucleotidyl cyclases for optogenetic applications. *J. Biol.*

Chem. **285**(53),41501-8. doi: 10.1074/jbc.M110.177600.

Sawchenko, P.E., Swanson, L.W., and Vale, W.W. (1984). Corticotropin-releasing factor:

co-expression within distinct subsets of oxytocin-, vasopressin-, and neurotensin-

immunoreactive neurons in the hypothalamus of the male rat. *J. Neurosci.* **4**(4),1118-29.

doi: 10.1523/JNEUROSCI.04-04-01118.1984.

Shipston, M.J. (2022). Glucocorticoid action in the anterior pituitary gland: Insights from

corticotroph physiology. *Curr. Opin. Endocr. Metab. Res.* **25**,100358. doi:

10.1016/j.coemr.2022.100358.

Simmons, D.M., and Swanson, L.W. (2009). Comparison of the spatial distribution of

seven types of neuroendocrine neurons in the rat paraventricular nucleus: toward a global

3D model. *J. Comp. Neurol.* **516**(5),423-41. doi: 10.1002/cne.22126.

Sinars, C.R., Cheung-Flynn, J., Rimerman, R.A., Scammell, J.G., Smith, D.F., and

Clardy, J. (2003). Structure of the large FK506-binding protein FKBP51, an Hsp90-binding

protein and a component of steroid receptor complexes. *Proc. Natl. Acad. Sci. U. S. A.*

100(3),868-73. doi: 10.1073/pnas.0231020100.

Stierl, M., Stumpf, P., Udwari, D., Gueta, R., Hagedorn, R., Losi, A., Gärtner, W.,

Petereit, L., Efetova, M., Schwarzel, M., et al. (2011). Light modulation of cellular

cAMP by a small bacterial photoactivated adenylyl cyclase, bPAC, of the soil bacterium

Beggiatoa. *J. Biol. Chem.* **286**(2),1181-8. doi: 10.1074/jbc.M110.185496.

Stocco, D.M. (2000). The role of the StAR protein in steroidogenesis: challenges for the

future. *J. Endocrinol.* **164**(3),247-53. doi: 10.1677/joe.0.1640247.

Swaminathan, A., Gliksberg, M., Anbalagan, S., Wigoda, N., and Levkowitz, G. (2023).

Stress resilience is established during development and is regulated by complement

factors. *Cell Rep.* **42**(1),111973. doi: 10.1016/j.celrep.2022.111973.

Swanson, L.W. (1991). Biochemical switching in hypothalamic circuits mediating

responses to stress. *Prog. Brain Res.* **87**,181-200. doi: 10.1016/s0079-6123(08)63052-6.

Swanson, L.W., and Sawchenko, P.E. (1983). Hypothalamic integration: organization of

the paraventricular and supraoptic nuclei. *Annu. Rev. Neurosci.* **6**,269-324. doi:

10.1146/annurev.ne.06.030183.001413.

Tan, J.X.M., Ang, R.J.W., and Wee, C.L. (2022). Larval zebrafish as a model for

mechanistic discovery in mental health. *Front. Mol. Neurosci.* **15**,900213. doi:

10.3389/fnmol.2022.900213.

Teicher, M.H., Samson, J.A., Anderson, C.M., and Ohashi, K. (2016). The effects of

childhood maltreatment on brain structure, function and connectivity. *Nat. Rev. Neurosci.*

17,652–666. doi: 10.1038/nrn.2016.111.

Theodoridi, A., Dinarello, A., Badenetti, L., Pavlidis, M., Dalla Valle, L., and

Tsalafouta, A. (2021). Knockout of the *hsd11b2* Gene Extends the Cortisol Stress

Response in Both Zebrafish Larvae and Adults. *Int. J. Mol. Sci.* **22**(22),12525. doi:

10.3390/ijms222212525.

Timmermans, S., Souffriau, J., and Libert, C. (2019). A General Introduction to

Glucocorticoid Biology. *Front. Immunol.* **10**,1545. doi: 10.3389/fimmu.2019.01545.

Tokarz, J., Möller, G., Hrabě de Angelis, M., and Adamski, J. (2015). Steroids in teleost

fishes: A functional point of view. *Steroids* **103**,123-44. doi:

10.1016/j.steroids.2015.06.011.

Tsalafouta, A., Papandroulakis, N., Gorissen, M., Katharios, P., Flik, G., and Pavlidis,

M. (2014). Ontogenesis of the HPI axis and molecular regulation of the cortisol stress

response during early development in Dicentrarchus labrax. *Sci. Rep.* **4**,5525. doi:

10.1038/srep05525.

Unger, J.L., and Glasgow, E. (2003). Expression of isotocin-neurophysin mRNA in

developing zebrafish. *Gene Expr. Patterns* **3**(1),105-8. doi: 10.1016/s1567-

133x(02)00064-9.

Veenstra-VanderWeele, J., and Warren, Z. (2015). Intervention in the context of

development: pathways toward new treatments. *Neuropsychopharmacology* **40**,225–237.

doi: 10.1038/npp.2014.232.

vom Berg-Maurer, C.M., Trivedi, C.A., Bollmann, J.H., De Marco, R.J., and Ryu, S.

(2016). The Severity of Acute Stress Is Represented by Increased Synchronous Activity

and Recruitment of Hypothalamic CRH Neurons. *J. Neurosci.* **36**(11),3350-62. doi:

10.1523/JNEUROSCI.3390-15.2016.

Weger, M., Diotel, N., Weger, B.D., Beil, T., Zaucker, A., Eachus, H.L., Oakes, J.A., do

Rego, J.L., Storbeck, K.H., Gut, P., et al. (2018). Expression and activity profiling of

the steroidogenic enzymes of glucocorticoid biosynthesis and the fdx1 co-factors in

zebrafish. *J. Neuroendocrinol.* **30**(4),e12586. doi: 10.1111/jne.12586.

Weinstock, M. (2008). The long-term behavioural consequences of prenatal stress.

Neurosci. Biobehav. Rev. **32**(6),1073-86. doi: 10.1016/j.neubiorev.2008.03.002.

Wendelaar Bonga, S.E. (1997). The stress response in fish. *Physiol. Rev.* **77**(3),591-625.

doi: 10.1152/physrev.1997.77.3.591.

Westerfield, M. (2000). *The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish*

(Danio rerio). 4th ed. Eugene: Univ. of Oregon Press.

Whitnall, M.H., Mezey, E., and Gainer, H. (1985). Co-localization of corticotropin-

releasing factor and vasopressin in median eminence neurosecretory vesicles. *Nature*

317(6034),248-50. doi: 10.1038/317248a0.

Wilson, K.S., Tucker, C.S., Al-Dujaili, E.A., Holmes, M.C., Hadoke, P.W., Kenyon,

C.J., and Denvir, M.A. (2016). Early-life glucocorticoids programme behaviour and

metabolism in adulthood in zebrafish. *J. Endocrinol.* **230**(1),125-42. doi: 10.1530/JOE-

15-0376.

Yeh, C.M., Glöck, M., and Ryu, S. (2013). An optimized whole-body cortisol

quantification method for assessing stress levels in larval zebrafish. *PLoS One*

8(11),e79406. doi: 10.1371/journal.pone.0079406.

Zannas, A.S., Wiechmann, T., Gassen, N.C., and Binder, E.B. (2016). Gene-Stress-

Epigenetic Regulation of FKBP5: Clinical and Translational Implications.

Neuropsychopharmacology **41**(1),261-74. doi: 10.1038/npp.2015.235.

Ziv, L., Muto, A., Schoonheim, P.J., Meijsing, S.H., Strasser, D., Ingraham, H.A.,

Schaaf, M.J., Yamamoto, K.R., and Baier, H. (2013). An affective disorder in

zebrafish with mutation of the glucocorticoid receptor. *Mol. Psychiatry* **18**(6),681-91.

doi: 10.1038/mp.2012.64.

Figure legends

Figure 1. Forced swimming at 5 dpf reduces a larva's cortisol response to a subsequent exposure to the same type of stress. (**A**,**B**) Comparison steps and time points for measuring baseline whole-body cortisol levels (**A**) and whole-body cortisol trajectory post-vortex (**B**) in 6 dpf larvae with or without ELC exposure. (**C**,**D**) Baseline whole-body cortisol levels (**C**) and whole-body cortisol trajectory post-vortex (**D**) in control (black) and ELC (orange) larvae. (**C**) p=0.90 by Mann-Whitney test; N=6 per group. Box and whiskers (min to max), all data points shown. (**D**) 10' and 20': ****p<0.0001, 30': **p=0.003, 40': p=0.125, 60': p=0.99 by Bonferroni's test after a two-way ANOVA on square root-transformed cortisol levels; N=6 per group. Mean ± S.E.M. The red line represents the average baseline whole-body cortisol levels from (**C**).

Figure 2. Peptidergic NPO cell numbers in wildtype larvae with or without ELC exposure. (**A**) Comparison of cell numbers in the NPO with or without ELC exposure. (**B**) Maximum intensity projections of confocal stacks show NPO cells expressing *oxt* (green), *avp* (red), or *crh* (blue) in control (above) or ELC (below) larvae. Double-colour comparisons of each pairwise combination are also shown. Scale bar: 25 μm. (**C**) Comparison of coexpression levels in the NPO with or without ELC exposure. (**A**,**C**) Orange: ELC larvae, N=20; black: control larvae, N = 27; all data points shown; p values by Mann-Whitney tests. (See also Results.)

Figure 3. Transcript abundance of genes involved in cortisol metabolism, steroidogenesis, and stress modulation in control and ELC larvae. (**A**-**L**) Fold change in expression of selected genes relative to control reference samples, measured under baseline (basal) conditions and at 30, 60, or 120 minutes after high-strength vortex exposure in

control (black) and ELC (orange) larvae; N = 5 per group. p values for group and time factors from two-way ANOVAs are shown in the top-left corner of each panel. Asterisks indicate Bonferroni's test results following the ANOVAs (*p<0.05, **p<0.01, ***p<0.001). **(M)** Density grid summarizing fold change comparisons in expression levels of the 12 target genes. Asterisks on the right denote results for the group and time factors from two-way ANOVAs. Asterisks within ELC panels indicate Bonferroni's test results following the ANOVAs. (See Results for details.)

Figure 4. ELC alters GC_R to a second vortex during the refractory period. (**A**) Sampling scheme for measuring GC_R to homotypic stress in 6 dpf larvae with or without ELC. (**B**) Whole-body cortisol following two 3-minute vortices applied within a 30-minute refractory period in control (black) and ELC (orange) larvae. Control larvae did not show cortisol levels above baseline in response to a second 3-minute vortex, while ELC larvae did. N=6 per group. ****$p < 0.0001$ by unpaired two-tailed t-test. (**C**) Baseline whole-body cortisol levels in ELC larvae relative to controls (data from **Fig. 1C**), depicted for the sake of interpreting panels (**D**) and (**E**). $p = 0.40$ by one-sample t-test against fold change of '1'. (**D**) GC_R to 1st and 2nd vortices relative to baseline in control and ELC larvae. Data from **Fig. 1C**, **Fig. 1D** (10 min post-vortex), and (**B**). N=6 per group. p values indicate Bonferroni's test results following a two-way ANOVA. (**E**) GC_R to the 2nd vortex relative to GC_R to the 1st vortex in control (black) and ELC (orange) larvae (data from **D**). $p < 0.0001$ by unpaired two-tailed t-test. (**B**-**E**) Box and whiskers (min to max), (**C**-**E**) all data points shown.

Figure 5. Proposed model illustrating the initial adaptation of the HPI axis following ELC. Prolonged early-life activation of the HPI axis, induced by increased environmental challenge, diminishes the NPO's capacity to influence downstream HPI axis activity. Baseline conditions show enhanced cortisol inactivation, which can persist upon stress

921   onset. There is an increase in cortisol synthesis capacity and elevated levels of stress

922   modulators. These adaptations enhance the larva's capability to regulate GC levels,

923   preventing excess cortisol while facilitating its production in response to familiar stress.

925 **Figure 2**

Figure 3

 Figure 4

     Table 1 Design primer sequences.

Gene name	Gene id	Primer sequence (5´→ 3´)
avp	ENSDARG00000058567	F: CCCAGCCGGAGCCCATCAGA
		R: CCATGCAGACCTGCGCCTCC
pomca	ENSDARG00000043135	F: CGCAGACCCATCAAGGTGTA
		R: CGTTTCGGCGGATTCCT
nr3c2	ENSDARG00000102082	F: CCCATTGAGGACCAAATCAC
		R: AGTAGAGCATTTGGGCGTTG
mc2r	ENSDARG00000054949	F: CTTGGTGATTGTGGCTGTG
		R: GATAAAGCAGTACATGGGAGAG
nr3c1	ENSDARG00000025032	F: ACAGCTTCTTCCAGCCTCAG
		R: CCGGTGTTCTCCTGTTTGAT
fkbp5	ENSDARG00000028396	F: CCGACTGCTGTCTTTACCT
		R: CATGCTGTTTCACAATCTTGC
oxt	ENSDARG00000042845	F: CGGCCTGCTACATCTCAAAC
		R: TGCCTTCACCACAGCAGATA
crhr1	ENSDARG00000003989	F: TTTATCCTGCGAAATGCCAC
		R: GCACCAGATCACATTGCTC
cyp11c1	ENSDARG00000042014	F: TGTGCTGAAGGTGATTCTCG
		R: GCTCATGCACATTCTGAGGA
crh	ENSDARG00000027657	F: ATAGGTGGTGGGTTGTTCC
		R: CTACATTCATACGGCGGTG
star	ENSDARG00000006137	F: TCAAATTGTGTGCTGGCATT
		R: CCAAGTGCTAGCTCCAGGTC
hsd11b2	ENSDARG00000001975	F: TGCTGCTGGCTGTACTTCAC
		R: TGCATCCAACTTCTTTGCTG