1   **Title**
2
3   **Local synthesis of Reticulon-1C lessens the outgrowth of injured axons and Spastin**
4   **activity**
5   **Authors**
6

Alejandro Luarte[1,2,3]*, Javiera Gallardo[8], Daniela Corvalán[1,2], Ankush Chakraborty [1,2], Cláudio
Gouveia-Roque[5], Francisca Bertin[11], Carlos Contreras[1,2], Juan Pablo Ramírez [1,2], André Weber
[20], Waldo Acevedo[8], Werner Zuschratter[19], Rodrigo Herrera-Molina[9,10], Úrsula Wyneken [1,2,3] ,
Andrea Paula-Lima [16,17], Tatiana Adasme-Rocha[18], Jorge Toledo[21], Rodrigo Vergara[22] , Antonia
Figueroa[1,2], Carolina González[15], Christian González-Billault[4, 12, 13, 14], Ulrich Hengst [5,6], and
Andrés Couve [4]
13
14   **Affiliations**
15
1 Faculty of Medicine, Universidad de los Andes, Santiago 7620001, Chile
2 Program in Neuroscience, Center for Biomedical Research, and Innovation (CiiB),
Universidad de los Andes, Santiago 7620001, Chile
3 IMPACT, Center of Interventional Medicine for Precision and Advanced Cellular Therapy,
Santiago 7620001, Chile
4 Department of Neuroscience, Faculty of Medicine, Universidad de Chile, Santiago, Chile.
5 The Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia
University Vagelos College of Physicians and Surgeons, New York, NY 10032, USA
6 Department of Pathology and Cell Biology, Columbia University Vagelos College of
Physicians and Surgeons, New York, NY 10032, USA
7 Instituto de Química, Facultad de Ciencias, Pontificia Universidad Católica de Valparaíso,
Valparaíso 2373223, Chile
8 Centro de Medicina Regenerativa, Facultad de Medicina, Clínica Alemana-Universidad del
Desarrollo, Santiago, Chile
9 Centro Integrativo de Biología y Química Aplicada, Universidad Bernardo O'Higgins,
8307993 Santiago, Chile
10 Department of Pharmacology & Physiology, George Washington University, Ross Hall,
Washington, DC, USA
11 Facultad de Medicina Veterinaria y Agronomía, Universidad de Las Américas, sede
Providencia, Manuel Montt 948, 7500975, Santiago, Chile
12 Department of Biology, Faculty of Sciences, Universidad de Chile, Santiago, Chile
13 Public Nutrition Unit, Institute for Nutrition and Food Technologies, Universidad de Chile,
Santiago, Chile
14 The Buck Institute for Research on Aging, Novato, USA
15 Latin American Brain Health Institute (BrainLat), Universidad Adolfo Ibañez, Santiago,
Chile
16 Biomedical Neuroscience Institute and Department of Neurosciences, Faculty of Medicine,
Universidad de Chile, Santiago 8380000, Chile
17 Institute for Research in Dental Sciences (ICOD), Faculty of Dentistry, Universidad de Chile,
Santiago 8380544, Chile
18 Oficina de apoyo a la investigación clínica, Hospital Clínico Universidad de Chile, Santiago
8380494, Chile
19 Leibniz Institute for Neurobiology, 39118 Magdeburg, Germany
20 Photonscore GmbH, Brenneckestr. 20, 39118 Magdeburg, Germany
21 Health Sciences Department, Universidad de Aysén, Coyhaique, Chile.
22 Departamento de Kinesiología, Facultad de Artes y Educación Física, Universidad
Metropolitana de Ciencias de la Educación
53
54   *Corresponding author at: Faculty of Medicine, Universidad de los Andes, Santiago 7620001,
55   Chile. aluarte@uandes.cl

Abstract

The regenerative potential of developing cortical axons following injury depends on intrinsic mechanisms, such as axon-autonomous protein synthesis, that are still not fully understood. An emerging factor in this regenerative process is the bi-directional interplay between microtubule dynamics and the delicate morphology of the axonal endoplasmic reticulum (ER). Therefore, we hypothesize that locally synthesized ER structural proteins regulate microtubule dynamics and the regenerative response of cortical axons. This hypothesis is supported by RNA data-mining, which identified *Reticulon-1* as the sole mRNA coding for an ER-shaping protein across eight axonal transcriptomes and found it to be downregulated following injury. Using compartmentalized microfluidic chambers, we demonstrate that local knockdown of *Reticulon-1* mRNA increases the outgrowth while reducing the distal tubulin levels of injured axons. Furthermore, we show by live-cell imaging that axonal *Reticulon-1* knockdown restores the loss of microtubule growth rate and length caused by injury. Interestingly, axonal inhibition of the microtubule-severing protein Spastin fully prevents the effects of axonal *Reticulon-1* knockdown on outgrowth as well as distal tubulin levels, while not affecting most of the analyzed microtubule dynamic parameters. Further, we demonstrate that the Reticulon-1C isoform is locally synthesized within axons and provide evidence supporting its role in attenuating Spastin-mediated microtubule severing. These findings uncover a mechanism by which axonal protein synthesis provides fine control over the microtubule cytoskeleton in response to injury.

Introduction

Developing cortical axons can regenerate after injury but lose this capacity at later stages (Cooke et al., 2022). The transition from a regenerative to a non-regenerative state highlights the importance of identifying the factors that may underlie this shift. The regenerative potential of developing axons depends on extrinsic and intrinsic (cell autonomous) factors that regulate their outgrowth (He & Jin, 2016). The structure of the axonal endoplasmic reticulum (ER) is a determinant intrinsic factor for regulating outgrowth during development and after injury (Aridor & Fish, 2009; Deitch & Banker, 1993; Krijnse-Locker et al., 1995; Nozumi et al., 2009; Petrova et al., 2020; Rao et al., 2016). Indeed, ER proteins are locally synthesized in axons and may contribute to regeneration (Pacheco et al., 2020; Willis et al., 2005). However, the role of axonal protein synthesis in shaping ER function after injury remains largely unexplored.

Unlike the ribosome-rich somato-dendritic ER, the structure of the axonal ER predominantly comprises continuous tubules of smooth ER, maintained by specific ER-shaping proteins (Voeltz et al., 2006; Wu et al., 2017). As such, the mostly tubular axonal ER controls calcium handling, lipid provision, and potentially transmembrane protein supply, which contribute to axon elongation (Cornejo et al., 2017; de Juan-Sanz et al., 2017; González et al., 2016; Luarte et al., 2018). More recently, a bi-directional relationship of the axonal ER with microtubule dynamics has emerged as critical for regulating outgrowth. For instance, the physical coupling of the axonal ER to growing microtubules through STIM1 determines its positioning into filopodia in growth cones to control the guidance of developing axons (Pavez et al., 2019). In turn, the axonal ER controls microtubule dynamics, which is required for axonal specification and proper outgrowth in response to injury (Farías et al., 2019; Rao et al., 2016).

Different lines of evidence support that the protein components of the axonal ER may interact with proteins that regulate microtubule dynamics. For instance, the interaction of ER structural proteins with microtubule severing enzymes may contribute to sculpt the axonal cytoskeleton required for outgrowth (Rao et al., 2016; Stone et al., 2012). As such, the microtubule-severing protein Spastin is known to interact with different ER-shaping proteins (Evans et al., 2006; Mannan et al., 2006), but the contribution of axonal protein synthesis to this process has not been addressed. Here, we hypothesized that the local synthesis of ER structural proteins controls microtubule dynamics and the elongation of injured axons. Using a variety of techniques, we report that local synthesis of the ER-shaping protein Reticulon-1 (Rtn-1) controls microtubule dynamics and the outgrowth of injured cortical axons depending on Spastin-mediated microtubule severing.

Results

Rtn-1 mRNA is consistently identified in axonal transcriptomes

113
114
115
116
We hypothesized that the local synthesis of an ER structural protein controls microtubule
dynamics and the elongation of injured axons. As a first approach to test our hypothesis, we
looked for mRNAs coding for ER structural proteins in reported axonal transcriptomes from
different species, developmental stages, and days *in vitro* (DIV) as indicated in Table I. To
enhance the robustness of our analysis, we focused on transcripts that were consistently identified
across conditions through Benjamini–Hochberg (Ben) analysis. We identified 104 common
mRNAs across all studies (Figure 1A) and classified them under the 'Cellular Component'
category using Gene Ontology (GO) analysis. We found that *Rtn-1* mRNA was the only transcript
coding for an ER structural protein that is shared across all the samples (Figure 1B). Note that
several of the identified transcripts in the category 'endoplasmic reticulum' code for cytoplasmic
ribosomal components, which indeed can be attached to the axonal ER (Koppers et al., 2024) and
be locally synthesized in axons (Shigeoka et al., 2019). On a similar approach, but combining
data from dendrites and axons, it was found that *Reticulon-3* mRNA is present in 16 out of 20
studies, further suggesting a wider presence of other mRNAs coding for ER structural proteins in
axons (von Kügelgen & Chekulaeva, 2020). Interestingly, one of the studies presented in (Figure
1B) showed that *Rtn-1* mRNA levels in cortical axons are decreased by 5.26 ± 0.06-fold after
axotomy (Taylor et al., 2009). Thus, *Rtn-1* mRNA is consistently identified in axons, and it is
down-regulated upon injury.

135
Next, we studied the distribution of Rtn-1 protein in axons from cultured rat cortical
neurons at early developmental stages. To this end, we assessed the fluorescence intensity of Rtn-
1 along the terminal 50 µm of distal axons at 1, 3, and 7 DIV. Axonal identity was confirmed by
β3-tubulin and the absence of the dendritic marker microtubule-associated protein 2 (MAP2).
Rtn-1 fluorescence was concentrated in the most distal axonal region corresponding to the growth
cone (GC), which is significantly more pronounced at 7 DIV compared to 1 DIV (Supplementary
Figure 1A-C). Rtn-1 enrichment within GCs diverged from the staining pattern of KDEL, a
luminal ER marker, which did not exhibit such progressive accumulation (Supplementary Figure
1A-C). The distribution of Rtn-1 agreed with prior studies that localized this protein at the base
of the GC's hand-like structure, as outlined by the F-actin cytoskeleton stained with phalloidin
(Nozumi et al., 2009) (Supplementary Figure 1D). This distribution supports a model where Rtn-
1 contributes to the dense network of ER tubules characteristically found in the GCs, an
observation well-documented in earlier studies (Deitch & Banker, 1993)
149

Axonal synthesis of Rtn-1 lessens outgrowth after injury

To elucidate the role of Rtn-1 synthesis in the outgrowth of injured axons, we first validated knockdown (KD) of this protein by Western blot (WB) in cultured cortical neurons. Neuronal Rtn-1 isoforms include isoform A (Rtn-1A), comprising 776 amino acids, and isoform C (Rtn-1C) with 208 amino acids. Both isoforms share a highly conserved C-terminal region known as the Reticulon homology domain (RHD), differing only in their 20 N-terminal amino acids (Chiurchiù et al., 2014; Roebroek et al., 1993). Thus, we transfected 7 DIV dissociated cortical neurons with siRNAs designed to target the conserved RHD domain (i.e., siRNAs that target all Rtn-1 isoforms) or a control sequence (scrambled, Scr). Forty-eight hours later, samples were analyzed using a pan-Rtn-1 antibody raised against the conserved RHD region. Two distinct Rtn-1 bands were observed: a higher molecular weight band near 180 kDa, likely representing Rtn-1A (Van De Velde et al., 1994), and a lower band near 24 kDa, corresponding to the apparent molecular weight of Rtn-1C (Roebroek et al., 1993; Van De Velde et al., 1994) (Figure 2A). When quantifying the 24 kDa band, both siRNAs significantly reduced Rtn-1 levels compared to Scr (Figure 2A). We also examined the 180 kDa band and found that siRNA 1 reduced expression to a mean of 0.41 relative to Scr, showing a strong trend that did not reach statistical significance (p = 0.05; N = 3; Wilcoxon test compared to 1, data not shown). In contrast, siRNA 2 further reduced expression to a mean of 0.29, which was statistically significant (p = 0.04; N = 3; Wilcoxon test compared to 1, data not shown). Together, these results indicate that, when considering both isoforms, siRNA 2 achieves the stronger KD. Therefore, unless otherwise indicated, for all subsequent experiments siRNA 2 was used to downregulate Rtn-1. Immunofluorescence staining further revealed a marked decrease in Rtn-1 intensity in the cell body and distal axons of neurons co-transfected with an RFP-expressing plasmid and siRNA 2 under the same recently described conditions (Figure 2B).

Once we validated the siRNA 2, we aimed at performing KD of Rtn-1 expression selectively within the axonal compartment (axonal Rtn-1 KD) in 7 DIV microfluidic chamber cultures. This compartmentalized approach has been validated in multiple studies to locally control neuronal protein synthesis (Baleriola et al., 2014; Batista et al., 2017; Zhang et al., 2013).Twenty-four hours later, we found that axonal Rtn-1 KD did not alter somatic Rtn-1 levels yet elicited a significant ~ 25% decrease in somatic β3-tubulin, which were quantified as the intensity fold change for both proteins compared to Scr (Figure 2C). Compatible with Rtn-1 local synthesis, axonal Rtn-1 KD led to a significant ~ 46% decrease in its fluorescence intensity within the distal 50 μm of axons, which was depicted by the corrected fluorescence intensity profile and quantified as the intensity fold change compared to Scr (Figure 2D and E). A slight but significant ~ 16% decrease in β3-tubulin fluorescence intensity fold change was also observed after axonal

187   Rtn-1 KD compared to Scr (Figure 2D and E). These results support that local synthesis of Rtn-1
188   can be locally downregulated in the axonal compartment and suggest an interplay with the
189   organization of the microtubule cytoskeleton.
190
To investigate the role of local Rtn-1 synthesis in axonal outgrowth, we subjected 8 DIV
cortical axons to axotomy by vacuum aspiration in microfluidic chambers. To this end, we
simultaneously performed Rtn-1 KD in either axonal or somato-dendritic compartments and
measured outgrowth twenty-four hours later (Figure 3A). To measure outgrowth, we took
advantage of the semi-circle configuration of the NeuroAnatomy FIJI plugin and performed Sholl
analysis (McCurdy et al., 2019; Sholl, 1953). We quantified the summation of the intersections
of binarized axons with concentric semi-circles of 2 μm increasing radius starting from the
microgroove exit and centered at axonal fields of equivalent size (Figure 3B, top). Interestingly,
we observed a significant increase in outgrowth (higher number of intersections summation)
within the 500-1000 μm and 1000-1500 μm ranges with axonal Rtn-1 KD compared to Scr
(Figure 3B, bottom). Furthermore, this response was specific for the axonal compartment, as
somatic Rtn-1 KD showed a non-significant effect with a slight opposite trend (Figure 3C).
203
To discard off-target effects of siRNA used for axonal Rtn-1 KD, we validated a third
and completely different oligonucleotide sequence (siRNA 3) specifically directed against Rtn-1
and tested its capacity to reproduce the observed phenotype with local KD in injured axons. We
found by WB that this sequence significantly decreased Rtn-1C levels, while Scr and non-
transfected neurons did not. Also, no significant changes of β3-tubulin levels were observed
(Supplementary Figure 2A-B). To test the ability of siRNA 3 to promote outgrowth, we
selectively treated axotomized axons in microfluidic chambers and measured outgrowth twenty-
four hours after injury. Axonal KD performed with siRNA 3 resembled the overall outgrowth
tendency observed with siRNA 2 (Figures 3A-B). It was observed a significant increase in the
summation of axonal intersections within the 0-500 μm and 500-1000 μm ranges after axonal
Rtn-1 KD compared to Scr (Supplementary Figure 2C-D). Together, these results suggest that the
increased outgrowth of injured axons observed with axonal Rtn-1 KD is not likely due to siRNA
off-target effects.
217
To further characterize the role of axonally synthesized Rtn-1 in injured axons, we
assessed the morphology of GCs in the same experimental setting described above. To this end,
we used β3-tubulin immune staining and phalloidin incubation to reveal the actin cytoskeleton.
Axonal Rtn-1 KD did not modify the outline (perimeter) and the area of GCs. However, axonal
Rtn-1 KD produced GCs characterized by a more elongated and compact actin cytoskeleton,
reflected on a significant increase in the elongation ratio (calculated as the length of the major

axis divided by the length of the minor axis of the best-fitting ellipse to the actin boundary) and longest shortest path (LSP, which reflects the maximum length of the skeletonized actin outline), alongside a reduced number of filopodia (Figure 3D). Collectively, these morphometric parameters revealed that axonal Rtn-1 KD yielded GCs resembling a spade-like morphology that has been previously related to increased outgrowth (Kaethner & Stuermer, 1992). Compatible with this idea, somatic Rtn-1 KD did not promote outgrowth nor modified any of these morphological parameters, except for an increase in the GC LSP (Figure 3D). Together, these results suggest that axonal synthesis of Rtn-1 is normally acting to repress the outgrowth of injured axons.

Axonal Rtn-1 KD decreases β3-tubulin levels in injured axons

The previously described changes in β3-tubulin elicited by axonal Rtn-1 KD before injury (Figure 2D), lead us to examine β3-tubulin and Rtn-1 fluorescence intensity under this condition in axotomized axons. Different from what we observed in intact axons (Figure 2D and E), twenty-four hours of axonal Rtn-1 KD did not alter Rtn-1 fluorescence significantly, but it induced a ~ 20 % reduction in β3-tubulin intensity compared to Scr (Figure 4A). In contrast, somatic Rtn-1 KD did decrease Rtn-1 fluorescence in axons by ~ 33% (Figure 4B), supporting the notion that most of axonal Rtn-1 protein originates in the soma. Of note, somatic Rtn-1 KD also decreased β3-tubulin intensity by ~ 19% in axons (Figure 4B).

Consistent with a potential compensatory response of the cell body upon injury, axonal Rtn-1 KD leads to increased somatic Rtn-1 fluorescence compared to Scr (Figure 4C). Interestingly, an isoform-specific increase in Rtn-1C levels has been previously described upon the injury of cortical neurons (Fan et al., 2018). Despite we still do not know the mechanism, the somatic increase that we observe may partially explain the lack of changes in axon Rtn-1 levels after twenty-four hours of axonal KD (Figure 4A), which is potentially countered by somatic Rtn-1 synthesis. Intriguingly, resembling the somatic increase of Rtn-1, β3-tubulin levels were also elevated in the cell bodies following axonal Rtn-1 KD compared to Scr (Figure 4C). In contrast, somatic Rtn-1 KD resulted in a significant reduction in the somatic fluorescence intensity for both proteins compared to Scr (Figure 4D). Together, these results indicate that after axonal injury, the levels of Rtn-1 in the somatic and axonal compartments tend to change in a manner that parallels β3-tubulin, which suggests a mechanistic relationship between these two proteins.

Spastin activity regulates β3-tubulin and axon re-growth under local Rtn-1 KD

The significant alterations in β3-tubulin levels after Rtn-1 KD in the context of injury, led us to explore the role of Rtn-1 interactors that can modify the microtubule cytoskeleton. Notably, a previous study had shown by yeast two hybrid assay that a protein sequence from Rtn-1C directly interacts with the microtubule-severing protein Spastin (Mannan et al., 2006). Therefore, we explored the localization of Spastin and Rtn-1C in distal axons before and after injury.

To this end, we stained 9 DIV non-injured and injured axons (submitted to axotomy at 8 DIV) against Spastin (all isoforms) and Rtn-1C to perform Stimulated Emission Depletion (STED) super-resolution microscopy. In both conditions, Rtn-1C and Spastin showed enrichment of the staining in the GC (Figure 5A, left panel). The distribution of Rtn-1C along the axonal shaft is fine and continuous, consistent with its localization in ER tubules, which appear decorated with clusters of Spastin. In contrast to the more central positioning of Rtn-1C in the axonal shaft, Spastin displays a broader distribution in both conditions (Figure 5A, left panel). This difference is also seen in the GCs, where Rtn-1C is more concentrated at the center of these domains (Figure 5A, left panel). In injured axons, distal Rtn-1C staining appears less compact, adopting a more tubulated pattern (Figure 5A, left panel). Moreover, following injury, Rtn-1C staining seems slightly retracted from the GC compared to the non-injured condition, as also suggested by its relative intensity profile (Figure 5A, right panel). Spastin distribution along the distal axon remains similar between conditions (Figure 5A, right panel). However, in injured axons, Spastin appears less diffuse and more concentrated in clusters than in non-injured axons (Figure 5A left and right panel). No significant changes were observed when analyzing the total fluorescence intensity of both proteins (Supplementary Figure 4). These findings suggest that both proteins are closely apposed in the GC and along the axonal shaft. Although we did not directly test their direct molecular association, these results are consistent with Rtn-1C and Spastin sharing a similar subcellular localization, potentially enabling their functional interaction in distal axons.

To further investigate the role of these proteins, we subjected 8 DIV cortical axons to axotomy by vacuum aspiration in microfluidic chambers. Under these conditions, we aimed at elucidating the potential connection of Rtn-1 axonal synthesis KD with Spastin's microtubule-severing activity. To this end, we took advantage of a Spastazoline (SPTZ), a well-characterized and specific inhibitor of the ATPase activity of Spastin (i.e., microtubule severing)(Cupido et al., 2019). Under these conditions, we monitored the changes in the fluorescence intensity of β3-tubulin, Rtn-1C, and Spastin in distal axons forty-eight hours after injury. Note that these experiments were performed in a two-fold longer period compared to those shown in Figure 4. This extended time was taken to i) reveal more clearly the changes in β3-tubulin content and ii)

allow the proper expression of the EB3-GFP plasmid which is used for the live-cell imaging described later in Figure 5. We found that axonal Rtn-1 KD resulted in a ~ 60% reduction in β3-tubulin levels within the distal 50μm of axons, compared to controls (Scr + DMSO) (Figure 5B). However, co-incubation with 50 nM SPTZ (applied four hours after axotomy) completely abolished the β3-tubulin reduction elicited by axonal Rtn-1 KD, while it did not alter Rtn-1C and Spastin levels (Figure 5B). This rescue of β3-tubulin levels during axonal Rtn-1 KD by a specific drug (SPTZ), makes it unlikely an off-target effect of the siRNA over *β3-tubulin* mRNA. If this was the case, this potential off-target effect should be still operating in the presence of SPTZ. Also discarding unspecific effects over other proteins, SPTZ administration alone did not significantly modify β3-tubulin levels, nor those of Rtn-1C and Spastin (Figure 5B). Together, these observations suggest that axonal Rtn-1 KD correlates with higher Spastin microtubule severing.

Next, we aimed at exploring whether the increased outgrowth elicited by axonal Rtn-1 KD after injury depended on Spastin's microtubule severing. To this end, we studied the effects of axonal Rtn-1 KD on outgrowth forty-eight hours after axotomy by Sholl analysis. In contrast with the effects observed after twenty-four hours of axotomy (Figure 3A), after forty-eight hours of axonal Rtn-1 KD there was only a trend (non-statistically significant) to increase outgrowth in the 1000-1500 μm range compared to controls (Figure 5C and D). However, local co-incubation of 50 nM SPTZ with axonal Rtn-1 KD abolished this trend, producing a statistically significant outgrowth decrease compared to the axonal Rtn-1 KD + DMSO condition (Figure 5 C and D). Local SPTZ application in axons treated with a Scr sequence displayed a trend to increase outgrowth compared to all the conditions, particularly in the 1000-1500 μm segment, but any of these comparisons reached statistical significance (Figure 5C and D). Together, these results suggest that locally synthesized Rtn-1 normally acts to suppress the outgrowth of injured axons, a process that could involve the microtubule-severing activity of Spastin.

Considering the observed shifts in microtubule content and the previously reported role of Spastin in controlling microtubule stability (Aiken & Holzbaur, 2024; Kuo et al., 2019; Teixeira Lopes, 2018), we aimed at quantifying microtubule dynamics by live-cell imaging before and after injury. To this end, cortical neurons cultured in microfluidic chambers were transfected at 7 DIV with a plasmid coding for the microtubule tip-binding protein EB3 fused to the fluorescent construct Emerald GFP (EB3-GFP), which labels the growing plus-ends of polymerizing microtubules (Stepanova et al., 2003). Next, axotomy was performed at 8 DIV microfluidic chambers, and immediately, axons were locally transfected with a Rtn-1 siRNA or Scr sequence. Non-injured 8 DIV axons were transfected in parallel. Four hours after transfection, non-injured and injured axons were either treated with DMSO (vehicle) or 50 nM SPTZ. Then forty-eight

hours since injury (at 10 DIV), axons were directly imaged in microfluidic chambers to examine the dynamic behavior of EB3-GFP puncta (comets).

We first analyzed the effect of axonal Rtn-1 KD on non-injured axons. Interestingly, we found that axonal Rtn-1 KD elicited a statistically significant increase in microtubule growth rate, while it did not significantly modify microtubule track length and lifetime compared to controls (Scr + DMSO) (Figure 5E). To reveal the role of Spastin's microtubule severing in modifying these parameters, we co-incubated axons with 50 nM SPTZ. As such, SPTZ treatment did not prevent the increase in microtubule growth rate elicited by axonal Rtn-1 KD nor modified comet's lifetime but induced a significant increase in track length when compared to controls (Figure 5E).

Next, we analyzed the effect of axonal Rtn-1 KD on comet parameters after injury. In contrast with the milder increase observed in non-injured axons, axonal Rtn-1 KD augmented by nearly two-fold the growth rate and track length of microtubules, while it did not modify comet lifetime compared controls (Figure 5E, left and right panels). Of note, compared to non-injured axons, growth rate and track length are significantly decreased on axotomized axons (from 0.09 to 0.06 μm/s, and from 4.6 μm to 2.6 μm; p<0.0001, Mann Whitney test). These results are compatible with axonal damage promoting a more unorganized or less processivity of microtubule growth (Rolls et al., 2020; Yap et al., 2017). SPTZ co-treatment did not prevent the increase in microtubule growth rate and track length elicited by axonal Rtn-1 KD after injury compared to controls, nor modified comet's lifetime (Figure 5E, right panels). However, SPTZ treatment along with axonal Rtn-1 KD did induce a significant increase in comet lifetime when compared to axonal Rtn-1 KD + DMSO as shown in right panels of Figure 5E. Together, these results suggest that axonal Rtn-1 synthesis controls microtubule dynamics in both non-injured and injured axons, for the most part independently of Spastin-mediated microtubule severing.

Collectively, these results suggest that Rtn-1 would counteract Spastin function locally in axons, impacting on outgrowth. Previous evidence points to a specific role for the Rtn-1C isoform in this potential interplay (Mannan et al., 2006). However, it is still unclear which Rtn-1 isoform is locally produced in axons.

The Reticulon-1 isoform C (Rtn-1C) is locally synthesized in cortical axons

We sought to characterize the isoform expression of *Rtn-1* mRNA and protein in both axons and cell bodies. Because microfluidic chambers yield only limited cellular material, we adopted an alternative culture approach using "neuronballs." This method enables the segregation of an axon-enriched fraction by mechanically separating axons from somato-dendritic structures. The axonal compartment of desomatized neuronballs was identified by IF against medium chain neurofilament (NF-M), while the somato-dendritic compartment was stained with microtubule-associated protein 2 (MAP2) (Figure 6A). RNA was then extracted from both the axonal and somato-dendritic fractions and submitted to reverse transcription PCR (RT-PCR). The purity of the axonal fraction was evidenced by the absence of amplification for *Map2*, *Histone h1(*general nuclei marker), and the astrocyte-specific glial fibrillary acidic protein (*Gfap*) mRNAs. In contrast, we found positive amplification for *β-actin* transcript, which is known to be present in axonal compartment (Figure 6A). These fractions were further analyzed with *Rtn-1* specific primers targeted against C-terminal domain or the N-terminal domain (to reveal *Rtn-1A*). In contrast to the somato-dendritic fraction, axons did not amplify for *Rtn-1A* mRNA, suggesting that *Rtn-1C* transcript is the predominant isoform in this compartment (Figure 6A).

Next, we investigated the protein composition of the axonal and somato-dendritic fractions of neuronballs by WB. Consistent with the sample purity observed for mRNAs, the axonal fraction was positive for Tau and β3-tubulin, with no detection of the dendritic marker MAP 2 (Figure 6A). For Rtn-1 identification, we employed a pan-Reticulon-1 antibody targeting the conserved C-terminal Reticulon homology domain (RHD). Importantly, the axonal fraction showed exclusivity for the lower molecular weight band, indicative of Rtn-1C presence, mirroring RT-PCR results (Figure 6A). Immunofluorescence analysis of Rtn-1C in axons from 8 DIV microfluidic chambers, displayed a distribution pattern consistent with that of the pan-Reticulon-1 (Rtn-1 antibody), as illustrated in Figure 2 (Figure 6B). In contrast, Rtn-1A presence was limited to a small subset of axons, exhibiting a peak of localization near the growth cone base (Figure 6B). Thus, resembling the distribution of transcripts, Rtn-1C protein appears as the predominant isoform in the axonal compartment.

In Figure 2 we provided evidence compatible with the axonal synthesis of Rtn-1 in cortical axons. In addition, we showed that the predominant mRNA and protein in cortical axons correspond to the Rtn-1C isoform (Figure 6A and B). However, whether Rtn-1C is indeed locally synthesized in cortical axons has not been proved.

To demonstrate the local production of Rtn-1C in axons, we employed a proximity
ligation assay integrated with puromycylation (puro-PLA) (Dieck et al., 2015). During this assay,
we selectively administered a puromycin pulse (15 minutes) to the axonal or somato-dendritic
compartment of microfluidic chambers, allowing its covalent incorporation into nascent
polypeptides. This enabled the detection of newly synthesized Rtn-1C peptides through
fluorescent signals produced by the nanometric proximity (<40 nm distance) of puromycin and
Rtn-1C antibodies (Figure 6C). Positive amplification was observed in the somato-dendritic
compartment of microfluidic chambers with local puromycin treatment as dots or puncta (Figure
6D, left panel). However, virtually no signal was detected in the cell bodies when puromycin was
added to the axonal compartment, showing the effective compartmentalization of this approach
(Figure 6D, left panel). Positive signal in the cell bodies was abrogated when the protein synthesis
inhibitor emetine was previously added, further supporting that these signals reflect protein
synthesis (Figure 6D, left panel). Supporting its axonal synthesis, fluorescent puncta were
observed in axons, which was significantly reduced by previous emetine treatment (Figure 6D,
right panel). Interestingly, when the puromycin pulse was applied exclusively to the somatic
compartment, positive signal was still detected in axons in a significantly lower amount compared
to direct and puromycin treatment, but higher than the emetine/puromycin condition (Figure 6D,
right panel). This suggests that at least a fraction of puromycylated peptides synthesized in the
cell bodies may rapidly (within 15 minutes) reach the axons through as-yet uncharacterized
mechanisms. Nevertheless, the significantly stronger signal detected in the axonal compartment
with direct puromycin treatment is fully consistent with local Rtn-1C synthesis in axons.
429

To further validate puro-PLA detection, we tested the Rtn-1C antibody specificity by WB
and IF, and whether puro-PLA signals depended on local depletion of *Rtn-1* mRNA (axonal Rtn-
1 KD). Due to their low background for neuronal components, we used COS-7 cells expressing
the rat Rtn-1C protein fused to GFP (Rtn-1C-GFP) and analyzed them by WB, using the same
Rtn-1C antibody of the puro-PLA experiment. As expected, a single band of the predicted
molecular weight for the recombinant protein was observed in the lane of transfected cells, while
no band was apparent in non-transfected cells (Supplementary Figure 3A). Also, revealing an
isoform-specific immunodetection, the ER pattern depicted by Rtn-1C-GFP fully overlapped with
the IF using the Rtn-1C antibody (Supplementary Figure 3B). Similar results were observed for
Rtn-1A-GFP expressing COS-7 cells stained against Rtn-1A antibody (Supplementary Figure
3B). Of note, when we performed crossed antibody staining of Rtn-1 isoforms on transfected
cells, no positive staining was found (Data not shown). We also tested the specificity of this
antibody by transfecting cells with siRNA 2 sequence. As expected, siRNA 2 induced a
significant decrease of Rtn-1C band compared to Scr (Supplementary Figure 3C). We also
validated the presence of *Rtn-1* mRNA in axons of similar conditions (9 DIV microfluidic

chamber) as shown by supplementary Figure 3D. Finally, 9 DIV axons were locally treated in
microfluidic chambers with siRNA 2 for twenty-four hours, which produced a significant
decrease of positive puro-PLA signal compared to Scr (Supplementary Figure 3D, right).
Together, these results confirm that the quantified puro-PLA signal is reflecting the local
translation of *Rtn-1C* mRNA in axons.
450
451   **Rtn-1C but not Rtn-1A reduces the microtubule severing activity of Spastin**
452
We established that Rtn-1C is locally synthesized in cortical axons, while Rtn-1A
synthesis should predominantly occur in the cell body. Therefore, we sought to understand the
functional implications of such compartmentalization. We chose COS-7 cells as a heterologous
system because of their increased co-transfection efficiency, flat and extended morphology, and
low background of neuronal proteins. These features provided an advantageous context to
visualize ER protein distribution and functional relationships.
459
Thus, we first over-expressed in COS-7 cells an RFP-tagged version of the most abundant
Spastin isoform during neuronal development, which corresponds to M87 isoform. Of note, in
contrast to M1 isoform, M87 isoform is expected to be soluble and devoid of the ER-shaping
membrane domain (Solowska et al., 2017). Next, the plasmid coding for isoform M87 was co-
transfected with Rtn-1C-GFP, Rtn-1A-GFP or GFP alone. Spastin-RFP displayed a punctate
distribution when co-expressed with GFP, indicative of its potential interaction with endosomal
components, as reported elsewhere (Allison et al., 2013) (Figure 7A, upper panels). Under this
condition, Spastin puncta strongly co-distributed with some clusters of soluble GFP. It has been
described that under certain circumstances soluble proteins such as GFP might be bulk up taken
by late endosomal structures (Sahu et al., 2011), which are also known to be positive for M87
Spastin (Allison et al., 2019). To rule out that, in our case, this co-distribution is due to
fluorescence crosstalk, we independently acquired images for each fluorophore in co-transfected
cells. We observed that COS-7 cells co-transfected with Spastin-RFP and GFP displayed the same
previously described puncta for both channels, discarding crosstalk (Supplementary Figure 5).
Further discarding unspecific channel bleeding, some cells that exclusively express Spastin-RFP,
harbor puncta that are exclusively seen in one channel, as indicated in supplementary Figure 5.
Similarly, some strong GFP puncta (indicated in the image) are not detected in the RFP channel
(Supplementary Figure 5). Complementary, cells co-transfected with Rtn-1A-RFP and GFP show
distinct distribution patterns without overlapping puncta, suggesting that the punctate distribution
of GFP is associated exclusively with Spastin co-expression and not explained by unspecific
bleeding (Supplementary Figure 5).
481

Of note, Spastin-RFP pattern was drastically altered when co-expressed with Rtn-1C-GFP (Figure 7B, middle panels). As such, and potentially compatible with their previously reported direct interaction, Spastin closely followed the Rtn-1C pattern distribution (Figure 7B, middle panels). Conversely, Rtn-1A-GFP showed minimal overlap with Spastin-RFP (Figure 7B, lower panels). Quantification of protein colocalization by Mander's coefficients substantiated these observations (Figure 7C).

Notably, the co-expression of Spastin-RFP with Rtn-1C-GFP maintained tubulin levels unaltered when compared to non-transfected cells from each experimental condition, in marked contrast with their GFP and Rtn-1A-GFP counterpart (Figure 7B and C). Furthermore, the filamentous distribution of tubulin was clearly preserved with Spastin-RFP/ Rtn-1C-GFP co-expression, which is a strong indicative of limited severing of microtubule filaments (Figure 7B, middle panels). These effects cannot be attributed to differences in transfection efficiency, as fluorescence intensity of Spastin-RFP and GFP constructs did not significantly differ across conditions (Supplementary Figure 5). Therefore, these results suggest that the tight co-distribution of Rtn-1C with Spastin is associated with reduced microtubule severing activity, which is not observed for the Rtn-1A isoform.

The examination of the sequences of Rtn-1C and 1A reveals a unique N-terminal variance as the point of differentiation between both isoforms, as they share their RHD region (Figure 7D). Considering the limited colocalization of Rtn-1A with Spastin (Figure 7B, lower panels), RHD appears not to be sufficient for their interaction. This observation hints at the exclusive 20 N-terminal amino acids of Rtn-1C as being critical for Spastin binding. Therefore, we performed *in silico* modeling of the Rtn-1C N-terminal and anticipated its affinity with the microtubule-binding domain (MTBD) of Spastin. This domain showed larger interaction energies compared to the endosome (Microtubule Interacting and Trafficking domain, MIT) and microtubule severing domain (ATPases Associated with diverse cellular Activities, AAA) (Figure 7E). Furthermore, the affinity of a Rtn-1A equivalent region for these domains was also less thermodynamically favored compared to Rtn-1C (Figure 7E). These results are compatible with previous evidence showing that deletion of Spastin's MIT domain, and most of the MTBD region, abrogates Spastin-Rtn-1C co-precipitation(Mannan et al., 2006). This suggests that Rtn-1C binding to Spastin MTBD is competing with the binding of this domain to microtubules. Mechanistically, this would prevent the ring conformation around microtubule filaments required for the severing activity of Spastin(Roll-Mecak & Vale, 2008). Overall, these findings reveal that Rtn-1C, but not Rtn-1A, closely associates with Spastin, leading to a reduction in its microtubule severing activity.

Discussion

This study uncovers a mechanism whereby downregulation of the local synthesis of the ER protein Rtn-1 promotes the outgrowth of injured cortical axons and microtubule dynamics. Collectively, our results suggest that this enhancement depends on the shielding effect that axonally synthesized Rtn-1C exerts over Spastin microtubule severing. Using this mechanism, cortical neurons may precisely and locally control outgrowth after injury. This finely tuned response after damage is intriguing given that its halting effect on outgrowth may seem, at first sight, detrimental for proper regeneration. However, we may speculate that limiting microtubule turnover at injury site helps axons to accumulate molecular components required for a successful injury repair. In any case, it would be interesting to explore whether local synthesis of Rtn-1 offers a potential therapeutic pathway to enhance regeneration by unleashing its shielding effect over Spastin locally in axons.

Control of Rtn-1 synthesis as an adaptive mechanism for axonal injury.

It would also be interesting to further explore whether developing neurons decrease axonal *Rtn-1* mRNA translation as a mechanism to promote outgrowth during development and after damage. For instance, a previous study using axotomized 10 DIV rat cortical neurons in microfluidic chambers found a decrease on *Rtn-1* mRNA levels compared to non-injured axons (Taylor et al., 2009). This reported reduction may also imply that injured axons do not harbor enough *Rtn-1* mRNA to support local synthesis, which would modify the interpretation of our post-injury results. Still, the observed similarity between specific phenotypes before and after injury, including decrease in distal β3-tubulin levels and increased microtubule growth rate are compatible with Rtn-1C synthesis occurring before and after injury. A more definite answer might be provided by performing puro-PLA measurements of Rtn-1C synthesis before and after injury, which is an ongoing work.

Currently, it is unknown to what extent a similar mechanism is retained in adult cortical neurons. Likewise, we do not know whether this phenomenon is fully operative *in vivo,* even though *Rtn-1* mRNA is consistently found *in vitro* and *ex vivo*. Of note, *Rtn-1* mRNA is dynamically regulated in the axonal translatome of retinal ganglion cells *in vivo* (Shigeoka et al., 2016), but the functional implications of axonal Rtn-1 synthesis in a more physiological context are yet to be elucidated. Interestingly, growing axonal tips from injured nerves in Lamprey are nearly 13-fold enriched in *Spastin* mRNA, suggesting that other constituent of our described mechanism may be locally synthesized after injury (Jin et al., 2022).

The presence of the axonal ER (particularly the smooth ER) is thought to be critical for promoting the regeneration of injured neurites by providing membranes and controlling calcium and microtubule dynamics required for outgrowth (Luarte et al., 2018). This has been supported by evidence showing the distribution of smooth ER markers, including ER-shaping proteins such as Reticulons, REEPs, Atlastin, and Spastin isoforms, preferentially at the distal tip of regenerating neurites (Rao et al., 2016; Stone et al., 2012). Indeed, Reticulons share RHD with REEPs, some Spastin isoforms, and Atlastins, which makes them suitable to shape the mostly tubular smooth ER (Chiurchiù et al., 2014; Iwahashi et al., 2007). Highlighting their functional relevance, forcing the localization of smooth ER into axonal tips by the over-expression of the ER membrane protein Protrudin enhances outgrowth of CNS axons (Petrova et al., 2020).

Different studies have shown that the presence of the axonal ER is associated with the fine-tuning of microtubules stability to sustain axonal extension (Farías et al., 2019; Lee et al., 2009). Our results support a model where axonal outgrowth after injury is restricted by reducing the microtubule growth processivity (reflected by track length), which might be controlled by locally synthesized Rtn-1C. As such, axonal KD of *Rtn-1C* mRNA may set an optimal range of microtubule growth, resembling the beneficial role observed with the treatment of low-dose microtubule stabilizers in axonal regeneration (Brizuela et al., 2015; B. Liu et al., 2023). These results would explain why we observed a trend (yet non-significant) of SPTZ to increase regeneration when applied alone locally in axons (Figure 5C-D). The opposite trend observed with the SPTZ+ Rtn-1 siRNA condition highlights the specificity of this response, pointing that microtubule stability has a relatively narrow optimal range to promote outgrowth (Sengottuvel et al., 2011).

Some studies have pointed to a non-canonical role for Rtn-1C synthesis by signaling in the nucleus, potentially through DNA binding and histone deacetylase inhibition (Nepravishta et al., 2010, 2012). It is tempting to speculate that these still emerging roles may also contribute to the observed phenotypes. Of note, different axonally synthesized proteins exert transcriptional control after in response to injury or local cues (Twiss et al., 2016).

The differential injury response in axonal outgrowth observed after somatic or axonal Rtn-1 KD, emphasizes the unique molecular and structural environments of axons versus cell bodies. This difference might be explained by at least two scenarios. First, we used a Rtn-1 siRNA that is directed against the conserved RHD sequence; thus, it also targets the Rtn-1A isoform. For instance, we observed limited co-distribution of Rtn-1A with Spastin compared to Rtn-1C, and its N- terminal recruits other signaling pathways (Kaya et al., 2013). Therefore, the observed differential response might be due to the predominant presence of Rtn-1C in axons, while both isoforms are present in the cell bodies. Secondly, increased microtubule severing after axonal

589 Rtn-1 KD might promote outgrowth when coupled to predominantly outward-directed growing
590 filaments that characterize axons, but not with the radial or mixed orientation of microtubules in
591 the cell bodies and dendrites(Baas et al., 2016). Interestingly, global KD of Rtn-1 in developing
592 cortical neurons decreases axonal outgrowth (Nozumi et al., 2009), further supporting the
593 relevance understanding this compartmentalized response.

594 The role of Spastin's microtubule severing on outgrowth after injury

The positive relationship that we observed between increased Spastin-severing activity
and the outgrowth of injured axons may seem counter-intuitive. However, microtubule severases,
such as Spastin or Katanin, are known to increase the microtubule filament mass and growth
under certain conditions. Indeed, under certain conditions Spastin can stabilize newly produced
microtubule tips, paradoxically yielding a net increase in microtubule mass (Kuo & Howard,
2021). Furthermore, Spastin tends to concentrate at the tip of growing microtubules to behave as
a dual enzyme that cleaves but also stabilizes them after cutting (Kuo et al., 2019). This
stabilization property may also partially explain why we observed that the axonal inhibition of
Spastin by SPTZ alone tends to promote outgrowth after injury (Figure 5C-D).

Collectively, our results might be compatible with axonal Rtn-1 KD releasing Spastin
molecules to sever more microtubules, and thus, potentially promote a net increase in filament
growing ends and mass. In agreement with this speculation, inhibition of axonal tyrosine ligase
(which favor dynamic microtubule ends) impedes outgrowth in injured axons from dorsal root
ganglion (Song et al., 2015). Analogously, when Fidgetin (a severing enzyme of microtubule
labile ends) is depleted in neurons, axonal regeneration is improved (Matamoros et al., 2019).

Like other studies, we observed that increased axonal outgrowth correlates with dynamic
microtubule parameters such as growth rate (Liz et al., 2014). However, others have shown that
the dynamic properties of microtubules, including microtubule growth rate and track length, can
adapt to different outgrowth scenarios, with low capacity to predict axonal extension (Kiss et al.,
2018). Thus, controlling dynamic parameters of microtubules is not the only factor that
determines the outgrowth capacity of injured axons. The exact mechanisms by which locally
synthesized Rtn-1C controls dynamic parameters of microtubules remains unsolved as they seem
not to rely on Spastin severing activity. Other Spastin-independent interactions of Rtn-1 may
depend on its presence on the tubular ER, which includes the control of functional contacts with
mitochondria and autophagosomes (D' Eletto et al., 2019; Reali et al., 2015), but their
implications for microtubule dynamic parameters and axonal extension also remain to be solved.

Decreasing axonal Rtn-1C translation or increasing Spastin microtubule severing locally
in axons may constitute an emerging mechanism for promoting injury recovery. For instance,
local application of recombinant Spastin together with polyethylene glycol increases recovery

after sciatic nerve injury (Y. F. Lin et al., 2019). Furthermore, local augmentation of axonal

Spastin by the pharmacological local regulation of 14-3-3 promotes the recovery of spinal cord

injury (Q. Liu et al., 2024) . Indeed, axonal regeneration is particularly sensitive to the absence of

Spastin (Stone et al., 2012). Similarly, results in drosophila show that Spastin gene dosage is

particularly relevant for the regenerative response of axons (Stone et al., 2012) .

Implications for length-dependent axonopathies and injury response

The failure of ER shaping protein-function potentially participates in the etiology of

multiple length-dependent axonopathies such as hereditary spastic paraplegias (HSPs)

(Blackstone et al., 2011). Here, we have described an emerging mechanism relating Rtn-1C with

the activity of Spastin, which is the most frequently mutated isoform in HSP (Hazan et al., 1999;

Mannan et al., 2006). As a second point of analysis, most of the studies performed to understand

axonopathies and axonal damage have focused on a "cell-body-centered view" for protein

synthesis and ER function. However, increasing evidence supports the relevance of axonal

autonomous response to injury or cues from the local brain microenvironment (Jung et al., 2014;

Lopez-Verrilli et al., 2013; Zhang et al., 2013, 2015). Similarly, our work underscores a highly

local injury adaptation, which is in harmony with the astonishing extension of central nervous

system's axons (H.-M. Lin et al., 2018). These results may provide understanding of axonal

pathologies and injury response with a special focus on the autonomous response of cortical

axons.

Materials and Methods

Animals and surgical procedures. Pregnant Sprague–Dawley rats at embryonic day 18 were

obtained from the Central Animal Facility of Universidad Católica de Chile and euthanized in

accordance with the guidelines of the Agencia Nacional de Investigación y Desarrollo de Chile

(ANID). All the experimental the procedures were approved by the Institutional Bioethics

Committee of Universidad de los Andes (CEC2022106).

Neuron cell cultures. Cortical neurons were cultured from 18-day embryos like previously

reported (Deitch & Banker, 1993). Dissociated primary cortical neurons (20,000-30,000 cells)

were seeded on glass coverslips coated with 0.1 mg/mL poly-L-lysine (Sigma-Aldrich, P4707).

Microfluidic chambers were prepared as previously reported (Taylor et al., 2005). Chambers

harboring axonal and somatic compartment connected with 750 μm long microgrooves were

mounted over a squared 25 mm glass coverslip previously coated with 0.1 mg/mL poly-D-lysine

(Corning, 354210) and 2 μg/mL laminin (Sigma, L2020). Plating medium was completely

replaced with growth medium 24 hours after seeding. Plating medium contained MEM (Gibco,

15630080), 10 % horse serum (Gibco, 16050122), 0.6 % D-glucose (Sigma Aldrich, G8769), 0.1

% Glutamax (Gibco, 35050061), 2 % penicillin/ 0.1 % streptomycin (Life
Technologies,15140122), while growth medium contained Neurobasal (Gibco, 21103049), 3 %
B27 (Gibco, 17504044), 0.1 % Glutamax, penicillin/streptomycin 0.1 %). An amount of 100 µL
of fresh growth medium was added with a frequency of 48 hours to the somatic compartment to
prevent loss of water. Neuronball cell culture was performed as previously reported (Sasaki et al.,
2010).Neurons were suspended on growth medium supplemented with 10 % horse serum. A total
amount of 30,000 cells was seeded as hanging drops of 20 µL during 3 days inside the cover of
100 mm dishes filled with water. Floating spheres were gently washed three times to discard non-
attached cells and then seeded on poly-L-lysine coated culture plates. Each well contained 400
µL of growth medium previously supplemented with 1 µM cytosine β-D-arabinofuranoside
hydrochloride (AraC, Sigma-Aldrich, C1768) to prevent glial growth. Neurites were allowed to
grow until 7 days after seeding. The periphery of each neuronballs was carefully scraped with a
pipette and washed three times with cold PBS to further remove any remaining cell bodies. Cell-
body-containing spheres were extracted by vacuum suction.
675
Immunofluorescence. Cells were washed twice with PBS and fixed with 4% paraformaldehyde
(PFA, Sigma Aldrich, P6148) and 4% sucrose in PBS for 20 min at room temperature. The same
procedure was applied to COS-7 cells. For assessment of growth cone morphology, neurons were
fixed for 20 min at room temperature by adding an equal volume of 8% PFA and 22% sucrose in
PBS directly to the culture medium. After fixation, cells were washed twice with PBS and
sequentially incubated for 10 min each at room temperature with: (i) permeabilization solution
(0.5% NP-40 [Abcam, ab142227] and 0.5% w/v Bovine Serum Albumin [Winkler, BM-0150] in
PBS); (ii) quenching solution (50 mM NH_4Cl in PBS); and (iii) blocking solution (0.5% w/v
Bovine Serum Albumin and 5% Fetal Bovine Serum [Sigma Aldrich, F2442] in PBS). Primary
antibodies, diluted in 0.5% w/v Bovine Serum Albumin in PBS, were incubated overnight at 4
°C. Coverslips were then washed three times (5 min each) with the same solution and incubated
with secondary antibodies, diluted in the same buffer, for 1 h at room temperature. Finally,
samples were washed with PBS and distilled water and mounted using VECTASHIELD®.
689
Growth cone morphometry. Growth cones (GCs) were imaged after β3-tubulin and F-actin
labeling with phalloidin using identical acquisition settings across conditions. Analyses were
performed in FIJI (ImageJ). The actin outline was thresholded (same parameters for all images)
to isolate single binarized GCs at distal axon tips (axon shaft excluded). From the binarized GC
mask we computed: area (µm²), perimeter (µm), and elongation ratio (major axis/minor axis of
the best-fit ellipse to the actin boundary). For skeleton-based metrics, the binarized GC mask was
skeletonized (FIJI Skeletonize) to obtain a one-pixel-wide graph of the actin scaffold. We then
used Analyze Skeleton (2D/3D) to extract (i) LSP (Longest Shortest Path), defined as the

maximum geodesic distance (in μm) between any two skeleton nodes within the GC—i.e., the
longest of all pairwise shortest paths along the skeleton—and (ii) number of filopodia, defined as
the count of terminal branches (end-points in the skeleton) that extend from the GC body and
exceed a minimum length of 1.0 μm (same length threshold applied to all conditions). When
needed, small spurious branches produced by noise were pruned before counting. All
measurements were performed blinded to condition.

Colocalization analysis. COS-7 cells were transfected with indicated plasmids for twenty-four
hours and imaged using a confocal microscope (Fluoview FV1000, Olympus) equipped with a
UPLSAPO 60×/1.35 NA objective and 4× digital zoom. Images were acquired with a Kalman
filter (two iterations). For each condition, an average of 8–12 confocal sections (0.2 μm thickness
each) were analyzed per cell. Colocalization between RFP and GFP signals was quantified using
Manders' coefficients (M1 and M2) with the JaCoP plugin (Just Another Colocalization Plugin)
in ImageJ (FIJI). Identical thresholding parameters were applied to each channel across all
conditions.

Fluorescence intensity analysis of COS-7. COS-7 cells were imaged with a Leica TCS SP8
confocal microscope using a 40×/1.30 NA oil-immersion objective. For each field, z-stacks of
15–17 optical sections (1 μm step size) were acquired, and summed intensity projections were
generated for analysis. Acquisition parameters (laser power, detector gain, and offset) were kept
constant across all conditions. Cells were manually segmented in FIJI using α-tubulin
immunostaining to delineate them. For each ROI, mean fluorescence intensity of β3-tubulin, GFP,
and RFP channels was measured, and background signal (from a cell-free region of the same
image) was subtracted. Corrected mean intensity was multiplied by the cell area to calculate total
fluorescence per cell. Fold change values were obtained by normalizing to the mean of the control
group.

Fluorescence intensity analysis of somas and axons. Images were acquired using an Olympus
BX61WI microscope in wide-field configuration equipped with a 40×/1.30 NA oil-immersion
objective and a Hamamatsu ORCA-R2 CCD camera, controlled by Cell^R software (Olympus,
Tokyo, Japan). Exposure time, gain, binning, and illumination settings were kept constant across
all conditions. For somatic quantifications, cell bodies were manually segmented in FIJI using
β3-tubulin staining as a guide, restricting the region of interest (ROI) to the soma and excluding
processes. For each cell, the mean background intensity (measured in a cell-free region of the
same image) was subtracted from the mean somatic intensity, and total fluorescence (integrated
intensity) was calculated by multiplying by the corresponding cell area. Fold change per cell was
determined by normalizing to the mean value of the corresponding control group. For axonal

quantifications, a 1-µm-wide segmented line was drawn starting at the axon tip and extending 50 µm proximally to extract the intensity profile (FIJI). Background intensity was subtracted from each axon profile using values measured in axon-free areas of the same image. For quantifying protein changes, the integrated intensity across the entire 0–50 µm segment was calculated and expressed as fold change relative to controls. To minimize intensity variability across different experimental dates, while preserving the relative differences between conditions, we plotted the corrected intensity profiles. For β3-tubulin, a per-axon baseline was defined by subtracting the intensity at 0 µm (the axon tip) from the entire profile. For Rtn-1, the baseline was defined as the mean pixel intensity within the distal 5 µm of the profile (45–50 µm from the tip), which was subtracted from the entire trace of each axon. These normalizations were applied solely for visualization purposes and were not used in fold change calculations.

Stimulated emission depletion (STED) microscopy analysis of axons. Super-resolution imaging was performed on TCS SP8-3X STED microscope system (Leica Microsystems) equipped with a HC APO CS2 100x/1.40 oil objective. Distal axons of microfluidic chambers were imaged using excitation and depletion laser lines appropriate for each fluorophore (see Reagents in supplementary material). Images were digitized in 1024 × 1024 pixels format file with 5x zoom factor. These settings achieved 18.93 nm resolution in the XY plane and a 220 nm resolution in Z. Four confocal slices from each axon were sum-projected and analyzed. No deconvolution or post-processing algorithms were applied. Instead, all quantitative and qualitative analyses were performed directly on the raw STED images. Acquisition parameters (laser power, detector gain, pixel size, and dwell time) were kept constant across conditions. For quantification, the total fluorescence intensity of the last 15 µm of each distal axon was measured after background subtraction. Two analyses were performed: (i) intensity fold change was calculated by comparing non-axotomized and axotomized axons; and (ii) the relative intensity, which was obtained by dividing the intensity of each pixel by the total intensity of the corresponding staining.

Transfections. All transfections were performed Lipofectamine™ 3000 (Thermo Fisher, L3000001) following manufacturer's instructions but without using p3000 reagent. For neurons and COS-7 cells, RNA or DNA complexes were added to chambers diluted on pure Neurobasal medium. For axonal transfection of siRNAs, medium on the axonal compartment was carefully removed and replaced with 50 µL of the transfection mixture containing 20 pmol of the corresponding oligonucleotide plus 50 µL of fresh growth media. ~300 µL of conditioned medium remained at the somatic compartment, generating positive flux towards the axons. Conversely, for somatic transfection the medium present at the somatic compartment was carefully removed and replaced with 50 µL of the transfection mixture plus 100 µL of fresh growth media, this time generating positive flux towards the soma. For validating siRNA efficiency, 200,000 neurons

were seeded on 35 mm plates and transfected with 40 pmol of corresponding oligos and submitted
to Western blot after forty-eight hours. For colocalization, Western blot, and immunofluorescence
analyses, COS-7 cells were seeded at ~1–2 × 10^5 per 25 mm coverslip and cultured for 24 h before
transfection with 500 ng of pCMV3-C-GFPSpark plasmid encoding either Rtn-1C-GFP or Rtn-
1A-GFP, together with pCMV3-C-ORFSpark encoding Spastin-RFP (all plasmids are detailed in
the Reagents section of the Supplementary material).

Axotomy and Sholl analysis. At 8 DIV, axons were removed from the axonal compartment of
microfluidic chambers by vacuum aspiration performed five times with 100 µL of pure
Neurobasal medium. Subsequently, all media was removed, and the transfection mixture was
added either to the axonal or somatic compartment. For quantification of axonal outgrowth, β3-
tubulin–stained axons were identically thresholded across conditions, binarized, and curated to
fully trace axonal trajectories. The Sholl analysis was performed in FIJI using the NeuroAnatomy
plugin. Concentric semi-circles were applied at 2 µm radial intervals, starting from the ends of
the microgrooves and centered on axonal fields of equivalent size. The analysis extended to a
maximum radial distance of either 1200 µm or 1500 µm, depending on the experiment.

Probe-labeling of primary antibodies for puromycylation proximity ligation assay (puro-
PLA). Monoclonal antibodies mouse anti-Reticulon-1C and mouse anti-puromycin (3RH1,
Kerafast) were directly labeled with the Duolink® In Situ Probemaker™ PLUS (1002748995,
Sigma Aldrich) and MINUS (1002748992, Sigma Aldrich), respectively, following
manufacturer's instructions. Both antibodies with a concentration of 1 µg/µL were labeled
overnight at room temperature with the corresponding buffer and probes to a final volume of 20
µL. Labeled antibodies were further diluted with 20 µL of storage solution.

Puro-PLA staining protocol. Puro-PLA was under conditions similar to well-established
protocols (Dieck et al., 2015). Briefly, a 15-min pulse of 1.8 µM puromycin (P8833, Sigma-
Aldrich) was applied exclusively to the axonal or somatic compartment of microfluidic chambers
containing 9 DIV cortical neurons. As an additional control, this incubation was performed after
90 min of local pretreatment with 40 nM emetine (E2375, Sigma-Aldrich) in distilled water
(vehicle), added either to the somatic or axonal compartment to block protein synthesis.
Immediately after incubations, cells were washed twice with PBS and fixed with 4% PFA and
4% sucrose in PBS for 20 min at room temperature, followed by three PBS washes (5 min each).
Cells were then permeabilized with 0.25% Triton X-100 in PBS for 10 min, washed three times
with PBS (5 min each), blocked with 5% goat serum in PBS for 1 h, and incubated overnight at 4
°C with probe-labeled antibodies (anti–Reticulon-1C, 1:50; anti-puromycin, 1:200) diluted in the
same blocking solution. Detection of locally synthesized Reticulon-1C was performed following

the manufacturer's instructions, except that no labeled secondary antibody step was included.

Afterward, cells were incubated for 1 h at room temperature with anti-β3-tubulin conjugated to

Alexa Fluor 488 and mounted with Duolink® In Situ Mounting Medium with DAPI (Sigma-

Aldrich). Images were acquired as previously described using a Zeiss LSM 800 confocal

microscope with a 63× oil objective. PLA-positive puncta present within a β3-tubulin mask that

defined the axonal area were quantified using FIJI software.

RT-PCR from neuronballs and microfluidic chambers. For axonal RNA isolation from

neuronballs, at least 200 balls were desomatized and processed with RNAqueous™-Micro Total

RNA Isolation Kit (AM193, Thermo Fisher), following the manufacturer's instructions.

Approximately 20–30 axotomized neuronballs (somatic RNA) were processed using the same

protocol. Reverse transcription was performed from 100 ng of RNA as input using the High-

Capacity cDNA Reverse Transcription Kit (4368814, Thermo Fisher). For negative controls,

axonal samples were submitted to the same protocol, but without adding reverse transcriptase.

Resulting cDNA products were treated with DNase according to the manufacturer's instructions

(EN0521, Thermo Fisher). For PCR, cDNA was amplified using the GoTaq® Green Master Mix,

2X (M712, Promega), with the following cycling parameters: initial denaturation at 95 °C for 5

min; 35 cycles of denaturation at 95 °C for 30 s, annealing at 57 °C for 30 s, and extension at 72

°C for 45 s; followed by a final extension at 72 °C for 10 min. PCR products were loaded on a

2% agarose gel and stained with GelRed® Nucleic Acid Gel Stain (Biotium, 41003), following

the manufacturer's recommendations. We performed the RNA isolation from the axonal/somatic

compartments of microfluidic chambers (7-8 chambers) as previously reported (Taylor et al.,

2005).For the subsequent steps, we followed the same procedures as in neuronballs, but

performing 42 PCR cycles.

Western blot. Proteins were isolated using RIPA buffer (50 mM Tris-HCl, pH 7.4, 150 mM

NaCl, 0.25% sodium deoxycholate, 1% NP-40, 1% SDS, 1 mM EDTA) supplemented with

protease and phosphatase inhibitors (Thermo Fisher, A32963). Extracted proteins were quantified

using the Bicinchoninic Acid (BCA) method (Thermo Fisher, 23225) and separated on

polyacrylamide gels under denaturing conditions (SDS-PAGE; Cleveland et al., 1977). For each

sample, 20 μg of protein was loaded per lane for both complete neuronal or cell homogenates,

and axonal or somatic fractions of neuronballs. Proteins were resolved on 15% acrylamide gels

by electrophoresis at 70 V for 45 min followed by 100 V for 45 min. Gels were transferred to

nitrocellulose membranes (Bio-Rad) at a constant current of 350 mA for 90 min. After transfer,

membranes were blocked with 5% (w/v) skim milk in PBS for one hour at room temperature

under constant agitation and then incubated overnight at 4 °C with the corresponding primary

antibody diluted in PBS. Membranes were incubated with the appropriate secondary antibody for

one hour at room temperature. When indicated, total protein loading was normalized using No-

Stain™ total protein labeling reagent (Thermo Fisher, A44449) according to the manufacturer's

instructions. Bands and lanes were quantified by densitometry using Photoshop 7.0 (Adobe).

Live imaging of microtubule plus-ends. Cortical neurons were cultured in microfluidic

chambers and transfected with EB3-Emerald GFP plasmid at 7 DIV. At 8 DIV, neurons were

axotomized by vacuum aspiration, and siRNAs were transfected into the axonal compartment

.The Spastin inhibitor Spastazoline (HY-111548, MedChemExpress) was added to the axonal

compartment 4–5 h after axotomy at a final concentration of 50 nM. Forty-four hours after

transfection, axons expressing EB3-GFP were randomly selected and imaged in Tyrode's solution

(125 mM NaCl, 2 mM KCl, 2 mM CaCl$_2$, 2 mM MgCl$_2$, 30 mM glucose, 25 mM HEPES, pH 7.4)

at 32 °C. Cells were equilibrated for 10 min before recording. Movies were acquired for 3 min at

2 frames per second (fps). To maintain hydrostatic pressure towards the axonal side, the volume

difference between compartments was carefully preserved throughout the experiment. Imaging

was performed on an inverted microscope (Observer.D1, Zeiss) equipped with an EMCCD

camera (Evolve 512, Photometrics) controlled by VisiView® software (Visitron Systems GmbH).

A 63× oil-immersion objective and a GFP single-band exciter ET filter set (excitation 470/40,

emission 59022m, dichroic 59022BS) were used. The mEmerald-EB3-C-20 plasmid was a gift

from Michael Davidson (Addgene plasmid #54076; http://n2t.net/addgene:54076; RRID:

Addgene_54076).

Analysis of microtubule dynamics. All acquired time series were filtered using the Gaussian

Blur in FIJI (applied to temporal frames). Mobile components of the series were isolated by

applying a 4-sigma Gaussian blur, while static elements were extracted by applying a 50-sigma

blur. To enhance detection of moving structures, static components were subtracted from mobile

components using the *Image Calculator* function in FIJI. The resulting filtered image series were

then used to generate kymographs with the FIJI plugin *Kymograph Builder*. Kymographs were

constructed from the distal portion of axons by tracing a 1-µm-wide line, with a total length

ranging from 30 to 50 µm. Growth cones were excluded from the analysis. All analyzed

parameters were quantified as previously described (Stepanova et al., 2010). EB3-GFP comets

were analyzed from kymographs in FIJI. Individual dashes corresponding to microtubule growth

events were manually traced using the Segmented Line tool. The angle of each dash was measured

relative to the x-axis. The horizontal projection of the dash, obtained by multiplying its length by

the cosine of the angle, represents the distance traveled by the comet in micrometers (track

length). The vertical projection, obtained by multiplying the dash length by the sine of the angle,

was proportional to comet lifetime. To obtain comet lifetime in seconds, vertical projection was

divided by the total y-axis length of the recording kymograph and multiplied by the total

acquisition time (180 seconds for a 3-minute movie). Growth rate was calculated by dividing the
track length by the comet lifetime for each dash.

Molecular docking between Spastin (M87) and Rtn-1C. First, Spastin M87 and Rtn-1A models
of *Rattus norvegicus* were retrieved from AlphaFold Protein Structure Database (accession
numbers AF-A0A0G2K590 and AF-Q64548, respectively). Then, M87 spastin model were
generated from the removal of the first 86 amino acids of the M1 Spastin model, whereas Rtn-1C
3D-structure was built with MODELLER version 10.4 using an Rtn-1A model as template. For
this, amino acid sequences of Rtn-1C were retrieved from UniProtKB/Swiss-Prot (accession
numbers Q64548). Additionally, the first 20 amino acids (of the N-terminal region) of the Rtn-
1C model and region 200-317 containing the microtubule-binding domain of Spastin M87 were
refined using ModLoop server (Fiser & Sali, 2003). Subsequently, we evaluated the interaction
of the Rtn-1C N-terminal with three domains of Spastin M87- endosome or MIT (region 116-
194), microtubule-binding (region 270-328) and microtubule-severing o AAA ATPase (region
342-599) - using the HADDOCK docking protein-protein web server(Van Zundert et al., 2016).
Among many models, low-energy conformations for each complex were extracted from
HADDOCK 2.2 server, as previously described (Acevedo et al., 2018) The binding energy
(ΔG_bind) of the complexes were predicted using the PRODIGY web server at a temperature of
310 K, and chain identifiers for Rtn-1 and Spastin M87 were specified as Interactor 1 and 2,
respectively. Finally, graphical analysis of molecular docking studies was performed using VMD
1.9.2 software(Humphrey et al., 1996) .

Statistical Analysis. Unless otherwise stated, data are presented as mean ± standard error.
Normality was assessed using the Shapiro–Wilk test. For normally distributed data, comparisons
between two groups were performed using two-tailed Student's t-test; for non-normally
distributed data, the two-tailed Mann–Whitney *U t*est was applied. Western blot fold change
values were compared to a theoretical mean of 1 using the Wilcoxon signed-rank test. For
comparisons among multiple groups, one-way ANOVA followed by Tukey's post hoc test was
used for normally distributed data, whereas the Kruskal–Wallis test followed by Dunn's post hoc
test was applied for non-parametric data. Statistical significance was defined as $p < 0.05$.
Fluorescence intensity experiments in Figures 2 and 4 were analyzed at the single-cell level using
linear mixed-effects (LME) models, with experimental date included as a random effect.
Statistical tests were performed using GraphPad Prism (GraphPad Software, San Diego, CA,
USA) and R (v4.0 or later) with the lme4 and lmerTest packages.

References

Acevedo, W., Ramírez-Sarmiento, C. A., & Agosin, E. (2018). Identifying the interactions between natural, non-caloric sweeteners and the human sweet receptor by molecular docking. *Food Chemistry*, *264*, 164–171. https://doi.org/10.1016/J.FOODCHEM.2018.04.113

Aiken, J., & Holzbaur, E. L. F. (2024). Spastin locally amplifies microtubule dynamics to pattern the axon for presynaptic cargo delivery. *BioRxiv*, *123*(3), 332a. https://doi.org/10.1101/2023.08.08.552320

Allison, R., Edgar, J. R., & Reid, E. (2019). Spastin MIT Domain Disease-Associated Mutations Disrupt Lysosomal Function. *Frontiers in Neuroscience*, *13*, 486039. https://doi.org/10.3389/FNINS.2019.01179/BIBTEX

Allison, R., Lumb, J. H., Fassier, C., Connell, J. W., Martin, D. Ten, Seaman, M. N. J., Hazan, J., & Reid, E. (2013). An ESCRT-spastin interaction promotes fission of recycling tubules from the endosome. *Journal of Cell Biology*, *202*(3), 527–543. https://doi.org/10.1083/JCB.201211045/VIDEO-2

Aridor, M., & Fish, K. N. (2009). Selective Targeting of ER Exit Sites Supports Axon Development. *Traffic*, *10*(11), 1669–1684. https://doi.org/10.1111/j.1600-0854.2009.00974.x

Baas, P. W., Rao, A. N., Matamoros, A. J., & Leo, L. (2016). Stability properties of neuronal microtubules. *Cytoskeleton (Hoboken, N.J.)*, *73*(9), 442. https://doi.org/10.1002/CM.21286

Baleriola, J., Walker, C. A., Jean, Y. Y., Crary, J. F., Troy, C. M., Nagy, P. L., & Hengst, U. (2014). Axonally synthesized ATF4 transmits a neurodegenerative signal across brain regions. *Cell*, *158*(5), 1159–1172. https://doi.org/10.1016/J.CELL.2014.07.001

Batista, A. F. R., Martínez, J. C., & Hengst, U. (2017). Intra-axonal Synthesis of SNAP25 Is Required for the Formation of Presynaptic Terminals. *Cell Reports*. https://doi.org/10.1016/j.celrep.2017.08.097

Bigler, R. L., Kamande, J. W., Dumitru, R., Niedringhaus, M., & Taylor, A. M. (2017). Messenger RNAs localized to distal projections of human stem cell derived neurons. *Scientific Reports 2017 7:1*, *7*(1), 1–11. https://doi.org/10.1038/s41598-017-00676-w

Blackstone, C., Kane, C. J. O., Reid, E., O'Kane, C. J., & Reid, E. (2011). Hereditary spastic paraplegias: membrane traffic and the motor pathway. *Nature Reviews. Neuroscience*, *12*(1), 31–42. https://doi.org/10.1038/nrn2946

Brizuela, M., Blizzard, C. A., Chuckowree, J. A., Dawkins, E., Gasperini, R. J., Young, K. M., & Dickson, T. C. (2015). The microtubule-stabilizing drug Epothilone D increases axonal sprouting following transection injury in vitro. *Molecular and Cellular Neuroscience*, *66*(PB), 129–140. https://doi.org/10.1016/j.mcn.2015.02.006

Chiurchiù, V., Maccarrone, M., & Orlacchio, A. (2014). The role of reticulons in neurodegenerative diseases. *Neuromolecular Medicine*, *16*(1), 3–15. https://doi.org/10.1007/s12017-013-8271-9

Cooke, P., Janowitz, H., & Dougherty, S. E. (2022). Neuronal Redevelopment and the Regeneration of Neuromodulatory Axons in the Adult Mammalian Central Nervous System. *Frontiers in Cellular Neuroscience*, *16*, 872501. https://doi.org/10.3389/FNCEL.2022.872501/BIBTEX

Cornejo, V. H., Luarte, A., & Couve, A. (2017). Global and local mechanisms sustain axonal proteostasis of transmembrane proteins. *Traffic*, *18*(5), 255–266. https://doi.org/10.1111/tra.12472

Cupido, T., Pisa, R., Kelley, M. E., & Kapoor, T. M. (2019). Designing a chemical inhibitor of the AAA protein spastin using active site mutations. *Nature Chemical Biology*, *15*(5), 444. https://doi.org/10.1038/S41589-019-0225-6

D' Eletto, M., Risuglia, A., Oliverio, S., Mehdawy, B., Nardacci, R., Bordi, M., & Di Sano, F. (2019). Modulation of autophagy by RTN-1C: role in autophagosome biogenesis. *Cell Death & Disease*, *10*(12). https://doi.org/10.1038/S41419-019-2099-7

de Juan-Sanz, J., Holt, G. T., Schreiter, E. R., de Juan, F., Kim, D. S., & Ryan, T. A. (2017). Axonal Endoplasmic Reticulum Ca2+ Content Controls Release Probability in CNS Nerve Terminals. *Neuron*, *93*(4), 867-881.e6. https://doi.org/10.1016/J.NEURON.2017.01.010

Deitch, J. S., & Banker, G. A. (1993). An electron microscopic analysis of hippocampal neurons developing in culture: early stages in the emergence of polarity. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *13*(10), 4301–4315. https://doi.org/10.1523/jneurosci.1544-09.2009

Dieck, S. tom, Kochen, L., Hanus, C., Bartnik, I., Nassim-Assir, B., Merk, K., Mosler, T., Garg, S., Bunse, S., Tirrell, D. A., & Schuman, E. M. (2015). Direct visualization of identified and newly synthesized proteins in situ. *Nature Methods*. https://doi.org/10.1038/nmeth.3319

Evans, K., Keller, C., Pavur, K., Glasgow, K., Conn, B., & Lauring, B. (2006). Interaction of two hereditary spastic paraplegia gene products, spastin and atlastin, suggests a common pathway for axonal maintenance. *Proceedings of the National Academy of Sciences of the United States of America*. https://doi.org/10.1073/pnas.0510863103

Fan, X. xuan, Hao, Y. ying, Guo, S. wen, Zhao, X. ping, Xiang, Y., Feng, F. xue, Liang, G. ting, & Dong, Y. wei. (2018). Knockdown of RTN1-C attenuates traumatic neuronal injury through regulating intracellular Ca2+ homeostasis. *Neurochemistry International*, *121*, 19–25. https://doi.org/10.1016/J.NEUINT.2018.10.018

Farías, G. G., Fréal, A., Tortosa, E., Stucchi, R., Pan, X., Portegies, S., Will, L., Altelaar, M., & Hoogenraad, C. C. (2019). Feedback-Driven Mechanisms between Microtubules and the Endoplasmic Reticulum Instruct Neuronal Polarity. *Neuron*. https://doi.org/10.1016/j.neuron.2019.01.030

Farias, J., Holt, C. E., Sotelo, J. R., & Sotelo-Silveira, J. R. (2020). Axon microdissection and transcriptome profiling reveals the in vivo RNA content of fully differentiated myelinated motor axons. *RNA (New York, N.Y.)*, *26*(5), 595–612. https://doi.org/10.1261/RNA.073700.119

González, C., Cánovas, J., Fresno, J., Couve, E., Court, F. A., & Couve, A. (2016). Axons provide the secretory machinery for trafficking of voltage-gated sodium channels in peripheral nerve. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(7), 1823–1828. https://doi.org/10.1073/PNAS.1514943113

Gumy, L. F., Yeo, G. S. H., Tung, Y. C. L., Zivraj, K. H., Willis, D., Coppola, G., Lam, B. Y. H., Twiss, J. L., Holt, C. E., & Fawcett, J. W. (2011). Transcriptome analysis of embryonic and adult sensory axons reveals changes in mRNA repertoire localization. *RNA (New York, N.Y.)*, *17*(1), 85–98. https://doi.org/10.1261/RNA.2386111

Hazan, J., Fonknechten, N., Mavel, D., Paternotte, C., Samson, D., Artiguenave, F., Davoine, C.-S., Cruaud, C., Dürr, A., Wincker, P., Brottier, P., Cattolico, L., Barbe, V., Burgunder, J.-M., Prud'homme, J.-F., Brice, A., Fontaine, B., Heilig, R., & Weissenbach, J. (1999). Spastin, a new AAA protein, is altered in the most frequent form of autosomal dominant spastic paraplegia. *Nature Genetics*, *23*(3), 296–303. https://doi.org/10.1038/15472

He, Z., & Jin, Y. (2016). Intrinsic Control of Axon Regeneration. *Neuron*, *90*(3), 437–451. https://doi.org/10.1016/J.NEURON.2016.04.022

Humphrey, W., Dalke, A., & Schulten, K. (1996). VMD: Visual molecular dynamics. *Journal of Molecular Graphics*, *14*(1), 33–38. https://doi.org/10.1016/0263-7855(96)00018-5

Iwahashi, J., Hamada, N., & Watanabe, H. (2007). Two hydrophobic segments of the RTN1 family determine the ER localization and retention. *Biochemical and Biophysical Research Communications*, *355*(2), 508–512. https://doi.org/10.1016/j.bbrc.2007.02.001

Jin, L. Q., Zhou, Y., Li, Y. S., Zhang, G., Hu, J., & Selzer, M. E. (2022). Transcriptomes of Injured Lamprey Axon Tips: Single-Cell RNA-Seq Suggests Differential Involvement of MAPK Signaling Pathways in Axon Retraction and Regeneration after Spinal Cord Injury. *Cells*, *11*(15), 2320. https://doi.org/10.3390/CELLS11152320/S1

Jung, H., Gkogkas, C. G., Sonenberg, N., & Holt, C. E. (2014). Remote control of gene function by local translation. In *Cell*. https://doi.org/10.1016/j.cell.2014.03.005

Kaethner, R. J., & Stuermer, C. A. 0. (1992). Dynamics of Terminal Arbor Formation and Target Approach of Retinotectal Axons in Living Zebrafish Embryos: A Time-Lapse Study of Single Axons. *The Journal of Neuroscience*, *72*(8), 32573271.

Kaya, L., Meissner, B., Riedl, M. C., Muik, M., Schwarzer, C., Ferraguti, F., Sarg, B., Lindner, H., Schweigreiter, R., Knaus, H. G., Romanin, C., & Bandtlow, C. E. (2013). Direct association of the reticulon protein RTN1A with the ryanodine receptor 2 in neurons. *Biochimica et Biophysica Acta*, *1833*(6), 1421. https://doi.org/10.1016/J.BBAMCR.2013.02.012

Kiss, A., Fischer, I., Kleele, T., Misgeld, T., & Propst, F. (2018). Neuronal growth cone size-dependent and -independent parameters of microtubule polymerization. *Frontiers in Cellular Neuroscience*, *12*, 195. https://doi.org/10.3389/FNCEL.2018.00195/FULL

Koppers, M., Özkan, N., Nguyen, H. H., Jurriens, D., McCaughey, J., Nguyen, D. T. M., Li, C. H., Stucchi, R., Altelaar, M., MacGillavry, H. D., Kapitein, L. C., Hoogenraad, C. C., & Farías, G. G. (2024). Axonal endoplasmic reticulum tubules control local translation via P180/RRBP1-mediated ribosome interactions. *Developmental Cell*, *0*(0). https://doi.org/10.1016/j.devcel.2024.05.005

Krijnse-Locker, J., Parton, R. G., Fuller, S. D., Griffiths, G., & Dotti, C. G. (1995). The organization of the endoplasmic reticulum and the intermediate compartment in cultured rat hippocampal neurons. *Molecular Biology of the Cell*, *6*(10), 1315–1332. https://doi.org/10.1091/mbc.6.10.1315

Kuo, Y. W., & Howard, J. (2021). Cutting, Amplifying, and Aligning Microtubules with Severing Enzymes. *Trends in Cell Biology*, *31*(1), 50. https://doi.org/10.1016/J.TCB.2020.10.004

Kuo, Y. W., Trottier, O., Mahamdeh, M., & Howard, J. (2019). Spastin is a dual-function enzyme that severs microtubules and promotes their regrowth to increase the number and mass of microtubules. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(12), 5533–5541. https://doi.org/10.1073/PNAS.1818824116/SUPPL_FILE/PNAS.1818824116.SM03.AVI

Lee, M., Paik, S. K., Lee, M. J., Kim, Y. J., Kim, S., Nahm, M., Oh, S. J., Kim, H. M., Yim, J., Lee, C. J., Bae, Y. C., & Lee, S. (2009). Drosophila Atlastin regulates the stability of muscle microtubules and is required for synapse development. *Developmental Biology*, *330*(2), 250–262. https://doi.org/10.1016/J.YDBIO.2009.03.019

Lin, H.-M., Kuang, J.-X., Sun, P., Li, N., Lv, X., & Zhang, Y.-H. (2018). Reconstruction of Intratelencephalic Neurons in the Mouse Secondary Motor Cortex Reveals the Diverse Projection Patterns of Single Neurons. *Frontiers in Neuroanatomy*, *12*. https://doi.org/10.3389/fnana.2018.00086

Lin, Y. F., Xie, Z., Zhou, J., Chen, H. H., Shao, W. W., & Lin, H. D. (2019). Effect of exogenous spastin combined with polyethylene glycol on sciatic nerve injury. *Neural Regeneration Research*, *14*(7), 1271–1279. https://doi.org/10.4103/1673-5374.251336

Liu, B., Liu, S., & Sun, D. (2023). Low-Dose Taxol Promotes Neuronal Axons Extension and Functional Recovery after Spinal Cord Injury. *Mediators of Inflammation*, *2023*, 5604103. https://doi.org/10.1155/2023/5604103

Liu, Q., Yang, H., Luo, J., Peng, C., Wang, K., Zhang, G., Lin, H., & Ji, Z. (2024). 14-3-3 protein augments the protein stability of phosphorylated spastin and promotes the recovery of spinal cord injury through its agonist intervention. *ELife*, *12*. https://doi.org/10.7554/eLife.90184

Liz, M. A., Mar, F. M., Santos, T. E., Pimentel, H. I., Marques, A. M., Morgado, M. M., Vieira, S., Sousa, V. F., Pemble, H., Wittmann, T., Sutherland, C., Woodgett, J. R., & Sousa, M. M. (2014). Neuronal deletion of GSK3β increases microtubule speed in the growth cone and enhances axon regeneration via CRMP-2 and independently of MAP1B and CLASP2. *BMC Biology*, *12*(1), 1–19. https://doi.org/10.1186/1741-7007-12-47/FIGURES/8

Lopez-Verrilli, M. A., Picou, F., & Court, F. A. (2013). Schwann cell-derived exosomes enhance axonal regeneration in the peripheral nervous system. *GLIA*. https://doi.org/10.1002/glia.22558

Luarte, A., Cornejo, V. H., Bertin, F., Gallardo, J., & Couve, A. (2018). The axonal endoplasmic reticulum: One organelle—many functions in development, maintenance, and plasticity. In *Developmental Neurobiology*. https://doi.org/10.1002/dneu.22560

Maciel, R., Bis, D. M., Rebelo, A. P., Saghira, C., Züchner, S., & Saporta, M. A. (2018). The human motor neuron axonal transcriptome is enriched for transcripts related to mitochondrial function and microtubule-based axonal transport. *Experimental Neurology*, *307*, 155. https://doi.org/10.1016/J.EXPNEUROL.2018.06.008

Mannan, A. U., Boehm, J., Sauter, S. M., Rauber, A., Byrne, P. C., Neesen, J., & Engel, W. (2006). Spastin, the most commonly mutated protein in hereditary spastic paraplegia interacts with Reticulon 1 an endoplasmic reticulum protein. *Neurogenetics*, *7*(2), 93–103. https://doi.org/10.1007/s10048-006-0034-4

Matamoros, A. J., Tom, V. J., Wu, D., Rao, Y., Sharp, D. J., & Baas, P. W. (2019). Knockdown of Fidgetin Improves Regeneration of Injured Axons by a Microtubule-Based Mechanism. *Journal of Neuroscience*, *39*(11), 2011–2024. https://doi.org/10.1523/JNEUROSCI.1888-18.2018

McCurdy, E. P., Chung, K. M., Benitez-Agosto, C. R., & Hengst, U. (2019). Promotion of Axon Growth by the Secreted End of a Transcription Factor. *Cell Reports*, *29*(2), 363. https://doi.org/10.1016/J.CELREP.2019.08.101

Nepravishta, R., Bellomaria, A., Polizio, F., Paci, M., & Melino, S. (2010). Reticulon RTN1-CCT peptide: A potential nuclease and inhibitor of histone deacetylase enzymes. *Biochemistry*, *49*(2), 252–258. https://doi.org/10.1021/BI9012676/SUPPL_FILE/BI9012676_SI_001.PDF

Nepravishta, R., Polizio, F., Paci, M., & Melino, S. (2012). A metal-binding site in the RTN1-C protein: new perspectives on the physiological role of a neuronal protein. *Metallomics*, *4*(5), 480–487. https://doi.org/10.1039/C2MT20035J

Nijssen, J., Aguila, J., Hoogstraaten, R., Kee, N., & Hedlund, E. (2018). Axon-Seq Decodes the Motor Axon Transcriptome and Its Modulation in Response to ALS. *Stem Cell Reports*, *11*(6), 1565. https://doi.org/10.1016/J.STEMCR.2018.11.005

Nozumi, M., Togano, T., Takahashi-Niki, K., Lu, J., Honda, A., Taoka, M., Shinkawa, T., Koga, H., Takeuchi, K., Isobe, T., & Igarashi, M. (2009). Identification of functional marker proteins in the mammalian growth cone. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(40), 17211–17216. https://doi.org/10.1073/pnas.0904092106

Pacheco, A., Merianda, T. T., Twiss, J. L., & Gallo, G. (2020). Mechanism and role of the intra-axonal Calreticulin translation in response to axonal injury. *Experimental Neurology*. https://doi.org/10.1016/j.expneurol.2019.113072

Pavez, M., Thompson, A. C., Arnott, H. J., Mitchell, C. B., D'Atri, I., Don, E. K., Chilton, J. K., Scott, E. K., Lin, J. Y., Young, K. M., Gasperini, R. J., & Foa, L. (2019). STIM1 is required for remodeling of the endoplasmic reticulum and microtubule cytoskeleton in steering growth cones. *Journal of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.2496-18.2019

Petrova, V., Pearson, C. S., Ching, J., Tribble, J. R., Solano, A. G., Yang, Y., Love, F. M., Watt, R. J., Osborne, A., Reid, E., Williams, P. A., Martin, K. R., Geller, H. M., Eva, R., & Fawcett, J. W. (2020). Protrudin functions from the endoplasmic reticulum to support axon regeneration in the adult CNS. *Nature Communications 2020 11:1*, *11*(1), 1–15. https://doi.org/10.1038/s41467-020-19436-y

Rao, K., Stone, M. C., Weiner, A. T., Gheres, K. W., Zhou, C., Deitcher, D. L., Levitan, E. S., & Rolls, M. M. (2016). Spastin, atlastin, and ER relocalization are involved in axon but not dendrite regeneration. *Molecular Biology of the Cell*. https://doi.org/10.1091/mbc.E16-05-0287

Reali, V., Mehdawy, B., Nardacci, R., Filomeni, G., Risuglia, A., Rossin, F., Antonioli, M., Marsella, C., Fimia, G. M., Piacentini, M., & Di Sano, F. (2015). Reticulon protein-1C is a key component of MAMs. *Biochimica et Biophysica Acta*, *1853*(3), 733–745. https://doi.org/10.1016/J.BBAMCR.2014.12.031

Roebroek, A. J. M., Van De Velde, H. J. K., Van Bokhoven, A., Broers, J. L. V, Ramaekers, F. C. S., & Van De Ven, W. J. M. (1993). Cloning and expression of alternative transcripts of a novel neuroendocrine-specific gene and identification of its 135-kDa translational product. *Journal of Biological Chemistry*, *268*(18), 13439–13447.

Roll-Mecak, A., & Vale, R. D. (2008). Structural basis of microtubule severing by the hereditary spastic paraplegia protein spastin. *Nature*. https://doi.org/10.1038/nature06482

Rolls, M. M., Thyagarajan, P., & Feng, C. (2020). Microtubule dynamics in healthy and injured neurons. *Developmental Neurobiology*, *81*(3), 321. https://doi.org/10.1002/DNEU.22746

Rotem, N., Magen, I., Ionescu, A., Gershoni-Emek, N., Altman, T., Costa, C. J., Gradus, T., Pasmanik-Chor, M., Willis, D. E., Ben-Dov, I. Z., Hornstein, E., & Perlson, E. (2017). ALS Along the Axons - Expression of Coding and Noncoding RNA Differs in Axons of ALS models. *Scientific Reports*, *7*. https://doi.org/10.1038/SREP44500

Saal, L., Briese, M., Kneitz, S., Glinka, M., & Sendtner, M. (2014). Subcellular transcriptome alterations in a cell culture model of spinal muscular atrophy point to widespread defects in axonal growth and presynaptic differentiation. *RNA (New York, N.Y.)*, *20*(11), 1789–1802. https://doi.org/10.1261/RNA.047373.114

Sahu, R., Kaushik, S., Clement, C. C., Cannizzo, E. S., Scharf, B., Follenzi, A., Potolicchio, I., Nieves, E., Cuervo, A. M., & Santambrogio, L. (2011). Microautophagy of Cytosolic Proteins by Late Endosomes. *Developmental Cell*, *20*(1), 131–139. https://doi.org/10.1016/J.DEVCEL.2010.12.003

Sasaki, Y., Welshhans, K., Wen, Z., Yao, J., Xu, M., Goshima, Y., Zheng, J. Q., & Bassell, G. J. (2010). Phosphorylation of Zipcode Binding Protein 1 Is Required for Brain-Derived Neurotrophic Factor Signaling of Local -Actin Synthesis and Growth Cone Turning. *Journal of Neuroscience*, *30*(28), 9349–9358. https://doi.org/10.1523/JNEUROSCI.0499-10.2010

Sengottuvel, V., Leibinger, M., Pfreimer, M., Andreadaki, A., & Fischer, D. (2011). Taxol Facilitates Axon Regeneration in the Mature CNS. *Journal of Neuroscience*, *31*(7), 2688–2699. https://doi.org/10.1523/JNEUROSCI.4885-10.2011

Shigeoka, T., Jung, H., Jung, J., Turner-Bridger, B., Ohk, J., Lin, J. Q., Amieux, P. S., & Holt, C. E. (2016). Dynamic Axonal Translation in Developing and Mature Visual Circuits. *Cell*, *166*(1), 181–192. https://doi.org/10.1016/j.cell.2016.05.029

Shigeoka, T., Koppers, M., Wong, H. H. W., Lin, J. Q., Cagnetta, R., Dwivedy, A., de Freitas Nascimento, J., van Tartwijk, F. W., Ströhl, F., Cioni, J. M., Schaeffer, J., Carrington, M., Kaminski, C. F., Jung, H., Harris, W. A., & Holt, C. E. (2019). On-Site Ribosome Remodeling by Locally Synthesized Ribosomal Proteins in Axons. *Cell Reports*, *29*(11), 3605-3619.e10. https://doi.org/10.1016/j.celrep.2019.11.025

Sholl, D. A. (1953). Dendritic organization in the neurons of the visual and motor cortices of the cat. *Journal of Anatomy*, *87*(Pt 4), 387. https://pmc.ncbi.nlm.nih.gov/articles/PMC1244622/

Solowska, J. M., Rao, A. N., & Baas, P. W. (2017). Truncating mutations of SPAST associated with hereditary spastic paraplegia indicate greater accumulation and toxicity of the M1 isoform of spastin. *Molecular Biology of the Cell*, *28*(13), 1728. https://doi.org/10.1091/MBC.E17-01-0047

Song, W., Cho, Y., Watt, D., & Cavalli, V. (2015). Tubulin-tyrosine ligase (TTL)-mediated increase in tyrosinated α-tubulin in injured axons is required for retrograde injury signaling and axon regeneration. *Journal of Biological Chemistry*, *290*(23), 14765–14775. https://doi.org/10.1074/jbc.M114.622753

Stepanova, T., Slemmer, J., Hoogenraad, C. C., Lansbergen, G., Dortland, B., De Zeeuw, C. I., Grosveld, F., Van Cappellen, G., Akhmanova, A., & Galjart, N. (2003). Visualization of Microtubule Growth in Cultured Neurons via the Use of EB3-GFP (End-Binding Protein 3-Green Fluorescent Protein). *Journal of Neuroscience*, *23*(7), 2655–2664. https://doi.org/10.1523/JNEUROSCI.23-07-02655.2003

Stepanova, T., Smal, I., Van Haren, J., Akinci, U., Liu, Z., Miedema, M., Limpens, R., Van Ham, M., Van Der Reijden, M., Poot, R., Grosveld, F., Mommaas, M., Meijering, E., & Galjart,

N. (2010). History-dependent catastrophes regulate axonal microtubule behavior. *Current Biology*. https://doi.org/10.1016/j.cub.2010.04.024

Stone, M. C., Rao, K., Gheres, K. W., Kim, S., Tao, J., La Rochelle, C., Folker, C. T., Sherwood, N. T., & Rolls, M. M. (2012). Normal Spastin Gene Dosage Is Specifically Required for Axon Regeneration. *Cell Reports*, *2*(5), 1340–1350. https://doi.org/10.1016/j.celrep.2012.09.032

Taylor, A. M., Berchtold, N. C., Perreau, V. M., Tu, C. H., Li Jeon, N., & Cotman, C. W. (2009). Axonal mRNA in Uninjured and Regenerating Cortical Mammalian Axons. *Journal of Neuroscience*, *29*(15), 4697–4707. https://doi.org/10.1523/JNEUROSCI.6130-08.2009

Taylor, A. M., Blurton-Jones, M., Rhee, S. W., Cribbs, D. H., Cotman, C. W., & Jeon, N. L. (2005). A microfluidic culture platform for CNS axonal injury, regeneration and transport. *Nature Methods*. https://doi.org/10.1038/nmeth777

Teixeira Lopes, A. (2018). *Spastin regulates microtubule dynamics and is required for normal motor and cognitive functions*.

Twiss, J. L., Kalinski, A. L., Sachdeva, R., & Houle, J. D. (2016). Intra-axonal protein synthesis – a new target for neural repair? *Neural Regeneration Research*, *11*(9), 1365. https://doi.org/10.4103/1673-5374.191193

Van De Velde, H. J. K., Roebroek, A. J. M., Senden, N. H. M., Ramaekers, F. C. S., & Van De Ven, W. J. M. (1994). NSP-encoded reticulons, neuroendocrine proteins of a novel gene family associated with membranes of the endoplasmic reticulum. *Journal of Cell Science*, *107*(9), 2403–2416. https://doi.org/10.1242/JCS.107.9.2403

Van Zundert, G. C. P., Rodrigues, J. P. G. L. M., Trellet, M., Schmitz, C., Kastritis, P. L., Karaca, E., Melquiond, A. S. J., Van Dijk, M., De Vries, S. J., & Bonvin, A. M. J. J. (2016). The HADDOCK2.2 Web Server: User-Friendly Integrative Modeling of Biomolecular Complexes. *Journal of Molecular Biology*, *428*(4), 720–725. https://doi.org/10.1016/J.JMB.2015.09.014

Voeltz, G. K., Prinz, W. A., Shibata, Y., Rist, J. M., & Rapoport, T. A. (2006). A class of membrane proteins shaping the tubular endoplasmic reticulum. *Cell*, *124*(3), 573–586. https://doi.org/10.1016/j.cell.2005.11.047

von Kügelgen, N., & Chekulaeva, M. (2020). Conservation of a core neurite transcriptome across neuronal types and species. *Wiley Interdisciplinary Reviews: RNA*, *11*(4), e1590. https://doi.org/10.1002/WRNA.1590;CTYPE:STRING:JOURNAL

Willis, D., Ka, W. L., Zheng, J. Q., Chang, J. H., Smit, A., Kelly, T., Merianda, T. T., Sylvester, J., Van Minnen, J., & Twiss, J. L. (2005). Differential transport and local translation of cytoskeletal, injury-response, and neurodegeneration protein mRNAs in axons. *Journal of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.4235-04.2005

Wu, Y., Whiteus, C., Xu, C. S., Hayworth, K. J., Weinberg, R. J., Hess, H. F., & De Camilli, P. (2017). Contacts between the endoplasmic reticulum and other membranes in neurons. *Proceedings of the National Academy of Sciences of the United States of America*. https://doi.org/10.1073/pnas.1701078114

Yap, Y. C., King, A. E., Guijt, R. M., Jiang, T., Blizzard, C., Breadmore, M. C., & Dickson, T. C. (2017). Mild and repetitive very mild axonal stretch injury triggers cystoskeletal mislocalization and growth cone collapse. *PLOS ONE*, *12*(5), e0176997. https://doi.org/10.1371/JOURNAL.PONE.0176997

Zhang, Y., Chopp, M., Liu, X. S., Kassis, H., Wang, X., Li, C., An, G., & Zhang, Z. G. (2015). MicroRNAs in the axon locally mediate the effects of chondroitin sulfate proteoglycans and cGMP on axonal growth. *Developmental Neurobiology*, *75*(12), 1402–1419. https://doi.org/10.1002/dneu.22292

Zhang, Y., Ueno, Y., Liu, X. S., Buller, B., Wang, X., Chopp, M., & Zhang, Z. G. (2013). The MicroRNA-17-92 cluster enhances axonal outgrowth in embryonic cortical neurons. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *33*(16), 6885–6894. https://doi.org/10.1523/JNEUROSCI.5180-12.2013

Acknowledgments: This research was partially supported by HPC OCÉANO (FONDEQUIP N° EQM170214) and the supercomputing infrastructure of the NLHPC (CCSS210001). Centro Nacional de Inteligencia Artificial CENIA, FB210017, BASAL, ANID.

Funding:
ANID projects Fondecyt Initiation 11220601 (A.L.); EMBO scientific exchange grant 9218; EMBO short-stay Fellowship 8263.

Author contributions.
Examples: AL, JPR
Conceptualization: AL, AC, RHM.
Methodology: AL, TA, APL, CG, CC, FB, JG, WA, AW, ACH, DC, JT,RV.
Investigation: AL, WZ, RHM, UH, CGR, ACH, AF, JPR.
Visualization: AL, CGR, AC, JT.
Supervision: AL, AC, UH, WZ, CG, APL.
Writing—original draft: AL, AC.
Writing—review & editing: AL, AC, CG, RHM, UW, CGR.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data needed to evaluate the conclusions in the paper will be available upon request.

Tables and Figures

1294

Study	Neuron cell type	Model to obtain axonal fraction
(Saal et al., 2014)	Mice motor neurons	Spinal cord motor neurons cultured in microfluidic chambers (7 DIV).
(Rotem et al., 2017)	Mice motor neurons	Spinal cord motor neurons cultured in porous membrane inserts (10 DIV).
(Maciel et al., 2018)	iPSC-derived human motor neurons	Motor neurons were cultured on porous membrane inserts (10 DIV)
(Bigler et al., 2017)	Glutamatergic neurons differentiated from human embryonic stem cells (hESCs)	hESCs differentiated for 59 DIV in microfluidic chambers.
(Nijssen et al., 2018)	Human and mice motor neurons.	Spinal cord and human stem cell derived motor neurons were cultured in microfluidic chambers (7 and 14 DIV, respectively).
(Farias et al., 2020)	Rat mature myelinated motor neurons	Axoplasm microdissection of motor axons (*Ex vivo*).
(Gumy et al., 2011)	Rat embryonic/adult sensory neurons from dorsal root ganglion (DRG)	Primary neurons were cultured in microfluidic chambers (4 DIV).
(Taylor et al., 2009)	Uninjured and regenerating rat cortical neurons	Primary neurons cultured in microfluidic chambers (10 DIV).

1295 **Table I.** Analyzed publications examining the transcriptome of an axon-enriched neuronal

1296 fraction. Days *in vitro* (DIV).

1297

1298

Figure 1. Analysis of diverse axonal transcriptomes reveals that *Rtn-1* mRNA is consistently

identified. **A.** Scheme illustrates studies of axonal transcriptomes and 104 shared mRNAs

identified through a Benjamini–Hochberg (Ben) analysis. **B.** Table lists the category "Cellular

Component" (Gene Ontology, GO) to classify the proteins coded by the 104 common mRNAs.

The endoplasmic reticulum (ER) category only harbors *Rtn-1* (bold) as the only common mRNA

coding for an ER structural protein.

1305

1306

Figure 2. Axonal synthesis of Rtn-1 can be locally regulated in cortical neurons. A. *Left:*

Western blot bands of Rtn-1 and β3-tubulin (β3-tub) confirms the knockdown (KD) efficacy of

siRNAs specifically designed against *Rtn-1* mRNA (Rtn-1 siRNA). 7 DIV Cortical neurons were

transfected with Rtn-1 siRNAs and a control sequence (scrambled, Scr) for forty-eight hours. 20

µg of protein were loaded per lane, and β3-tubulin was used as loading control. *Right:* Graphs

quantify the fold change of the 24 kDa Rtn-1 band after treatments with siRNAs 1-2 compared to

Scr. Data are displayed as mean ± standard error of N = 6 biological replicates. Wilcoxon test

compared to a theoretical mean of 1; ∗p < 0.05; ∗∗p < 0.01. **B.** Immunofluorescence images of

the soma and distal axon of neurons stained against β3-tub and co-transfected with a plasmid

encoding RFP and siRNA 2 or Scr under the same conditions indicated in A. Representative

images from N = 4 biological replicates. Bar soma and axons = 20 µm **C.** *Left:* schematic

representation of the axonal KD strategy based on controlling volume differences between the

compartment of microfluidic chambers. Under these conditions, 7 DIV cortical neurons were

selectively submitted to axonal treatments for twenty-four hours. *Right:* representative images and quantitative analysis of Rtn-1 and β3-tub intensity fold change in the cell bodies after axonal treatments. Bar = 10 μm. Data are displayed as mean ± standard deviation (SD) of 229-300 neurons from N = 3 biological replicates. Linear mixed-effects model with experimental date included as a random effect; ∗∗∗p < 0.001. **D.** Immunofluorescence showing the intensity changes of Rtn-1 and β3-tubulin along the last 50 μm of axons locally treated with Rtn-1 siRNA or Scr. Bar =10 μm **E.** *Left:* corrected fluorescence intensity profiles of Rtn-1 and β3-tubulin from the last 50 μm of distal axons locally treated with Rtn-1 siRNA (red line) or Scr (black line). Intensity profiles depict the mean ± standard error from all the analyzed axons. *Right:* Bar graphs display the mean ± standard deviation of the fluorescence intensity fold change of 87-102 axons from N = 3 biological replicates. Linear mixed-effects model with experimental date included as a random effect; ∗p < 0.05; ∗∗∗p < 0.001.

Figure 3. Axonal Rtn-1 knockdown (KD) promotes outgrowth and modify growth cone (GC) cytoskeleton after injury. A. Immunofluorescence against β3-tubulin in axons growing in microfluidic chambers twenty-four hours after injury (vacuum aspiration axotomy) comparing axonal and somatic Rtn-1 KD performed with a specific siRNA (Rtn-1-siRNA) or control sequence (scrambled, Scr). Bar = 50 µm. **B.** *Top:* schematic of the Sholl analysis performed with the NeuroAnatomy plugin (FIJI). The number of intersections of binarized axons (*left*) with concentric semi-circles of increasing radius starting in the microgroove at the center of each axonal field were quantified (*right*). *Bottom:* mean plots of the Sholl analyses performed after axonal Rtn-1 KD (red line) compared to Scr (black line). Bar graphs show the summation (sum.) of the total intersections within the indicated distance bins. **C.** Same analysis as in panel B but performed for somatic Rtn-1 KD. Data are displayed as mean ± standard error of 23-24 axonal fields from N = 6 biological replicates for axonal Rtn-1 KD; and 18-20 axonal fields from N = 3

biological replicates for somatic Rtn-1 KD. Two-tailed Mann–Whitney *U* test; **p < 0.01; ****p < 0.0001.**C.** Images show axonal growth cones after axonal or somatic Rtn-1 KD compared to Scr labeled against β3-tubulin (red) and Phalloidin (green) for staining the F-actin outline. Bar = 5 μm. Graphs depict the corresponding differences in GC area, perimeter, elongation ratio (or aspect ratio), number of filopodia, and longest shortest path (LSP), which corresponds to maximal distance calculated from the skeletonized outline (shown with the red line of the schematic drawings below). Data are displayed as mean ± standard error of 145-153 axons for axonal KD; 227-148 axons for somatic KD from N = 3 biological replicates. Two-tailed Mann–Whitney *U* test; ∗∗p < 0.01; ∗∗∗∗p < 0.0001.

Figure 4. Axonal and somatic β3-tubulin levels are dependent on the local synthesis of Rtn-1 after injury. A. *Left:* immunofluorescence images and their corresponding corrected fluorescence intensity profiles along the last 50 μm of distal axons stained against Rtn-1 and β3-tubulin after axonal transfection with Rtn-1 siRNA (red; axonal Rtn-1 KD) or a control sequence (black; scrambled, Scr). Intensity profiles depict the mean ± standard deviation from all the analyzed axons. Bar = 10 μm. *Right:* bar graphs quantify the Rtn-1 and β3-tubulin (β3-tub) fluorescence intensity fold change. Data are displayed as the mean ± standard deviation of 206–

242 axons from N = 3 biological replicates. Linear mixed-effects model with experimental date included as a random effect; ***p < 0.001.**B.** Same analysis as in panel A but performed for somatic Rtn-1 KD. Data are displayed as mean ± standard error of 133–161 axons from N = 3 biological replicates. Linear mixed-effects model with experimental date included as a random effect; ***p < 0.001. **C.** Images depict neuronal cell bodies stained for Rtn-1 and β3-tub after axonal Rtn-1 KD or Scr. Bar graphs depict Rtn-1 and β3-tub fluorescence intensity fold change displayed as mean ± standard error of 358–382 neurons from N = 3 biological replicates. Linear mixed-effects model with experimental date included as a random effect; ***p < 0.001. Bar = 10 µm. **D.** Same analysis as in panel C but performed for somatic Rtn-1 KD. Data are displayed as mean ± standard error of 252–237 neurons from N = 3 biological replicates. Linear mixed-effects model with experimental date included as a random effect; ***p<0.001. Bar = 10 µm.

1375

Figure 5. Axonal Rtn-1 KD after injury controls distal tubulin levels and outgrowth depending on Spastin microtubule severing. **A.** *Left:* representative fluorescent super resolution STED images of distal non-injured and injured cortical axons (forty-four hours post-axotomy) in microfluidic chambers, which were stained against Rtn-1C (green), and Spastin (red), axon outlines are depicted (white lines). *Right:* Plots depict the relative fluorescence intensity profile (normalized against total distal intensity) of Rtn-1C and Spastin in non-injured (black) and injured (red) axons. Data are displayed as mean ± standard error of 9-13 axons from N = 4 biological

replicates. Bar = 5 µm **B.** *Top:* representative confocal images of distal axons stained against β3-

tubulin (β3-tub, yellow), Rtn-1C (green), and Spastin (red) in microfluidic chambers forty-four

hours post-axotomy under indicated treatments: control (Scrambled, Scr), axonal Rtn-1 KD (Rtn-

1 siRNA), and Spastin pharmacological inhibition with 50 nM Spastazoline (SPZT) or vehicle

(DMSO). Bar = 10 µm. *Bottom:* quantifications of fluorescence intensity fold change of indicated

proteins in the last 50 µm normalized to Scr + DMSO. Data are displayed as mean ± standard

error of 24-40 axons for condition from N = 3 biological replicates. Kruskal–Wallis followed by

Dunn's post hoc test; ∗∗∗$p < 0.001$.**C.** Illustrative axonal fields in microfluidic chambers forty-

four hours post-axotomy stained against β3-tub under the treatments shown in B. Bar = 200 µm.

D. *Left:* mean plots of the Sholl analyses performed with colored lines indicating the analyzed

conditions. *Right:* bar graphs show the summation (Sum.) of the total intersections within the

indicated distance bins. Data are displayed as mean ± standard error of 22-24 axonal fields for

condition from N = 3 biological replicates. One-way ANOVA followed by Tukey's post hoc test;

∗$p < 0.05$. **E.** *Left:* axonal kymographs showing microtubule dynamics in neurons with indicated

treatments forty-four hours after axotomy, dashes represent the traces of EB3-GFP (end-binding

protein 3 fused to Emerald GFP). Scale bars represent 10 µm and 30 seconds, respectively. *Right:*

bar graphs show microtubule dynamic parameters in non-injured and injured axons with indicated

treatments. Data are displayed as mean ± standard error of 146-377 comets/dashes for condition

from N = 3 biological replicates. Kruskal–Wallis followed by Dunn's post hoc test; ∗$p < 0.05$;

∗∗$p < 0.01$; ∗∗∗∗$p < 0.001$.

1403

Figure 6. **Rtn-1C is locally synthesized in cortical axons**. **A.** *Left:* immunofluorescence of

neuronballs stained against neurofilament medium chain (NF-M, grayscale and green) and

Microtubule-associated protein 2 (MAP2, grayscale and red) with nuclei counterstaining (DAPI,

blue). Top panel shows intact neuronballs, bottom panel shows them after mechanic removal of

the cell bodies. *Middle:* RT- PCR bands of somatic (soma), axonal (Axon), and (-) (reaction of

axonal RNA without reverse transcriptase) to reveal mRNAs coding for MAP2, glial fibrillary

acidic protein (GFAP), Histone H1, β-actin, Rtn-1C, and Rtn-1A. *Right:* Western blot analysis of

the corresponding fractions using a pan-Rtn-1 antibody to reveal Rtn-1A and Rtn-1C isoforms,

β3-tubulin (β3-tub), and Tau. Bar = 500 μm. Images and bands are representative of two

biological replicates. **B.** Confocal images showing the expression of Rtn-1C and Rtn-1A (green)

and β3-tub (red) in the distal axonal compartment of microfluidic chambers. Bar = 10 μm. Images

are representative of two biological replicates **C.** Schematic of the microfluidic chamber setup

used to perform proximity ligation assay coupled to puromycylation (puro-PLA) is depicted

below. Local puromycin (puro) incorporation into newly synthesized peptides allows

amplification when puro is at nanometric distance from the N-terminal. **D.** puro-PLA positive

spots are shown in green at somatic and axonal compartments. β3-tub labeling depicts cell bodies

and axons. Insets show higher magnification of puro-PLA signals. Bar graphs quantify dots per

cell bodies and per axonal area after local co-treatments of puro with the protein synthesis

inhibitor emetine (Eme) or vehicle (Veh), or puro at the opposite compartment. Bar = 5 μm;

inserts= 1 μm. Data are displayed as mean ± standard deviation of 11-12 somatic and 16-33 axonal

fields from N = 4 biological replicates. Kruskal–Wallis followed by Dunn´s post hoc test; $*p <$

0.05; $**p < 0.01$; $***p < 0.001$; $**** p < 0.0001$.

1426

1427

1428

1429

Figure 7. Rtn-1C co-distributes with Spastin and attenuates its microtubule-severing

activity. A. Representative images of COS-7 cells expressing Spastin-RFP (fire red) together with

GFP (green), Rtn-1A-GFP, or Rtn-1C-GFP, and immunostained against tubulin (yellow). Bar =10

μm. **B.** Bar graphs show the indicated Mander's coefficients to quantify the colocalization

between Spastin-RFP and GFP or Rtn-1 isoforms. Data are displayed as mean ± standard error

from 9 cells per condition from N = 2 biological replicates. One-way ANOVA with Tukey's post

hoc test; *$p < 0.05$, **$p < 0.01$, ****$p < 0.0001$. **C.** Tubulin intensity fraction normalized to non-

transfected cells (NT) in cells expressing Spastin-RFP with GFP, Rtn-1A-GFP, or Rtn-1C-GFP

compared to non-transfected cells. Data are displayed as mean ± standard error from 145 non-transfected cells, 33 Spastin-RFP/GFP cells, 35 Spastin-RFP/Rtn-1A-GFP cells, and 32 Spastin-RFP/Rtn-1C-GFP cells from N = 2 biological replicates. One-way ANOVA with Tukey's post hoc test; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. **D.** Domain organization of Spastin isoforms (M1 and M87) and Rtn-1 isoforms (Rtn-1A and Rtn-1C), including the Reticulon Homology Domain (RHD), Microtubule Interacting and Trafficking domain (MIT), Microtubule-Binding Domain (MTBD), and ATPases Associated with diverse cellular Activities (AAA) domain (severing domain). **E.** Predicted binding affinities (ΔG binding, kcal/mol) between Spastin M87 and Rtn-1 isoforms at different domains, highlighting strong interactions with Rtn-1C. **F.** Structural model (VMD snapshots) of the predicted interaction between the MTBD (yellow) of Spastin M87 (blue) and the N-terminal region (green) of Rtn-1C (red).

Supplementary figure 1. Rtn-1 is progressively enriched in the growth cone (GC) of developing cortical neurons. A. *Top:* immunofluorescence images showing the distribution of Rtn-1 and microtubule-associated protein 2 (MAP2) in cultured cortical neurons at 1, 3, and 7 days *in vitro* (DIV). Bar = 10 μm. *Bottom:* co-labeling of Rtn-1 or KDEL (green) with β3-tubulin (β3-tub, red) depicts the endoplasmic reticulum in distal axons. Bar = 10 μm **B.** Intensity profiles indicate the relative intensity (Rel. intensity, normalized against the total fluorescence) of Rtn-1 and KDEL in distal axons. Data are displayed as mean ± standard error of 80–100 axons from N = 3 biological replicates. **C.** Bar graphs quantify the percentage of total Rtn-1 and KDEL

fluorescence present in the first 10 μm of axons (GC) at 1, 3 and 7 DIV. Data are displayed as mean ± standard error of 80–100 axons from N = 3 biological replicates. ∗p < 0.05; n.s., not significant; One-way ANOVA followed by Tukey's post hoc test. **D.** Representative image of Rtn-1 distribution (green) at the GC identified with the F-actin outline (grays) stained with phalloidin. Bar = 5 μm.

Supplementary figure 2. Axonal knockdown (KD) of Rtn-1 with a different siRNA reproduced increased outgrowth. A. Representative Western blot of Rtn-1C and β3-tubulin (β3-tub) in three independent 7 day *in vitro* (DIV) cortical neuron cultures (N1–N3) transfected with siRNA 3 directed against Rtn-1 (Rtn-1 siRNA 3) and a control sequence (scrambled, Scr) for forty-eight hours. 9 DIV, non-transfected (NT) cells were also analyzed. Total protein staining was used as a loading control. **B.** Bar graphs quantify the fold change in β3-tub (left) and Rtn-1C (right) levels relative to NT. Data are displayed as the mean ± standard error from N = 3 biological replicates. Wilcoxon test compared to a theoretical mean of 1; ∗p < 0.05 **C.** Representative images of axonal fields stained against β3-tub in microfluidic chambers after axonal KD with Rtn-1 siRNA 3 or Scr. Bar = 50 μm.**D.** *Left:* mean plots of the Sholl analyses performed after axonal Rtn-1 KD (red) compared to Scr (black). *Right:* Bar graphs indicate the summation (sum.) of intersections within the defined distance bins. Data are displayed as the mean ± standard error of

23-30 axonal fields from N =3 biological replicates. Unpaired Student's t Test; *p < 0.05; **p < 0.01.

Supplementary figure 3. Validation of Rtn-1C antibody specificity and puro-PLA signal by axonal knockdown (KD) of Rtn-1. A. *Top:* Schematic of the epitope (red bar) within the N-terminal region of Rtn-1C (top). *Bottom:* Western blot analysis of COS-7 cells non-transfected (lane 1) or transfected with Rtn-1C-GFP (lane 2). **B.** Immunofluorescence validation of α-Rtn-1C and α-Rtn-1A antibodies in COS-7 cells transfected with Rtn-1C-GFP (left) or Rtn-1A-GFP (right). Bar = 20 μm.**C.** *Left:* Western blot of 7 days *in vitro* (DIV) cortical neurons transfected for forty-four hours with siRNA targeting Rtn-1 (siRNA 2) or control sequence (scrambled, Scr). *Right:* Bar graphs quantify the intensity fold change of Rtn-1C versus Scr (normalized to β3-tubulin). Data are displayed as mean ± standard error from N = 3 biological replicates. *p < 0.05; Wilcoxon test compared to a theoretical mean of 1.**D.** *Left:* RNA was isolated from 8 DIV microfluidic chamber compartments and submitted to RT- PCR. Bands depict somatic (S), axonal (A) and negative control corresponding to axonal samples without reverse transcriptase (-). *Top right:* 8 DIV cortical axons were locally treated with siRNA 2 and Scr for twenty-four hours and then submitted to puromycylation proximity ligation assay (puro-PLA) to reveal local Rtn-1C

synthesis (green puncta) along β3-tubulin–positive axons (red). Bar = 5 μm Bar insert = 2.5 μm. *Bottom right:* Bar graphs quantify the number of positive dots per axonal area. Data are displayed as the mean ± standard deviation of 15-16 axonal fields from N = 3 biological replicates. Mann–Whitney test; ***p < 0.001.

Supplementary figure 4. Distal axonal levels of Spastin and Rtn-1C are not altered by axotomy. Quantification of the fluorescence intensity for Spastin (left) and Rtn-1C (right) in distal (last 15 μm) non-injured (black) and injured (red) axons in microfluidic chambers as described in Figure 5. Data are displayed as mean ± standard error of 9-13 axons from N = 3 biological replicates. No significant differences were detected between conditions.

Supplementary figure 5. Single-channel acquisition confirms punctate distribution of Spastin-RFP and GFP in co-transfected COS-7 cells. *Top:* COS-7 cells were co-transfected with Spastin-RFP (left, 555 channel) and GFP (right, 488 channel). Spastin-RFP displays a

punctate distribution that strongly co-distributes with GFP puncta. Independent acquisition of each channel demonstrates that this overlap is not due to fluorescence crosstalk, since some puncta appear exclusively in one channel (arrowheads). *Bottom:* Cells transfected with Rtn-1A-RFP and GFP show distinct distribution patterns supporting absence of crosstalk. Bar = 5 μm.

Supplementary figure 6. **Expression levels of Spastin and Rtn-1 constructs are comparable across analyzed cells.** Quantification of fluorescence intensity for RFP (left) and GFP (right) signals in cells co-expressing Spastin-RFP with GFP, Rtn-1C-GFP, or Rtn-1A-GFP, corresponding to the experimental conditions described in Figure 7. Data are displayed as mean ± standard error of 33 Spastin-RFP/GFP cells; 35 Spastin-RFP/Rtn-1A-GFP cells; and 32 Spastin-RFP/Rtn-1C-GFP cells from N = 2 biological replicates. Kruskal–Wallis followed by Dunn's post hoc test. No significant differences were found across conditions.