Excess calcium to transition sheep and goats

Effects of excess dietary calcium supplementation during the transition phase on

calcium and bone metabolism of dairy sheep and goats and growth response of their

suckling offspring

Daniel Brugger[1], Annette Liesegang[1]

Institute of Animal Nutrition and Dietetics, Vetsuisse-Faculty, University of Zurich

Annette Liesegang, Institut für Tierernährung und Diätetik, Winterthurerstrasse 270, 8057

Zurich, Switzerland, +41 44 635 88 23, aliese@nutrivet.uzh.ch

Abstract

This study investigated the effects of excess dietary Ca supplementation during the transition

phase on Ca and bone metabolism in dairy sheep and goats, as well as the growth response of

their suckling offspring. Multiparous dairy sheep (n=6/group) and goats (n=5/group) were

randomly assigned to a control and treatment group. The experimental group received a high

Ca diet (1.3% DM) based on hay and concentrate three weeks *ante partum*, compared to the

control group (0.6% DM). Experimental feeding ended with parturition and animals were

henceforth fed according to recommendations for lactating sheep and goats and kept together

with their suckling offspring. The observation period spanned from 21 d *ante partum* to 56 d

post partum. Animals were subject to continuous veterinary surveillance and monitored for

signs of milk fever. Data collection comprised quantitative and functional parameters of Ca

and bone metabolism as well as birth weights and daily weight gain of suckling lambs and

kids. Data was analyzed by repeated measures and endpoint mixed models. The results of

quantitative markers indicated that excess dietary Ca *ante partum* significantly increased fecal

Ca concentrations until parturition by 2 $\log_{10}$-orders of magnitude, suggesting efficient

physiological mechanisms to manage Ca overload by increasing fecal excretion. No

significant differences were observed in serum Ca levels, urinary Ca excretion, or bone

mineral density (BMD) between control and experimental groups at any point in time of

observation, indicating stable Ca homeostasis despite the dietary excess. However, species-

specific differences were noted, with sheep generally showing significantly higher serum and

colostral Ca levels, as well as BMD, compared to goats. Serum calcitriol as well as markers of

bone formation and resorption were monitored, revealing significant increases in serum

osteocalcin *post partum* in goats and sheep fed high Ca *ante partum.* All other serum markers

including calcitriol were not affected by the feeding regime but species of animal. Vitamin D

receptor abundance in goats' colonic mucosa were numerically lower during parturition when

high Ca was fed. All repeated measures were significantly affected by time except for urinary

Ca and bone specific alkaline phosphatase activity in serum. Offspring of sheep and goats in

the high Ca group exhibited significantly reduced average daily weight gain compared to

control, despite similar species-specific birth weights. We conclude that while dairy sheep and

goats can effectively manage high dietary Ca intake without overt signs of hypocalcemia or

milk fever, there are notable impacts on offspring growth and potential long-term

physiological effects that require further research. These findings contribute to the

understanding of mineral nutrition in periparturient dairy ruminants and highlight the need for

balanced dietary strategies to optimize health and productivity in both dams and their

offspring.

Key words: Mineral, Homeostasis, Hypocalcemia, Milk Fever

Nonstandard abbreviations: bAP = bone-specific alkaline phosphatase activity; BMD = bone

mineral density; calcitriol = 1,25-$(OH)_2$ vitamin D; CTX = crosslaps; CpG = cytosine linked

to guanine in 5'$\rightarrow$3' direction of a DNA molecule; ICTP = crosslinked carboxyterminal

teleopeptide of type I collagen; qCT = quantitative computer tomography; SAS = statistical

analysis software; VDR = vitamin D receptor

Introduction

Calcium is essential for structure and function of the animal organism and already in 1905 an experimental model of dietary Ca deficiency was successfully applied in dairy cows (Babcock, 1905).

To avoid a drop in circulatory Ca, the organism has three options to stabilize the system, by mobilizing Ca from the skeleton, increasing absorption capacity at the gut barrier, and limiting endogenous losses via the kidneys and gastrointestinal tract (Wilkens and Muscher-Banse, 2020). With respect to the regulation of Ca metabolism, the animal species under study is an important factor to consider. For example, sheep and goats have been demonstrated to respond quite differently to Ca-restricted feeding with the first tending quicker to bone mobilization while the latter will first attempt to compensate via more efficient absorption from the small intestinal mucosa (Wilkens et al., 2012). A relevant regulation of renal Ca excretion and reabsorption in response to changes in the endogenous Ca status has been reported for mice (Hoenderop et al., 2002, van Cromphaut et al., 2003, Ko et al., 2009), but was not confirmed in ruminants including sheep, goats (Herm et al., 2015) and cows (Wild et al., 2021). Quantitative adaption to fluctuations in the endogenous Ca status are regulated by a complex system of endocrine factors, that control the regulation of absorption, excretion and bone remodeling, the detailed discussion of which exceeds the capacities of this introduction (see the review by Martín-Tereso and Martens (2014) and Wilkens and Muscher-Banse (2020)).

In lactating animals, proper Ca supply is especially important since milk supplies high levels of Ca to the offspring on a daily basis (Hernández-Castellano et al., 2020). In this context, the transition phase during the last weeks *prepartum* is critical. In high-yielding dairy cows, the sudden onset of lactation after birth in combination with impaired feed intake around parturition puts animals in danger of developing milk fever if the mobilization of bone stores

lacks capacity to bridge this gap in dietary intake. This is typically accompanied by

hypocalcaemia, mostly also by hypophosphatemia and often hypomagnesemia (Liesegang et

al., 2007, Goff, 2008, Martín-Tereso and Martens, 2014). Symptoms develop within 48 h

postpartum and may include listlessness, coma and sternal recumbence. The situation for

ewes and nannies is different. Due to their lower milk yield, meat sheep breeds develop milk

fever rarely *post partum* though a higher tendency for carrying twin or even triplet fetuses

may increase the risk of developing a hypocalcaemia during the last trimester given an

increasing fetal Ca requirement during the last 4 weeks *ante partum* due to the calcification of

the fetal skeleton. Sheep typically show preceding anorexia and stiff gait before sternal

recumbence. Dairy sheep breeds and most goat breeds combine a high milk yield potential

with a high probability for carrying multiple fetuses, consequently clinical events of

hypocalcemia *pre-* and *postpartum* are likely. High-yielding dairy goats and sheep develop a

milk fever comparable to dairy cows but much later at 1-3 weeks postpartum. If prepartum

symptoms arise they are less pronounced than the postpartum complex (Oetzel, 1988).

Dietary and metabolic factors that affect the incidence of milk fever are various and the

underlying literature is dominated by studies in cows. Amongst relevant factors are the

DCAD, Mg, P, and Ca in the *ante partum* diet (Lean et al., 2006). There has been some debate

regarding the proper dietary Ca concentration *ante partum*, with some sources suggesting that

diet Ca at 1-1.5% DM or even 2-3% does not affect the incidence of milk fever whereas

others recommend going below 1%. An earlier meta-analysis on the matter attempted to bring

some order in the chaos. Authors developed a model with data from 51 trials that estimates the

incidence of milk fever as a function of breed of cow as well as *prepartum* dietary DCAD, Ca,

P, Mg, and diet exposure time. They concluded that the response of milk fever incidence to

dietary Ca *ante partum* follows a bell curve, which suggests that concentrations between 1.1-

1.5% DM should be avoided, whereas higher or lower levels gradually reduce the incidence

of milk fever (Lean et al., 2006).

Studies applying high dietary Ca levels in the feeding of periparturient animals are rare,

especially with sheep and goats. Therefore, the present study was conducted, which aimed to

collect such data and attempted to test the conclusions of Lean et al. (2006) under controlled

conditions. Therefore, periparturient sheep and goats of our in-house herd of dairy sheep and

dairy goats were either given diets with 0.6% DM or 1.3% DM of Ca during the last three

weeks *ante partum* and observed until 56 d of lactation. Our working hypothesis was that

goats and sheep supplied higher diet Ca (equivalent to 30-36 g/d) will show signs of

hypocalcemia whereas the lower supplied animals do not. It further allowed us to observe the

response of mineral and bone metabolism to dietary Ca overload in two species with distinct

strategies to maintain Ca homeostasis.

Material and Methods

Animals and Housing

This study was conducted at the research and laboratory facilities of the Institute of Animal Nutrition and Dietetics, Vetsuisse-Faculty, University of Zurich, Switzerland. Ten and twelve multiparous gestating dairy sheep and goats (East Friesian Dairy Sheep, Saanen goats), respectively, aged between 4 and 6 years were housed in individual pens with concrete floor and wood shavings as bedding material, from four weeks before the estimated parturition date. Both species were housed in rooms separate from each other. The goats and sheep were housed individually until their delivery date, afterwards with their respective lambs and kids in lambing/kidding pens. The average body weight just before birth of goats and sheep was 95.2 ± 1.5 kg and 101 ± 2.22 kg, respectively, and 78.3 ± 2.22 kg and 81.0 ± 2.31 kg directly after birth, respectively.

Diets and Experimental design

Animals were randomly divided in two groups according to their ear tag numbers by a random sequence generator (https://www.random.org/sequences/): one control and one experimental group per species. The effective sample size for sheep and goats was five and six per group, respectively. The feeding trial started three weeks before the estimated day of parturition. The diets were adapted to the specific energy and nutrient requirements during gestation and lactation for sheep and goats (Agroscope, 2017). The control groups received a balanced diet provided by hay and a commercial high energy concentrate (UFA 766, Herzogenbuchsee, Switzerland). Additionally, for the experimental groups until parturition, the same concentrate was applied but mixed up daily with 2.5 times (16.8 g Ca /day) the recommended daily calcium requirement as $CaCO_3$ (Table 1). From parturition onwards, all animals received lactation feeding according to good practice. Animals were fed twice per day restrictive amounts of hay and concentrate adapted to their requirements *ante* and *post*

partum, and had *ad libitum* access to drinking water. The composition of the commercial

concentrate (UFA 766) as communicated by the manufacturer, is shown in

Supplementary Table 1.

Zootechnical performance

Sheep and Goats were weighed closely before and directly after parturition. Due to the

suckling of the offspring, daily milk yield could not be assessed. The lambs and kids were

weighed at parturition and once between 33-59 days *post partum*, to determine the average

daily weight gain during the suckling phase. The variation in the time frame between first and

second weighing was due to management issues. This variation in observational periods was

considered as a control variable in the analysis of the growth performance data (see

subsection on statistical analysis below).

Sample collection and handling

For this experiment, a closed batch of hay and concentrates was ordered and three

representative samples each were taken according to good practice (VDLUFA, 2012) in order

to investigate the feed value by chemical analysis. Feed samples were directly subject to

chemical analysis after sample collection.

Blood was collected before morning feeding on days -21, -7, -6, -5, -4, -3, -2, -1, 0

(parturition), 1, 2, 3, 4, 5, 6, 7, 14, 28, and 56 in serum monovettes (9 and 6 mL, Vacuette®,

Greiner Bio-One Vacuette Schweiz GmbH) and subject to serum collection (1580 x g, 10

min) after allowing coagulation at room temperature. The serum was stored at -80°C until

further usage. Urine was collected in a beaker on days -21, -14, -7, 0, 7, and 14 before

morning feeding and stored at   -20°C until further usage. Fecal grab samples of each

animal were collected at days -21, -2, -1, and 0 before morning feeding and stored at -20°C

until further usage. Goats were subject to colonic mucosal biopsies by veterinary personal on

days -21, -7, 0, and 7. An endoscope (PCF-20, Olympus) equipped with biopsy forceps was

inserted 40 cm into the large intestine via the anus to collect 4 samples (1.5-2.0 mm diameter)

per animal*time. Two of said samples were immediately put into 4% neutral buffered

formalin and the remaining two samples were snap frozen in liquid nitrogen and stored at -

80°C.

<u>Sample analysis</u>

Feed was analyzed for dry matter, crude nutrients, fiber fractions, HCl-insoluble ash, and

mineral profile (Ca, P, Mg) according to standard procedures (VDLUFA, 2012).

Blood serum was analyzed for Ca, osteocalcin, $1,25\text{-}(OH)_2$ vitamin D (calcitriol), bone-

specific alkaline phosphatase (bAP), crosslinked carboxyterminal teleopeptide of type I

collagen (ICTP), and crosslaps (CTX) as described by Liesegang et al. (2007).

Feces and colostrum were analyzed for total Ca, and urine for total Ca and creatinine. The

urinary Ca excretion was expressed as ratio to urinary creatinine since the material could not

be sampled quantitatively. These analyzes and calculation have been described in detail by

Liesegang et al. (2007).

Goat colonic mucosal biopsies were subject to semi-quantitative immunohistochemical

determination of vitamin D receptor (VDR) abundance as originally described by Liesegang

et al. (2008) and Sidler-Lauff et al. (2010) in sufficient detail. For sheep, no suitable antibody

was available at the time of study.

<u>Bone mineral density</u>

Bone mineral density (BMD) was determined by quantitative computer tomography (qCT);

the respective equipment and approach has been described earlier in detail (Liesegang et al.,

2007).

<u>Statistical analysis</u>

The individual animal was the smallest experimental unit. All data analysis was performed

with SAS 9.4 (SAS Institute Inc.) applying the procedure MIXED. Repeated measures mixed

models were applied to analyze the effects of the fixed variables "group", "species", "time",

and interactions considering the individual animal nested under the respective feeding group

as a random factor. Repeated measures mixed modelling involved in any case the mapping of

covariance structures appropriate for the respective data structure following guidance by

Wang and Goonewardene (2004). Due to the limited availability of suitable antibodies, the

VDR protein expression over time was only analyzed in colonic biopsies of goats, therefore

no species effect could be estimated. The colostral Ca concentration of sheep and goats was

analyzed with an endpoint mixed model with the fixed effect of "group", "species", and

interaction, again including the individual animal nested under the respective feeding group as

random effect. In case the residuals of respective mixed models missed the precondition of

normality, the analysis was repeated after Log_{10}-transformation of data and displayed as such.

Data was presented in any case as least square mean values with Tukey-adjusted 95%

confidence limits. $P \leq 0.05$ was defined as threshold for a TYPE I statistical error.

The sample size of the study was designed to meet a minimum statistical power of $1-\beta = 0.8$

at $\alpha = 0.05$ (McDonald, 2014) for the primary outcome variables (quantitative and functional

measures of Ca metabolism). Earlier datasets on small ruminant Ca metabolism from our

institute served as datasets for prospective power analysis (Liesegang et al., 2007, Liesegang,

2008).

Results

<u>Animal health and development</u>

All sheep and goats remained healthy throughout the entire study period with no signs of pathology and hypocalcemia in particular, based on continuous veterinary surveillance. Sheep showed five cases of triplets, four cases of twins and one single birth (n = 22 lambs in total) whereas the majority of goats gave birth to twins with one case of triplets and two cases of single births (n = 23 kids in total). The sex ratio of sheep and goat offspring was (m/f) 0.71 and 1.30, respectively.

All lambs and kids were born with species-dependent birth weights (5.12 vs. 4.69 kg for lambs and kits, respectively; P = 0.06) irrespective of sex (P = 0.16) and unaffected by the mothers' dietary Ca supply *ante partum* (P = 0.94) (Figure 1A). However, lambs and kids of mothers fed high Ca *ante partum* showed significantly lower daily weight gain from milk intake than control animals (0.16 vs. 0.14 kg/d and 0.27 vs. 0.23 kg/d for lambs and kids, respectively, P = 0.02) irrespective of sex and species, which in both cases had distinct effects on the animals' growth performance (males grew faster than females; kids grew faster than lambs) (P = 0.07 and P < 0.0001, respectively) (Figure 1B).

<u>Quantitative markers of calcium metabolism</u>

Analyzed fecal Ca concentration during the observation period ranged between 1.46 and 5.19 % DM in sheep as well as 1.17 and 14.9 % DM in goats whereas the ranges for urinary Ca excretion were 0.005 and 0.41 mmol/mmol creatinine in sheep as well as 0.004 and 1.55 mmol/mmol creatinine in goats. Fecal and urinary data had to be log-transformed to meet the precondition of normality, accounting for fecal Ca ranges of 0.16 and 0.71 $\log_{10}$(% DM) for sheep as well as 0.07 and 1.17 $\log_{10}$(% DM) for goats. The respective log-transformed urinary

Ca excretion ranges were -2.31 and -0.38 $\log_{10}$(mmol/mmol creatinine) in sheep as well as -

2.42 and 0.19 $\log_{10}$(mmol/mmol creatinine) in goats.

Excess supply with dietary Ca *ante partum* significantly increased fecal Ca concentration

until parturition by at least two Log_{10}-orders of magnitude (P < 0.0001) irrespective of animal

species, which expressed a distinct effect on total analyzed fecal Ca (P = 0.05) (Figure 2A). In

contrast, the feeding regime had no effect on the urinary calcium excretion (P = 0.56), which

was only different with respect to the animal species (goats expressing higher values than

sheep) (P = 0.0003) and remained rather stable from 3 weeks *ante partum* to 14 d *post*

partum.

Serum Ca was not affected by the *ante partum* feeding group (P = 0.83) and gradually and

significantly (P < 0.0001) decreased independent of animal species towards parturition to

gradually increase again during lactation (Figure 3A). In any case, sheep showed higher

serum Ca levels than goats (P = 0.0003). The colostral Ca concentration was significantly

lower in goats compared to sheep (P = 0.002) but was not affected by the *ante partum* Ca

feeding regime (P = 0.86) (Figure 3B).

Figure 4 highlights the bone mineral density of dairy sheep and dairy goats from transition to

56 d in milk, as affected by Ca feeding during the last three weeks *ante partum*, time and

animal species. Excess Ca feeding *ante partum* did not affect bone mineral density (P = 0.63),

which significantly declined towards parturition to gradually increase *post partum* until d 56

of lactation (P < 0.0001). On average, sheep showed higher bone density than goats (P <

0.0001).

<u>Serum calcitriol and markers of bone formation</u>

Serum calcitriol was significantly changed with respect to time (P < 0.0001) and animal

species (P < 0.0001) but not by the *ante partum* feeding regime (P = 0.75), although numerical

differences between species-dependent treatment groups were evident (Figure 5A). Serum

levels in sheep gradually reduced towards parturition and fluctuated throughout whereas in

goats serum levels gradually increased until 4 d *post partum* to turnaround and reduce until

56 d *post partum*. Sheep showed higher serum levels in the first week of observation after

which the ratio changed for the benefit of the goats throughout the rest of the study.

Serum activity of bAP ranged between 38 and 251 U/L in sheep as well as 21 and 117 U/L in

goats. After Log_{10}-transformation to meet the precondition of normality, these ranges were

1.58 and 2.40 Log_{10}(U/L) in sheep as well as 1.32 and 2.07 Log_{10}(U/L) in goats. Feeding

group and time expressed no significant effect on bAP throughout the whole experiment (P =

0.59 and 0.12, respectively) (Figure 5B). Sheep showed in any case significantly higher serum

activities than goats ($P < 0.0001$).

Serum osteocalcin remained rather stable and comparable between species von 5 d *ante*

partum to 7 d *post partum*, after which it significantly and gradually increased in both species

($P < 0.0001$), with goats showing highest values and steepest increase over time until d 56

post partum (P = 0.007) (Figure 5C). Most notably, this increase in serum osteocalcin was

significantly higher in those sheep and especially goats who received excess dietary Ca during

the last three weeks *ante partum* (P = 0.02).

Serum markers of bone resorption

Serum ICTP showed no marked changes in both species until parturition, afterwards it

significantly increased over the first week post partum to gradually reduce until 56 d *post*

partum ($P < 0.0001$) (Figure 6A). Average serum levels over time were significantly higher

for sheep than goats ($P < 0.0001$). At no time did the feeding *ante partum* express any

significant effect on serum ICTP levels of dairy sheep and goats (P = 0.98).

Sheep showed markedly higher serum CTX levels during this study than goats ($P < 0.0001$)

(Figure 6 B). In both species serum levels significantly increased until the mid of week 1 *post*

partum. Afterwards, levels gradually declined until 56 d *post partum* (P < 0.0001). No

significant effect of the *ante partum* Ca feeding was observed (P = 0.59).

Caprine colonic abundance of mucosal vitamin D receptor

Figure 7 highlights the statistical evaluation of VDR immune reactivity in caprine colonic

mucosa until parturition, in goats fed 0.6% DM and 1.3% DM dietary Ca during transition.

Figure 8 presents exemplary immunohistochemical images in addition. Vitamin D receptor

abundance in caprine colonic mucosa was not significantly affected by *ante partum* feeding (P

= 0.19) or time (P = 0.1), respectively, though showed numerically reduced abundance from

14 d *ante partum* to parturition in goats fed the high Ca diet.

Discussion

An earlier meta-analysis concluded that the risk for hypocalcaemia in dairy cows significantly increases if dietary Ca during the peripartal phase is between 1.1-1.5% DM (Lean et al., 2006). This hypothesis has yet not been tested in dairy sheep and goats, therefore the present study investigated whether Ca fed at 1.3% DM (compared to 0.6% DM) during the last three weeks *ante partum* increases the incidence of hypocalcaemia *ante* and/or *post partum* by dysregulating Ca homeostasis.

Effects on quantitative Ca metabolism and the incidence of hypocalcaemia

Sheep and goats showed in any case significant differences in the expression of observed measures of quantitative Ca metabolism (fecal Ca, urinary Ca excretion, colostrum Ca, serum Ca, BMD). Differences in magnitude of these parameters between both species have also been observed in several earlier comparative studies on the matter. Our observation of higher values in sheep with respect to urinary Ca excretion and colostral Ca and bone mineral density are in line with earlier studies (Liesegang et al., 2006, 2007, Liesegang, 2008). We made the same observation with serum Ca, which agrees to Kohler et al. (2013) and Liesegang (2008), though Wilkens et al. (2014) observed the opposite with goats expressing higher values than sheep. The reason for this discrepancy is unclear at the moment.

The feeding showed mostly no or just numerical effects on quantitative measures. Increasing dietary Ca feeding during the peripartal phase significantly increased fecal Ca concentration by at least 2fold during the last three weeks but did not significantly affect serum and colostral Ca concentration and urinary Ca excretion in both species at any given point during the study. Most notably, no signs of hypocalcemia or any pathology were noted *ante* or *post partum*, respectively. This suggests that the organism was able to dampen Ca absorption and/or recirculate passively absorbed surplus Ca back into the gastrointestinal tract. Furthermore, the generous peripartal Ca supply did not impair the animals' ability to

compensate the sudden increase in Ca losses due to the onset of lactation, though the opposite

is expected for dairy cows based on the meta-analysis by Lean et al. (2006). Calcitriol, the

active form of vitamin D in the mammal organism, improves jejunal Ca absorption when

dietary Ca supply is deficient. Conversely, the efficiency of jejunal Ca absorption is markedly

lower when the dietary Ca supply is sufficient (Abdel-Hafeez et al., 1982). It may be

speculated that a dietary excess supply with Ca may further impair active calcitriol-dependent

Ca absorption. The numerically reduced abundance of colonic mucosal VDR in our treatment

goats *ante partum* is in support of this notion and highlights the adjustment of the homeostatic

regulation towards a dampening of Ca absorption capacity in response to high Ca during the

peripartal phase. Consequently, bone mineral density was not affected by the dietary treatment

at any time of observation. Literature on the effects of Ca oversupply in sheep and goats is

scarce, particularly with respect to the peripartal phase. The only study on sheep by

Braithwaite (1979), however, agrees with our observations. The colleague studied gradual

differences in dietary Ca intake in wethers and observed increasing fecal Ca losses while

fractional Ca absorption remained stable at ~10% when feeding 0.6, 1.2, and 2.4% Ca in

dietary DM, respectively. In parallel, bone Ca resorption dropped inversely to the increase in

Ca intake and passively absorbed surplus was increasingly redirected into the gastrointestinal

tract. Given the comparable response of our goats in the present study, we hypothesize that the

counter-regulation of quantitative Ca homeostasis to dietary Ca excess is comparable between

both species, apart from the above discussed differences in baseline values of observed

parameters. Neither sheep nor goats showed any relevant response in urinary calcium

excretion in response to the feeding regime, confirming once more results from earlier studies

suggesting livestock ruminants do not clear calcium surplus via the kidneys (Herm et al.,

2015, Wild et al., 2021).

Taken together, sheep and goats were able to stabilize quantitative Ca homeostasis and did not

express signs of hypocalcemia or milk fever neither *ante* nor *post partum*, despite a drastic

increase in dietary Ca during the last weeks *ante partum* from 0.6 to 1.3% DM.

<u>Effects on serum markers of bone formation and resorption</u>

We observed a core set of serum bone formation and resorption markers, which have been

earlier established in ruminants transitioning from gestation to lactation (Liesegang et al.,

2000, Liesegang et al., 2007). Observed measures of bone formation and resorption showed

changes over time, that have earlier been attributed to the species-dependent adaption towards

parturition and the onset and further course of lactation (Liesegang et al., 2006, 2007,

Liesegang, 2008, Kohler et al., 2013, Wilkens et al., 2014). However, none of the observed

parameters responded to the feeding treatment during the last weeks of gestation and mostly

also *post partum*. Consequently, the BMD was not affected and suggests the feeding

intervention with 1.3% DM Ca in the diet caused no obvious dysregulation of Ca and bone

homeostasis in neither species.

However, serum osteocalcin showed a significant gradual increase in concentration from d7 to

d56 of lactation in animals receiving the high dietary Ca *ante partum*. This response was true

for both species but most pronounced in the dairy goats. Serum osteocalcin is considered a

marker of bone formation, with increasing levels pointing towards rising embedding of

calcium in the skeleton (Liesegang et al., 2000). This observation was puzzling given the lack

of response of other markers of bone formation (calcitriol, bAP) and resorption (ICTP, CTX))

and especially since the effect occurred 8d after the suspension of the high Ca feeding. To the

best of our knowledge such a response pattern has yet not been reported for serum osteocalcin

and it raises questions on its physiological purpose. If the rise in serum osteocalcin indeed

suggests increasing bone formation activity from d 8 to d 56 of lactation, this did not affect

the bone mineral density (or concentration; data not shown). However, the necessity for

increased bone formation in the high Ca group during lactation appears questionable,

especially since the challenge feeding was stopped 8d before the rise in serum levels.

Biomedical research efforts in recent years suggest a hormonal role of uncarboxylated

osteocalcin, involving the regulation of systemic energy metabolism (Lee et al., 2007), muscle

energy metabolism (Mera et al., 2016), brain development (Obri et al., 2018), and others. At

this point, the meaning of these results is up to speculation, hence, more targeted research on

the matter is urgently needed.

<u>Effects on offspring development</u>

High Ca treatment of mothers *ante partum* was associated with decreased average daily

weight gain of suckling lambs and kids. Since the offspring showed no differences in birth

weight that were associated to the feeding of the mothers *ante partum*, it can be concluded

that this response was due to effects on either the physiological status of the mothers or their

offspring or both. The aforementioned differences in serum osteocalcin are pointing towards a

distinct physiological conditioning of mothers in the treatment group. This may have affected

the quantity and quality of secreted milk and therefore the growth response of their offspring,

but this needs yet to be confirmed. Furthermore, high Ca feeding during the last three weeks

post partum and the associated physiological adaption of the mothers may have promoted

fetal programming in the offspring that impaired their digestive capacity for calcium and

maybe other nutrients therefore negatively affecting their growth response. The possibility of

mineral nutrition during gestation to induce fetal programming in ruminants either for the

benefit or disadvantage of the offspring has recently been reviewed by Anas et al. (2023) for

cattle, highlighting a multitude of already established interactions as well as exiting future

routes for research on the matter. Furthermore, an earlier study in rats tested the effects of

varying dietary Ca supply during gestation including excess supply at 1.2% DM (Li et al.,

2018). They observed abnormal expression of hepatic and adipose genes in the offspring in

response thereby promoting dyslipidemia and hepatic lipid accumulation. Unfortunately, no

data on calcium homeostasis in rat mothers and their offspring was obtained. Nevertheless, it

may be argued that excess supply with a mineral over several weeks may induce epigenetic

changes in the mother, which have the potential to be inherited to the offspring. For example,

piglets exposed to excess zinc supply over 14 days post-weaning show not only

downregulation of active zinc transport at the gut mucosa (Martin et al., 2013) but in fact a

methylation of CpG islands of the respective transporter gene, which further impairs its

transcription (Karweina et al., 2015). This has yet not been shown for calcium but is has been

demonstrated that Ca^{2+} signaling pathways are involved in epigenetic modifications (Sharma

et al., 2014). It may therefore be interesting to have a closer look on the epigenetic

consequences of dietary Ca-excess as well as opposite feeding regimes during gestation and

their potential to epigenetically imprint Ca homeostatic regulation and development of the

offspring.

<u>The geographical region as a potential confounding factor</u>

Depending on the geographic location, Ca levels in the environment are quite variable. In

Switzerland, animals are often confronted with high levels of native Ca in diets, originating

from the Ca-rich geological formations, which shape the chemical composition of soil in

certain areas. For example an earlier survey of Schlegel et al. (2016) found a mean Ca

concentration of 7 g/kg DM in 236 samples of mixed grassland from Posieux, Switzerland,

and our own experience with hay qualities delivered to our animal facility in the Zurich area

are in line with this observation. This leads to quite relevant intake levels of native Ca from

feed particularly when grass-based roughage is fed, even when mineral Ca supplements are

not applied. Since the organism attempts to maintain Ca homeostasis by fine-tuning

absorption, losses, and mobilization of skeletal reserves according to the environmental

conditions (Martín-Tereso and Martens, 2014, Wilkens and Muscher-Banse, 2020), it may be

argued that sheep and goats raised predominantly on Ca-rich roughage may be better suited to

deal with dietary Ca overload than such in other areas. However, it may also suggest that

during Ca excess due to active supplementation of inorganic Ca to such raw materials, a

metabolic imprinting as hypothesized for the offspring from the present study may more

likely promote the observed adverse effects on daily weight gain of suckling lambs and goat

kids. Therefore, the present results must be considered with caution when it comes to the

general translation into dairy sheep and goat feeding systems. More data is urgently needed

also from regions with less Ca in the soil.

Conclusion

The present study did not find any signs of hypocalcaemia in sheep and goats challenged with

excessive Ca during the transition phase. However, serum osteocalcin showed increasing

concentrations in lactating sheep and goats fed high Ca *ante partum* compared to the

respective control. Interestingly, though no obvious negative effects of the feeding regime

could be observed for the mothers, the offspring of sheep and goats fed high Ca during the

transition phase did grow slower over time compared to control, despite comparable birth

weights. This may have originated from either effects on the physiological status of the

mothers, thereby affecting milk yield and quality, or the offspring or both. The hypothesis of a

potential adverse effect of high dietary Ca *ante partum* on the digestive capacity of the

offspring via epigenetic imprinting may be an exciting avenue for future research. Our

animals were reared in a Ca-rich environment since their birth, therefore, we cannot rule out

that the present findings are exclusive for this situation. Follow-up studies are urgently needed

and should also comprise observations in production regions with less Ca in the soils.

Notes

Raw datasets can be requested via E-mail to the corresponding author.

This study received no external funding.

We thank Dr. med. vet. Sybil Lüthi for excellent technical assistance in sample and data collection. We further thank the technical staff in the animal and laboratory facilities of the Institute of Animal Nutrition and Dietetics, Vetsuisse-Faculty, University of Zurich, Switzerland.

All procedures used in the experiment involving animals were in accordance with the Swiss guidelines for animal welfare at the time of study and were registered and approved by the Cantonal Veterinary Office (ZH 9/2006).

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

Abdel-Hafeez, H. M., M. Manas-Almendros, R. Ross, A. D. Care, and D. H. Marshall. 1982. Effects of dietary phosphorus and calcium on the intestinal absorption of Ca in sheep. Brit. J. Nutr. 47: 69-77.

Agroscope. 2017. Fütterungsempfehlungen für Wiederkäuer (Grünes Buch). Vol. https://www.agroscope.admin.ch/agroscope/de/home/services/dienste/futtermittel/fuetterungsempfehlungen-wiederkaeuer.html. Agroscope.

Anas, M., W. J. S. Diniz, A. C. B. Menezes, L. P. Reynolds, J. S. Caton, C. R. Dahlen, and A. K. Ward. 2023. Maternal mineral nutrition regulates fetal genomic programming in cattle: A review. Metabolites 13: 593.

Babcock, S. M. 1905. The addition of salt to the ration of dairy cows. University of Wisconsin Experiment Station, 22nd Annual Report.

Braithwaite, G. D. 1979. The effect of dietary intake of calcium and phosphorus on their absorption and retention by mature Ca-replete sheep. J. Agric. Sci. 92: 337-342.

Goff, J. P. 2008. The monitoring, prevention, and treatment of milk fever and subclinical hypocalcemia in dairy cows. The Veterinary Journal 176 (1): 50-57.

Herm, G., A. S. Muscher-Banse, G. Breves, B. Schröder, and M. R. Wilkens. 2015. Renal mechanisms of calcium homeostasis in sheep and goats. J. Anim. Sci. 93: 1608-1621.

Hernández-Castellano, L. E., L. L. Hernandez, and R. M. Bruckmaier. 2020. Review: Endocrine pathways to regulate calcium homeostasis around parturition and the prevention of hypocalcemia in perparturient dairy cows. Animal 14 (2): 330-338.

Hoenderop, J. G., O. Dardenne, M. van Abel, A. W. van der Kemp, C. H. van Os, R. St. Arnaud, and R. J. Bindels. 2002. Modulation of renal Ca^{2+} transport protein genes by dietary Ca^{2+} and 1,25-dihydroxyvitamin D_3 in 25-hydroxyvitamin D_3-1alpha-hydroxylase knockout mice. FASEB J. 16: 1398-1406.

482 Karweina, D., S. Kreuzer-Redmer, U. Müller, T. Franken, R. Pieper, U. Baron, S. Olek, J.

483    Zentek, and G. A. Brockmann. 2015. The zinc concentration in the diet and the length

484    of the feeding period affect the methylation status of the ZIP4 zinc transporter gene in

485    piglets. PLoS ONE 11: e0143098.

486 Ko, S. H., G. S. Lee, T. T. Vo, E. M. Jung, K. C. Choi, K. W. Cheung, J. W. Kim, J. G. Park,

487    G. T. Oh, and E. B. Jeung. 2009. Dietary calcium and 1,25-dihydroxyvitamin D3

488    regulate transcription of calcium transporter genes in calbindin-D9k knockout mice.

489    Journal of Reproduction and Development 55: 137-142.

490 Kohler, M., F. Leiber, H. Willems, L. Merbold, and A. Liesegang. 2013. Influence of altitude

491    on vitamin D and bone metabolism of lactating sheep and goats. J. Anim. Sci. 91:

492    5259-5268.

493 Lean, I. J., P. J. DeGaris, D. M. McNeil, and E. Block. 2006. Hypocalcemia in dairy cows:

494    meta-analysis and dietary cation anion difference theory revisited. J. Dairy Sci. 89 (2):

495    669-684.

496 Lee, N. K., H. Sowa, E. Hinoi, M. Ferron, J. D. Ahn, C. Confavreux, R. Dacquin, P. J. Mee,

497    M. D. McKee, D. Y. Jung, Z. Zhang, J. K. Kim, F. Mauvais-Jarvis, P. Ducy, and G.

498    Karsenty. 2007. Endocrine regulation of energy metabolism by the skeleton. Cell 130

499    (3): 456-469.

500 Li, P., X. Chang, X. Fan, C. Fan, T. Tang, R. Wang, and K. Qi. 2018. Dietary calcium status

501    during maternal pregnancy and lactation affects lipid metabolism in mouse offspring.

502    Scientific Reports 8: 16542.

503 Liesegang, A. 2008. Influence of anionic salts on bone metabolism in periparturient dairy

504    goats and sheep. J. Dairy Sci. 91: 2449-2460.

505 Liesegang, A., J. Risteli, and M. Wanner. 2006. The effects of first gestation and lactation on

506    bone metabolism in dairy goats and milk sheep. Bone 38: 794-802.

Liesegang, A., J. Risteli, and M. Wanner. 2007. Bone metabolism of milk goats and sheep
during second pregnancy and lactation in comparison to first lactation. J. Anim.
Physiol. Anim. Nutr. 91: 217-225.

Liesegang, A., M.-L. Sassi, J. Risteli, R. Eicher, M. Wanner, and J.-L. Riond. 2000.
Biochemical markers of bone formation and resoprtion around parturition and during
lactation in dairy cows with high and low standard milk yields. J. Dairy Sci. 83 (8):
1773-1781.

Liesegang, A., K. Singer, and A. Boos. 2008. Vitamin D receptor amounts across different
segments of the gastrointestinal tract in Brown Swiss and Hosltein Friesian cows of
different age. J. Anim. Physiol. Anim. Nutr. 92: 316-323.

Martín-Tereso, J. and H. Martens. 2014. Calcium and magnesium physiology and nutrition in
relation to the prevention of milk fever and tetany (dietary management of
macrominerals in preventing disease). Veterinary Clinics of North America: Food
Animal Practice 30 (3): 643-670.

Martin, L., U. Lodemann, A. Bondzio, E. M. Gefeller, W. Vahjen, J. R. Aschenbach, J. Zentek,
and R. Pieper. 2013. A high amount of dietary zinc changes the expression of zinc
transporters and metallothionein in jejunal epithelial cells in vitro and in vivo but does
not prevent zinc accumulation in jejunal tissue of piglets. J. Nutr. 143 (8): 1205-1210.

McDonald, J. H. 2014. Power Analysis. Pages 40-44 in Handbook of Biological Statistics, 3[rd]
ed. J. H. McDonald, ed. Sparky House Publishing, Baltimore, MD, USA.

Mera, P., K. Laue, M. Ferron, C. Confavreux, J. Wei, M. Galán-Díez, A. Lacampagne, S. J.
Mitchell, J. A. Mattison, Y. Chen, J. Bacchetta, P. Szulc, R. N. Kitsis, R. de Cabo, R.
A. Friedman, C. Torsitano, T. E. McGraw, M. Puchowicz, I. Kurland, and G. Karsenty.
2016. Osteocalcin signaling in myofibers is necessary and sufficient for optimum
adaptation to exercise. Cell. Metab. 23 (6): 1078-1092.

Obri, A., L. Khrimian, G. Karsenty, and F. Oury. 2018. Osteocalcin in the brain: from

embryonic development to age-related decline in cognition. Nat. Rev. Endocrinol. 14

(3): 174-182.

Oetzel, G. R. 1988. Parturient paresis and hypocalcemia in ruminant livestock. Vet. Clin.

North Am. Food Anim. Pract. 4: 351-364.

Schlegel, P., U. Wyss, Y. Arrigo, and H. D. Hess. 2016. Mineral concentrations of fresh

herbage from mixed grassland as influenced by botanical composition, harvest time

and growth stage. Anim. Feed Sci. Technol. 219: 226-233.

Sharma, A., H. Nguyen, C. Geng, and H. Lou. 2014. Calcium-mediated histone modifications

regulate alternative splicing in cardiomyocytes. Proc. Natl. Acad. Sci. USA 111 (46).

Sidler-Lauff, K., A. Boos, M. Kraenzlin, and A. Liesegang. 2010. Influence of different

calcium supplies and a single vitamin D injection on vitamin D receptor- and calbindin

D9k-immunoreactivities in the gastroinestinal tract of goat kids. J. Anim. Sci. 88 (11):

3598-3610.

van Cromphaut, S. J., K. Rummens, I. Stockmans, E. van Herck, F. A. Dijcks, A. G.

Ederveen, P. Carmeliet, R. Bouilon, and G. Carmeliet. 2003. Intestinal calcium

transporter genes are upregulated by estrogens and the reproductive cycle through

vitamin D receptor-independent mechanisms. J. Bone Miner. Res. 18: 1725-1736.

VDLUFA. 2012. VDLUFA-methods book III: The chemical analysis of feedstuffs. VDLUFA-

Verlag, Darmstadt, Germany.

Wang, Z. and L. A. Goonewardene. 2004. The use of MIXED models in the analysis of

animal experiments with repeated measures data. Can. J. Anim. Sci. 84 (1): 1-11.

Wild, K. J., W. Siegert, W. M. Windisch, K.-H. Südekum, and M. Rodehutscord. 2021. Meta-

analysis-based estimates of efficiency of calcium utilization by ruminants. Animal 15

(8): 100315.

Wilkens, M., J. Richter, D. R. Fraser, A. Liesegang, G. Breves, and B. Schröder. 2012. In
contrast to sheep, goats adapt to dietary calcium restriction by increasing intestinal
absorption of calcium. Comparative Biochemistry and Physiology Part A, Molecular
& Integrative Physiology 163 (3-4): 396-406.

Wilkens, M. R., A. Liesegang, J. Richter, D. R. Fraser, G. Breves, and B. Schröder. 2014.
Differences in peripartal plasma parameters related to calcium homeostasis of dairy
sheep and goats in comparison to cows. J. Dairy Res. 81 (3): 325-332.

Wilkens, M. R. and A. S. Muscher-Banse. 2020. Review: Regulation of gastrointestinal and
renal transport of calcium and phosphorus in ruminants. Animal 14: s29-s43.

Table 1. Chemical composition and daily intake levels of feedstuffs fed during transition and lactation
of dairy sheep and goats.

Chemical analysis, per kg DM	Hay (2nd cut)	Concentrate	Concentrate + $CaCO_3$
Energy, MJ NEL	5.90	7.00	6.70
Dry matter, g	928	884	896
Crude protein, g	155	133	115
Crude fiber, g	263	69.3	64.2
NDF, g	654	267	194
ADF, g	357	90.4	70.2
ADL, g	38.4	18.9	13.2
Crude fat, g	28.0	61.5	49.2
Crude ash, g	81.1	54.2	170
HCl-insoluble ash, g	13.3	4.30	4.60
Ca, g	5.90	6.90	62.3
Mg, g	2.00	1.90	1.40
P, g	3.30	4.60	4.30

Component, kg DM/d	Sheep			Goats	
	Control	Treatment		Control	Treatment
		ante partum			
Hay (2nd cut)	2.0	2.0		2.0	2.0
Concentrate[1]	0.3	0.3		0.3	0.3
$CaCO_3$	-	0.042		-	0.042
Total DM fed	2.3	2.342		2.3	2.342
Nutrient intake, g/d					
Crude protein	350	344		350	344
Energy (MJ NEL)	13.9	13.8		13.9	13.8
Ca	13.9	30.5		13.9	30.5
P	7.98	7.89		7.98	7.89
Mg	4.57	4.42		4.57	4.42
		post partum			
Hay (2nd cut), kg/d	2.4	2.4		2.4	2.4
Concentrate, kg/d	0.5	0.5		0.7	0.7
$CaCO_3$, g/d	-	-		-	-
Total DM fed	2.9	2.9		3.1	3.1
Nutrient intake, g/d					
Crude protein	438	429		465	452
Energy (MJ NEL)	17.7	17.5		19.1	18.8
Ca	17.6	45.3		19.0	57.8
P	10.2	10.1		11.1	11.0
Mg	5.75	5.5		6.13	5.78

571   Commercial concentrate UFA 765. For the composition of the concentrate based on manufacturer
572   instructions, please see Supplementary Table 1.

Figure 1. Effects of varying dietary Ca (0.6 vs. 1.3% DM) of dairy sheep and goats during the last three weeks *ante partum* as well as sex and species of offspring on the birth weight (A) and daily weight gain during the suckling phase (B). Effective sample size was five sheep and six goats per group, which gave birth to 22 and 23 lambs and kids in total, respectively. Data is presented as least square mean values with Tukey-adjusted upper 95% confidence limits based on linear mixed model analysis. "Age" refers to the time of second weighing during the suckling phase, which differed between individual animals due to management reasons (range: 33-59 d *post partum*) and was therefore introduced as a continuous control variable to the statistical analysis of daily weight gain data. G*S and G*Sex refer to statistical interactions between feeding group and species of animal or sex of offspring, respectively. P ≤ 0.05 was defined as statistical type I error threshold.

Figure 2. Effects of varying dietary Ca (0.6 vs. 1.3% DM) of dairy sheep and goats during the

last three weeks *ante partum* as well as time and animal species on the fecal calcium

concentration (A) and urinary calcium excretion (B) from three weeks *ante partum* to

parturition and 14 d *post partum*, respectively. Effective sample size was five sheep and six

goats per group. Data is presented as Log_{10}-transformed least square mean values with Tukey-

adjusted upper or lower 95% confidence limits, respectively, based on repeated measures

linear mixed model analysis. G*T, G*S and S*T refer to statistical interactions between

feeding group and animal species or time, respectively, as well as animal species and time. $P \leq$

0.05 was defined as statistical type I error threshold.

Figure 3. Effects of varying dietary Ca (0.6 vs. 1.3% DM) of dairy sheep and goats during the last three weeks *ante partum* as well as time and animal species on serum calcium (A) and colostral calcium (B) from three weeks *ante partum* to parturition and 56 d *post partum*, respectively. Effective sample size was five sheep and six goats per group. Data is presented as least square mean values with Tukey-adjusted upper 95% confidence limits based on repeated measures linear mixed model analysis and regular linear mixed model analysis, respectively. G*T, G*S and S*T refer to statistical interactions between feeding group and animal species or time, respectively, as well as animal species and time. P ≤ 0.05 was defined as statistical type I error threshold.

Figure 4. Effects of varying dietary Ca (0.6 vs. 1.3% DM) of dairy sheep and goats during the

last three weeks *ante partum* as well as time and animal species on bone mineral density from

two weeks *ante partum* to 56 d *post partum*. Effective sample size was five sheep and six

goats per group. Data is presented as least square mean values with Tukey-adjusted upper

95% confidence limits based on repeated measures linear mixed model analysis. G*T, G*S

and S*T refer to statistical interactions between feeding group and animal species or time,

respectively, as well as animal species and time. P ≤ 0.05 was defined as statistical type I error

threshold.

Figure 5. Effects of varying dietary Ca (0.6 vs. 1.3% DM) of dairy sheep and goats during the last three weeks *ante partum* as well as time and animal species on serum 1,25(OH)$_2$ Vitamin D (A), serum bone specific alkaline phosphatase activity (B), and serum osteocalcin (C) from three weeks and 5 d *ante partum*, respectively, to 56 d *post partum*. Effective sample size was five sheep and six goats per group. Data is presented as least square mean values (Log10-scale in case of bone specific alkaline phosphatase activity) with Tukey-adjusted upper 95% confidence limits based on repeated measures linear mixed model analysis. G*T, G*S and S*T refer to statistical interactions between feeding group and animal species or time, respectively, as well as animal species and time. P ≤ 0.05 was defined as statistical type I error threshold.

Figure 6. Effects of varying dietary Ca (0.6 vs. 1.3% DM) of dairy sheep and goats during the

last three weeks *ante partum* as well as time and animal species on serum ICTP (A) and

serum CTX (B) from three weeks and 3 d *ante partum*, respectively, to 56 d *post partum*.

Effective sample size was five sheep and six goats per group. Data is presented as least square

mean values (Log10-scale in case of bone specific alkaline phosphatase activity) with Tukey-

adjusted upper 95% confidence limits based on repeated measures linear mixed model

analysis. G*T, G*S and S*T refer to statistical interactions between feeding group and animal

species or time, respectively, as well as animal species and time. ICTP, crosslinked

carboxyterminal teleopeptide of type I collagen; CTX, crosslaps. $P \leq 0.05$ was defined as

statistical type I error threshold.

Figure 7. Effects of varying dietary Ca (0.6 vs. 1.3% DM) of dairy sheep and goats during the

last three weeks *ante partum* as well as time on caprine vitamin D receptor abundance in

colonic mucosa from two weeks *ante partum* to 7 d *post partum*. Effective sample size was

six goats per group. Data is presented as least square mean values with Tukey-adjusted upper

95% confidence limits based on repeated measures linear mixed model analysis. G*T, refers

to statistical interaction between feeding group and time. P ≤ 0.05 was defined as statistical

type I error threshold. See Figure 8 for exemplary immunohistochemical images.

Figure 8. Exemplary immunohistochemical staining of the caprine Vitamin D Receptor in the

colon of goats at parturition, fed 0.6 % DM (left) or 1.3 % DM (right) dietary calcium during

the last three weeks prepartum. Arrows mark SE, surface epithelium; SG, surface gland

sections; IG, intermediate gland sections; BG, basal gland sections. Samples were taken at

parturition with an endoscope equipped with biopsy forceps 40 cm into the large intestine

from the anus. Goats fed excessive Ca prepartum showed reduced immune reactivity to VDR

staining, though this could not be statistically confirmed. Figure 7 shows the statistical

analysis of associated immune reactivity.