High calcium to sheep and goats antepartum

Antepartum high dietary supply of calcium affects bone homeostasis and offspring growth

in dairy sheep and dairy goats

D. Brugger[1], A. Liesegang[1]

[1]Institute of Animal Nutrition and Dietetics, Vetsuisse-Faculty, University of Zurich

Annette Liesegang, Institut für Tierernährung und Diätetik, Winterthurerstrasse 270, 8057

Zurich, Switzerland, +41 44 635 88 23, aliese@nutrivet.uzh.ch

Interpretive Summary

Many studies have focused on deficient Ca supply to small dairy ruminants, and little is known about high dietary Ca during the last days antepartum. Here we show that multiparous dairy goats and sheep were able to balance their Ca homeostasis despite the high dietary Ca supply with no consequences for their health. However, we observed that animals fed high dietary Ca showed a gradual increase in serum osteocalcin postpartum and their suckling offspring showed reduced average daily weight gain. The functional background and practical consequences are yet unknown and may direct future research on the matter.

Abstract

This study investigated the effects of high dietary Ca supplementation during the final 21 days antepartum on Ca and bone homeostasis in dairy sheep and goats, and the growth response of their suckling offspring. Multiparous dairy sheep (n=5/group, 10 animals total) and goats (n=6/group, 12 animals total) were randomly assigned to 2 experimental groups. Feeding occurred restrictively (2.3 kg DM/d*animal^{-1} antepartum for both species; 2.9 and 3.1 kg DM/d*animal^{-1} for sheep and goats postpartum, respectively) according to recommendations with the exception of Ca during antepartum, where 1 group received a high Ca diet with CaCO$_3$ (1.3% in DM; 2.49fold recommendations) based on hay and concentrate 3 weeks antepartum, and the other group received no extra Ca on top (0.6% in DM; 1.15fold recommendations). Experimental feeding ended with parturition and animals were henceforth fed according to Swiss recommendations for lactating sheep and goats and kept together with their suckling offspring. The observation period spanned from 21 days antepartum to 56 days postpartum. Animals were subject to continuous veterinary surveillance and monitored for signs of milk fever. Data collection comprised quantitative and functional parameters of Ca and bone homeostasis as well as birth weights and daily weight gain of suckling lambs and kids. Data was analyzed by repeated measures and endpoint mixed models. The results of quantitative markers indicated that high dietary Ca antepartum significantly increased fecal Ca concentrations until parturition, suggesting efficient physiological mechanisms to manage Ca overload by increasing fecal excretion. No significant differences were observed in serum Ca levels, urinary Ca excretion, or bone mineral density between antepartum Ca feeding groups at any point in time of observation, indicating stable Ca homeostasis despite the dietary challenge. Species-specific differences in quantitative markers were noted including serum and colostral Ca levels as well as bone mineral density and mostly reflected results from earlier comparative studies. Serum 1,25-(OH)$_2$ vitamin D (**calcitriol**) as well as markers of bone formation and

resorption were monitored, revealing significant increases in serum osteocalcin postpartum in goats and sheep fed high Ca antepartum. All other serum markers including calcitriol were not affected by the feeding regime but again by the species of animal, in line with previous works. All repeated measures were significantly affected by time except for urinary Ca and bone specific alkaline phosphatase activity in serum. Offspring of sheep and goats in the high Ca group exhibited significantly reduced average daily weight gain compared to the 0.6 % in DM Ca group, despite similar birth weights. We conclude that while dairy sheep and goats effectively managed high dietary Ca intake without overt signs of hypocalcemia or milk fever under the present experimental conditions, there were notable impacts on offspring growth and potential long-term physiological effects that require further research. These findings contribute to the understanding of mineral nutrition in late gestating dairy goats and sheep and stresses the need for further research on balanced dietary strategies to optimize health and productivity of lactating animals and their offspring.

Key words: Transition phase, Calcium Homeostasis, Small Ruminant, Maternal Nutrition

Nonstandard abbreviations: bAP = bone-specific alkaline phosphatase activity; BMD = bone mineral density; calcitriol = 1,25-$(OH)_2$ vitamin D; CTX = crosslaps;; ICTP = crosslinked carboxyterminal teleopeptide of type I collagen; VDR = vitamin D receptor

Introduction

To avoid a drop in circulatory Ca, the organism has 3 options to stabilize the system, by increasing absorption capacity at the gut barrier, limiting endogenous losses via the gastrointestinal tract and kidneys, and mobilizing Ca from the skeleton (Wilkens and Muscher-Banse, 2020). With respect to the regulation of Ca homeostasis, the animal species under study is a key factor to consider. For example, sheep and goats have been demonstrated to respond quite differently to Ca-restricted feeding with the sheep tending quicker to bone mobilization while the latter will initially attempt to compensate via more efficient absorption from the small intestinal mucosa (Wilkens et al., 2012a). A relevant regulation of renal Ca excretion and reabsorption in response to changes in the endogenous Ca status has been reported for mice (Hoenderop et al., 2002, van Cromphaut et al., 2003, Ko et al., 2009), but was not confirmed in ruminants including sheep, goats (Herm et al., 2015) and cows (Wild et al., 2021). Quantitative adaption to fluctuations in the endogenous Ca status are regulated by a complex system of endocrine factors, that control the regulation of absorption, excretion and bone remodeling, the detailed discussion of which exceeds the capacities of this introduction (see the reviews by Martín-Tereso and Martens (2014) and Wilkens and Muscher-Banse (2020)).

In lactating dairy ruminants, a proper endogenous Ca status is especially important since milk supplies high levels of Ca to the offspring on a daily basis and the high yields and associated Ca losses with milk of modern genotypes disproportionately increase Ca demand compared to the mere needs of the offspring, which do not differ much shortly before and after parturition (Hernández-Castellano et al., 2020). In this context, the last weeks antepartum are critical. In high-yielding dairy cows, the sudden onset of lactation after birth in combination with impaired feed intake around parturition puts animals in danger of developing milk fever if the mobilization of bone stores lacks capacity to bridge this gap in dietary intake. This is typically accompanied by hypocalcemia, mostly also by hypophosphatemia and often hypomagnesemia

(Liesegang et al., 2007, Goff, 2008, Martín-Tereso and Martens, 2014). Symptoms develop within 48 h postpartum and may include listlessness, coma, and sternal recumbence. The situation for sheep and goats is different. Due to their lower milk yield, meat sheep breeds develop milk fever rarely postpartum though a higher tendency for carrying twin or even triplet fetuses may increase the risk of developing a hypocalcemia during the last trimester given an increasing fetal Ca requirement during the last 4 weeks antepartum due to the calcification of the fetal skeleton. Sheep typically show preceding anorexia and stiff gait before sternal recumbence. Dairy sheep breeds and most goat breeds combine a high milk yield potential with a high probability for carrying multiple fetuses, consequently clinical events of hypocalcemia antepartum and postpartum are likely. High-yielding dairy goats and sheep develop a milk fever comparable to dairy cows but much later at 1-3 weeks postpartum. If antepartum symptoms arise they are less pronounced than the postpartum complex (Oetzel, 1988).

Dietary and metabolic factors that affect the incidence of milk fever are various and the underlying literature is dominated by studies in cows. Amongst relevant factors are the DCAD, Mg, P, and Ca in the antepartum diet (Lean et al., 2006). There has been some debate regarding the proper dietary Ca concentration antepartum, with some sources suggesting that diet Ca at 1-1.5% in DM or even 2-3% does not affect the incidence of milk fever whereas others recommend going below 1%. An earlier meta-analysis on the matter attempted to bring some order in the chaos. The authors developed 2 models using data from 51 trials, updating the findings of an earlier meta-analysis by Oetzel (1991). They concluded that dietary calcium exhibits a non-linear relationship with milk fever incidence, with both exceptionally low and very high dosages showing protective effects. Model I included antepartum DCAD, as well as diet Ca, P, Mg, and diet exposure time as variables while Model II included antepartum dietary K and S in replacement of DCAD. Both models suggested a quadratic effect of dietary Ca on the pathogenesis of milk fever (odds ratios 0.131 and 0.115, respectively). Simulations with

Model I based on dietary Ca varying from 0 to 2.5 % in DM and assuming all other variables remain stable at levels associated with low milk fever risk, yielded a bell curve of predicted milk fever incidence (%) over varying diet Ca with a peak of ~+3.8% incidence increase at 1.35 % in DM dietary Ca. Based on this simulation it has been suggested that dietary Ca concentrations between 1.1-1.5% in DM should be avoided, whereas higher or lower levels gradually reduce the incidence of milk fever (Lean et al., 2006). These findings would explain the inconsistency between available studies on the interaction of dietary Ca dosage and milk fever incidence. This raises an interesting question about the situation in other livestock ruminants.

Studies on high dietary calcium (Ca) levels in gestating sheep and goats are limited, and specific guidelines for managing dietary Ca to prevent milk fever in these species do not exist, particularly in Switzerland. Our experimental design was informed by Lean et al. (2006), who found that in cows, dietary Ca levels between 1.1-1.5% in DM (vs. 0.6% in DM) in antepartum feed could increase milk fever incidence by up to 3.8%. This study aimed to evaluate the effects of high dietary Ca in dairy sheep and goats during the last 21 days antepartum on Ca homeostasis and bone metabolism and to provide initial data on potential risks extending into lactation. Given the naturally high environmental Ca in alpine regions, including Switzerland, where average Ca concentrations in forage can reach 7 g/kg DM (Schlegel et al., 2016), oversupply with dietary Ca is not an unlikely event. Therefore, gestating sheep and goats from our in-house dairy herd were fed diets containing either 0.6% or 1.3% Ca in DM during the final 3 weeks antepartum, providing 1.15 and 2.54 times the Swiss daily Ca intake recommendations (Agroscope, 2017). These levels represented a lower Ca intake (<1.1% in DM) to potentially reduce milk fever risk and a higher intake approximating the hypocalcemia risk peak noted by Lean et al. (2006). At the same time, it avoided dietary Ca deficiency in all animals, which would not have benefited our research. Observations continued until 56 days

postpartum, during which animals received standard lactation diets with recommended levels

of Ca according to current Swiss recommendations (Agroscope, 2017). We hypothesized that

goats and sheep fed on the higher Ca diet antepartum (equivalent to 30.5 g/day) would show

signs of hypocalcemia at some point during the observation period, unlike those on the lower

Ca diet. This design also allowed for comparing responses in mineral and bone metabolism to

dietary Ca overload in 2 species with distinct homeostatic strategies of dealing with variations

in Ca supply (Wilkens et al., 2012b).

Material and Methods

Animals and Housing

This study was conducted at the research and laboratory facilities of the Institute of Animal Nutrition and Dietetics, Vetsuisse-Faculty, University of Zurich, Switzerland. A total of 10 and 12 multiparous gestating dairy sheep and goats (East Friesian Dairy Sheep, Saanen goats), respectively, aged between 4 and 6 years were housed in individual pens with concrete floor and wood shavings as bedding material, from 4 weeks before the estimated parturition date. The imbalance in animal numbers between species was due to miscarriages of 2 sheep before the onset of the study. Both species were housed in rooms separate from each other. The goats and sheep were housed individually until their delivery date, afterwards with their respective lambs and kids in lambing/kidding pens. The average body weight just before birth of goats and sheep was 95.2 ± SD1.5 kg and 101 ± SD2.22 kg, respectively, and 78.3 ± SD2.22 kg and 81.0 ± SD2.31 kg directly after birth, respectively.

Diets and Experimental Design

Animals were randomly divided in 2 experimental groups according to their ear tag numbers by a random sequence generator (https://www.random.org/sequences/). The effective sample size for sheep and goats was 5 and 6 per group, respectively. The feeding trial started 3 weeks before the estimated day of parturition. The diets were adapted to the specific Swiss recommendations for energy and nutrient intake during gestation and lactation for sheep and goats (Agroscope, 2017). The 0.6 % in DM Ca groups received a balanced diet provided by grass hay (2nd cut) and a commercial high energy concentrate (UFA 765, Herzogenbuchsee, Switzerland). Additionally, for the 1.3 % in DM Ca groups until parturition, the same concentrate was applied but mixed up daily with 2.5 times (16.8 g Ca /day) the recommended daily Ca supply from $CaCO_3$ (Table 1). From parturition onwards, all animals received lactation feeding to meet current Swiss feeding recommendations which accounted for 17.6 and 19.0 g

Ca per day for sheep and goats, respectively (Table 1) (Agroscope, 2017). Throughout the study,

animals were fed twice per day restrictive amounts of hay (2 kg/animal*d⁻¹ antepartum and 2.4

kg/animal*d⁻¹ postpartum for both species) and concentrate (0.3 kg/animal*d⁻¹ antepartum for

both species and 0.50 and 0.70 kg/animal*d⁻¹ postpartum for sheep and goats, respectively)

adapted to recommendations for antepartum and postpartum feeding (with the exception of Ca

antepartum), and had ad libitum access to drinking water (Table 1). The restrictive feeding

approach allowed for better feed intake control during single housing of animals. The

composition of the commercial concentrate (UFA 765) as communicated by the manufacturer

UFA (Union des Fédération Agricoles), is shown in Supplementary Table 1.

Collection and handling of samples and growth parameters

Colostrum was hand milked from each animal immediately after parturition, and yield ranged

between 0.5 – 1.2 L per animal in both species. After sampling 50 mL for subsequent analysis,

colostrum was distributed equally over nursing bottles and supplied to the neonates.

Sheep and goats were weighed before morning feeding the day before and directly after

parturition. Due to the suckling of the offspring, daily milk yield could not be assessed. The

lambs and kids were weighed directly after parturition and once between 33-59 days

postpartum, to determine the average daily weight gain during the suckling phase. The variation

in the time between 1st and 2nd weighing was because lambs and kids were weaned at different

ages to be subject to a different research purpose. To account for this variation in observational

periods, we included the time interval between the 1st weighing at birth and the 2nd weighing

(the age of each lamb and kid at 2nd weighing) as a continuous variable in the statistical analysis

of weight gain. This approach controls for potential age effects that could bias the estimation of

treatment effects (see the subsection on statistical analysis below).

For this experiment, a closed batch of grass hay (2nd cut) and concentrates was ordered and 3

representative samples each were taken according to published standard methods 1.3

("Regulations of the German Agricultural Society on the Sampling of Feed and Sample Treatment") and 1.5 ("Sampling of Hay") (VDLUFA, 2012) in order to investigate the feed value by chemical analysis. Feed samples were directly subject to chemical analysis after sample collection.

Blood was collected in daily frequency during 28 days before estimated parturition and on days 1, 2, 3, 4, 5, 6, 7, 14, 28, and 56 postpartum, in any case before morning feeding at 8 a.m. in serum monovettes (9 and 6 mL, Vacuette®, Greiner Bio-One Vacuette Schweiz GmbH) and subject to serum collection (1580 x g, 10 min) after allowing coagulation at room temperature. The serum was stored at -80°C until further usage. After the actual parturition, samples of days -21, -4, -3, -2, -1, 0 (parturition), 1, 2, 3, 4, 5, 6, 7, 14, 28, and 56 were reserved for further analysis.

Urine was collected in a beaker upon spontaneous urination, in parallel to the blood sampling antepartum and on days 7 and 14 postpartum, and homogenized to be separated into 2 equal subsamples. The 1st of the subsamples was acidified (12 mol/L HCl, pH 1-2) for later Ca determination, the other was subject to the later determination of creatinine. Both subsamples were stored at -20°C. After parturition, samples of days -21, -14, -7, 0, 7, and 14 were reserved for further usage. Fecal grab samples of each animal were collected in parallel to blood sampling antepartum and days -21, -2, -1, and 0 were subject to further analysis. Samples were stored at -20°C until further usage.

All goats were subject to colonic mucosal biopsies by veterinary personnel on days -21, -7, 0, and 7. An endoscope (PCF-20, Olympus) equipped with biopsy forceps was inserted 40 cm into the large intestine via the anus to collect 4 samples (1.5-2.0 mm diameter) per animal*time^{-1}; 2 of said samples were immediately put into 4% neutral buffered formalin and the remaining 2 samples were snap frozen in liquid nitrogen and stored at -80°C.

Feed analysis

Feed was analyzed according to VDLUFA (2012) with the following methods. For DM according to Method 3.1 ("Moisture, Water") by oven drying at 105°C for 4 h. Net energy for lactation (Method 25.1 "Determination of Gas Production according to the Hohenheim Gas Test") was assessed by quantification of cumulative gas production from rumen fluid over 24h at 39°C and regression estimation of NEL considering cumulative gas production and analyzed crude nutrients and fiber fractions (Society of Nutrition Physiology (GfE), 2008a, b). Crude protein determination (Method 4.1.1 "Determination of Crude Protein") involved the Kjeldahl method of H_2SO_4 extraction in the presence of a catalyst, alkalization with NaOH, titration of NH_3 and multiplying the assessed nitrogen concentration with a factor of 6.25. Starch and ash corrected neutral detergent fiber determination (aNDFom) (Method 6.5.1 "Determination of Neutral-Detergent-Fiber after amylase digestion (aNDF) as well as after amylase digestion and ashing (aNDFom)") involved treatment with heat-stable amylase for 1h, subsequent cooking in neutral detergent solution, and ashing of the solid residue. Ash-corrected acid detergent fiber (ADFom) (Method 6.5.2 "Determination of acid-detergent fiber (ADF) and of acid detergent fiber after ashing (ADFom)") was assessed by 1h cooking in acid detergent solution and ashing of the solid residue. Acid Detergent Lignin (Method 6.5.3 "Determination of Acid Detergent Lignin (ADL)") was quantified by incubating the starch-corrected ADF residue for 3h at room temperature in 72% H_2SO_4 and subsequent ashing of the residue. The determination of the ether extract (Method 5.1.2 "Determination of Crude Fat") involved the pre-extraction in hot HCl and subsequent extraction of the dry residue in petroleum ether. Crude ash (Method 8.1 "Determination of Crude Ash") was measured by dry ashing at 550 °C for 4 h, and HCl-insoluble ash (Method 8.2 "Determination of Hydrochloric Acid-insoluble Ash") added an aliquot of the crude ash residue with boiling HCl. The dietary mineral profile (Ca, P, Mg) (Method 10 "Determination of selected elements in plant material and feedstuffs with ICP-OES") involved the sample ashing in a furnace, extraction with HCl with a deviation from the

protocol by measuring solution not via ICP-OES but by flame atomic absorption spectrometry (flame-AAS) (ContrAA 700, Analytik Jena).

Analysis of blood serum parameters of bone and mineral homeostasis

Blood serum was analyzed for osteocalcin, 1,25-$(OH)_2$ vitamin D (**calcitriol**), bone-specific alkaline phosphatase (**bAP**), crosslinked carboxyterminal teleopeptide of type I collagen (**ICTP**), and crosslaps (**CTX**).

Serum osteocalcin was quantified with a commercial EIA test kit (MikroVue Osteocalcin EIA Kit; Quidel Corporation). Intra- and interassay CVs were 5.00% and 5.75% with a sensitivity of 0.45 ng/mL. Serum calcitriol determination was conducted with an RIA kit (1,25VitD RIA, sheep-anti 1,25VitD antibody; Immunodiagnostics Systems GmbH). 9.0, 9.3 <8pmol/L Intra- and interassay CVs were 9.0% and 9.3% with a sensitivity an 8 pmol/L. Serum bAP was analyzed with an immunoassay via a monoclonal anti-bAP and para-nitrophenylphosphate as a substrate (Alkphase-BTM^{TM}; Metra Biosystems). Intra- and interassay CVs were 0.58 and 0.52. Serum ICTP was measured by a RIA immunoassay (UniQ® ICTP RIA, DiaSource) with an intra- and interassay CV of 5.6 and 10.0 and a sensitivity of 0.6 µg/L. Serum CTX was assayed by a commercial ELISA assay (Serum CrossLaps (CTX-I) ELISA; Immunodiagnostics Systems GmbH) with intra- and interassay CVs of 1.7% and 2.5% at a sensitivity of 0.02 ng/mL.

For CTX, ICTP, and calcitriol, data gaps at -4 days and for CTX additionally for -21 days, as depicted in Figures 5 and 6, respectively, were due to compromised sub-samples. All other sampling decisions were intentional, based on a balance between data relevance, workload, and monetary considerations.

Analysis of Ca and creatinine in biological matrices

Serum Ca was measured via flame-AAS after microwave wet digestion of 2 mL serum per technical replicate in 65% HNO_3 and 30% H_2O_2 (Ethos, MLS GmbH) against a dilution

gradient (0.2, 0.4, 0.6, 0.8, 1.0 mg/L) of a commercial AAS-certified standard (ROTI®Star 1

000 mg/l Ca, in 2 % HNO_3; Roth).

Dried Feces (105°C, 4 h) was milled (0.5 mm) and 500 mg per technical replicate were subject

to Ca analysis as described for serum above.

Colostrum were defatted by centrifugation (4000 x g, 4°C, 30 min), and subject to the same

procedures of sample preparation and Ca analyzes as described for feces and blood serum.

Urine was analyzed for total Ca with the method described for serum and further analyzed for

creatinine by a kinetic color test using a commercial kit (Crea Jaffe DIA00540, Diatools AG)

on the Cobas Mira autoanalyzer with an intra- and interassay CV of 0.83% and 0.85% at a

sensitivity of 18 μmol/L. The urinary Ca excretion was expressed as ratio to urinary creatinine

since the material could not be sampled quantitatively. Creatinine is generally excreted at a

relatively constant rate, making it a useful reference compound to normalize mineral

concentrations for differences in urine dilution caused by hydration status, collection timing or

other individual factors (Sullivan et al., 2012).

Vitamin D receptor immunoreactivity

Goat colonic mucosal biopsies were subject to semi-quantitative immunohistochemical

determination of vitamin D receptor (**VDR**) abundance as originally described by Liesegang et

al. (2008) and Sidler-Lauff et al. (2010). Caprine colonic VDR was labelled after rinsing the

tissue with TBS buffer and incubation for 80 min with a biotinylated rat monoclonal antibody

(9A7γ, ThermoScientific) at a 1:200 dilution in Tris-buffered saline against unspiked Tris-

buffered saline on an adjacent colonic section as negative control and duodenal cross sections

of pigs (obtained from the in-house slaughtering facility of the Vetsuisse-Faculty, University of

Zurich) (aa 89-105) as positive control based on earlier reported identical cross-reactivity

(Milde et al., 1989). For sheep, no suitable antibody was available at the time of study.

Nuclear immunoreactivity was assessed with a binocular microscope (DMLB®, Leica) with a

digital camera (DC480, Leica) (magnification 400x, 5-7 microscopic fields). For each sample,

500 cells of the following cell types were evaluated for nuclear staining intensities: surface

epithelial cells, intermediate glandular/crypt epithelial cells, superficial glandular/crypt

epithelial cells, basal glandular/crypt epithelial cells. Staining intensities of nuclei were scored

0, 0.5, 1, 2, 3 for negative, very weak, weak, intermediate, and strong, which is correlated to

the absence of brown staining (only blue counter staining), light brown, brown, and dark brown,

respectively. From this an immune reactivity score was calculated with the following formula:

Immune Reactivity

$$= 0 \times n(\text{negative}) + 0.25 \times n(\text{very weak}) + 1 \times n(\text{weak})$$

$$+ 4 \times n(\text{intermediate}) + 9 \times n(\text{strong})$$

Where n is the number of cells within 500 cells expressing a respective staining intensity, with

a range of calculated immune reactivity scores between 0 (no specific staining) to 4500 (all 500

cells with strong staining).

<u>Bone mineral density</u>

Bone mineral density (**BMD**) was measured with peripheral quantitative computer tomography

(Stratec XCT 2000 bone scanner, Stratec Medizinaltechnik GmbH) in the middle of the left

metatarsus after length determination with a caliper. Measurements were taken in the mid

diaphysis and cortical BMD was automatically calculated (cortical mode 2, >640 mg/cm^3

cortical bone threshold).

<u>Statistical analysis</u>

The individual animal was the smallest experimental unit. All data analysis was performed with SAS 9.4 (SAS Institute Inc.) applying the procedure MIXED. Repeated measures mixed models were applied to analyze the effects of the fixed variables "diet Ca", "species", "time", and interactions considering the individual animal nested under the respective feeding group as a random factor. Repeated measures mixed modelling involved in any case the mapping of covariance structures appropriate for the respective data structure following guidance by Wang and Goonewardene (2004). Due to the uneven spacing of times of observation for respective datasets, covariance structures like first-order autoregressive (AR(1)) and related structures (e.g., Toeplitz), which require an equal spacing of time, were not considered. The Bayesian Information Criterion (BIC) served as model fit statistic, and we chose the candidate yielding the lowest BIC. The respectively mapped covariance structure for different repeated measures datasets were as follows: compound symmetry (serum Ca, urinary Ca), heterogeneous compound symmetry (BMD), first-order ante-dependent (serum bAP, serum ICTP, serum CTX, serum calcitriol, serum osteocalcin, fecal Ca). Due to the limited availability of suitable antibodies, the VDR protein expression over time was only analyzed in colonic biopsies of goats, therefore no species effect could be estimated.

The colostral Ca concentration of sheep and goats was analyzed with an endpoint mixed model with the fixed effect of "diet Ca", "species", and interaction (diet Ca*species), again including the individual animal nested under the respective feeding group as random effect. Birth weights and daily weight gain of the offspring were also analyzed with an endpoint mixed model accounting for the fixed effects of "diet Ca", "species", "sex" and interactions (diet Ca*species, diet Ca*sex) including the individual animal nested under respective mother as random effect. For the analysis of average daily weight gain, the animals' age at the 2nd weighing was included as a continuous control variable to account for potential age-related bias in estimating average daily weight gain. In case the residuals of respective mixed models missed the precondition of

normality, the analysis was repeated after Log_{10}-transformation of data and displayed as such.

This applied to the following datasets: fecal Ca, urinary Ca, serum bAP, serum ICTP, serum

CTX. Data was presented in any case as least square mean values with Tukey-adjusted 95%

confidence limits (Cl[upper, lower]). $P \leq 0.05$ was defined as threshold for a Type I Statistical

Error.

Earlier datasets were used for prospective power analyses during study planning. Therefore,

effect sizes of ante and postpartum adaption of assessed serum parameters were determined

with the respective datasets of Liesegang et al. (2007). Effects size of VDR immune reactivity

in the goat colon to dietary Ca variation were determined according to the respective dataset of

Sidler-Lauff et al. (2010). The BMD effect sizes of their fluctuation from gestation to lactation

of milk goats and sheep were estimated with the respective data of Liesegang et al. (2007). The

effect sizes and minimum number of replicates required to achieve statistical significance (with

a minimum statistical power of $1-\beta = 0.8$ at $\alpha = 0.05$) (McDonald, 2014) were estimated using

R version 4.4.2 (The R Project for Statistical Computing). This analysis employed the SIMR

package following the approach outlined byGreen and MacLeod (2016), with the modification

of using the previously published datasets mentioned earlier instead of simulated datasets.

Based on these estimations, the anticipated minimum power of ≥ 0.8 was reached in any case

assuming 6 replicates per feeding group, applying the models described above.

Results

Animal health and development

All sheep and goats remained healthy throughout the entire study period with no signs of pathology and hypocalcemia, based on continuous veterinary surveillance. Sheep showed 5 cases of triplets, 4 cases of twins and 1 single birth (n = 24 lambs in total) whereas most goats gave birth to twins with 1 case of triplets and 2 cases of single births (n = 23 kids in total). The sex ratio of sheep and goat offspring was (m/f) 0.71 and 1.30, respectively.

All lambs and kids were born with species-dependent birth weights (4.62 Cl[4.22, 5.01] vs. 5.15 Cl[4.80, 5.51] kg for kits vs lambs; $P = 0.06$) irrespective of sex (4.71 Cl[4.35, 5.07] vs. 5.06 Cl[4.71, 5.41] kg for female vs. male; $P = 0.16$) and unaffected by the mothers' dietary Ca supply antepartum (4.89 Cl[4.53, 5.23] vs. 4.88 Cl[4.54, 5.22] kg at 0.6% vs. 1.3% in DM, respectively; $P = 0.94$) (Figure 1A). However, lambs and kids of mothers fed high Ca antepartum showed significantly lower daily weight gain from milk intake than animals fed low Ca (0.18 Cl[0.17, 0.20] vs. 0.22 Cl[0.20, 0.24] kg/d at 1.3% vs. 0.6% in DM, respectively; $P = 0.02$) irrespective of sex and species, which expressed distinct effects on the animals' growth performance (0.19 Cl[0.17, 0.21 vs. 0.21 Cl[0.19, 0.23] kg/d for female vs. male; $P = 0.06$; 0.25 [0.23, 0.27] vs. 0.15 [0.13, 0.17] kg/d for kids vs. lambs; $P < 0.0001$) (Figure 1B). Supplementary Table 2 highlights all LSmeans and associated Tukey-adjusted 95% confidence limits of birth weights and daily weight gain of offspring as related to species, sex, and dietary Ca supply of the mothers antepartum.

Quantitative markers of calcium metabolism

Analyzed fecal Ca concentrations during the observation period ranged between 1.46 and 5.19 % in DM in sheep as well as 1.17 and 14.9 % in DM in goats whereas the ranges for urinary

Ca excretion were 0.005 and 0.41 mmol/mmol creatinine in sheep as well as 0.004 and 1.55

mmol/mmol creatinine in goats. Fecal and urinary data had to be log_{10}-transformed to meet the

precondition of normality, accounting for fecal Ca ranges of 0.16 and 0.71 log_{10}(% in DM) for

sheep as well as 0.07 and 1.17 log_{10}(% in DM) for goats. The respective log-transformed urinary

Ca excretion ranges were -2.31 and -0.38 log_{10}(mmol/mmol creatinine) in sheep as well as

-2.42 and -0.19 log_{10}(mmol/mmol creatinine) in goats. Supplementary Table 3 highlights all

LSmeans and associated Tukey-adjusted 95% confidence limits of fecal Ca concentrations and

urinary Ca excretion, expressed in the original measurement scale as related to species, time,

and dietary Ca supply antepartum.

Higher supply with dietary Ca antepartum significantly increased fecal log_{10}-Ca concentrations

until parturition by 1.43fold vs. 2.87fold in sheep vs. goats and in any case significantly above

the control level (0.21 Cl[0.07, 0.34] vs. 0.62 [0.48, 0.75] and 0.35 Cl[0.22, 0.48] vs. 0.66

Cl[0.53, 0.79] log_{10}(% in fecal DM) in 0.6% vs. 1.3% in DM of sheep and goats, respectively;

$P < 0.0001$) (Figure 2A). On average, sheep showed lower fecal Ca than goats (0.41 Cl[0.36,

0.46] vs. 0.47 Cl[0.43, 0.52] log10(% in fecal DM) for sheep vs. goats; $P = 0.05$) but the

response patterns over time were different. Sheep in the low Ca group slightly declined fecal

Ca until parturition to 0.64fold of the level at -21 days antepartum, while sheep in the in the

high Ca group responded opposite to this as mentioned before (1.43fold increase until

parturition). In contrast, both control and high Ca goats increased the levels until parturition but

in different magnitudes (1.75fold vs. 2.87fold for low vs. high Ca). This difference in time

patterns between species was reflected by a strong statistical interaction between species and

time ($P < 0.0001$).

In contrast, the feeding regime had no effect on the urinary Ca excretion (-1.04 Cl[-1.22, -0.86]

vs. -0.96 Cl[-1.14, -0.79] log10(mmol/mmol creatinine) at 0.6% vs. 1.3% in DM; $P = 0.56$)

(Figure 2B). Values fluctuated over time but a clear trend in adaptation of urinary losses was

not evident ($P = 0.61$). The only significant effect was in comparison between sheep and goats

irresponsive of treatment or time, with goats excreting on average 3.26fold the levels of sheep

(-1.26 Cl[-1.44, -1.07]vs. -0.75 Cl[-0.91, -0.58] $\log_{10}$(mmol/mmol creatinine) in sheep vs.

goats; $P = 0.0003$).

Figure 3 shows the response of serum Ca and colostral Ca to varying Ca feeding antepartum.

Serum Ca was not affected by the antepartum feeding (2.47 Cl[2.35, 2.59] vs. 2.49 Cl[2.37,

2.61] mmol/L at 0.6% vs. 1.3% in DM; $P = 0.80$) and gradually and significantly ($P < 0.0001$)

decreased independent of animal species towards the initial 4 days postpartum to gradually

increase to previous levels during the further course of the observation period (Figure 3A). In

any case, sheep showed on average 1.16fold higher serum Ca levels than goats (2.29 Cl[2.17,

2.40] vs. 2.67 Cl[2.54, 2.79] mmol/L in goats vs. sheep; $P = 0.0003$).

The colostral Ca concentration (Figure 3B) was 2.35fold higher in sheep than in goats (1.44

Cl[0.69, 2.19] vs. 3.38 Cl[2.56, 4.20] g/L in goats vs. sheep; $P = 0.002$) but was not affected by

the antepartum Ca feeding regime (2.36 Cl[1.58, 3.14] vs. 2.46 Cl[1.67, 3.24] g/L at 0.6% vs

1.3% Ca in DM; $P = 0.86$) (Figure 3B).

Supplementary Table 4 highlights all LSmeans and associated Tukey-adjusted 95% confidence

limits of serum Ca and colostral Ca as related to species, time, and dietary Ca supply

antepartum.

Figure 4 highlights the BMD of dairy sheep and dairy goats from -14 day antepartum to 56 days

postpartum, as affected by Ca feeding during the last 3 weeks antepartum, time and animal

species. Higher Ca feeding antepartum did not affect BMD (868 Cl[845, 891] vs. 877 Cl[854,

900] mg/cm^3 at 0.6% vs. 1.3% in DM; $P = 0.63$), which significantly declined towards

parturition from 932 Cl[904, 960] to 864 Cl[807, 922] mg/cm^3 in sheep and 845 Cl[817, 873]

to 782 Cl[738, 826] mg/cm^3 in goats, to gradually increase postpartum until day 56 of lactation

where it roughly approached in both species the initial level at the beginning of the observation

(Sheep: 944 Cl[920, 967] mg/cm^3, Goats: 846 Cl[823, 868] mg/cm^3 $P < 0.0001$). On average,

sheep showed 1.08fold higher BMD than goats (839 Cl[808, 853] vs. 913 Cl[888, 938] mg/cm^3

in goats vs. sheep; $P < 0.0001$).

Supplementary Table 5 highlights all LSmeans and associated Tukey-adjusted 95% confidence

limits of BMD as related to species, time, and dietary Ca supply antepartum.

<u>Serum calcitriol and markers of bone formation</u>

Serum calcitriol, bAP and osteocalcin of sheep and goats over time in response to varying Ca

feeding antepartum are shown in Figure 5. Serum calcitriol was 1.98fold higher in goats than

in sheep (70.2 Cl[56.8, 83.6] vs. 139 Cl[128, 149] pmol/L in sheep vs. goats; $P < 0.0001$) but

not different between antepartum feeding groups (103 Cl[90.1, 117] vs. 105 Cl[92.3, 119]

pmol/L at 0.6% vs. 1.3% in DM; $P = 0.75$) (Figure 5A). Serum levels significantly changed

over time ($P < 0.0001$) which interacted with species ($P < 0.0001$) as follows. In sheep, serum

levels reduced towards parturition and fluctuated throughout, whereas in goats serum levels

increased until 4 days postpartum to turnaround and approach initial levels until 56 days

postpartum.

Serum activity of bAP ranged between 38 and 251 U/L in sheep as well as 21 and 117 U/L in

goats. After Log$_{10}$-transformation to meet the precondition of normality, these ranges were 1.58

and 2.40 Log$_{10}$(U/L) in sheep as well as 1.32 and 2.07 Log$_{10}$(U/L) in goats. Feeding group (1.76

Cl[1.67, 1.85] vs. 1.80 Cl[1.70, 1.90] log$_{10}$(U/L) at 0.6% vs. 1.3% in DM; $P = 0.59$) expressed

no significant effect on bAP (Figure 5B). This was also true for time, though control goats

showed a transient numerical drop by ~22% of serum activity around parturition ($P = 0.12$).

Sheep showed on average 1.25fold higher serum activities than goats (1.58 Cl[1.49,, 1.68] vs.

1.97 Cl[1.888, 2.07] log$_{10}$(U/L) in goats vs. sheep; $P < 0.0001$).

Serum osteocalcin remained stable and comparable between species from -5 days antepartum

to 5 days and 7 days postpartum, respectively, after which it gradually increased on average by

2.27fold in sheep and 3.23fold in goats until 56 days postpartum ($P < 0.0001$) (Figure 5C). This

difference in magnitude was significant between species ($P = 0.007$). Most notably, this increase

in serum osteocalcin was significantly higher in those sheep and especially goats who received

higher dietary Ca during the last 3 weeks antepartum (29.9 Cl[5.62, 54.1] vs. 39.6 Cl[15.3, 63.8]

and 42.2 Cl[17.8, 66.6] vs. 80.5 [58.4, 103] mmol/L at 0.6% vs. 1.3% in DM in sheep and goats,

respectively; $P = 0.02$).

Supplementary Table 6 highlights all LSmeans and associated Tukey-adjusted 95% confidence

limits of serum calcitriol, bAP, and osteocalcin, expressed in the original measurement scale

also in the case of bAP, as related to species, time, and dietary Ca supply antepartum.

<u>Serum markers of bone resorption</u>

Serum concentrations of ICTP ranged between 4.87 and 90.2 µg/L in sheep as well as 5.75 and

57.6 µg/L in goats, while CTX ranged 0.50 and 10.3 ng/mL in sheep as well as 0.35 and 8.72

ng/mL in goats. After Log_{10}-transformation to meet the precondition of normality, these ranges

were for ICTP 0.69 and 1.95 log_{10}(µg/L) as well as -0.30 and 1.01 log_{10}(µg/L) and for CTX

0.76 and 1.76 log_{10}(ng/mL) as well as -0.46 and 0.94 log_{10}(ng/mL). Supplementary Table 7

highlights all LSmeans and associated Tukey-adjusted 95% confidence limits of serum ICTP

and CTX, expressed in the original measurement scale, as related to species, time, and dietary

Ca supply antepartum. Serum ICTP showed no marked changes in both species until parturition,

afterwards it significantly increased over the initial 2 days postpartum (1.49fold vs. 1.60fold in

sheep vs. goats) and held a plateau until 7 days postpartum from which on the levels gradually

reduced over the following weeks to approach initial values at 56 days postpartum ($P < 0.0001$)

(Figure 6A). Average serum levels were significantly higher for sheep than goats (1.19 Cl[1.14,

1.23] vs. 1.30 Cl[1.26, 1.35] $\log_{10}(\mu g/L)$ in goats vs. sheep; $P < 0.0001$) with a significant

interaction between species and time reflecting fluctuations in differences between species over

given times ($P = 0.001$). The feeding antepartum did not express any significant effect on serum

ICTP levels of dairy sheep and goats (1.24 Cl[1.20, 1.29] vs. 1.25 [1.20, 1.29] $\log_{10}(\mu g/L)$ at

1.3% vs. 0.6% in DM; $P = 0.98$).

Sheep showed markedly higher serum CTX levels during this study than goats ($P < 0.0001$)

(Figure 6 B). In sheep serum levels significantly increased from -3 days antepartum until 2 days

postpartum and held a plateau until day 7 postpartum with a subsequent decline during the

following weeks approaching initial values at day 57 postpartum. In contrast goats showed not

much response until parturition after which it peaked at 2 days postpartum and fluctuated

around that plateau for the next 14 days to decline until day 56 with the high Ca group showing

the steepest decline. These responses over time were significant ($P < 0.0001$) but did not

statistically interact with other variables like diet Ca or species. Neither on average (0.35

Cl[0.27, 0.43] vs. 0.38 [0.30, 0.46] $\log_{10}(\mu g/L)$ at 1.3% vs. 0.6% in DM; $P = 0.59$) nor over

time (Diet Ca*Time: $P = 0.64$) did the feeding antepartum affect serum CTX levels in a

significant manner.

Caprine colonic abundance of mucosal vitamin D receptor

Figure 7 highlights the statistical evaluation of VDR immune reactivity in caprine colonic

mucosa in goats fed 0.6% in DM and 1.3% in DM dietary Ca antepartum. Figure 8 presents

exemplary immunohistochemical images from both groups at parturition. Supplementary Table

8 highlights all LSmeans and associated Tukey-adjusted 95% confidence limits of VDR

immune reactivity, as related to time and dietary Ca supply antepartum.

Vitamin D receptor abundance in caprine colonic mucosa was not significantly affected by diet

Ca antepartum (2268 Cl[1856, 2680] vs. 2669 Cl[2214, 3125] immune reactivity at 1.3% vs.

0.6% in DM; $P = 0.19$). In both groups, the scores showed a curvilinear trend over time which

could not be statistically confirmed ($P = 0.1$).

Discussion

An earlier meta-analysis concluded that the risk for hypocalcemia in dairy cows significantly increases if dietary Ca during the antepartum phase is between 1.1-1.5% in DM (Lean et al., 2006). This hypothesis has yet not been tested in dairy sheep and goats, therefore the present study investigated whether Ca fed at 1.3% in DM (compared to 0.6% in DM) during the last 3 weeks antepartum increases the incidence of hypocalcemia antepartum and/or postpartum by dysregulating Ca homeostasis.

<u>Effects on quantitative Ca homeostasis and the incidence of hypocalcemia</u>

Sheep and goats showed in any case significant differences in the expression of observed measures of quantitative Ca homeostasis (fecal Ca, urinary Ca excretion, colostrum Ca, serum Ca, BMD). Differences in magnitude of these parameters between both species have also been observed in several earlier comparative studies. Our observation of higher values in sheep with respect to urinary Ca excretion and colostral Ca and BMD are in line with earlier studies (Liesegang et al., 2006, 2007, Liesegang, 2008). We made the same observation with serum Ca, which agrees to Kohler et al. (2013) and Liesegang (2008), though Wilkens et al. (2014) observed the opposite with goats expressing higher values than sheep. The reason for this discrepancy is unclear at the moment. Interestingly, both species exhibited a drop in serum Ca at 3 and 4 days postpartum, respectively. Similar response patterns have recently been described in dairy cows, particularly for Zn balance and serum levels (Daniel et al., 2023). It is possible that the serum Ca pattern observed in the present study mirrors a transient drop in Ca retention at the onset of lactation, comparable to the patterns observed for Zn in dairy cows. However, due to lack of data this interpretation remains speculative for the present experimental conditions and requires confirmation in future studies.

The feeding showed mostly no or just numerical effects on quantitative measures. Increasing dietary Ca feeding during the antepartum phase significantly increased fecal Ca concentration

by at least 2fold during the last 3 weeks but did not significantly affect serum and colostral Ca concentration and urinary Ca excretion in both species at any given point during the study. Most notably, no signs of hypocalcemia or any pathology were noted antepartum or postpartum, respectively, although according to Oetzel (1988), dairy sheep and goats are expected to develop milk fever symptoms comparable to cows but at 1-3 weeks postpartum. However, since 1988, advancements in milk yields and the associated calcium losses suggest that feeding regimes promoting milk fever may now be more likely to trigger the condition and potentially earlier than originally described by Oetzel. Literature on the effects of Ca oversupply in sheep and goats is scarce, particularly with respect to the antepartum phase. However, the only study on sheep by Braithwaite (1979)agrees with our observations. The colleague studied gradual differences in dietary Ca intake in adult sheep and observed increasing fecal Ca losses. Fractional Ca absorption (% of intake) was at 31.8 % at the lowest Ca supply (0.22 % in DM) and dropped to a plateau around ~9-10% with gradually increasing diet Ca to 0.6, 1.2, and 2.4% in dietary DM, respectively. In parallel, bone Ca resorption dropped inversely to the increase in Ca intake and inevitable unregulated absorption of surplus Ca was increasingly redirected into the gastrointestinal tract. Given the response of our sheep and goats in the present study, it may be concluded that for our animals and under the present study conditions, we can reject our initial hypothesis of increased milk fever risk when feeding higher Ca in the range proposed by Lean et al. (2006) for dairy cows. It remains to be tested what the outcome would be when feeding a different Ca source, for example a source with a higher water solubility like $CaCl_2$ (Goff and Horst, 1993). Neither sheep nor goats showed any relevant response in urinary Ca excretion in response to the feeding regime over time, confirming once more results from earlier studies suggesting livestock ruminants do regulate Ca primarily via the gastrointestinal tract (Herm et al., 2015, Wild et al., 2021). We assume that the animals were able to dampen uptake from the gastrointestinal tract and, in parallel, effectively rerouted unavoidable unregulated Ca uptake towards excretory pathways, as described by Braithwaite (1979). Future studies must

identify the precise physiological mechanisms behind this effective compensation of dietary Ca

overload. See in this context also our discussion below, on potentially confounding factors

associated with our study. Furthermore, the functional basis for the differences in fecal Ca

accumulation between the 2 species remains to be investigated. Initially, goats exhibited lower

fecal Ca levels at -21 days antepartum, but this trend reversed until parturition. The underlying

causes are currently speculative and may involve differences in dry matter digestibility or the

regulation of Ca absorption at the gut barrier. Future studies should explore the effects of high

Ca feeding on gastrointestinal pH, as well as the digestive and absorptive capacities for dry

matter and minerals, to provide a more comprehensive understanding of this phenomenon.

<u>Effects on serum calcitriol and markers of bone formation and resorption</u>

We observed a core set of serum bone formation and resorption markers, which have been

earlier established in ruminants from gestation into lactation (Liesegang et al., 2000, Liesegang

et al., 2007). Observed measures of bone formation and resorption showed changes over time,

that have earlier been attributed to the species-dependent adaption towards parturition and the

onset and further course of lactation (Liesegang et al., 2006, 2007, Liesegang, 2008, Kohler et

al., 2013, Wilkens et al., 2014). However, none of the observed parameters showed a significant

response to the feeding treatment during the final weeks of the antepartum and the majority of

the postpartum period. Consequently, the BMD was not affected and suggests the feeding

intervention with 1.3% in DM Ca in the diet caused no obvious dysregulation of Ca and bone

homeostasis in either species as compared to the lower Ca feeding. However, serum osteocalcin

showed a significant gradual increase in concentration from d7 to d56 of lactation in animals

receiving the high dietary Ca antepartum. This response was true for both species but most

pronounced in the dairy goats. Serum osteocalcin is considered a marker of bone formation,

with increasing levels pointing towards rising embedding of Ca in the skeleton (Liesegang et

al., 2000). This observation was puzzling given the lack of response of calcitriol as well as markers of bone formation (calcitriol, bAP) and resorption (ICTP, CTX)) and especially since the effect occurred 8d after the suspension of the high Ca feeding. To the best of our knowledge such a response pattern has not yet been reported for serum osteocalcin and it raises questions on its physiological purpose. The rise in serum osteocalcin was not associated with changes in BMD (or mineral concentration; data not shown) nor did other markers of bone formation show an orchestrated response that would suggest any feeding-associated changes in bone metabolism. Therefore, it may be necessary to have a look at osteocalcin from a different perspective. Biomedical research efforts in recent years suggest a hormonal role of uncarboxylated osteocalcin beyond bone metabolism, involving among others the regulation of systemic energy metabolism (Lee et al., 2007), muscle energy metabolism (Mera et al., 2016), and brain development (Obri et al., 2018). At this point, the meaning of these results is up to speculation and more targeted research on the matter is urgently needed.

Effects on offspring development

High Ca treatment of mothers antepartum was associated with decreased average daily weight gain of suckling lambs and kids. Since the offspring showed no differences in birth weight that were associated to the feeding of the mothers antepartum, we rule out quantitative effects on placental nutrient transfer. The most plausible explanation for reduced offspring growth during the suckling phase is an impairment of milk yield and/or quality. Feed intake was not affected in the present study and all animals consumed the necessary dietary dry matter according to recommendations. At the same time, we noted differences in serum osteocalcin in the higher Ca groups, pointing towards a distinct physiological conditioning. This likely has affected the quantity and/or quality of secreted milk and therefore the growth response of their offspring, but this needs yet to be confirmed. To the best of our knowledge there is yet no data available

in dairy sheep and goats that allows us a more informed discussion on this matter and future studies must put a focus on the potential effects of the present experimental conditions on milk yield and quality as well as lactation physiology. In addition, a further hypothesis may deserve some attention in future studies. High Ca feeding during the last 3 weeks antepartum and the associated physiological adaption of the mothers may have promoted fetal programming in the offspring that impaired their digestive capacity for Ca (and maybe other nutrients) postpartum and growth response. The possibility of mineral nutrition during gestation to induce fetal programming in ruminants either for the benefit or disadvantage of the offspring has recently been reviewed by Anas et al. (2023) for cattle, highlighting a multitude of already established interactions as well as future routes for research on the matter. In this context, an earlier study in rats tested the effects of varying dietary Ca supply during gestation including higher supply at 1.2% in DM (Li et al., 2018). The colleagues observed abnormal expression of hepatic and adipose genes in the offspring as well as dyslipidemia and hepatic lipid accumulation in response. Unfortunately, no data on Ca homeostasis in rat mothers and their offspring was obtained. Nevertheless, it may be argued that high supply of a mineral over several weeks may induce epigenetic changes in the mother, which could be inherited to the offspring. It may therefore be interesting to have a closer look on the epigenetic consequences of high dietary Ca as well as opposite feeding regimes during gestation and their potential to imprint Ca homeostatic regulation and development of the offspring postpartum.

The geographical region and other potentially confounding factors

Depending on the geographic location, Ca levels in the environment are quite variable. In Switzerland, animals are often confronted with high levels of native Ca in diets, originating from the Ca-rich geological formations, which shape the chemical composition of soil in certain areas. For example an earlier survey of Schlegel et al. (2016) found a mean Ca concentration

of 7 g/kg DM in 236 samples of mixed grassland from Posieux, Switzerland, and our own experience with grass hay qualities delivered to our animal facility in the Zurich area are in line with this observation. This leads to quite relevant intake levels of native Ca from feed particularly when grass-based roughage is fed, even when mineral Ca supplements are not applied. Since the organism attempts to maintain Ca homeostasis by fine-tuning absorption, losses, and mobilization of skeletal reserves according to the environmental conditions (Martín-Tereso and Martens, 2014, Wilkens and Muscher-Banse, 2020), it may be argued that sheep and goats raised predominantly on Ca-rich roughage may be better suited to deal with high dietary Ca challenge. However, it may also suggest that during the feeding of high dietary Ca due to active supplementation of inorganic Ca to such raw materials, a metabolic imprinting as hypothesized earlier may more likely promote the observed adverse effects on daily weight gain of suckling lambs and goat kids. Therefore, the present results must be considered with caution when it comes to the general translation into dairy sheep and goat feeding systems. More data is urgently needed also from regions with less Ca in the soil.

This study is among the earliest to compare high Ca responses between sheep and goats. While many biomarkers in our findings align with previous research on bone metabolism and calcium homeostasis principles within and across these species, our study is too preliminary to draw definitive conclusions regarding species-specific effects under the current treatment conditions. Given that "species" is an intrinsic characteristic, it cannot be effectively randomized within a single study limited to 1 herd per species. However, our findings represent an initial step toward further research in this area, underscoring the need for not only replication of these experimental conditions but also replication across multiple herds to strengthen the reliability of observed responses.

The current Swiss feeding recommendations for sheep and goats suggest comparable dietary Ca levels during the last 3 weeks antepartum (Agroscope, 2017). Based on these guidelines, our

study applied the same diet to both species. While these recommendations likely exceed net requirements due to built-in safety margins, it remains an open question whether current genotypes in both species respond similarly to a standardized diet. To enhance future research, it would be beneficial to revisit and, if needed, revise the calcium feeding guidelines for dairy sheep and goats individually, ensuring they reflect species-specific physiological needs more accurately.

Neither sex of offspring nor litter size were balanced in the present study, therefore it may be argued whether this imbalance may have biased the results on daily weight gain during the suckling phase. However, sex was included as a variable in the statistical model and multiple births outweighed single births in both species (only 1 single birth in sheep and 2 in goats). In addition, the random baseline of each animal nested under the respective mother was added as a random factor. We are therefore confident the results truly reflect a physiological response from the varying antepartum Ca feeding of the mothers. Nevertheless, this is the 1st experiment reporting this phenomenon and further replication of the study ideally in other labs is needed to further confirm our findings.

Conclusion

The present study did not find any signs of hypocalcemia in sheep and goats challenged with higher Ca during the last 21 days antepartum. However, serum osteocalcin showed increasing concentrations in lactating sheep and goats fed higher Ca antepartum compared to the respective lower Ca group. Interestingly, although no obvious negative effects of the feeding regime could be observed for the mothers, the offspring of sheep and goats fed higher Ca antepartum did grow slower over time compared to the lower Ca groups, despite comparable birth weights. This likely has originated from effects on the physiological status of the mothers, thereby affecting milk yield and quality. Nevertheless, the additional hypothesis of a potential adverse

effect of high dietary Ca antepartum on the digestive capacity of the offspring via epigenetic

imprinting may be an exciting avenue for future research. Our animals were reared in a Ca-rich

environment since their birth; therefore, we cannot rule out that the present findings are

exclusive for this situation. Follow-up studies are urgently needed and should also comprise

observations in production regions with less Ca in the soils.

Notes

Raw datasets can be requested via E-mail to the corresponding author.

This study received no external funding.

We thank Dr. med. vet. Sybil Lüthi for excellent technical assistance in sample and data

collection. We further thank the technical staff in the animal and laboratory facilities of the

Institute of Animal Nutrition and Dietetics, Vetsuisse-Faculty, University of Zurich,

Switzerland.

All procedures used in the experiment involving animals were in accordance with the Swiss

guidelines for animal welfare at the time of study and were registered and approved by the

Cantonal Veterinary Office (ZH 9/2006).

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

References

Agroscope. 2017. Fütterungsempfehlungen für Wiederkäuer (Grünes Buch). Vol. https://www.agroscope.admin.ch/agroscope/de/home/services/dienste/futtermittel/fuett erungsempfehlungen-wiederkaeuer.html. Agroscope.

Anas, M., W. J. S. Diniz, A. C. B. Menezes, L. P. Reynolds, J. S. Caton, C. R. Dahlen, and A. K. Ward. 2023. Maternal mineral nutrition regulates fetal genomic programming in cattle: A review. Metabolites 13:593.

Braithwaite, G. D. 1979. The effect of dietary intake of calcium and phosphorus on their absorption and retention by mature Ca-replete sheep. J. Agric. Sci. 92:337-342.

Daniel, J. B., D. Brugger, S. van der Drift, D. van der Merwe, N. Kendall, W. Windisch, J. Doelman, and J. Martín-Tereso. 2023. Zinc, copper, and manganese homeostasis and potential trace metal accumulation in dairy cows: longitudinal study from late lactation to subsequent mid-lactation. The Journal of Nutrition 153(4):1008-1018.

Goff, J. P. 2008. The monitoring, prevention, and treatment of milk fever and subclinical hypocalcemia in dairy cows. The Veterinary Journal 176(1):50-57.

Goff, J. P. and R. L. Horst. 1993. Oral administration of calcium salts for the treatment of hypcalcemia in cattle. J. Dairy Sci. 76(1):101-108.

Green, P. and C. J. MacLeod. 2016. SIMR: an R package for power analysis of generalized linear mixed models by simulation. Methods in Ecology and Evolution 7:493-498.

Herm, G., A. S. Muscher-Banse, G. Breves, B. Schröder, and M. R. Wilkens. 2015. Renal mechanisms of calcium homeostasis in sheep and goats. J. Anim. Sci. 93:1608-1621.

Hernández-Castellano, L. E., L. L. Hernandez, and R. M. Bruckmaier. 2020. Review: Endocrine pathways to regulate calcium homeostasis around parturition and the prevention of hypocalcemia in perparturient dairy cows. Animal 14(2):330-338.

Hoenderop, J. G., O. Dardenne, M. van Abel, A. W. van der Kemp, C. H. van Os, R. St. Arnaud,

and R. J. Bindels. 2002. Modulation of renal Ca^{2+} transport protein genes by dietary

Ca^{2+} and 1,25-dihydroxyvitamin D_3 in 25-hydroxyvitamin D_3-1alpha-hydroxylase

knockout mice. FASEB J. 16:1398-1406.

Ko, S. H., G. S. Lee, T. T. Vo, E. M. Jung, K. C. Choi, K. W. Cheung, J. W. Kim, J. G. Park, G.

T. Oh, and E. B. Jeung. 2009. Dietary calcium and 1,25-dihydroxyvitamin D3 regulate

transcription of calcium transporter genes in calbindin-D9k knockout mice. Journal of

Reproduction and Development 55:137-142.

Kohler, M., F. Leiber, H. Willems, L. Merbold, and A. Liesegang. 2013. Influence of altitude

on vitamin D and bone metabolism of lactating sheep and goats. J. Anim. Sci. 91:5259-

5268.

Lean, I. J., P. J. DeGaris, D. M. McNeil, and E. Block. 2006. Hypocalcemia in dairy cows:

meta-analysis and dietary cation anion difference theory revisited. J. Dairy Sci.

89(2):669-684.

Lee, N. K., H. Sowa, E. Hinoi, M. Ferron, J. D. Ahn, C. Confavreux, R. Dacquin, P. J. Mee, M.

D. McKee, D. Y. Jung, Z. Zhang, J. K. Kim, F. Mauvais-Jarvis, P. Ducy, and G. Karsenty.

2007. Endocrine regulation of energy metabolism by the skeleton. Cell 130(3):456-469.

Li, P., X. Chang, X. Fan, C. Fan, T. Tang, R. Wang, and K. Qi. 2018. Dietary calcium status

during maternal pregnancy and lactation affects lipid metabolism in mouse offspring.

Scientific Reports 8:16542.

Liesegang, A. 2008. Influence of anionic salts on bone metabolism in periparturient dairy goats

and sheep. J. Dairy Sci. 91:2449-2460.

Liesegang, A., J. Risteli, and M. Wanner. 2006. The effects of first gestation and lactation on

bone metabolism in dairy goats and milk sheep. Bone 38:794-802.

Liesegang, A., J. Risteli, and M. Wanner. 2007. Bone metabolism of milk goats and sheep

during second pregnancy and lactation in comparison to first lactation. J. Anim. Physiol.

Anim. Nutr. 91:217-225.

Liesegang, A., M.-L. Sassi, J. Risteli, R. Eicher, M. Wanner, and J.-L. Riond. 2000. Biochemical

markers of bone formation and resoprtion around parturition and during lactation in

dairy cows with high and low standard milk yields. J. Dairy Sci. 83(8):1773-1781.

Liesegang, A., K. Singer, and A. Boos. 2008. Vitamin D receptor amounts across different

segments of the gastrointestinal tract in Brown Swiss and Hosltein Friesian cows of

different age. J. Anim. Physiol. Anim. Nutr. 92:316-323.

Martín-Tereso, J. and H. Martens. 2014. Calcium and magnesium physiology and nutrition in

relation to the prevention of milk fever and tetany (dietary management of

macrominerals in preventing disease). Veterinary Clinics of North America: Food

Animal Practice 30(3):643-670.

McDonald, J. H. 2014. Power Analysis. Pages 40-44 in Handbook of Biological Statistics, 3rd

ed. J. H. McDonald, ed. Sparky House Publishing, Baltimore, MD, USA.

Mera, P., K. Laue, M. Ferron, C. Confavreux, J. Wei, M. Galán-Díez, A. Lacampagne, S. J.

Mitchell, J. A. Mattison, Y. Chen, J. Bacchetta, P. Szulc, R. N. Kitsis, R. de Cabo, R. A.

Friedman, C. Torsitano, T. E. McGraw, M. Puchowicz, I. Kurland, and G. Karsenty.

2016. Osteocalcin signaling in myofibers is necessary and sufficient for optimum

adaptation to exercise. Cell. Metab. 23(6):1078-1092.

Milde, P., J. Merke, E. Ritz, M. R. Haussler, and E. W. Rauterberg. 1989. Immunohistochemical

detection of 1,25-dihydroxyvitamin D_3 receptors and estrogen receptors by monoclonal

antibodies: comparison of four immunoperoxidase methods. The Journal of

Histochemistry and Cytochemistry 37:1609-1617.

Obri, A., L. Khrimian, G. Karsenty, and F. Oury. 2018. Osteocalcin in the brain: from embryonic

development to age-related decline in cognition. Nat. Rev. Endocrinol. 14(3):174-182.

Oetzel, G. R. 1988. Parturient paresis and hypocalcemia in ruminant livestock. Vet. Clin. North

Am. Food Anim. Pract. 4:351-364.

Oetzel, G. R. 1991. Meta-analysis of nutritional risk factors for milk fever in dairy cattle. J.

Dairy Sci. 74:3900-3912.

Schlegel, P., U. Wyss, Y. Arrigo, and H. D. Hess. 2016. Mineral concentrations of fresh herbage

from mixed grassland as influenced by botanical composition, harvest time and growth

stage. Anim. Feed Sci. Technol. 219:226-233.

Sidler-Lauff, K., A. Boos, M. Kraenzlin, and A. Liesegang. 2010. Influence of different calcium

supplies and a single vitamin D injection on vitamin D receptor- and calbindin D9k-

immunoreactivities in the gastroinestinal tract of goat kids. J. Anim. Sci. 88(11):3598-

3610.

Society of Nutrition Physiology (GfE). 2008a. Communications of the Commitee for

Requirement Standards of the Society of Nutrition Physiology: New equations for

predicting metabolisable energy of grass and maize products for ruminants. Proc. Soc.

Nutr. Physiol. 17:191-198.

Society of Nutrition Physiology (GfE). 2008b. Prediction of metabolisable energy of compound

feeds for ruminants. Proc. Soc. Nutr. Physiol. 17:199-204.

Sullivan, K., S. Freeman, E. Van Heugten, K. Ange-van Heugten, B. Wolfe, and M. H. Poore.

2012. Impact of two types of complete pelleted, wild ungulate feeds and two pelleted

feed to hay ratios on the development of urolithogenic compounds in meat goats as a

model for girrafes. J. Anim. Physiol. Anim. Nutr. 97(3):566-576.

van Cromphaut, S. J., K. Rummens, I. Stockmans, E. van Herck, F. A. Dijcks, A. G. Ederveen,

P. Carmeliet, R. Bouilon, and G. Carmeliet. 2003. Intestinal calcium transporter genes

are upregulated by estrogens and the reproductive cycle through vitamin D receptor-

independent mechanisms. J. Bone Miner. Res. 18:1725-1736.

VDLUFA. 2012. VDLUFA-methods book III: The chemical analysis of feedstuffs. VDLUFA-

Verlag, Darmstadt, Germany.

Wang, Z. and L. A. Goonewardene. 2004. The use of MIXED models in the analysis of animal

experiments with repeated measures data. Can. J. Anim. Sci. 84(1):1-11.

Wild, K. J., W. Siegert, W. M. Windisch, K.-H. Südekum, and M. Rodehutscord. 2021. Meta-

analysis-based estimates of efficiency of calcium utilization by ruminants. Animal

15(8):100315.

Wilkens, M., J. Richter, D. R. Fraser, A. Liesegang, G. Breves, and B. Schröder. 2012a. In

contrast to sheep, goats adapt to dietary calcium restriction by increasing intestinal

absorption of calcium. Comparative Biochemistry and Physiology Part A, Molecular &

Integrative Physiology 163(3-4):396-406.

Wilkens, M. R., A. Liesegang, J. Richter, D. R. Fraser, G. Breves, and B. Schröder. 2014.

Differences in peripartal plasma parameters related to calcium homeostasis of dairy

sheep and goats in comparison to cows. J. Dairy Res. 81(3):325-332.

Wilkens, M. R. and A. S. Muscher-Banse. 2020. Review: Regulation of gastrointestinal and

renal transport of calcium and phosphorus in ruminants. Animal 14:s29-s43.

Wilkens, M. R., J. Richter, D. R. Fraser, A. Liesegang, G. Breves, and B. Schröder. 2012b. In

contrast to sheep, goats adapt to dietary calcium restriction by increasing intestinal

absorption of calcium. Comparative Biochemistry and Physiology Part A, Molecular &

Integrative Physiology 163:396-406.

Table 1. Chemical composition and daily intake levels of feedstuffs fed during the last 21 days antepartum until 56 days of lactation of dairy sheep and goats

Chemical analysis	Grass hay (2nd cut)	Concentrate	[1]Concentrate + CaCO3
Energy, MJ NEL	5.90	7.00	6.70
Dry matter, g/kg DM	928	884	896
Crude protein, g/kg DM	155	133	115
[1]aNDFom, g/kg DM	654	267	194
[2]ADFom, g/kg DM	357	90.4	70.2
ADL, g/kg DM	38.4	18.9	13.2
Ether extract, g/kg DM	28.0	61.5	49.2
Crude ash, g/kg DM	81.1	54.2	170
HCl-insoluble ash, g/kg DM	13.3	4.30	4.60
Ca, g/kg DM	5.90	6.90	62.3
Mg, g/kg DM	2.00	1.90	1.40
P, g/kg DM	3.30	4.60	4.30

Component, kg DM/d	Sheep			Goats	
	Low Ca	High Ca		Low Ca	High Ca
			antepartum		
Grass hay (2nd cut)	2.00	2.00		2.00	2.0
[3,4]Concentrate	0.30	0.3		0.30	0.3
CaCO3	-	0.042		-	0.042
Total DM fed	2.30	2.342		2.30	2.342
[5]Nutrient intake, g/d					
Crude protein	350	345		350	345
Energy (MJ NEL)	13.9	13.8		13.9	13.8
Ca	13.9	30.5		13.9	30.5
P	7.98	7.89		7.98	7.89
Mg	4.57	4.42		4.57	4.42
			postpartum		
Grass hay (2nd cut), kg/d	2.40	2.40		2.40	2.40
[3,4]Concentrate, kg/d	0.50	0.50		0.70	0.70
CaCO3, g/d	-	-		-	-
Total DM fed	2.90	2.90		3.10	3.10
[3]Nutrient intake, g/d					
Crude protein	439	439		465	465
Energy (MJ NEL)	17.7	17.7		19.1	19.1
Ca	17.6	17.6		19.0	19.0
P	10.2	10.2		11.1	11.1
Mg	5.75	5.75		6.13	6.13

Commercial concentrate UFA 765 by supplier UFA (Union des Fédérations Agricoles). For the composition of the concentrate based on manufacturer instructions, please see Supplementary Table 1.

[1]aNDFom, starch and ash corrected NDF after treatment with heat-stable amylase and ashing of the solid residue according to Method 6.5.1 (VDLUFA, 2012)

[2]ADFom, ash corrected ADF after ashing of the solid residue according to Method 6.5.2
(VDLUFA, 2012)

[3]860 g of commercial concentrate + 140 g CaCO3.

[4]During ==antepartum== feeding either the basal concentrate from the manufacturer or the
artificially Ca-enriched concentrate were fed to ensure comparative feeding of 0.6 % in DM
Ca vs. 1.3 % in DM Ca until parturition. During subsequent lactation feeding, no comparative
feeding was applied therefore only the basal concentrate was fed to all animals at highlighted
daily amounts.

[5]Daily nutrient intake was calculated from daily supplied dry feed (hay, concentrate) and
analyzed nutrient concentrations.

Figure 1. Effects of varying dietary Ca (0.6 vs. 1.3% in DM) of dairy sheep and goats during

the last 3 weeks antepartum as well as sex and species of offspring on the birth weight (A) and

daily weight gain during the suckling phase (B). Effective sample size was 5 sheep and 6 goats

per group, which gave birth to 24 and 23 lambs and kids in total, respectively. Data is presented

as least square mean values with standard errors based on linear mixed model analysis. "Age"

refers to the time of 2nd weighing during the suckling phase, which differed between individual

animals due to management reasons (range: 33-59 days postpartum) and was therefore

introduced as a continuous control variable to the statistical analysis of daily weight gain data.

Ca*S and Ca*Sex refer to statistical interactions between diet Ca and species of animal or sex

of offspring, respectively. $P \leq 0.05$ was defined as statistical type I error threshold.

Figure 2. Effects of varying dietary Ca (0.6 vs. 1.3% in DM) of dairy sheep and goats during

the last 3 weeks antepartum as well as time and animal species on the fecal Ca concentration

(A) and urinary Ca excretion (B) from 3 weeks antepartum to parturition and 14 days

postpartum, respectively. Effective sample size was 5 sheep and 6 goats per group. Data is

presented as Log_{10}-transformed least square mean values with standard errors, respectively,

based on repeated measures linear mixed model analysis. Ca*T, Ca*S and S*T refer to

statistical interactions between diet Ca and animal species or time, respectively, as well as

animal species and time. $P \leq 0.05$ was defined as statistical type I error threshold.

Figure 3. Effects of varying dietary Ca (0.6 vs. 1.3% in DM) of dairy sheep and goats during

the last 3 weeks ==antepartum== as well as time and animal species on serum Ca (A) and colostral

Ca (B) from 3 weeks ==antepartum== to parturition and 56 days ==postpartum==, respectively. Effective

sample size was 5 sheep and 6 goats per group. Data is presented as least square mean values

with standard errors based on repeated measures linear mixed model analysis and regular linear

mixed model analysis, respectively. Ca*T, Ca*S and S*T refer to statistical interactions

between diet Ca and animal species or time, respectively, as well as animal species and time. *P*

≤ 0.05 was defined as statistical type I error threshold.

Figure 4. Effects of varying dietary Ca (0.6 vs. 1.3% in DM) of dairy sheep and goats during

the last 3 weeks antepartum as well as time and animal species on BMD from 2 weeks

antepartum to 56 days postpartum. Effective sample size was 5 sheep and 6 goats per group.

Data is presented as least square mean values with standard errors based on repeated measures

linear mixed model analysis. Ca*T, Ca*S and S*T refer to statistical interactions between diet

Ca and animal species or time, respectively, as well as animal species and time. $P \leq 0.05$ was

defined as statistical type I error threshold.

Figure 5. Effects of varying dietary Ca (0.6 vs. 1.3% in DM) of dairy sheep and goats during the last 3 weeks antepartum as well as time and animal species on serum calcitriol (A), serum bAP (B), and serum osteocalcin (C) from 3 weeks and 5 days antepartum, respectively, to 56 days postpartum. Effective sample size was 5 sheep and 6 goats per group. Data is presented as least square mean values (Log$_{10}$-scale in case of bAP activity) with standard errors based on repeated measures linear mixed model analysis. Ca*T, Ca*S and S*T refer to statistical interactions between diet Ca and animal species or time, respectively, as well as animal species and time. $P \leq 0.05$ was defined as statistical type I error threshold. Due to compromised sub-samples for analysis of calcitriol at -4 days antepartum, we cannot provide the respective data.

Figure 6. Effects of varying dietary Ca (0.6 vs. 1.3% in DM) of dairy sheep and goats during

the last 3 weeks antepartum as well as time and animal species on serum ICTP (A) and serum

CTX (B) from 3 weeks and 3 days antepartum, respectively, to 56 days postpartum. Effective

sample size was 5 sheep and 6 goats per group. Data is presented as least square mean values

(Log$_{10}$-scale in case of bAP activity) with standard errors based on repeated measures linear

mixed model analysis. Ca*T, Ca*S and S*T refer to statistical interactions between diet Ca and

animal species or time, respectively, as well as animal species and time. . $P \leq 0.05$ was defined

as statistical type I error threshold. Due to compromised sub-samples of analysis of CTX at -21

days antepartum and for ICTP and CTX at -4 days antepartum, we cannot provide the respective

data.

913

Figure 7. Effects of varying dietary Ca (0.6 vs. 1.3% in DM) of dairy sheep and goats during

the last 3 weeks ==antepartum== as well as time on caprine vitamin D receptor abundance in colonic

mucosa from 2 weeks ==antepartum== to 7 days ==postpartum==. Effective sample size was 6 goats per

group. Data is presented as least square mean values with standard errors based on repeated

measures linear mixed model analysis. Ca*T, refers to statistical interaction between diet Ca

and time. $P \leq 0.05$ was defined as statistical type I error threshold. See Figure 8 for exemplary

immunohistochemical images.

Figure 8. Exemplary immunohistochemical staining of the caprine VDR in the colon of goats

at parturition, fed 0.6 % in DM (left) or 1.3 % in DM (right) dietary Ca during the last 3 weeks

antepartum. Arrows mark SE, surface epithelium; SG, surface gland sections; IG, intermediate

gland sections; BG, basal gland sections. Samples shown in these photos were taken at

parturition with an endoscope equipped with biopsy forceps 40 cm into the large intestine from

the anus. Goats fed high Ca antepartum showed reduced immune reactivity to VDR staining,

though this could not be statistically confirmed. Figure 7 shows the statistical analysis of VDR

immune reactivity over time from -14 days antepartum to 7 days postpartum.