Pemivibart is less active against recent SARS-CoV-2 JN.1 sublineages

Qian Wang[1,4], Yicheng Guo[1,4], Jerren Ho[1], David D. Ho[1,2,3*]

[1]Aaron Diamond AIDS Research Center, Columbia University Vagelos College of Physicians and Surgeons, New York, NY 10032, USA

[2]Department of Microbiology and Immunology, Columbia University Vagelos College of Physicians and Surgeons, New York, NY 10032, USA

[3]Division of Infectious Diseases, Department of Medicine, Columbia University Vagelos College of Physicians and Surgeons, New York, NY 10032, USA

[4]These authors contributed equally

*Correspondence: dh2994@cumc.columbia.edu (D.D.H.)

Abstract

Protection from COVID-19 vaccination is suboptimal in many immunocompromised individuals. In March 2024, the Food and Drug Administration issued an Emergency Use Authorization for pemivibart (Permagard/VYD222), an engineered human monoclonal antibody, for pre-exposure prophylaxis in this vulnerable population. However, SARS-CoV-2 has since evolved extensively, resulting in multiple Omicron JN.1 sublineages. We therefore evaluated the in vitro neutralizing activity of pemivibart against the prevalent forms of JN.1, including KP.2, KP.3, KP.2.3, LB.1, and, importantly, KP.3.1.1, which is now expanding most rapidly. A panel of VSV-based pseudoviruses representing major JN.1 sublineages was generated to assess their susceptibility to pemivibart neutralization in vitro. Structural analyses were then conducted to understand the impact of specific spike mutations on the virus-neutralization results. Pemivibart neutralized both JN.1 and KP.2 in vitro with comparable activity, whereas its potency was decreased slightly against LB.1, KP.2.3, and KP.3 but substantially against KP.3.1.1. Critically, the 50% inhibitory concentration of pemivibart against KP.3.1.1 was ~6 µg/mL, or ~32.7 fold higher than that of JN.1. Structural analyses suggest that Q493E and the S31-deletion mutations in viral spike contribute to the antibody evasion, with the latter having a more pronounced effect. Our findings show that pemivibart has lost substantial neutralizing activity in vitro against KP.3.1.1, the most rapidly expanding lineage of SARS-CoV-2 today. Close monitoring of its clinical efficacy is therefore warranted. These results also highlight the imperative to expand our arsenal of preventive agents to protect millions of immunocompromised individuals who could not respond robustly to COVID-19 vaccines.

Main text:

Pemivibart (Permagard; VYD222) was authorized in March 2024 for emergency use as pre-exposure prophylaxis against COVID-19 for immunocompromised individuals who could not respond robustly to vaccines[1]. This human monoclonal antibody was derived from ADG-2, an antibody directed to the SARS-CoV-2 spike [receptor-binding domain (RBD) class 1/4 region] that has previously demonstrated protective efficacy against SARS-CoV-2 infection[2]. However, ADG-2 lost virus-neutralizing activity against the Omicron variant and its subsequent subvariants. Nine mutations (5 in the heavy chain and 4 in the light chain) were introduced to yield pemivibart, which demonstrated greater breadth in neutralizing recent SARS-CoV-2 strains[3]. For example, the engineered antibody could neutralize JN.1, the subvariant that dominated the first half of 2024, with 50% inhibitory concentrations (IC_{50}) of 0.075 μg/mL and 0.063 μg/mL in pseudovirus and authentic virus neutralization assays in vitro, respectively[3].

SARS-CoV-2 JN.1 subvariant first emerged in late 2023 and rapidly became dominant globally. In the past 6 months, JN.1 has continued to evolve, giving rise to multiple sublineages with unique spike mutations (**Figure S1A**). KP.2 was the first significant progeny to appear, but it was later outcompeted by KP.3, with both sublineages gradually displacing the original JN.1 (**Figure S1B**). More recently, KP.3.1.1, KP.2.3, and LB.1 have emerged, each independently developing a deletion of S31 (S31Δ) in the N-terminal domain (NTD) of spike (**Figure S1A**). Importantly, KP.3.1.1 is now the fasting growing sublineage worldwide (**Figure S1B**).

We therefore assessed the impact of recent SARS-CoV-2 evolution on the neutralizing activity of pemivibart. Pseudoviruses for JN.1, KP.2, KP.3, KP.2.3, KP.3.1.1, and LB.1 were constructed and subjected to neutralization assays as previously described[4]. Pemivibart neutralized both JN.1 and KP.2 in vitro with comparable activity, whereas its potency was decreased slightly against LB.1, KP.2.3, and KP.3 but substantially against KP.3.1.1 (**Figure 1A**). Critically, the IC_{50} of pemivibart against KP.3.1.1 was ~6 μg/mL, or ~32.7 fold higher than the IC_{50} for JN.1 (**Figure 1B**). We note that our virus neutralization assays yielded results concordant to those reported by Invivyd, the manufacturer of pemivibart, against multiple SARS-CoV-2 strains (**Figure S2**).

Comparing the results of KP.2 versus KP.3 (**Figure 1B**), it is apparent that Q493E mutation has a greater impact on pemivibart activity than R346T mutation. Although Q493E does not reside within the epitope of ADG-2[2] (and presumably pemivibart), it may increase resistance to pemivibart due to an enhanced binding affinity for the ACE2 receptor facilitated by an epistatic effect with the F456L mutation[5]. Comparing the results of KP.3 versus KP.3.1.1 (**Figure 1B**), it is also apparent that S31Δ mutation has an even larger impact (~9 fold) on the engineered antibody. Interestingly, multiple sublineages have converged to this mutation independently (**Figure S1A**). How S31Δ confers resistance to pemivibart, and presumably to serum antibodies in the population as well, is not immediately clear given it is quite distal to the typical SARS-CoV-2 neutralizing

epitopes. We note, however, that this deletion results in the creation of a potential N-linked
glycosylation site (**Figure S3A**), as well as the loss of hydrogen bonding with and F58, F59 and
S60 (**Figure S3B**). Both changes could alter the conformation at the bottom of the NTD and
thereby affect the up-and-down motion of the RBD (**Figure S3C**) that could negatively impact
antibodies directed to so-called class 1 and class 1/4 regions of this domain. The precise structural
mechanism notwithstanding, our findings show a substantial loss of pemivibart activity against
KP.3.1.1, the most rapidly expanding SARS-CoV-2 subvariant currently. Therefore, close
monitoring of the clinical efficacy of this newly authorized monoclonal antibody is warranted.

Figure and legend

Figure 1. Neutralization of pemivibart against JN.1 sublineages.

A. Neutralization curves of Pemivibart against the indicated pseudotyped viruses.
B. Spike mutations in the S1 region, along with the corresponding neutralization IC_{50} values and
fold-changes for the indicated JN.1 sublineages. Δ, deletion.
90

References:

1.   FDA Roundup: March 22, 2024. 2024. (Accessed 07/25/2024, at https://www.fda.gov/news-events/press-announcements/fda-roundup-march-22-2024.)

2.   Rappazzo CG, Tse LV, Kaku CI, et al. Broad and potent activity against SARS-like viruses by an engineered human monoclonal antibody. Science 2021;371:823-9.

3.   CDER Scientific Review Documents Supporting Emergency Use Authorizations for Drug and Biological Therapeutic Products | COVID-19. 2024. (Accessed 08/10/2024, at https://www.fda.gov/drugs/coronavirus-covid-19-drugs/cder-scientific-review-documents-supporting-emergency-use-authorizations-drug-and-biological.)

4.   Wang Q, Mellis IA, Bowen A, et al. Recurrent SARS-CoV-2 spike mutations confer growth advantages to select JN.1 sublineages. bioRxiv 2024:2024.05.29.596362.

5.   Taylor AL, Starr TN. Deep mutational scanning of SARS-CoV-2 Omicron BA.2.86 and epistatic emergence of the KP.3 variant. bioRxiv 2024.

Supplementary Methods

Cell lines

Vero-E6 (CRL-1586) cells and HEK293T (CRL-3216) cells were obtained from ATCC and
cultured at 37°C with 5% CO_2 in Dulbecco modified Eagle medium (DMEM) supplemented with
10% fetal bovine serum (FBS) and 1% penicillin-streptomycin. Morphology of each cell line was
confirmed visually before use. All cell lines tested mycoplasma negative.

SARS-CoV-2 variant spike plasmids

Spike-expressing plasmids for JN.1, KP.2, and KP.3 were previously generated[1]. To make the
spike-expressing plasmids for KP.2.3, KP.3.1.1, and LB.1 for pseudovirus production, spike
mutations were introduced to the JN.1 spike-expressing construct using the QuikChange II XL
site-directed mutagenesis kit (Agilent). Spike mutations of each subvariant on top of JN.1 are
shown in **Figure S1A**.

Pseudovirus production

SARS-CoV-2 pseudoviruses were produced in a vesicular stomatitis virus (VSV) background, in
which the native VSV glycoprotein was replaced by SARS-CoV-2 variant spikes, as previously
described[2]. Briefly, the spike-expressing construct was transfected into HEK293T cells using
1mg/mL PEI MAX (Polysciences Inc). 24 hours later, the transfected cells were infected with
VSV-G pseudotyped ΔG-luciferase (G*ΔG-luciferase, Kerafast) for 2 hours, then washed three
times with culture medium before being cultured in fresh medium for an additional 24 hours. Anti-
VSVG (I1) antibody[3] was added to deplete non-pseudotyped viruses. Pseudoviruses were then
harvested, clarified by centrifugation, aliquoted and stored at -80°C before use.

Pseudovirus neutralization

Each SARS-CoV-2 pseudovirus was titrated on Vero-E6 cells to standardize the viral infectious
dose before use in neutralization assays. Serially diluted antibodies (5-fold dilution starting from
10 µg/mL) were added to 96-well plates. Then, pseudoviruses were added and incubated with the
serial antibody dilutions at 37 °C for 1 hour. Wells containing only pseudoviruses were included
as controls in each plate. Vero-E6 cells were then added at a density of 4×10^4 cells per well and
incubate at 37 °C overnight. Cells were lysed and luminescence was measured using the Luciferase
Assay System (Promega) and SoftMax Pro v.7.0.2 (Molecular Devices) according to the
manufacturers' instructions. Data were analyzed using GraphPad Prism v.9.3.

Acknowledgements

This study was supported by funding from the NIH SARS-CoV-2 Assessment of Viral Evolution
(SAVE) Program (subcontract no. 0258-A709-4609 under federal contract no. 75N93021C00014)
and the Gates Foundation (project INV019355) to D.D.H. and internal startup funding UR014016
from Columbia University to Y.G. We thank all who contributed their data to GISAID.

Author Contributions

D.D.H. conceived of this project. Q.W. and J.H. performed experiments. Y.G. conducted
bioinformatic analyses. Q.W., Y.G., and D.D.H. analyzed the results and wrote the manuscript.

Declaration of Interests

D.D.H. co-founded TaiMed Biologics and RenBio, serves as a consultant for WuXi Biologics
and Brii Biosciences, and is a board director at Vicarious Surgical. The remaining authors have
no competing interests to declare.

Supplementary Figures

Figure S1. Overview of circulating SARS-CoV-2 JN.1 sublineages.

A. Evolutionary pathway of JN.1 sublineages. Δ, deletion.

B. Global frequencies of major SARS-CoV-2 forms during the indicated time period. Sequence

frequencies were obtained from the Global Initiative on Sharing All Influenza Data (GISAID).

Figure S2. Correlation of pseudovirus neutralization IC$_{50}$ values of pemivibart reported by

Invivyd and values generated as a part of this study.

Figure S3. Predicted effects of the S31Δ mutation on the JN.1 spike
A. The S31Δ mutation may introduce a glycosylation site at N30.
B. Structural details of the S31-surrounding region in the NTD. Black dash lines indicate
hydrogen bonds. PDB: 8Y5J.
C. Overall view of ADG-2/pemivibart binding site on the RBD in the up position and S31Δ on
the external surface of NTD.

Supplementary References
97
1.   Wang Q, Mellis IA, Bowen A, et al. Recurrent SARS-CoV-2 spike mutations confer
growth advantages to select JN.1 sublineages. bioRxiv 2024:2024.05.29.596362.
2.   Liu L, Wang P, Nair MS, et al. Potent neutralizing antibodies against multiple epitopes
on SARS-CoV-2 spike. Nature 2020;584:450-6.
3.   Lefrancios L, Lyles DS. The interactionof antiody with the major surface glycoprotein of
vesicular stomatitis virus. I. Analysis of neutralizing epitopes with monoclonal antibodies.
Virology 1982;121:157-67.
105