Endoplasmic reticulum stress inhibition preserves mitochondrial function and cell survival during early onset of isoniazid-induced oxidative stress

Truong Thi My Nhung[1,#], Nguyen Ky Phat[2,#], Trinh Tam Anh[2], Tran Diem Nghi[1], Nguyen Quang Thu[2], Ara Lee[3], Nguyen Tran Nam Tien[2], Nguyen Ky Anh[4], Kimoon Kim[3], Duc Ninh Nguyen[5], Dong-Hyun Kim[2], Sang Ki Park[1], Nguyen Phuoc Long[2,*]

[1]Department of Life Sciences, Pohang University of Science and Technology, Pohang 37673, Republic of Korea

[2]Department of Pharmacology and PharmacoGenomics Research Center, Inje University College of Medicine, Busan 47392, Republic of Korea

[3]Department of Chemistry, Pohang University of Science and Technology (POSTECH), Pohang 37673, Republic of Korea

[4]Faculty of Pharmacy, Ton Duc Thang University, Ho Chi Minh City 700000, Vietnam

[5]Section for Comparative Pediatrics and Nutrition, Department of Veterinary and Animal Sciences, University of Copenhagen, 1870, Frederiksberg, Denmark

[#]These authors contributed equally to this work

*Corresponding author:

Nguyen Phuoc Long (email: phuoclong@inje.ac.kr)

Abstract

A comprehensive understanding of isoniazid (INH)-mediated hepatotoxic effects is

essential for developing strategies to predict and prevent severe liver toxicity in tuberculosis

treatment. Our study utilized multi-omics profiling to investigate the toxic effects of INH,

revealing significant involvement of endoplasmic reticulum (ER) stress, mitochondrial

impairment, redox imbalance, and altered metabolism. Followed-up mechanistic studies revealed

that INH triggered the generation of cytosolic reactive oxygen species (ROS) and the activation of

the Nrf2 signaling pathway prior to mitochondrial ROS accumulation. Subsequently, INH

disrupted mitochondrial function by impairing respiratory complexes I-IV and caused

mitochondrial membrane proton leaks without affecting ATP synthase activity, together leading

to mitochondrial depolarization and reduced ATP production. These disturbances enhanced

mitochondrial fission and mitophagy. While much attention has been given to mitochondrial

dysfunction and oxidative stress in INH-induced hepatotoxicity, our findings highlight the

potential of inhibiting ER stress during early INH exposure to mitigate cytosolic and mitochondrial

oxidative stress. We further revealed the critical role of Nrf2 signaling in protecting liver cells

under INH-induced oxidative stress by maintaining redox homeostasis and enabling metabolic

reprogramming via regulating the expression of antioxidant genes and cellular lipid abundance.

We also identified other antioxidant pathways (e.g., selenocompound metabolism, HIF-1 signaling

pathway, and pentose phosphate pathway) as potential alternative mechanisms besides Nrf2

signaling in response to INH-induced oxidative stress. In conclusion, our research emphasizes the

importance of ER stress, redox imbalance, metabolic changes, and mitochondrial dysfunction

underlying INH-induced hepatotoxicity.

Keywords: isoniazid, multi-omics, endoplasmic reticulum stress, oxidative stress, mitochondria

dysfunction, complex I-IV impairment

Introduction

Tuberculosis (TB) remains one of the leading causes of mortality, accounting for approximately 1.3 million deaths in 2022 [1]. TB treatment is a long-term and complicated procedure, with a standard regimen for drug-susceptible TB requiring at least six months and often associated with various adverse drug reactions [2-4]. Isoniazid (INH), a first-line anti-TB drug, plays a vital role in this regimen [5]. However, severe adverse events, particularly hepatotoxicity, are a significant concern after prolonged use of INH [6]. Therefore, it is crucial to gain more insights into the mechanistic understanding of INH-induced hepatotoxicity.

Several studies highlight the emerging role of mitochondrial dysfunction in the mechanisms underlying INH-induced hepatotoxicity [7]. INH has been shown to cause mitochondrial dysfunction, including impaired mitochondrial complexes, reduced mitochondrial respiration, and diminished energy production [8]. Specifically, INH impairs mitochondrial complexes I, II, and III [9] and mediates the generation of reactive oxygen species (ROS), leading to mitochondrial oxidative stress and apoptosis [10]. Additionally, INH affects mitochondrial biogenesis and dynamics [11]. However, conflicting reports exist, with some studies suggesting that INH does not affect mitochondrial complex I and II activity [8]. While INH alone has been found to reduce whole liver glutathione (GSH) levels and increase lipid peroxidation, it does not alter mitochondrial GSH, oxidized glutathione (GSSG) content, or lipid peroxidation [10], nor does it induce liver cell death [8, 12, 13]. These contradictions raise questions about whether mitochondrial oxidative stress and dysfunction are the primary mechanisms through which INH triggers liver damage.

Nuclear factor erythroid-derived factor 2-related factor 2 (Nrf2) is crucial for cellular

defense against oxidative stress and for regulating antioxidant and cytoprotective genes. In the

normal cell state, Nrf2 is kept inactive in the cytoplasm by Keap1 [14, 15]. However, under

oxidative stress, ROS modify Keap1, Nrf2 moves to the nucleus [16], where it binds to DNA and

triggers the production of detoxifying enzymes like glutathione S-transferase (GST), heme

oxygenase-1 (Hmox1), and superoxide dismutase (SOD). These enzymes neutralize ROS and

repair damaged proteins, reducing oxidative damage and enhancing cell survival. Nrf2 also helps

regenerate antioxidants like glutathione. It is reported that Nrf2 can upregulate the expression of

SLC7A11, a component of glutamate-cystine antiporter (xCT), which increases cystine uptake,

inducing glutathione synthesis [17]. In zebrafish, expression of *Nrf2* and its target genes (*HMOX1*,

NQO1, *GCLM*, *GCLC*) were found to be upregulated upon INH exposure [18]. However, the role

of Nrf2 signaling in INH-induced liver toxicity has yet to be fully elucidated.

Our study provided a more comprehensive analysis of INH-associated hepatotoxicity.

Previous research has often examined isolated mechanisms in different biological systems,

resulting in fragmented insights. In contrast, our study utilizes multi-omics analysis to offer an

integrated view of the metabolic alterations and their association with oxidative stress induced by

INH toxicology. We captured the complex interplay between key molecular events, including

endoplasmic reticulum (ER) stress, redox imbalance, mitochondrial dysfunction, and cellular

metabolic reprogramming. Furthermore, by investigating INH hepatotoxicity in Nrf2-knockout

(Nrf2-KO) cells, we highlighted the critical role of Nrf2 signaling and identified alternative

protective mechanisms against INH-induced oxidative stress. Our findings present a more

complete picture of the molecular mechanisms underlying INH-associated hepatotoxicity and

suggest potential targets for both preventive and therapeutic interventions for INH-induced liver

injury.

Material and Methods

Reagents and Plasmid Construction

Detailed information regarding antibodies, chemicals, and plasmid construction is provided in

Supplementary Material and Methods.

Cell culture and transfection

HepG2, Hep3B, and Huh-7 cell lines were cultured in MEM or DMEM media

supplemented with 1% (*v/v*) penicillin-streptomycin (Hyclone, SH40003.01), 1% (*v/v*) sodium

pyruvate (Gibco, 11360070), and 10% (*v/v*) FBS (Sigma, 12003C) and maintained in a 5% CO_2

humidified incubator at 37°C. Cells were transfected using Lipofectamine 2000 in OptiMEM2000,

following the manufacturer's protocol, and media were replaced 12 hours post-transfection.

Generation of the Nrf2 KO HepG2 cell line

Nrf2-KO cells were created using single-guide RNAs (sgRNAs) inserted into the

gRNA_Cloning Vector (Addgene, #41824). The sgRNA sequences were

GCGACGGAAAGAGTATGAGC and TATTTGACTTCAGTCAGCGA. The efficiency of these

sgRNAs was verified by observing GFP-expressing cells using pCAG-EGxxFP (Addgene,

#50716), which contained 500 bp of genomic DNA, including the gRNA target sites. Following

sgRNAs and Cas9 transfection, cells were subjected to a 4-d selection period using Geneticin, with

the dosage determined by an antibiotic kill curve (0.5 mg/mL). Subsequently, cells were plated at

single-cell density and allowed to develop for 10-14 d. KO clones were identified and selected

through immunoblotting analysis.

IC20 and IC50 determination

Cells were seeded in 96-well plates at 4×10^4 cells per well and treated with varying concentrations of Isoniazid (500 µM to 75 mM) for 72 hours, with media refreshed every 24 hours. After treatment, 10 µL of CCK-8 reagent was added to each well, and OD at 460 nm was measured to determine cell viability. IC20 and IC50 values were calculated using dose-response curves for each cell line.

Cellular and molecular imaging assays

For detailed descriptions of other methods, including ROS detection, immunoblotting, laser scanning confocal microscopy, lipid droplet staining, and immunocytochemistry, please refer to the **Supplementary Material and Methods section**.

MitoXpress Xtra oxygen consumption assay

Oxygen consumption was monitored using the MitoXpress Xtra Oxygen Consumption Assay, with 1 µM MitoXpress added to each well and sealed with mineral oil. Using a microplate reader (Infinite M200, TECAN), fluorescence was measured at 37°C, with signals recorded for 30 minutes to establish basal respiration, followed by sequential addition of oligomycin (1 µg/mL), FCCP (2 µM), and rotenone/antimycin A (1 µM each), recording signals for 30 minutes after each addition. Measurements were normalized to 1000 cells and used to determine basal, ATP-linked, maximal, and non-mitochondrial respiration rates by comparing oxygen consumption rate (OCR) before and after each treatment.

RNA extraction, cDNA synthesis and RT-qPCR

Cell pellets were subjected to RNA isolation using TRIzol reagent (Virginia Tech Bio-

Technology, #TS200-001). cDNA was synthesized using Invitrogen Superscript IV Reverse

Transcriptase (Invitrogen, #18090050) and oligo-dT primers (Invitrogen, #18418020). qPCR was

conducted with FastStart SYBR Green Master Mix (Roche, #04913850001) on a StepOnePlus™

Real-Time PCR System (Applied Biosystems, #15360337), using GAPDH for normalization via

the $2^{-\Delta\Delta Ct}$ method. The following primers were used for housekeeping gene and Nrf2 target genes:

GAPDH, CTCCTGCACCACCAACTGCT and GGGCCATCCACAGTCTTCTG; *GCLC*,

TCCAGGTGACATTCCAAGCC and GAAATCACTCCCCAGCGACA; *HMOX1*,

AGTCTTCGCCCCTGTCTACT and GCTGGTGTGTAGGGGATGAC; *NQO1*,

GCACTGATCGTACTGGCTCA and CCACCACCTCCCATCCTTTC; *SLC7A11*,

TTTTCTGAGCGGCTACTGGG and CAGCTGGTAGAGGAGTGTGC.

High-throughput cellular omics experiments

Cellular extraction of metabolites, lipids, and proteins for metabolomics, lipidomics, and

proteomics data acquisition, as well as data processing and normalization, are described in detail

in the **Supplementary Material and Methods**.

Statistical analysis

Quantitative data are shown as mean ± standard deviation (SD) unless specified differently.

Sample sizes differed according to the experimental conditions. Statistical tests were carried out

using GraphPad Prism 10.2.3. Two-sided Student's t-test was used to compare the means between

the two groups. To compare means across more than two groups, one-way or two-way ANOVA

with Tukey's post hoc test was applied. A significance level of 0.05 was used.

The log-transformed median-normalized omics data were subject to univariate analyses.

The *limma* method [19] was used for single-factor differential analysis to identify differential

proteins, while the unpaired two-sided t-test was performed to identify differential metabolites and

lipids. Regarding two-factor differential analysis, ExpressAnalyst [20] (for proteomics data) and

MetaboAnalyst 6.0 [21] (for metabolomics and lipidomics data), which implemented the limma

functionality, were employed. The false discovery rate cut-off of 0.05 was applied to consider the

significant differential metabolites/lipids/proteins.

We conducted functional analysis to gain biological insights into molecular pathways from

significantly differential proteins, metabolites, or lipids. Detailed information regarding functional

analysis can be retrieved from **Supplementary Material and Methods.**

Results

INH exposure induces ROS and apoptosis in HepG2, Hep3B, and Huh-7 cells

To explore the cytotoxicity of INH in hepatocytes, we assessed cell viability in three

different liver cell lines (HepG2, Hep3B, Huh-7) at various concentrations of INH ranging from

0.5 to 75 mM (Fig. 1A). Using these genetically and functionally diverse cell lines ensures that

our findings on INH's hepatotoxicity are robust and generalizable, confirming that the observed

cytotoxic patterns are not restricted to the specific characteristics of a single cell type. After a 72-

h incubation, the IC20 and IC50 values of INH for each cell line were determined, representing

20% and 50% reductions in cell viability compared to the control group. IC20 and IC50 were

respectively 17 and 32.5 mM for HepG2 WT cells, 15 and 60 mM for Hep3B cells, and 25 and 45

mM for Huh-7 cells. We further examined how IC20 and IC50 doses of INH affected the

expression of apoptosis-related proteins, including Bax, Bcl-2, cleaved Caspase-9, and cleaved

Caspase-3. Using immunoblotting analysis, we observed that increasing concentrations of INH for

72 h elevated the levels of Bax, cleaved Caspase-9, and cleaved Caspase-3, accompanied by a

corresponding decrease in Bcl-2 levels in all three hepatic cell lines (Fig. 1B). These results

indicate that INH induces apoptosis in these liver cell lines in a dose-dependent manner.

Figure 1. Isoniazid (INH) impaired cell viability and induced apoptosis, associated with elevated cellular ROS levels. (A) Dose-response curve of INH in HepG2, Hep3B, and Huh-7 cells for 72 h. (B) INH exposure influenced the expression of apoptosis-related proteins. Representative immunoblots of Bax, Bcl-2, cleaved Capase-9, and cleaved Caspase-3 proteins after 72-h INH exposure with concentrations that inhibit 20% and 50% cell growth in three cell lines. β-actin served as protein loading control. (C) Representative immunofluorescence images of

DHE staining (left) and quantitative data (right) after INH treatment for 72h in the indicated cells.

Data are presented as means ± SD from three independent experiments. Statistical differences

between groups were determined by one-way ANOVA with Tukey post-tests for multiple

comparisons, ****P-value < 0.0001.

It was reported that INH-induced apoptosis is related to oxidative stress [22, 23], so we

utilized dihydroethidium (DHE) staining to investigate the impact of INH on ROS production in

these three liver cell lines treated with varying concentrations of INH for 72 h. As shown in Fig.

1C, INH-treated samples in all cell lines exhibited significantly increased ROS production

compared to the control group. Altogether, the findings from apoptosis-related protein analysis

and DHE staining suggest that oxidative stress is associated with the cytotoxic effects triggered by

INH.

193   Multi-omics reveals a comprehensive landscape regarding alterations of multiple cellular

194   processes under INH exposure

INH treatments for 72 h significantly altered the proteome of HepG2 WT cells compared

to controls (Fig. 2A). Specifically, 1376 and 1855 differential proteins were identified in the 17

mM and 32.5 mM INH treatments versus control comparisons, respectively (Supplementary Fig.

S1A-B). Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis

revealed that 32.5 mM INH treatment affected ER unfolded protein response ("Protein processing

in endoplasmic reticulum"), redox homeostasis ("Cysteine and methionine metabolism,"

"Glutathione metabolism," "Ferroptosis"), mitochondrial metabolic pathways ("TCA cycle,"

"Pyruvate metabolism," "Oxidative phosphorylation," and "Fatty acid metabolism"), and ROS-

related pathways ("Chemical carcinogenesis - reactive oxygen species") (Fig 2. B-E,

Supplementary Table S1). The 17 mM INH-treated group followed a similar pattern of molecular

alterations but to a lesser extent (Fig 2. B-E). The differential proteins were also enriched in Gene Ontology Cellular Component (GO:CC) terms related to ER and mitochondrial complexes (Supplementary Table S2). The abundance of multiple proteins related to ER stress response, including ERO1A, HSPA5, HSP90B1, PDIA3, PDIA4, CANX, CALR, DNAJB11, ERLEC1, HYOU1, and ERP29, were increased (Fig. 2B, Supplementary Fig. S2A), suggesting the involvement of the unfolded protein response (UPR).

Fig. 2. Proteomics analysis suggested that INH treatment induced endoplasmic reticulum stress, mitochondrial dysfunction, oxidative stress, altered redox state, and perturbations of

cellular metabolism in HepG2 cells. (A) Principal component analysis scores plot of proteome

profiles of INH-treated and untreated WT HepG2 cells. (B-E) Heatmaps of significantly

differential proteins between INH-treated and untreated WT HepG2 cells enriched into significant

KEGG pathways related to endoplasmic reticulum (B), mitochondria (C), redox balance (D), and

reactive oxygen species generation (E). Abbreviation: WT, wild-type; FDR, false discovery rate;

KEGG, Kyoto Encyclopedia of Genes and Genomes.

We also observed significant enrichment of mitochondrial protein complexes in both

oxidative phosphorylation and ROS-related pathways (Fig. 2C and E, Supplementary Fig. S2B).

In particular, the decreased abundance of numerous subunits of complex I (NDUFA2, NDUFA4,

NDUFA9, NDUFS1, NDUFS2, NDUFS3, NDUFS4, NDUFV1, among others), II (SDHA,

SDHB, SDHC), III (UQCRFS1), IV (COX6B1, COX7A2, COX7C) suggests impaired

mitochondrial electron transport chain function, which is essential for ROS induction. Conversely,

the increased expression levels of complex V subunits (ATP5F1A, ATP5F1B, ATP5F1C,

ATP6V1B2, ATP6V1E1, among others), TCA cycle enzymes (MDH1, MDH2, PDHB, IDH3A,

IDH3G, IDH2, PC, ACO1, DLAT, DLD, OGDH, and SUCLG1), enzymes involved in the

conversion of pyruvate to acetyl-CoA (PDHB, PDHA1, and DLAT), β-oxidation enzymes

(ACSL1, ACSL5, CPT1A, CPT2, HADHA, HADHB, ACADS, and ACADVL) were observed.

Remarkably, these molecules are all involved in ATP production and maintenance of energy

homeostasis. Moreover, we found higher abundance of proteins for glutathione synthesis and

antioxidant defense (GPX1, GSTK1, GSTO1, G6PD, GSR, and GCLC), ferroptosis (ACSL1 and

ACSL5), detoxifying enzymes in response to oxidative stress (SOD2, AKR1C2, AKR1C3, NQO1,

MGST1, MGST2, and GSTM3). These findings suggest the alterations in glutathione production,

lipid peroxidation, and redox imbalance due to oxidative stress caused by INH treatment over 72

h (Fig. 2D-E). Collectively, proteomic data suggest the changes in ER stress response,

mitochondrial complex I-IV dysfunction, along with metabolic responses to stress conditions

induced by INH.

Since metabolic alterations were found at the proteome level, we further performed high-

throughput metabolic profiling to examine the consequences of INH treatment in cellular

metabolism via metabolomics and lipidomics. In the principal component analysis scores plot, the

metabolome of the HepG2 WT cells receiving INH treatment showed significant separation from

the control group (Fig. 3A). Compared to the control, the 17 mM INH-treated group had 20

significantly differential metabolites, while 32.5 mM INH-treated group had 32 significantly

differential metabolites (Supplementary Fig. S3A-B). Despite the increased abundance of TCA

cycle enzymes and proteins associated with oxidative stress response, metabolites related to the

TCA cycle (e.g., citrate, isocitrate, malate, and beta-nicotinamide adenine dinucleotide) and

glutathione metabolism (e.g., GSH, GSSG, glutamine) were found to be significantly decreased in

32.5 mM INH-treated group (Fig. 3B). The 17 mM INH-treated group showed a similar pattern

with a lesser extent in significant metabolites (Fig. 3B). In addition, results of pathway analysis

showed that 17 mM INH treatment was associated with a KEGG pathway related to redox balance

("Glutathione metabolism"), and 32.5 mM INH treatment was associated with a pathway related

to mitochondria ["Citrate cycle (TCA cycle)"] (Supplementary Table S3). These findings

suggested a disruption in the TCA cycle and an altered redox state due to the toxic effects of INH

treatment for 72 h, even with the enhanced expression of TCA enzymes and antioxidant proteins,

which might be the compensatory mechanisms for maintaining cellular homeostasis. Besides, in

both 17 mM and 32.5 mM INH-treated groups, the abundance of carnitine and acylcarnitine with

shorter chain [CAR(2:0)] were decreased, whereas the abundance of acylcarnitines with longer

chain [CAR(16:0) and CAR(14:0)] was increased compared to controls (Fig. 3B).

Fig. 3. Metabolomics and lipidomics analyses suggested cellular perturbations related to

mitochondrial dysfunction, oxidative stress, and lipid droplet formation in HepG2 cells

under INH exposure. (A) PCA scores plot of metabolome profiles of INH-treated and untreated

WT HepG2 cells. (B) Heatmap of significantly differential metabolites between INH-treated WT

HepG2 cells and controls. (C) PCA scores plot of lipidome profiles of INH-treated and untreated

WT HepG2 cells. (D) Heatmap presents differences in total lipid subclass levels between INH-

treated WT HepG2 cells and controls. (E) PCA heatmap of lipid ontology enrichment analysis of

significantly differential lipids between INH-treated WT HepG2 cells and controls. (F)

Representative images of INH-treated HepG2 cells stained with BODIPY 409/503 (left) and

quantification of lipid droplet (LD) accumulation in cells (right). Both concentrations of INH

increased LD number, size (area), and intensity in HepG2 cells after 72 h of treatment. Data are

shown as means $\pm$ SD, analyzed from more than 100 cells per group. Statistical differences

between groups were determined by one-way ANOVA with Tukey post-tests for multiple

comparisons; ****$P < 0.0001$. Abbreviation: PCA, principal component analysis, WT, wild-type;

FDR, false discovery rate; PC, Phosphatidylcholine; BMP, Bismonoacylglycerophosphate; PC(O-

), Ether-linked phosphatidylcholine; CL, Cardiolipin; SHexCer, Sulfatide; PG,

Phosphatidylglycerol; SM, Sphingomyelin; PI, Phosphatidylinositol; PS, Phosphatidylserine; PE,

Phosphatidylethanolamine; Cer, Ceramide; LPC, Lysophophatidylcholine; LPE,

Lysophosphatidylethanolamine; LPC(O-), Ether-linked lysophosphatidylcholine; PE(O-), Ether-

linked phosphatidylethanolamine; LPE(O-), Ether-linked lysophosphatidylethanolamine; NAE,

N-acyl ethanolamines; FA, Fatty acid; TG(O-), Ether-linked triacylglycerol; HexCer,

Hexosylceramide; DG, Diacylglycerol; DG(O-), Ether-linked diacylglycerol; CE, Cholesteryl

ester; TG, Triacylglycerol.

Regarding the lipidomics analysis, 17 mM and 32.5 mM INH-treated groups showed a

distinct lipid profile compared to the control group (Fig. 3C) with 281 and 358 significantly

differential lipids, respectively (Supplementary Fig. S3C-D). The total lipid levels of cholesteryl

ester (CE), diacylglycerol (DG), triacylglycerols (TGs), and fatty acids (FAs) were generally

higher in the treatment group, where the 32.5 mM clearly showed a more considerable extent of

significant lipid species (Fig. 3D, Supplementary Fig. S4). On the contrary, total lipid levels of

sphingomyelin (SM), phosphatidylethanolamines (PEs), phosphatidylserines (PSs),

phosphatidylinositols (PIs), phosphatidylcholines (PCs), cardiolipins (CLs), and ether-linked

phosphatidylcholines [PC(O-)] were lower in the treatment groups and also exhibited a dose-

dependent pattern (Fig. 3D, Supplementary Fig. S4). The enrichment analysis indicated a dose-

dependent reduction in the abundance of lipid species associated with ER, mitochondria, and

membrane components in INH-treated groups (Fig. 3E). Moreover, the INH-treated groups also

displayed an upregulation of polyunsaturated fatty acids (PUFAs) and lipid species related to the

formation of lipid droplets (Fig. 3E). The elevation in the abundance of PUFAs, FAs, and

acylcarnitine, combined with the reduction of carnitine level, indicated the disruption of the β-

oxidation process. Additionally, these observations suggested that the carnitine shuttle pathway,

which plays a key role in the carnitine and long-chain fatty acids transport through mitochondrial

membranes for β-oxidation, was impaired. The increased expression levels of proteins involved in

converting pyruvate to acetyl-CoA might compensate for the shortage of acetyl-CoA due to

disrupted β-oxidation. In agreement with our lipidomics data, we observed an increase in the

number, size, and intensity of lipid droplets in INH-treated cells after 72 h using BODIPY staining.

However, no significant difference was noted between the two concentrations (Fig. 3F).

Overall, our results from multi-omics data analysis indicated that exposing HepG2 WT

cells to INH was associated with 4 critical molecular events: ER stress, mitochondrial dysfunction,

disrupted redox imbalance and oxidative stress, and significantly altered metabolism.

INH induces mitochondrial dysfunction and stress responses in HepG2 WT cells

We first explored the effects of 72 h of INH treatment on mitochondria by quantifying

mitochondrial morphology in HepG2 WT cells (Fig. 4A-E). INH significantly decreased the

mitochondrial content and area (Fig. 4B-C), reduced the aspect ratio (Fig. 4D), and increased the

number of individual mitochondria (Fig. 4E). These data suggest that INH enhanced mitochondrial

fission and mitophagy.

Fig. 4. INH exposure resulted in mitochondrial dysfunction after 72 h. (A-E) Mitochondrial

morphology analysis in HepG2 cells treated with INH for 72 h. (A) Representative images of

mitochondrial networks using TOM20-mScarlet in HepG2 cells treated with 0 mM, 17 mM, and

32.5 mM INH. Magnified views of the boxed regions in the main images (bottom left) highlight

changes in mitochondrial structure. Scale bar: 10 µm. (B) Quantification of mitochondrial content

is expressed as the percentage of cytosol occupied by mitochondria. (C) Measurement of average

mitochondrial area. (D) Aspect ratio analysis, representing the ratio of major to minor axes of the

mitochondria. (E) The absolute number of mitochondria per cell. (F) TMRM fluorescence intensity

over time was normalized to baseline, indicating mitochondrial membrane potential in the presence

of oligomycin and FCCP. (G) Mito-AT1.03 YFP/CFP fluorescence ratio, indicating ATP levels in

mitochondria. (H) Representative images and quantification of mitoSypHer-red fluorescence

intensity, showing mitochondrial pH. Scale bar: 2 mm. (I-L) Mitochondrial oxygen consumption

rate determined using mitoXpress of HepG2 after 72-h INH treatment including (I) basal

respiration rate, (J) ATP production rate, (K) proton leak, and (L) maximal respiration compacity.

(M) Monitoring mitochondrial Ca^{2+} uptake in INH-treated HepG2 cells using GCaMP6mt sensor

before and after ATP addition (left). Data for the maximum peak of GCaMP6mt signals are shown

(right). (N) Quantification of mitoKeima-Red using fluorescence intensity ratio (550/440) in WT

HepG2 cells after 72 h of INH exposure. Data are presented as means ± SD, analyzed from more

than 100 cells per group. Statistical differences between groups were determined using one-way

ANOVA with Tukey's post-test for multiple comparisons; n.s., not significant, *P-value <0.05,

**P-value < 0.01, ***P-value < 0.001, ****P-value < 0.0001.

We monitored mitochondrial membrane potential (Δψm) using TMRM fluorescence (Fig.

4F). INH exposure for 72 h caused immediate depolarization of Δψm upon treatment with the ATP

synthase inhibitor oligomycin and a sharp drop with the protonophore FCCP. These results suggest

that INH impairs mitochondrial respiration, and to maintain Δψm, mitochondria rely primarily on

ATP hydrolysis by complex V. We further found the depletion of mitochondrial ATP production

under 72-hour INH treatment using a mitochondria-targeted ATP probe (mitoAT1.03) (Fig. 4G).

In addition, mitochondrial acidification under INH exposure was indicated by the reduction in

mito-SypHer red signals (Fig. 4H), likely due to increased proton accumulation within the matrix.

To determine the impact of INH on mitochondrial function, we measured mitochondrial

oxygen consumption rates (OCRs) using mitoXpress at 72 h after treatment. Fig. 4I-L shows that

INH-treated groups exhibited lower basal respiration rates, reduced ATP production, and

decreased maximal respiratory capacity despite increased proton leakage. These findings suggest

that INH induces uncoupling of the mitochondrial membrane proton leak without promoting ATP

synthesis and causes dysfunction in mitochondrial complexes I-IV, while ATP synthase activity

remains unaffected. Additionally, we observed increased Ca^{2+} transfer from the ER to

mitochondria (Fig. 4M). In support of mitochondrial morphology analysis, we detected an increase

of mitoKeima-Red, a mitophagy marker, in the INH-treated groups (Fig. 4N).

These findings highlight the impact of INH on mitochondrial membrane integrity and

functions, including impairment of mitochondrial complex I-IV, uncoupling of the mitochondrial

membrane proton leak causing mitochondrial depolarization, and increased mitochondrial fission

and mitophagy.

ER stress is related to cellular ROS generation and altered redox balance during early INH

exposure

To investigate if mitochondrial ROS is the critical factor impairing mitochondrial function

and causing cellular oxidative stress, cellular ROS was monitored using cellROX and

mitochondrial ROS using mitoSOX from 12 to 72 h after INH treatment. A significant increase in

cellular ROS was observed at 12 h (Fig. 5A). In comparison, mitochondrial ROS levels increased

only after 48 h (Fig. 5B). This suggests that INH initially increases cytosolic ROS, followed by a

later increase in mitochondrial ROS, indicating that mitochondrial complex dysfunction may not

be the primary cause of INH-induced cellular oxidative stress.

Fig. 5. Cytosolic ROS levels rapidly increased after INH exposure, followed by a subsequent

rise in mitochondrial ROS. (A-B) Time-dependent effects of INH treatment on cellular and

mitochondrial ROS levels in HepG2 cells. (A) Representative images of HepG2 cells treated with

0 mM, 17 mM, and 32.5 mM INH for 0, 12, 24, 48, and 72 h, stained with CellROX Green to

detect cellular ROS (left). Quantification of CellROX fluorescence intensity over time is shown

on the right. Scale bar: 2 mm. (B) Representative images of HepG2 cells treated with 0 mM, 17

mM, and 32.5 mM INH for 0, 12, 24, 48, and 72 h, stained with MitoSOX Red to visualize

mitochondrial ROS (left). The right panel shows the quantification of MitoSOX fluorescence

intensity over time. Scale bar: 2 mm. (C) Quantification of the cytoGRX1-roGFP2 fluorescence

intensity ratio (405/488 nm) in HepG2 cells treated with 0 mM, 17 mM, and 32.5 mM INH for 0,

12, 24, 48, and 72 h, indicating changes in cytosolic redox state (GSSG/GSH ratio) over time. (D)

Measurement of the mitoGRX1-roGFP2 fluorescence intensity ratio (405/488 nm) in HepG2 cells

treated with 0 mM, 17 mM, and 32.5 mM INH for 0, 12, 24, 48, and 72 h, showing alterations in

the mitochondrial redox state (GSSG/GSH ratio) over time. (E-H) Mitochondrial oxygen

consumption rate in HepG2 cells was determined using mitoXpress after 24-h INH treatment. The

analysis includes (E) basal respiration rate, (F) ATP production rate, (G) maximal respiration rate,

and (H) proton leak. (I) Cytosolic ROS levels at 24 h, determined using cytoHyPer-Red signals,

in INH-treated cells alone or in combination with different compounds: mitoTEMPO 15 μM,

TUDCA 1mM, and VAS2870 10 μM. (J) Mitochondrial ROS levels at 48 h, monitored using

mitochondrial HyPer-Red, in cells treated with INH alone or in mitoTEMPO 15 μM, TUDCA

1mM, and VAS2870 10 μM. (K) Western blot analysis of the indicated proteins following INH

treatment for 24 h. Data are displayed as means ± SD, based on analysis of more than 100 cells

per group. Statistical differences between groups were determined using one-way ANOVA with

Tukey's post-test for multiple comparisons; n.s., not significant, ****P-value < 0.0001.

To further explore the impact of increased ROS on glutathione oxidation, the ratio of

oxidized glutathione (GSSG) to reduced glutathione (GSH) was measured using cytoplasmic and

mitochondrial ratio metric glutathione sensors (Fig. 5C-D) from 12 to 72 h after INH treatment.

Supporting the ROS time-lapse data, cytosolic GSSG/GSH was significantly higher from 12 h,

while it only markedly increased in mitochondria from 48 h. Additionally, measurements of OCR

24 h after INH exposure did not show significant changes in basal respiration rate, ATP production,

maximal respiratory capacity, or proton leak in both the INH-treated and control groups (Fig. 5E-

H). This suggests that mitochondrial dysfunction is not the primary cause of cellular oxidative

stress triggered by INH.

To identify the primary source of cytosolic ROS early increase, inducing the antioxidant

response at 24 h, we inhibited mitochondrial ROS, ER stress, and its associated ROS, and

membrane-associated NADPH oxidases using mitoTEMPO, TUDCA, and VAS2870, respectively

[24]. As illustrated in Figure 5I-J, the ER stress inhibitor TUDCA successfully reduced both

cytoplasmic and mitochondrial ROS at 24 and 48 h after INH exposure, indicating that the ROS

primarily originated from ER stress. We further observed a significant increase in the protein levels

of peIF2, Ero1-Lα, and calnexin 24 h after INH treatment, as demonstrated by the immunoblotting

analysis (Fig. 5K). This suggests that the PERK/eIF2 pathway is activated early during INH

exposure, further connecting ER stress to the observed rise in ROS levels.

Collectively, these findings indicate that INH treatment primarily triggers cytosolic ROS

induction related to ER stress. This induction is followed by a subsequent rise in mitochondrial

ROS, which may result from the diffusion of ROS from the cytosol or mitochondrial complex

dysfunction, ultimately altering glutathione oxidation.

Nrf2 activation is the key antioxidant defense in response to INH-induced oxidative stress

In the presence of cellular ROS, transcription factor Nrf2 (also known as nuclear factor

erythroid 2-related factor 2, NFE2L2) can be activated and translocated into the nucleus to enhance

the expression of multiple antioxidant genes [25]. To assess the activation of Nrf2 triggered by

INH, we examined the expression of multiple reported Nrf2 targets using quantitative PCR

(qPCR), including *HMOX1*, *SLC7A11*, *GCLC*, and *NQO1* from 24 h to 72 h (Fig. 6 A-D). We

observed the upregulation of mRNA expression of these Nrf2 targets in INH 32.5 mM treatment

from 24 h and remarkably increased at 72 h in both 17 mM and 32.5 mM treatment groups.

Furthermore, we observed that Nrf2 was increasingly localized to the nucleus 24 h after exposure

to INH, indicating enhanced activation of Nrf2 signaling pathways at this time point (Fig. 6E).

Fig. 6. Oxidative stress triggered by INH exposure induces Nrf2 activation. (A-D) Normalized

mRNA expression levels of Nrf2-targeted genes in HepG2 cells treated with 0 mM, 17 mM, and

32.5 mM INH for 24, 48, and 72 h. (A) *HMOX1*, (B) *SLC7A11*, (C) *GCLC,* and (D) *NQO1* mRNA

expression measured using qPCR. (E) Representative immunofluorescence images of HepG2 cells

treated with 0 mM, 17 mM, and 32.5 mM INH for 24 h, stained for Nrf2 (red) and Hoechst (blue).

Merged images show nuclear localization of Nrf2 in INH-treated groups. Scale bar: 10 µm. The

graph on the right quantifies the mean intensity of nuclear Nrf2 fluorescence. (F) Normalized

CellROX fluorescence intensity in WT and Nrf2 KO HepG2 cells treated with 0 mM, 17 mM, and

32.5 mM INH for 72 h. (G) Quantification of mitochondrial ROS levels in WT and Nrf2 KO cells

using MitoSOX fluorescence intensity after 72 h INH treatment. (H) *HMOX1* mRNA expression

levels in WT and Nrf2 KO cells measured by qPCR. Data are shown as means ± SD. Statistical

differences between groups were determined using one-way ANOVA with Tukey's post-test for

multiple comparisons; n.s., not significant, *P-value < 0.05, **P-value < 0.01, ***P-value < 0.001,

****P-value < 0.0001.

To further validate the role of Nrf2 in INH-induced oxidative stress, we generated Nrf2-

KO cells using the CRISPR/Cas9 system, as shown in Supplementary Fig. S5. In these Nrf2-KO

cells, we observed a significant increase in both cytosolic ROS (Fig. 6F) and mitochondrial ROS

(Fig. 6G) when exposed to INH for 72 h compared to WT groups. This increase in ROS levels

indicates heightened oxidative stress due to the absence of Nrf2. Moreover, we found that Nrf2

was essential for the upregulation of the antioxidant gene *HMOX1* in cells exposed to INH (Fig.

6H), further emphasizing the role of Nrf2 in regulating the cellular response to oxidative stress.

These findings demonstrate that Nrf2 signaling is crucial for cellular adaptation to INH-

induced oxidative stress, highlighting its role in upregulating key antioxidant defenses to

counteract ROS damage.

Multi-omics unveils the critical role of Nrf2 in antioxidation defense and cell survival under

INH exposure

We investigated the effects of Nrf2 KO on HepG2 cells treated with INH for 72 h. The Nrf2 KO cells demonstrated increased susceptibility to INH, exhibiting lower cell viability compared to WT HepG2 cells under the same INH concentration (Supplementary Fig. S6, Supplementary Fig. S7A). This data suggests that Nrf2 plays a protective role in mitigating INH-induced cytotoxicity.

We further investigated the molecular profile differences between Nrf2-KO and WT HepG2 cells under 72-hour INH treatment, finding substantial differences in their proteomics profiles, with 1745 differential proteins identified between the two cell types (Supplementary Fig. S7, Supplementary Fig. S7B, Supplementary Fig. S7C). Pathway enrichment analysis on these proteins revealed a disturbance in the apoptosis pathway, suggesting that KO cells were more vulnerable to INH treatments than WT cells (Supplementary Table S4). Furthermore, we observed perturbation in biological processes associated with mitochondrial function, including those related to the TCA cycle ("Citrate cycle (TCA cycle)", "Pyruvate metabolism," "Oxidative phosphorylation" and "Glycolysis / Gluconeogenesis") and β-oxidation ("Fatty acid metabolism," "Fatty acid degradation," and "PPAR signaling pathway"). In addition, pathways related to ER stress ("Protein processing in endoplasmic reticulum" and "Protein export") were shown to be changed (Supplementary Table S4). The GO:CC terms relevant to mitochondrial respiratory complexes and ER were found to be significant (Supplementary Table S5). These disturbances indicated the mitochondrial dysfunction and ER stress induced by INH treatment, which could be more severe in Nrf2-KO HepG2 cells. Moreover, compared to WT cells, KO-cells exhibited perturbations in multiple pathways related to antioxidative response. They included "Glutathione

metabolism," "Cysteine and methionine metabolism," "Ferroptosis," "HIF-1 signaling pathway,"

"Selenocompound metabolism," "Pentose phosphate pathway," "Sulfur metabolism," "Ascorbate

and aldarate metabolism," and "Glycine, serine and threonine metabolism" (Supplementary Table

S4). Noticeably, except for the three former pathways, the six latter pathways were not

significantly altered when comparing INH-treated WT cells with untreated WT cells. Given the

loss of protection against oxidative stress from the Nrf2 signaling system, the newly founded

significant pathways might represent alternative mechanisms that help KO cells manage oxidative

stress. However, these mechanisms are insufficient to maintain cellular homeostasis. Besides, we

also observed significant alterations in pathways related to endogenous cellular metabolism, i.e.,

amino acids metabolism, carbon metabolism, and lipid metabolism (Supplementary Table S4).

We captured a subtle difference in the metabolome between KO and WT cells under the

same INH treatments with only 9 significantly differential metabolites (Supplementary Fig. S7D-

F). The pathway analysis results showed metabolic perturbation related to redox balance, i.e.,

"Glutathione metabolism." Indeed, we found that, under INH treatments, abundances of GSH and

GSSG decreased in KO cells compared to WT cells (Supplementary Fig. S7F). We observed

substantial alterations in the lipid profiles of KO compared to WT cells upon INH treatments, with

409 significantly differential lipids between these two types of cells (Supplementary Fig. S7G-H).

It should be noted that basal levels of lipids in KO cells were substantially different from those of

WT cells (Supplementary Fig. S7I). The levels of complex lipids such as glycerophospholipids,

sphingolipids, and ceramide were significantly lower. The levels of lipid species involved in lipid

droplet formation, regardless of INH treatment, were higher than WT. However, no significant

differences were observed in the levels of droplet-related lipids between KO-cell groups. In

agreement with our lipidomics data (Supplementary Fig. S7I), we observed similar patterns with

insignificant changes in the number, size, and fluorescence intensity of lipid droplets between KO-

cell groups (Supplementary Fig. S7J).

Collectively, our findings emphasized the central role of Nrf2 in protecting HepG2 cells

against INH-induced stress. Additionally, we suggested alternative mechanisms that may be

involved in cell defense against INH-induced oxidative stress besides Nrf2.

Discussion

In this study, we extensively elucidated the mechanism by which INH induces

hepatotoxicity in liver cells, expanding beyond the previously understood focus on mitochondrial

oxidative stress and dysfunction (Fig. 7). Our findings reveal that INH-induced cellular ROS,

associated with ER stress, occurs prior to the increase in mitochondrial ROS and subsequent

mitochondrial dysfunction. Furthermore, prolonged INH exposure results in mitochondrial

dysfunction related to the impairment of mitochondrial complexes I-IV and a proton leak in the

mitochondrial membrane, without affecting mitochondrial ATP synthase. Additionally, INH

prompted metabolic reprogramming in response to disruptions in the TCA cycle and beta-

oxidation, leading to the accumulation of fatty acids and polyunsaturated fatty acids.

Fig. 7. Mechanisms of INH-induced cellular oxidative stress and mitochondrial dysfunction

in HepG2 cells

INH exposure initially induced cytosolic ROS rise, causing antioxidant responses including

glutathione system (GSH/GSSG) and Nrf2-targeted antioxidant genes (*HMOX1*, *SLC7A11*,

GCLC, *NQO1*), enhancing cellular defense mechanisms. Mitochondrial ROS has not risen during

this period, and mitochondrial homeostasis is adequately maintained. By inhibiting ER stress

during this stage using TUDCA, INH-induced cytosolic ROS can be eliminated. However,

prolonged INH exposure results in antioxidant defense becoming overwhelmed, resulting in

increased mitochondrial ROS in combination with mitochondrial complex I-IV defects and

uncoupling mitochondrial membrane proton leaks that lead to mitochondrial Ca^{2+} overload, proton

accumulation, mitochondrial depolarization, impaired ATP production, and insufficient TCA

cycle. These changes resulted in mitochondrial fission, mitophagy and ultimately, apoptosis, as

indicated by the activation of caspase-3 and caspase-9 and BAX translocation. In addition, there

are elevated levels of complex lipids with polyunsaturated fatty acyl chains, fatty acids, and lipid

droplet formation, indicating a global lipid alteration of the liver cell due to prolonged INH

exposure.

We identified defects in mitochondrial complexes I-IV. On the other hand, an increased

expression of ATP synthase (mitochondrial complex V) was found to be a compensatory

mechanism by which the cell attempts to maintain ATP production. These results corroborate a

previous study indicating that hydrazine, a major reactive metabolite of INH, can inhibit

mitochondrial complex II activity [8]. In addition, INH induced a mitochondrial proton leak,

supporting the "uncoupling to survive" hypothesis [26], which should be further verified. This

mechanism generally exists to minimize oxidative damage by reducing the proton motive force

and mitochondrial superoxide production [27]. Our proteomics data also revealed an increase in

the protein abundance of aspartate-glutamate carriers (SLC25A12, SLC25A13), dicarboxylate

carriers (SLC25A4, SLC25A5), and the phosphate carrier (SLC25A3) following INH treatment.

Evidence supports that these members of mitochondrial solute carriers can facilitate mitochondrial

proton leak [28].

Mitochondrial oxidative stress and dysfunction have long been regarded as the primary

factors in INH-induced hepatotoxicity [7]. Our study provides evidence that mitochondrial

superoxide levels increase concurrently with mitochondrial dysfunction after prolonged INH

exposure. Interestingly, cellular ROS levels, but not the mitochondrial ROS levels, and the

activation of the Nrf2 pathway were observed as early responses to INH treatment. In line with

our findings, Chowdhury A *et al.* reported that INH 50 mg/kg alone induced hepatic oxidative

stress in mice. However, they did not observe an increase in oxidative stress in the mitochondria

of these mice [10]. Using TUDCA, an ER stress inhibitor, we found that ER stress and ER-

associated ROS are the primary sources of cellular ROS elevation in early INH exposure time.

However, TUDCA functions as a chemical chaperone, simultaneously mitigating ER protein

misfolding and ER redox imbalance [29]. Inside the ER, ROS are mainly produced from the

oxidative protein folding process, which is mediated by protein disulfide isomerase (PDIs) and ER

oxidoreductin-1 (Ero1) proteins [30], and NOX4–a member of the NADPH oxidase (NOX) family

[31]. Regarding the activation of PERK/eIF2 signaling, we also observed the upregulation of ER

oxidoreductin-1 alpha (ERO1A) in early INH treatment. ERO1A generates hydrogen peroxide

(H_2O_2) as a byproduct while reoxidizing PDI, transferring disulfide bonds (-S-S-) to newly

synthesized proteins, which are essential for their correct folding. Moreover, hydrazine is a potent

reducing agent that can donate its electrons to the disulfide bonds of proteins or GSSG, converting

them into two free thiol groups (-SH) [32, 33]. Thus, INH or hydrazine may directly affect the ER

protein unfolding/misfolding process and induce ER redox imbalance. Under ER stress, the

accumulation of unfolded or misfolded proteins increases protein folding load, leading to excessive

ROS (H_2O_2) production as a byproduct of disulfide bond formation. This process also depletes

GSH, which is crucial to refolding misfolded proteins. Consequently, the overproduction of ROS

and GSH depletion leads to oxidative stress [34, 35].

Upon INH exposure, the TCA cycle was impaired, partially reflected by the depletion of

its intermediates. The mechanism of how INH impairs the TCA cycle is possibly due to the

reaction between INH and keto acids, key intermediates of the TCA cycle [36]. In addition, a

previous study showed that hydrazine can induce the depletion of TCA cycle intermediates [37].

Our data also suggested the disruption of β-oxidation, another critical process for maintaining

cellular energy homeostasis. One potential mechanism for this disruption is the suppression of the

carnitine shuttle process, which plays a vital role in transporting long-chain fatty acids to the

mitochondrial matrix [38]. Specifically, the later step of transferring acylcarnitines to the

mitochondrial matrix may be insufficient, as indicated by increments in the abundance of

acylcarnitines with longer chains and fatty acids, as well as a decrease in the level of carnitine. The

impairments of mitochondrial TCA cycle, β-oxidation, and oxidative phosphorylation may lead to

energy homeostasis disorder.

Interestingly, the current study highlights several adaptations to the INH-induced energy

crisis, implicating potential intervention strategies to prevent and ameliorate hepatotoxicity. We

observed increased abundances of complex V protein subunits, TCA cycle enzymes, and β-

oxidation enzymes, suggesting a compensatory effort to enhance ATP production as well as

support energy and biosynthetic demands. Additionally, impairment of β-oxidation might lead to

a deficiency in acetyl-CoA. Therefore, the increase in the conversion of pyruvate to acetyl-CoA

may ensure consistent supply for the TCA cycle and lipid biosynthesis. Indeed, the expression

levels of multiple enzymes involved in pyruvate metabolism were upregulated. Moreover, the

abundances of CP1A (carnitine palmitoyltransferase 1A) and CPT2 (carnitine palmitoyltransferase

2), two main enzymes involved in carnitine-based fatty acids transport [39], were upregulated.

This might be an effort to rescue the carnitine shuttle pathway to facilitate the β-oxidation process.

Besides, lipid synthesis and turnover maintain membrane integrity and produce crucial signaling

molecules. Together, these changes demonstrate a sophisticated cellular adaptation, enabling

compensation for impaired mitochondrial function and ensuring cell survival and functionality.

Importantly, our study underscores the pivotal role of Nrf2 signaling in antioxidation

defense and cellular survival in HepG2 cells subjected to INH treatment. Notably, cells lacking

Nrf2 are particularly vulnerable to INH-induced oxidative stress, highlighting the critical

importance of Nrf2 in cellular defense mechanisms. We identified alternative pathways involved

in cellular antioxidant defense in Nrf2 KO conditions, such as selenocompound metabolism, the

HIF-1 signaling pathway, and the pentose phosphate pathway. Selenocompound metabolism is

critical for counteracting oxidative stress through selenoproteins like glutathione peroxidases

(GPxs) and thioredoxin reductases (TrxRs), which use selenium to neutralize ROS, thus protecting

cells from damage. Hypoxia-inducible factor-1 (HIF-1) regulates genes linked to oxidative stress,

inflammation, and cell death [40], and is associated with liver toxicity in various contexts,

including drug-induced liver injury [41]. A case-control study indicates that the HIF-1α gene is

linked to a higher risk of anti-tuberculosis drug-induced liver injury (ATB-DILI) [42], and it has

been shown that anti-TB drug-induced hepatocyte ferroptosis is controlled by the HIF-

1α/SLC7A11/GPx4 signaling pathway [43]. Given the crucial roles of selenoproteins and HIF-1

signaling in antioxidant defenses, there is limited evidence on how these factors contribute to the

progression of ATB-DILI, particularly INH-DILI.

Nrf2 plays a significant role in regulating lipid metabolism, especially in the context of

GSH synthesis in the liver. According to a study by Asantewaa et al. (2024), GSH synthesis in the

liver is essential for maintaining lipid abundance [44]. The repression of Nrf2 leads to increased

lipid production and storage. In our study, Nrf2 KO cells displayed higher levels of lipid droplets

and overall lipid storage compared to WT cells, regardless of INH treatments. Our data are

consistent with the previous finding demonstrating that the repression of lipid storage and

triglyceride levels is contingent on Nrf2 activation [44].

There are a few limitations and grounds for future studies. Even though widely used for

toxicological investigation, molecular alterations of HepG2 cells may not fully represent the

responses of normal liver cells. Additionally, our study helped uncover some potentially important

questions regarding INH-induced toxicity for future investigations: (1) The detailed mechanism

underlying INH-induced ER stress and the involvement of NOX4 in this molecular event, (2) the

exact mechanisms by which INH or its metabolites impair mitochondrial complexes and the role

of uncoupling proteins in mitochondrial dysfunction, and (3) the crosstalk between GSH and Nrf2

signaling in modulating cellular lipidome upon INH treatment. Future research should also validate

our findings in animal or more physiologically relevant models, such as *in vitro* 3D hepatic

microtissues, to better understand the *in vivo* relevance. Investigating the interactions between INH

metabolites and mitochondrial complexes and exploring therapeutic strategies targeting Nrf2 and

alternative antioxidant pathways could provide deeper insights and potential host-directed

preventative therapy for INH-induced liver injury.

In conclusion, this study provides significant insights into the mechanisms of INH-induced

hepatotoxicity, highlighting the complex interplay between ER stress, oxidative stress,

mitochondrial dysfunction, and metabolic reprogramming. We demonstrated that ER stress is the

primary source of ROS generation, initiating cellular oxidative stress in early INH exposure

followed by mitochondrial oxidative stress in the later stages. The anti-ER-stress treatment can

suppress the INH-induced ROS formation and protect cells against oxidative stress, which is the

potential for developing preventative and alleviative therapy towards INH-induced liver injury.

Our findings also emphasize the central role of Nrf2 signaling in the antioxidant response against

INH-derived oxidative stress.

Acknowledgment

Fig.7 was created using Biorender.com

Funding

This study was supported by the National Research Foundation of Korea (NRF) grant

funded by the Korean government (MSIT) (grant No. 2018R1A5A2021242). The funding

organizations were not involved in the study design, data acquisition, data analysis, data

interpretation, or the content presented in the manuscript.

CRediT authorship contribution statement

Truong Thi My Nhung: Data curation, Methodology, Investigation, Formal analysis,

Visualization, Validation, Writing – Original Draft, Writing – Review & Editing. **Nguyen Ky**

Phat: Data curation, Methodology, Investigation, Formal analysis, Visualization, Validation,

Writing – Original Draft, Writing – Review & Editing. **Trinh Tam Anh**: Methodology, Formal

analysis, Visualization, Validation, Writing – Review & Editing. **Tran Diem Nghi**: Methodology,

Formal analysis, Visualization, Validation, Writing – Review & Editing. **Nguyen Quang Thu**:

Data curation, Formal analysis, Visualization, Validation, Writing – Review & Editing. **Ara Lee**:

Methodology, Formal analysis, Software, Writing – Review & Editing. **Nguyen Tran Nam Tien**:

Methodology, Formal analysis, Software, Writing – Review & Editing. **Nguyen Ky Anh**:

Methodology, Formal analysis, Writing – Review & Editing. **Kimoon Kim:** Supervision,

Methodology, Validation, Resources, Writing – Review & Editing. **Duc Ninh Nguyen:**

Methodology, Validation, Writing – Review & Editing. **Dong Hyun Kim**: Supervision,

Resources, Writing – Review & Editing, Funding acquisition. **Sang Ki Park**: Methodology,

Investigation, Resources, Supervision, Writing – Review & Editing, Funding acquisition. **Nguyen**

Phuoc Long: Conceptualization, Methodology, Investigation, Formal analysis, Validation,

Resources, Supervision, Writing – Original draft, Writing – Review & Editing, Funding

acquisition.

Declaration of competing interest

The authors declare no known competing financial interests that may influence the results

of this work.

Reference

1. World Health Organization Global tuberculosis report 2023. World Health Organization. 2023.

2. World Health Organization, WHO consolidated guidelines on tuberculosis: Module 4: Treatment - Drug-susceptible tuberculosis treatment. 2022: Geneva.

3. Forget E.J. and Menzies D. Adverse reactions to first-line antituberculosis drugs. Expert Opin Drug Saf. 2006;5(2):231-49.

4. Nahid P., Dorman S.E., Alipanah N., Barry P.M., Brozek J.L., Cattamanchi A., et al. Official American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America Clinical Practice Guidelines: Treatment of Drug-Susceptible Tuberculosis. Clin Infect Dis. 2016;63(7):e147-e95.

5. Horsburgh C.R., Jr., Barry C.E., 3rd, and Lange C. Treatment of Tuberculosis. N Engl J Med. 2015;373(22):2149-60.

6. Nolan C.M., Goldberg S.V., and Buskin S.E. Hepatotoxicity associated with isoniazid preventive therapy: a 7-year survey from a public health tuberculosis clinic. JAMA. 1999;281(11):1014-8.

7. Boelsterli U.A. and Lee K.K. Mechanisms of isoniazid-induced idiosyncratic liver injury: emerging role of mitochondrial stress. J Gastroenterol Hepatol. 2014;29(4):678-87.

8. Lee K.K., Fujimoto K., Zhang C., Schwall C.T., Alder N.N., Pinkert C.A., et al. Isoniazid-induced cell death is precipitated by underlying mitochondrial complex I dysfunction in mouse hepatocytes. Free Radic Biol Med. 2013;65:584-94.

9. Ahadpour M., Eskandari M.R., Mashayekhi V., Haj Mohammad Ebrahim Tehrani K., Jafarian I., Naserzadeh P., et al. Mitochondrial oxidative stress and dysfunction induced by isoniazid: study on isolated rat liver and brain mitochondria. Drug Chem Toxicol. 2016;39(2):224-32.

10. Chowdhury A., Santra A., Bhattacharjee K., Ghatak S., Saha D.R., and Dhali G.K. Mitochondrial oxidative stress and permeability transition in isoniazid and rifampicin induced liver injury in mice. J Hepatol. 2006;45(1):117-26.

11. Zhang T., Ikejima T., Li L., Wu R., Yuan X., Zhao J., et al. Impairment of Mitochondrial Biogenesis and Dynamics Involved in Isoniazid-Induced Apoptosis of HepG2 Cells Was Alleviated by p38 MAPK Pathway. Front Pharmacol. 2017;8:753.

12. Schwab C.E. and Tuschl H. In vitro studies on the toxicity of isoniazid in different cell lines. Hum Exp Toxicol. 2003;22(11):607-15.

13. Shen C., Meng Q., Zhang G., and Hu W. Rifampicin exacerbates isoniazid-induced toxicity in human but not in rat hepatocytes in tissue-like cultures. Br J Pharmacol. 2008;153(4):784-91.

14. Cullinan S.B., Gordan J.D., Jin J., Harper J.W., and Diehl J.A. The Keap1-BTB protein is an adaptor that bridges Nrf2 to a Cul3-based E3 ligase: oxidative stress sensing by a Cul3-Keap1 ligase. Mol Cell Biol. 2004;24(19):8477-86.

15. Kobayashi A., Kang M.I., Okawa H., Ohtsuji M., Zenke Y., Chiba T., et al. Oxidative stress sensor Keap1 functions as an adaptor for Cul3-based E3 ligase to regulate proteasomal degradation of Nrf2. Mol Cell Biol. 2004;24(16):7130-9.

16. Zhang D.D. and Hannink M. Distinct cysteine residues in Keap1 are required for Keap1-dependent ubiquitination of Nrf2 and for stabilization of Nrf2 by chemopreventive agents and oxidative stress. Mol Cell Biol. 2003;23(22):8137-51.

17. Sasaki H., Sato H., Kuriyama-Matsumura K., Sato K., Maebara K., Wang H., et al.
Electrophile response element-mediated induction of the cystine/glutamate exchange
transporter gene expression. J Biol Chem. 2002;277(47):44765-71.
18. Jia Z.L., Cen J., Wang J.B., Zhang F., Xia Q., Wang X., et al. Mechanism of isoniazid-
induced hepatotoxicity in zebrafish larvae: Activation of ROS-mediated ERS, apoptosis
and the Nrf2 pathway. Chemosphere. 2019;227:541-50.
19. Ritchie M.E., Phipson B., Wu D., Hu Y., Law C.W., Shi W., et al. limma powers
differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Res. 2015;43(7):e47.
20. Liu P., Ewald J., Pang Z., Legrand E., Jeon Y.S., Sangiovanni J., et al. ExpressAnalyst: A
unified platform for RNA-sequencing analysis in non-model species. Nat Commun.
2023;14(1):2995.
21. Pang Z., Lu Y., Zhou G., Hui F., Xu L., Viau C., et al. MetaboAnalyst 6.0: towards a
unified platform for metabolomics data processing, analysis and interpretation. Nucleic
Acids Res. 2024;52(W1):W398-W406.
22. Ramachandran A., Visschers R.G.J., Duan L., Akakpo J.Y., and Jaeschke H. Mitochondrial
dysfunction as a mechanism of drug-induced hepatotoxicity: current understanding and
future perspectives. J Clin Transl Res. 2018;4(1):75-100.
23. Bhadauria S., Mishra R., Kanchan R., Tripathi C., Srivastava A., Tiwari A., et al. Isoniazid-
induced apoptosis in HepG2 cells: generation of oxidative stress and Bcl-2 down-
regulation. Toxicol Mech Methods. 2010;20(5):242-51.
24. Romani P., Nirchio N., Arboit M., Barbieri V., Tosi A., Michielin F., et al. Mitochondrial
fission links ECM mechanotransduction to metabolic redox homeostasis and metastatic
chemotherapy resistance. Nat Cell Biol. 2022;24(2):168-80.
25. Tonelli C., Chio I.I.C., and Tuveson D.A. Transcriptional Regulation by Nrf2. Antioxid
Redox Signal. 2018;29(17):1727-45.
26. Brand M.D. Uncoupling to survive? The role of mitochondrial inefficiency in ageing. Exp
Gerontol. 2000;35(6-7):811-20.
27. Lambert A.J. and Brand M.D. Superoxide production by NADH:ubiquinone
oxidoreductase (complex I) depends on the pH gradient across the mitochondrial inner
membrane. Biochem J. 2004;382(Pt 2):511-7.
28. Skulachev V.P. Uncoupling: new approaches to an old problem of bioenergetics. Biochim
Biophys Acta. 1998;1363(2):100-24.
29. Abdel-Ghaffar A., Elhossary G.G., Mahmoud A.M., Elshazly A.H.M., Hassanin O.A.,
Saleh A., et al. Potential prophylactic effect of chemical chaperones for alleviation of
endoplasmic reticulum stress in experimental diabetic cataract. Bulletin of the National
Research Centre. 2019;43(1):71.
30. Ramming T., Okumura M., Kanemura S., Baday S., Birk J., Moes S., et al. A PDI-catalyzed
thiol-disulfide switch regulates the production of hydrogen peroxide by human Ero1. Free
Radic Biol Med. 2015;83:361-72.
31. Laurindo F.R., Araujo T.L., and Abrahao T.B. Nox NADPH oxidases and the endoplasmic
reticulum. Antioxid Redox Signal. 2014;20(17):2755-75.
32. Furst A., Berlo R.C., and Hooton S. Hydrazine as a Reducing Agent for Organic
Compounds (Catalytic Hydrazine Reductions). Chemical Reviews. 1965;65(1):51-68.
33. Yew W.W., Chang K.C., and Chan D.P. Oxidative Stress and First-Line Antituberculosis
Drug-Induced Hepatotoxicity. Antimicrob Agents Chemother. 2018;62(8).

34. Cao S.S. and Kaufman R.J. Endoplasmic reticulum stress and oxidative stress in cell fate
decision and human disease. Antioxid Redox Signal. 2014;21(3):396-413.
35. Chen X., Shi C., He M., Xiong S., and Xia X. Endoplasmic reticulum stress: molecular
mechanism and therapeutic targets. Signal Transduct Target Ther. 2023;8(1):352.
36. Li F., Miao Y., Zhang L., Neuenswander S.A., Douglas J.T., and Ma X. Metabolomic
analysis reveals novel isoniazid metabolites and hydrazones in human urine. Drug Metab
Pharmacokinet. 2011;26(6):569-76.
37. Bollard M.E., Keun H.C., Beckonert O., Ebbels T.M., Antti H., Nicholls A.W., et al.
Comparative metabonomics of differential hydrazine toxicity in the rat and mouse. Toxicol
Appl Pharmacol. 2005;204(2):135-51.
38. Longo N., Frigeni M., and Pasquali M. Carnitine transport and fatty acid oxidation.
Biochim Biophys Acta. 2016;1863(10):2422-35.
39. Knottnerus S.J.G., Bleeker J.C., Wust R.C.I., Ferdinandusse S., L I.J., Wijburg F.A., et al.
Disorders of mitochondrial long-chain fatty acid oxidation and the carnitine shuttle. Rev
Endocr Metab Disord. 2018;19(1):93-106.
40. Kaminsky-Kolesnikov Y., Rauchbach E., Abu-Halaka D., Hahn M., Garcia-Ruiz C.,
Fernandez-Checa J.C., et al. Cholesterol Induces Nrf-2- and HIF-1alpha-Dependent
Hepatocyte Proliferation and Liver Regeneration to Ameliorate Bile Acid Toxicity in
Mouse Models of NASH and Fibrosis. Oxid Med Cell Longev. 2020;2020:5393761.
41. Chu Q., Gu X., Zheng Q., and Zhu H. Regulatory mechanism of HIF-1alpha and its role in
liver diseases: a narrative review. Ann Transl Med. 2022;10(2):109.
42. Chong Y., Zhu H., Ren Q., Ma X., and Feng F. Interaction between the HIF-1alpha gene
rs1957757 polymorphism and CpG island methylation in the promoter region is associated
with the risk of anti-tuberculosis drug-induced liver injury in humans: A case-control study.
J Clin Pharm Ther. 2022;47(7):948-55.
43. Liu Y., Chen W., Cen Y., Zhao X., Chen Z., Liang Y., et al. Hepatocyte ferroptosis
contributes to anti-tuberculosis drug-induced liver injury: Involvement of the HIF-
1alpha/SLC7A11/GPx4 axis. Chem Biol Interact. 2023;376:110439.
44. Asantewaa G., Tuttle E.T., Ward N.P., Kang Y.P., Kim Y., Kavanagh M.E., et al.
Glutathione synthesis in the mouse liver supports lipid abundance through NRF2
repression. Nat Commun. 2024;15(1):6152.