Scalable and universal prediction of cellular phenotypes

Yuge Ji[1,2]†, Alejandro Tejada-Lapuerta[1,3]†, Niklas A. Schmacke[1], Zihe Zheng[1], Xinyue Zhang[3], Simrah Khan[1,4], Ina Rothenaigner[5], Juliane Tschuck[5], Kamyar Hadian[5], Fabian J. Theis[1,2,3]*

[1]Institute of Computational Biology, Helmholtz Munich, Germany.

[2]TUM School of Life Sciences Weihenstephan, Technical University of Munich, Germany.

[3]TUM School of Computing, Information and Technology, Technical University of Munich, Munich.

[4]Ludwig Maximilians University, Faculty of Biology.

[5]Research Unit Signaling and Translation, Helmholtz Munich, Neuherberg, Germany.

† These authors contributed equally to this work

*Corresponding author. Email: fabian.theis@helmholtz-munich.de

Abstract: Biological systems can be understood by perturbing individual components and studying the system's response. Cell biology experiments are defined by the applied treatment, cellular state, and the assayed phenotype. Given the vast number of possible combinations, testing every scenario is impractical. We present Prophet, a transformer-based computational model for cellular phenotype prediction. Prophet learns a representation of the cell biology experiment space, enabling it to predict the outcomes of untested small molecule or genetic perturbations in new cellular contexts across diverse phenotypes including gene expression, cell viability, and cell morphology. Its scalable architecture facilitates training across independent assays, using transfer learning to enhance performance across phenotypes. In vitro validation shows Prophet's potential to guide experimental design, making it a valuable tool for accelerating biological discovery.

Introduction

High-throughput screening and multiplexed phenotypic readouts have revolutionized biological experimentation. They enable simultaneous testing of thousands of compounds across numerous biological contexts, accelerating the pace of discovery. For instance, small molecule and CRISPR screens have been conducted to systematically interrogate the biological mechanisms of cancer resistance (1, 2), identify new drug targets (3, 4), capture synergistic interactions (5), and deconvolve drug mechanisms of action (6–9). While statistical and AI-based analyses have facilitated discoveries in each screen, a key limitation of past models is the lack of integration across screens (10, 11), despite considerable overlap in experiments—for instance, the same cell line treated with different compounds or genetic perturbations, or the same experiment measured with different readouts. Learning across all of these screens could provide notable increases in the efficiency of identifying effective and cell-context specific compounds while simultaneously making findings more robust across technical variation (12).

Given the immense combinatorial space inherent to biological experimentation, accurate in silico modeling is needed to guide the selection of relevant future experiments and reduce the number of early-stage experiments needed to yield new biological insights(13). While previous methods offer tailored solutions for specific experimental contexts (11, 14–18), we propose a general approach to treatment response prediction which can fully leverage the vast data resources now available. Inspired by the success of transformers in solving complex problems across various domains by modeling data as a set of tokens to be transformed towards an outcome (19–21), we develop a novel approach to cellular response prediction: modeling the experimental space using set-based representations. We universally decompose each experiment into a unique combination of three fundamental elements—the cellular state, the treatments being performed, and the intended phenotypic readout. This approach forms the basis for our model, Prophet (Predictor of phenotypes), a transformer-based regression model that learns the relationships between these factors. Prophet's framework enables it to be pretrained on 4.7 million experiments across a broad spectrum of phenotypes from multiple independent datasets, producing accurate predictions that empower downstream applications, such as nominating additional chemotherapeutic drugs for BRAF-mutant melanoma and reducing the number of experiments needed for viability screens by at least 60x. Together, its scaling behavior and strong predictive capacity establish Prophet as a robust general model for treatment response in biological systems.

Results

A unified framework for experimental cell biology

We consider the outcome of a cell biology experiment as the composite of three essential components: the cell state the experiment is performed in, the applied treatment, and the phenotypic readout used to measure the effect of the perturbation (Fig. 1a, b). The cellular state represents the basal state before treatment. For instance, most drug screens are performed on cell lines that we characterize as the cell state. The treatment is the specific intervention or perturbation applied to the cells during the experiment. Commonly, these are genetic perturbations (e.g. CRISPR/Cas9 knockouts, CRISPRi/a) or drug perturbations with small molecules. Finally, the phenotypic readout is the measurable outcome used to assess the effect of the applied treatment on the cell state; for example a morphological measurement such as cell size, or the IC50 of a compound. To model the relationship between these elements in a

generalizable way, we developed Prophet, a supervised transformer-based deep learning model (Fig. 1b, c). To train it, we collected 9 perturbational datasets to create the largest compendium of publicly available screening datasets to date: SCORE(3, 22), LINCS(23), JUMP(8), GDSC(3, 22, 24), GDSC2(25), CTRP(26), PRISM(1), Horlbeck et al.(27) and Saunders et al.(28) (fig. S1, Table S1). For example, we consider one measurement of the IC50 of the drug isoniazid in HeLa cells a single experiment that is passed to Prophet as one datapoint. Measuring cell morphology after the same treatment in the same cell line would be a different data point, and so would measuring the IC50 of the same drug in another cell line (Fig. 1c). We hypothesized that when trained across a sufficient amount of experimental data, a single model could be employed to make predictions for any compound or genetic treatment, and for any cell line or cell type (Fig. 1).

Prophet's architecture consists of 8 transformer encoder units (29, 30) with 8 attention heads per layer and a feed-forward network with a hidden dimensionality of 1,024 to generate a 512-dimensional embedding of each experiment (Fig. 1c, fig. S2). Prophet leverages existing knowledge of cellular states and treatments by projecting prior knowledge-based representations into a common token space using neural networks as tokenizers. The readout representations are modeled as learnable embeddings, directly projected in the token space. Through self-attention, Prophet effectively captures relationships between treatments, cellular states, and phenotypic outcomes, facilitating nuanced learning and prediction. The output of the transformer is fed into a feed-forward network that predicts the phenotype of interest. Prophet is trained end-to-end using supervised learning to minimize mean square error (MSE) between the predicted and measured phenotypes.

To make prior knowledge available for training Prophet, we create representation vectors for each axis of the experimental space, encoding the cellular environment the experiment is conducted in, and the treatments that are applied, to serve as input to the model.

To represent cell state, as most small molecule and CRISPR screens are performed in cell lines, we use bulk RNA sequencing data from the Cancer Cell Line Encyclopedia (CCLE) (31), which covers 1,406 cell lines. For in vivo treatments, we assume each cell type in a tissue represents one cell state and encode them using their RNA expression profile after pseudobulk aggregation.

For treatments, we use two different methods of generating embeddings depending on whether the treatment is a small molecule or genetic perturbation. For small molecules, we create descriptive embeddings by concatenating their RDKit2D and Morgan fingerprints (32, 33). For genetic perturbations, we build a vector that represents the target gene as the concatenation of a vector capturing its transcriptomic variability and a vector that represents its genomic sequence (Methods). We obtain the transcriptomic variability vector by transposing the CCLE "cell line x gene expression" matrix and finding a low dimensional representation of the genes using PCA. We derive the genomic sequence vector from Enformer's embeddings (34) for the entire coding sequence of the gene.

Since there is no prior knowledge suitable to represent an embedding of most phenotypes, we model them as a learnable embedding through one hot encodings and a linear layer. However, if there exists a suitable prior knowledge-based representation of the phenotype, it can be used as input to a tokenizer network, similar to treatment and cell state representations.

Prophet is capable of jointly modeling diverse cellular phenotypes used as readouts in experiments. We train Prophet to predict cell viability, compound IC50, Cell Painting morphology features (35), mRNA transcript abundance, and cell type proportion. Its transformer-based architecture allows Prophet to handle an arbitrary number of perturbations, from singletons to any combination, so that datasets with single treatments and combinations can be modeled under the same learning framework. Additionally, utilizing prior knowledge-based embeddings enables Prophet to predict in silico outcomes for experiments performed in unseen conditions, extrapolating to unseen perturbations or cellular states.

Transfer learning improves in silico experimental prediction

We assessed Prophet's efficacy in predicting the outcome of unseen experiments. We conducted five-fold cross-validation with three validation seeds (Methods, fig. S5a) and compared the average coefficient of determination (R2) and the Spearman correlation of Prophet against three baseline architectures: random forest, a multilayer perceptron, and linear regression. For each method we report the difference in R2 and Spearman correlation to the mean estimator (Methods). The mean estimator serves as a global approximation of a treatment or cell state response based on prior experiments (fig. S5a, Methods) and is considered a robust baseline in drug response prediction (11).

We first trained specialized models on each dataset individually, with the same architecture but a smaller number of parameters to prevent overfitting (Fig. 2a, Methods). We call these models "Prophet-individual" (for hyperparameter tuning, see Supplementary Methods, Table S2). These Prophet-individual models generally outperform baseline methods across the board (Fig. 2b). Then, to benchmark Prophet's transfer learning capabilities, we conducted experiments training jointly on two datasets (Prophet-two-dataset). We combined training on GDSC2(25), a small molecule combinatorial perturbation screen reporting IC50 values, with PRISM(1), a much larger single-treatment screen relying on a CellTiterGlo viability readout. PRISM contains 8 of the 61 compounds and 76 of the 110 cell lines in GDSC2. Performance for both unseen treatments and unseen cell lines in GDSC2 improved by 3% and 28%, respectively, after joint training with PRISM. These results suggested Prophet could benefit from joint training on additional datasets and prompted us to increase the amount of datasets used during pretraining.

We then pretrained our largest Prophet model on 4.7 million experimental conditions, encompassing 26k treatments, 1,550 cell states, and three different readouts (IC50, viability and RNA expression) across seven small molecule and genetic perturbation screens (fig. S2a, b, Table S1). Afterwards, we fine-tuned the model for up to 20,000 steps with a batch size of up to 256 observations and a low learning rate (Fig. 2a, Methods, Table S3). This approach follows the same strategy as current foundation models (19, 36, 37), which are pre-trained on large amounts of data and then fine-tuned for specific datasets using the pre-trained model as a backbone.

We find that pretraining Prophet on all datasets and fine tuning for the dataset of interest consistently matches or outperforms baseline models, including Prophet-individual models, across all datasets, with baseline models frequently failing to outperform the mean estimator (Fig. 2b). Prophet demonstrates notable performance not only in predicting responses to small molecule monotherapies (GDSC, CTRP, PRISM) but also in predicting the outcomes of unseen genetic perturbations (SCORE). In addition, Prophet successfully predicts the effect of the combination of two drugs and the combination of two CRISPR perturbations (GDSC

combinations and Horlbeck, respectively). To showcase Prophet's ability to extrapolate to unseen scenarios we predict the IC50 of Mitoxantrone, a clinically approved compound indicated for multiple cancers (38, 39), which does not appear in any dataset in training. Still, Prophet's de novo predicted IC50 for Mitoxantrone reflects its efficacy in different cancer types (Fig. 2c), and

5   is corroborated by the IC50 values measured in GDSC (24) as well as other publications (40, 41). Furthermore, in addition to viability, Prophet accurately predicts diverse and complex measurements including cell morphology (JUMP) and transcriptomics (LINCS) (Fig. 2b). This generalizability serves as evidence of Prophet's ability to enable transfer learning, leveraging information across multiple screens. As an example, pretraining Prophet and fine-tuning it on

10 GDSC2 outperforms Prophet-individual by 70% in predicting unseen treatments and by 30% in predicting experiments on unseen cell states. Most notably, it also outperforms a Prophet model trained on PRISM and GDSC2 (Prophet-two-datasets) by 28% in predicting unseen treatments and by 29% in predicting experiments on unseen cell states (Fig. 2e).

Prophet's transfer learning capability is not limited to phenotypes seen during the pre-training

15 stage. We did not pre-train Prophet on any morphological measurement, but Prophet fine-tuned on JUMP outperformed both the Prophet-individual model trained only on JUMP and the baseline (Fig. 2b). We observed the same behavior with unseen RNA readouts: the mRNA abundance readout of the gene TNFRSF21 from LINCS was not included in the pre-training corpus; nonetheless, the fine-tuned version of Prophet is 25% better than the baseline and 10%

20 better than a single-dataset Prophet model trained uniquely on TNFRSF21 readouts (Fig. 2d).

The prediction of the most likely treatments to be a hit, defined as an intervention that results in a desired biological effect, can reduce the number of total screening experiments needed. Critically, Prophet's performance is not limited to accurate prediction of values close to the mean but Prophet also accurately predicts extreme values—hence, possible hits (fig. S5b). Therefore,

25 we can use Prophet to generate in silico outcomes for treatment or cell state combinations not yet tested, and use these values to suggest experiments which are most likely to produce a desired phenotype. As a case example, we take Prophet's in silico simulated viability values and compute the increase in the number of hits over randomly selecting compounds from the GDSC library. We define a hit as a compound that induces greater than 2x fold-change in viability as

30 compared to DMSO in a given cell line (Supplementary Methods). Using Prophet to predict viability for drugs in unseen cell lines in GDSC shows an improvement in R2 of .04 over the mean estimator (Fig. 2b). However, this increase translates into a 13x increase in the number of hits when performing experiments suggested by Prophet. Here, Prophet achieves an area-under the precision-recall curve (AUPRC) of .81 (Fig. 2f). When identifying hits with a more stringent

35 viability fold-change of at least 4x, the potential of Prophet is even more apparent: here, Prophet leads to a reduction in performed experiments of 60x compared to identifying hits from randomly selected compounds (Supplementary Methods). Prophet was conservative in its predictions and had no false positives among its top-ranked 1% of hits. In addition to precision-recall, we further evaluate using Spearman rank (fig. S5c). Prophet outperforms the baseline

40 models and the mean in all instances in which it also did the same using R2. This demonstrates that Prophet not only accurately predicts phenotypic values but also effectively captures the underlying ranking of responses, indicating its ability to make reliable comparisons across different experimental conditions.

To investigate how the uncertainty of Prophet varies with the number of conditions (cell states

45 and treatments) seen during training, we trained an ensemble of Prophet models. We found a

clear trend: the more treatments and cell states seen by Prophet, the higher the confidence in the predictions (Fig. 2g). Likewise, we also studied how the performance of Prophet changes as a function of the number of conditions seen during training: we trained Prophet-individual models to predict the effect of either unseen treatments or unseen cell states, including 5-80% of the data along the respective axis. The amount of data required to outperform the mean estimation varies per dataset, but as expected we observe that Prophet's predictions improve with additional training data (fig. S4, S5).

Prophet encounters challenges in outperforming the mean estimator in certain scenarios, such as predicting the effects of observed treatments in unobserved cell lines within SCORE, or forecasting the impacts of untested small molecules in known cell lines within LINCS. Further investigation revealed that the correlation between experimental replicates is a crucial metric for evaluating the predictability of their outcomes. We conducted a statistical analysis of the correlation between replicates (Table S4). This analysis revealed that, along the problematic axis, replicates of the same experimental conditions are less similar to one another than to the replicates of any other random condition. In such cases of noisy data with a strong batch effect the mean estimator is the best possible predictor.

Prophet learns a universal landscape of biological experiments

Drug screens are limited in scope, either by the number of compounds tested or the variety of cell states. Hence, they provide a limited view of the profile of a compound, encompassing just the effects in the tested cell states. Including more cell states would provide a more robust and informative view of the pharmacological profile of a drug (1). Now, with Prophet, we can examine the treatment effect in silico across hundreds of cell lines at once. We represent each small molecule—or combinations of small molecules—as a vector in which each dimension is the Prophet-predicted viability value in each cancer cell line (Fig. 3a). This vector represents an "activity-based fingerprint" for each compound. In this high-dimensional representation, compounds with similar sensitivity profiles across cancer cell lines cluster together when viewed using a 2D representation (Fig. 3b). We can draw comparisons between these shared response patterns for unbiased annotations of mechanisms of action (Fig. 3c), identification of new therapeutic candidates (Fig. 3d, e), or isolation of tissue-specific mechanisms (Fig. 3f, fig. S7a).

Compounds which target pathways of cell proliferation.

We used Prophet to generate viability predictions for 8,620 small molecule treatments in 979 cell lines (Fig. 3a). We concatenated the 979 cell line values per drug to construct an activity-based fingerprint, which together form a viability-contextualized small molecule manifold (Fig. 3b). We focus on four distinct clusters in this space (Fig. 3c, fig. S6a) containing a total of 64 compounds that target specific regulators of cell proliferation: two from the MAPK signaling cascade (containing compounds targeting MEK and EGFR), and two from the PI3K/AKT/mTOR pathway. Two of the 64 compounds have never been seen in the training data (fig. S6b). With each of the 8,620 compounds tested in an average of 365 out of 979 cell lines, Prophet accurately completed 75.2% of the cell line ⛐ drug experimental space, grouping compounds in an unbiased manner after limited or no training examples.

Prophet's identified clusters are also biologically precise: after annotating the 64 compounds (Table S7, Supplementary Methods, fig. S6a), we found only one potential misclassification: the

HSP90 inhibitor tanespimycin, which Prophet identifies as similar in effect to mTOR inhibitors(42). Additionally, we noted one misannotation: Ibrutinib, typically labeled as a BTK inhibitor, has been shown to bind to the EGFR mutant kinase in non-small cell lung cancer (NSCLC) (43, 44). We suggest that Ibrutinib also has a similar effect in other cancer types. The joint clustering could be a pure effect of chemical similarity—however, we find compounds show considerable dissimilarity in chemical space (fig. S6c, Supplementary Methods). This indicates that the experiment representations learned by Prophet are driven by the phenotypic effects of small molecule treatments rather than their chemical structures, underscoring the utility of this manifold in capturing biologically relevant information which could not be identified using structural similarity alone.

Therapeutic treatment candidates in BRAF-V600E mutant melanoma.

We wondered whether we can identify compounds which phenocopy existing, clinically approved treatments. To this end, we examined Prophet's predictions for four compounds already clinically approved for BRAF V600E-mutant melanoma. From the full 979 cell line viability space above, we create a subspace of 23 melanoma cell lines categorized as either BRAF V600E-mutant or -wild-type (Fig. 3e). 74% of these experiments were unseen in training. Such cancers respond well to various MEK and BRAF inhibitors (45–47). This subspace reveals a cluster of compounds containing three clinically approved monotherapies (46, 48, 49) and two combination therapies (47, 50) which are highlighted in bold (Fig. 3e). 45% of the other compounds in this cluster have been indicated for BRAF V600E melanoma in in vitro experiments (Table S8), suggesting that the remaining compounds are relevant therapeutic candidates.

We further examine the predicted viability per cell line. Two out of 6 wild-type cell lines were on average sensitive to compounds in this cluster, compared to 9 out of 17 BRAF V600E-mutants (Fig. S8a). Among these, resistance to MEK or RAF inhibition despite the BRAF V600E mutation has been documented in certain cell lines, such as A2058's resistance to MEK inhibitor PD0325901 (51), a behavior our model accurately predicts (fig. S8b) even though the current mechanism of resistance remains unclear.

Predicting brain and CNS-specific monotherapies.

A similar in silico generated subspace of perturbation effects in brain and central nervous system (CNS)-derived cell lines reveals a cluster of neurological compounds, enriched for putative calcium channel blockers (p-value<10-7, Fig. 3f, Supplementary Methods) which did not cluster when looking at global tissue responses (fig. S7c). Calcium signaling promotes growth and tumorigenesis of brain cancers (52). Recent studies demonstrate that calcium channel inhibition suppresses brain tumor growth, decreases the expression of oncogenes and increases expression of tumor suppressor genes, even proposing to repurpose existing drugs to treat brain cancers (53, 54).

Experimental space contextualized by RNA expression.

While the compound space detailed above was constructed using Prophet-predicted viability values, Prophet can also construct a compound space using other readouts, such as RNA expression learned from the LINCS dataset. We make predictions for BIRC5 expression after

treatment in the same set of compounds and cell lines (fig. S7d), expanding the original by 5,846 small molecules and 922 cell lines. This new space captures patterns of BIRC5 expression after compound treatment (fig. S7e). This enables us to identify a set of MEK inhibitors affecting BIRC5 expression and reveals a new cluster of compounds acting on BIRC5 (fig. S7f). Although

5   BIRC5 expression after perturbation has not been reported in the literature for many of these compounds, the cluster includes a survivin inhibitor and Auranofin, which has been reported to downregulate BIRC5 in osteosarcoma(55). Both of these compounds were not previously tested in LINCS.

10   *Identifying EGFR sensitivity in non-small cell lung cancer.*

Applying drug activity fingerprint concept to cell lines, we can also generate a fingerprint for cell lines which we can use to investigate the response signatures of biological systems to a specific array of drugs. As an example, we identify non-small cell lung cancer (NSCLC) cell lines that

15   react to EGFR inhibition. We subset the experimental space to NSCLC cell lines and compute an average EGFR inhibition sensitivity score per cell line (Fig. 3g, Supplementary Methods). EGFR mutant cell lines, with a constitutively active EGFR receptor, are often especially sensitive to EGFR inhibitors (56). Prophet identifies a gradient in the cell lines with respect to sensitivity to EGFR inhibitors, and is successful at scoring five known putative EGFR mutant or EGFR-

20   sensitive cell lines as EGFR sensitive (H1975, H1650, HCC4006, HCC827 and CALU3) (57–59) (Fig. 3h).

In vitro experimental design and validation

To further demonstrate the accuracy of Prophet's predictions, we experimentally validated a subset of the model's predictions. To do so, we used a setup with existing pre-built 384-well

25   plates, each with 352 unique drug perturbations applied to 9 cancer cell lines (Table S9). In total, there were 16 pre-built plates. This setup represents a scenario in which limitations in experiment size constrain the compounds and cell lines that can be tested.

Aiming to pick a plate with maximally diverse drug responses across cell lines, we used Prophet

30   to make in silico predictions for the outcomes of the compounds in each plate in the nine available cell lines. We used the predicted effects on cell viability to score each plate in two categories—dissimilarity across compounds and similarity within clusters (fig. S9a, Supplementary Methods). We then tested the selected plate and compared the results to Prophet's predictions. Prophet successfully predicted the overall outcome for the entire plate,

35   with a median of 0.53 Spearman correlation (Fig. 4c), despite not having seen any related data during training. Comparing the predicted viability scores for each compound of the selected plate with viability measurements in PRISM, the dataset with the best coverage of the compounds and cell lines out of which 8/9 cell lines and 41/352 compounds were also tested, resulted in a median of 0.07 Spearman correlation (Fig. 4c).

For the two known MEK inhibitors tested, GDC-0623 and AZD8330, Prophet correctly predicted no significant change in cell viability after treatment specifically in MCF7, but none of the other cell lines, with high certainty (Fig. 4d, fig. S9b). Our experimental validation confirmed this observation (Fig. 4d), showing that both MEK inhibitors produced no significant changes in

45   viability in MCF7 cells, but significant changes in other cell lines. This finding is supported by previous work documenting MCF7 resistance to some MEK inhibitors, including trametinib(60),

but not other MEK inhibitors such as selumetinib and pimasertib(61). Prophet independently uncovers this highly specific resistance despite not previously seeing either compound in MCF7 cells in training, and not seeing GDC-0623 at all.

Prophet predicts in vivo phenotypes

5 Prophet's capabilities extend to in vivo experimental settings. Saunders et al. 2023 (28) generated a single-cell transcriptomic atlas of the effect of 23 genetic knockouts—including combinations—on 1,812 developing zebrafish embryos across 5 timepoints (Fig. 5a). Their analysis includes using the single-cell transcriptomics data to compute relative cell type frequencies for each embryo (Fig. 5b) and comparing the genetic perturbations using similarities
10 and differences in cell type proportion change. In this high-dimensional space, genetic perturbations clustered together based on the affected zebrafish developmental lineage. Given our framework, we asked if Prophet could learn these cell type proportion changes and make predictions about additional knockouts for specific lineages.

15 Given that this is the only in vivo dataset and that no common prior knowledge exists for it within our dataset compendium, we did not use any pretrained Prophet version, but trained a single Prophet model from scratch (Methods). We then used the predicted relative cell type frequencies to construct a zebrafish cell type proportion experimental space, extending it with predictions of single knockouts, and analyzed perturbation-dependent deviations relative to wild-
20 type embryos (Fig. 5c).

Despite only 23 perturbations in the dataset, the cell type proportion experimental space learnt by Prophet qualitatively matches the one obtained by the authors in Fig. 2c of the original publication (Fig. 5d, top panel), where it can be seen that different gene knockouts provoke
25 different responses in terms of cell type proportion in the zebrafish embryos. The gene knockout fingerprints learnt by Prophet group genes according to their involvement in the mesoderm, nervous system and neural crest lineages (Fig. 5d, bottom panel). We also run predictions for singleton knock outs of zebrafish transcription factors (TFs): for example, znf711 and znf740b from the zf-C2H2 family and dbpb, bach1b from the bZIP family, to produce additional
30 candidates for further experiments.

As an example, we run Prophet to get predictions of gene targets that reduce the cell type proportion of anterior migratory muscle at 36 hours post-fertilization (hpf)(Supplementary Methods). We measure the strength of a knockout as fold change using the control embryo cell
35 type proportion at 36hpf as a reference. Among the top 10 gene targets predicted (Fig. 5e), we find that Rsp02 and Hopx rank first and third, respectively; for both genes, previous studies describe their key role in mesoderm development (62); (63, 64). Other genes in the top-5 (Bmp1 and Wnt8b) are part of families of genes also involved in mesoderm development(65–67). All ten of the Prophet-predicted strongest gene knockouts are not part of the original dataset.
40 Prophet's predictions also report an uncertainty obtained from the variance of predictions of an ensemble of models (Methods). This informs experimentalists of Prophet's confidence in a prediction, allowing them to make an informed decision about whether or not to include high uncertainty TFs in a panel.

45 Finally, Prophet can also provide fine-grained predictions across timepoints by incorporating the time of the readout into the readout token. The proportion of anterior migratory muscle cells

varies across time points given an F0 knockout in a zebrafish embryo in the genes Rspo2 and Hopx (Fig. 5f). These detailed temporal predictions can assist biologists to make more informed decisions about which subsequent experiments to conduct, such as focusing on particular stages for gene expression analysis or investigating downstream effects during critical developmental windows.

This example showcases how Prophet is flexible enough to extend beyond cell lines and in vitro experiments to in vivo phenotypes. In the future, Prophet could be provided with genetically orthologous data in human developmental models to bridge the gap between controlled laboratory experiments and complex in vivo systems, helping to translate findings from basic research into humans and potential therapeutic applications.

This example showcases how our predictive framework is flexible enough to extend beyond cell lines and *in vitro* experiments to *in vivo* phenotypes. In the future, Prophet could be provided with genetically orthologous data in human developmental models to bridge the gap between controlled laboratory experiments and complex *in vivo* systems, helping to translate findings from basic research into humans and potential therapeutic applications.

Discussion

Prophet introduces a novel and scalable approach for phenotypic experimental biology, leveraging transfer learning to effectively predict outcomes across a diverse range of biological experiments. It significantly outperforms simpler architectures and the mean baseline in prediction on unseen experimental conditions (Fig. 2b, fig. S5c). Prophet learns a biologically meaningful experimental space that can be explored to generate new hypotheses. For instance, Prophet's precise predictions allow users to make screens 60x smaller while identifying the same number of hits (Fig. 2f). Activity-based compound fingerprints made possible by Prophet accurately recover known biological pathways, with predictions validating in vitro. Notably, Prophet's capabilities extend beyond cell lines to encompass predictions for in vivo settings (Fig. 5), laying the groundwork for systematic phenotypic drug discovery even in complex biological systems. These evaluations underscore Prophet's potential as a first step towards a foundation model for experimental biology, highlighted by its scaling properties, transfer learning capabilities, and performance on extensive datasets gathered.

Limitations of the current approach are, firstly, the simplified assumption of the three-axis experimental space (Fig. 1). There exist other elements that define an experiment, such as a temporal dimension or the dosage employed, as well as more subtle variations such as technical variation or protocol choice. Prophet considers the time point at which an experiment is measured to be a specification of the readout, without considering the ordinal nature of the variable. Drug dosage could be considered similarly, and these should add ordinal dimensions to experimental space as a topic of future development. Similarly, Prophet could take as input directly the genetic sequences or chemical structures of the treatments, without any preprocessing, or extend to antibody treatments with the appropriate representation and dataset. While Prophet can currently be applied to genetic perturbation datasets, fully leveraging its transfer learning capabilities will require incorporation of more extensive phenotypic datasets of genetic perturbations. With future genetic intervention data and joint datasets, Prophet is posed to effectively translate knowledge between genetic screens and small molecule assays.

While Prophet successfully transfers learning between independent assays and phenotypes, a challenge that remains unsolved is how to transfer learning across scales—between in vitro, in vivo and patient assays. Mastering this transfer is crucial for improving clinical outcomes in drug discovery because more complex functional assays, using organoids, patient samples, or in vivo experiments can often be limited in size. If supplied with this data, the Prophet architecture and framework would enable more accurate predictions by transferring information from scalable in vitro perturbation screens, up the chain of complexity to different biological contexts.

Prophet represents a new concept in phenotypic drug discovery, demonstrating that transfer learning between independent assays is not only possible but also enhances results. Its analysis capabilities and general architecture position it as a step towards establishing a foundation model for perturbation biology: a comprehensive unified model that integrates diverse types of information under a single learning paradigm. Prophet's methodology allows it to be used as a backbone for a lab-in-the-loop system to propose the next set of perturbational experiments, guiding phenotypic drug discovery and navigating the vast experimental space. These attributes make Prophet a powerful tool for driving progress in biological research and experimental design.

References and Notes:

1. S. M. Corsello, R. T. Nagari, R. D. Spangler, J. Rossen, M. Kocak, J. G. Bryan, R. Humeidi, D. Peck, X. Wu, A. A. Tang, V. M. Wang, S. A. Bender, E. Lemire, R. Narayan, P. Montgomery, U. Ben-David, C. W. Garvie, Y. Chen, M. G. Rees, N. J. Lyons, J. M. McFarland, B. T. Wong, L. Wang, N. Dumont, P. J. O'Hearn, E. Stefan, J. G. Doench, C. N. Harrington, H. Greulich, M. Meyerson, F. Vazquez, A. Subramanian, J. A. Roth, J. A. Bittker, J. S. Boehm, C. C. Mader, A. Tsherniak, T. R. Golub, Discovering the anti-cancer potential of non-oncology drugs by systematic viability profiling. Nat Cancer 1, 235–248 (2020).

2. W. Yang, J. Soares, P. Greninger, E. J. Edelman, H. Lightfoot, S. Forbes, N. Bindal, D. Beare, J. A. Smith, I. R. Thompson, S. Ramaswamy, P. A. Futreal, D. A. Haber, M. R. Stratton, C. Benes, U. McDermott, M. J. Garnett, Genomics of Drug Sensitivity in Cancer (GDSC): a resource for therapeutic biomarker discovery in cancer cells. Nucleic Acids Res. 41, D955–61 (2013).

3. F. M. Behan, F. Iorio, G. Picco, E. Gonçalves, C. M. Beaver, G. Migliardi, R. Santos, Y. Rao, F. Sassi, M. Pinnelli, R. Ansari, S. Harper, D. A. Jackson, R. McRae, R. Pooley, P. Wilkinson, D. van der Meer, D. Dow, C. Buser-Doepner, A. Bertotti, L. Trusolino, E. A. Stronach, J. Saez-Rodriguez, K. Yusa, M. J. Garnett, Prioritization of cancer therapeutic targets using CRISPR-Cas9 screens. Nature 568, 511–516 (2019).

4. C. Bock, P. Datlinger, F. Chardon, M. A. Coelho, M. B. Dong, K. A. Lawson, T. Lu, L. Maroc, T. M. Norman, B. Song, G. Stanley, S. Chen, M. Garnett, W. Li, J. Moffat, L. S. Qi, R. S. Shapiro, J. Shendure, J. S. Weissman, X. Zhuang, High-content CRISPR screening. Nature Reviews Methods Primers 2, 1–23 (2022).

5. M. P. Menden, D. Wang, M. J. Mason, B. Szalai, K. C. Bulusu, Y. Guan, T. Yu, J. Kang, M. Jeon, R. Wolfinger, T. Nguyen, M. Zaslavskiy, AstraZeneca-Sanger Drug Combination DREAM Consortium, I. S. Jang, Z. Ghazoui, M. E. Ahsen, R. Vogel, E. C. Neto, T. Norman, E. K. Y. Tang, M. J. Garnett, G. Y. D. Veroli, S. Fawell, G.

Stolovitzky, J. Guinney, J. R. Dry, J. Saez-Rodriguez, Community assessment to advance computational prediction of cancer drug combinations in a pharmacogenomic screen. Nat. Commun. 10, 2674 (2019).

6. S. N. Chandrasekaran, J. Ackerman, E. Alix, D. Michael Ando, J. Arevalo, M. Bennion, N. Boisseau, A. Borowa, J. D. Boyd, L. Brino, P. J. Byrne, H. Ceulemans, C. Ch'ng, B. A. Cimini, D.-A. Clevert, N. Deflaux, J. G. Doench, T. Dorval, R. Doyonnas, V. Dragone, O. Engkvist, P. W. Faloon, B. Fritchman, F. Fuchs, S. Garg, T. J. Gilbert, D. Glazer, D. Gnutt, A. Goodale, J. Grignard, J. Guenther, Y. Han, Z. Hanifehlou, S. Hariharan, D. Hernandez, S. R. Horman, G. Hormel, M. Huntley, I. Icke, M. Iida, C. B. Jacob, S. Jaensch, J. Khetan, M. Kost-Alimova, T. Krawiec, D. Kuhn, C.-H. Lardeau, A. Lembke, F. Lin, K. D. Little, K. R. Lofstrom, S. Lotfi, D. J. Logan, Y. Luo, F. Madoux, P. A. Marin Zapata, B. A. Marion, G. Martin, N. J. McCarthy, L. Mervin, L. Miller, H. Mohamed, T. Monteverde, E. Mouchet, B. Nicke, A. Ogier, A.-L. Ong, M. Osterland, M. Otrocka, P. J. Peeters, J. Pilling, S. Prechtl, C. Qian, K. Rataj, D. E. Root, S. K. Sakata, S. Scrace, H. Shimizu, D. Simon, P. Sommer, C. Spruiell, I. Sumia, S. E. Swalley, H. Terauchi, A. Thibaudeau, A. Unruh, J. Van de Waeter, M. Van Dyck, C. van Staden, M. Warchoł, E. Weisbart, A. Weiss, N. Wiest-Daessle, G. Williams, S. Yu, B. Zapiec, M. Żyła, S. Singh, A. E. Carpenter, JUMP Cell Painting dataset: morphological impact of 136,000 chemical and genetic perturbations, bioRxiv (2023)p. 2023.03.23.534023.

7. S. R. Srivatsan, J. L. McFaline-Figueroa, V. Ramani, L. Saunders, J. Cao, J. Packer, H. A. Pliner, D. L. Jackson, R. M. Daza, L. Christiansen, F. Zhang, F. Steemers, J. Shendure, C. Trapnell, Massively multiplex chemical transcriptomics at single-cell resolution. Science 367, 45–51 (2020).

8. S. N. Chandrasekaran, B. A. Cimini, A. Goodale, L. Miller, M. Kost-Alimova, N. Jamali, J. G. Doench, B. Fritchman, A. Skepner, M. Melanson, A. A. Kalinin, J. Arevalo, M. Haghighi, J. C. Caicedo, D. Kuhn, D. Hernandez, J. Berstler, H. Shafqat-Abbasi, D. E. Root, S. E. Swalley, S. Garg, S. Singh, A. E. Carpenter, Three million images and morphological profiles of cells treated with matched chemical and genetic perturbations. Nat. Methods 21, 1114–1121 (2024).

9. D. Wang, J. Hensman, G. Kutkaite, T. S. Toh, A. Galhoz, GDSC Screening Team, J. R. Dry, J. Saez-Rodriguez, M. J. Garnett, M. P. Menden, F. Dondelinger, A statistical framework for assessing pharmacological responses and biomarkers using uncertainty estimates. Elife 9 (2020).

10. D. M. Camacho, K. M. Collins, R. K. Powers, J. C. Costello, J. J. Collins, Next-Generation Machine Learning for Biological Networks. Cell 173, 1581–1592 (2018).

11. A. Partin, T. S. Brettin, Y. Zhu, O. Narykov, A. Clyde, J. Overbeek, R. L. Stevens, Deep learning methods for drug response prediction in cancer: Predominant and emerging trends. Front. Med. 10, 1086097 (2023).

12. C. Pacini, J. M. Dempster, I. Boyle, E. Gonçalves, H. Najgebauer, E. Karakoc, D. van der Meer, A. Barthorpe, H. Lightfoot, P. Jaaks, J. M. McFarland, M. J. Garnett, A. Tsherniak, F. Iorio, Integrated cross-study datasets of genetic dependencies in cancer. Nat. Commun. 12, 1661 (2021).

13. D. Yao, L. Binan, J. Bezney, B. Simonton, J. Freedman, C. J. Frangieh, K. Dey, K. Geiger-Schuller, B. Eraslan, A. Gusev, A. Regev, B. Cleary, Scalable genetic screening for regulatory circuits using compressed Perturb-seq. Nat. Biotechnol., doi: 10.1038/s41587-023-01964-9 (2023).

14. T. Chu, T. T. Nguyen, B. D. Hai, Q. H. Nguyen, T. Nguyen, Graph Transformer for Drug Response Prediction. IEEE/ACM Trans. Comput. Biol. Bioinform. 20, 1065–1072 (2023).

15. W. Peng, T. Chen, W. Dai, Predicting Drug Response Based on Multi-Omics Fusion and Graph Convolution. IEEE J Biomed Health Inform 26, 1384–1393 (2022).

16. Y.-C. Tang, A. Gottlieb, Explainable drug sensitivity prediction through cancer pathway enrichment. Sci. Rep. 11, 3128 (2021).

17. Y. Zhu, Z. Ouyang, W. Chen, R. Feng, D. Z. Chen, J. Cao, J. Wu, TGSA: protein-protein association-based twin graph neural networks for drug response prediction with similarity augmentation. Bioinformatics 38, 461–468 (2022).

18. Y. L. Chow, S. Singh, A. E. Carpenter, G. P. Way, Predicting drug polypharmacology from cell morphology readouts using variational autoencoder latent space arithmetic. PLoS Comput. Biol. 18, e1009888 (2022).

19. A. Dosovitskiy, L. Beyer, A. Kolesnikov, D. Weissenborn, X. Zhai, T. Unterthiner, M. Dehghani, M. Minderer, G. Heigold, S. Gelly, J. Uszkoreit, N. Houlsby, An Image is Worth 16x16 Words: Transformers for Image Recognition at Scale, arXiv [cs.CV] (2020). http://arxiv.org/abs/2010.11929.

20. T. Hayes, R. Rao, H. Akin, N. J. Sofroniew, D. Oktay, Z. Lin, R. Verkuil, V. Q. Tran, J. Deaton, M. Wiggert, R. Badkundri, I. Shafkat, J. Gong, A. Derry, R. S. Molina, N. Thomas, Y. Khan, C. Mishra, C. Kim, L. J. Bartie, M. Nemeth, P. D. Hsu, T. Sercu, S. Candido, A. Rives, Simulating 500 million years of evolution with a language model, bioRxiv (2024)p. 2024.07.01.600583.

21. T. B. Brown, B. Mann, N. Ryder, M. Subbiah, J. Kaplan, P. Dhariwal, A. Neelakantan, P. Shyam, G. Sastry, A. Askell, S. Agarwal, A. Herbert-Voss, G. Krueger, T. Henighan, R. Child, A. Ramesh, D. M. Ziegler, J. Wu, C. Winter, C. Hesse, M. Chen, E. Sigler, M. Litwin, S. Gray, B. Chess, J. Clark, C. Berner, S. McCandlish, A. Radford, I. Sutskever, D. Amodei, Language Models are Few-Shot Learners. Adv. Neural Inf. Process. Syst. abs/2005.14165 (2020).

22. G. P. Way, T. Natoli, A. Adeboye, L. Litichevskiy, A. Yang, X. Lu, J. C. Caicedo, B. A. Cimini, K. Karhohs, D. J. Logan, M. H. Rohban, M. Kost-Alimova, K. Hartland, M. Bornholdt, S. N. Chandrasekaran, M. Haghighi, E. Weisbart, S. Singh, A. Subramanian, A. E. Carpenter, Morphology and gene expression profiling provide complementary information for mapping cell state. cels 13, 911–923.e9 (2022).

23. A. Subramanian, R. Narayan, S. M. Corsello, D. D. Peck, T. E. Natoli, X. Lu, J. Gould, J. F. Davis, A. A. Tubelli, J. K. Asiedu, D. L. Lahr, J. E. Hirschman, Z. Liu, M. Donahue, B. Julian, M. Khan, D. Wadden, I. C. Smith, D. Lam, A. Liberzon, C. Toder, M. Bagul, M. Orzechowski, O. M. Enache, F. Piccioni, S. A. Johnson, N. J. Lyons, A. H. Berger, A. F. Shamji, A. N. Brooks, A. Vrcic, C. Flynn, J. Rosains, D. Y. Takeda, R. Hu, D. Davison, J. Lamb, K. Ardlie, L. Hogstrom, P. Greenside, N. S.

Gray, P. A. Clemons, S. Silver, X. Wu, W.-N. Zhao, W. Read-Button, X. Wu, S. J. Haggarty, L. V. Ronco, J. S. Boehm, S. L. Schreiber, J. G. Doench, J. A. Bittker, D. E. Root, B. Wong, T. R. Golub, A Next Generation Connectivity Map: L1000 Platform and the First 1,000,000 Profiles. Cell 171, 1437–1452.e17 (2017).

24.   F. Iorio, T. A. Knijnenburg, D. J. Vis, G. R. Bignell, M. P. Menden, M. Schubert, N. Aben, E. Gonçalves, S. Barthorpe, H. Lightfoot, T. Cokelaer, P. Greninger, E. van Dyk, H. Chang, H. de Silva, H. Heyn, X. Deng, R. K. Egan, Q. Liu, T. Mironenko, X. Mitropoulos, L. Richardson, J. Wang, T. Zhang, S. Moran, S. Sayols, M. Soleimani, D. Tamborero, N. Lopez-Bigas, P. Ross-Macdonald, M. Esteller, N. S. Gray, D. A. Haber, M. R. Stratton, C. H. Benes, L. F. A. Wessels, J. Saez-Rodriguez, U. McDermott, M. J. Garnett, A Landscape of Pharmacogenomic Interactions in Cancer. Cell 166, 740–754 (2016).

25.   P. Jaaks, E. A. Coker, D. J. Vis, O. Edwards, E. F. Carpenter, S. M. Leto, L. Dwane, F. Sassi, H. Lightfoot, S. Barthorpe, D. van der Meer, W. Yang, A. Beck, T. Mironenko, C. Hall, J. Hall, I. Mali, L. Richardson, C. Tolley, J. Morris, F. Thomas, E. Lleshi, N. Aben, C. H. Benes, A. Bertotti, L. Trusolino, L. Wessels, M. J. Garnett, Effective drug combinations in breast, colon and pancreatic cancer cells. Nature 603, 166–173 (2022).

26.   A. Basu, N. E. Bodycombe, J. H. Cheah, E. V. Price, K. Liu, G. I. Schaefer, R. Y. Ebright, M. L. Stewart, D. Ito, S. Wang, A. L. Bracha, T. Liefeld, M. Wawer, J. C. Gilbert, A. J. Wilson, N. Stransky, G. V. Kryukov, V. Dancik, J. Barretina, L. A. Garraway, C. S.-Y. Hon, B. Munoz, J. A. Bittker, B. R. Stockwell, D. Khabele, A. M. Stern, P. A. Clemons, A. F. Shamji, S. L. Schreiber, An interactive resource to identify cancer genetic and lineage dependencies targeted by small molecules. Cell 154, 1151–1161 (2013).

27.   M. A. Horlbeck, A. Xu, M. Wang, N. K. Bennett, C. Y. Park, D. Bogdanoff, B. Adamson, E. D. Chow, M. Kampmann, T. R. Peterson, K. Nakamura, M. A. Fischbach, J. S. Weissman, L. A. Gilbert, Mapping the Genetic Landscape of Human Cells. Cell 174, 953–967.e22 (2018).

28.   L. M. Saunders, S. R. Srivatsan, M. Duran, M. W. Dorrity, B. Ewing, T. H. Linbo, J. Shendure, D. W. Raible, C. B. Moens, D. Kimelman, C. Trapnell, Embryo-scale reverse genetics at single-cell resolution. Nature 623, 782–791 (2023).

29.   A. Vaswani, N. Shazeer, N. Parmar, J. Uszkoreit, L. Jones, A. N. Gomez, L. Kaiser, I. Polosukhin, Attention Is All You Need, arXiv [cs.CL] (2017). http://arxiv.org/abs/1706.03762.

30.   J. Devlin, M.-W. Chang, K. Lee, K. Toutanova, BERT: Pre-training of Deep Bidirectional Transformers for Language Understanding, arXiv [cs.CL] (2018). http://arxiv.org/abs/1810.04805.

31.   J. Barretina, G. Caponigro, N. Stransky, K. Venkatesan, A. A. Margolin, S. Kim, C. J. Wilson, J. Lehár, G. V. Kryukov, D. Sonkin, A. Reddy, M. Liu, L. Murray, M. F. Berger, J. E. Monahan, P. Morais, J. Meltzer, A. Korejwa, J. Jané-Valbuena, F. A. Mapa, J. Thibault, E. Bric-Furlong, P. Raman, A. Shipway, I. H. Engels, J. Cheng, G. K. Yu, J. Yu, P. Aspesi Jr, M. de Silva, K. Jagtap, M. D. Jones, L. Wang, C. Hatton, E. Palescandolo, S. Gupta, S. Mahan, C. Sougnez, R. C. Onofrio, T. Liefeld, L.

MacConaill, W. Winckler, M. Reich, N. Li, J. P. Mesirov, S. B. Gabriel, G. Getz, K. Ardlie, V. Chan, V. E. Myer, B. L. Weber, J. Porter, M. Warmuth, P. Finan, J. L. Harris, M. Meyerson, T. R. Golub, M. P. Morrissey, W. R. Sellers, R. Schlegel, L. A. Garraway, The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature 483, 603–607 (2012).

32. G. Landrum, RDKit: A software suite for cheminformatics, computational chemistry, and predictive modeling. http://www.rdkit.org/RDKit_Overview.pdf.

33. H. L. Morgan, The Generation of a Unique Machine Description for Chemical Structures-A Technique Developed at Chemical Abstracts Service. J. Chem. Doc. 5, 107–113 (1965).

34. Ž. Avsec, V. Agarwal, D. Visentin, J. R. Ledsam, A. Grabska-Barwinska, K. R. Taylor, Y. Assael, J. Jumper, P. Kohli, D. R. Kelley, Effective gene expression prediction from sequence by integrating long-range interactions. Nat. Methods 18, 1196–1203 (2021).

35. M.-A. Bray, S. Singh, H. Han, C. T. Davis, B. Borgeson, C. Hartland, M. Kost-Alimova, S. M. Gustafsdottir, C. C. Gibson, A. E. Carpenter, Cell Painting, a high-content image-based assay for morphological profiling using multiplexed fluorescent dyes. Nat. Protoc. 11, 1757–1774 (2016).

36. A. Kolesnikov, L. Beyer, X. Zhai, J. Puigcerver, J. Yung, S. Gelly, N. Houlsby, Big Transfer (BiT): General Visual Representation Learning, arXiv [cs.CV] (2019). http://arxiv.org/abs/1912.11370.

37. X. Zhai, A. Kolesnikov, N. Houlsby, L. Beyer, Scaling Vision Transformers, arXiv [cs.CV] (2021). http://arxiv.org/abs/2106.04560.

38. I. C. Henderson, J. C. Allegra, T. Woodcock, S. Wolff, S. Bryan, K. Cartwright, G. Dukart, D. Henry, Randomized clinical trial comparing mitoxantrone with doxorubicin in previously treated patients with metastatic breast cancer. J. Clin. Oncol. 7, 560–571 (1989).

39. E. J. Feldman, High-dose mitoxantrone in acute leukaemia: New York Medical College experience. Eur. J. Cancer Care 6, 27–32 (1997).

40. S. Vibet, K. Mahéo, J. Goré, P. Dubois, P. Bougnoux, I. Chourpa, Differential subcellular distribution of mitoxantrone in relation to chemosensitization in two human breast cancer cell lines. Drug Metab. Dispos. 35, 822–828 (2007).

41. K. Takara, K. Yamamoto, M. Matsubara, T. Minegaki, M. Takahashi, T. Yokoyama, K. Okumura, Effects of α-adrenoceptor antagonists on ABCG2/BCRP-mediated resistance and transport. PLoS One 7, e30697 (2012).

42. P. G. Richardson, C. S. Mitsiades, J. P. Laubach, S. Lonial, A. A. Chanan-Khan, K. C. Anderson, Inhibition of heat shock protein 90 (HSP90) as a therapeutic strategy for the treatment of myeloma and other cancers. Br. J. Haematol. 152, 367–379 (2011).

43. H. Wu, A. Wang, W. Zhang, B. Wang, C. Chen, W. Wang, C. Hu, Z. Ye, Z. Zhao, L. Wang, X. Li, K. Yu, J. Liu, J. Wu, X.-E. Yan, P. Zhao, J. Wang, C. Wang, E. L. Weisberg, N. S. Gray, C.-H. Yun, J. Liu, L. Chen, Q. Liu, Ibrutinib selectively and

irreversibly targets EGFR (L858R, Del19) mutant but is moderately resistant to EGFR (T790M) mutant NSCLC Cells. Oncotarget 6, 31313–31322 (2015).

44. A. Wang, X.-E. Yan, H. Wu, W. Wang, C. Hu, C. Chen, Z. Zhao, P. Zhao, X. Li, L. Wang, B. Wang, Z. Ye, J. Wang, C. Wang, W. Zhang, N. S. Gray, E. L. Weisberg, L. Chen, J. Liu, C.-H. Yun, Q. Liu, Ibrutinib targets mutant-EGFR kinase with a distinct binding conformation. Oncotarget 7, 69760–69769 (2016).

45. K. T. Flaherty, I. Puzanov, K. B. Kim, A. Ribas, G. A. McArthur, J. A. Sosman, P. J. O'Dwyer, R. J. Lee, J. F. Grippo, K. Nolop, P. B. Chapman, Inhibition of mutated, activated BRAF in metastatic melanoma. N. Engl. J. Med. 363, 809–819 (2010).

46. G. S. Falchook, G. V. Long, R. Kurzrock, K. B. Kim, T. H. Arkenau, M. P. Brown, O. Hamid, J. R. Infante, M. Millward, A. C. Pavlick, S. J. O'Day, S. C. Blackman, C. M. Curtis, P. Lebowitz, B. Ma, D. Ouellet, R. F. Kefford, Dabrafenib in patients with melanoma, untreated brain metastases, and other solid tumours: a phase 1 dose-escalation trial. Lancet 379, 1893–1901 (2012).

47. R. B. Corcoran, C. E. Atreya, G. S. Falchook, E. L. Kwak, D. P. Ryan, J. C. Bendell, O. Hamid, W. A. Messersmith, A. Daud, R. Kurzrock, M. Pierobon, P. Sun, E. Cunningham, S. Little, K. Orford, M. Motwani, Y. Bai, K. Patel, A. P. Venook, S. Kopetz, Combined BRAF and MEK Inhibition With Dabrafenib and Trametinib in BRAF V600-Mutant Colorectal Cancer. J. Clin. Oncol. 33, 4023–4031 (2015).

48. A. Hauschild, P. A. Ascierto, D. Schadendorf, J. J. Grob, A. Ribas, F. Kiecker, C. Dutriaux, L. V. Demidov, C. Lebbé, P. Rutkowski, C. U. Blank, R. Gutzmer, M. Millward, R. Kefford, T. Haas, A. D'Amelio Jr, E. Gasal, B. Mookerjee, P. B. Chapman, Long-term outcomes in patients with BRAF V600-mutant metastatic melanoma receiving dabrafenib monotherapy: Analysis from phase 2 and 3 clinical trials. Eur. J. Cancer 125, 114–120 (2020).

49. G. V. Long, K. T. Flaherty, D. Stroyakovskiy, H. Gogas, E. Levchenko, F. de Braud, J. Larkin, C. Garbe, T. Jouary, A. Hauschild, V. Chiarion-Sileni, C. Lebbe, M. Mandalá, M. Millward, A. Arance, I. Bondarenko, J. B. A. G. Haanen, J. Hansson, J. Utikal, V. Ferraresi, P. Mohr, V. Probachai, D. Schadendorf, P. Nathan, C. Robert, A. Ribas, M. A. Davies, S. R. Lane, J. J. Legos, B. Mookerjee, J.-J. Grob, Dabrafenib plus trametinib versus dabrafenib monotherapy in patients with metastatic BRAF V600E/K-mutant melanoma: long-term survival and safety analysis of a phase 3 study. Ann. Oncol. 30, 1848 (2019).

50. Z. Eroglu, A. Ribas, Combination therapy with BRAF and MEK inhibitors for melanoma: latest evidence and place in therapy. Ther. Adv. Med. Oncol. 8, 48–56 (2016).

51. F. Xing, Y. Persaud, C. A. Pratilas, B. S. Taylor, M. Janakiraman, Q.-B. She, H. Gallardo, C. Liu, T. Merghoub, B. Hefter, I. Dolgalev, A. Viale, A. Heguy, E. De Stanchina, D. Cobrinik, G. Bollag, J. Wolchok, A. Houghton, D. B. Solit, Concurrent loss of the PTEN and RB1 tumor suppressors attenuates RAF dependence in melanomas harboring (V600E)BRAF. Oncogene 31, 446–457 (2012).

52. A. Maklad, A. Sharma, I. Azimi, Calcium Signaling in Brain Cancers: Roles and Therapeutic Targeting. Cancers 11 (2019).

53. Y. Zhang, N. Cruickshanks, F. Yuan, B. Wang, M. Pahuski, J. Wulfkuhle, I. Gallagher, A. F. Koeppel, S. Hatef, C. Papanicolas, J. Lee, E. E. Bar, D. Schiff, S. D. Turner, E. F. Petricoin, L. S. Gray, R. Abounader, Targetable T-type Calcium Channels Drive Glioblastoma. Cancer Res. 77, 3479–3490 (2017).

54. M. Sedeeq, A. Maklad, T. Dutta, Z. Feng, R. Wilson, N. Gueven, I. Azimi, T-Type Calcium Channel Inhibitors Induce Apoptosis in Medulloblastoma Cells Associated with Altered Metabolic Activity. Mol. Neurobiol. 59, 2932–2945 (2022).

55. E. Topkas, N. Cai, A. Cumming, M. Hazar-Rethinam, O. M. Gannon, M. Burgess, N. A. Saunders, L. Endo-Munoz, Auranofin is a potent suppressor of osteosarcoma metastasis. Oncotarget 7, 831–844 (2016).

56. Lynch Thomas J., Bell Daphne W., Sordella Raffaella, Gurubhagavatula Sarada, Okimoto Ross A., Brannigan Brian W., Harris Patricia L., Haserlat Sara M., Supko Jeffrey G., Haluska Frank G., Louis David N., Christiani David C., Settleman Jeff, Haber Daniel A., Activating Mutations in the Epidermal Growth Factor Receptor Underlying Responsiveness of Non–Small-Cell Lung Cancer to Gefitinib. N. Engl. J. Med. 350, 2129–2139.

57. T. Okabe, I. Okamoto, K. Tamura, M. Terashima, T. Yoshida, T. Satoh, M. Takada, M. Fukuoka, K. Nakagawa, Differential constitutive activation of the epidermal growth factor receptor in non-small cell lung cancer cells bearing EGFR gene mutation and amplification. Cancer Res. 67, 2046–2053 (2007).

58. A. Bairoch, The Cellosaurus, a Cell-Line Knowledge Resource. J. Biomol. Tech. 29, 25–38 (2018).

59. F. Morgillo, T. Cascone, E. D'Aiuto, E. Martinelli, T. Troiani, P. Saintigny, R. De Palma, J. V. Heymach, L. Berrino, C. Tuccillo, F. Ciardiello, Antitumour efficacy of MEK inhibitors in human lung cancer cells and their derivatives with acquired resistance to different tyrosine kinase inhibitors. Br. J. Cancer 105, 382–392 (2011).

60. X. Wen, Y.-C. Ou, G. Bogatcheva, G. Thomas, A. Mahadevan-Jansen, B. Singh, E. C. Lin, R. Bardhan, Probing metabolic alterations in breast cancer in response to molecular inhibitors with Raman spectroscopy and validated with mass spectrometry. Chem. Sci. 11, 9863–9874 (2020).

61. Y. Gu, M. Helenius, K. Väänänen, D. Bulanova, J. Saarela, A. Sokolenko, J. Martens, E. Imyanitov, S. Kuznetsov, BRCA1-deficient breast cancer cell lines are resistant to MEK inhibitors and show distinct sensitivities to 6-thioguanine. Sci. Rep. 6, 28217 (2016).

62. A. H. Reis, S. Y. Sokol, Rspo2 antagonizes FGF signaling during vertebrate mesoderm formation and patterning. Development 147 (2020).

63. A. H. Reis, S. Y. Sokol, Rspo2 inhibits TCF3 phosphorylation to antagonize Wnt signaling during vertebrate anteroposterior axis specification. Sci. Rep. 11, 13433 (2021).

64. C. E. Friedman, Q. Nguyen, S. W. Lukowski, A. Helfer, H. S. Chiu, J. Miklas, S. Levy, S. Suo, J.-D. J. Han, P. Osteil, G. Peng, N. Jing, G. J. Baillie, A. Senabouth, A. N. Christ, T. J. Bruxner, C. E. Murry, E. S. Wong, J. Ding, Y. Wang, J. Hudson, H. Ruohola-Baker, Z. Bar-Joseph, P. P. L. Tam, J. E. Powell, N. J. Palpant, Single-Cell

Transcriptomic Analysis of Cardiac Differentiation from Human PSCs Reveals HOPX-Dependent Cardiomyocyte Maturation. Cell Stem Cell 23, 586–598.e8 (2018).

65. A. Manfrin, Y. Tabata, E. R. Paquet, A. R. Vuaridel, F. R. Rivest, F. Naef, M. P. Lutolf, Engineered signaling centers for the spatially controlled patterning of human pluripotent stem cells. Nat. Methods 16, 640–648 (2019).

66. S. M. Morgani, J. J. Metzger, J. Nichols, E. D. Siggia, A.-K. Hadjantonakis, Micropattern differentiation of mouse pluripotent stem cells recapitulates embryo regionalized cell fate patterning. Elife 7 (2018).

67. R. C. Lindsley, J. G. Gill, M. Kyba, T. L. Murphy, K. M. Murphy, Canonical Wnt signaling is required for development of embryonic stem cell-derived mesoderm. Development 133, 3787–3796 (2006).

68. I. Loshchilov, F. Hutter, Decoupled Weight Decay Regularization, arXiv [cs.LG] (2017). http://arxiv.org/abs/1711.05101.

69. I. Loshchilov, F. Hutter, Sgdr: Stochastic gradient descent with warm restarts. arXiv preprint arXiv:1608. 03983 (2016).

70. X. Glorot, Y. Bengio, "Understanding the difficulty of training deep feedforward neural networks" in Proceedings of the Thirteenth International Conference on Artificial Intelligence and Statistics, Y. W. Teh, M. Titterington, Eds. (PMLR, Chia Laguna Resort, Sardinia, Italy, 13--15 May 2010)vol. 9 of Proceedings of Machine Learning Research, pp. 249–256.

71. D. Szklarczyk, A. L. Gable, D. Lyon, A. Junge, S. Wyder, J. Huerta-Cepas, M. Simonovic, N. T. Doncheva, J. H. Morris, P. Bork, L. J. Jensen, C. von Mering, STRING v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. Nucleic Acids Res. 47, D607 (2019).

72. T. M. Norman, M. A. Horlbeck, J. M. Replogle, A. Y. Ge, A. Xu, M. Jost, L. A. Gilbert, J. S. Weissman, Exploring genetic interaction manifolds constructed from rich single-cell phenotypes. Science 365 (2019).

73. J. M. Replogle, R. A. Saunders, A. N. Pogson, J. A. Hussmann, A. Lenail, A. Guna, L. Mascibroda, E. J. Wagner, K. Adelman, G. Lithwick-Yanai, N. Iremadze, F. Oberstrass, D. Lipson, J. L. Bonnar, M. Jost, T. M. Norman, J. S. Weissman, Mapping information-rich genotype-phenotype landscapes with genome-scale Perturb-seq. Cell 185 (2022).

74. J. Du, P. Jia, Y. Dai, C. Tao, Z. Zhao, D. Zhi, Gene2vec: distributed representation of genes based on co-expression. BMC Genomics 20, 7–15 (2019).

75. S. A. Aleksander, J. Balhoff, S. Carbon, J. M. Cherry, H. J. Drabkin, D. Ebert, M. Feuermann, P. Gaudet, N. L. Harris, D. P. Hill, R. Lee, H. Mi, S. Moxon, C. J. Mungall, A. Muruganugan, T. Mushayahama, P. W. Sternberg, P. D. Thomas, K. Van Auken, J. Ramsey, D. A. Siegele, R. L. Chisholm, P. Fey, M. C. Aspromonte, M. V. Nugnes, F. Quaglia, S. Tosatto, M. Giglio, S. Nadendla, G. Antonazzo, H. Attrill, G. Dos Santos, S. Marygold, V. Strelets, C. J. Tabone, J. Thurmond, P. Zhou, S. H. Ahmed, P. Asanitthong, B. D. Luna, M. N. Erdol, M. C. Gage, K. M. Ali, K. Y. C. Li, M. Long, A. Michalak, A. Pesala, A. Pritazahra, S. C. C. Saverimuttu, R. Su, K. E.

Thurlow, R. C. Lovering, C. Logie, S. Oliferenko, J. Blake, K. Christie, L. Corbani, M. E. Dolan, H. J. Drabkin, D. P. Hill, L. Ni, D. Sitnikov, C. Smith, A. Cuzick, J. Seager, L. Cooper, J. Elser, P. Jaiswal, P. Gupta, P. Jaiswal, S. Naithani, M. Lera-Ramirez, K. Rutherford, V. Wood, J. L. De Pons, M. R. Dwinell, G. T. Hayman, M. L. Kaldunski, A. E. Kwitek, S. J. F. Laulederkind, M. A. Tutaj, M. Vedi, S. J. Wang, P. D'Eustachio, L. Aimo, K. Axelsen, A. Bridge, N. Hyka-Nouspikel, A. Morgat, S. A. Aleksander, J. M. Cherry, S. R. Engel, K. Karra, S. R. Miyasato, R. S. Nash, M. S. Skrzypek, S. Weng, E. D. Wong, E. Bakker, T. Z. Berardini, L. Reiser, A. Auchincloss, K. Axelsen, G. Argoud-Puy, M. C. Blatter, E. Boutet, L. Breuza, A. Bridge, C. Casals-Casas, E. Coudert, A. Estreicher, F. M. Livia, M. Feuermann, A. Gos, N. Gruaz-Gumowski, C. Hulo, N. Hyka-Nouspikel, F. Jungo, P. Le Mercier, D. Lieberherr, P. Masson, A. Morgat, I. Pedruzzi, L. Pourcel, S. Poux, C. Rivoire, S. Sundaram, A. Bateman, E. Bowler-Barnett, H. Bye-A-Jee, P. Denny, A. Ignatchenko, R. Ishtiaq, A. Lock, Y. Lussi, M. Magrane, M. J. Martin, S. Orchard, P. Raposo, E. Speretta, N. Tyagi, K. Warner, R. Zaru, A. D. Diehl, R. Lee, J. Chan, S. Diamantakis, D. Raciti, M. Zarowiecki, M. Fisher, C. James-Zorn, V. Ponferrada, A. Zorn, S. Ramachandran, L. Ruzicka, M. Westerfield, The Gene Ontology knowledgebase in 2023. Genetics 224 (2023).

76. M. Ashburner, C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis, K. Dolinski, S. S. Dwight, J. T. Eppig, M. A. Harris, D. P. Hill, L. Issel-Tarver, A. Kasarskis, S. Lewis, J. C. Matese, J. E. Richardson, M. Ringwald, G. M. Rubin, G. Sherlock, Gene Ontology: tool for the unification of biology. Nat. Genet. 25, 25–29 (2000).

77. CZI Single-Cell Biology Program, S. Abdulla, B. Aevermann, P. Assis, S. Badajoz, S. M. Bell, E. Bezzi, B. Cakir, J. Chaffer, S. Chambers, J. Michael Cherry, T. Chi, J. Chien, L. Dorman, P. Garcia-Nieto, N. Gloria, M. Hastie, D. Hegeman, J. Hilton, T. Huang, A. Infeld, A.-M. Istrate, I. Jelic, K. Katsuya, Y. J. Kim, K. Liang, M. Lin, M. Lombardo, B. Marshall, B. Martin, F. McDade, C. Megill, N. Patel, A. Predeus, B. Raymor, B. Robatmili, D. Rogers, E. Rutherford, D. Sadgat, A. Shin, C. Small, T. Smith, P. Sridharan, A. Tarashansky, N. Tavares, H. Thomas, A. Tolopko, M. Urisko, J. Yan, G. Yeretssian, J. Zamanian, A. Mani, J. Cool, A. Carr, CZ CELL×GENE Discover: A single-cell data platform for scalable exploration, analysis and modeling of aggregated data, bioRxiv (2023)p. 2023.10.30.563174.

78. D. van der Meer, S. Barthorpe, W. Yang, H. Lightfoot, C. Hall, J. Gilbert, H. E. Francies, M. J. Garnett, Cell Model Passports—a hub for clinical, genetic and functional datasets of preclinical cancer models. Nucleic Acids Res. 47, D923 (2019).

79. F. A. Wolf, P. Angerer, F. J. Theis, SCANPY: large-scale single-cell gene expression data analysis. Genome Biol. 19, 15 (2018).

80. L. McInnes, J. Healy, J. Melville, UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction, arXiv [stat.ML] (2018). http://arxiv.org/abs/1802.03426.

81. A. Tsherniak, F. Vazquez, P. G. Montgomery, B. A. Weir, G. Kryukov, G. S. Cowley, S. Gill, W. F. Harrington, S. Pantel, J. M. Krill-Burger, R. M. Meyers, L. Ali, A. Goodale, Y. Lee, G. Jiang, J. Hsiao, W. F. J. Gerath, S. Howell, E. Merkel, M.

Ghandi, L. A. Garraway, D. E. Root, T. R. Golub, J. S. Boehm, W. C. Hahn, Defining a Cancer Dependency Map. Cell 170, 564–576.e16 (2017).

82. L. Heumos, Y. Ji, L. May, T. Green, X. Zhang, X. Wu, J. Ostner, S. Peidli, A. Schumacher, K. Hrovatin, M. Mueller, F. Chong, G. Sturm, A. Tejada, E. Dann, M. Dong, M. Bahrami, I. Gold, S. Rybakov, A. Namsaraeva, A. A. Moinfar, Z. Zheng, E. Roellin, I. Mekki, C. Sander, M. Lotfollahi, H. B. Schiller, F. J. Theis, Pertpy: an end-to-end framework for perturbation analysis, bioRxiv (2024)p. 2024.08.04.606516.

83. M. Ester, H. P. Kriegel, J. Sander, X. Xu, A density-based algorithm for discovering clusters in large spatial databases with noise. KDD (1996).

84. M. Dumble, M.-C. Crouthamel, S.-Y. Zhang, M. Schaber, D. Levy, K. Robell, Q. Liu, D. J. Figueroa, E. A. Minthorn, M. A. Seefeld, M. B. Rouse, S. K. Rabindran, D. A. Heerding, R. Kumar, Discovery of Novel AKT Inhibitors with Enhanced Anti-Tumor Effects in Combination with the MEK Inhibitor. PLoS One 9 (2014).

85. B. R. Davies, H. Greenwood, P. Dudley, C. Crafter, D. H. Yu, J. Zhang, J. Li, B. Gao, Q. Ji, J. Maynard, S. A. Ricketts, D. Cross, S. Cosulich, C. C. Chresta, K. Page, J. Yates, C. Lane, R. Watson, R. Luke, D. Ogilvie, M. Pass, Preclinical pharmacology of AZD5363, an inhibitor of AKT: pharmacodynamics, antitumor activity, and correlation of monotherapy activity with genetic background. Mol. Cancer Ther. 11 (2012).

86. D. Sarker, J. E. Ang, R. Baird, R. Kristeleit, K. Shah, V. Moreno, P. A. Clarke, F. I. Raynaud, G. Levy, J. A. Ware, K. Mazina, R. Lin, J. Wu, J. Fredrickson, J. M. Spoerke, M. R. Lackner, Y. Yan, L. S. Friedman, S. B. Kaye, M. K. Derynck, P. Workman, J. S. de Bono, First-in-human Phase I study of Pictilisib (GDC-0941), a potent pan-class I phosphatidylinositol-3-kinase (PI3K) inhibitor, in patients with advanced solid tumors. Clin. Cancer Res. 21, 77 (2015).

87. M. C. Pietanza, T. J. Lynch, P. N. Lara, J. Cho, R. H. Yanagihara, N. Vrindavanam, N. M. Chowhan, S. M. Gadgeel, N. A. Pennell, R. Funke, B. Mitchell, H. A. Wakelee, V. A. Miller, XL647--a multitargeted tyrosine kinase inhibitor: results of a phase II study in subjects with non-small cell lung cancer who have progressed after responding to treatment with either gefitinib or erlotinib. J. Thorac. Oncol. 7 (2012).

88. H. A. M. Gomaa, A. T. Ali, M. Abdel Gabbar, M. A. Kandeil, The Effect of Canertinib on Sensitivity of Cytotoxic Drugs in Tamoxifen-Resistant Breast Cancer Cells In Vitro. Int. J. Genomics Proteomics 2018 (2018).

89. Recent advancements of 4-aminoquinazoline derivatives as kinase inhibitors and their applications in medicinal chemistry. Eur. J. Med. Chem. 170, 55–72 (2019).

90. E. Jasek-Gajda, H. Jurkowska, M. Jasińska, J. A. Litwin, G. J. Lis, Combination of ERK2 inhibitor VX-11e and voreloxin synergistically enhances anti-proliferative and pro-apoptotic effects in leukemia cells. Apoptosis 24, 849 (2019).

91. A. Maiti, K. Naqvi, T. M. Kadia, G. Borthakur, K. Takahashi, P. Bose, N. G. Daver, A. Patel, Y. Alvarado, M. Ohanian, C. D. DiNardo, J. E. Cortes, E. J. Jabbour, G. Garcia-Manero, H. M. Kantarjian, F. Ravandi, Phase II Trial of MEK Inhibitor Binimetinib (MEK162) in RAS-mutant Acute Myeloid Leukemia. Clin. Lymphoma Myeloma Leuk. 19 (2019).

92. R. J. Sullivan, J. R. Infante, F. Janku, D. J. L. Wong, J. A. Sosman, V. Keedy, M. R. Patel, G. I. Shapiro, J. W. Mier, A. W. Tolcher, A. Wang-Gillam, M. Sznol, K. Flaherty, E. Buchbinder, R. D. Carvajal, A. M. Varghese, M. E. Lacouture, A. Ribas, S. P. Patel, G. A. DeCrescenzo, C. M. Emery, A. L. Groover, S. Saha, M. Varterasian, D. J. Welsch, D. M. Hyman, B. T. Li, First-in-Class ERK1/2 Inhibitor Ulixertinib (BVD-523) in Patients with MAPK Mutant Advanced Solid Tumors: Results of a Phase I Dose-Escalation and Expansion Study. Cancer Discov. 8 (2018).

93. J. F. Ohren, H. Chen, A. Pavlovsky, C. Whitehead, E. Zhang, P. Kuffa, C. Yan, P. McConnell, C. Spessard, C. Banotai, W. T. Mueller, A. Delaney, C. Omer, J. Sebolt-Leopold, D. T. Dudley, I. K. Leung, C. Flamme, J. Warmus, M. Kaufman, S. Barrett, H. Tecle, C. A. Hasemann, Structures of human MAP kinase kinase 1 (MEK1) and MEK2 describe novel noncompetitive kinase inhibition. Nat. Struct. Mol. Biol. 11, 1192–1197 (2004).

94. M. E. Burkard, M. McKean, J. R. Ahnert, N. B. Mettu, J. C. Jones, J. G. Misleh, W. W. Ma, K.-H. Lim, E. Gabriela Chiorean, M. J. Pishvaian, S. M. Gadgeel, H. A. McKean, B. Kreider, D. Knoerzer, A. Groover, M. L. Varterasian, J. A. Box, C. Emery, R. J. Sullivan, A two-part, phase II, multi-center study of the ERK inhibitor ulixertinib (BVD-523) for patients with advanced malignancies harboring MEK or atypical BRAF alterations (BVD-523-ABC). J. Clin. Oncol., doi: 10.1200/JCO.2022.40.16_suppl.TPS3172 (2022).

95. A. Chaikuad, E. Tacconi, J. Zimmer, Y. Liang, N. S. Gray, M. Tarsounas, S. Knapp, A unique inhibitor binding site in ERK1/2 is associated with slow binding kinetics. Nat. Chem. Biol. 10, 853 (2014).

96. C. Du, X. Guan, J. Yan, Two-pore channel blockade by phosphoinositide kinase inhibitors YM201636 and PI-103 determined by a histidine residue near pore-entrance. Communications biology 5 (2022).

97. S. M. Guichard, J. Curwen, T. Bihani, C. M. D'Cruz, J. W. Yates, M. Grondine, Z. Howard, B. R. Davies, G. Bigley, T. Klinowska, K. G. Pike, M. Pass, C. M. Chresta, U. M. Polanska, R. McEwen, O. Delpuech, S. Green, S. C. Cosulich, AZD2014, an Inhibitor of mTORC1 and mTORC2, Is Highly Effective in ER+ Breast Cancer When Administered Using Intermittent or Continuous Schedules. Mol. Cancer Ther. 14 (2015).

98. L. Salphati, T. P. Heffron, B. Alicke, M. Nishimura, K. Barck, R. A. Carano, J. Cheong, K. A. Edgar, J. Greve, S. Kharbanda, H. Koeppen, S. Lau, L. B. Lee, J. Pang, E. G. Plise, J. L. Pokorny, H. B. Reslan, J. N. Sarkaria, J. J. Wallin, X. Zhang, S. E. Gould, A. G. Olivero, H. S. Phillips, Targeting the PI3K pathway in the brain--efficacy of a PI3K inhibitor optimized to cross the blood-brain barrier. Clin. Cancer Res. 18 (2012).

99. L. Ciuffreda, D. Del Bufalo, M. Desideri, C. Di Sanza, A. Stoppacciaro, M. R. Ricciardi, S. Chiaretti, S. Tavolaro, B. Benassi, A. Bellacosa, R. Foà, A. Tafuri, F. Cognetti, A. Anichini, G. Zupi, M. Milella, Growth-inhibitory and antiangiogenic activity of the MEK inhibitor PD0325901 in malignant melanoma with or without BRAF mutations. Neoplasia 11, 720–731 (2009).

100. V. Roulstone, M. Pedersen, J. Kyula, D. Mansfield, A. A. Khan, G. McEntee, M. Wilkinson, E. Karapanagiotou, M. Coffey, R. Marais, A. Jebar, F. Errington-Mais, A. Melcher, R. Vile, H. Pandha, M. McLaughlin, K. J. Harrington, BRAF- and MEK-Targeted Small Molecule Inhibitors Exert Enhanced Antimelanoma Effects in Combination With Oncolytic Reovirus Through ER Stress. Mol. Ther. 23, 931 (2015).

101. J. Tsai, J. T. Lee, W. Wang, J. Zhang, H. Cho, S. Mamo, R. Bremer, S. Gillette, J. Kong, N. K. Haass, K. Sproesser, L. Li, K. S. M. Smalley, D. Fong, Y.-L. Zhu, A. Marimuthu, H. Nguyen, B. Lam, J. Liu, I. Cheung, J. Rice, Y. Suzuki, C. Luu, C. Settachatgul, R. Shellooe, J. Cantwell, S.-H. Kim, J. Schlessinger, K. Y. J. Zhang, B. L. West, B. Powell, G. Habets, C. Zhang, P. N. Ibrahim, P. Hirth, D. R. Artis, M. Herlyn, G. Bollag, Discovery of a selective inhibitor of oncogenic B-Raf kinase with potent antimelanoma activity. Proceedings of the National Academy of Sciences 105, 3041–3046 (2008).

102. J. T. Lee, L. Li, P. A. Brafford, M. van den Eijnden, M. B. Halloran, K. Sproesser, N. K. Haass, K. S. M. Smalley, J. Tsai, G. Bollag, M. Herlyn, PLX4032, a Potent Inhibitor of the B-Raf V600E Oncogene, Selectively Inhibits V600E-positive Melanomas. Pigment Cell Melanoma Res. 23, 820 (2010).

103. E. J. Morris, S. Jha, C. R. Restaino, P. Dayananth, H. Zhu, A. Cooper, D. Carr, Y. Deng, W. Jin, S. Black, B. Long, J. Liu, E. Dinunzio, W. Windsor, R. Zhang, S. Zhao, M. H. Angagaw, E. M. Pinheiro, J. Desai, L. Xiao, G. Shipps, A. Hruza, J. Wang, J. Kelly, S. Paliwal, X. Gao, B. S. Babu, L. Zhu, P. Daublain, L. Zhang, B. A. Lutterbach, M. R. Pelletier, U. Philippar, P. Siliphaivanh, D. Witter, P. Kirschmeier, W. R. Bishop, D. Hicklin, D. G. Gilliland, L. Jayaraman, L. Zawel, S. Fawell, A. A. Samatar, Discovery of a novel ERK inhibitor with activity in models of acquired resistance to BRAF and MEK inhibitors. Cancer Discov. 3, 742–750 (2013).

104. S. J. Park, S. W. Hong, J. H. Moon, D. H. Jin, J. S. Kim, C. K. Lee, K. P. Kim, Y. S. Hong, E. K. Choi, J. S. Lee, J. L. Lee, T. W. Kim, The MEK1/2 inhibitor AS703026 circumvents resistance to the BRAF inhibitor PLX4032 in human malignant melanoma cells. Am. J. Med. Sci. 346 (2013).

105. U. A. Germann, B. F. Furey, W. Markland, R. R. Hoover, A. M. Aronov, J. J. Roix, M. Hale, D. M. Boucher, D. A. Sorrell, G. Martinez-Botella, M. Fitzgibbon, P. Shapiro, M. J. Wick, R. Samadani, K. Meshaw, A. Groover, G. DeCrescenzo, M. Namchuk, C. M. Emery, S. Saha, D. J. Welsch, Targeting the MAPK Signaling Pathway in Cancer: Promising Preclinical Activity with the Novel Selective ERK1/2 Inhibitor BVD-523 (Ulixertinib). Mol. Cancer Ther. 16 (2017).

106. T. D. Lebedev, E. R. Khabusheva, S. R. Mareeva, K. A. Ivanenko, A. V. Morozov, P. V. Spirin, P. M. Rubtsov, A. V. Snezhkina, A. V. Kudryavtseva, M. I. Sorokin, A. A. Buzdin, V. S. Prassolov, Identification of cell type–specific correlations between ERK activity and cell viability upon treatment with ERK1/2 inhibitors. J. Biol. Chem. 298 (2022).

107. S. Su, G. Chhabra, M. A. Ndiaye, C. K. Singh, T. Ye, W. Huang, C. N. Dewey, V. Setaluri, N. Ahmad, PLK1 and NOTCH Positively Correlate in Melanoma and Their

Combined Inhibition Results in Synergistic Modulations of Key Melanoma Pathways. Mol. Cancer Ther. 20, 161–172 (2021).

108. L. N. Micel, J. J. Tentler, A. C. Tan, H. M. Selby, K. L. Brunkow, K. M. Robertson, S. L. Davis, P. J. Klauck, T. M. Pitts, E. Gangolli, R. Fabrey, S. M. O'Connell, P. W. Vincent, S. G. Eckhardt, Antitumor activity of the MEK inhibitor TAK-733 against melanoma cell lines and patient-derived tumor explants. Mol. Cancer Ther. 14 (2015).

109. M. J. Sale, K. Balmanno, J. Saxena, E. Ozono, K. Wojdyla, R. E. McIntyre, R. Gilley, A. Woroniuk, K. D. Howarth, G. Hughes, J. R. Dry, M. J. Arends, P. Caro, D. Oxley, S. Ashton, D. J. Adams, J. Saez-Rodriguez, P. D. Smith, S. J. Cook, MEK1/2 inhibitor withdrawal reverses acquired resistance driven by BRAFV600E amplification whereas KRASG13D amplification promotes EMT-chemoresistance. Nat. Commun. 10, 1–22 (2019).

110. Y. Cheng, H. Tian, Current Development Status of MEK Inhibitors. Molecules : A Journal of Synthetic Chemistry and Natural Product Chemistry 22 (2017).

111. H. Y. Lim, J. Heo, H. J. Choi, C.-Y. Lin, J.-H. Yoon, C. Hsu, K.-M. Rau, R. T. P. Poon, W. Yeo, J.-W. Park, M. H. Tay, W.-S. Hsieh, C. Kappeler, P. Rajagopalan, H. Krissel, M. Jeffers, C.-J. Yen, W. Y. Tak, A Phase II Study of the Efficacy and Safety of the Combination Therapy of the MEK Inhibitor Refametinib (BAY 86-9766) Plus Sorafenib for Asian Patients with Unresectable Hepatocellular Carcinoma. Clin. Cancer Res. 20, 5976–5985 (2014).

112. T. Tegnebratt, E. Ruge, S. Bader, N. Ishii, S. Aida, Y. Yoshimura, C.-H. Ooi, L. Lu, N. Mitsios, V. Meresse, J. Mulder, M. Pawlak, M. Venturi, J. Tessier, S. Stone-Elander, Evaluation of efficacy of a new MEK inhibitor, RO4987655, in human tumor xenografts by [18F] FDG-PET imaging combined with proteomic approaches. EJNMMI Res. 4 (2014).

Acknowledgments: Madeleine Duran and Cole Trapnell for advice on zebrafish analysis and the list of annotated zebrafish transcription factors. Paul Bertin for suggesting the use of an accuracy score for biological interpretability, additional potential model architectures, and innumerable other helpful insights. Szen Toh for making suggestions for both datasets and embeddings, helping coordinate with the Garnet lab which authored so many of these datasets, and commenting on the initial draft of the manuscript. Tim Treis for invaluable advice on the cell painting datasets. Luke Zappia for helpful and extensive commentary on the figure designs. Stefanie Brandner for excellent technical assistance in the small molecule library experiment. Riccardo Trozzo, Ekaterina Zhigalova, and Juan José Montero Valderrama for biological advice on CRISPR perturbations. Stefan Peidli, Tessa Green, and Camille Fourneaux for advising on content and clarity in the initial draft of the manuscript.

Funding:

European Research Council, grant DeepCell – 101054957

Chan Zuckerberg Initiative Foundation (CZIF; grant CZIF2022-007488 (Human Cell Atlas Data Ecosystem)

Author contributions:

Conceptualization: ATL, YJ, FJT

Methodology: ATL, YJ, FJT, SK

Investigation: SK, YJ, ATL, IR, JT, KH

Visualization: YJ, ATL, XZ, ZZ

Funding Acquisition: FJT

5   Supervision: FJT

Writing – original draft: YJ, ATL, XZ, ZZ

Y.J. and A.T.L. contributed equally and have the right to list their name first in their curriculum vitae. All authors read, corrected, and approved the manuscript.

Competing interests: F.J.T. consults for Immunai Inc., Singularity Bio B.V., CytoReason
10   Ltd, Cellarity, Curie Bio Operations, LLC and has an ownership interest in Dermagnostix GmbH and Cellarity.

Data and materials availability: The code for Prophet can be found at https://github.com/theislab/prophet. Data obtained from in vitro validation can be found at https://figshare.com/articles/dataset/Prophet_viability_plate_xlsx/26516002.

15   **Supplementary Materials:**

Materials and Methods

Supplementary Text

Figs. S1 to S9

Tables S1 to S9

Fig. 1 | Prophet is a universal phenotype predictor. (**A**) Prophet models the experimental space along three axes, namely cellular state, treatments, and phenotypic readout. Each point in the experimental space is mapped to its corresponding phenotypic readout. We illustrate an example of how Prophet samples two different points of the experimental space. The two points differ in the biological system these experiments were performed in: HaCaT and A375 cells.

This flexibility enables Prophet to be trained across datasets. (**B**) Cellular states and treatments are passed to Prophet as prior knowledge-based representations. Phenotypes can be either modeled as a learnable embedding or grounded on prior knowledge. (**C**) Prophet relies on a transformer-based architecture that leverages self-attention and permutation equivariance to represent the experimental context as a set and obtain a meaningful representation that maps to the phenotypic readout. It consists of tokenizer networks that take prior knowledge to create tokens, the transformer to which those tokens are fed and a small regression head that uses the output of the transformer to get a readout prediction. Prophet can predict any quantitative phenotype, including cell viability, gene expression and cell morphology.

Fig. 2 | Prophet's performance and scaling across diverse datasets. (**A**) Prophet-individual models (top) are Prophet models trained only on a single dataset. On the other hand, the full Prophet model (bottom) is pretrained using all datasets and then fine tuned on the dataset of interest. (**B**) Improvement in the R2 score over the mean baseline of the models evaluated: linear regression, multilayer perceptron (MLP), Random Forest, single-assay Prophet (Prophet-individual) and fine-tuned Prophet (Prophet). (**C**) Prophet accurately predicts the IC_{50} value of

the unseen compound mitoxantrone, a clinically approved compound, against different cancers. (**D**) The fluorescent RNA readout of the gene TNFRSF21 is not part of the pre-training corpus. However, fine-tuning Prophet on it still yields better results than the mean estimator and than training a Prophet-individual model only on this dataset. (**E**) In GDSC^2, fine-tuned Prophet outperforms Prophet-individual, the mean estimator and also a Prophet-two-dataset model trained jointly in GDSC^2 and PRISM. Notably, Prophet is fine-tuned for just 20,000 steps with a batch size of 256 observations – much fewer optimization steps than the other models, which highlights the effectiveness of the pre-training stage. (**F**) The area-under the precision-recall curve (AUPRC) is 0.81 for *hit* identification in GDSC - defining each *hit* as a compound that causes at least a 2x fold change in cell viability. (**G**) The variance in the R2 score obtained by Prophet is reduced as more parts of the experimental space - more cell states and more treatments - are sampled during training.

Fig. 3 | Prophet learns a map of the experimental space. (**A**) Schematic of how Prophet generates a complete experimental space *in silico*. For the treatment space, Prophet predicts the IC_{50} for each compound or compound combination across 979 cell lines. Each single or

combinatorial treatment is then represented as a 979-dimensional vector, in which each dimension is Prophet's IC_{50} prediction for that compound in one cell line. We call this treatment representation an "activity-based fingerprint". (**B**) Two dimensional visualization (UMAP) of activity-based fingerprints of 8,620 small molecule treatments, each represented as one point.

5 (**C**) Compounds with differential effects can cluster into four different biological signaling pathways involved in cell proliferation: MAPK, EGFR, PI3K/AKT and mTOR (Table S6). (**D**) Prophet can create experimental subspaces by constraining the cell lines or compounds. For the melanoma and brain cancer subspaces, only cell lines representing the respective tissues are included. (**E**) IC_{50} treatment subspace for melanoma cell lines, with a zoom in on a cluster of

10 compounds containing known cancer treatments. Each dot is a treatment represented as a 23-dimensional vector in which each dimension refers to the predicted IC_{50} value in one of the 23 selected melanoma cell lines. Clinically approved monotherapies for *BRAF* V600E mutant melanoma are highlighted in bold. (**F**) IC_{50} treatment space for a subset of brain and central nervous system cancer cell lines, where calcium channel blockers (zoom in) are clustered

15 together and exhibit a differential effect. (**G**) To model cell line space with Prophet, each cell line is represented as a vector in which each dimension is Prophet's IC_{50} prediction for one compound in that cell line. (**H**) Here we model the activity of EGFR inhibitor treatments in non-small cell lung cancer (NSCLC) cell lines. Each point is a cell line represented as a vector in which each dimension is the predicted IC_{50} value for each treatment in that cell line for a set of

20 NSCLC cell lines and EGFR inhibitors. The cell lines HC4006, NCIH1975, NCIH1650, HCC827 and CALU3 are known to be sensitive to EGFR inhibitors. Points are colored by predicted IC_{50} sensitivity (Methods) to EGFR inhibitors. Lower IC_{50} values imply higher sensitivity and higher cell death.

Fig. 4 | In vitro validation of selected Prophet predictions. (**A**) Workflow accommodating the constraints of an in-house small molecule screening setup. (**B**) Proportion of the performed screen with completely new experiments. Cell line and drug experiments not seen by Prophet in any training dataset shown in gray. Experiments which were measured at least once, with any readout, are shown in red. Each column represents one compound. The two compounds also in panel D are highlighted. (**C**) Spearman correlation per compound of the real vs. predicted viability values across cell lines (teal, right axis) for the 352 compounds we experimentally validated. In yellow are the same correlations, but with the real viability values measured from PRISM. (**D**) Absolute CellTitreGlo fluorescence values from the in vitro experiments (three replicates) vs. Prophet's predicted values (6 ensembles). Boxes show the quartiles, and whiskers show 1.5 times the interquartile range from the box. Prophet provides tight uncertainty bounds for these predictions, and more loose uncertainty for the U2OS cell line.

Fig. 5 | Prophet extrapolates relative cell type abundance in zebrafish development. (**A**) Summary of the original experimental setup from Saunders et al. 2023. (**B**) How the single-cell x gene matrix was converted into cell type proportions per embryo. The latter is the phenotypic readout predicted by Prophet. Adapted from Saunders et al. 2023. (**C**) Schematic of how Prophet generates a complete experimental space *in silico*. Prophet predicts the cell type frequencies of one zebrafish embryo for each genetic KO combination across 124 cell types to construct a genetic KO "fingerprint". (**D**) Prophet's learnt experimental space of genetic KO "fingerprints".. Depicted are: combinatorial predictions which include a knockout seen in the original publication (top panel, colored by the seen knockout; bottom panel, colored by the lineage of the seen knockout) and singleton predictions for zebrafish transcription factors (gray) not seen in the original publication. Prophet's experimental space recapitulates the findings obtained by the authors in the original publication and the separation between genes involved in different developmental lineages. (**E**) Top-10 gene targets that suppress the appearance of the "anterior migratory muscle" cell type as predicted by Prophet, ranked by their average effect on the fold-change of the cell type proportion as compared to control. Rspo2 and Hopx are known regulators of mesoderm development, while Wnt8b and Bmp1a belong to families of genes also involved in the same lineage. (**F**) The predicted absolute cell type proportion of anterior migratory muscle at 5 timepoints for the top 1 and 3 predicted gene targets, with control as comparison.